

4

AD-A210 045



AL-CP-89-01


AD:

**PROCEEDINGS OF THE 10 MAY 1989  
ANTIPROTON TECHNOLOGY WORKSHOP**

**A compilation of presentation materials from the workshop held at Brookhaven National Laboratory, jointly sponsored in accordance with the AL/DoE Memorandum of Agreement for Applied Research In Energy Storage support from Brookhaven National Laboratory**

May 1989

Editor: Gerald D. Nordley



**Approved for Public Release**

cb 5

Distribution is Unlimited. The AFAL Technical Services Office has reviewed this report, and it is releasable to the National Technical Information Service, where it will be available to the general public, including foreign nationals.

**Astronautics Laboratory (AFSC)**

Air Force Space Technology Center  
Space Division, Air Force Systems Command  
Edwards Air Force Base,  
California 93523-5000

89


031

## NOTICE

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related government procurement operation, the government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

## FOREWORD

This special report comprises the presentations provided by speakers at the Antiproton Technology Workshop held at Brookhaven National Laboratory (BNL) 10 May 1989 jointly sponsored under the Astronautics Laboratory (AFSC) / Department of Energy-BNL Memorandum of Agreement for support of Applied Research In Energy Storage (ARIES). This special report has been reviewed and approved in accordance with the distribution statement on the cover and on the DD form 1473.

  
GERALD D. NORDLEY, Major, USAF  
Research Staff Manager, ARIES  
Air Force Astronautics Laboratory

  
SEYMOUR BARON  
Associate Director  
Brookhaven National Laboratory

FOR THE DIRECTOR

  
ROBERT C. CORLEY  
Chief, Astronautical Sciences Division

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release. Distribution is unlimited.			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) AL-CP-89-001		5. MONITORING ORGANIZATION REPORT NUMBER(S)			
6a. NAME OF PERFORMING ORGANIZATION Astronautics Laborator,	6b. OFFICE SYMBOL (if applicable) LSX	7a. NAME OF MONITORING ORGANIZATION			
6c. ADDRESS (City, State, and ZIP Code) Edwards Air Force Base, CA 93523-5000		7b. ADDRESS (City, State, and ZIP Code)			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO. 62302F	PROJECT NO. 5730	TASK NO. 00	WORK UNIT ACCESSION NO. 5E
11. TITLE (Include Security Classification) (U) PROCEEDINGS OF THE 10 MAY 1989 ANTIPROTON TECHNOLOGY WORKSHOP					
12. PERSONAL AUTHOR(S) Nordley, Gerald D., Editor					
13a. TYPE OF REPORT Special	13b. TIME COVERED FROM 89/5/10 TO 89/5/10	14. DATE OF REPORT (Year, Month, Day) 89/05		15. PAGE COUNT	
16. SUPPLEMENTARY NOTATION A compilation of presentation materials from the workshop held at Brookhaven National Laboratory, jointly sponsored IAW AL/DoE Memorandum of Agreement for Applied Research in Energy					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Antiproton Beams, Imaging, NDA/NDE, Radiotherapy, Condensed Antimatter, CP Violation		
20	07				
20	08				
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>This workshop, held at Brookhaven National Laboratory, 10 May 1989, was a follow-on to the Antiproton Science and Technology Workshops held at the RAND Corporation in Santa Monica through October 1987 following the Air Force Project Forecast II initiative in Antiproton Technology. The workshop was attended by about 50 researchers from a wide variety of disciplines, including medicine, particle physics, and the aerospace industry. New, more efficient technology for a variety of scientific, medical, and industrial uses could result from antiproton experiments proposed by workshop participants. Antiprotons are particles of antimatter which release highly penetrating radiation when they are stopped in normal matter. According to presentations at the Antiproton Technology Workshop this radiation can be used, in very small quantities, to image objects and determine their composition and density. In larger amounts, the radiation could be used to kill cancer tumors or produce highly localized heating and shock waves. DOE plans are contingent on potential user support.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Gerald D. Nordley, Maj, USAF		22b. TELEPHONE (Include Area Code) (805) 275-5653	22c. OFFICE SYMBOL LSX		

Block 16.

Storage support from Brookhaven National Laboratory.



Accession For	
100	<input checked="" type="checkbox"/>
200	<input type="checkbox"/>
300	<input type="checkbox"/>
400	<input type="checkbox"/>
500	<input type="checkbox"/>
600	<input type="checkbox"/>
700	<input type="checkbox"/>
800	<input type="checkbox"/>
900	<input type="checkbox"/>
Disc	<input type="checkbox"/>
A-1	<input type="checkbox"/>

# PROCEEDINGS OF THE ANTIPROTON TECHNOLOGY WORKSHOP\*

HELD AT BROOKHAVEN NATIONAL LABORATORY 10 MAY 1989

## EXECUTIVE SUMMARY

### 1. Background

New, more efficient technology for a variety of scientific, medical, and industrial uses could result from antiproton experiments proposed by a workshop of government, industry and academic researchers at Brookhaven National Laboratory, Wednesday 10 May 1989. Antiprotons are particles of antimatter which release highly penetrating radiation when they are stopped in normal matter. According to presentations at the Antiproton Technology Workshop this radiation can be used, in very small quantities, to image objects and determine their composition and density. In larger amounts, the radiation could be used to kill cancer tumors or produce highly localized heating and shock waves. The Alternate Gradient Synchrotron, or "AGS", located at Brookhaven is one of the few particle accelerators in the world capable of making the number of antiprotons needed to perform the experiments.

The workshop was a follow-on to the Antiproton Science and Technology Workshops held at the RAND Corporation in Santa Monica through October 1987 following the Air Force Project Forecast II initiative in Antiproton Technology. The workshop was attended by about 50 researchers from a wide variety of disciplines, including medicine, particle physics, and the aerospace industry.

### 2. Workshop Results

Aerospace uses include the detection of physical or chemical flaws in the manufacture of composite materials, with implications for increased aviation safety and lighter, less expensive rockets.

An existing market of about \$100 billion a year in medical imaging and radiotherapy has attracted the interest of private investors. Demonstrations of rapid, low radiation imaging of hard tissues and killing cancer tumors might prove the viability of a new, privately funded accelerator to provide antiprotons for medical and industrial uses.

Atomic chemists want to make antihydrogen to see if it obeys the same physical rules as ordinary hydrogen. Antihydrogen would be made by combining antiprotons with the anti-electron, or positron, the first form of antimatter discovered back in 1935.

Physical scientists are interested in radiation effects and small but intense shock waves that could be produced by pulsed antiproton beams. Protection of spacecraft from solar storms and meteor impacts are among many uses of radiation and shock data.

Particle physicists are interested in broken symmetries in particle reactions which one might expect to have identical outcomes, but don't. Such reactions help tell us how the universe was made and what its ultimate destiny might be.

Antiproton Workshop members came from organizations as diverse as the Lahey Clinic in Boston, the Astronautics Laboratory at Edwards Air Force Base, General Dynamics Corporation in Fort Worth, and the University of Illinois. The workshop agenda is provided as table 1. Workshop attendance is provided as table 2.

The only source of antiprotons suitable for many of the experiments discussed is the European accelerator in Switzerland, which has long waiting lines for experimenters. The researchers generally agreed that an antiproton source in the United States, perhaps based on the Brookhaven AGS accelerator, Fermilab's accelerator, or the booster ring planned for the Superconducting Supercollider will make United States science and technology significantly more competitive in areas discussed. Significant informational activities concerning antiproton technology continue within DOE. Potential user interest expressed as serious proposals is a significant determinant of DOE support.

\*The workshop was jointly sponsored by the Astronautics Laboratory and Brookhaven National Laboratory (DOE).

Table 1. Antiproton Technology Working Group Final Agenda

0830	Informal discussions		
0900	Welcome and Administrative Remarks	Dr S. Baron, BNL; Maj G. Nordley, AL	
0915	IMAGING AND ANALYSIS - T. KALOGEROPOLOUS, SYRACUSE		
0915	Stopping Power of MeV Proton and Antiproton Beams	R. A. Lewis	The Pennsylvania State U.
0930	Recent Simulation Results of ASTER	Robert Muratore	Syracuse University
0945	Pbar Testing of Hydrogen Effects in Sealed Carbon-Carbon Composites	Harris Carter	Gen Dynamics Ft Worth
1000	Potential for Antiprotons in Radiation Oncology	M. Leibenhaut, MD	Lahey Clinic Medical Cen.
1015	Prospects for a Commercial Antiproton Source	Brian Von Herzen	Antimatter Technology Corp
1030	Break		
1045	ANTIHYDROGEN AND CONDENSED MATTER PHYSICS - CHRIS BRASIER, U. DAYTON (AL)		
1045	Prospects for Exciting Extreme States in Nuclear Matter with Intense Antiproton Beams	E. D. Minor	The Pennsylvania State U.
1100	Status of AL Studies Relating to Condensed Antimatter	Gerald Nordley	Astronautics Lab (AFSC)
1115	Electromagnetic Traps for Atomic Antihydrogen	Isaac Silvera	Harvard University
1130	Antihydrogen Production	Arthur Rich	University of Michigan
1200	LUNCH: BNL CAFETERIA		
1215	Luncheon Speech: HQ DoE Antiproton Activities	David Goodwin	Dept of Energy
1300	OPTION: AGS TOUR OR INFORMAL DISCUSSIONS		
1400	ENERGY DEPOSITION AND RELEASE - GERALD SMITH, PENN STATE		
1400	Antiproton Catalyzed Fusion	T. Kalogeropoulos	Syracuse University
1415	Antiproton Induced Fusion Reaction	W. S. Toothacker	The Pennsylvania State U.
1430	Options for a Laboratory Microfusion Facility (LMF)	Bruno Augenstein	The RAND Corporation
1445	Modeling Antiproton - Plasma Interactions	John Callas	Jet Propulsion Laboratory
1500	Concepts for Experimental Determination of Radiation Shielding and Metal Clad Pellet Performance	Brice Cassenti	UTRC - Hartford
1515	Break		
1530	PARTICLE PHYSICS - D. C. PEASLEE UNIVERSITY OF MARYLAND		
1530	Introduction to CP Violation Studies with Pbars	D. C. Peaslee	University of Maryland
1545	Test of CP Non-conservation in Pbar-P to $\Xi$ bar- $\Xi$	A. M. Nathan	University of Illinois
1600	Studies of CP Violation with Pure $K_0$ $K_0$ bar Beams from Pbars	James Miller	Boston University
1615	Search for CP Violation in Pbar-P to $J/\psi$	Gerald A. Smith	The Pennsylvania State U.
1630	Studies of Rare Modes of Pbar-P Annihilation	C. B. Dover	Brookhaven N.L.
1645	Antiproton Production Calculation by the Multistring Model VENUS Computer Code	H. Takahashi	Brookhaven N. L.
1700	Closing Remarks	G. Nordley	AL (AFSC)

Table 2. Attendees at the Antiproton Technology Workshop

Brookhaven National Laboratory Dr. Hiroshi Takahashi Bldg 130 Upton NY 11973	Syracuse University Prof T. Kalogeropoulos Dept of Physics Syracuse NY 13244-1130	Los Alamos National Laboratory Dr Nick King P-15 P.O. Box 1663 Los Alamos NM 87545	RAND Corporation Dr Bruno Augenstein 1700 Main St Santa Monica CA 90406-2138	Kip Tarpley 4313 Knox Ave, #206 College Park MD 20740
Brookhaven National Laboratory Dr James R. Powell Dept of Nuclear Energy Upton NY 11973	Syracuse University Robert Muratore Dept of Physics Syracuse NY 13244-1130	United Technologies Research Center Dr Bruce N. Cassenti MS 18 Silver Lane East Hartford CT 06108	McDonnell Douglas Astronautics V.E. (Bill) Haloulakos MS 13-3 5301 Bolsa Ave Huntington Beach CA 92647	Rockwell International, Rocketdyne Div Mr. Jim McClanahan M/C WB389 6633 Canoga Ave Canoga Park CA 91304
Brookhaven National Laboratory Orto W. Lazareth Bldg 701 Upton NY 11973	Pennsylvania State University Dr Gerald A. Smith 303 Osmond Lab University Park PA 16802	Harvard University Dr Isaac Silveira, Dept of Physics 44 Oxford St Cambridge MA 02138	Antimatter Technology Corp Brian Van Herzen 2379 Kalamianoaale Ave Hilo HI 96720	Charles R. Pellegrino 360 Shore Rd, 31 Long Beach NY 11561
Brookhaven National Laboratory Hans Ludewig Bldg 701 Upton NY 11973	Pennsylvania State University R.A. Lewis 303 Osmond Laboratory University Park PA 16802	University of Virginia Prof Stephen T. Thornton Dept of Physics Charlottesville VA 22901	General Dynamics Harris Carter P.O. Box 748 Fort Worth TX 77251	
Brookhaven National Laboratory Dr Peter Hausstein Bldg 555A Upton NY 11973	Pennsylvania State University E.D. Minor 303 Osmond Laboratory University Park PA 16802	University of Michigan R.A. Rich Physics Dept Ann Arbor MI 48109-1120	Rice University Prof B.E. Bonner Bonner Nuclear Laboratory Houston TX 77251-1892	
U.S. Dept of Energy Dave Goodwin ER20.1/GTN High Energy & Nuclear Physics Washington DC 20545	Pennsylvania State University W.S. Toothacker 303 Osmond Laboratory University Park PA 16802	University of Wisconsin Dan Reeder Dept of Physics 1150 University Ave Madison WI 53706	AL/LSX Maj Gerald D. Nordley Edwards AFB CA 93523-5000	
University of Maryland Prof D.C. Peaslee Dept of Physics & Astronomy College Park MD 20742-3015	Brookhaven National Laboratory Dr C. Dover Physics Dept Upton NY 11973	NASA Lewis Research Center Michael LaPointe 21000 Brookpark Rd Cleveland OH 44135	AL/LSX Dr Patrick Carrick, UDRI Edwards AFB CA 93523-5000	
University of Illinois Dr George H. Miley, Nuc Eng Lab 103 S. Goodwin Ave Urbana IL 61801-2984	Brookhaven National Laboratory M. Divadeenam Bldg 902 Upton NY 11973	Dr Robert L. Forward Forward Unlimited P.O. Box 2783 Malibu CA 90265-7783	AL/LSX Dr Chris Brazier, UDRI Edwards AFB CA 93523-5000	
University of Illinois A.M. Nathan Dept of Physics Champaign IL 61801-2984	Brookhaven National Laboratory Dr Mark Sakitt Physics Dept, Bldg 510A Upton NY 11973	Jet Propulsion Laboratory John Callas MS 248-100 4800 Oak Grove Dr Pasadena CA 91109	Eastern New Mexico University Dr John M. Kenney Dept of Chemistry Portales NM 88130	
				Eastern New Mexico University Dr M. Inga Kenney Dept of Chemistry Portales NM 88130



## CONTENTS

<u>Presentation</u>	<u>Page</u>
Stopping Power of MeV Proton and Antiproton Beams R. A. Lewis, The Pennsylvania State U.	1*
Recent Simulation Results of ASTER Robert Muratore, Syracuse University	2
Pbar Testing of Hydrogen Effects in Sealed Carbon-Carbon Composites Harris Carter, Gen Dynamics Ft Worth	22
Potential for Antiprotons in Radiation Oncology Mark Leibenhaut, MD, Lahey Clinic Medical Cen.	30
Prospects for a Commercial Antiproton Source Brian Von Herzen, Antimatter Technology Corp	46
Prospects for Exciting Extreme States in Nuclear Matter with Intense Antiproton Beams E. D. Minor, The Pennsylvania State U.	59
Status of AL Studies Relating to Condensed Antimatter Gerald Nordley, Astronautics Lab (AFSC)	71
Electromagnetic Traps for Atomic Antihydrogen Isaac Silvera, Harvard University	83
Antihydrogen Production Arthur Rich, University of Michigan	97
Headquarters DoE Antiproton Activities David Goodwin, Dept of Energy	114
Antiproton Catalyzed Fusion T. Kalogeropoulos, Syracuse University	126
Antiproton Induced Fusion Reaction W. S. Toothacker, The Pennsylvania State U.	127*
Options for a Laboratory Microfusion Facility (LMF) Bruno Augenstein, The RAND Corporation	128
Modeling Antiproton - Plasma Interactions John Callas, Jet Propulsion Laboratory	135
Concepts for Experimental Determination of Radiation Shielding and Metal Clad Pellet Performance Brice Cassenti, UTRC - Hartford	142
Introduction to CP Violation Studies with Pbars D. C. Peaslee, University of Maryland	162

Test of CP Non-conservation in $P\bar{P}$ to $\Xi\bar{\Xi}$ A. M. Nathan , University of Illinois	167
Studies of CP Violation with Pure $K_0 K_0\bar{}$ Beams from Pbars James Miller, Boston University	168*
Search for CP Violation in $P\bar{P}$ to $J/\Psi$ Gerald A. Smith, The Pennsylvania State U.	169
Studies of Rare Modes of $P\bar{P}$ Annihilation C. B. Dover, Brookhaven N.L.	180
Antiproton Production Calculation by the Multistring Model VENUS Computer Code H. Takahashi, Brookhaven N. L.	195

\* Copies of viewgraphs were unavailable at the time of compilation (17 May 1989). They may be inserted if recieved later.

**STOPPING POWER OF MeV PROTON  
AND ANTIPROTON BEAMS**

**R. A. LEWIS**

**LABORATORY FOR ELEMENTARY PARTICLE SCIENCE  
THE PENNSYLVANIA STATE UNIVERSITY  
UNIVERSITY PARK, PA**

Note: We regret that copies of the transparencies used in Dr Lewis' excellent presentation were not available for inclusion in the proceedings.

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**RECENT SIMULATION RESULTS OF ASTER**

**ROBERT MURATORE**

**DEPARTMENT OF PHYSICS  
SYRACUSE UNIVERSITY**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

R e c e n t

S i m u l a t i o n

R e s u l t s

o f A S T E R

Robert Muratore  
Syracuse University

R E C E N T   S I M U L A T I O N

R E S U L T S   O F   A S T E R

**Robert Muratore**

Department of Physics, Syracuse University

201 Physics Building, Syracuse, New York 13244-1130 USA

*Abstract.* ASTER, an imaging technique proposed several years ago, is now ready to be built. ASTER uses antiprotons to form direct three dimensional images of the target density profile. Useful images can be obtained with less than one million antiprotons, well within current production levels. ASTER has potential advantages over other imaging techniques, including flexibility, speed, lower dose, and less ambiguity. Simulations show that the scattering of antiprotons by target nuclei reduces the correlation of image and target, but increasing the number of antiprotons used by less than an order of magnitude overcomes this effect.

WHEN COMPLICATED TECHNOLOGY is used in medicine, reassuring names are attached to the machines and techniques. One speaks of CAT scans, PET, and MRI (née NMR). Today I will talk about an imaging technique which has been discussed before at these meetings, ASTER, named after the wildflower. Since I am limited to about ten minutes, I will keep my talk simple. Here is the outline:

**I. ASTER is ripe.**

It is my contention that this flower has formed its fruit, and that not only is this fruit ripe for picking, but neither is it spoiled, as some have suggested.

*A. ASTER uses antiprotons to image densities, and enough antiprotons are currently produced.*

I will begin by reviewing ASTER.<sup>1,2,3,4,5</sup> ASTER is an acronym for Antiprotonic STERiography. In the ASTER imaging technique, still on the drawing board,

a beam of antiprotons are sent into a target. Collisions with electrons slow the antiprotons down, according to the well known stopping power

$$\frac{dE}{dx} = D\rho \frac{Z}{A} \frac{1}{\beta^2} \left[ \ln \frac{2m_e c^2 \beta^2}{I(1-\beta^2)} - \beta^2 \right],$$

where  $E$  is the kinetic energy of the particle,  $x$  is the distance traversed,  $Z$  is the proton number,  $A$  is the atomic mass,  $\beta$  is the speed relative to the speed of light,  $D$  is a constant approximately equal to  $0.30707 \text{ MeV cm}^2/\text{g}$ ,  $\rho$  is the density,  $m_e$  is the electron mass, and  $I$  is an empirical function of  $Z$  which represents the average ionization potential of *all* electrons in an atom.<sup>6</sup>

The important features are the inverse square relation of the stopping power and the speed, which results in the Bragg peak, and the direct dependence on the density.

ASTER (a third definition here) means star (as in \*). When the antiprotons have come to rest, they annihilate on a nucleon. Outward from the annihilation site stream various particles. In a bubble chamber photo, this event looks like a star (Fig. 1). Among the particles produced are charged pions. These are of sufficient energy to exit a target the size of the human body, and of sufficient mass to be deflected just a small amount before emerging. By detecting the directions of these pions and tracing their paths back to the intersection point with the antiproton path, the annihilation site can be determined precisely.

In this way, the range as a function of energy,  $R(E)$ , can be determined for the target, and  $R(E)$  can be mapped to  $\rho(R)$ , a density profile.

Simulations of ASTER imaging confirm the estimates of the number of antiprotons needed for a scan,  $N$ :

$$N \sim \frac{\text{volume}}{\Delta x \Delta y \Delta z} \times \left( \frac{\sigma_V \rho}{\Delta x \delta \rho} \right)^2$$

where the antiprotons are assumed to be travelling initially in the  $x$  direction,  $\Delta x$ ,  $\Delta y$ , and  $\Delta z$  are the step sizes with which the beam is incremented in the various directions,  $\sigma_V$  is the error in determining the vertex, and  $\delta \rho/\rho$  is the contrast resolution. To image a slice of  $10 \times 10 \times 0.5 \text{ cm}^3$  requires  $2 \times 10^5$  antiprotons, for 1% contrast resolution and 1.5 mm spatial resolution within the slice. To image a whole

organ might require 20 slices, or  $4 \times 10^6$  antiprotons, well within current production levels. The corresponding dose is about  $200 \mu\text{Gy} = 0.02 \text{ rads}$ . Considering the biological effect of protons, the dose is about a tenth of the natural average annual background in the United States.<sup>6</sup>

*B. ASTER has advantages over other imaging techniques.*

ASTER appears to be lower in dose for comparable images than x-ray CT, as shown in a comparison of an ASTER simulation (Fig. 2) of the imaging of a Plexiglas and water phantom and the actual x-ray CT image of the same phantom. The phantom is an 8 cm diameter Plexiglas disk inside a 10 cm diameter Plexiglas cylinder filled with water. In the 3 mm thick disk the letter E is engraved to a depth of 1.5 mm. In the simulation, this cylinder was immersed in a rectangle filled with water. An x-ray CT scan (Fig. 3) was made of the cylinder in the plane containing the engraved disk. The dose imparted by the ASTER simulation was  $100 \mu\text{Gy}$ , over two orders of magnitude less than that imparted by the CT scan, approximately 30 mGy.

The table in Fig. 4 gives an overview of ASTER with other techniques. No one technique seems better than all the others for every situation. Similarly, ASTER will be complementary to the other techniques. Nonetheless, ASTER has potential for lower dose, higher resolution, faster scans, and imaging of elements as well as density. Perhaps most importantly, ASTER avoids the uncertainties introduced by back-projection techniques. Finally, ASTER is a flexible technique, as the following discussion shows.

*C. The scattering of antiprotons does not spoil the image quality.*

There has been some question as to whether the scattering of the antiprotons off the nuclei will irretrievably lower the resolution of ASTER. In water, the antiproton beam spreads out with a width of  $\sigma_y = 0.0195R^{0.966}$ . There is a well defined centroid, so resolution can be maintained by increasing the number of antiprotons used. In heterogeneous media, one can imagine that some of the antiproton paths will sample regions of different density, hopelessly convoluting the relation of stopping position to density profile. However, this is not the case, as I will show by considering individual antiproton paths in water, and by showing successful images



of highly heterogeneous targets.

In terms of individual paths, it is reasonable that transverse scattering will not ruin ASTER images. This is because the average density in a small region is obtained from the difference in the mean stopping positions of two cohorts of antiprotons with nearly the same energy. In engineering terms, one would say that one is looking at the difference between two integrals, and integration suppresses the noise.

Fig. 5 shows the paths of many antiprotons in water. The horizontal (longitudinal) and vertical (transverse) scales are the same, and the three dimensional paths have been projected onto the plane. Next, I considered only the antiprotons stopping in a small transverse bin. If the initial energy of the antiprotons is varied just a bit, the antiprotons still sample the same region in space. That is, a group of paths stopping about  $R$  tends to sample the same portion of the target as a group of paths stopping about  $R + \Delta R$ . This is shown in Figs. 6, 7, 8, and 9. This is true even though I have included the finite beam width in the Monte Carlo.

The sampling of the same region in space by the antiprotons is a statistical phenomenon. Therefore, it suggests that the scattering problem can be overcome by increasing the number of antiprotons, a method already required by the straggling. To test this, I simulated the imaging of a "random" target, which was the most heterogeneous thing I could think of. ASTER is a very flexible tool, and the imaging can be oriented to best advantage. I imaged this random slice longitudinally, so that each antiproton travelled in the slice. And I imaged the random slice transversally, so that the antiprotons travelled through a centimeter of water before encountering the slice perpendicular to the plane. The transverse orientation is shown in Fig. 10, the random target in Fig. 11, and the transverse image in Fig. 12.

The heterogeneous nature of this target convolutes the longitudinal image more than the transverse image. So for a given number of antiprotons, the transverse image will be better.

The quality of the image can be shown by correlating the image and the target. I define a correlation number  $C^{-1}$ , where

$$C^2 \equiv \sum_j \sum_k (\rho_{1jk} - \rho_{2jk})^2 / n,$$

$\rho_{1jk}$  and  $\rho_{2jk}$  are the densities of the target and image at the  $jk$ th pixel, and  $n$  is the number of pixels used for comparison. The correlation increases as the number of antiprotons increases for both the transverse and longitudinal imaging (Fig. 13). The correlations match when about four to nine times as many antiprotons are used in the longitudinal case as in the transverse case. So increasing the number of antiprotons by an order of magnitude will overcome severe heterogenous effects. After this increase is made, ASTER still imparts an order of magnitude less dose than x-ray CT.

If the decrease in correlation is due to the heterogeneous convolutions, than the effect will begin to show up in the transverse case as the slice is lowered deeper into the water, so that the antiproton beam is more spread out when it reaches the slice. This is shown in Fig. 13 by the decrease in correlation between transverse image and target as the depth increases.

#### References

- 1 deGuzman, Allan F. Ph.D. dissertation, Syracuse University, 1986.
- 2 Gray, L., and Kalogeropoulos, T. E. *IEEE Trans Nucl Sci NS-29* (1982) 1051-1057.
- 3 Kalogeropoulos, T.; Archambeau, J.; Bassano, D.; Bennett, G.; Gottchalk, B.; Gray, L.; Koehler, A.; Muratore, R.; and Urie, M. In *Proceedings of the RAND Workshop on Antiproton Science and Technology, October 6-9, 1987*, edited by B. W. Augenstein, B. E. Bonner, F. E. Mills, and M. M. Nieto. Teaneck, NJ: World Scientific, 1988.
- 4 Kalogeropoulos, Theodore E., and Muratore, Robert. *Nuclear Instruments and Methods* (1989) *in press*.
- 5 Muratore, Robert. Ph.D. dissertation, Syracuse University, 1988.
- 6 Particle Physics Group. *Phys Lett B204* (1988) 1-486.



ASTFRT.flx.11:

18-NOV-1988 22:12:03.39

total # pbars stopping =	194811	
total # pbars injected =	212973	
	file =	E.dat.3
height above slice (cm) =	1.000	
slice width along beam (cm) =	0.500	
segment length (cm) =	0.050	
percent error in density =	1.000	
white, black densities (g/cm**3) =	1.100	1.050
horiz, vert magnifications =	1.000	1.000

50  
 $2 \times 10^{-4}$  Gy




Fig.2




Fig. 3

Table: IMAGING TECHNIQUES					
system	CT <sup>a</sup>	MRI	ultrasound <sup>a</sup>	PET	ASTER
detected	transmitted x-rays	rf from induced emf	acoustic echoes	$\gamma$ -rays	$\pi^{\pm}$ (& x-rays)
imaged	electron density	induced nuclear magnetization	discontinuities in speed of sound	tagged chemical concentration	electron density (& elements)
structure	"	"	density, elasticity	biochemical	"
inference	transform	tran. or ver.	transform	transform	vertex
source	x-ray tube	precession	transducer	decay	$\bar{p}$
detector	x-ray det.	rf coil	transducer	$\gamma$ -ray det.	drift ch.
spatial resolution	0.5 mm	2 mm	2 mm	~ 1 mm	< 0.5 mm
temporal resolution	0.5 s	0.1 to 100 s	0.01 s	10 to 1000 s	< 0.01 s
dose	0.03 Gy	(nonionizing)	(nonionizing)	varies	0.0001 Gy
<sup>a</sup> Fullerton, Gary D., and Zagzebski, James A., eds. <i>Medical Physics of CT &amp; Ultrasound</i> . New York: American Institute of Physics, 1980.					

Fig. 4

follow.for.26

3-MAY-1989 11:00:43.39

diamtr	rhom	R	dR	magy	ncatch	up	down
0.100	1.000	7.500	0.050	1.000	20	10.000	-10.000




Fig.5

follow.for.26

3-MAY-1989 11:10:19.23

diamtr	rhom	R	dR	magy	ncatch	up	down
0.100	1.000	7.500	0.050	10.000	15	0.100	0.000




Fig.6



follow.for.26

3-MAY-1989 11:11:45.57

diamtr	rhom	R	dR	magy	ncatch	up	down
0.100	1.000	7.250	0.050	10.000	15	0.100	0.000




Fig.7

follow.for.26

3-MAY-1989 11:24:48.78

diamtr	rhom	R	dR	magy	ncatch	up	down
0.100	1.000	6.000	0.100	2.000	8	-0.300	-0.400




Fig.8

follow.for.26

3-MAY-1989 11:27:25.99

diamtr	rhom	R	dR	magy	ncatch	up	down
0.100	1.000	6.500	0.100	2.000	8	-0.300	-0.400





Fig.9



$\bar{P}$

Fig. 10

ASTERT.flx.11

1-NOV-1988 10:11:26.53

file = random2.dat.1

white, black densities (g/cm\*\*3) = 1.050 0.950  
horiz, vert magnifications = 1.000 1.000




Fig. 11


ASTERT.flx.11

2-NOV-1988 13:20:56.24

file = astert.dat.7

target = random2.dat.1

white, black densities (1/cm\*\*3) = 1.050 0.950  
horiz, vert magnifications = 1.000 1.000



$$\frac{\delta p}{\rho} = 1\%$$

$$\text{depth} = 1 \text{ cm}$$

$$\text{slice thickness} = 0.2 \text{ cm}$$

$$N_p = 4.7 \times 10^5$$

$$\left(\frac{N_p}{\text{pixel}}\right) = 175$$

$$\text{segment length} = 0.2 \text{ cm}$$

Fig.12




Fig.13

***Pbar TESTING OF HYDROGEN EFFECTS  
IN SEALED CARBON-CARBON COMPOSITES***

***HARRIS CARTER***

***GENERAL DYNAMICS FORT WORTH  
FORT WORTH, TX***

**PRESENTED AT THE ANTI-PROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**



UNCLASSIFIED

## Rationale and Concept for NDE Application of Antiprotons ( $\bar{u}$ )

- Early applications might include non-destructive evaluation (NDE) of aerospace materials
- Advanced carbon-carbon structures present
  - ✓ Special NDE problems (e.g. need for "backscatter" rather than transmission)
  - ✓ Special features suggesting  $\bar{p}$ 's for NDE (vs ultrasound or x-rays)

**Assumption:** Useful Sources will be  $\bar{p}$ 's trapped at  $\leq$  Kev energies or stabilized in chemical complexes

**Suggested NDE mode:** Use low energy  $\bar{p}$ 's as portable source of  $\Pi^-$  mesons

**Purpose:** Determine atomic ratios O/C and H/C deep in Carbon - Carbon structures

Annihilation at source:  $\bar{p} + p \rightarrow \Pi^0 + \Pi^+ \Pi^-$  (K.E.  $\sim$  250 Mev)

**Reactions:**

Reactions in Target:  $\Pi^-$  (stopped) +  $p \rightarrow n + \gamma$  (129 Mev)

$\Pi^-$  (stopped) +  $O \rightarrow \Pi^-$  O mesic atom + 178 Kev x-ray

UNCLASSIFIED

BN06752

UNCLASSIFIED

© 1989 GDFW

**ADVANCED CARBON-CARBON: EDGE VIEW SHOWING  
CRACKS AND VOIDS**



**ADVANCED CARBON-CARBON: HIGHER MAGNIFICATION SHOWING  
INCOMPLETE REACTION IN CONVERSION COAT**



0.1 mm


B18087

© 1984 GDFW


UNCLASSIFIED

# Need for NDE to Sample Atomic Ratios in Thick C-C Structures (u)

a) Residual Oxygen and Hydrogen Indicate incomplete pyrolysis



b) Hydrogenation of C-C in high temp H<sub>2</sub> environment is possible reversion mechanism



• Energetic  $\pi^-$  from  $\bar{p}$  + p annihilation could reveal (a) poor cure or (b) in use hydrogenation at points deep in Carbon-Carbon structure by  $\gamma$  and x-ray backscatter


UNCLASSIFIED

BN06753

UNCLASSIFIED (C) 1989 GDFW

UNCLASSIFIED

# Proposed Deep-Target Chemical Diagnostics using $\Pi^-$ from $\bar{p}$ Annihilation (u)



• In practical case, x-ray counts from  $10^8$   $\bar{p}$  might not exceed  $10^3$  ; but:


- ✓ Effective clutter would be low due to gating & low count rate
- ✓ X-rays from, say,  $\Pi^-$  O could be easily counted and identified

UNCLASSIFIED

BN06754

UNCLASSIFIED © 1984 6DFW

RANGE OF NEGATIVE PIONS IN C-C;  
NON-RELATIVISTIC ESTIMATE.



KINETIC ENERGY (MEV)

© 1984 GDFW

UNCLASSIFIED

# X-Rays from $2P \rightarrow 1S$ Transitions in $\Pi^-$ Mesic Atoms and Attenuation in Carbon-Carbon ( $\mu$ )

$\Pi^-$ Atom	X-ray energy (KEV)	Mass abs coef in C-C ( $\rho = 1.5 \text{ gm/cc}$ ): $\mu(\text{cm}^{-1})$
H	2.4	$\sim 150$
C	100	.23
O	178	.19
(Fluorescent X-ray from normal O atom)	(.65)	( $\sim 1000$ )

29

C(100 Kev) = 1/3.1  
O(178 Kev) = 1/2.6

X-ray line degradation through 5 cm of C-C:

- Oxygen, implying poor cure, could be identified in thick (e.g., 5 cm) C-C structures
- X-rays from  $\Pi^-$  H would not be observed; presence of H might be inferred from  $\Pi^-$  C reduction due to competitive orbital capture of  $\Pi^-$  by H

UNCLASSIFIED

BN06751

UNCLASSIFIED

**THEORETICAL POTENTIAL OF ANTIPROTONS IN  
RADIATION ONCOLOGY**

**Mark H. Liebenhaut, M.D.**

**Department of Radiation Therapy  
Lahey Clinic Medical Center  
Boston MA**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**



### Potential Radiation Damage

A) Direct Effects - Damage DNA Itself

B) Indirect Effects - Form Free Radicals

OH•

DNA•

DNA• + O<sub>2</sub> → DNAOO•


Oxygen Enhancement Ratio =  $\frac{\text{Dose Hypoxic}}{\text{Dose in Oxygen}}$

x-rays OER = 2.5-3  
neutrons OER = 1.6


$$\text{Relative Biological Effect} = \frac{\text{Dose}_{250}}{\text{Dose}_r}$$

RBE varies with: 1) system or tumor studied  
2) amount of damage in that system

**FIG. 6-11.** Variation of the OER and the RBE as a function of the LET of the radiation involved. The data were obtained by using T<sub>1</sub> kidney cells of human origin, irradiated with various naturally occurring  $\alpha$ -particles or with deuterons accelerated in the Hammersmith cyclotron. Note that the rapid increase of RBE and the rapid fall of OER both occur at about the same LET, namely about 100 keV/ $\mu$ . (Redrawn from Barendsen GW: in Proceedings of the Conference on Particle Accelerators in Radiation Therapy. US Atomic Energy Commission, Technical Information Center, LA-5180-C, October 1972, pp 120-125)



Source: Hall, Eric J., *Radiobiology for the Radiologist*, 2nd. ed., Harper & Row, Philadelphia, pg. 109.



**Figure 9.3.** Central axis depth dose distribution for different quality photon beams. Field size, 10 x 10 cm; SSD = 100 cm for all beams except for 3.0 mm Cu HVL, SSD = 50 cm. Data are from Reference 13 and the Appendix.

Source: Khan, Faiz M., *The Physics of Radiation Therapy*, Williams & Wilkins, Baltimore, pg. 161.

ELECTRON BEAM CENTRAL AXIS DEPTH DOSE





Figure 14.8. Comparison of central axis depth dose distributions of the Sagittaire linear accelerator (continuous curves) and the Siemen's betatron (dashed curve). [Reprinted with permission from: Tapley (35).]

Source: Khan, pg. 314.



Source: Hall, pg. 296.




FIG. 15-6. Depth-dose curve for 187-MeV protons from the Uppsala synchrocyclotron. The dose reaches a sharp peak at a depth of about 23 cm. (Redrawn from Larsson B: Br J Radiol 34:143-151, 1961)

Source: Hall, pg. 311.








FIG. 15-7. Illustrating the way in which the Bragg peak for a proton beam can be spread out. Curve A is the depth-dose distribution for the primary beam of 160-MeV protons at the Harvard cyclotron, which has a half-width of only 0.6 cm. Beams of lower intensity and shorter range, as illustrated by curves B, C, D, and E, can be added to give a composite curve S, which results in a uniform dose over 2.8 cm. The broadening of the peak is achieved by passing the beam through a rotating wheel with sectors of varying thickness. (Redrawn from Koehler AM, Preston WM: Radiology 104:191-195, 1972)

Source: Hall, pg. 312.




**FIG. 15-8.** Cross-section of the dose distribution that can be obtained in the treatment of an imaginary carcinoma of the cervix, using a four-field technique with  $^{60}\text{Co}$   $\gamma$ -rays, 11-MeV x-rays from a betatron, and 160-MeV protons from the Harvard cyclotron. (From Koehler AM, Preston WM: Radiology 104:191-195, 1972)

Source: Hall, pg. 313.




A




B

Source: Hall, pg. 24.




**FIG. 15-10.** Depth-dose curve for carbon ions in which the Bragg peak has been spread out over 10 cm by the use of a ridge filter. The ions had an initial energy of 400 MeV/nucleon, corresponding to a total energy of 4.8 GeV. The spread-out peak is located between 12 and 22 cm deep. The lower curve represents the physical absorbed dose. The upper curve represents the biologically effective dose; it is, in fact, the product of dose and RBE, calculated at the level of 50% cell survival. (By courtesy of Dr. J.D. Chapman)

Source: Hall, pg. 318.




**Figure 3.** Variation of energy deposition by beams of protons and antiprotons with depth in an absorber. Each curve normalised to 1 at a depth of  $0.5 \text{ g cm}^{-2}$ .

Source: Sullivan, A. H., "A Measurement of the local energy deposition by antiprotons coming to rest in tissue-like material," *Phys. Med. Biol.*, 1985; 30, 1297-1303.



Source: Gray, L. and Kalogeropoulos, T.E., " Possible Biomedical Applications of Antiproton Beams: Focused Radiation Transfer," Radiation Research, 1984: 97, 246-252.



DOSE DISTRIBUTION ADVANTAGE →

Source: Hall, pg. 320.

**PROSPECTS FOR A COMMERCIAL  
ANTIPROTON SOURCE**

**BRIAN VON HERZEN**

**ANTIMATTER TECHNOLOGY CORPORATION  
HILO, HI**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**



**Prospects for a Commercial  
Antiproton Source**

**Brian Von Herzen, Ph.D.**

**Antimatter**  
TECHNOLOGY CORPORATION

## **Objective:**

**To develop the production facilities, transport systems, and equipment needed to apply antimatter to problems in medicine, aerospace, and academic research.**

# Current Efforts

- to obtain complete funding for the production, distribution, and application of antimatter.
- to develop a cost-effective source for antiprotons.
- to develop a portable system capable of storing antiprotons and delivering them to remote sites.
- to develop the necessary imaging, diagnostic and therapeutic equipment for medical applications.

# Funding Being Sought

**\$195 million for a dedicated production facility, or an extension of an existing facility**

**\$15 million to develop a portable storage device capable of transporting antiprotons to remote sites.**

**\$35 million for medical applications, including imaging, diagnosis, medical clinics, and development.**

-----  
**\$245 million needed for commercial break-even**

**Antimatter**  
TECHNOLOGY CORPORATION

# Possible Antiproton Sources

- Collaboration with Brookhaven
- Collaboration with Fermilab
- Collaboration with a future facility
  - Advanced Hadron Facility at Los Alamos
  - Triumf in Canada
  - SSC Collaboration
- Dedicated production facility

# Transport and Storage Systems

- Medical Tabletop ring by Prof. Robert Wilson
- Design studies in RAND proceedings by researchers at UCLA and Los Alamos
  - Penning traps
  - Superconducting storage rings
- Molecular storage of antimatter

**Antimatter**  
TECHNOLOGY CORPORATION

# Imaging and Treatment

- Proton therapy at Harvard Cyclotron, and Mass General Hospital (4500 patients treated).
- Antiprotons are thought to be much more effective than protons, leading to reduced mortality.
- Imaging experiments at BNL (Kalogeropoulos *et al.*)
- Acceptance of particle treatment by the medical community (Loma Linda medical cyclotron installed).

# Potential Markets

- Cancer Treatment
- Medical Imaging
- Non-destructive Testing



# Cancer Treatment

- \$40 billion spent per year on cancer treatment
- 1 million new cases of cancer each year
- Over half of the patients receive radiation therapy.
- Antiprotons are the most selective particles in being able to deliver radiation to the tumor while leaving overlying tissues unharmed.
- A ten percent market penetration in the short-term could be expected to produce revenues of over \$1 billion per year.

**Antimatter**  
TECHNOLOGY CORPORATION

# Medical Imaging Market

- The medical imaging market is even larger than the cancer market (\$50 billion).
- CT scans produce too much radiation.
- Magnetic resonance imaging has limitations to certain types of tissue.
- Benefits from combined imaging and therapy.
- No really satisfactory techniques exist for mammography.

**Antimatter**  
TECHNOLOGY CORPORATION

# Non-Destructive Evaluation

- **Aerospace applications for critical components**
  - **Turbines**
  - **Composites**
  - **Structural members**
  - **Inspection of aging aircraft**
- **Aerospace spends 2% of sales on non-destructive test**
- **Aerospace sales amount to over \$50 billion per year**
- **Aerospace NDE is already a billion dollar market**
- **Electronic Industry**
  - **Inspection of solder joints**
  - **Automated annealing of cold solder joints**

**Antimatter**  
TECHNOLOGY CORPORATION

# References

- Lee, Y.Y. and D.I. Lowenstein, "Low Energy Antiproton Possibilities at BNL," Proceedings of the RAND Workshop on Antiproton Science and Technology, Oct. 6-9, 1987, World Scientific.
- Cline, D., "A Storage Ring for Antimatter Transport," RAND Proceedings.
- Goldman, T., "An Advanced Hadron Facility: Prospects and Applicability to Antiproton Production," RAND Proceedings.
- Howe, S.D., M. Hynes and A. Picklesimer, "Portable Pbars, Traps that Travel," RAND Proceedings.
- Kalogeropoulos, T., *et al.*, "Biomedical Potential of Antiprotons," RAND Proceedings.
- Greszczuk, L.B., "Potential Applications of Antiprotons for Inspection and Processing of Composites," RAND Proc.
- Wilson, R.R., "Radiological use of fast protons," Radiology 47:487, 1946.
- American Cancer Society, Annual Cancer Statistics for 1988.

**PROSPECTS FOR EXCITING EXTREME STATES  
IN NUCLEAR MATTER WITH INTENSE  
ANTIPROTON BEAMS**

**E. D. MINOR**

**DEPARTMENT OF PHYSICS  
THE PENNSYLVANIA STATE UNIVERSITY**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

EXCITING EXTREME STATES  
IN  
NUCLEAR MATTER  
USING  
INTENSE ANTIPROTON BEAMS


AUTHORS: E.D. Minor  
T.A. Armstrong  
R. Bishop  
V. Harris  
R.A. Lewis  
G.A. Smith


} Pennsylvania  
State  
University

SUPPORTED BY: Air Force Office of Scientific Res.  
Air Force Systems Command  
USAF grant  
U.S. National Science Foundation

WORKSHOP  
ON  
ANTIPROTON TECHNOLOGY

MAY 10, 1989  
BNL





95% of all  
stopped pN  
⇒ 2.8 π's

**Binary Annihilation**  
pN

$$E_{\pi} = \frac{2(938) - 5(140)}{5}$$

≈ 215 MeV KE.

$\sqrt{s} = 1266 \text{ MeV}$   
for πN

$10125 \sqrt{s} \approx 1300$   
for (2,2)



# NUCLEAR RESPONSE TO EXCITATION ENERGY $E^*$

- if  $E^*/A \lesssim 2-3$  MeV, the nucleus de-excites by thermal evaporation



(compound nucleus, N. BOHR)


- if  $E^*/A \gtrsim 2-3$  MeV, the nucleus fragments



(multi-fragmentation)

- if  $E^*/A \gg 8$  MeV, the nucleus disintegrates  
p, d, t, ...

annihilation at rest





● PS183  
● Hernandez and Oset

cf: J. Cugnon, P. Deney, J. Vandermaellou, Preprint, Liège.  
E. Hernandez, E. Oset. Valencia. Private communication


PS183 { CARBON  $\langle N_{\pi^{\pm}} \rangle = 2.59 \pm 0.04$   
URANIUM  $\langle N_{\pi^{\pm}} \rangle = 2.32 \pm 0.02$

Ref: EXP: P.L. McGaughey et al. Phys. Rev. Lett. 56(1986) 2156.  
 INC: J. Cugnon, P. Demaye, J. Vandrommenh. Preprint




McGaughey et al.

PS 183



f: INC: E. Hernandez, E. Orset. Valencia. Private communication.

{ PS183 (at rest)  
 { McGaughey et al. (608 MeV/c) }



Integrated Multiplicities:

$\langle N_{\pi^+} \rangle$  { stopped  $\bar{p}$  66  
 { 608 MeV/c  $\bar{p}$

Uranium

0.93


0.69

Carbon

1.12

0.96

ENERGY-WEIGHTED SPECTRA



$\pi^0 \rightarrow 2\gamma$   
 $\eta \rightarrow 5\gamma$   
 $\eta \rightarrow 3\pi^0 \rightarrow 6\gamma$   
 $\eta \rightarrow \pi^0\gamma\gamma \rightarrow 4\gamma$   
 $\eta \rightarrow \pi^+\pi^-\pi^0 \rightarrow 4\gamma$

$\omega \rightarrow \pi^+\pi^-\pi^0 \rightarrow 4\gamma$   
 $\omega \rightarrow \pi^0\gamma \rightarrow 3\gamma$

$\rho\bar{\rho}$  production.  
 $\langle M \rangle = 0.07 \pm 0.01$   
 $\langle \omega \rangle = 0.28 \pm 0.11$   
 $\langle \pi^\pm \rangle \approx 3.1$   
 $\langle \pi^0 \rangle \approx 1.8$

Refs: Adiels, Ljet et al. Phys. Lett. 152B (1986) 405.  
 Roy, J. Proc. of the Fourth Int. Symp. on N $\bar{N}$  Interactions. May 2-4, 1975. Syracuse, N.Y.  
 J. Cugnon, P. Denaye, J. Vandermouillon. Liège. Preprint

# ENERGY TRANSFER

$$E_{\text{TRANS}} = 1876.6 - \sum_{i=\{\pi^{\pm}, \alpha\}} \langle n_i \rangle \langle E_i \rangle$$

	$\langle M_E \rangle$	$\langle E_E \rangle$ MeV	$\langle M_{E\pm} \rangle$	$\langle E_{E\pm} \rangle$ MeV	$E_{\text{TOT}}$ MeV	$E_{\text{TRAN}}$ MeV
CARBON	$3.62 \pm 0.22$	$196 \pm 1$	$2.59 \pm 0.03$	$385 \pm 2$	$1706 \pm 45$	$170 \pm 45$
URANIUM	$2.73 \pm 0.18$	$185 \pm 1$	$2.32 \pm 0.02$	$377 \pm 2$	$1380 \pm 34$	$497 \pm 34$

## COMPARISONS:

CARBON: E. Hernandez, E. Oset (Prediction)  
( $n=4$ )

$E_{\text{TRAN}}$

228 MeV

P. Jassolotte et al. (Prediction) OXYGEN 260 MeV

URANIUM: E. Hernandez, E. Oset (Pred)

491 MeV

P. Jassolotte et al. (Pred)

380 MeV

$E^*/A$   
 $\leq 2.3 \text{ MeV}$   
 $\approx 2.3 \text{ MeV}$   
 $> 2.3 \text{ MeV}$


Future Behaviour  
Thermal evaporation  
multi-fragmentation  
nuclear disintegration: p, d, t, ...

Refs: Armstrong, T.A., et al. Zeit. f. Phys. A332 (1959).


E. Hernandez and E. Oset. Valencia. Private communication

Jassolotte, P., et al. Nucl. Phys. A484 (1988) 542.

# EXCITATION ENERGY $E^*$



$E^*$  scales by a factor of 5 between  $\bar{p}$  annihilation at rest and  $\bar{p}$  at 2.0 GeV



$\sim 20 \text{ MeV} = \text{Total ejectile K.E.}$

## $\bar{p}$ -Uranium annihilation

$$E^* = 1876.6 - E_\pi - W_{EJ} = 226 \text{ MeV} \quad \text{(at rest)} \\ \text{subcritical!}$$

But, at 2 GeV:  $E^* = 5 \times E^*(\text{rest}) = 1,133 \text{ MeV}$


$$E^*/A = 4.76 \text{ MeV}$$

! Multi-fragmentation!

# WHAT'S THE POINT?

## ENERGY DISTR. (A=27 FRAGS)

A.S. Botvina et al. / Statistical simulation




Botvina, A.S. et al.  
Nucl. Phys. A475 (1987) 663


Nucleus

Fragment

Range (TAR6 THICKNESS), T



$10^4 \text{ \AA}$



$0.5 \times 10^4 \text{ \AA}$

← The Stopping Power and Range of Ions and Solids. J.F. Ziegler, J.P. Biersack, J. Littler and R. Pagan. New York, 1985.

Cross Section,  $\sigma$

$0.36 \text{ b}$

$1.3 \text{ b}$

$\frac{\sigma \cdot \beta}{A^{2/3}} \approx 40 \text{ mb}$

Mean free path,  $\lambda$

$46 \text{ cm.}$

$1.2 \text{ cm.}$

$\lambda = \left( \sigma \frac{N_0}{A \rho} \right)^{-1}$

Yield

$2.2 \text{ per } 10^6 \text{ p's}$

$4.2 \text{ per } 10^6 \text{ p's}$

$Y = \frac{T}{\lambda} R_p$




**STATUS OF ASTRONAUTICS LABORATORY  
STUDIES RELATING TO CONDENSED  
ANTIMATTER**

**GERALD NORDLEY**

**APPLIED RESEARCH IN ENERGY STORAGE OFFICE  
ASTRONAUTICS LABORATORY (AFSC)  
EDWARDS AFB, CA**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

# ANTIMATTER PROGRAM AREAS




## ANTIPROTON PRODUCTION

OVERVIEW

## ANTIMATTER STORAGE

OVERVIEW


HYDROGEN CLUSTER IONS  
SOLID ANTIHYDROGEN STORAGE



## ANTIMATTER ANNIHILATION

OVERVIEW

FUTURE DEVELOPMENT




# ANTIPROTON PRODUCTION

## MOTIVATION

U S SOURCE FOR U S EXPERIMENTS  
NATIONAL SCIENCE AND TECHNOLOGY BASE  
CERN BUSY, UNEASY ON DEFENSE WORK  
ENABLING FOR DEFENSE RELATED USES


EVALUATE FEASIBILITY OF SCALE-UP  
ENABLING FOR FORECAST II PROPULSION  
GOAL OF \$1M/mg  
NO OTHER KNOWN "CUSTOMER"



## ANTIMATTER STORAGE

### MOTIVATION


- DEVELOP PORTABLE ANTI-PROTON STORAGE
  - ENABLE REMOTE EXPERIMENTS
  - ESTABLISH TECHNOLOGY BASE
  - FACILITATE NEAR TERM USES
- UNDERSTAND CLUSTER ION SCIENCE
  - PROPELLANT MOLECULE GROWTH
  - CONTACT-FREE NUCLEATION
  - CLUSTER TO SOLID TRANSITION ENERGY
- HIGH DENSITY ANTIHYDROGEN STORAGE
  - TECHNOLOGICAL SPINOFF
  - ENABLING FOR FORECAST II PROPULSION






# Contact-Free Storage....

7-4-17-5 D33



## PENNING TRAP ION TRAPS: PENNING TRAP




### PAYOFF:

- Effective Storage
- Portable
- Enables Near Term Uses
  - NDE of Nozzles, Fuels, Propulsion Materials
  - High Energy Physics at Universities
  - Medical Research

TR538a

# CLUSTER ION EXPERIMENT



# SCHEMATIC

## HYDROGEN CLUSTER ION PROJECT AFAL/AFOSR

### SCIENTIFIC OBJECTIVE

- o Determine Associative and Dissociative Pathways for Trapped Cluster Ions
  - Initially, develop technique to follow reaction pathways for  $H_5^+$  systems using a dense non-neutral plasma as a "pseudo-wall" to absorb the energy of association:  
$$M^+ + H_3^+ + H_2 \Rightarrow H_5^+ + M^{++}$$
$$M^+ + H_5^+ + H_2 \Rightarrow H_7^+ + M^{++}$$
  - Larger  $H_n^+$  systems as experience is gained
- o Assess Potential of Bulk Antihydrogen Nucleation via Cluster Ions

### APPROACH

1. Fill the trap with cluster ions of a selected size
2. Introduce molecular hydrogen, allowing it to interact with the trapped ions
  - use proven electromagnetic cooling techniques to maintain temperature
3. Measure size and radial distribution of products through NMR and/or mass spec




## **PAYOFF/POTENTIAL APPLICATIONS**

- o Bridge between theory and experiment on cluster ion formation & growth
  - Explain semi-empirical models for free expansion nozzles
  - Create new models for contact free processes
- o Illuminate third body cooling processes in non neutral plasmas (pseudo walls)
- o Provides versatile experimental apparatus in an active field
  - Cluster Ion Beam Experiments (Y.T. Lee et al., UCB)
  - Models for anti-cluster ion nucleation and growth (Saxon, SRI; Turner, AFIT)
  - Contribute to understanding cluster ion formation and deposition processes relevant to ion-matrix HEDM systems (Bae and Cosby, SRI)
  - Related to surface impact studies (Friedman, BNL and others)
- o Explore measurement techniques in ion trap environment

# ANTIHYDROGEN ICE STORAGE

1 K  
RADIATIVE  
HEAT SINK




U. HAWAII ( DR JIM GAINES )  
BROOKHAVEN NL (DR JIM POWELL)

- Preliminary results
- HOW STABLE IS SOLID ANTIHYDROGEN EXPOSED TO ANNIHILATION REACTION ?
  - 10 mg iceball will withstand 300 Annihilations/s
- WHAT TYPE OF VACUUM DOES THIS IMPLY ?
  - Number density Similar to Interstellar Space  
Based on: 100m/s atom velocity  
100% cross section  
Classical law
- WHAT ELSE DO WE NEED TO KNOW ?
  - Real cross sections
  - Very high Vacuum Material Behavior
  - Synergistic effects

# AIR FORCE ANTIPROTON TECHNOLOGY

## Path of Antiproton Technology Development



Antimatter Publications 1988 - Supported at Least in Part by AFAL

- General

- (1) B. Augenstein, B. Bonner, F. Mills, and M. Nieto, ed., Antiproton Science and Technology, World Scientific, New Jersey, 1988
- (2) Nordley, G., "Air Force Antimatter Technology Program" in Proceedings of the AFAL Workshop on Intense Positron Beams, Kells and Ottenwhife, ed., World Scientific, New Jersey, 1988
- (3)\* Forward, R.L. and Davis, J., "Mirror Matter", Wiley Scientific, New York, 1988

- Antiproton Production

- (4) Takahashi, Hiroshi, and Werner, Klaus, "Antiproton and Antineutron Production by Relativistic Heavy Ion Collision", BNL technical note (Journal article in preparation)

- Antiproton Storage

- (5)\* Talbi, D. and Saxon, R., "Theoretical Study of Excited Singlet States of  $H_3^+$ : Potential Surfaces and Transition Moments", submitted to J. Chem. Phys. in February 1988
- (6) Bahns, J.T., "Condensation and Storage of Hydrogen Cluster Ions", UDRI Technical Report, April 1988

- Antiproton Annihilation and Applications

- (7)\* Nordley, G., "Application of Antimatter Electric Power to Interstellar Propulsion", submitted to JBIS July 1988, preprint available from AFAL

\* Formally reviewed publications

**ELECTROMAGNETIC TRAPS**  
**FOR**  
**ATOMIC HYDROGEN OR ANTIHYDROGEN**

**ISAAC F SILVERA**

**DEPT. OF PHYSICS  
HARVARD UNIVERSITY  
CAMBRIDGE MA 02138**

PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989

Electromagnetic Traps  
for  
Atomic Anti hydrogen  
(or Hydrogen)

---

Isaac F Silvera  
Dept of Physics  
Harvard University  
Cambridge Ma 02138

Three types of traps:


1. Static Magnetic Trap

2. Laser Trap

3. Microwave Trap

I shall discuss hydrogen  
as the problems are the  
same for antihydrogen

# Hyperfine States of Hydrogen $|m_s m_i\rangle$



electron spin =  $1/2 \hbar : \vec{S}$

nuclear spin =  $1/2 \hbar : \vec{I}$

Total spin :  $F = I + S = 1 \text{ or } 0$



For static magnetic fields  
Maxwell's equations do  
not allow a field maximum  
in Free space.


A Field Minimum is allowed

No restriction on AC fields  
(microwave or laser)

# STATIC MAGNETIC TRAP (H. Kess)

uses field minimum

QUADRUPOLE  
MAGNET



Trap potential  $U \gg kT$



relaxation : magnetic dipole  
spin-exchange

$$\frac{dn}{dt} = -Gn^2$$

Rates are very rapid  
But  $n = 10^{14}/\text{cm}^3$   $\tau_{1/2} = 1\text{sec}$


$$\frac{1}{n} - \frac{1}{n_0} = -\frac{t}{\tau_{1/2}}$$

$\tau_{1/2} = 1/G$

MIT (Kleppner + Grestak et al)  
Amsterdam (Walraven et al)

Trap  $\sim 10^{11}$  atoms  
Densities  $10^{12} - 10^{13}/\text{cm}^3$

# Laser cooling of hydrogen




Laser light can cool and trap (Phillips et al; Chu et al Pritchard) ENS group)

Cool to quantum limit  $T_{min} = \frac{\Gamma}{2} = 2.2 \text{ mK}$  Lyman- $\alpha$

$\Gamma \equiv$  spontaneous emission rate



Dipole Trap Potential  $U$



$\omega_0 =$  transition frequency  
 $\omega =$  laser frequency

$\uparrow$  Deepest potential at  $\delta = -\Gamma$   
 But severe heating of gas due to spont. emission

For Na Chu et al used  $\delta \geq 10^3 \Gamma$  to minimize heating  
 but potential - few mK well



LASER SPONTANEOUS COOLING

Laser power is a major problem

For 1216 Å sources are pulsed, non-linear harmonic generators

Current sources  $\sim 10^{10} \frac{\text{photons}}{\text{sec}}$  (1 Watt =  $6 \times 10^{17} \frac{\text{photons}}{\text{sec}}$ )

Very Large improvements possible

but still pulsed at  $\sim 30 \text{ Hz}$

$$t_{\text{off}} = 33 \text{ msec}$$

Trapped atoms rapidly expand during off time

trap dimensions  $\sim$  microns

$\bar{v}$  at 2.2 mK  $\sim$  3 meters/sec


off time expansion

$$\Delta x = \bar{v} \times t_{\text{off}} = 3 \text{ m/sec} \times 33 \text{ msec} \\ \approx 10^8 \text{ cm} !$$

# New Trap: Microwave Trap

Agosta, Silvera, Vertchev + Stoof

Just like laser trapping, but no spontaneous emission



Dressed atoms: Diagonalize atomic states and radiation field ( $N$ -photons)

Dressed states

$$|1\rangle = \cos\theta |c, N-1\rangle + \sin\theta |b, N\rangle$$

low field seeker

$$|2\rangle = -\sin\theta |c, N-1\rangle + \cos\theta |b, N\rangle$$

high field seeker

State  $|2\rangle$  seeks highest microwave field


Confocal resonator cavity



$$U_{\text{trap}} = \frac{\hbar}{2} \omega_{\text{rabi}}$$


$$\omega_{\text{rabi}} \approx \mu_{bc} B_{\text{microwave}}$$

$$\therefore U_{\text{trap}} = \frac{\hbar}{2} \mu_{bc} B_{\text{microwave}}$$



$$|1\rangle = \cos\theta |c, N-1\rangle + \sin\theta |b, N\rangle$$

$$|2\rangle = -\sin\theta |c, N-1\rangle + \cos\theta |b, N\rangle$$



Plot of the admixture  
of  $|c\rangle$  or  $|b\rangle$  states  
into  $|1\rangle$  or  $|2\rangle$  states.

FIG. 1B



bare state relaxation



state  $|2\rangle$  : trapped




FIG. 2

## Advantages of $\mu$ -wave trap

No spontaneous heating  $\tau_{\text{spont}} \approx 10^7$  years

Microwave Technology well developed

## Difficulties

- Require microwave fields of several hundred gauss  
at  $\nu \approx 50$  GHz.

- Has not yet been built (Article  
To appear  
PRL next week)

**Loading** - A long story for another ~~session~~ time

# **ANTIHYDROGEN PRODUCTION**

**ARTHUR RICH**

**DEPARTMENT OF PHYSICS  
UNIVERSITY OF MICHIGAN  
ANN ARBOR, MI**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

Anti-Hydrogen: Formation and Applications

Presented by: Arthur Rich  
Physics Dept.  
University of Michigan





# Anti-Hydrogen: Formation and Applications

## I - Introduction

## II - Methods of $\bar{H}$ Production

### A - Overview

### B - Specific Methods



## III - $\bar{H}$ - Applications

## IV - Conclusions

COLLABORATORSUNIVERSITY OF MICHIGAN - EXPERIMENTAL

RALPH CONTI - POSITRONIUM ( $P_s$ )  $1^3S_1$  DECAY RATE ( $\lambda_T$ ) FINE STRUCTURE ( $n=2$ )  
TRANSITIONS AND CP TEST; ANTI-HYDROGEN ( $\bar{H}$ ) FORMATION

WILLIAM FRIEZE -  $e^+$  -  $P_s$  CONDENSED MATTER RESEARCH;  $e^+$  IMAGING;  $\bar{H}$

DAVID GIDLEY -  $\lambda_T$  (Gas and Vacuum);  $e^+$  -  $P_s$  CONDENSED MATTER  
AND POLARIZED  $e^+$  SURFACE MAGNETISM RESEARCH;  $e^+$  IMAGING;  $\bar{H}$

HENRY GRIFFIN (CHEMISTRY) - INTENSE  $e^+$  SOURCE DEVELOPMENT ( $\bar{H}$ )

JEFFREY NICO -  $\lambda_T$  (Vacuum)

MARK SKALSEY - WEAK INTERACTION TESTS via PRECISION BETA DECAY  
POLARIZATION MEASUREMENT, INTENSE  $e^+$  SOURCE DEVELOPMENT

TOM STEIGER - INTENSE  $e^+$  SOURCE DEVELOPMENT ( $\bar{H}$ )

JAMES VAN HOUSE - SEARCH FOR  $e_-$  HELICITY IN OPTICALLY ACTIVE  
MOLECULES USING POLARIZED  $e^+$  BEAMS;  $e^+$  IMAGING;  $\bar{H}$

PAUL ZITZEWITZ -  $\lambda_T$  (Vacuum);  $\bar{H}$ ; OPTIMIZATION OF POLARIZED  $e^+$  BEAMS

UNIVERSITY OF MICHIGAN - THEORETICAL

WILLIAM FORD

ROBERT LEWIS

YUKIO TOMOZAWA

GORDON KANE

LEONARD SANDER

EDWARD YAO

UNIV. OF TORONTO

DEREK PAUL

PRINCETON UNIV.

FRANK CALAPRICE

## WEAK INTERACTIONS

CERN-HEIDELBERG-DARMSTADT

HELMUT POTTH, et al.

ANTI-HYDROGEN

WAKE FOREST UNIV.

ROGER HEGSTROM

ORIGIN OF BIOLOGICAL ACTIVITY

GM RESEARCH LAB

WESTRUM CAPEHART

SURFACE MAGNETISM



## H APPLICATIONS

1. TCP Tests
  - Hyperfine Structure
  - Lamb Shift
  - Fine Structure
  - Electronic Structure (Rydberg) and  $m_{\bar{p}}$  (inertial),  $\mu_{\bar{p}}$
2. Production of Polarized  $\bar{p}$ 
  - i. Transfer of Polarization from an Initially Polarized  $e^+$  Beam -  

$$e^+(\uparrow) + \bar{p} \rightarrow \bar{H}(\uparrow) \rightarrow e^+(\uparrow) + \bar{p}(\uparrow)$$
  - ii. Optical pumping, Resonant Ionization, Lamb Shift Spin Filter -  

$$e^+ + \bar{p} \rightarrow \bar{H}; \quad \bar{H} + nh\nu \rightarrow \bar{H}(\uparrow) \rightarrow \bar{p}(\uparrow) + e^+(\uparrow)$$
3. Astrophysics
  - $$\bar{H} + H \rightarrow [\bar{p}p + Ps; \quad \bar{p} + p + Ps; \quad \bar{p}p + e^+ + e^-; \quad \bar{p} + p + e^+ + e^-]$$
  - $$\bar{H} + H \rightarrow \bar{H} + H$$
4.  $\bar{H}$  - Gravity Interaction
  - $$m_{\bar{p}} (\text{inertial})/m_p (\text{inertial}) - [\omega(\bar{H} - hfs)/\omega(H - hfs)]$$
  - $$m_{\bar{p}} \text{ gravitational -}$$
5. Atomic Physics
  - $$\bar{H} + \text{Matter} \rightarrow \bar{p} + e^+ + \text{Matter (stripping)}$$
  - $$\bar{H} + \text{Matter} \rightarrow \bar{H} + \text{Matter (elastic and inelastic scattering)}$$
6.  $\bar{H}$  - Energy storage

# ANTI-PROTON PRODUCTION



*Gabriele, Kells et al.*

*Deutsch et al. (e) ...*


*Gabriele (c)*



Pilot experiment for antihydrogen production  
at LEAR

Ann Arbor - BNL - CERN - Heidelberg - Karlsruhe Collaboration

Proposal 188



H Production by Radiative Recombination - Projected Rates (1 pass)


Assume equal area, non-relativistic,  $\bar{p}$  and  $e^+$  beams. Then

$$R(e^+ + \bar{p} \rightarrow \bar{H} + h\nu) \approx n(e^+) \langle \sigma v_T(e^+ - \bar{p}) \rangle (N(\bar{p})\eta)$$

where:  $n(e^+) = e^+ / \text{cm}^3$  in  $(e^+, \bar{p})$  overlap region

*10 Ci <sup>22</sup>Na 1 Remoderation*

*e<sup>+</sup> Source*



$$n = \frac{R(e^+/s)}{A(L = v \times 1 \text{ sec})} \sim \frac{(4 \times 10^{11}) \times (10^{-4})}{10^{-2} \text{ cm}^2 \times 10^{10} \text{ cm}} \sim 1 \text{ cm}^{-3}$$

$\frac{\alpha}{c^2} \omega^3 a_0^2 = \frac{\alpha}{c^2} \left(\frac{\omega c}{a_0}\right)^2 a_0^2 = \alpha^2 \frac{c}{a_0}$   
 $\leftarrow \gamma \quad \Delta T \rightarrow a_0 / v_T$

$$\sigma_i(\infty \rightarrow 1) = 2\pi \left[ (\pi a_0^2) \left( \frac{\alpha^4}{\beta_T} \right) \left( \frac{\alpha}{\beta_T} \right) \right] = \frac{2\pi^2 \alpha \pi_e^2}{\beta_T^2} \text{ cm}^2$$

$$\sigma = \sum \sigma_i \sim 3\sigma_1 \sim 0.5 \pi_e^2 / \beta_T^2$$

$$\langle \sigma v_T \rangle \sim \frac{\alpha \pi_e^2 c}{\langle \beta_T \rangle} \sim \frac{(4 \times 10^{-30}) \times (3 \times 10^{10})}{6 \times 10^{-4}} \sim 2 \times 10^{-12} \text{ cm}^2/\text{s}$$

$\alpha_T$   
 $T_T \sim 0.1 \text{ eV}$

$N(\bar{p}) =$  number of  $\bar{p}$  stored in LEAR  $\approx 10^{11}$  (Cooled -  $10^{10}$ )

$\eta =$  fraction of  $\bar{p}$  ring overlapped by  $e^+$  beam  $\approx 4\text{m}/80\text{m} \approx 0.05$

$$R \sim 1 \times (3 \times 10^{-12}) \times (5 \times 10^9) \sim 10^{-2}/\text{s} \sim 30 \text{ hr}^{-1} \text{ ( } 3 \text{ h}^{-1} \text{ )}$$

## H Production by Radiative Recombination

### Using Recirculating e<sup>+</sup>

$$R_S(e^+ + \bar{p} \rightarrow \bar{H}) \approx n_S(e^+) \alpha_I (N(\bar{p}) \eta) \quad \begin{matrix} 3 \times 10^{-12} & 10^{11} & 0.05 \\ \uparrow & \uparrow & \text{(fraction of } \bar{p} \text{ ring occup. by } e^+) \end{matrix}$$


$$\alpha_I \equiv \langle \sigma v_I \rangle \approx \frac{\pi r_e^2}{\beta^2} v_I \sim 10^{-19} \times 10^7$$

10 Ci Remoderate and store-16

$$n_S \equiv \frac{R(e^+/s)}{A(L = 10^3 \text{ cm})} = \frac{(4 \times 10^{11}) \times 10^{-5}}{10^{-2} \times 10^3} \sim 4 \times 10^6 \text{ cm}^{-3}$$

0.1 eV

e<sup>+</sup> - accumulated, cooled, pulsed and recirculated (10 m e<sup>+</sup> storage ring)



$$R_S(e^+ + \bar{p} \rightarrow \bar{H}) = (4 \times 10^6) \times (3 \times 10^{-12}) \times (5 \times 10^9) \sim 6000 / \text{s}$$

**H Production by Radiative Recombination :  
Recirculating e<sup>+</sup> and Stimulated Recombination**


$$R_S(e^+ + \beta \rightarrow H) \approx \underbrace{n_3(e^+)}_{3 \times 10^{-12}} \alpha \underbrace{N(\bar{p})}_{5 \times 10^7} G$$

*Laser recomb. gain factor*

*Recomb. and storage for 1 sec*

*e<sup>+</sup> storage*  $n_3 \approx \frac{R(e^+/s)}{A(L=10^3 \text{ cm})} = \frac{(4 \times 10^{11}) \times 10^{-5}}{10^{-2} \times 10^3} \sim 4 \times 10^5 \text{ cm}^{-3}$

e<sup>+</sup> - accumulated, cooled, pulsed and recirculated (10 m. storage ring)



G = Stimulated recombination gain factor ( $e^+ + \beta + h\nu \rightarrow H + 2h\nu$ ) =

$$= \frac{\pi^2 I (\text{MW/cm}^2)}{2 (T_e/0.1 \text{ eV})}$$

If  $\lambda_{\text{Lab}} = 521 \text{ nm}$  ( $\lambda_{\text{CM}} = \lambda(1 - \beta)\gamma = 364 \text{ nm}$  for  $\infty \rightarrow 2$ )

$\frac{14111}{m=2}$   
 $\frac{14111}{m=1}$

then  $G \sim 16 I$  for  $T_e \sim 0.1 \text{ eV}$




Commercial 20 MW/cm<sup>2</sup>  $\text{CO}_2$  laser (250 Hz, 20 ns pulse)  $G \sim 10^3$

Finally - If pulsed e<sup>+</sup> matches laser (stored pulse)

$$R_S(e^+ + \beta \rightarrow H) = \underbrace{(4 \times 10^5)}_{n_3} \times \underbrace{(3 \times 10^{-12})}_{\alpha} \times \underbrace{(5 \times 10^7)}_{N(\bar{p})} (G) \sim 6000 \text{ G/s}$$

CW - (CO/CO<sub>2</sub> ~ 20 W),  $n > 10$ ,  $G \sim 10^3$  but  
re-ionization and field ionization problems

# Positron Accumulator and Pulsers



$V_{g2}$   $\epsilon_M(W) \times \dots (\epsilon_{RM})$   $\epsilon_{RM}$   $20\text{ms}$   $20\text{mA}$   $5\text{kV}$


# of  $e^+$  / pulse  $\sim 4 \times 10^{11} \times (3 \times 10^{-4})^2 \times 20 \times 10^{-3} \sim 2 \times 10^5$  / pulse

" / sec  $\sim 2 \times 10^5 \times 50 \sim 10^7$  / A

$\times \epsilon_{RM}(\text{Ne}) \sim 10^2 \epsilon_{RM}(W)$


## Positron recirculator

### a) circular device



$E \times B$   
 circumference: 15 m, vacuum:  $10^{-12}$  Torr  
 magnetic field 200-600 G,  $e^+$  energy: 20-100 keV

### b) linear device



H APPLICATIONS

1. TCP Tests

- Hyperfine Structure
- Lamb Shift
- Fine Structure

}  $N \sim 10 \text{ cycles/m at } \beta \sim 0.3$

Electronic Structure (Rydberg) and  $m_{\bar{p}}$  (inertial),  $\mu_{\bar{p}}$

2. Production of Polarized  $\bar{p}$

i. Transfer of Polarization from an Initially Polarized  $e^+$  Beam -

$$e^+(\uparrow) + \bar{p} \rightarrow \bar{H}(\uparrow) \rightarrow e^+(\uparrow) + \bar{p}(\uparrow)$$

$$P_z(e^+) \quad P_z(e^+) \sim \frac{1}{2} P_z(e^+), \quad P_z(\bar{p}) \sim \frac{1}{2} P_z(e^+)$$

ii. Optical pumping, Resonant Ionization, Lamb Shift Spin Filter -

$$e^+ + \bar{p} \rightarrow \bar{H}; \quad \bar{H} + nh\nu \rightarrow \bar{H}(\uparrow) \rightarrow \bar{p}(\uparrow) + e^+(\uparrow)$$

3. Astrophysics

$$\bar{H} + H \rightarrow [\bar{p}p + Ps; \quad \bar{p} + p + Ps; \quad \bar{p}p + e^+ + e^-; \quad \bar{p} + p + e^+ + e^-]$$

$$\bar{H} + H \rightarrow \bar{H} + H$$

4.  $\bar{H}$  - Gravity Interaction

$$m_{\bar{p}} \text{ (inertial)} / m_p \text{ (inertial)} - [\omega(\bar{H} - hfs) / \omega(H - hfs)]$$

$m_{\bar{p}}$  gravitational -

5. Atomic Physics

$$\bar{H} + \text{Matter} \rightarrow \bar{p} + e^+ + \text{Matter (stripping)}$$

$$\bar{H} + \text{Matter} \rightarrow \bar{H} + \text{Matter (elastic and inelastic scattering)}$$

6.  $\bar{H}$  - Energy storage





## Polarized $\bar{p}$ - Projected Rates

Rate

$$R_S(e^+ + \bar{p} \rightarrow \bar{H}) = \underbrace{(4 \times 10^5)}_{n_S} \times \underbrace{(3 \times 10^{-12})}_{\alpha_F} \times \underbrace{(5 \times 10^8)}_{N(\bar{p}) \eta} \overset{\text{Current Laser gain } (G \sim 10^2 - 10^3)}{\sim 600 \text{ G/s}}$$

$\xrightarrow{(5 \times 10^9)} \sim 6000 \text{ G/s}$   
*Future (?)*


Polarization

- 1) If no attempt is made to maximize  $P(e^+)$ ,  $P(e^+) \sim 0.15$
- 2) To increase  $P(e^+)$  use Be absorber to reduce low energy  $e^+$  from beta spectrum (recall  $P_{\text{Loag}} = \langle v \rangle / c$ )
- 3) Maximize  $P^2 I \Rightarrow P(e^+) = 0.5$  but  $n_S = 1.5 \times 10^5$  (200 G/4)  
(2000 G/4)

Result

$$R_S(\bar{H}) \sim R_S(\bar{p}) \sim (2 \times 10^7 - 2 \times 10^8) \text{ G/day at } P(\bar{p}) \simeq \frac{1}{2} P(e^+) \simeq 0.25$$

### Polarized $\bar{p}$ - Projected Rates and Uses



$$A_{||} = \frac{1}{P_{\bar{p}}P_p} \left[ \frac{\sigma_T(\uparrow\bar{p}\uparrow p) - \sigma_T(\downarrow\bar{p}\uparrow p) - \sigma_T(\uparrow\bar{p}\downarrow p) + \sigma_T(\downarrow\bar{p}\downarrow p)}{\sigma_T(\uparrow\bar{p}\uparrow p) + \sigma_T(\downarrow\bar{p}\uparrow p) + \sigma_T(\uparrow\bar{p}\downarrow p) + \sigma_T(\downarrow\bar{p}\downarrow p)} \right]$$

where  $P_{\bar{p}}$  and  $P_p$  are the longitudinal polarizations of the antiproton beam and proton target, respectively, and the  $\sigma_T$  are the measured total cross sections with the sense of polarizations indicated by the arrows. A similar asymmetry  $A_{\perp}$  could be measured for the  $p$  and  $\bar{p}$  polarized transversely to the incident  $\bar{p}$  direction. Each asymmetry could be measured to a precision

$$\delta(A) = \frac{1}{\sqrt{N_{event}} P_{\bar{p}}P_p}$$

where  $N_{event}$  is the total number of  $\bar{p}$  interacting in the target. If the target thickness is chosen so that 20% of the incident  $\bar{p}$  interact in the target (a sufficiently small fraction to avoid degradation of the asymmetry by multiple scattering events) and given  $P_{\bar{p}} = 0.25$  and  $P_p \sim 0.12$ , ( $P_p$  is a typical value for the effective proton polarization in a hydrocarbon target with 70% hydrogen proton polarization) then in one day of running one can attain

$$\delta(A) = \pm \frac{1}{\sqrt{0.2 \times 3 \times 10^7} \cdot 0.25 \times 0.12} = \pm 0.01^*$$

\*  $A_{||}(pp) = 0.15$  ( $P_{Lab} = 1.6 \text{ GeV}/c$ )

## CONCLUSIONS

### I - HISTORY ( $e^+$ )

1932 - 72            **HEROIC PERIOD**

1972 - 85             $R(\text{slow } e^+) \sim (10^4 - 10^6)/\text{sec}$

$n(e^+/\text{cm}^3) \sim 1$

1985  $\rightarrow$              $R \sim 10^8/\text{s}$

$n \sim 10^6/\text{cm}^3$

### II - FUTURE

**INCREASE IN R, n  $\Rightarrow$**

1) **IMPROVED QUANTUM ELECTRODYNAMICS AND SYMMETRY TESTS.**

2)  **$e^+$  PLASMA.**

3)  **$e^+$  IMAGING.**

4) **ANTI-HYDROGEN.**

5) **BOSE-EINSTEIN CONDENSATION OF POSITRONIUM.**

$$(\lambda_{deB} \sim m^{-1/3})$$

**HEADQUARTERS DOE ANTIPROTON ACTIVITIES**

**DAVE GOODWIN**

**OFFICE OF HIGH ENERGY AND NUCLEAR PHYSICS  
U.S. DOE ER-20.1/GTN  
WASHINGTON, DC 20545**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**WORKSHOP ON ANTI-PROTON TECHNOLOGY**

**BROOKHAVEN, MAY 10, 1989**

**"HEADQUARTERS DOE ANTI-PROTON ACTIVITIES"**

**DAVE GOODWIN**

**SPECIAL ASST. TO THE ASSOCIATE DIRECTOR OF THE  
OFFICE OF HIGH ENERGY AND NUCLEAR PHYSICS (OHENP)**

**U.S. DOE, ER-20.1/GTN, WASH., D.C. 20545  
(301) 353-4037 (FAX 5079), FTS 233-4037 (FAX 5079)**

**EFFORTS TO OBTAIN RESOURCES FROM PROGRAM OFFICES**

**AVERAGE SINCE 10/87 WORKSHOP:**

- o ONE DOCUMENT PER WEEK**
- o DISCUSS WITH 2 PEOPLE PER DAY**

**SUPERCONDUCTING SUPER COLLIDER (SSC) STUDY**

**MARCH 16, 1989 LETTER FROM DIEBOLD TO SCHWITTERS:  
1995 PHYSICS FROM MEDIUM ENERGY BOOSTER (100 TO 200 GEV)  
10/2-4/89 OPEN HOUSE IN TEXAS**

**JIM BENSINGER (BRANDEIS), ROOM 2089C, SSC/URA, c/o LBL,  
90/4040, BERKELEY, CA 94720, (415) 486-4772 EXT. 6083,  
FTS 451-4772 EXT. 6083, FAX (415) 486-6796**

**PROVIDED: U. OF MICH. LETTER, 10/88 AND 2/89 RAND REPORTS  
AND 14 REPLIES TO SURVEY LETTER**

**DOCUMENTS: 5 (INCL. FY91 FUNDING ISSUE FOR SSC/BROOKHAVEN)**

**STATUS: OPEN**

SUPERCOMPUTERS

- FY88: WORKSHOP ANNOUNCEMENT, NO PROPOSALS**
- FY89: RFTPS TO 54 WORKSHOP ATTENDEES  
ONE PROPOSAL: TAKAHASHI (280 "HOURS"; \$100+K)  
UP TO 60+HOURS/MONTH AVAILABLE**
- FY90: 5/15/89 SUPERCOMPUTER ALLOCATIONS COMMITTEE MEETING:  
90,000 HOURS FOR HIGH ENERGY PHYSICS (HEP)/SSC,  
NUCLEAR PHYSICS (NP), BASIC ENERGY SCIENCES (BES) &  
HEALTH ENVIRONMENTAL RESEARCH (HER); EXCL. 27,000  
HOURS FOR FUSION  
RFTPS TO MOST ATTENDEES OF BOTH WORKSHOPS AND  
MIRROR MATTER NOTICE  
TAKAHASHI: 380 HOURS (< \$150K)**

**DOCUMENTS: 2**

**STATUS: OPEN**



10/11/88 SURVEY LETTER

319 HEP/SSC  
210 NP  
271 DOD  
54 WORKSHOP ATTENDEES  
1 U. OF N.C.  
387 MIRROR MATTER  
1,242

119

14 REPLIES (MIRROR MATTER SUMMARIZES): NONE NEGATIVE  
BES & HER: 14 REPLIES, 10/88 RAND REPORT & 5/89 WORKSHOP  
NOTICE WITH 2/89 RAND REPORT  
15TH REPLY: LIPTHANE

DOCUMENTS: 9

STATUS: "CLOSED"

TECHNOLOGY TRANSFER

3/28-29/89: DOE TECHNOLOGY TRANSFER WORKING GROUP:

- LOS ALAMOS, BERKELEY, OAK RIDGE, PACIFIC NORTHWEST, DEFENSE PROGRAMS (DP) AND FUSION

5/9/89: ER TECHNOLOGY TRANSFER STEERING GROUP

- BROOKHAVEN, BERKELEY, OAK RIDGE, ARGONNE, PACIFIC NORTHWEST, BES, HER AND FUSION

4/4/89: \$5 MIL./YR. FOR "SMALL SCIENCE" IMAGING/ANALYSIS AND ENERGY DEPOSITION WITH BES, HER, DP, NIH AND MINORITY EDUCATION (\$20 MIL./YR.)

FY90 REQUEST FOR SMALL SCIENCE & SUGGESTION FOR MINORITY EDUCATION

DOCUMENTS: 3

STATUS: OPEN

**SMALL BUSINESS INNOVATION RESEARCH (SBIR)**

**RECALL MIRROR MATTER: \$50K, \$500K**

**FY88: ELECTRON COOLING (D. LARSON)**

**2 TRANSPORTERS (W. WING)**

**LASER COOLING (" " )**

**RELATIVISTIC SELF-COLLIDER: RESCOL (B. MAGLICH)**

**FY89: ANTIPROTON TOPIC (OHENP SIGN): "PREMATURE"**

**ELECTRON COOLING (D. LARSON)**

**FY90 SBIR**

**HEP/SSC/NP MAY REDUCE FROM 6 TOPICS TO 3 - 5**

**5/4/89: ANTI-PROTON TOPIC**

**PROVIDED: 15 SURVEY REPLIES, 10/88 & 2/89 RAND REPORTS,  
SSC STUDY, SUMMARY FOR HUNTER & BROOKHAVEN PAPER**

**WILL NEED REVIEWERS**

**STATUS: OPEN**

**DOCUMENTS: 2**

**WEEKLY/BIWEEKLY REPORTS**

**WEEKLY: FROM OHENP THRU OFFICE OF ENERGY RESEARCH (OER)  
TO SECRETARY & ALL DOE PROGRAM & OPERATIONS  
OFFICES**

**BIWEEKLY: FROM OHENP TO ALL ER OFFICES (INCL. HEP, SSC,  
NP, BES, HER AND FUSION)**

**MONTHLY: TO ALL DOE OPS OFFICES AND ER OFFICES**

**10/87 WORKSHOP: WEEKLY, BIWEEKLY AND MONTHLY**

**12/1/87 AFAL BRIEFING (OHENP, HEP, BES & HER ATTEND):  
2 WEEKLY, BIWEEKLY, MONTHLY, NOTICES AND MINUTES (TO  
OER, ALL HEP/SSC & NP STAFF & BES, HER, FUSION & DP)**

WEEKLY/BIWEEKLY REPORTS

7/26/88 AFAL/AFOSR MEETING: WEEKLY AND BIWEEKLY

3/24/89 SDIO & 3/27/89 NASA MEETINGS: WEEKLY & BIWEEKLY

5/89 WORKSHOP: 1ST OF 2 WEEKLY, 1ST OF 2 BIWEEKLY & NOTICE  
TO OHENP, ALL HEP/SSC & NP STAFF & BES, HER  
FUSION & DP, WITH 2/89 RAND REPORT

RAND ASSOCIATED PRESS ITEM: WEEKLY AND BIWEEKLY

G-2: WEEKLY, BIWEEKLY, MONTHLY & MEMOS (INCL. 14 SURVEY  
REPLIES)

LEAR: WEEKLY & BIWEEKLY (4/17/89 LETTER FROM 3RD VERY  
RELIABLE SOURCE): STATUS OF LEAR AFTER 1992 ?

ANNUAL REPORT TO CONGRESS

DOCUMENTS: 28

STATUS: CONTINUING ACTION

**FY88 JASON STUDY**

**OPER MEETING WITH JASON**

**DOCUMENTS: 3**

**STATUS: CLOSED**

**KAON**

125

**(K̄AONS, ANTIPROTONS, ŌTHERS STRONGLY INTERACTING PARTICLES  
& N̄EUTRINOS)**

**NUCLEAR SCIENCE ADVISORY COMMITTEE (NSAC) REPORT**

**PROVIDED NP WITH 14 SURVEY REPLIES**

**AFTER CEBAF & RHIC**

**DOCUMENTS: 2**

**STATUS: "OPEN"**

PROPOSED USAF/HUNTER MEETING

SUMMARY TO HUNTER (SIGNED BY NP) PROVIDED TO OHENP, ALL  
HEP/SSC & NP STAFF & BES, HER, FUSION & DP

DOE POINT-OF-CONTACT

DOCUMENTS: 2 STATUS: "CLOSED"

TEXAS ACCELERATOR CENTER (TAC)

\$3 MIL. FROM CONGRESS

TAC PARTICIPATION SUGGESTED IN TRIP REPORT ON 10/87  
WORKSHOP (PROVIDED TO OER, OHENP, ALL HEP/SSC & NP STAFF &  
BES, HER, FUSION & DP)

AS OF 8/1/89: NO FURTHER HEP/SSC FUNDING

DOCUMENTS: 1 STATUS: "OPEN"



ACCELERATOR PRODUCED TRITIUM (APT)

PROPOSED ANSWER TO SENATE QUESTION INCLUDED ANTI-PROTONS  
JASON & 2/89 RAND REPORTS TO DP (INCL. CONGRESSIONAL  
LIASON)

ENERGY RESEARCH ADVISORY BOARD (ERAB) SUB-PANEL & GAO

EXYDER (XY SELF-COLLIDER):

0 \$17.628 MIL. FOR 2 1/2 YR., 10 KG/YR. OF ANTI-PROTONS  
INFO. TO DP INCL. ANTI-PROTONS

0 2/4/88 MIGMA MEETING WITH HEP, BES & FUSION (1 G/YR)

DOCUMENTS: 9

STATUS: "CLOSED"

SMALL/DISADVANTAGED BUSINESS (8A)

UP TO \$170K/YR.

DOCUMENTS: 7 (IN 1 OF 5 STATUS REPORTS TO AFAL/ANTI-M)

STATUS: "OPEN" (NEED PROPOSAL)

RAND REPORT ON REMOTE POWER

WITH 2/89 RAND REPORT TO: DP, NUCLEAR MATERIALS PRODUCTION  
& SP-100

DOCUMENTS: 3

STATUS: "OPEN"

ANNUAL MEETING OF NUCLEAR PHYSICS LAB DIRECTORS

RECOMMENDED AGENDA ITEM ON ANTI-PROTONS

DOCUMENTS: 1

STATUS: OPEN

INSTITUTIONAL PLANNING REVIEWS

RECOMMENDED AGENDA ITEM ON ANTIPROTONS FOR FY88 BROOKHAVEN  
REVIEW

FY89: 9/13/89 FERMILAB & 7/17-18/89 BROOKHAVEN REVIEWS

ALSO: 8/15-24/89 "PHYSICS AT FERMILAB IN THE 1990S"

DOCUMENTS: 6

STATUS: OPEN

THE PLANETARY SOCIETY

PROVIDE REFS FOR 9/85 AFAL REPORT & WORKSHOP PROCEEDINGS

DOCUMENTS: 1

STATUS: OPEN

OER & OHENP INFORMED

OER: 8 WEEKLY, 7 BIWEEKLY, 3 MONTHLY, JASON MEETING,  
SUMMARY FOR HUNTER, WORKSHOP: RAND & TRIP REPORTS, AFAL  
BRIEFING MINUTES AND 7 INFORMAL DISCUSSIONS

130

OHENP: SIGN 8 WEEKLY, 7 BIWEEKLY, SBIR & AFAL BRIEFING  
(ATTEND); SUMMARY FOR HUNTER, 3 MONTHLY, WORKSHOP: RAND &  
TRIP REPORTS, AFAL BRIEFING MINUTES, 5/89 WORKSHOP NOTICE  
WITH 2/89 RAND REPORT, FY90 FUNDS, SSC, BROOKHAVEN,  
14 SURVEY REPLIES, 7 PAPERS/ETC. & 22 INFORMAL DISCUSSIONS

HEP/SSC & NP INFORMED

HEP/SSC: 7 BIWEEKLY, 3 MONTHLY; ALL STAFF: SUMMARY FOR HUNTER, WORKSHOP TRIP REPORT, AFAL BRIEFING MINUTES ("ATTEND") & 5/89 WORKSHOP NOTICE WITH 2/89 RAND REPORT; WORKSHOP RAND REPORT, FERMILAB, BROOKHAVEN, SSC, 14 SURVEY REPLIES, 40 PAPERS/ETC., WEEKLY STAFF MEETINGS & INFORMAL DISCUSSION WITH 1 PERSON EVERY OTHER DAY

131

NP: 7 BIWEEKLY, 3 MONTHLY; ALL STAFF: SUMMARY FOR HUNTER (SIGN), WORKSHOP TRIP REPORT, AFAL BRIEFING MINUTES & 5/89 WORKSHOP NOTICE WITH 2/89 RAND REPORT; KAON/SSC, ANNUAL MEETING, WORKSHOP: RAND REPORT, 14 REPLIES, 1 PAPER, WEEKLY STAFF MEETINGS & INFORMAL DISCUSSION WITH 1 PERSON EVERY OTHER DAY

DP & FUSION INFORMED

DP: TECHNOLOGY TRANSFER, CLASSIFIED RAND REPORT, REMOTE POWER, APT/EXYDER, SUMMARY FOR HUNTER, AFAL BRIEFING, WORKSHOP: RAND & TRIP REPORTS, 4 PAPERS/ETC. (INCL. SDIO MEETING WEEKLY) AND 10 INFORMAL DISCUSSIONS

FUSION: 7 BIWEEKLY, 3 MONTHLY, SUMMARY FOR HUNTER, WORKSHOP TRIP REPORT, AFAL BRIEFING MINUTES, 5/89 WORKSHOP WITH 2/89 RAND REPORT AND 7 INFORMAL DISCUSSIONS

**FUNDING CONSORTIUM**

**DOE**

**DOD (E.G., USAF AND SDIO)**

**NSF**

**NIH**

**NASA**

**PRIVATE (E.G., ANTI-M AND DOD CONTRACTORS)**

**EUROPE (ITALY AND LEAR)**

**PROPOSAL**

**CAPTIAL: SSC OR \$15 MIL. FOR BROOKHAVEN  
\$ 10 MIL./YR. OPERATING (AT LEAST \$1 MIL./YR. FOR EACH):**

- (1) PRODUCTION**
- (2) PRODUCTION R&D**
- (3) TRANSPORTERS**
- (4) TRANSPORTER R&D**
- (5) "APPLIED" R&D (E.G., IMAGING, ANALYSIS &  
NON-PROPULSION ENERGY DEPOSITION)**



**PROPOSAL (CONTINUED)**

- (6) "ANTIGRAVITY"**
- (7) OTHER HENP R&D**
- (8) OTHER BASIC R&D ("NON-PROPULSION")**
- (9) PROPULSION R&D**


# **ANTIPROTON CATALYZED FUSION**

**T. E. KALOGEROPOLOUS**

**DEPARTMENT OF PHYSICS  
SYRACUSE UNIVERSITY**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

# $\bar{P}$ CATALYSIS



$\underbrace{\hspace{10em}}_{\text{MESONS} + 3 \text{ NUCLEONS}}$   
 Charge = +1

## EXPECTED RATE (ORDER(?) ESTIMATE)

$$R_{\bar{P}}^n \approx R_{\pi^-} \cdot \frac{\tau_{\bar{P}}^n}{\tau_{\pi^-}} \cdot \left( \frac{v_{\pi^-}}{v_{\bar{P}}^n} \right)^3$$

$\swarrow$   $\searrow$   $\searrow$   
 $10^2$   $(10^{-11} - 10^{-13}) / 10^{-6}$   $\left( \frac{m_{\bar{P}}}{m_{\pi}} \cdot \frac{1}{\eta^2} \right)^3 = 10^3 / \eta^6$


For  $n=2$

$R_{\bar{P}} \approx (1-10^{-2})\%$


# EVIDENCE

- COMPLETE STUDY OF 3223  $\bar{P}$  ANNIHILATIONS  
(PRL 33, 1631, 1974) RESULTED IN  
SEVEN "ZOO" EVENTS



- "ZOO" EVENTS ARE NOT:





OR




- TWO INTERESTING EVENTS



FRAME 285126 EVENT (TWO PROTONS)





<u>Track</u>	<u>θ</u>	<u>ε</u>	<u>D</u>	<u>P<sub>x</sub></u>	<u>P<sub>y</sub></u>	<u>P<sub>z</sub></u>	<u>E</u>	<u>COMMENTS</u>
π <sup>-</sup> (1)	13.6±.4	42.1±.3	368±26	196±14	178±13	-254±18	394±24	
π <sup>-</sup> (2)	33.2±.2	102.6±.2	193±8	-32±1	158±7	106±4	238±6	
π <sup>-</sup> (3)	42.0±.4	255.6±.4	415±30	-77±6	-299±22	-278±20	438±28	
π <sup>+</sup> (4)	42.7±.7	1.7±.7	253±18	186±13	6±2	-172±12	289±16	
π <sup>+</sup> (5)	53.5±.3	344.6±.4	255±15	146±9	-40±3	205±12	291±13	
P(6)	-21.0±7.3	331.2±3.9	169	139	-76	-61	958	STOPS (1.8cm)
P(7)	50.5±.3	179.9±.3	533±35	<u>-339±22</u>	<u>-1±2</u>	<u>411±27</u>	<u>1079±17</u>	OUT (26 cm)
				221±31	-74±26	-43±42	3682±45	



$$\left. \begin{array}{l} \\ \end{array} \right\} MM = m_n$$

FRAME 285082 EVENT (THREE PROTONS)



Track	E	c	R	$\underline{P_x}$	$\underline{P_y}$	$\underline{P_z}$	E	COMMENTS
$\pi^-(1)$	$31.4 \pm 1$	$109.4 \pm 0.06$	$404 \pm 16$	$-114 \pm 5$	$325 \pm 13$	$210 \pm 8$	$428 \pm 15$	
$\pi^-(2)$	$45.6 \pm 2$	$301.4 \pm 2$	$257 \pm 13$	$94 \pm 5$	$-154 \pm 8$	$-184 \pm 9$	$293 \pm 11$	
$\pi^-(3)$	$-33.0 \pm 0.09$	$221.0 \pm 0.06$	$194 \pm 7$	$-123 \pm 4$	$-107 \pm 4$	$-106 \pm 4$	$239 \pm 6$	
$\pi^+(4)$	$51.8 \pm 1.6$	$99.7 \pm 0.2$	$210 \pm 14$	$-22 \pm 2$	$128 \pm 9$	$165 \pm 11$	$252 \pm 12$	
$\pi^+(5)$	$10.7 \pm 1$	$177.4 \pm 0.05$	$270 \pm 7$	$-265 \pm 7$	$12 \pm 0$	$50 \pm 1$	$304 \pm 6$	
P(6)	$-30.6 \pm 2$	$99.6 \pm 0.1$	$441 \pm 15$	$-24 \pm 1$	$379 \pm 13$	$-224 \pm 8$	$1036 \pm 6$	OUT (33cm)
P(7)	$30.9 \pm 4$	$26.1 \pm 0.2$	129	$123 \pm 1$	$60 \pm 1$	$82 \pm 1$	952	STOPS (1.6cm)
P(8)	$-34.7 \pm 5$	$335.3 \pm 4$	250	$187 \pm 1$	$-86 \pm 1$	$-142 \pm 2$	$971 \pm 0$	STOPS (7.7cm)
				$-144 \pm 11$	$557 \pm 24$	$-149 \pm 18$	$4475 \pm 25$	

$$\bar{p} + {}^4\text{He}/2d \rightarrow 3\pi^- + 2\pi^+ + 3p + M\pi^2 (-0.3 \pm 0.04 \text{ GeV}^2/c^4)$$

NOT A  $\pi$ !

SPECIAL SCAN FOR "ZOO" EVENTS

- FILM FROM DISTANT RUNNING PERIOD THAN THAT FOR "COMPLETE STUDY"

- TOTAL  $\bar{P}$  EVENTS SCANNED 8800  
"ZOO" EVENTS FOUND 21

$$BR(ZOO) = (0.24 \pm 0.05)\%$$

- COMPARES WELL WITH THAT OF COMPLETE SCAN:


$$BR(ZOO) = (0.22 \pm 0.08)\%$$

---

ARE THESE EVENTS EXAMPLES OF  
 $\bar{P}$ - FUSION CATALYSIS?

---

• SECONDARY INTERACTIONS : NO!



1) Rate of '200' events  
 $\sim 100$  times the expected  
 rate for secondary  
 interactions

2)  $KE(P_1) + KE(P_2) > 140 \text{ MeV}$   
 which is not satisfied  
 in the two special events

• CHAMBER CONTAMINANTS SUCH AS N/O ...

- Typical results of chamber gas in % :

$O_2$  (0.05);  $D_2O$  (0.4);  $H_2O$  (0.04);  $HDO$  (0.06);

$N_2$  (0.2) .

OTHERS "UNMEASURABLE".

- BUT in  $D_2$ -liquid these contaminants  
 should freeze out!



• WHAT ABOUT  ${}^4\text{He}$ ?

- From The event which fits  $\bar{p} + {}^4\text{He}$  we estimate from ~~its~~ <sup>its</sup> a priori probability a contamination from  ${}^4\text{He}$  of  $\sim 10\%$  (!)
- This is not in line with BR(200).

CONCLUSIONS

- (1)  $\bar{p}$ -FUSION CATALYSIS IN LIQUID  $\text{D}_2$  OCCURS WITH A RATE  $\lesssim 1/100\bar{p}$
- (2) IF POSSIBLE  ${}^4\text{He}$  CONTAMINATION AT A LEVEL  $\gg 1\%$  COULD BE EXCLUDED THEN A GOOD CANDIDATE EVENT FOR  $\bar{p}/\text{D}_2$  FUSION INDUCED REACTION HAS BEEN OBSERVED AMONG 3000 ANNIHILATIONS.
- (3) SEARCH FOR  $\bar{p}$  FUSION CATALYSIS AND ITS PROPERTIES IS A GOOD COMPLIMENTARY RESEARCH TO  $\mu^-$ -CATALYSIS.

# **ANTIPROTON INDUCED FUSION REACTION**

**W. S. TOOTHACKER**

**LABORATORY FOR ELEMENTARY PARTICLE SCIENCE  
THE PENNSYLVANIA STATE UNIVERSITY  
UNIVERSITY PARK, PA**

**Note: We regret that copies of the transparencies used in Dr Toothacker's excellent presentation were not available for inclusion in the proceedings.**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**OPTIONS FOR A LABORATORY  
MICROFUSION FACILITY (LMF)**

**BRUNO AUGENSTEIN**

**THE RAND CORPORATION  
SANTA MONICA, CA**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

---

**SPECIAL DOE LABORATORY?**

**BROOKHAVEN MEETING,**

**MAY 10, 1989**

**WORKSHOP ON ANTIPROTON TECHNOLOGY**

---

# DOE LABORATORY MICROFUSION FACILITY (LMF) FOR TESTING

- Valid operational requirement
- Why have an LMF?
  - Simulate underground explosions in laboratory setting
- Against what contingencies?
  - Nuclear test ban: complete, or more restricted than current limitations
- No "assured design" yet for meeting LMF requirements

## WHAT CAN DO WE DO WITH AN LMF?

---

- Maintain nuclear design competence (complement, validate computations) — *ideally, with many tests per year*
  - Equation of state; opacity; energy flows; design principles
  - Exploratory research on new concepts
  - Effects simulation
- "Laboratory" facility → maximum energy releases in ~1/10 - 1 ton HE range
- Keep core proficiency program going
  - Prevent technological surprise, breakout

# TWO OPTIONS FOR LMF

## CONCEPTUAL STAGE (DOE)

- Very high energy laser, particle beam
  - Ignite small TN pellet
- TN energy release
  - 200-1000 MJoules
- Effects simulation
  - Radiation, EMP, etc.
- Projected cost goals
  - \$700 to 1000 million
- Construction projections
  - 5-6 years?
  - Available late 90s?

## ALTERNATIVE POSSIBILITY

- Use antiproton source
  - Based on J. Solem paper ('87 Proceedings)
  - Additional classified paper (Solem, Mayer, Augenstein)
  - Other related aspects (Pennsylvania State Group - G. Smith, et al)

*(Both options have two distinct regions:  
1. Ignite fuel, very local region;  
2. Conditions for sustained propagation of ignition.  
2. > > 1. in experimental preparations)*

# ANTIPROTON OPTION FOR LMF

---

- Initial experimentation
  - Standard initial tools
    - Antiproton source ( $\sim 10^{14}$  -  $10^{16}$  antiprotons/year) — use a small fraction
    - Portable storage
  - Significant Technical issues
    - Space/time compression of antiproton bunch; target configuration; precision diagnostics (must be made by normal means, not by sp. ts.)
    - Parameterized by Solem; Pennsylvania State group
- Broad-scale experimentation (comparable to laser, PB option plans)
  - Source roughly equivalent to Large Hadron Facility capability
  - Progress in storage, extraction, post-extraction compression



## LMF RECOMMENDATIONS

- DOE sponsor serious look at antiproton option
  - Comparable ground rules to laser, PB option
- Two cost bases
  - Piggy-back on Large Hadron Facility source, if firmed up; or,
  - Stand-alone source comparable to Large Hadron Facility source
- Evaluate range of simulation experimentation possible *(see for LMF allow's basic science spots. and spots. in other fields - e.g., propulsion, power generation)*
  - Availability for other kinds of experimental uses
- Criteria for antiproton option viability (initial screening)
  - Comparable range of uses
  - Highest cost option (stand-alone source)  $\leq$  laser, PB projections *(I mean for belief on highest cost much less than laser, PB costs)*

**MODELING ANTIPROTON - PLASMA  
INTERACTIONS**

**( ANTIMATTER THRUSTER MODELING )**

**JOHN L. CALLAS**

**JET PROPULSION LABORATORY  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
PASADENA, CA**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**Antimatter for Spacecraft Propulsion**


**ANTIMATTER THRUSTER MODELING**

**JPL**

**JOHN L. CALLAS**

**JET PROPULSION LABORATORY  
CALIFORNIA INSTITUTE OF TECHNOLOGY**

# JPL MISSION $\Delta V$ vs SPECIFIC IMPULSE




Alpha Centauri →  
(50 Years)

TAU (50 Years) →  
Mars (40 Days) →


CNSR, MRSR →

# JPL ANTIMATTER THRUSTER CONCEPTS


**SOLID CORE**  
 $I_{sp} = 800-1000 \text{ lbf-s/lbm}$   
 Efficiency < 70%




**GAS CORE**  
 $I_{sp} = 1000-2500 \text{ lbf-s/lbm}$   
 Efficiency < 50%



**PLASMA CORE**  
 $I_{sp} = 5000-100000 \text{ lbf-s/lbm}$   
 Efficiency << 50%




**BEAM CORE**  
 $I_{sp} = 10^7 \text{ lbf-s/lbm}$   
 Efficiency < 60%



# **JPL ANTIMATTER THRUSTER ISSUES**

- **dE/dx in Plasmas**
- **Photon Attenuation**
- **Bremsstrahlung Radiation (Electrons, Plasma)**
- **Synchrotron Radiation (Electron, Plasma)**
- **Nuclear Processes (Charge Exchange, Fission)**
- **Particle Decays (Neutrino Losses)**
- **Confinement (Annihilation Products, Plasma)**
- **Technology (Magnets, Shielding)**

Bmax = 27.6 Teslas  
Bmin = 9.3 Teslas



0.5M

# **JPL PRELIMINARY MODELING RESULTS**

- **Very Low Efficiency (<1%)**
- **Ultra High Loss Mechanisms (Neutrinos, Bremsstrahlung)**
- **Poor Annihilation Product Confinement (Charged and Neutral)**
- **Very Challenging Technology (Magnets, Shielding)**



**CONCEPTS FOR THE EXPERIMENTAL  
DETERMINATION OF RADIATION SHIELDING  
AND METAL CLAD PELLET PERFORMANCE**

**BRICE CASSENTI**

**UNITED TECHNOLOGIES RESEARCH CENTER  
HARTFORD, CT**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**Concepts for the Experimental Determination of  
Radiation Shielding and Metal Clad Pellet  
Performance**

**Brice Cassenti**



## RADIATION SOURCE

- Annihilation reaction



- Neutral pion decay



## Additional Radiation Sources

- Charge Exchange




- Fission




- Etc.

# GAMMA RAY INTERACTIONS

- Compton scattering




- Pair creation




# ELECTRON / POSITRON INTERACTIONS

• Gamma ray emission




$$e^- \xrightarrow{Z} \gamma + e^-$$

• Ionization



$$e^+ + e^- \xrightarrow{Z} e^+ + e^-$$

• Annihilation




$$e^+ + e^- \rightarrow \gamma + \gamma$$

# GAMMA RAY ENERGY DISTRIBUTION


Energy distribution

$$\frac{1}{N_0} \frac{dN}{dE}$$



# GAMMA RAY INTENSITY VARIATION

- Tungsten
- $\pi^0$  decay




Relative gamma  
ray intensity

Energy, MeV




# ELECTRON / POSITRON INTENSITY

- Tungsten
- $\pi^0$  decay



# ENERGY ABSORPTION

- Tungsten
- $\pi^0$  Decay




RA2197TX.008

Enabling procedure (Donoghue, 1986)

$$\bar{p}p \rightarrow \dots$$

3

# ENERGY ABSORPTION




RA2197TX.009

## **SAMPLE CALCULATION**


- **OTV mission**
  - 10 ton payload
  - 5.5 km / s velocity increment
  - 8 mg annihilated
- **Shield mass**
  - 3.5 tons
- **Maximum temperature rise**
  - 500 deg C

# JPL ANTIMATTER INTERACTION EXPERIMENT




from Callas: Antimatter Spacecraft Propulsion Experiments...

# JPL ANTIMATTER INTERACTION EXPERIMENT




from Callas: Antimatter Spacecraft Propulsion Experiments...

# PELLET ROCKET CONCEPTUAL DESIGN




# PELLET CONTAINEMENT






# JPL ANTIMATTER INTERACTION EXPERIMENT



# PROGRAM OUTLINES



## Summary

- Radiation shielding
- Pellet performance
- JPL Experiment
- Simulations

### References

Cassenti, B. N.: Radiation Shield Analysis for Antimatter Rockets. Paper No. AIAA-87-1813, presented at the AIAA/SAE/ASME/ASEE 23rd Joint Propulsion Conference, San Diego, June 29-July 2, 1987.

Hasegawa, A.: Magnetically Insulated Inertial Fusion: A New Approach to Controlled Thermonuclear Fusion. *Physical Review Letters*, Vol. 56, No. 2, pp. 139-142, 1986.

Max, C. E., W. M. Manheimer and J. J. Thomson: Enhanced Transport Across Laser Generated Magnetic Fields. *Phys. Fluids*, Vol. 21, pp. 128-139, 1978.

Kammash, T. and D. L. Galbraith: A Novel Fusion Scheme for Space Propulsion. Paper No. AIAA-87-2154, presented at the AIAA/ASME/ASEE 23rd Joint Propulsion Conference, San Diego, June 29-July 2, 1987.

Kammash, T. and D. L. Galbraith: Mars Missions with the MICF Fusion Propulsion System. Paper No. AIAA-88-2926, presented at the AIAA/ASME/ASEE 24th Joint Propulsion Conference, Boston, July 11-13, 1988.

**INTRODUCTION TO CP VIOLATION STUDIES**  
**WITH  $P$ bars**

**D. C. PEASLEE**

**DEPARTMENT OF PHYSICS AND ASTRONOMY**  
**UNIVERSITY OF MARYLAND**

PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989

# CP Violation

Kaon decay (Fitch + Cronin, 1964)

$$K_S \rightarrow 2\pi, \quad K_L \rightarrow 3\pi, \quad \Gamma_L \sim .002 \Gamma_S$$

Limitations:

1) Two parameters only,  $\epsilon$  and  $\epsilon'$ .  
No other CP in (u, d, s) at present.



2) Fundamental question unresolved:

i)  $|\Delta S|=2$  only (superweak)

ii)  $\left\{ \begin{array}{l} |\Delta S|=1 \text{ (milliweak)} \\ \text{and } |\Delta S|=2 \text{ by second-order} \end{array} \right.$

Hadron decay (T.D. Lee, 1966; Overseth + Pakvasa, 1969)

Non-leptonic,



Compare: decay rate,  $\Gamma - \bar{\Gamma}$   
 $\alpha(\vec{\sigma} \cdot \vec{p})$ ,  $\alpha + \bar{\alpha}$   
 $\beta(\vec{\sigma} \times \vec{\sigma}' \cdot \vec{p})$ ,  $\beta + \bar{\beta}$

In principle, 3 new measures of CP

$\Rightarrow$  All are  $|\Delta S| = 1$

Enabling procedure (Donoghue, 1986)

$$\begin{aligned} \bar{p} p &\rightarrow \Xi \Xi \\ &\rightarrow \bar{\pi} \Lambda \end{aligned}$$

Production: strong interaction, preserves CP

$\therefore$  Perfect anticorrelation of initial states,

$$\sigma_m = -\bar{\sigma}_m, \quad p_m = -\bar{p}_m$$

Eliminates major uncertainties.

Estimates for K-M, Higgs, Left-right:

$$\underline{\Xi \rightarrow \pi \Lambda}$$

$$|\beta + \bar{\beta}| \sim 10^{-4} \text{ to } 10^{-3} \sim 3 |\alpha + \bar{\alpha}|$$

$$\underline{\Lambda \rightarrow \pi p}$$

$$|\beta + \bar{\beta}| \sim 10^{-4} \text{ to } 10^{-3} \sim 7 |\alpha + \bar{\alpha}|$$



## Implementation

④

- 1) Don't attempt  $(\Gamma - \bar{\Gamma}) / (\Gamma + \bar{\Gamma}) \sim 10^{-6}$  to  $10^{-5}$
- 2)  $|\alpha + \bar{\alpha}|_{\Omega} < |\alpha + \bar{\alpha}|_{\Xi}$  (estimate)
- 3)  $|\beta + \bar{\beta}|_{\Xi}$  only feasible,  $\Lambda$  decay analyzes  $\sigma_m'$

Conflicting aims:

- i)  $\sigma_{\Omega} \sim 10 \sigma_{\Xi}$ , but CP parameters largest (estimated) for  $\Xi$ .
- ii) For  $\Xi$ ,  $|\beta + \bar{\beta}| > |\alpha + \bar{\alpha}|$ ; but  $|\alpha + \bar{\alpha}|$  simpler, perhaps higher acceptance.

**TEST OF NON-CONSERVATION IN  
IN  $P$  bar- $P$  to  $\Xi$  bar -  $\Xi$**

**A. M. NATHAN**

**DEPARTMENT OF PHYSICS  
UNIVERSITY OF ILLINOIS  
CHAMPAIGN, IL**

Note: We regret that reproducible copies of the transparencies used in Dr Nathan's excellent presentation were not available for inclusion in the proceedings.

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**STUDIES OF CP VIOLATION WITH  
PURE  $K_0 - K_0$  bar BEAMS FROM  $P$  bars**

**JAMES MILLER**

**DEPARTMENT OF PHYSICS  
BOSTON UNIVERSITY**

Note: We regret that reproducible copies of the transparencies used in Dr Miller's excellent presentation were not available for inclusion in the proceedings.

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**SEARCH FOR CP VIOLATION IN**

$$**Pbar P \rightarrow J/\Psi \rightarrow \Lambda^0 \Lambda^0 bar**$$


**G. A. SMITH**

**LABORATORY FOR ELEMENTARY PARTICLE SCIENCE  
THE PENNSYLVANIA STATE UNIVERSITY  
UNIVERSITY PARK, PA**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

SEARCH FOR CP-VIOLATION

IN






G.A. SMITH  
(PENN STATE)

WORKSHOP  
ON  
ANTIPROTON TECHNOLOGY

MAY 10, 1989  
BNL

# HIGH RESOLUTION HEAVY QUARK SPECTROSCOPY WITH ANTI-PROTONS

NOTE REMARKABLE SIMILARITY BETWEEN SPECTRA OF CHARMONIUM AND POSITRONIUM!



# R704 (ISR)





Table 2  
 $\bar{p}p \rightarrow J/\psi \rightarrow e^+e^-$  runs



Run	Date	$\sqrt{s}$ (MeV/c <sup>2</sup> )	$\int L dt$ (nb <sup>-1</sup> )	Events	Mass (MeV/c <sup>2</sup> )
1	11 May 1983	3096.7-3101.0	10.0	16	3097.05 $\pm$ 0.22 $\pm$ 0.21
2	19 July 1983	3095.8-3097.5	8.0	13	3097.35 $\pm$ 0.22 $\pm$ 0.30
3	26 July 1983	3096.4-3097.6	18.0	14	3095.69 $\pm$ 0.35 $\pm$ 0.32
4	3 August 1983	3096.4-3097.0	21.5	34	3096.66 $\pm$ 0.25 $\pm$ 0.19
5	22 March 1984	3096.1-3098.0	63.5	81	3096.79 $\pm$ 0.14 $\pm$ 0.11
6	5 April 1984	3096.7	20.0	35	3096.64 $\pm$ 0.34 $\pm$ 0.64

C. Baglin et al, Nucl. Phys. B286, 592 (1987)



# R704 (ISR)



NO COPLANARITY CUT



WITH COPLANARITY CUT  $\sqrt{s} = 2981 \text{ MeV}$

M[e<sup>+</sup>e<sup>-</sup>] (GeV)



TOTAL ENERGY (GeV)


$\sqrt{s} = 3097 \text{ MeV}$






# FERMILAB E-760

FERMILAB - FERRARA - GENOA - IRVINE  
NORTHWESTERN - PENN STATE - TORINO



NAT ELECTROMAGNETIC  
CALORIMETER


$^3S_1 \rightarrow e^+e^-$   
 $^1S_0 \rightarrow \gamma\gamma$   
 $^3P \rightarrow ^3S_1, \gamma$   
 $\quad \quad \quad \rightarrow e^+e^-$   
etc.




**FINE, HYPERFINE STRUCTURE OF P-STATES?**

$\Delta M = a \langle L \cdot S \rangle + b \langle T \rangle + c \langle S_1 \cdot S_2 \rangle$


$a \sim 35 \text{ MeV} ?$   
 $b \sim 40 \text{ MeV} ?$   
 $c = 0$  Confinement in scalar gluonic field?




57-603-2 4-6



11-11-57 11-11-57 11-11-57 11-11-57



**FERMILAB**  
**FERRARA**  
**GENOVA**  
**IRVINE**  
**NORTHWESTERN**  
**PENN STATE**  
**TORINO**



**FERMILAB**  
**ANTI-PROTON SOURCE**

$$\mathcal{L}_{E760} \approx 10^{31} \text{ cm}^{-2} \text{ s}^{-1} \text{ (Gas jet)}$$



$$\mathcal{L}_{LEAR} \approx 1.5 \times 10^{29} \text{ cm}^{-2} \text{ s}^{-1} \text{ (} 2 \times 2 \text{ GEV/c}^2 \text{)}$$

(POST ACOL)

$$\lesssim 4436 \text{ MEV}$$

# LOOKING AT CP INVARIANCE AND QUANTUM MECHANICS IN $J/\psi \rightarrow \Lambda \bar{\Lambda}$ DECAY

M.H. TIXIER ET. AL. PHYS. LETT. 212B, 523 (1988)



$$\frac{\partial \sigma}{\partial \cos \theta \partial \Omega' \partial \Omega''} \quad \begin{array}{l} 1) \text{ CP INVARIANT} \\ 2) \text{ NO ASSUMPTION ON} \\ \text{REL. } \Lambda, \bar{\Lambda} \text{ POLARIZATION} \end{array}$$

$$\propto 2 \left[ 1 - \left( \frac{P_\Lambda^2}{E_\Lambda^2} \right) \sin^2 \theta \right] (1 - \alpha_\Lambda a_n b_n)$$

$$+ \left( \frac{P_\Lambda^2}{E_\Lambda^2} \right) \sin^2 \theta \left[ 1 - \alpha_\Lambda^2 (\vec{a} \cdot \vec{b} - 2a_x b_x) \right]$$

- 2)  $P_\Lambda^2/E_\Lambda^2 = 0.48 (J/\psi), \alpha_\Lambda^2 = 0.412$
- b)  $\hat{n}$  orthogonal to  $\vec{N}_{y'} (J/\psi \text{ polarization})$  and  $\hat{x}$




Fig. 7. The  $\vec{a} \cdot \vec{b}$  distribution. MC simulation and data

TO TEST CP, SET  $\alpha_\Lambda^2 = -\alpha_\Lambda \alpha_{\bar{\Lambda}}$ . ASSUMING VALIDITY OF Q.M.,  $\alpha_{\bar{\Lambda}}$  IS EXTRACTED FROM  $\vec{a} \cdot \vec{b}$  DISTRIBUTION  $\Rightarrow$

$\alpha_{\bar{\Lambda}} = 0.01 \pm 0.10 (-0.07 \pm 0.09, P. BARNES ET. AL. PHYS. LETT. 117B, 1986)$

# FURTHER TESTS OF CP INVARIANCE IN $J/\psi \rightarrow \Lambda^0 \bar{\Lambda}^0$

- 1) GOAL:  $10^{-4}$  ERROR IN A) J. DONOGHUE ET AL,  
PR D34 833 (1986)
- 2) PRESENT LIMITS:  $\sim 10^{-1}$  2) L. WOLFENSTEIN,  
ANN. REV. NUCL. PART.  
SCI 36, 137 (1986)
- 3) FIRST STEP  $\Rightarrow 10^{-2}$

$$E-760: \text{RATE} = (5 \times 10^{31} \text{ cm}^{-2} \text{ s}^{-1}) (10^{-30} \text{ cm}^2) (0.13 \times 10^{-2})$$

$$\times (0.64) (0.64) = 0.027 \text{ SEC}^{-1}$$

$\text{BR}_{J/\psi \rightarrow \Lambda \bar{\Lambda}}$   
 $\Lambda \rightarrow p \bar{\pi} \quad \bar{\Lambda} \rightarrow \bar{p} \pi^+$

EXTRAPOLATE DM2 RESULTS ( $2 \times 10^3$  EVTS.)  
TO  $10^{-2}$  LEVEL ( $2 \times 10^5$  EVENTS)

$$\frac{2 \times 10^5 \text{ EVTS}}{0.027 \text{ SEC}^{-1}} \Rightarrow 7.4 \times 10^6 \text{ SEC (2.8 months)}$$

WILL PROBABLY GET DONE FOR FREE  
(SYSTEMATIC ERRORS ??)

- 4) NEXT STEP  $\Rightarrow 10^{-3}$

REQUIRES  $2 \times 10^7 J/\psi \rightarrow \Lambda^0 \bar{\Lambda}^0$ , OR  $\sim 10^{10} J/\psi$ 's !!

$J/\psi$  FACTORY @  $\mathcal{L} \approx 10^{33} \text{ cm}^{-2} \text{ SEC}^{-1}$

BEST BET IS PROBABLY  $e^- e^+$   
LINEAR COLLIDER?

**RARE MODES OF NUCLEON-ANTINUCLEON  
ANNIHILATION**

**CARL B. DOVER**

**DEPARTMENT OF PHYSICS  
BROOKHAVEN NATIONAL LABORATORY  
UPTON, NY**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

RARE MODES OF  
 $N\bar{N}$  ANNIHILATION

CARL B. DOVER  
BROOKHAVEN LAB



PRODUCTION OF

$J^{\pi C}$  EXOTICS IN  $\bar{N}N \rightarrow \text{mesons}$

PRODUCTION OF

$J^{\pi C}$  EXOTICS IN  $\bar{N}N \rightarrow \text{mesons}$

What is a  $J^{\pi C}$  exotic meson?

$J$  = total angular momentum

$\pi$  = parity ( $\pm$ )

$C$  = charge conjugation parity ( $\pm$ )

$J^{\pi C}$  exotic quantum numbers cannot correspond to a quark-antiquark ( $Q\bar{Q}$ ) system as for ordinary mesons ( $\pi, \eta, \rho, \omega, \dots$ )

Examples:  $J^{\pi C} = 0^{--}, 0^{+-}, 1^{-+}$  are exotic

$\leadsto$  must have more complicated structure than  $Q\bar{Q}$

$\leadsto Q^2\bar{Q}^2, Q\bar{Q}g, \dots$

$\uparrow$   
excitation of the gluonic field

## THEORETICAL PREDICTIONS FOR MASSES OF EXOTICS

consider  $Q\bar{Q}g$  "hybrids"  $(0^{-+} \text{ or } 1^{--})_{Q\bar{Q}} \otimes \left\{ \begin{array}{l} TE (1^{+-}) \\ TM (1^{--}) \end{array} \right.$

$\uparrow$   $\uparrow$   
 $^1S_0$   $^3S_1$

lowest lying are  $^3S_1 \otimes TE = \left\{ \begin{array}{l} 1^{-+} (1^-) \quad \rho_g \\ 1^{-+} (0^+) \quad \omega_g \end{array} \right.$

### Mass Estimates and Decays:

- 1) MIT Bag Model (Barnes, Close, deViron, NP B224, 241 (1983))

$$M(\rho_g) \approx 1.4 \text{ GeV}, \quad M(\omega_g) \approx 1.55 \text{ GeV}$$

(Chanowitz, Sharpe NP B222, 211 (1983) give larger M's but  $\approx$  same splitting)

- 2) Flux Tube Model (Isgur, Kokoski, Paton, PRL 54, 869 (1985))

$$M(\rho_g) \approx M(\omega_g) \approx 1.9 \text{ GeV} \quad [\text{also } 0^{+-}(0^-, 1^+), 2^{+-}(0^-, 1^+)]$$

Selection Rules:  $\rho_g \rightarrow \pi^\pm B^\mp, \pi^0 D$ ;  $\rho_g \not\rightarrow \pi\eta, \pi\rho, \rho\omega, K^*$

- 3) QCD Sum Rules (Latorre et al, Z. Phys. C34, 347 (1987))

$$M(\rho_g) \approx 1.6 - 2.1 \text{ GeV}$$

$\rho_g \rightarrow \pi\rho, K^* \bar{K}$  large,  $\rho_g \rightarrow \pi\eta, \pi\eta'$  suppressed

Candidate for a  $J^{\pi C}$  exotic resonance:

D. Alde et al, Phys. Lett. B205, 397 (1988)

reaction  $\pi^- p \rightarrow X^0 n$   
 $\searrow \pi^0 \eta (l=1)$

$X^0$  seen as interference effect (asymmetry)  
with  $\pi^- p \rightarrow A_2^0 n$   
 $\searrow \pi^0 \eta (l=2)$

$X^0$  is  $J^{\pi C}(IG) = 1^{-+}(1^-)$  exotic

$M = 1406 \pm 20 \text{ MeV}$ ,  $\Gamma = 180 \pm 30 \text{ MeV}$

Note:  $X$  should be accessible in  $\bar{N}N \rightarrow \pi X$

$\rightarrow$  look for optimum channels

## Experimental Searches for Exotics

### 1) LEAR at CERN

JETSET, CRYSTAL BARREL, OBELIX

### 2) Brookhaven AGS E818 Chung et al

$$\pi^- p \rightarrow \rho_0^- p \text{ at } 12 \text{ GeV}/c$$

Decay chain:  $\rho_0^- \rightarrow \pi^- f_2 \rightarrow \rho^+ \pi^-$   
 $\rho^+ \rightarrow K^+ \bar{K}^0 \rightarrow \pi^+ \pi^-$

overall reaction  $\pi^- p \rightarrow K^+ \pi^- 3\pi^- p$

### 3) KEK (Japan)

$$\pi^- p \rightarrow \rho_0^- p \text{ at } 6 \text{ GeV}/c$$

Decay chain:  $\rho_0^- \rightarrow \pi^- \eta \rightarrow 2\gamma$

Production of  $J^{\pi C}$  exotics in  $\bar{p}p \rightarrow \pi^0 X^0$

$J^{\pi C}(I^G)$ of $X$	$\bar{N}N(L=0) \rightarrow \pi^0 X^0(\ell_f)$	$\bar{N}N(L=1) \rightarrow \pi^0 X^0(\ell_f)$
$0^{--}(0^-)$	${}^3{}^3S_1(\ell_f=1)$	—
$0^{--}(1^+)$	${}^1{}^3S_1(\ell_f=1)$	—
$0^{--}(0^+, 1^-)$	—	—
$0^{+-}(0^-)$	—	${}^3{}^1P_1(\ell_f=1)$
$0^{+-}(1^+)$	—	${}^1{}^1P_1(\ell_f=1)$
$0^{+-}(0^+, 1^-)$	—	—
$1^{-+}(0^+) \omega_g$	${}^3{}^1S_0(\ell_f=1)$	${}^3{}^3P_1(\ell_f=0,2), {}^3{}^3P_2(\ell_f=2)$
$1^{-+}(1^-) \rho_g$	${}^1{}^1S_0(\ell_f=1)$	${}^1{}^3P_1(\ell_f=0,2), {}^1{}^3P_2(\ell_f=2)$
$1^{-+}(0^-, 1^+)$	—	—

C conservation is strong constraint!  
see  $X^0$  with both  $I=0, 1$

Production of  $J^{\pi C}$  exotics in  $\bar{p}p \rightarrow \pi^{\pm} X^{\mp}$

$J^{\pi C} (I^G)$ of $X$	$\bar{p}p (L=0) \rightarrow \pi^{\pm} X^{\mp} (\ell_f)$	$\bar{p}p (L=1) \rightarrow \pi^{\pm} X^{\mp} (\ell_f)$
$0^{--} (1^-)$	${}^3{}^3S_1 (\ell_f=1)$	${}^1{}^3P_0 (\ell_f=0), {}^1{}^3P_2 (\ell_f=2)$
$0^{--} (1^+)$	${}^1{}^3S_1 (\ell_f=1)$	${}^3{}^3P_0 (\ell_f=0), {}^3{}^3P_2 (\ell_f=2)$
$0^{+-} (1^-)$	${}^1{}^1S_0 (\ell_f=0)$	${}^1{}^3P_1, {}^3{}^1P_1 (\ell_f=1)$
$0^{+-} (1^+)$	${}^3{}^1S_0 (\ell_f=0)$	${}^3{}^3P_1, {}^1{}^1P_1 (\ell_f=1)$
$1^{-+} (1^-)$	${}^1{}^1S_0 (\ell_f=1), {}^3{}^3S_1 (\ell_f=1)$	${}^1{}^3P_1, {}^3{}^1P_1 (\ell_f=0,2), {}^1{}^3P_2 (\ell_f=)$
$1^{-+} (1^+)$	${}^3{}^1S_0 (\ell_f=1), {}^1{}^3S_1 (\ell_f=1)$	${}^3{}^3P_1, {}^1{}^1P_1 (\ell_f=0,2), {}^3{}^3P_2 (\ell_f=)$

no constraint from  $C$   
 $I=0$   $X$  forbidden



'roduction of  $J^{\pi C}$  exotics in  $\bar{p}n \rightarrow \pi^- X^0$

$J^{\pi C} (I^G)$ of $X$	$\bar{p}n (L=0) \rightarrow \pi^- X^0$	$\bar{p}n (L=1) \rightarrow \pi^- X^0$
$0^{--} (0^-, 1^-)$	${}^3S_1 (L_f=1)$	—
$0^{--} (0^+, 1^+)$	—	${}^3P_0 (L_f=0), {}^3P_2 (L_f=2)$
$0^{+-} (0^-, 1^-)$	—	${}^3P_1 (L_f=1)$
$0^{+-} (0^+, 1^+)$	${}^3S_0 (L_f=0)$	${}^3P_1 (L_f=1)$
$1^{-+} (0^-, 1^-)$	${}^3S_1 (L_f=1)$	${}^3P_1 (L_f=0, 2)$
$1^{-+} (0^+, 1^+)$	${}^3S_0 (L_f=1)$	${}^3P_1 (L_f=0, 2), {}^3P_2 (L_f=2)$

no constraint from  $C$   
see both  $I=0, 1$

## NON-STRANGE DECAY MODES OF $J=0, 1$ EXOTICS

consider  $PS+PS, PS+V, PS+T, VV, PS+S$

" $\eta$ " =  $\{\eta, \eta'\}$ , " $\omega$ " =  $\{\omega, \varphi\}$ , " $f$ " =  $\{f, f'\}$ , " $\sigma$ " =  $\{\sigma, \sigma^*\}$ , " $D$ " =  $\{D, E\}$

$J^{\pi C}(I^G)$	Allowed Decays $X^0 \rightarrow M_1 M_2 (L)$
$0^{--}(0^-)$	$\pi^0 \rho^0, \pi^\pm \rho^\mp, \eta \omega (L=1)$
$0^{--}(0^+)$	$\pi^\pm A_2^\mp (L=2), \pi^\pm \delta^\mp (L=0)$
$0^{--}(1^-)$	$\pi^\pm \rho^\mp (L=1)$
$0^{--}(1^+)$	$\pi^0 \omega, \eta \rho^0 (L=1), \pi^\pm A_2^\mp (L=2), \pi^\pm \delta^\mp (L=0)$
$0^{+-}(0^-)$	$\pi^0 B^0, \pi^\pm B^\mp (L=1)$
$0^{+-}(0^+)$	$\pi^+ \pi^- (L=0), \rho^+ \rho^- (L=0, 2), \pi^\pm A_1^\mp (L=1)$
$0^{+-}(1^-)$	$\pi^\pm B^\mp (L=1)$
$0^{+-}(1^+)$	$\pi^+ \pi^- (L=0), \rho^+ \rho^- (L=0, 2), \pi^\pm A_1^\mp (L=1), \pi^0 H (L=1)$
$1^+(0^-)$	$\pi^\pm \rho^\mp (L=1), \pi^\pm B^\mp (L=0, 2)$
$1^+(0^+)$ $\omega_8$	$\pi^0 \pi^0, \pi^\pm \pi^\mp (L=1), \eta \eta (L=1), \rho^0 \rho^0, \rho^+ \rho^- (L=1, 3)$ $\omega \omega (L=1, 3), \pi^0 A_1^0, \pi^\pm A_1^\mp (L=0, 2), \pi^0 A_2^0, \pi^\pm A_2^\mp (L=2)$
$1^+(1^-)$ $\rho_8$	$\pi^0 \eta (L=1), \pi^\pm \rho^\mp (L=1), \rho^0 \omega (L=1, 3), \pi^0 D (L=0, 2)$ $\pi^0 f (L=2), \pi^\pm B^\mp (L=0, 2)$
$1^+(1^+)$	$\pi^+ \pi^- (L=1), \rho^+ \rho^- (L=1, 3), \pi^\pm A_1^\mp (L=0, 2)$ $\pi^\pm A_2^\mp (L=2)$

Search for  $1^{-+}(1^-)$  exotic in  $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$

SIGNAL:  $\bar{p}p (^1S_0) \rightarrow \pi^0 X^0 (J_f=1)$   
 $\quad \quad \quad \searrow \pi^0 \eta (L=1)$

BACKGROUND:  $\bar{p}p (^1S_0) \rightarrow \pi^0 A_2^0 (J_f=2)$   
 $\quad \quad \quad \searrow \pi^0 \eta (L=2)$

$\bar{p}p (^1S_0) \rightarrow \pi^0 \delta^0 (J_f=0)$   
 $\quad \quad \quad \searrow \pi^0 \eta (L=0)$

also  $\sigma \eta (J_f=0)$ ,  $f \eta (J_f=2)$  couple to  $\pi^0 \pi^0 \eta$

Experimentally, look for  $\pi^0 \pi^0 \eta \rightarrow \underline{6\gamma}$  (CRYSTAL BARREL)

## Exotics in $\bar{p}p$ Annihilation:

### A) All neutral modes

$$1) \bar{p}p \rightarrow \pi^0 \omega_8 \rightsquigarrow 2\pi^0 (L_f=1)$$

$\pi^0 f, \pi^0 \sigma$  backgr.

$$2) \bar{p}p \rightarrow \pi^0 \omega_8 \rightsquigarrow \eta\eta (L_f=1)$$

$$3) \bar{p}p \rightarrow \pi^0 \rho_3^0 \rightsquigarrow \pi^0 \eta (L_f=1)$$

$$4) \bar{p}p \rightarrow \pi^0 \rho_3^0 \rightsquigarrow \pi^0 f (L_f=2) \rightsquigarrow 2\pi^0$$

### B) Charged Modes:

$$1) \bar{p}p \rightarrow \pi^\pm \rho_3^\mp \rightsquigarrow \pi^\mp B^0 \rightsquigarrow \pi^0 \omega \rightsquigarrow \pi^0 \gamma$$

CRYSTAL BARRE:

$$2) \bar{p}p \rightarrow \pi^\pm \rho_3^\mp \rightsquigarrow \pi^\mp D^0 \rightsquigarrow \pi^0 \pi^0 \eta$$

$$3) \bar{p}p \rightarrow \pi^\pm \rho_3^\mp \rightsquigarrow \pi^\mp \rho^0 \rightsquigarrow \pi^+ \pi^-$$

## Outlook:

$N\bar{N}$  annihilation very promising  
as a means of producing exotic  
mesons  $X$  in reactions  $N\bar{N} \rightarrow \pi X$

## Difficulties:

- 1)  $\Gamma_X$  large  $\rightarrow$  new states are broad resonance  
 $\rightarrow$  detailed amplitude analysis  
need to extract interference  
effects
- 2) branching ratios for  $N\bar{N} \rightarrow \pi X$  likely to be small  
 $\rightarrow$  no good theoretical estimate

## Advantages:

- 1)  $N\bar{N}$  annihilation potentially rich in  
gluonic excitations
- 2)  $N\bar{N}$  at rest provides control of initial  
quantum numbers ( $L=0$  or  $L=1$ )
- 3)  $\bar{p}p \rightarrow \pi^0 X^0, \pi^\pm X^\mp$ ;  $\bar{p}n \rightarrow \pi^- X^0$  gives quantum number  
filtration

**ANTIPROTON PRODUCTION CALCULATION  
BY THE MULTISTRING MODEL  
"VENUS" COMPUTER CODE**

**HIROSHI TAKAHASHI**

**DEPARTMENT OF NUCLEAR ENERGY  
BROOKHAVEN NATIONAL LABORATORY  
UPTON, NY**

**PRESENTED AT THE ANTIPROTON TECHNOLOGY WORKSHOP  
HELD AT BROOKHAVEN NATIONAL LABORATORY  
10 MAY 1989**

**ANTIPROTON PRODUCTION CALCULATION BY MULTISTRING  
MODEL VENUS COMPUTER CODE**

**RINOSHI TAKAHASHI  
KLAUS WERNER  
JAMES POWELL**

**Brookhaven National Laboratory**

**WORKSHOP ANTIPROTON TECHNOLOGY  
MAY 10, 1989**

**Brookhaven National Laboratory  
Upton, New York, 11973**

## Fermi Lab Source Performance

F. E. Mills. N I M P A 271 (1988) 176

Proton Energy  $E_{lab} = 120 \text{ GeV}$

Antiproton Energy  $= 8 \text{ GeV}$

Target : Cu

Cross Section Missing Factor  $\approx 2.5$

Mo Gormley.



## Antiproton Source (CERN, Fermi Lab)

Hojvat and Van Ginneken

Empirical formula

$$\frac{E}{\sigma_{\text{obs}}} \frac{d^3\sigma}{dP^3} = [k(1-x_R)^m e^{1/P^2} (-3P^2)]$$

$$\times [1 + 24S^{-2} e^{th(8x_R)}] [a e^{th(bP^2)} e^{th(-cx_R)}]$$

$$x_R = E/E_{\text{max}}$$

$$k = 0.065, \quad m = 8.0$$

Target	a	b	c
H	1.00	0.0	0.0
W	1.69	1.38	1.79
Pb	1.73	1.37	1.83

# VENUS

A multistring model

for ultrarelativistic heavy ion collision

---

- model for 'ordinary' collisions  
(no plasma)
- pp extrapolation (test: PA)
- string fragmentation consistent  
with  $e^+e^-$ ,  $\nu P$ ,  $\bar{\nu}P$ ,  $\mu P$  data
- Monte Carlo formulation  
(event generator)
- motivated by Regge theory  
(like DPM)


# Step Structure

## I) Multiple Scattering


Geometry +  $\sigma_{NN}$  determine NN coll



## II) individual NN collision



## III) String fragmentation



Multi String Model. Venus Code






Figure 1

# Feynman Field Fragmentation



elementary vertices :



Splitting function (using quark counting)

$$f(x) \sim x^\alpha (1-x)^{2n-1}$$

$n$ : number of spectators

$$\alpha = \begin{cases} 3/2 & \text{for baryon prod} \\ 0 & \text{for meson prod} \end{cases}$$

Table 4.1 The Particle Yield For Proton-Proton Collision (%)

Number of Hits:	15000	15000
Lab Energy of Projectile (GeV.)	200	1000
$\bar{p}$	4.53 (5.4)*	11.5 (16.2)*
$\bar{n}$	4.39	11.2
p	128.7	132.3
n	62.3	67.7
$\pi^-$	262.4	399.2
$\pi^+$	325.1	461.7
$\gamma$	735.8	1052.5
$k^-$	15.1	28.4
$k^+$	24.4	41.4
$\Sigma^+$	4.39	6.08
$\bar{\Sigma}^-$	0.37	1.36
$\Lambda^-$	2.0	6.3
$\Lambda^+$	14.1	22.3
$e^-$	4.4	6.0
$e^+$	4.4	6.0

\* ( ) is calculated by Hojovat and Van Ginneken's empirical formula.

Table 4.2 The Particle Yield For Proton-Pb Collision (%)

Number of Hits:	5822	5773
Lab Energy of Proton (GeV.)	200	1000
$\bar{p}$	7.07(9.0)*	21.38 (29.0)*
$\bar{n}$	6.93	22.37
p	213.59	227.94
n	218.83	226.64
$\pi^-$	751.96	1154.70
$\pi^+$	765.16	1168.16
$\gamma$	1998.35	2921.7
$\kappa^-$	27.75	65.70
$\kappa^+$	49.51	89.51
$\Sigma^+$	28.37	45.27
$\bar{\Sigma}^-$	3.48	9.41
$\Sigma^-$	6.10	9.41
$\bar{\Sigma}^+$	0.66	2.66
$\Lambda^-$	0.64	3.02
$\Lambda^+$	5.54	9.39
$e^-$	11.33	16.45
$e^+$	11.33	16.45

\* ( ) is calculated by Hojovat & Van Ginneken's empirical formula.




Figure 3.1







Figure 3.2



Iso. Kinestika.  
 quark Cascade Recombination Model  
 Analytic

Figure 3.3




Figure 3.4




Figure 3.5




Figure 3.6




Figure 3.9

Table 4.3 The Particle Yield For Si-Si (%)

	9831	9858
Number of Hits:	200	1000
Lab Energy of Projectile (GeV./A)		
$\bar{p}$	26.88 (82.5)*	79.86 (260.5)*
$\bar{n}$	27.41 (81.7)	79.20 (258.4)
p	680.24	710.38 (2238.7)
n	684.46	706.08 (2229.6)
$\pi^-$	2931.60	4353.5 (15226.2)
$\pi^+$	2935.06	4358.07 (15200.8)

\* ( )s are at central collision (b=0)

Table 4.4 The Particle Yield For O-Pb Collision (%)

Number of Hits: Lab Energy of Projectile (GeV./A)	7000	876
	200	1000
$\bar{p}$	40.04 (78.6)*	124.77
$\bar{n}$	37.84 (77.8)	122.14
p	153.5 (316.3)	1181.73
n	1235.35 (348.1)	1234.24
$\pi^-$	5298.5 (15216.2)	7899.2
$\pi^+$	5421.3 (14835.1)	7756.39

\* ( )s are at central collision ( $b=0$ ).






Figure 4.1




Figure 4.6




Figure 4.2




Figure 4.3





Figure 4.5



$$x = \frac{p_{||}}{p_{||\text{max}}}$$

Figure 4.7

## Central Collision

Xf spectrum concentrated to low Xf region

Recent Cern Experiment of Ox-Pb Collision

Pion Production 240 at 200 Gev Hydrodynamic  
107 Venus Calculation  
300 Venus Central

**Successive Collisions Increase P production**

specular Nucleus  $\sim 2$  times to Proton collision  
Multiple collision  $\pi \rightarrow \bar{P}$ .

## Modification of Venus Code

- Important Sampling Method
- Russian Roulette,

Careful Choice of Parameter is required

## Two Calculations of Collision Events

- Creation of the string
- The fragmentation of Strings
- Convolution of two functions

## More Sophisticated model

- Flux Tube model
- Quark Gluon Plasma Formation

Dynamics of Plasma expansion and Hadronization  
Heins et al Theory

- Fragmentation model using string model.
- Baldin et al. C-Cu (3.8 GeV/A)  $\bar{P}/\pi^-$  ratio. ~~100~~<sup>60</sup> times  
of P. nucleus collision.