

AD-A210 565

4

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 107

Quasi-Diffusion Between Phonon and Roton Gases in Two- and Three-Dimensional Liquid Helium

by

C. I. Um, C. W. Jun, H. J. Shin and Thomas F. George

Prepared for Publication

in

Journal of Low-Temperature Physics

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

July 1989

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

DTIC  
ELECTE  
JUL 24 1989  
S B D

89 7 24 067

**REPORT DOCUMENTATION PAGE**

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-107		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Quasi-Diffusion Between Phonon and Roton Gases in Two- and Three-Dimensional Liquid Helium			
12. PERSONAL AUTHOR(S) C. I. Um, C. W. Jun, H. J. Shin and Thomas F. George			
13a. TYPE OF REPORT	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) July 1989	15. PAGE COUNT 22
16. SUPPLEMENTARY NOTATION Prepared for publication in the Journal of Low-Temperature Physics			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	QUASI-DIFFUSION
			LIQUID HELIUM
			PHONON GAS
			THREE-DIMENSIONAL
			ROTON GAS
			TEMPERATURE VARIATIONS
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  Temperature variations of the quasi-diffusion constant and thermal diffusion ratio between phonon and roton gases in two- and three-dimensional liquid helium are evaluated explicitly through the scattering of phonons by rotons. The diffusion constants in both dimensions decrease exponentially over the whole temperature range, while the thermal diffusion ratios increase and then decrease as temperature increases in the range from ~ 0.6 to ~ 1.1 K.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson		22b. TELEPHONE (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL

Quasi-diffusion between phonon and roton gases in two- and  
three-dimensional liquid helium

Chung-In Um and Chul-Won Jun  
Department of Physics  
College of Science  
Korea University  
Seoul 136-701, Korea

Hyun-Joon Shin  
Research Institute of Industrial Science and Technology  
1 Gyedong-dong, Pohang City  
Kyoungbuk 790-300, Korea

Thomas F. George  
Departments of Chemistry and Physics  
239 Fronczak Hall  
State University of New York at Buffalo  
Buffalo, New York 14260

Temperature variations of the quasi-diffusion constant and thermal diffusion ratio between phonon and roton gases in two- and three-dimensional liquid helium are evaluated explicitly through the scattering of phonons by rotons. The diffusion constants in both dimensions decrease exponentially over the whole temperature range, while the thermal diffusion ratios increase and then decrease as temperature increases in the range from  $\sim 0.6$  to  $\sim 1.1$  K.

PACS Nos.: 61.20.p, 66.10Cb, 67.40.Pm

## 1. Introduction

Since Landau's phenomenological theory of the superfluids,<sup>1</sup> it is well known that liquid helium at low frequencies, long wavelengths and low temperatures can be described by Landau's two-fluid equations.<sup>2</sup> These equations involve not only the normal and superfluid density and thermodynamic functions, but also the various sounds and transport kinetic coefficients in liquid  $^3\text{He}$  and  $^4\text{He}$ . Concerning the kinetic coefficients, Landau and Khalatnikov<sup>3</sup> first investigated the thermal conductivity and viscosity, which were also studied later by others.<sup>4</sup> Diffusive processes<sup>5</sup> in liquid  $^3\text{He}$  and  $^3\text{He}$ - $^4\text{He}$  mixtures have been extensively studied: Reuppeiner et al<sup>6</sup> measured the thermal conductivity ( $K$ ), thermal diffusion ratio ( $K_T$ ) and mass diffusivity ( $D$ ) as a function of  $^3\text{He}$  number concentration, and Behringer and Meyer<sup>7</sup> investigated diffusive relaxation processes in the normal and superfluid phase in liquid  $^3\text{He}$ - $^4\text{He}$  mixtures. All these quantities mentioned above are closely dependent on the elementary excitation, which is appropriate to describe the given Bose system. Motivated by these studies of  $^3\text{He}$ - $^4\text{He}$  mixtures, in the present paper we evaluate the temperature dependence of the diffusion constant ( $D$ ) and thermal diffusive ration ( $K_T$ ) between the phonon and roton gases in consideration of the scattering of phonons by rotons in two- and three-dimensional liquid  $^4\text{He}$ .

Recently we have obtained microscopically the Landau-type excitation spectrum, which is (anomalous) phonon-like at low momentum and roton-like for large momenta in the ring diagram approximation.<sup>8</sup> Using this excitation spectrum we have successfully analyzed the sounds,<sup>9</sup> sound attenuations,<sup>10</sup> kinetic coefficients,<sup>11</sup> etc., in two- and three-dimensional liquid  $^4\text{He}$ .

Through the calculations we can consider the excitation spectra in both dimensions:

$$\xi(p) = c_0 p [1 + \gamma p^2 - \delta p^4 + \dots] \quad (2D) \quad (1.1)$$

$$\xi(p) = c_0 p [1 + \delta_1 p^2 - \delta_2 p^4 + \dots] \quad (3D) \quad (1.2)$$

and

$$\xi(p) = \Delta + \frac{(p-p_0)^2}{2\mu} \quad (2D \text{ and } 3D) \quad (1.3)$$

where  $p$  is the roton momentum,  $c_0$  is the sound velocity, and  $\Delta$ ,  $p_0$  and  $\mu$  are the roton parameters (energy gap, momentum and effective mass, respectively). The subscript zero refers to  $T = 0$  K. All coefficients in Eqs. (1.1) and (1.2) are positive, which are determined by the potential parameters. Here, we have adopted a soft potential with a Lennard-Jones type tail, which helps to make a smooth connection of the attractive part with the soft repulsive core. In the next section we evaluate the differential phonon-roton scattering cross section, and calculations of the diffusion constant and thermal diffusion ratio are described in Sec. 3. Finally, the results and discussion are presented in Sec. 4.

## 2. Phonon-roton differential scattering cross section

To obtain the two-dimensional phonon-roton scattering cross section, in this section we consider a roton in the presence of the phonon field. We can treat this roton as a particle in a moving liquid  $^4\text{He}$ . Then there appears an additional term  $-\vec{p} \cdot \vec{v}$ , which can be written in symmetric form<sup>12</sup>

$$-\frac{1}{2}(\vec{p} \cdot \vec{v} + \vec{v} \cdot \vec{p}) \quad (2.1)$$

QUALITY  
CONTROLLED

<input checked="" type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>

Codes

/or

Dist	Special
A-1	

where  $\vec{P}$  and  $\vec{v}$  are the momentum and velocity operators.<sup>9</sup> The phonon field changes the density of the medium, and thus we may expand the roton energy [Eq. (1.3)] in terms of the density  $\rho'$  to second order ( $\rho' = \rho - \rho_0$ ) as

$$H_r = H_{r_0} + \frac{\partial}{\partial \rho} (\Delta + \frac{1}{2\mu} (P-P_0)^2) \rho' + \frac{1}{2!} \frac{\partial^2}{\partial \rho^2} (\Delta + \frac{1}{2\mu} (P-P_0)^2) \rho'^2 + \dots \quad (2.2)$$

where  $H_{r_0}$  is given by Eq. (1.3). Since the value of most rotons is close to  $P_0$ , we may neglect  $(P-P_0)$  and replace  $P$  by  $P_0$ . We may also drop the term  $(\frac{\partial \Delta}{\partial \rho} \rho')$ , which is much smaller than (2.1). Then the interaction energy between phonons and rotons can be written as

$$V = - \frac{1}{2} (\vec{P} \cdot \vec{v} + \vec{v} \cdot \vec{P}) + \frac{1}{2} \left\{ \frac{\partial^2 \Delta}{\partial \rho^2} + \frac{1}{\mu} \left( \frac{\partial P_0}{\partial \rho} \right)^2 \right\} \rho' \quad (2.3)$$

where the terms in the second bracket of Eq. (2.3) have magnitudes on the order of  $10^{-1}$  to 1 in three-dimensional liquid helium.<sup>13</sup>

When the roton changes momentum  $\vec{P}$  to  $\vec{P}'$ , it absorbs a phonon with momentum  $\vec{p}$  and emits a phonon with momentum  $\vec{p}'$ . Here we may consider two intermediate processes: (I)  $\vec{P} + \vec{p} \rightarrow \vec{P}' = \vec{P} + \vec{p} - \vec{p}'$  and (II)  $\vec{P} - \vec{p}' \rightarrow \vec{P}' = \vec{P} - \vec{p}' + \vec{p}$ . Since the roton momentum is much larger than that of the phonon, we may view this interaction as similar to that of heavy and light particles. The momentum and energy conservation law in collision processes can be written as

$$cp + \frac{1}{2\mu} (P-P_0)^2 = cp' + \frac{1}{2\mu} (|\vec{P} + \vec{p} - \vec{p}'| - P_0)^2 \quad (2.4)$$

Under the conditions  $p, p' \ll P_0$  and  $E - cp \ll 3 \mu c^2$ , Eq. (2.4) becomes

$$p - p' = \frac{1}{2\mu c P_0^2} (P_0 \hat{m} \cdot (p \hat{n} - p' \hat{n}'))^2$$

$$= \frac{P^2}{2\mu c} (\hat{m} \cdot (\hat{n} - \hat{n}'))^2, \quad (2.5)$$

where  $\hat{m}$ ,  $\hat{n}$  and  $\hat{n}'$  are the unit vectors directed along  $\vec{P}_0$ ,  $\vec{p}$  and  $\vec{p}'$ , respectively. Therefore energy conservation implies  $p \approx p'$ . This means that particles do not change the magnitude of momentum, but change its direction. Taking account of  $p, p' \ll P_0$  and  $P = P_0$ , the matrix element given in the differential cross section

$$d\sigma = \left(\frac{2\pi}{\hbar c_0}\right) |\langle F | H_3 | I \rangle|^2 \delta(E_F - E_I) \frac{d\vec{P}}{(2\pi\hbar)^2} \quad (2.6)$$

can be written in second-order perturbation as

$$\langle F | H_3 | I \rangle = \frac{P_0 P}{2\rho_0} \{ [(\hat{m} \cdot \hat{n}) + (\hat{m} \cdot \hat{n}')] (\hat{n} \cdot \hat{n}') + \frac{P_0}{\mu c} (\hat{m} \cdot \hat{n})^2 (\hat{m} \cdot \hat{n}')^2 + A \}, \quad (2.7)$$

$$A = \frac{\rho^2}{P_0 c} \left( \frac{\partial^2 \Delta}{\partial \rho^2} + \frac{1}{\mu} \left( \frac{\partial P_0}{\partial \rho} \right)^2 \right). \quad (2.8)$$

Substituting Eqs. (2-7)-(2.8) into Eq. (2.6) and performing the integration over  $p'$ , we obtain

$$d\sigma = \frac{P_0^2 p^3}{8\pi\hbar^3 \rho_0^2 c^2} \{ [(\hat{m} \cdot \hat{n}) + (\hat{m} \cdot \hat{n}')] (\hat{m} \cdot \hat{n}') + \frac{P_0^2}{\mu c} (\hat{m} \cdot \hat{n})^2 (\hat{m} \cdot \hat{n}')^2 + A^2 \} d\theta \quad (2.9)$$

Averaging Eq. (2.9) over all directions of roton momentum, we finally obtain

$$d\sigma = \frac{P_0^2 p^3}{8\pi\hbar^3 \rho_0^2 c^2} \left\{ (1 + \cos\psi) \cos^2\psi + \frac{1}{128} \left(\frac{P_0}{\mu c}\right)^2 (35\cos^4\psi + 3\sin^4\psi + 30\cos^2\psi \sin^2\psi) + \frac{1}{4} \frac{P_0 A}{\mu c} (3\cos^2\psi + \sin^2\psi) + A^2 \right\} d\psi \quad ,$$

where  $\psi$  is the angle between the incident and scattered phonons.

Through the similar calculations we can obtain the differential scattering cross section of phonons by rotons in bulk liquid  ${}^4\text{He}$ :<sup>14</sup>

$$d\sigma = \left(\frac{P_0 p}{4\pi\hbar^2 \rho_0 c}\right)^2 \left\{ \frac{2}{3}(1 + \cos\psi) \cos^2\psi + \frac{1}{105} \left(\frac{P_0}{\mu c}\right)^2 (1 + 8\cos^2\psi + \frac{3}{8}\cos\psi) + \frac{2A}{12} \left(\frac{P_0}{\mu c}\right) (1 + 2\cos^2\psi) + A^2 \right\} d\Omega \quad , \quad (2.11)$$

where  $d\Omega$  is the solid angle.

### 3. Diffusion constant and thermal diffusion ratio

Since the excitation spectrum in liquid  ${}^4\text{He}$  consists of phonons and rotons, we may treat liquid  ${}^4\text{He}$  as a mixed phonon-roton gas. In the temperature range from  $-0.7$  to  $-1.2$  K, where the scattering of phonons by rotons is dominant, we consider the diffusion of phonons and rotons and also



the thermal diffusion ratio under the existence of a temperature gradient. There thermal processes in the mixture of a phonon-roton gas are very similar to those of light particles in heavy particles. Since the roton momentum is much larger than that of the phonon, we may treat the interactions between phonons and rotons as similar to those between light and heavy particles. Therefore, we may assume that when a collision occurs only between a phonon and roton, the roton is at rest and the phonon momentum changes only its direction with the same magnitude.

Let  $n_p$  and  $n_r$  be the number densities of the phonon and roton, respectively, and  $\theta$  the angle between the direction of a phonon with momentum  $\vec{p}$  and the x-axis. Equation (2.11) represent the differential scattering cross section that the momentum  $\vec{p}$  of a phonon will be changed to the momentum  $\vec{p}'$  directed into the solid angle element  $d\Omega$ . The collision probability of a phonon per unit time becomes  $n_r c d\sigma$ , where  $c$  is the phonon velocity. Let the number of phonons in a given unit volume with momentum between  $p$  and  $p + dp$  and directed into the solid angle  $d\Omega$  be  $n(p, \theta, x) d^3p$ . Then the total number of phonons changing  $\vec{p}$  to  $\vec{p}'$  in  $d\Omega$  and conversely  $\vec{p}'$  to  $\vec{p}$  in  $d\Omega'$  are, respectively, given by

$$d^3p \int n_r c n(p, \theta, x) d\sigma, \quad (3.1)$$

$$d^3p \int n_r c n(p', \theta', x) d\sigma. \quad (3.2)$$

The difference between the above two processes,

$$d^3p n_r c \int [n(p', \theta', x) - n(p, \theta, x)] d\sigma, \quad (3.3)$$

represents the change in the number of phonons in the volume element  $d^3p$ , which must be equal to the total time derivative of  $n$ :

$$J_{\text{phr}}(n) = c \cos\theta \frac{\partial n}{\partial x} - n_r c \int [n(c', \theta', x) - n(p, \theta, x)] d\sigma' \quad (3.4)$$

If the temperature and concentration gradients vary very slowly, we can express the phonon distribution function as

$$n = n_0(p, x) + \delta n(p, \theta, x) \quad (3.5)$$

where  $\delta n$  is a small correction term to the equilibrium distribution function  $n_0$ . Now we assume that  $\delta n$  is linear in the gradient of the concentration and temperature, and has the form

$$\delta n = \cos\theta f(p, x) \quad (3.6)$$

Substitution of Eq. (3.6) and  $\cos\theta' = \cos\theta\cos\psi + \sin\theta\sin\psi\cos(\psi-\psi')$  in Eq. (3.4) and then integration over the solid angle yields

$$J_{\text{phr}}(n) = \cos\theta n_r c f(p, x) \frac{a^4 P_0^2 k_B^4 T^4}{4\pi h^4 \rho_0^2 c^6} \Gamma \quad (3.7)$$

$$\Gamma = \frac{4}{45} + \frac{1}{25} \left(\frac{P_0}{\mu c}\right)^2 + \frac{2}{9} \left(\frac{P_0 A}{\mu c}\right) + A_0^2$$

Here we have made use of the relation  $p = ak_B T/c$ , where  $a$  is a constant<sup>15</sup> (see the Appendix). From Eqs. (3.4) and (3.7), we obtain the following expression for  $f(p,x)$ :

$$f(p,x) = - \frac{1}{n_r} \frac{4\pi h^4 \rho_0 c^6}{a^4 P_0^2 \Gamma n_r k_B^4 T^4} \frac{\partial n}{\partial x} \quad (3.8)$$

The diffusion flux  $i$  along the  $x$ -axis is

$$i = \int n c \cos\theta d^3 p = c \int \cos^2\theta f(p,x) d^3 p \quad (3.9)$$

Substituting Eq. (3.8) into Eq. (3.9) and integrating over momentum  $p$  gives

$$i = \frac{4\pi h^4 \rho_0 c^7}{3a^4 P_0^2 \Gamma n_r k_B^4 T^4} \frac{\partial n}{\partial x} \quad (3.10)$$

Let the phonon concentration of the mixture be  $\eta = n_p/(n_0+n_r) = n_0/n$ . Then the diffusion flux  $i$  becomes

$$i = \frac{4\pi h^4 \rho_0 c^7}{3a^4 P_0^2 \Gamma n_r k_B^4 T^4} n \left[ \frac{\partial \eta}{\partial x} + \eta \frac{\partial}{\partial T} \ln(n) \frac{\partial T}{\partial x} \right] \quad (3.11)$$

Comparing Eq. (3.11) with the expression<sup>16</sup> for the diffusion flux

$$i = -nD(\vec{\nabla}\eta + \frac{K_T}{T} \vec{\nabla}T) \quad (3.12)$$

we obtain

$$D = \frac{4\pi\hbar\rho_0^2 c^7}{3a^4 P_0^2 \Gamma n_r k_B^4} \quad (3.13)$$

$$K_T = \eta T \frac{\partial}{\partial T} \ln(n) \quad (3.14)$$

Since the number densities of phonons and rotons per unit volume can be given by

$$n_p = 2! \zeta(3) 4\pi \left(\frac{k_B T}{2\pi\hbar c}\right)^3 \quad (3D) \quad (3.15)$$

$$n_r = \frac{2P_0^2 (\mu k_B)^{1/2}}{(2\pi\hbar^2)^{3/2}} T^{1/2} e^{-\Delta/k_B T} \quad (3D) \quad (3.16)$$

we obtain the diffusion coefficient and thermal diffusion ratio of phonons in bulk liquid helium:

$$D(T) = \frac{(2\pi)^{5/2} \rho_0^2 (\hbar c)^7}{3a^4 P_0^4 \mu^{1/2} \Gamma k_B^9} T^{-9/2} e^{\Delta/k_B T} \quad (3.17)$$

$$K_T = \frac{3 + \left(\frac{\zeta(3)}{\sqrt{2}}\right)^{-1} (\pi\mu)^{1/2} P_0^2 c^3 \left(\frac{1}{2} + \frac{\Delta}{k_B T}\right) (k_B T)^{-5/2} e^{-\Delta/k_B T}}{\left[1 + \left(\frac{\zeta(3)}{\sqrt{2}}\right)^{-1} (\pi\mu)^{1/2} P_0^2 c^3 (k_B T)^{-5/2} e^{-\Delta/k_B T}\right]^2} \quad (3.18)$$

Through similar calculations as done above, we obtain the diffusion coefficient and thermal diffusion ratio in thin helium films as

$$D(T) = \frac{(2^5 \pi)^{1/2} \rho_0^{2.5} c^6}{a^{13} P_0^3 \mu^{1/2} \Gamma k_B^{7/2}} T^{-7/2} e^{\Delta/k_B T}, \quad (3.19)$$

$$K(T) = \frac{2 + \{\zeta(2)\}^{-1} (2\pi\mu)^{1/2} P_0 c^2 \left(\frac{1}{2} + \frac{\Delta}{k_B T}\right) (k_B T)^{-3/2} e^{-\Delta/k_B T}}{[1 + \{\zeta(2)\}^{-1} (2\pi\mu)^{1/2} P_0 c^2 (k_B T)^{-3/2} e^{-\Delta/k_B T}]^2}, \quad (3.20)$$

where we used the number densities of phonons and rotons per unit surface given by

$$n_p = \frac{\zeta(2)}{2\pi} \left(\frac{k_B T}{\mu c}\right)^2, \quad (3.21)$$

$$n_r = \frac{P_0}{2\pi\mu^2} (2\pi\mu k_B T)^{1/2} e^{-\Delta/k_B T}, \quad (3.22)$$

and  $\Gamma'$  is given as

$$\Gamma' = \frac{1}{4} + \frac{9}{32} \left(\frac{P_0}{\mu c}\right)^2 + \frac{P_0 A}{\mu c} + 2A^2.$$

#### 4. Results and discussion

Since the elementary excitation spectrum in liquid helium consists of the combined phonon-roton gas, we may consider the diffusion of phonons into the roton gas, and also may include the thermal diffusion under the existence of the temperature gradient. We may treat the diffusion and thermal diffusion of the phonons in the roton gas as similar to those of light particles in heavy particle.. For convenience we neglected phonon-phonon or roton-roton scatterings, permitting only phonon-roton scatterings.

The thermal diffusion ratio represents the ratio of the thermal diffusion coefficient due to the temperature gradient to the diffusion coefficient given by the concentration gradient. The diffusion coefficient must be positive, but the diffusion ratio may be positive or negative. The negative value of the coefficient  $K_T$  means that the heavy particles will appear in the low-temperature region and the light particles in the high-temperature region. In  $^3\text{He}$ - $^4\text{He}$  mixtures<sup>17</sup> the ratio  $K_T$  becomes negative near the plait point and positive near transition temperature. Since thermal diffusion ratios in our calculations are positive, phonons and rotons are located in the low- and high-temperature regions, respectively.

The diffusion constant and thermal diffusion ratios of the phonon gas in bulk liquid helium and in thin helium films are given by Eqs. (3.17)-(3.20). The potential and roton parameters are chosen by the analysis of the excitation spectrum<sup>18</sup> for the bulk case and the specific data<sup>8</sup> for the thin helium films (Table I). For the density variations of the roton parameters, we have adopted the results obtained by Khalatnikov,<sup>14</sup> who analyzed the data of Dietrich et al<sup>19</sup> given by

$$\left(\frac{\rho}{\Delta}\right)\left(\frac{\partial\Delta}{\partial\rho}\right) = -1.142, \quad \left(\frac{\rho^2}{\Delta}\right)\left(\frac{\partial^2\Delta}{\partial\rho^2}\right) = -2.06, \quad \left(\frac{\rho}{P_0}\right)\left(\frac{\partial P_0}{\partial\rho}\right) = \frac{1}{3} .$$

The above parameters in thin helium films are not known and thus are assumed to have the same values in the bulk case. The constants  $a$  and  $a'$  in Eqs. (3.7) and (3.19) can be determined by the calculation of the collision time for the phonon-roton scatterings, which characterize the thermal conductivity and viscosity. The numerical values of these  $a$  and  $a'$  are 6.28 and 4.67, respectively.<sup>15</sup>

Figure 1 illustrates the diffusion constant  $D$  as a function of temperature in the bulk and for films. As temperature increases, the roton density increases more rapidly than the phonon density. Therefore, the mean free path of phonons becomes shorter and the coefficient decreases rapidly.

The temperature variations of the coefficients  $K_T$  are shown in Fig. 2. Below  $\sim 0.6$  K the coefficient is almost constant. In the temperature range from  $\sim 0.6$  to  $\sim 1.1$  K the coefficient ratio increases, which means that phonon-roton scatterings play a main role in the transport phenomena. This temperature range agrees well with the results obtained by Landau and Khalatnikov.<sup>12</sup> For temperature far below  $\sim 0.6$  K, the phonon population is dominant, and thus phonon-phonon interactions becomes important, while roton-roton interactions and five-phonon processes will be dominant above  $\sim 1.1$  K.

In thin helium films the temperature range in which phonon-roton scatterings are effective and dominant becomes  $\sim 0.3$  K  $< T < 0.8$  K. This range is in good agreement with the results obtained from the evaluation of kinetic coefficients in thin helium films.<sup>9</sup> All arguments mentioned with respect to the bulk helium will be valid.

#### Acknowledgments

This work was supported by the Special Research Fund (SRF), Ministry of Science and Technology, Republic of Korea. Support was also received by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

### Appendix

Making use of Eqs. (3.13)-(3.15), (3.21) and (A1) from Ref. 15, we can easily obtain the characteristic time  $\tau_{\text{ph-r}}^{-1}$  expressed as a function of the phonon momentum and temperature, respectively, as

$$\tau_{\text{ph-r}}^{-1} = N_r \frac{P_0 P^3}{8\hbar^3 \rho^2 c} \left[ \frac{1}{4} + \frac{9}{32} \left( \frac{P_0}{\mu c} \right)^2 + \frac{P_0}{\mu c} A + 2A^2 \right] , \quad (\text{A1})$$

and

$$\tau_{\text{ph-r}}^{-1} = \frac{6! \zeta(6)}{3! \zeta(3)} N_r \frac{P_0^2 k_B^3 T^3}{8\hbar^3 \rho^2 c^4} \left[ \frac{1}{4} + \frac{9}{32} \left( \frac{P_0}{\mu c} \right)^2 + \frac{P_0}{\mu c} A + 2A^2 \right] , \quad (\text{A2})$$

where  $N_r$  is the number of rotons per unit area. Setting (A1) equal to (A2), we get

$$pc = \left[ \frac{6! \zeta(6)}{3! \zeta(3)} \right]^{1/3} k_B T . \quad (\text{A3})$$

Therefore, we obtain  $a' = 4.67$  in two-dimensional liquid helium.

Through a similar calculation we can evaluate the characteristic time expressed by temperature and phonon momentum:

$$\tau_{\text{ph-r}}^{-1} = 4\pi c N_r \left( \frac{P_0 P^2}{4\pi\hbar^2 \rho c} \right)^2 \left[ \frac{4}{45} + \frac{1}{25} \left( \frac{P_0}{\mu c} \right)^2 + \frac{2}{9} \left( \frac{P_0}{\mu c} A \right) + A^2 \right] . \quad (\text{A4})$$

and



$$\tau_{\text{ph-r}}^{-1} = \frac{4\pi}{c} N_r \left( \frac{P_0 k_B T^2}{\hbar^2 \rho c^2} \right)^2 \left[ \frac{4}{45} + \frac{1}{25} \left( \frac{P_0}{\mu c} \right)^2 + \frac{2}{9} \left( \frac{P_0 A}{\mu c} \right) + A^2 \right] \quad (\text{A5})$$

From Eqs. (A4)-(A5) we obtain  $a = 2\pi$  in the bulk case.

## References

1. L. D. Landau, *J. Phys.*, U.S.S.R. 11, 91 (1947).
2. P. C. Hohenberg and P. C. Martin, *Ann. Phys. (NY)* 34, 291 (1965).
3. L. D. Landau and I. M. Khalatnikov, *Zh. Eksp. Teor. Fiz.* 19, 637, 709 (1949).
4. S. G. Hedge, R. A. Gordery and W. I. Glaberson, *Phys. Rev. B* 26, 2690 (1982); T. R. Kirkpatrick and J. R. Dorfman, *J. Low Temp. Phys.* 58, 339 (1985); G. Agnolet, S. L. Teitel and J. D. Reppy, *Phys. Rev. Lett.* 47, 1537 (1981).
5. E. D. Siggia and D. R. Nelson, *Phys. Rev. B* 15, 1427 (1977); D. Roe, G. Ruppeiner and H. Meyer, *J. Low Temp. Phys.* 27, 747 (1977); D. Roe and H. Meyer, *J. Low Temp. Phys.* 28, 349 (1977).
6. G. Ruppeiner, M. Ryschkewitsch and H. Meyer, *J. Low Temp. Phys.* 41, 179 (1980).
7. R. P. Behringer and H. Meyer, *J. Low Temp. Phys.* 46, 407 (1982); 46, 435 (1982).
8. A. Isihara and C. I. Um, *Phys. Rev. B* 19, 5725 (1979); A. Isihara, C. I. Um and S. T. Choh, *Physica (Utrecht)* 100B, 74 (1980); C. I. Um and W. H. Kahng, *J. Kor. Phys. Soc.* 13, 57 (1980).
9. C. I. Um, W. H. Kahng, S. T. Choh and A. Isihara, *Phys. Rev. B* 29, 5203 (1984); C. I. Um, H. K. Oh, W. H. Kahng and A. Isihara, *Phys. Rev. B* 34, 6151 (1986); 33, 7550 (1986); 28, 2509 (1983).
10. C. I. Um, W. H. Kahng, S. W. Nam and A. Isihara, *Physica* 114B, 191 (1982); A. Isihara, C. I. Um, C. W. Chum, W. H. Kahng and S. T. Choh, *Jpn. J. Appl. Phys.* 26, Suppl. 26-3, 297 (1987).
11. C. I. Um, C. W. Chun, W. H. Kahng and A. Isihara, *Jpn. J. Appl. Phys.* 26, Suppl. 26-3, 295 (1987).
12. I. M. Khalatnikov and D. M. Chernikova, *Zh. Eksp. Teor. Fiz.* 49, 1957 (1965); 50, 411 (1966); I. M. Khalatnikov, *An introduction to the theory of superfluidity* (Benjamin, New York, 1965), Chapt. 7; K. H. Bennemann and J. B. Ketterson, *The physics of liquid and solid helium* (Wiley, New York, 1976), Part 1, Chapt. 1.
13. D. Henshaw and A. Woods, *Phys. Rev.* 121, 266 (1961).
14. I. M. Khalatnikov and Yu. A. Matveev, *J. Low Temp. Phys.* 51, 99 (1983).
15. C. I. Um, C. W. Jun, W. H. Kahng and T. F. George, *Phys. Rev. B* 38, 8838 (1988).
16. L. D. Landau and E. M. Lifshitz, *Fluid mechanics* (Pergamon, New York, 1959), Chapt. 6, p. 224.

17. L. H. Cohen, M. L. Dingus and H. Meyer, J. Low Temp. Phys. 61, 79 (1985); Phys. Rev. Lett. 50, 1058 (1983).
18. C. I. Um, S. T. Choh and A. Isihara, Phys. Rev. B 23, 4498 (1981); R. J. Donnelly, J. A. Donnelly and R. N. Hills, J. Low Temp. Phys. 44, 471 (1981).
19. O. W. Dietrich, E. H. Graf, C. H. Huang and L. Passell, Phys. Rev. A 5, 1377 (1972).

Table I. Theoretical Parameters

	$\eta$	$\Delta/k_B$ (K)	$q_0$ ( $\text{\AA}^{-1}$ )	$\mu$	$C$ (m/s)
3D	$2.18 \times 10^{-2} \text{\AA}^{-3}$	8.616	1.930	$0.153m_{\text{He}}$	238.21
2D	$2.79 \times 10^{-2} \text{\AA}^{-2}$	4.12	1.02	$0.77m_{\text{He}}$	164.4

**Figure Captions**

1. Diffusion constants of phonons in the roton gas as a function of temperature. The solid and dotted lines represent the theoretical curves in two- and three dimensional liquid  ${}^4\text{He}$ .
2. Thermal diffusion ratio of phonons in the roton gas versus temperature. The solid and dotted lines represent the theoretical curves in two- and three-dimensional liquid  ${}^4\text{He}$ .

Fig. 1





Fig. 2



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Duda Naval Weapons Support Center Code SOC Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code LS2 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1



ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. <sup>Arnold</sup> Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburg  
Chemistry Building  
Pittsburg, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton 509 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853