

MARINE PHYSICAL LABORATORY

SCRIPPS INSTITUTION OF OCEANOGRAPHY

San Diego, California 92152

Statistics of a Chi-Square Random Variable
Obtained from Independent Gaussian Samples
with a Non-Zero Mean and Arbitrary Variance

Richard K. Brienzo

AD-A214 477

DTIC
ELECTE
NOV 22 1989
S B D

MPL TECHNICAL MEMORANDUM 405

MPL-U-40/88

January 1989

Approved for public release; distribution unlimited.

**BEST
AVAILABLE COPY**

89 11 21 052

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			Approved for public release; distribution unlimited.		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) MPL TECHNICAL MEMORANDUM 405 [MPL-U-40/88]			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Marine Physical Laboratory		6b. OFFICE SYMBOL (If applicable) MPL		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research Department of the Navy	
6c. ADDRESS (City, State, and ZIP Code) University of California, San Diego Scripps Institution of Oceanography San Diego, CA 92152			7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-80-C-0220	
8c. ADDRESS (City, State, and ZIP Code) Department of the Navy 800 North Quincy Street Arlington, VA 22217-5000			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) STATISTICS OF A CHI-SQUARE RANDOM VARIABLE OBTAINED FROM INDEPENDENT GAUSSIAN SAMPLES WITH A NON-ZERO MEAN AND ARBITRARY VARIANCE					
12. PERSONAL AUTHOR(S) Richard K. Brienzo					
13a. TYPE OF REPORT Tech Memo		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) January 1989	
15. PAGE COUNT 8					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	chi-square random variable		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The mean and variance of a chi-square random variable are generally given for the case in which the chi-square random variable is derived from a process having a zero mean and unit variance. In this report, the mean and variance of the random variable found by squaring and summing N samples of an independent Gaussian process with a non-zero mean and arbitrary variance is derived.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL W. S. Hodgkiss			22b. TELEPHONE (Include Area Code) (619) 534-1798		22c. OFFICE SYMBOL MPL

Statistics of a chi-square random variable obtained from independent Gaussian samples with a non-zero mean and arbitrary variance

Richard K. Brienzo

Marine Physical Laboratory
Scripps Institution of Oceanography
University of California, San Diego
San Diego, CA. 92152

ABSTRACT

The mean and variance of a chi-square random variable are generally given for the case in which the chi-square random variable is derived from a process having a zero mean and unit variance. In this report, the mean and variance of the random variable found by squaring and summing N samples of an independent Gaussian process with a non-zero mean and arbitrary variance is derived.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Statistics of a chi-square random variable obtained from independent Gaussian samples with a non-zero mean and arbitrary variance

1. Introduction

In this report, general expressions are derived for the mean and variance of a random variable that is found by squaring and summing N independent samples of a Gaussian process x_i , which has a non-zero mean and an arbitrary variance (i.e. $\sum_{i=1}^N x_i^2$).

A common structure which generates such a random variable is an energy detector (Figure 1). Each output sample is obtained by squaring and summing N input samples. This is the optimal detector for an unknown signal buried in white Gaussian noise, and is often used as a post-processor for other routines (for example, the output of a beamformer may be run through an energy detector to determine if a signal is present).




Figure 1. Energy detector.
Finds the energy in N samples of $x(n)$. Each output sample is produced by squaring and summing N input samples.

Chi-square random variables

Let z_1, z_2, z_3, \dots be normally distributed, independent random variables with zero mean and a variance of one, and define a new random variable

$$\chi_N^2 = z_1^2 + z_2^2 + z_3^2 + \dots + z_N^2 \quad (1)$$

The variable in (1) is called a chi-square random variable with N degrees of freedom. The density function of χ_N^2 approaches that of a normally distributed random variable for large N ($N > 30$), and is non-symmetric for smaller N . The mean and variance of the chi-square random variable in (1) is given by [1, page 105],

$$\mu_{\chi_N^2} = E\{\chi_N^2\} = N \quad (2)$$

$$\sigma_{\chi_N^2}^2 = \text{var}[\chi_N^2] = 2N \quad (3)$$

Details about the density and distribution functions may be found in [1, Section 4.2.2]. Random variables with a non-zero mean that are squared and summed have a non-central chi-square distribution [3,4].

A time series x can always be transformed to have a mean of zero and variance equal to one by defining the standardized variable z_i to be

$$z_i = \frac{x_i - \mu}{\sigma} \quad (4)$$

In some cases, it is desirable to find the statistics of the random variable formed by squaring and summing N values of x , which does not necessarily have a zero mean or variance equal to one. In the next section, general expressions for the mean and variance of a random variable that is obtained from squaring and summing independent Gaussian samples with a non-zero mean μ and arbitrary variance σ^2 are derived.

2. Derivation

Given the standardized random variable in (4), a chi-square random variable with N degrees of freedom is obtained by squaring the z_i and summing over N samples,

$$\begin{aligned} \chi_N^2 &= \sum_{i=1}^N z_i^2 \\ &= \frac{1}{\sigma^2} \left[\sum_{i=1}^N (x_i - \mu)^2 \right] \\ &= \frac{1}{\sigma^2} \left[\sum_{i=1}^N (x_i^2 - 2\mu x_i + \mu^2) \right] \end{aligned} \quad (5)$$

$$= \frac{1}{\sigma^2} \left[\sum_{i=1}^N x_i^2 - 2\mu \sum_{i=1}^N x_i + N\mu^2 \right] \quad (6)$$

$$\begin{aligned} E\{\chi_N^2\} &= \frac{1}{\sigma^2} E\left\{\sum_{i=1}^N x_i^2\right\} - \frac{2\mu}{\sigma^2} E\left\{\sum_{i=1}^N x_i\right\} + \frac{N\mu^2}{\sigma^2} \\ &= \frac{1}{\sigma^2} E\left\{\sum_{i=1}^N x_i^2\right\} - \frac{N\mu^2}{\sigma^2} \end{aligned}$$

Solving for the N squared and summed samples of x_i , and using Equation (2) gives

$$\begin{aligned} E\left\{\sum_{i=1}^N x_i^2\right\} &= N\sigma^2 + N\mu^2 \\ &= N(\sigma^2 + \mu^2) \end{aligned} \quad (7)$$

By definition, the variance of the chi-square random variable is given by

$$\text{var}[\chi_N^2] = E\{[\chi_N^2 - \mu_{\chi_N^2}][\chi_N^2 - \mu_{\chi_N^2}]\} \quad (8)$$

Using (2) and (6), and multiplying both sides by σ^4 gives

$$\sigma^4 \text{var}[\chi_N^2] = E\left\{\left[\sum_{i=1}^N x_i^2 - 2\mu \sum_{i=1}^N x_i + N\mu^2 - N\sigma^2\right]\left[\sum_{j=1}^N x_j^2 - 2\mu \sum_{j=1}^N x_j + N\mu^2 - N\sigma^2\right]\right\}$$

Replacing the expression on the left side with (3) and expanding yields

$$\begin{aligned} 2N\sigma^4 &= E\left\{\left(\sum_{i=1}^N x_i^2\right)^2 - 2\mu \sum_{i=1}^N x_i^2 \sum_{j=1}^N x_j + N\mu^2 \sum_{i=1}^N x_i^2 - N\sigma^2 \sum_{i=1}^N x_i^2 - 2\mu \sum_{i=1}^N x_i^2 \sum_{j=1}^N x_j \right. \\ &\quad + 4\mu^2 \left(\sum_{i=1}^N x_i\right)^2 - 2N\mu^3 \sum_{i=1}^N x_i + 2N\mu\sigma^2 \sum_{i=1}^N x_i + N\mu^2 \sum_{i=1}^N x_i^2 - 2N\mu^3 \sum_{i=1}^N x_i \\ &\quad \left. + N^2\mu^4 - N^2\mu^2\sigma^2 - N\sigma^2 \sum_{i=1}^N x_i^2 + 2N\mu\sigma^2 \sum_{i=1}^N x_i - N^2\mu^2\sigma^2 + N^2\sigma^4\right\} \\ &= E\left\{\left(\sum_{i=1}^N x_i^2\right)^2 - 4\mu \sum_{i=1}^N x_i^2 \sum_{j=1}^N x_j + 2N\mu^2 \sum_{i=1}^N x_i^2 - 2N\sigma^2 \sum_{i=1}^N x_i^2 - 4N\mu^3 \sum_{i=1}^N x_i \right. \\ &\quad \left. + 4N\mu\sigma^2 \sum_{i=1}^N x_i - 2N^2\mu^2\sigma^2 + N^2\mu^4 + N^2\sigma^4 + 4\mu^2 \left(\sum_{i=1}^N x_i\right)^2\right\} \end{aligned}$$

Using Equations (A1) - (A4) from the Appendix in the above equation results in

$$\begin{aligned}
 2N\sigma^4 &= E \left\{ \left(\sum_{i=1}^N x_i^2 \right)^2 \right\} - 4\mu(2N\mu\sigma^2 + N^2\mu\sigma^2 + N^2\mu^3) + 2N^2\mu^2(\sigma^2 + \mu^2) \\
 &\quad - 2N^2\sigma^2(\sigma^2 + \mu^2) - 4N^2\mu^4 + 4N^2\mu^2\sigma^2 - 2N^2\mu^2\sigma^2 + N^2\mu^4 \\
 &\quad + N^2\sigma^4 + 4\mu^2(N\sigma^2 + N^2\mu^2) \\
 &= E \left\{ \left(\sum_{i=1}^N x_i^2 \right)^2 \right\} - 8N\mu^2\sigma^2 - 4N^2\mu^2\sigma^2 - 4N^2\mu^4 + 2N^2\mu^2\sigma^2 + 2N^2\mu^4 \\
 &\quad - 2N^2\sigma^4 - 2N^2\mu^2\sigma^2 - 4N^2\mu^4 + 4N^2\mu^2\sigma^2 + N^2\mu^4 - 2N^2\mu^2\sigma^2 + N^2\sigma^4 \\
 &\quad + 4N\mu^2\sigma^2 + 4N^2\mu^4 \\
 &= E \left\{ \left(\sum_{i=1}^N x_i^2 \right)^2 \right\} - 4N\mu^2\sigma^2 - 2N^2\mu^2\sigma^2 - N^2\mu^4 - N^2\sigma^4
 \end{aligned}$$

Therefore,

$$E \left\{ \left(\sum_{i=1}^N x_i^2 \right)^2 \right\} = 2N\sigma^4 + 4N\mu^2\sigma^2 + 2N^2\mu^2\sigma^2 + N^2\mu^4 + N^2\sigma^4 \quad (9)$$

For a random variable y ,

$$\text{var}[y] = E\{y^2\} - (E\{y\})^2$$

therefore the variance of $\sum_{i=1}^N x_i^2$ is given by

$$\text{var} \left[\sum_{i=1}^N x_i^2 \right] = E \left\{ \left(\sum_{i=1}^N x_i^2 \right)^2 \right\} - \left(E \left\{ \sum_{i=1}^N x_i^2 \right\} \right)^2$$

Using Equations (9) and (7) results in

$$\begin{aligned}
 \text{var} \left[\sum_{i=1}^N x_i^2 \right] &= E \left\{ \left(\sum_{i=1}^N x_i^2 \right)^2 \right\} - N^2(\sigma^2 + \mu^2)^2 \\
 &= 2N\sigma^4 + 4N\mu^2\sigma^2 + 2N^2\mu^2\sigma^2 + N^2\mu^4 + N^2\sigma^4 - N^2\sigma^4 - 2N^2\mu^2\sigma^2 - N^2\mu^4
 \end{aligned}$$

$$= 2N\sigma^4 + 4N\mu^2\sigma^2 \quad (10)$$

3. Summary

If x is a Gaussian random process and has mean μ and variance σ^2 , then

$$E\left\{\sum_{i=1}^N x_i^2\right\} = N(\sigma^2 + \mu^2)$$

$$var\left[\sum_{i=1}^N x_i^2\right] = 2N\sigma^4 + 4N\mu^2\sigma^2$$

Note that when $\mu=0$ and $\sigma^2=1$ these equations reduce to (2) and (3).

Appendix: Expected values of various summations

$$1. \quad E \left\{ \sum_{i=1}^N x_i \right\} = N\mu \quad (A1)$$

$$2. \quad E \left\{ \sum_{i=1}^N x_i^2 \right\} = N(\sigma^2 + \mu^2) \quad (A2)$$

This is Equation (7) and was derived in the main text.

$$3. \quad E \left\{ \left(\sum_{i=1}^N x_i \right)^2 \right\} = N\sigma^2 + N^2\mu^2 \quad (A3)$$

Derivation

Let x_i be normally distributed random variables and define $\bar{x} = \sum_{i=1}^N x_i$, then \bar{x} is a normal random variable with mean $N\mu$ and variance $N\sigma^2$. The variance of \bar{x} may be written as

$$\text{var}[\bar{x}] = E\{\bar{x}^2\} - (E\{\bar{x}\})^2$$

so that

$$\begin{aligned} E\{\bar{x}^2\} &= E\left\{ \left(\sum_{i=1}^N x_i \right)^2 \right\} \\ &= \text{var}(\bar{x}) + (E\{\bar{x}\})^2 \\ &= N\sigma^2 + N^2\mu^2 \end{aligned}$$

$$4. \quad E \left\{ \sum_{i=1}^N x_i^2 \sum_{j=1}^N x_j \right\} = 2N\mu\sigma^2 + N^2\mu\sigma^2 + N^2\mu^3 \quad (A4)$$

Derivation

$$E \left\{ \sum_{i=1}^N x_i^2 \sum_{j=1}^N x_j \right\} = E \left\{ \sum_{i=1}^N \sum_{j=1}^N x_i^2 x_j \right\} = \sum_{i=1}^N \sum_{j=1}^N E \{ x_i^2 x_j \}$$

When $i = j$ this becomes the single sum $\sum_{i=1}^N E\{x_i^3\}$. From [2, page 162],

$$E\{x_i^3\} = 3\mu\sigma^2 + \mu^3$$

so

$$\sum_{i=1}^N E\{x_i^3\} = N(3\mu\sigma^2 + \mu^3) \quad i=j$$

When $i \neq j$

$$\sum_{i=1}^N \sum_{j=1}^N E\{x_i^2 x_j\} = \sum_{i=1}^N \sum_{j=1}^N E\{x_i^2\} E\{x_j\}$$

since the processes are independent. Using $E\{x_i\} = \mu$, and

$$E\{x_i^2\} = \text{var}\{x\} + \mu^2 = \sigma^2 + \mu^2$$

results in

$$\sum_{i=1}^N \sum_{j=1}^N E\{x_i^2\} E\{x_j\} = \sum_{i=1}^N \sum_{j=1}^N \mu(\sigma^2 + \mu^2)$$

These are summed over all i and j *except* for the case $i = j$, so there are $N(N-1)$ of them, giving

$$\sum_{i=1}^N \sum_{j=1}^N E\{x_i^2\} E\{x_j\} = N(N-1)\mu(\sigma^2 + \mu^2) \quad i \neq j$$

Adding the cases for $i = j$ and $i \neq j$ together gives

$$\begin{aligned} E\left\{\sum_{i=1}^N x_i^2 \sum_{j=1}^N x_j\right\} &= N(3\mu\sigma^2 + \mu^3) + N(N-1)\mu(\sigma^2 + \mu^2) \\ &= 2N\mu\sigma^2 + N^2\mu\sigma^2 + N^2\mu^3 \end{aligned}$$

References

- [1] Bendat, J.S., Piersol, A.G., *Random Data: Analysis and Measurement Procedures*, Wiley, New York (1971).
- [2] Papoulis, A., *Probability, Random Variables, and Stochastic Processes*, McGraw-Hill, New York (1965).
- [3] Patnaik, P.B., "The non-central χ^2 and F-distributions and their applications", *Biometrika* 36, 202-232 (1949).
- [4] Whalen, A.D., *Detection of Signals in Noise*, Academic Press, New York (1971).

Acknowledgements

This work was supported by the Office of Naval Research under Contract N00014-80-C-0220.