

AD-A218 292

## Differential Association of T-2 and T-2 Tetraol with Mammalian Cells

DISTRIBUTION STATEMENT A

Approved for public release  
Distribution UnlimitedDTIC  
ELECTE  
FEB 23 1990  
S D

JOHN L. MIDDLEBROOK and DENNIS L. LEATHERMAN

Department of Toxinology, Pathology Division, U.S. Army Medical Research Institute of Infectious Diseases, Frederick, Maryland

Accepted for publication May 22, 1989

## ABSTRACT

The interactions of T-2 and its metabolite T-2 tetraol (hereafter tetraol) with CHO (Chinese hamster ovary cells) and CHO ribosomes were studied. T-2 was about 300-fold more potent at inhibiting protein synthesis in CHO than was tetraol. Association of T-2 with CHO was highly specific and achieved a maximum at a concentration producing complete inhibition of protein synthesis. Association of tetraol with CHO was of low specificity, but the specific fraction did correlate with the dose-response curve for protein synthesis inhibition. Binding of both T-2 and tetraol to isolated CHO ribosomes was quantitatively similar and highly specific. With isolated ribosomes, each toxin competed effectively for the binding of the other. Using intact cells, tetraol competed for T-2 cell association, but not the converse. The kinetics at physiological temperature for total and specific T-2

cell association were much more rapid than those for tetraol. Furthermore, the rate of tetraol-cell association was indistinguishable from the rate for cellular uptake of tritiated water. At 0°C, there was a substantial association of T-2 with cells, whereas none was observed with tetraol. The kinetics of dissociation of both toxins from CHO were similar. We conclude that T-2 rapidly crosses the cell membrane of cells and binds to the intracellular target, the ribosomes. In contrast, tetraol is taken up by the cell much more slowly, and many more toxin molecules are found in the cell than there are ribosomes. It would appear that the main physical property of the toxins that brings about these results is the relative hydrophobicities of the molecules. Thus, we suggest that one factor determining the potency of a trichothecene toxin may be its lipophilicity.

Trichothecenes are a group of structurally related sesquiterpenoids produced by several species of fungi. Trichothecenes are demonstrably lethal to many animal species and have been implicated as the causative agents in human disease and death. Although the pathogenesis of toxicity in animals is complex and not well defined, it is clear that trichothecenes are potent inhibitors of eukaryotic protein synthesis (McLaughlin *et al.*, 1977).

Work from a number of laboratories has shown that trichothecenes block protein synthesis by binding to the 60S subunit of the eukaryotic ribosome (Barbacid and Vasquez, 1974; Wei *et al.*, 1974). Although these toxins apparently bind to a common site (Wei *et al.*, 1974; Cannon *et al.*, 1976) mechanistically, some appear to inhibit initiation, whereas others block elongation or termination. Generally, the initiation-inhibiting toxins are more potent than the elongation/termination inhibitors. The basis for these potency differences is not clear and could be due to differences in such events as entry into the cell, binding to ribosomes, metabolism, etc.

We recently reported a detailed study of the interaction of a trichothecene initiation inhibitor, T-2, with the mammalian

cell line, CHO, and isolated CHO-derived ribosomes (Middlebrook and Leatherman, 1989a,b). Our data suggested that T-2 crosses the cell membrane freely and that almost all cell-associated toxin is bound intracellularly to ribosomes. Although incubation of CHO with T-2 at physiological temperature produced a steady-state level of toxin cell binding, the steady state is apparently due to a balanced, but continuous, uptake and release process. Metabolism of the toxin, although rapid and extensive in animals (Pace *et al.*, 1985), was not a significant event over the course of our binding studies (Trusal, 1986). The small amount of metabolism which did occur lead to the same metabolites seen in animals, namely, HT-2 and tetraol.

Because the end metabolite, tetraol, may be quite important in animal (and presumably human) intoxications, we studied the binding and toxicity of that toxin to CHO and CHO ribosomes. On a molar basis, tetraol was much less toxic to the cells than T-2. In contrast, binding of T-2 and tetraol to ribosomes was essentially the same. This and other data suggest that entry into target cells may be a major factor determining potency (toxicity) of trichothecene toxins for cells. (Reprints sent)

## Methods

**Cells and cell culture.** Seed stock for the CHO line (K1 subclone) was obtained from the American Type Culture Collection (ATCC no.

Received for publication January 17, 1989.

**ABBREVIATIONS:** CHO, Chinese hamster ovary cells; tetraol, T-2 tetraol; HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; HBSS, Hanks' balanced salt solution.

CCI-61). Cells were maintained in 75-cm<sup>2</sup> T-flasks (Costar no. 3075) with Earle's minimal essential medium, 10% fetal bovine serum and 50  $\mu$ g/ml of gentamycin.

**Media and sera.** All media, vitamins, antibiotics and amino acids were obtained from Grand Island Biological Co. (Grand Island, NY). Fetal calf serum was obtained from Armour Pharmaceutical (Kankakee, IL).

**Toxins.** T-2 and tetraol were purchased from Calbiochem (La Jolla, CA); the other trichothecene toxins were obtained from Sigma Chemical Co. (St. Louis, MO). T-2 was tritium-labeled by New England Nuclear (Boston, MA) or Amersham/Searle (Arlington Heights, IL) using a previously published procedure (Wallace *et al.*, 1977). The toxin preparations had specific activities from 9.0 to 14.0 Ci/mmol and were equipotent to unlabeled T-2 in a protein synthesis inhibition assay. Radiolabeled tetraol was prepared from [<sup>3</sup>H]T-2 by a published procedure (Wei and Chu, 1985) and was the generous gift of Dr. R. Wanne-macher, Jr. (U.S. Army Medical Research Institute of Infectious Diseases, Frederick, MD). In many cases, it was necessary to cut the radiolabeled tetraol to achieve desired molar concentrations so the specific activity varies. All toxins were dissolved in methanol and diluted so that the maximal alcohol concentration exposure to cells was 0.1%. This concentration of alcohol had no detectable effect on toxin-cell association.

**Toxin-cell association assay.** Cells were seeded in 24-well tissue culture plates. On the day of experimentation ( $1-4 \times 10^5$  cells/well), the growth medium was replaced with 0.5 ml of Hanks' 199 supplemented with 10% fetal calf serum, 50  $\mu$ g/ml of gentamycin and 25 mM HEPES, pH 7.4 (complete H-199). Further incubations and manipulations at 37°C were carried out on top of a microscope slide warmer in a warmbox (both equilibrated to 37°C). This arrangement allowed much better temperature control than did a standard CO<sub>2</sub> incubator. Radiolabeled T-2 was added to the cells in 50- $\mu$ l volumes and incubation carried out under the conditions and for the times stipulated. To determine cell-associated toxin, cells were rinsed three times with HBSS and solubilized in 1.0 ml of 0.1 M NaOH. A 0.5-ml aliquot was added to a scintillation vial with 0.1 ml of 1.0 M HCl and 5.0 ml of Aquasol 2 (New England Nuclear). The sample was then counted in a Beckman 5801 liquid scintillation spectrophotometer.

**Protein synthesis assay.** Cells in complete H-199 were incubated with toxin for the times and under the conditions indicated. Protein synthesis was measured by the addition of 1  $\mu$ Ci/well of [<sup>3</sup>H]leucine (New England Nuclear, 110-150 Ci/mmol) and incubation at 37°C, usually for 30 min. The pulse was terminated by rinsing the cells twice with HBSS and adding 0.10 ml of 0.1 M NaOH. After 5 to 10 min at 37°C, a prenumbered 11-mm disc (Schleicher and Schuell, 740E) was added to each well to absorb the solubilized cells. Each disc was then transferred to a bottle of 10% trichloroacetic acid and the samples were processed in mass as follows: two rinses with 5% trichloroacetic acid, two rinses with 50:50 ethanol:acetone and one rinse with acetone. After drying, each disc was assayed for radioactivity in 2.0 ml of Liqua-fluor:toluene (New England Nuclear).

**Purification of ribosomes.** Ribosomes from CHO were purified as described by Gupta and Siminovitch (1976).

**Toxin-ribosome binding assay.** A previously described filter assay was used (Middlebrook and Leatherman, 1989b). Usually, 10 to 20  $\mu$ l of toxin(s) were added to 100  $\mu$ l of a 10 OD<sub>260</sub>/ml ribosomal suspension in buffer D (20 mM HEPES, pH 7.5; 120 mM KCl; 1.5 mM Mg acetate; 6 mM 2-mercaptoethanol) and incubation was carried out under the conditions stipulated. Binding was terminated by addition of the entire sample to a Whatman GF/F glass fiber filter, followed by four 4-ml washes. Each filter was then counted with Aquasol 2 in a liquid scintillation spectrophotometer.


**Sucrose density gradient analysis of polyribosome profiles.** The distribution of labeled, cell-associated toxin was analyzed in polyribosome profiles using a modification of the procedure described by Cundliffe *et al.* (1974). CHO in T-75 culture flasks were incubated to equilibrium at 37°C with site-saturating concentrations of labeled toxins (0.1  $\mu$ g/ml of T-2 toxin or 10  $\mu$ g/ml of tetraol). After washing

four times in ice-cold HBSS, the cells were scraped into TMNa buffer (10 mM Tris-Cl, pH 7.4; 15 mM MgCl<sub>2</sub>; 140 mM NaCl) and pelleted at 1000  $\times g$  for 10 min at 2°C. Cell pellets were lysed in TMNa buffer (on ice) containing 0.5% Nonidet P-40 (Sigma) and 200  $\mu$ l of lysate was layered onto 4.5-ml, 10 to 30% continuous sucrose gradients prepared in TMNa buffer. The gradients were centrifuged at 42,000 rpm in an SW 50.1 rotor (Beckman) for 2.25 hr at 2°C. Under these conditions, the 40S and 60S subunits, and the 80S free ribosomes were simultaneously resolved and most of the polyribosomal material was pelleted. After centrifugation, the gradients were fractionated from the top with a Buchler Auto-Densi-Flow II gradient collector (Searle) pumping at a rate of 0.6 ml/min with a chart speed of 30 mm/min. Absorbance at 254 nm was monitored continuously through an LKB 2238 Uvicord S II UV spectrophotometer. The polyribosome pellet was resuspended in TMNa buffer and a portion of this pellet was used to determine polysome absorbance. The remainder of the pellet and the gradient fractions were counted in a liquid scintillation counter using Aquasol 2 to determine polyribosome-bound toxin.


## Results

Comparative dose/response curves for the inhibition of CHO protein synthesis by T-2 and tetraol are shown in figure 1. After 3 hr incubation with cells, protein synthesis was inhibited 50% by 0.003  $\mu$ g/ml (6.2 nM) T-2. Tetraol also inhibited protein synthesis in CHO, but the curve was shifted to the right almost three orders of magnitude from that for T-2. It required 1  $\mu$ g/ml (3.4  $\mu$ M) to obtain a 50% inhibitory effect. Other than their positions, the two curves were similar. Both toxins induced a complete inhibition of protein synthesis and therefore appear to be full agonists with different potencies.

Binding isotherms for T-2 and tetraol at physiological temperature are depicted in figure 2. We observed binding of T-2 to CHO at much lower concentrations than for tetraol. In agreement with our previous work (Middlebrook and Leatherman, 1989a), we obtained an S-shaped curve with the midpoint at about 0.008  $\mu$ g/ml (17 nM). Saturation was seen at a con-



**Fig. 1.** Correlation of specific toxin-cell association with toxin-induced inhibition of protein synthesis in CHO. Inhibition of protein synthesis: CHO were incubated with the indicated concentrations of T-2 toxin ( $\Delta$ ) or tetraol ( $\bullet$ ) at 37°C. After 3 hr, 1.0  $\mu$ Ci of [<sup>3</sup>H]leucine was added to each well and incubation was continued for an additional 30 min at 37°C. The assay was terminated by washing with ice-cold HBSS and the cells were processed to determine incorporated leucine as described under "Methods." Toxin-cell association: these data are replotted using the data from figure 2 for the specific association of T-2 ( $\circ$ ) and tetraol ( $\bullet$ ) with CHO. All data points are the means of triplicate determinations.



**Fig. 2.** Total, specific and nonspecific association of T-2 toxin and tetraol with CHO. A: CHO were incubated at 37°C with the indicated concentrations of [<sup>3</sup>H]T-2 toxin with (□) and without (Δ) a 75-fold molar excess of unlabeled T-2 (triplicates). After 3 hr of incubation at 37°C, cells were processed to measure cell-associated toxin as described under "Methods." Toxin bound in the presence of excess competitor was subtracted from toxin bound in the absence of excess competitor to determine specifically bound toxin (○). B: CHO were incubated at 37°C for 3 hr with the indicated concentrations of [<sup>3</sup>H]tetraol with (■) and without (▲) a 75-fold molar excess of nonlabeled tetraol (triplicates). Cells were then processed for cell-associated toxin and specifically bound toxin (●) was determined as described above. Total and nonspecific counts are expressed as means  $\pm$  S.E.M. If not shown, S.E.M. was smaller than the symbol.

centration close to that producing complete inhibition of protein synthesis (fig. 1). Moreover, throughout the concentration range studied, inclusion of a 75-fold excess of unlabeled T-2 blocked almost all the cell association of the radiolabeled T-2, indicating specificity of the binding (fig. 2A).


Much higher concentrations of tetraol were required to observe toxin cell association. Tetraol total association did not appear to plateau at concentrations examined in these studies and reached much greater levels than did T-2. Unlike the isotherm for T-2, the inclusion of excess, unlabeled tetraol blocked only a small fraction of the total binding suggesting that, at 37°C, most of the tetraol-cell association was nonspe-

cific (fig. 2B). Although it was low, we calculated the level of specific tetraol-CHO association and compared the data with specific T-2-CHO association (fig. 1). When the resulting curves were plotted with the toxins' inhibition of protein synthesis, the specific binding and the pharmacological effects correlated closely (fig. 1).


Because the protein synthesis inhibitory effects of T-2 are due to toxin-ribosome binding (Barbacid and Vasquez, 1974; Wei *et al.*, 1974) we compared the binding of T-2 and tetraol to CHO ribosomes (fig. 3). Surprisingly, the binding of the two toxins was not substantially different. At a given concentration of toxin, there was a somewhat higher level of radiolabeled T-2 than tetraol association with CHO ribosomes. However, the ratio of T-2:tetraol ribosome binding decreased as the concentrations increased and an analysis of the data by the method of Scatchard (1949) indicated a slightly lower affinity of tetraol for the ribosome with the same apparent number of sites (data not shown). In marked contrast to the association of tetraol with cells, excess unlabeled tetraol competed completely for the binding of radiolabeled tetraol to ribosomes (fig. 3), demonstrating a specific interaction.

As another measure of toxin-ribosome binding, we examined the ability of T-2 and tetraol to compete with one another. Competition by the homologous and heterologous toxins was essentially the same, suggesting that the ribosomal binding affinities of the two toxins are close (data not shown). These results probably indicate that T-2 and tetraol bind to the same site on the ribosome, although allosteric effects of one toxin on binding of the other cannot be ruled out.

Cross competition experiments performed with intact CHO produced quite different results (fig. 4). In each case, we used radiolabeled toxin concentrations which were just saturating




**Fig. 3.** Binding of T-2 toxin and tetraol to CHO-derived ribosomes. Ribosome suspensions at  $10 \text{ OD}_{260}/\text{ml}$  were incubated at 4°C with the indicated concentrations of [<sup>3</sup>H]T-2 toxin with (■) or without (□) a 100-fold molar excess of unlabeled T-2 toxin and [<sup>3</sup>H]tetraol with (●) or without (○) a 100-fold molar excess of unlabeled tetraol. After 18 hr of incubation at 4°C ribosomes were processed for ribosome-associated toxin by the filtration method described under "Methods." Specifically bound T-2 (Δ) and specifically bound tetraol (▲) were determined as described in the legend to figure 2. Data points are the averages of duplicate determinations.



**Fig. 4.** T-2 and tetraol cross competition in CHO. A: T-2 toxin and tetraol competition for [<sup>3</sup>H]T-2 association with CHO; CHO were incubated at 37°C with 0.1 μg/ml of [<sup>3</sup>H]T-2 toxin and the indicated concentrations of unlabeled T-2 toxin (○) or unlabeled tetraol (●). After 3 hr, the cells were processed for cell-associated toxin as described under "Methods." B: T-2 toxin and tetraol competition for [<sup>3</sup>H]tetraol association with CHO; CHO were incubated at 37°C with 10 μg/ml of [<sup>3</sup>H]tetraol and the indicated concentrations of unlabeled tetraol (▲) or unlabeled T-2 toxin (△). After 3 hr of incubation with toxins, the cells were processed for cell-associated toxin as described under "Methods." Data points represent the means of triplicate determinations with S.E.M.

(specific binding), as judged by the data in figures 1 and 2. With radiolabeled T-2, we observed competition by both T-2 and tetraol which was essentially complete (fig. 4A). The shapes of the competition curves were similar and it appeared that T-2 was more effective at competition. As judged by those concentrations producing 50% competition, T-2 was 180-fold more potent than tetraol at blocking radiolabeled T-2 cell binding. In contrast, neither T-2 or tetraol competed effectively for CHO association of radiolabeled tetraol (fig. 4B). In both instances, the toxins produced a maximum of about 15% competition, which was not increased by concentrations of competitors producing 90% block of T-2 cell binding.

The kinetics of CHO association of radiolabeled T-2 and tetraol were compared with the cellular uptake of tritiated water (fig. 5). All three radiolabeled compounds were added to achieve the same final specific activity in the medium. It is evident that the association of T-2 with cells was much more rapid than



**Fig. 5.** Comparative kinetics for the association of T-2 toxin, tetraol, and H<sub>2</sub>O with CHO. CHO were incubated at 37°C with 0.35 μg/ml of [<sup>3</sup>H]tetraol (●), 0.35 μg/ml of [<sup>3</sup>H]T-2 toxin (▲) or [<sup>3</sup>H]H<sub>2</sub>O (□), each at 10 μCi/ml. After the indicated incubation times at 37°C, the cells were processed for cell-associated radioactivity as described under "Methods." Data points are the averages of duplicate determinations.

that of either tetraol or water. The zero time point with T-2, which consisted of adding toxin to the medium and immediately washing the cells, produced more specifically cell-bound counts than a 90-min incubation with a like concentration of tetraol. A maximal, steady-state binding of T-2 was achieved in approximately 10 min, and was maintained throughout the course of the experiment (fig. 5). On the other hand, tetraol-cell binding reached a steady state in about 1 hr, but it was only 2.2% of the T-2 binding plateau. Furthermore, the cell binding kinetic curve for tetraol was not significantly different from that obtained with tritiated water, which we used as a measure of fluid phase pinocytosis.

If association of T-2 and tetraol with CHO was studied when toxins were added at concentrations which produce 90 to 95% inhibition of protein synthesis, additional and important information was brought to light. Total and specific binding of T-2 to CHO at physiological temperature rapidly attained plateau levels which were quantitatively close to one another, 3.0 vs. 2.5 million toxin molecules/cell (fig. 6). If association was carried out at either 4° or 0°C, a longer time was required to reach the apparent plateau for both total and specific binding (fig. 6). Furthermore, the plateau level was about half that attained at 37°C (fig. 6), although extending the incubation at 4°C (but not 0°) to 24 hr resulted in an increase in the binding to 2.5 million molecules bound per cell (data not shown).

Association of pharmacologically equipotent (much higher molar) concentrations of tetraol with CHO is demonstrated by the data in figure 7. Several important differences were observed when compared to the data for T-2. First, many more molecules of tetraol bound to (or were taken up by) CHO than with T-2, viz., 20 vs. 3 million (compare fig. 7A and 6A). Second, at physiological temperature, it required 2 to 3 hr to reach maximal total and specific binding of tetraol compared to 0.5 hr for T-2. In addition, the total association of tetraol with cells was much higher than the specific association, a pattern opposite that for T-2. Third, the total association of tetraol with cells at 0° or 4°C compared to 37°C was much lower proportionately than the ratios seen with T-2. That is, with T-






Fig. 6. Effect of temperature on the total and specific association of T-2 toxin with CHO. A: Total association: [ $^3\text{H}$ ]T-2 toxin (0.1  $\mu\text{g}/\text{ml}$ ) was incubated with CHO for the indicated times at 0°C (□), 4°C (○) or 37°C (Δ). The cells were then processed for cell-associated toxin as described under "Methods." B: Specific association: the specific association of T-2 toxin with CHO at 0°C (■), 4°C (●) and 37°C (▲) has been determined from the data in A by subtraction of the nonspecific binding measured in the presence of a 75-fold molar excess of unlabeled toxin. Data points for total association are the means of triplicate determinations with S.E.M.

2 at both lower temperatures, there was about 50% of the total association seen at 37°C (fig. 6A). In contrast, with tetraol there was only 6 to 9% (4°C) or 2 to 3% (0°C) of the association at physiological temperature (fig. 7A). This temperature differential with T-2 and tetraol was even more pronounced when the specific association was considered. Once again, with T-2, the specific association at either 0° or 4°C (4–8 hr) was approximately 50% of the specific association at 37°C (fig. 6B). Statistically, there was no specific association of tetraol with CHO at 0°C, whereas we did observe specific association at 4°C, approximately 60 to 70% that at 37°C (fig. 7B).

The dissociation kinetics for cell-associated toxins are shown in figure 8. Cells were preincubated with concentrations of each toxin that produced similar pharmacological effects, i.e., 0.1  $\mu\text{g}/\text{ml}$  of T-2 and 10  $\mu\text{g}/\text{ml}$  of tetraol. Cells were then washed and




Fig. 7. Effect of temperature on the total and specific association of tetraol with CHO. A: Total association: [ $^3\text{H}$ ]tetraol (10  $\mu\text{g}/\text{ml}$ ) was incubated with CHO for the indicated times at 0°C (□), 4°C (○) or 37°C (Δ). The cells were then processed for cell-associated toxin as described under "Methods." B: Specific association: the specific association of tetraol with CHO at 0°C (■), 4°C (●) or 37°C (▲) has been determined from the data in A by subtraction of the nonspecific binding measured in the presence of a 75-fold molar excess of unlabeled toxin. Data points for total association are the means of triplicate determinations with S.E.M.

cell-associated radioactivity measured as a function of time. Clearly, the rates of T-2- and tetraol-CHO dissociation were similar (fig. 8). When analyzed by a semilogarithmic plot (fig. 8, inset), both curves in figure 8 fit straight lines (correlation coefficients of 0.93 and 0.98 for T-2 and tetraol, respectively), whose equations gave calculated half-times of 3.3 and 3.7 hr for T-2 and tetraol, respectively.

In an attempt to compare the subcellular locations of T-2 and tetraol, we bound cells with radiolabeled toxins, washed and lysed the cells and centrifuged the samples on sucrose gradients. The profiles of radioactivity seen with T-2-bound cells coincided with the absorbance peaks defining the location of ribosomes and polysomes (fig. 9A). The ribosome-associated toxin (87,000  $\pm$  2,800 dpm) represented 97% of the total radiolabeled toxin applied to the gradient (91,000  $\pm$  2,900 dpm) (value derived from duplicate gradients). In contrast, the radio-




Fig. 8. Dissociation of T-2 toxin and tetraol from CHO. CHO were incubated with 0.1  $\mu\text{g/ml}$  of [ $^3\text{H}$ ]T-2 toxin (solid symbols) or 10  $\mu\text{g/ml}$  of [ $^3\text{H}$ ]tetraol (open symbols) at 37°C. After 3 hr of incubation with toxins, the cells were washed three times with HBSS and fresh, toxin-free HBSS containing 10% fetal bovine serum was added to the cells. The incubation was continued at 37°C and, at the indicated times, the medium (●, ○) was removed from the monolayers and transferred to scintillation vials, adjusted to 0.1 N NaOH and counted in 5.0 ml of Aqualus 2. The cell monolayers (■, □) were solubilized and processed for cell-associated toxin as described under "Methods." All data points are the means of triplicate determinations with S.E.M. Inset: The data from figure 8 were analyzed by a semilogarithmic plot. The elimination half-time for T-2 (●) is 3.3 hr with coefficient of correlation ( $r$ ) = 0.93. The elimination half-time for tetraol (○) is 3.7 hr with coefficient of correlation ( $r$ ) = 0.98.




Fig. 9. Distribution of T-2 toxin and tetraol in CHO. A: T-2 toxin: CHO were incubated with 0.1  $\mu\text{g/ml}$  of [ $^3\text{H}$ ]T-2 toxin at 37°C. After 30 min, the cells were washed and processed for polyribosome profiles as described under "Methods." The solid and dashed lines denote absorbance and radioactivity, respectively. The arrows on both plots denote the position of the 80S free ribosome peak and the arrows on the abscissas indicate the direction of sedimentation. B: Tetraol: CHO were incubated with 10  $\mu\text{g/ml}$  of [ $^3\text{H}$ ]tetraol at 37°C. After 3.5 hr, the cells were washed and processed as described above. See text for further information.

activity profile for tetraol-bound cells (total  $15,000 \pm 160$  dpm) showed that most of the tetraol was on the top of the gradient with only a small fraction ( $2600 \pm 250$  dpm or 17%) in the ribosome/polysome region (fig. 9B). Thus it appears that virtually all the T-2 associated with cells was ribosome-bound, whereas a majority of cell-associated tetraol was free in the cytoplasm or bound to a very small organelle, vesicle or protein.

As a general rule, metabolizing systems of animals produce metabolites that are more polar than the parent compound. As a simple measure of the polarities of T-2 and tetraol, we measured their distribution coefficients between an aqueous buffer and benzene. The results indicated that T-2 and tetraol have diametrically opposed distribution coefficients. Although T-2 showed a preference for a nonpolar solvent by a ratio of

54:1, tetraol had a similar ratio of 1:270. If the distribution coefficients for that solvent system can be used as an estimate of lipophilicity, T-2 should be 14,000 times more lipophilic.

## Discussion

Trichothecene toxins are among the most potent nonprotein inhibitors of protein synthesis in eukaryotic cell systems. A good deal of evidence points toward toxin binding to ribosomes as the molecular event which brings about this inhibition. There is a large range of inhibitory potencies in the trichothecene family of toxins. Delineating the bases for this variation in potencies should provide insights into the mechanism of action of these toxins with cells and possibly animals.

There was almost three orders of magnitude difference between T-2 and its metabolite, tetraol, at inducing protein synthesis inhibition in CHO (fig. 1). The molecular basis for this potency difference does not appear to exist at the level of toxin-ribosome binding because: 1) there was only a 2- to 5-fold difference in cell-free ribosomal binding of the two toxins (fig. 3) and 2) competition experiments strongly suggest that the two toxins bind to a common ribosomal site (data not shown). Therefore, we focused on interactions with the cell as the probable cause for the large potency difference.

The data presented in figures 1, 4, 6 and 7 established five important features of tetraol-cell vs. T-2-cell association. First, there was no appreciable binding of tetraol to CHO at concentrations of toxin that produced maximal cell binding with T-2 (fig. 1). Second, whatever the extracellular (exposure) concentration of toxin, similar levels of specifically bound T-2 or tetraol produced the same pharmacological response (fig. 1). Third, both T-2 and tetraol competed completely for radiolabeled T-2 binding to cells, the latter being about 200-fold less effective (fig. 4A), whereas neither T-2 or tetraol competed for more than about 15% of radiolabeled tetraol cell binding (with a reversal in competitive potencies) (fig. 4B). Fourth, at maximally effective pharmacological concentrations, the kinetics of total T-2 cell binding quickly reached a plateau (fig. 6A), whereas the kinetics for total tetraol cell binding were much slower (relatively speaking) and did not appear to plateau (fig. 7A). Finally, the kinetics of release or out transport of T-2 and tetraol were, within experimental error, identical.

Based on the above features, we propose the following explanation or model to account for the differences in toxicity between T-2 and tetraol: T-2 diffuses across the cell membrane readily and, once inside the cell, binds rapidly to ribosomes. Thus, cell-associated T-2 is really a measure of ribosomally bound T-2 inside a limiting sack, e.g., the plasma membrane. This model is supported by our previous studies with T-2 and cells (Middlebrook and Leatherman, 1989a,b) and by the data in figure 9 showing the ribosome profiles. Tetraol, on the other hand, is unable to cross the cell membrane by diffusion, possibly because of its hydrophilic nature. Thus, tetraol is taken inside the cell by a much slower, nonsaturable process similar, if not identical, to fluid-phase pinocytosis. Once inside the cell, tetraol escapes the pinocytotic vesicles or lysosomes, binds to ribosomes and brings about inhibition of protein synthesis.

With such a model in mind, one can make certain predictions which appear to be borne out by our data. First, one would expect the rate of tetraol uptake to be the same as the uptake rate of other pinocytotically accumulated substances such as water or inulin. Indeed, as the data in figure 5 show, the kinetics of water and tetraol uptake were essentially the same. Second, the rate of uptake for tetraol should be much more temperature-dependent than that for T-2. Both the total uptake and specific binding of T-2 at 0° and 4°C were about 50% of those seen at physiological temperature (fig. 6). In contrast, the total uptake of tetraol at the lower temperatures was 10 to 15% of the 37°C uptake, whereas the specific uptake at 0° was statistically zero (fig. 7). There was a substantial specific uptake of tetraol at 4°C compared to physiological temperature, although it seemed to take much longer to attain that level (fig. 7). Finally, fluid-phase pinocytosis is a nonsaturable process so one would anticipate that the uptake of tetraol would be much more nonspecific

than that for T-2, a prediction reflected by the data in figures 2, 4, 6 and 7.

It is clear that trichothecene toxins can differ markedly in their affinities for ribosomes (Wei *et al.*, 1974) and that factor alone can account for some differences in their potencies, even if cellular uptake were similar. Sato and Ueno (1977) presented data which suggested that another factor determining toxicity for cells is the lipophilicity of individual trichothecenes. Without radiolabeled toxins, those workers were unable to define the molecular events underlying the observation that lipophilicity and toxicity are positively related. Although carried out with only two toxins, our experiments may indicate that the basis of the lipophilicity-toxicity correlation observed by Sato and Ueno (1977) is toxin entry into cells. It may be that differential uptake by target cells is an important, toxicity-determining property of other trichothecene toxins, but experiments with more trichothecenes would be required to state this as a general rule. However, it is probable that either ribosome affinity or lipophilicity could be the major potency determining factor for any given trichothecene or, in certain cases, both properties could contribute. Furthermore, the relationship between toxin-induced protein synthesis in cultured cells and toxin-induced lethality in animals is not clearly evident, so extrapolation to animal toxicity is not appropriate at this time.

#### References

- BARBACID, M. AND VASQUEZ, D.: Binding of [acetyl-<sup>14</sup>C]trichodermin to the peptidyl transferase centre of eukaryotic ribosomes. *Eur. J. Biochem.* **44**: 437-444, 1974.
- CANNON, M., JIMENEZ, A. AND VASQUEZ, D.: Competition between trichodermin and several other sesquiterpene antibiotics for binding to their receptor site(s) on eukaryotic ribosomes. *Biochem. J.* **160**: 137-145, 1976.
- CUNDLIFFE, E., CANNON, M. AND DAVIES, J.: Mechanism of inhibition of eukaryotic protein synthesis by trichothecene fungal toxins. *Proc. Natl. Acad. Sci. USA* **71**: 30-34, 1974.
- GUPTA, R. S. AND SIMINOVITCH, L.: The isolation and preliminary characterization of somatic cell mutants resistant to the protein synthesis inhibitor—emetine. *Cell* **9**: 213-219, 1976.
- MCLAUGHLIN, C. S., VAUGHAN, M. H., CAMPBELL, I. M., WEI, C. M., STAFFORD, M. E. AND HANSEN, B. S.: Inhibition of protein synthesis by trichothecenes. In *Mycotoxins in Human and Animal Health*, ed. by J. V. Rodericks, C. W. Hesseltine and M. A. Mehlman, pp. 263-273, Pathotox Publishers, Inc., Park Forest South, IL, 1977.
- MIDDLEBROOK, J. L. AND LEATHERMAN, D. L.: Specific association of T-2 toxin with mammalian cells. *Biochem. Pharmacol.*, in press, 1989a.
- MIDDLEBROOK, J. L. AND LEATHERMAN, D. L.: Binding of T-2 toxin to eukaryotic cell ribosomes. *Biochem. Pharmacol.*, in press, 1989b.
- PACE, J. G., WATTS, M. R., BURROWS, E. P., DINTERMAN, R. E., MATSON, C., HAUER, E. C. AND WANNEMACHER, R. W., JR.: Fate and distribution of <sup>3</sup>H-labeled T-2 mycotoxin in guinea pigs. *Toxicol. Appl. Pharmacol.* **80**: 377-385, 1985.
- SATO, N. AND UENO, Y.: Comparative toxicities of trichothecenes. In *Mycotoxins in Human and Animal Health*, ed. by J. V. Rodericks, C. W. Hesseltine and M. A. Mehlman, pp. 295-307, Pathotox Publishers, Inc., Park Forest South, IL, 1977.
- SCATCHARD, G.: The attractions of proteins for small molecules and ions. *Ann. N.Y. Acad. Sci.* **51**: 660-672, 1949.
- TRUSAL, L. R.: Metabolism of T-2 mycotoxin by cultured cells. *Toxicon* **24**: 597-603, 1986.
- WALLACE, E. M., PATHRE, S. V., MIROCHA, C. J., ROBISON, T. S. AND FENTON, S. W.: Synthesis of radiolabeled T-2 toxin. *J. Agric. Food Chem.* **25**: 836-838, 1977.
- WEI, C. M., CAMPBELL, I. M., MCLAUGHLIN, C. S. AND VAUGHAN, M. H.: Binding of trichodermin to mammalian ribosomes and its inhibition by other 12,13-epoxytrichothecenes. *Mol. Cell. Biochem.* **3**: 215-219, 1974.
- WEI, R. D. AND CHU, F. S.: Modification of in vitro metabolism of T-2 toxin by esterase inhibitors. *Appl. Environ. Microbiol.* **50**: 115-119, 1985.

Send reprint requests to: Dr. John L. Middlebrook, Department of Toxinology, Pathology Division, U.S. Army Medical Research Institute of Infectious Diseases, Frederick, MD 21701-5011.