

DD FORM 1473, JUN 86

2

ITATION PAGE			Form Approved OMB No. 0704-0188	
1a. REPORT Unc.		1b. RESTRICTIVE MARKINGS		
AD-A222 568		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited		
		2b. DECLASS N/A		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 287		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 911 0010		
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina Center for Stochastic Processes		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State, and ZIP Code) Statistics Department CB #3260, Phillips Hall Chapel Hill, NC 27599-3260		7b. ADDRESS (City, State, and ZIP Code) Bldg. 410 Bolling Air Force Base, DC 20332-6448		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (if applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85C 0144	
8c. ADDRESS (City, State, and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448		10. SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. A5
11. TITLE (Include Security Classification) Dichotomies for certain product measures and stable processes				
12. PERSONAL AUTHOR(S) Marques, Mauro S. de F. and Cambanis, S-				
13a. TYPE OF REPORT preprint		13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1990, March	15. PAGE COUNT 27
16. SUPPLEMENTARY NOTATION N/A				
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP		
XXXXXX	XXXXXXXXXX	XXXXXX		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) (Please see over)				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Professor Eytan Barouch		22b. TELEPHONE (Include Area Code) (202) 767-5026	22c. OFFICE SYMBOL AFOSR/NM	

19. Abstract

Abstract: Necessary and sufficient conditions for equivalence or singularity of certain product measures are given and applied to the problem of distinguishing a sequence of random vectors from affine transformations of itself. In particular sequences of independent stable random variables are considered and the singularity of sequences with different indexes of stability is proved. Using these results the dichotomy, "two processes are either equivalent or singular", is established for certain classes of stable processes, such as independently scattered measures and harmonizable processes. Also sufficient conditions for singularity and necessary conditions for absolute continuity are given for p^{th} order processes.

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



DICHOTOMIES FOR CERTAIN PRODUCT MEASURES
AND STABLE PROCESSES



by
Mauro S. de F. Marques
and
Stamatis Cambanis

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Date	
Project	

A-1

Technical Report No. 287

March 1990

219. J. Bather, Stopping rules and ordered families of distributions, Dec. 87. *Sequential Anal.*, 7, 1988, 111-126.
220. S. Cambanis and M. Kasjima, Two classes of self-similar stable processes with stationary increments, Jan. 88. *Stochastic Proc. Appl.*, 32, 1989, 305-329.
221. H.P. Hucke, C. Kallianpur and R.L. Karandikar, Smoothness properties of the conditional expectation in finitely additive white noise filtering, Jan. 88. *J. Multivariate Anal.*, 27, 1988, 261-269.
222. I. Mitoma, Weak solution of the Langevin equation on a generalized functional space, Feb. 88. (Revised as No. 238).
223. L. de Haan, S.I. Resnick, H. Rootzén and C. de Vries, Extremal behaviour of solutions to a stochastic difference equation with applications to arch-processes, Feb. 88. *Stochastic Proc. Appl.*, 32, 1989, 213-224.
224. O. Kallenberg and J. Szulga, Multiple integration with respect to Poisson and Lévy processes, Feb. 88. *Prob. Theor. Rel. Fields*, 89, 1989, 101-134.
225. D.A. Dawson and L.C. Corostiza, Generalized solutions of a class of nuclear space valued stochastic evolution equations, Feb. 88. *Appl. Math. Optimization*, to appear.
226. G. Samorodnitsky and J. Szulga, An asymptotic evaluation of the tail of a multiple symmetric α -stable integral, Feb. 88. *Ann. Probab.*, 17, 1989, 1503-1520.
227. J.J. Hunter, The computation of stationary distributions of Markov chains through perturbations, Mar. 88.
228. H.C. Ho and T.C. Sun, Limiting distribution of nonlinear vector functions of stationary Gaussian processes, Mar. 88. *Ann. Probab.*, to appear.
229. R. Brigola, On functional estimates for ill-posed linear problems, Apr. 88.
230. M.R. Leadbetter and S. Nandagopalan, On exceedance point processes for stationary sequences under mild oscillation restrictions, Apr. 88. *Proc. Oberwolfach Conf. on Extremal Value Theory*, J. Hüsiér and R. Reiss, eds., Springer, 1989, 69-80.
231. S. Cambanis, J. P. Nolan and J. Rosinski, On the oscillation of infinitely divisible processes, Apr. 88. *Stochastic Proc. Appl.*, 1990, to appear.
232. G. Hardy, C. Kallianpur and S. Ramasubramanian, A nuclear space-valued stochastic differential equation driven by Poisson random measures, Apr. 88.
233. D.J. Daley, T. Rolski, Light traffic approximations in queues (II), May 88. *Math. Operat. Res.*, to appear.
234. G. Kallianpur, I. Mitoma, R.L. Wolpert, Nuclear space-valued diffusion equations, July 88. *Stochastics*, 1989, to appear.
235. S. Cambanis, Admissible translates of stable processes: A survey and some new models, July 88.
236. E. Platen, On a wide range exclusion process in random medium with local jump intensity, Aug. 88.
237. R.L. Smith, A counterexample concerning the extremal index, Aug. 88. *Adv. Appl. Probab.*, 20, 1989, 661-663.
238. G. Kallianpur and I. Mitoma, A Langevin-type stochastic differential equation on a space of generalized functionals, Aug. 88.
239. C. Hudré, Harmonizability, V -boundedness, $(2, P)$ -boundedness of stochastic processes, Aug. 88. *Prob. Th. Rel. Fields*, 84, 1990, 39-54.
240. G.W. Johnson and G. Kallianpur, Some remarks on Hu and Meyer's paper and infinite dimensional calculus on finitely additive canonical Hilbert space, Sept. 88. *Th. Prob. Appl.*, to appear.
241. L. de Haan, A Brownian bridge connected with extreme values, Sept. 88. *Sankhya*, 1989, to appear.
242. O. Kallenberg, Exchangeable random measures in the plane, Sept. 88. *J. Theor. Probab.*, 3, 1990, 81-136.
243. E. Herry and S. Cambanis, Trapezoidal Monte Carlo integration, Sept. 88. *SIAM J. Numer. Anal.*, 27, 1990, 225-246.
244. L. Pitt, On a problem of H.P. McKean, Sept. 88. *Ann. Probab.*, 17, 1989, 1651-1657.
245. C. Hudré, On the linear prediction of multivariate $(2, P)$ -bounded processes, Sept. 88.
246. C. Hudré, Stochastic processes as Fourier integrals and dilation of vector measures, Sept. 88. *Bull. Amer. Math. Soc.*, 21, 1989, 281-285.
247. J. Migebeur, On the rate of convergence in Strassen's functional law of the iterated logarithm, Sept. 88. *Probab. Theor. Rel. Fields*, to appear.
248. G. Kallianpur and V. Perez-Abreu, Weak convergence of solutions of stochastic evolution equations on nuclear spaces, Oct. 88. *Proc. Trento Conf. on Infinite Dimensional Stochastic Differential Equations*, 1989, to appear.
249. R.L. Smith, Bias and variance approximations for estimators of extreme quantiles, Nov. 88.
250. H.L. Haré, Spectral coherence of nonstationary and transient stochastic processes, Nov. 88. 4th Annual ASSP Workshop on Spectrum Estimation and Modeling, Minneapolis, 1988, 387-390.
251. J. Leadbe, Maximum likelihood estimator for almost periodic stochastic processes models, Dec. 88.
252. M.R. Leadbetter and T. Hsing, Limit theorems for strongly mixing stationary random measures, Jan. 89. *Stochastic Proc. Appl.*, to appear.
253. M.R. Leadbetter, I. Weissman, L. de Haan, H. Rootzén, On clustering of high values in statistically stationary series, Jan. 89. *Proc. 4th Int. Meeting on Stat. Climatol.*, 217-222.
254. J. Leadbe, Least squares estimation in almost periodic point processes models, Feb. 89.
255. M.N. Vaidyanla, Orthogonal random vectors and the Hurwitz-Radon-Eckmann theorem, Apr. 89.

(9)

DICHOTOMIES FOR CERTAIN PRODUCT MEASURES
AND STABLE PROCESSES

Mauro S. de F. Marques
Departamento de Estatística
Universidade Estadual de Campinas
13081 Campinas, SP, Brasil

and Stamatis Cambanis
Department of Statistics
University of North Carolina
Chapel Hill, NC 27599-3260 USA

Abstract: Necessary and sufficient conditions for equivalence or singularity of certain product measures are given and applied to the problem of distinguishing a sequence of random vectors from affine transformations of itself. In particular sequences of independent stable random variables are considered and the singularity of sequences with different indexes of stability is proved. Using these results the dichotomy, "two processes are either equivalent or singular", is established for certain classes of stable processes, such as independently scattered measures and harmonizable processes. Also sufficient conditions for singularity and necessary conditions for absolute continuity are given for p^{th} order processes.

sub th
(KR)

AMS 1980 subject classifications: Primary 60G30.

Keywords: Equivalence and Singularity, Product Measures, Stable Processes

Research supported by the Air Force Office of Scientific Research Contract No. F49620 85C 0144.

1. INTRODUCTION

This paper investigates the equivalence and singularity of measures induced by non-Gaussian stable processes.

For two Gaussian processes the following dichotomy prevails: they are either mutually absolutely continuous (equivalent) or else they are singular (see, e.g. [6]). For non-Gaussian stable processes some results are available in [13] [30], [29] and [24].

In Section 2 an idea of LeCam [23] is developed further to provide a necessary and sufficient condition for equivalence and for singularity of certain product measures (Proposition 2.1). As an application, the results on the discrimination between a sequence of random vectors and its perturbation by rigid motions in [27], are extended to more general classes of perturbations (Corollary 2.2). Also necessary and sufficient conditions are given for the equivalence and for the singularity of certain sequences of independent stable random variables (Corollaries 2.3, 2.4); and the singularity of two sequences of independent symmetric stable variables with different indexes of stability is proved (Proposition 2.5).

In Section 3 an equivalence-singularity dichotomy is shown for certain symmetric stable processes (Proposition 3.2), including independently scattered measures (Proposition 3.1) and harmonizable processes (Corollary 3.3), and necessary and sufficient conditions for the two alternatives are given, identical to those in the Gaussian case. The singularity of an invertible symmetric stable process to its multiples is also proved (Corollary 3.4).

In Section 4, a necessary condition for equivalence of two Gaussian processes, namely the setwise equality of their reproducing kernel Hilbert spaces (RKHS), is extended to symmetric stable processes with the function space of the process introduced in [24] replacing the RKHS (Proposition 4.2).

Further, for p^{th} order processes with $1 < p < 2$, necessary conditions for absolute continuity and sufficient conditions for singularity are presented (Proposition 4.3) analogous to those in [12] for second order processes.

The following setting is considered. $X_i = (X_i(t) = X_i(t, \omega); t \in T)$, $i = 1, 2$, are stochastic processes on a probability space (Ω, \mathcal{F}, P) with parameter set T and real or complex values, i.e. values in $F = \mathbb{R}$ or \mathbb{C} . When $X_i(t) \in L_p(\Omega, \mathcal{F}, P) = L_p(P)$ for all $t \in T$ and some $p > 0$, X_i is called a p^{th} order process and its linear space $\mathcal{L}(X_i)$ is the $L_p(P)$ completion of the set of finite linear combinations of its random variables $\mathcal{L}(X_i) \stackrel{\Delta}{=} \text{sp}\{X_i(t); t \in T\}$. F^T denotes the set of all extended F -valued (i.e., real or complex valued) functions on T , $\mathcal{G} = \mathcal{G}(\overline{F}^T)$ the σ -field generated by the cylinder sets of \overline{F}^T , and μ_i (or μ_{X_i}) the distribution of the process X_i i.e. the probability induced on \mathcal{G} by X_i : $\mu_i(C) = P(\{\omega; X_i(\cdot, \omega) \in C\})$, $C \in \mathcal{G}$. We are interested in the Lebesgue decomposition of μ_2 with respect to μ_1 , and in particular in conditions for μ_1 and μ_2 to be singular ($\mu_1 \perp \mu_2$), for μ_2 to be absolutely continuous with respect to μ_1 ($\mu_2 \ll \mu_1$), and for μ_1 and μ_2 to be mutually absolutely continuous or equivalent ($\mu_1 \sim \mu_2$).

2. ON THE EQUIVALENCE AND SINGULARITY OF CERTAIN PRODUCT MEASURES

In this section we consider the case where $X_i = (X_{i,n}; n \in \mathbb{N})$, $i=1,2$, are sequences of independent random variables, or equivalently μ_1 and μ_2 are product measures on $F^{\mathbb{N}}$. The equivalence-singularity dichotomy of product measures was characterized in [18] in terms of the Hellinger distance of the marginal measures, which may be difficult to compute, e.g. for stable measures. The case of translates of product measures with identical marginals was solved in [25] under finite Fisher information. The sufficient condition for equivalence in [25] was extended in [23] to a more general scenario under LeCam's " ℓ " condition. Proposition 2.1 derives a nearly complete extension of a result of

Shepp in [25] under a condition closely related to LeCam's. As an application the equivalence-singularity dichotomy is established for a sequence of i.i.d. random vectors and an affine transformation of itself in Corollary 2.2 (extending the results in [27] about rigid motions), and for sequences of i.i.d. stable random variables in Corollaries 2.3 and 2.4.

Before stating the main results we need to introduce some concepts for which we refer to [28].

2.1 Preliminaries

The normalized Hellinger distance d of two probability measures P and Q on a measurable space (Ω, \mathcal{F}) is defined by

$$d^2(P, Q) = \frac{1}{2} \int_{\Omega} |(dP/d\nu)^{1/2} - (dQ/d\nu)^{1/2}|^2 d\nu,$$

where ν is any σ -finite measure dominating $P+Q$, i.e. $P+Q \ll \nu$ (e.g. $\nu = P+Q$); and d does not depend on ν .

Kakutani's theorem [18] states that if $(\mu_n; n \in \mathbb{N})$ and $(\lambda_n; n \in \mathbb{N})$ are sequences of probability measures with $\mu_n \sim \lambda_n$ and $\mu = \times_{n=1}^{\infty} \mu_n$ and $\lambda = \times_{n=1}^{\infty} \lambda_n$ are their product measures, then

$$(2.1) \quad \mu \sim \lambda \quad \bullet \quad \sum_{n=1}^{\infty} d^2(\mu_n, \lambda_n) < \infty \quad \text{and} \quad \mu \perp \lambda \quad \bullet \quad \sum_{n=1}^{\infty} d^2(\mu_n, \lambda_n) = \infty.$$

We consider the following setting. $(\Omega, \mathcal{F}, \nu)$ is a σ -finite measure space, and $\{P_{\theta}; \theta \in \Theta\}$ a family of probability measures on (Ω, \mathcal{F}) with $P_{\theta} \ll \nu$ and Θ an open subset of \mathbb{R}^k . Then $F: \Theta \rightarrow L_2(\Omega, \mathcal{F}, \nu) = L_2(\nu)$ defined by $F(\theta) = 2 [dP_{\theta}/d\nu]^{1/2}$ is said to be differentiable at θ , if there exists a map $DF(\theta) := DF(\cdot, \theta) : \Omega \rightarrow \mathbb{R}^k$ such that

$$\|DF(\theta)\|_{L_2(\Omega, \mathcal{F}, \nu; \mathbb{R}^k)}^2 = \int_{\Omega} \|DF(\omega, \theta)\|^2 \nu(d\omega) < \infty,$$

i.e. $DF(\theta) \in L_2(\Omega, \mathcal{F}, \nu; \mathbb{R}^k)$, and

$$\int_{\Omega} |F(\theta+h) - F(\theta) - \langle DF(\theta), h \rangle|^2 d\nu = o(\|h\|^2) \quad \text{as } \|h\| \rightarrow 0.$$

As usual F is said to be differentiable (on θ) if it is differentiable at each $\theta \in \Theta$. The Fisher information matrix is defined by

$$\mathcal{J}(\theta) = \int_{\Omega} DF(\theta) DF(\theta)' d\nu$$

(where $DF(\theta)'$ is the transpose of the column vector $DF(\theta)$). It is nonnegative definite, as $a' \mathcal{J}(\theta) a = \int_{\Omega} (a' DF(\theta))^2 d\nu$, and is positive definite if and only if the components of $DF(\theta)$ are linearly independent functions in $L_2(\nu)$.

2.2 Main result

As in [23] our purpose is to consider product measures

$$(2.2) \quad \mu = \times_{n=1}^{\infty} \mu_n, \quad \lambda = \times_{n=1}^{\infty} \lambda_n, \quad \text{where } \mu_n = P_{\theta} \quad \text{and} \quad \lambda_n = P_{\theta+h_n},$$

$\theta \in \Theta$ is fixed and $\theta + h_n \in \Theta$, $n=1,2,\dots$. Under LeCam's condition

$$"l" : \limsup_{0 < \|h\| \rightarrow 0} d^2(P_{\theta+h}, P_{\theta}) / \|h\| < \infty,$$

Proposition 2 in [23] shows that $\sum_{n=1}^{\infty} \|h_n\|^2 < \infty$ implies $\mu \sim \lambda$. Here we obtain an equivalence-singularity dichotomy along with necessary and sufficient conditions for the two alternatives, when $\mathcal{J}(\theta)$ is positive definite at θ and the following separation type condition (which is usually assumed in asymptotic statistical theory [16]) is satisfied,

$$(2.3) \quad \text{"for all sufficiently small } \delta > 0, \quad \inf_{\|h\| > \delta} d^2(P_{\theta+h}, P_{\theta}) > 0".$$

PROPOSITION 2.1. *Let μ and λ be as in (2.2), F be differentiable at θ and $\mathcal{J}(\theta)$ be positive definite.*

- 1) *If $0 < \|h_n\| \rightarrow 0$ as $n \rightarrow \infty$, then*

$$\mu \sim \lambda \quad \bullet \quad \sum_{n=1}^{\infty} \|h_n\|^2 < \infty, \quad \text{and} \quad \mu \perp \lambda \quad \bullet \quad \sum_{n=1}^{\infty} \|h_n\|^2 = \infty.$$

ii) If condition (2.3) is satisfied, then the conclusions of 1) hold for any sequence $(h_n; n \in \mathbb{N})$.

The sufficiency for equivalence follows from [23, Proposition 2], since L_2 -differentiability is clearly stronger than condition "ℓ", but we include a simple complete proof here.

Proof. Since F is differentiable at θ , as $0 < \|h\| \rightarrow 0$ we have

$$| \|F(\theta+h) - F(\theta)\|_{L_2(v)} - \|\langle DF(\theta), h \rangle\|_{L_2(v)} | = o(\|h\|).$$

Thus for any $\epsilon > 0$ there exists $\delta = \delta(\epsilon) > 0$ such that if $0 < \|h\| < \delta$,

$$\|h\|^{-1} \|\langle DF(\theta), h \rangle\|_{L_2(v)}^{-\epsilon} < \|h\|^{-1} \|F(\theta+h) - F(\theta)\|_{L_2(v)} < \|h\|^{-1} \|\langle DF(\theta), h \rangle\|_{L_2(v)}^{+\epsilon}.$$

But $\|\langle DF(\theta), h \rangle\|_{L_2(v)}^2 = \int_{\Omega} |\langle DF(\theta), h \rangle|^2 dv = h' \mathcal{J}(\theta) h$, implies that for all $h \neq 0$,

$$k(\theta) \leq \|h\|^{-1} \|\langle DF(\theta), h \rangle\|_{L_2(v)} \leq K(\theta)$$

where $k(\theta)$ and $K(\theta)$ are the smallest and the largest eigenvalues of $\mathcal{J}(\theta)$. Since $\mathcal{J}(\theta)$ is positive definite, $k(\theta) > 0$ and we can choose $0 < \epsilon < k(\theta)$ so that for all $0 < \|h\| < \delta$,

$$0 < L(\theta) < \|h\|^{-1} \|F(\theta+h) - F(\theta)\|_{L_2(v)} < U(\theta)$$

where $L(\theta) = k(\theta) - \epsilon$ and $U(\theta) = K(\theta) + \epsilon$. Thus since $d^2(P_{\theta}, P_{\theta'}) = \|F(\theta) - F(\theta')\|_{L_2(v)}^2 / 8$ we have for n large

$$0 < \frac{1}{8} L^2(\theta) \|h_n\|^2 < d^2(\mu_n, \lambda_n) < \frac{1}{8} U^2(\theta) \|h_n\|^2$$

and the result follows from (2.1).

ii) If (2.3) is satisfied and $h_n \rightarrow 0$, then there exist $\delta > 0$ and a subsequence $(n_j; j \in \mathbb{N})$ with $\|h_{n_j}\| > \delta$. It follows that

$$\sum_{n=1}^{\infty} d^2(\mu_n, \lambda_n) \geq \sum_{j=1}^{\infty} d^2(\mu_{n_j}, \lambda_{n_j}) \geq \sum_{j=1}^{\infty} \inf_{\|h\| > \delta} d^2(P_{\theta+h}, P_{\theta}) = \infty,$$

and from (2.1), $\mu \perp \lambda$. This combined with i) gives the result. \square

It should be mentioned that the differentiability of $F(\theta)$ is generally difficult to verify, but is implied by the classical Cramér-Wold and Hajek regularity conditions, which play an important role in statistical estimation theory and are in principle easy to check (see e.g. [28], §77). However, L_2 -differentiability is weaker than any of these classical conditions, and the definition of Fisher information presented here extends the classical one, namely $\mathcal{I}(\theta) = -E\{\partial^2 \ell_n(dP_{\theta}/d\nu)/\partial\theta^2\}$ under the usual conditions on $dP_{\theta}/d\nu$.

2.3 Examples

Affine Transformations in \mathbb{R}^k .

Suppose $(X_n; n \in \mathbb{N})$ is a sequence of i.i.d. random vectors in \mathbb{R}^k , $(A_n; n \in \mathbb{N})$ a sequence of $k \times k$ matrices and $(b_n; n \in \mathbb{N})$ a sequence of vectors in \mathbb{R}^k . In order to compare the sequence of random vectors $(X_n; n \in \mathbb{N})$ with $(A_n X_n + b_n; n \in \mathbb{N})$ we can take as parameter space θ any open subset of

$$\begin{aligned} \{\theta = (A, b); A = (a_{ij}); k \times k \text{ matrix, } b = (b_i) \in \mathbb{R}^k\} \\ \equiv \{\theta; \theta = (a_{11}, \dots, a_{1k}, \dots, a_{kk}, b_1, \dots, b_k)\} \equiv \mathbb{R}^{k^2+k} \equiv \mathbb{R}^{k^2} \times \mathbb{R}^k \end{aligned}$$

containing the point $(I, 0)$, with norm

$$\|\theta\|_{\mathbb{R}^{(k \times k) + k}}^2 = \|A\|_{\mathbb{R}^{k \times k}}^2 + \|b\|_{\mathbb{R}^k}^2 = \sum_{i,j=1}^k a_{ij}^2 + \sum_{i=1}^k b_i^2.$$

With P the common distribution of the i.i.d. random vectors X_n and $\theta = (A, b)$, we

define

$$(2.4) \quad P_{\theta}(B) = P_{(A,b)}(B) = P(\{AX_n + b \in B\})$$

and note that $P = P_{(I,0)}$. From Proposition 2.1 we have the following

COROLLARY 2.2. *Let the probability measures P_{θ} defined as in (2.4) be such that for an open set $\theta \subset \mathbb{R}^{k^2} \times \mathbb{R}^k$ with $(I,0) \in \theta$, the family $\{P_{\theta}; \theta \in \theta\}$ is dominated by some σ -finite measure ν on \mathbb{R}^k , $F(\theta)$ is differentiable at $(I,0)$ and $\mathcal{F}(I,0)$ is positive definite. If $A_n \rightarrow I$ and $b_n \rightarrow 0$ as $n \rightarrow \infty$ then*

$$(X_n) \sim (A_n X_n + b_n) \iff \sum_{n=1}^{\infty} \|b_n\|_{\mathbb{R}^k}^2 < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} \|I - A_n\|_{\mathbb{R}^{k \times k}}^2 < \infty,$$

and otherwise $(X_n) \perp (A_n X_n + b_n)$. Furthermore if condition (2.3) is satisfied, the above conclusions hold for all sequences (A_n, b_n) in θ .

Proof. Putting $\theta = (I,0)$ and $(A_n, b_n) = \theta + h_n$ we have $h_n = (A_n - I, b_n)$ and

$$\|h_n\|_{\mathbb{R}^{(k \times k) + k}}^2 = \|A_n - I\|_{\mathbb{R}^{(k \times k)}}^2 + \|b_n\|_{\mathbb{R}^k}^2.$$

The conclusion then follows from Proposition 2.1. □

Remarks. a) Since the space of $k \times k$ matrices is finite dimensional, any norm can be used in place of $\|\cdot\|_{\mathbb{R}^{k \times k}}$.

b) When $A_n = I$ for all n , Corollary 2.2 extends the result on translates in [25] from random variables ($k=1$) to random vectors ($k=2$).

c) Corollary 2.2 contains Theorems 1 and 2 in [27], which consider the case where A_n is a rotation, i.e. $A_n x + b_n$ is a rigid motion of $x \in \mathbb{R}^k$.

d) When the X_n 's are Gaussian random variables ($k=1$) with mean zero and variance one, Corollary 2.2 can be checked directly by computing Hellinger distances. However, the computation of Hellinger distance is not simple in

higher dimensions ($k \geq 2$) even for Gaussian random vectors.

Stable Sequences.

Here we denote by $f_{(\alpha, \beta, a, b)}$ the univariate stable density whose characteristic function $\int_{-\infty}^{\infty} \exp(iux) f_{(\alpha, \beta, a, b)}(x) dx$ is

$$\begin{aligned} & \exp\{-|au|^{\alpha} \exp[-i\pi\beta \operatorname{sgn}(u)/2] + ibu\}, \quad \text{if } \alpha \neq 1, \\ & \exp\{-|au| - i(2\beta/\pi)au \ln(|au|) + ibu\}, \quad \text{if } \alpha = 1. \end{aligned}$$

where $0 < \alpha \leq 2$, $|\beta| \leq \alpha \wedge (2-\alpha)$, $a > 0$ and $-\infty < b < \infty$ (see [10]). If $\beta = 0$ and $b = 0$, we have the symmetric α -stable case (SoS).

We establish the equivalence-singularity dichotomy for certain sequences of independent stable variables. Because results about L_2 -differentiability and the validity of the condition (2.3) at $\alpha = 1$ are not known, we consider only limiting values $\alpha \neq 1$.

COROLLARY 2.3. Let $(X_{1n}, n \in \mathbb{N})$ be a sequence of i.i.d. stable variables with density $f_{(\alpha_0, \beta_0, a_0, b_0)}$ and let $(X_{2n}; n \in \mathbb{N})$ be a sequence of independent stable variables where the density of each X_{2n} is $f_{(\alpha_n, \beta_n, a_n, b_n)}$ with $(\alpha_n, \beta_n, a_n, b_n) \rightarrow (\alpha_0, \beta_0, a_0, b_0)$ and $\alpha_0 \neq 1$. Then

$$(X_{1n}) \sim (X_{2n}) \Leftrightarrow \begin{cases} \sum_{n=1}^{\infty} (\alpha_n - \alpha_0)^2 < \infty, & \sum_{n=1}^{\infty} (\beta_n - \beta_0)^2 < \infty, \\ \sum_{n=1}^{\infty} (a_n - a_0)^2 < \infty, & \sum_{n=1}^{\infty} (b_n - b_0)^2 < \infty. \end{cases}$$

and otherwise $(X_{1n}) \perp (X_{2n})$.

Proof. Let θ be any open subset of $\{\theta = (\alpha, \beta, a, b); \alpha \in (0, 1) \cup (1, 2),$

$|\beta| < \alpha \wedge (2-\alpha), a > 0, -\infty < b < \infty\}$ containing the point $\theta_0 = (\alpha_0, \beta_0, a_0, b_0)$. It is known that the densities $\{f_{\theta}, \theta \in \theta\}$ satisfy the usual Cramér-Wald regularity conditions ([9], p. 952); hence $f_{\theta}^{1/2}$ is $L_2(\text{Leb})$ -differentiable at each $\theta \in \theta$ (see

e.g. [28], §77). Moreover the Fisher information matrix $\mathcal{J}(\theta_0)$ is positive definite [8, p. 954]. Therefore the assumptions of Proposition 2.1.1) hold at θ_0 . Since for $h_n = (\alpha_n - \alpha_0, \beta_n - \beta_0, a_n - a_0, b_n - b_0)$ we have $\|h_n\|_{\mathbb{R}^4}^2 = (\alpha_n - \alpha_0)^2 + (\beta_n - \beta_0)^2 + (a_n - a_0)^2 + (b_n - b_0)^2$, the result follows. \square

When all parameters except shift b are kept fixed, the separation condition (2.3) follows from the inequality in [16, Example 3, p. 57], and when $\beta = 0$ and $\alpha \in (0, 2]$ is fixed, it has been proved in [19]. Hence we have the following

COROLLARY 2.4. Let $(X_n; n \in \mathbb{N})$ be a sequence of i.i.d. standard SaS variables with density $f_{(\alpha, 0, 1, 0)}$ and $\alpha \in (0, 2]$, and let (a_n, b_n) and (a'_n, b'_n) be two sequences of pairs of real numbers with $a_n \neq 0$. Then

$$\begin{aligned} (a_n X_n + b_n) \sim (a'_n X_n + b'_n) &\iff (X_n) \sim ((a_n/a'_n)X_n + (b_n - b'_n)/a'_n) \\ &\iff \sum_{n=1}^{\infty} \{1 - |a_n/a'_n|\}^2 < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} \{(b_n - b'_n)/a'_n\}^2 < \infty, \end{aligned}$$

and otherwise $(a_n X_n + b_n) \perp (a'_n X_n + b'_n)$.

Proof: The first equivalence follows since the map $(x_n) \rightarrow ((x_n - b'_n)/a'_n)$ is invertible and the second follows from Corollary 2.2 since

$$(a_n/a'_n)X_n + (b_n - b'_n)/a'_n \text{ has density } f_{(\alpha, 0, a_n/a'_n, (b_n - b'_n)/a'_n)}. \quad \square$$

We next explore the tail behavior of a stable distribution to show that two infinite sequences of independent symmetric stable variables with two different indexes of stability are singular.

PROPOSITION 2.5. Let $X_i = (X_{in}; n \in \mathbb{N})$, $i=1, 2$ be two sequences of (nondegenerate) independent symmetric stable variables with index of stability α_i in $(0, 2]$ and scale parameters (a_{in}) . If $\alpha_1 \neq \alpha_2$ then $\mu_1 \perp \mu_2$.

Proof: Assume $\alpha_1 < \alpha_2 < 2$. For each $\gamma \in (0, 2)$ let Z_γ denote a SaS r.v. with

scale parameter 1. Thus $\mu_{in}(B) \stackrel{\Delta}{=} P(X_{in} \in B) = P(a_{in} Z_{\alpha_1} \in B)$. Since $c^\gamma P(|Z_\gamma| > c^\gamma) \rightarrow C_\gamma$ as $c \rightarrow \infty$ where C_γ is a positive constant (see e.g. [11]), given any $\epsilon > 0$, there exist $M_{\gamma, \epsilon}$ such that for $c > M_{\gamma, \epsilon}$,

$$(C_\gamma - \epsilon)c^{-\gamma} < P(|Z_\gamma| > a) < (C_\gamma + \epsilon)c^{-\gamma}.$$

From now on fix ϵ such that $0 < \epsilon < \min(C_{\alpha_1}, C_{\alpha_2})$.

Case 1. Assume

$$\sigma_n \stackrel{\Delta}{=} a_{1n}/a_{2n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Define $\psi: \mathbb{F}^{\mathbb{N}} \rightarrow \mathbb{F}^{\mathbb{N}}$ by $\psi(x) = (\psi_n(x) = x_n/a_{2n}; n \in \mathbb{N})$. It follows that ψ is an i.i.d. sequence of standard $S\alpha_2 S$ r.v.'s under μ_2 and under μ_1 an independent sequence of $S\alpha_1 S$ r.v.'s with scale parameter $a_{1n}/a_{2n} = \sigma_n$.

As before let d_v denote the total variation distance between probability measures. For $c > M(1 + \sup_n \sigma_n)$ where $M = \max(M_{\alpha_1}, M_{\alpha_2})$, we have

$$d_v(\mu_{1n} \psi_n^{-1}, \mu_{2n} \psi_n^{-1}) \geq P(|Z_{\alpha_2}| > c) - P(|\sigma_n Z_{\alpha_1}| > c) > (C_{\alpha_2} - \epsilon)c^{\alpha_2} - \sigma_n^{\alpha_1} (C_{\alpha_1} + \epsilon)c^{\alpha_1},$$

and thus

$$\liminf_{n \rightarrow \infty} d_v(\mu_{1n} \psi_n^{-1}, \mu_{2n} \psi_n^{-1}) \geq (C_{\alpha_2} - \epsilon)/c^{\alpha_2} > 0.$$

Since $d_v \leq 2d$ where d denotes the Hellinger distance (see e.g. [28]) we have

$$\sum_{n=1}^{\infty} d(\mu_{1n} \psi_n^{-1}, \mu_{2n} \psi_n^{-1}) = \infty \quad \text{and therefore by Kakutani's Theorem } \mu_1 \psi^{-1} \perp \mu_2 \psi^{-1},$$

which implies $\mu_1 \perp \mu_2$.

Case 2. Assume $\sigma_n \rightarrow 0$. Thus there exist $\delta > 0$ and a sequence $(n_k; k \in \mathbb{N})$ such that $\sigma_{n_k} \geq \delta$, i.e. $\sigma_{n_k}^{-1} \leq \delta^{-1}$. Define $\phi: \mathbb{F}^{\mathbb{N}} \rightarrow \mathbb{F}^{\mathbb{N}}$ by $\phi(x) = (\phi_k(x) = x_k/a_{1n_k}; k \in \mathbb{N})$. Then ϕ is an i.i.d. sequence of standard $S\alpha_1 S$ r.v.'s under μ_1 and under

μ_2 an independent sequence of $S\alpha_2 S$ r.v.'s with scale parameter $a_{2n_k}/a_{1n_k} = \sigma_{n_k}^{-1}$

For $c > M(1 + \delta^{-1})$ we have

$$\begin{aligned}
d_v(\mu_{1n_k} \phi_{n_k}^{-1}, \mu_{2n_k} \phi_{n_k}^{-1}) &\geq P(|Z_{\alpha_1}| > c) - P(|\sigma_{n_k}^{-1} Z_{\alpha_2}| > c) \\
&> (C_{n_k} - \epsilon)c^{\alpha_1} - \sigma_{n_k}^{-1}(C_{\alpha_2} + \epsilon)c^{\alpha_2} > (C_{n_k} - \epsilon)c^{\alpha_1} - \delta^{-1}(C_{\alpha_2} + \epsilon)c^{\alpha_2} \triangleq \delta'(c).
\end{aligned}$$

Since $\alpha_1 < \alpha_2$, we have $\delta'(c) > 0$ if and only if $c^{\alpha_2 - \alpha_1} > \delta^{-1}(C_{\alpha_2} + \epsilon)/(C_{\alpha_1} - \epsilon)$.

Thus, fixing $c > M(1 + \delta^{-1} + (C_{\alpha_2} + \epsilon)/(C_{\alpha_1} - \epsilon)^{1/(\alpha_2 - \alpha_1)})$ we obtain

$$\limsup_{n \rightarrow \infty} d_v(\mu_{1n} \phi_n^{-1}, \mu_{2n} \phi_n^{-1}) > \delta'(c) > 0$$

and the conclusion follows as in case 1.

If $\alpha_2 = 2$, the result can be shown with minor modifications in the proof. \square

3. DICHOTOMIES FOR CERTAIN SoS PROCESSES

For stochastic processes the equivalence-singularity dichotomy has been proved for product measures [18], for Gaussian processes ([10] and [14]), and for certain ergodic measures [20]. In [24], it was shown that this dichotomy prevails for translates of certain SoS processes. Such dichotomy for general SoS measures has been conjectured in [7] but the problem remains open. In this section we show that an equivalence-singularity dichotomy holds for certain SoS processes, e.g. independently scattered SoS measures and harmonizable SoS processes, and we give necessary and sufficient conditions for the two alternatives for all $\alpha \in (0, 2]$.

Recall that a random variable X is SoS with scale parameter $\|X\|_\alpha \in (0, \infty)$ if $E\{\exp(iuX)\} = \exp(-\|X\|_\alpha^\alpha |u|^\alpha)$, and a stochastic process $X = (X(t); t \in T)$ is SoS if all linear combinations $\sum_{k=1}^n a_k X(t_k)$ are SoS variables. When $\alpha = 2$ we have zero mean Gaussian variables and processes respectively. When $0 < \alpha < 2$, the tails of the distributions are heavier and only moments of order $p \in (0, \alpha)$ are finite.

We first prove a dichotomy for independently scattered SaS measures. Let I be an arbitrary set and \mathcal{F} a δ -ring of subsets of I with the property that there exists an increasing sequence $(I_n; n \in \mathbb{N})$ in \mathcal{F} with $\bigcup_n I_n = I$. A real stochastic process $Z = (Z(B); B \in \mathcal{F})$ is called an independently scattered SaS measure if for every sequence $(B_n; n \in \mathbb{N})$ of disjoint sets in \mathcal{F} , the random variables $\{Z(B_n); n \in \mathbb{N}\}$ are independent and whenever $\bigcup_n B_n \in \mathcal{F}$ then $Z(\bigcup_n B_n) = \sum_n Z(B_n)$ a.s., and for every $B \in \mathcal{F}$ the random variable $Z(B)$ is SaS, i.e.

$E\{\exp(iuZ(B))\} = \exp\{-m(B)|u|^\alpha\}$ where $m(B) = \|Z(B)\|_\alpha^\alpha$. Then m is a measure on \mathcal{F} which extends uniquely to a σ -finite measure on $\sigma(\mathcal{F})$, and is called the control measure of Z . The existence of an independently scattered SaS measure with a given control measure is a consequence of Kolmogorov's consistency theorem. When I is an interval of the real line and the control measure m is Lebesgue measure, then X has stationary independent increments,

$E\{\exp(iu[X(t)-X(t')])\} = \exp\{-|t-t'| |u|^\alpha\}$, and is called SaS motion on I .

The following notation will be used in Proposition 3.1. Recall that if a σ -finite measure space $(I, \sigma(\mathcal{F}), m)$ is such that $\sigma(\mathcal{F})$ contains all single points sets (e.g. I is a Polish space, $\sigma(\mathcal{F})$ its Borel sets, and \mathcal{F} the δ -ring of Borel sets with finite m -measure) then m can be decomposed into $m = m_a + m_d$ where m_a is purely atomic and m_d is diffuse (non-atomic) [21], and the set of atoms is at most countable, say $A = \{a_n\}$. Thus if $Z = (Z(B); B \in \mathcal{F})$ is an independently scattered SaS measure with control measure m , it can be decomposed into

$$Z = Z_a + Z_d,$$

where Z_a and Z_d are independent SaS independently scattered measures defined for all $B \in \mathcal{F}$ by $Z_a(B) = Z(A \cap B)$ and $Z_d(B) = Z(A^c \cap B)$, and have control measures m_a and m_d respectively. The atomic component has a series expansion

$Z_a(B) = \sum_n 1_B(a_n) Z(\{a_n\})$ which can be normalized by using the i.i.d. standard SaS variables $Z_n \stackrel{\Delta}{=} Z(\{a_n\}) m^{-1/\alpha}(\{a_n\})$ with $E\{\exp(iuZ_n)\} = \exp(-|u|^\alpha)$, as follows:

$$Z_a(B) = \sum_n 1_B(a_n) m^{1/\alpha}(\{a_n\}) Z_n .$$

PROPOSITION 3.1. For $i=1,2$, let $Z_i = (Z_i(B); B \in \mathcal{F})$ be an independently scattered $S\alpha_1 S$ measure with $\alpha_i \in (0,2]$ and control measure m_i which is not purely discrete with a finite number of atoms. Then $\mu_1 \sim \mu_2$ if and only if the following conditions are satisfied

- i) $\alpha_1 = \alpha_2$,
- ii) $m_{1d} = m_{2d}$,
- iii) m_1 and m_2 have the same set of atoms $A = \{a_n\}$ and

$$\sum_n [1 - m_1(\{a_n\})/m_2(\{a_n\})]^2 < \infty .$$

Furthermore if any of these conditions fails $\mu_1 \perp \mu_2$.

Proof. First suppose that m_1 and m_2 are not equivalent, e.g. $m_2 \not\ll m_1$. Then there exists $B \in \sigma(\mathcal{F})$ such that

$$\|Z_1(B)\|_{\alpha_1}^{\alpha_1} = m_1(B) = 0, \quad \text{and} \quad \|Z_2(B)\|_{\alpha_2}^{\alpha_2} = m_2(B) > 0.$$

Define $\Gamma_B: \mathbb{F}^{\mathcal{F}} \rightarrow \mathbb{F}$ by $\Gamma_B(x) = x(B)$. It follows that $\mu_1 \Gamma_B^{-1} \perp \mu_2 \Gamma_B^{-1}$ and thus $\mu_1 \perp \mu_2$. From now on we assume $m_1 \sim m_2$.

Suppose $\alpha_1 \neq \alpha_2$. Since m_1 and m_2 are not purely atomic with a finite number of atoms, we can choose an infinite sequence $(B_n; n \in \mathbb{N})$ of disjoint sets in \mathcal{F} such that $m_i(B_n) > 0$, $i = 1,2$. Define $\psi: \mathbb{F}^{\mathcal{F}} \rightarrow \mathbb{F}^{\mathbb{N}}$ by $\psi(x) = (\psi_n(x) = x(B_n); n \in \mathbb{N})$. Thus, for $i=1,2$, under μ_i , ψ is a sequence of independent $S\alpha_i S$ r.v.'s with $\|\psi_n\|_{\alpha_i}^{\alpha_i} = m_i(B_n)$. It follows from Proposition 2.5 that if $\alpha_1 \neq \alpha_2$, then $\mu_1 \psi^{-1} \perp \mu_2 \psi^{-1}$, so that $\mu_1 \perp \mu_2$. From now on we assume $\alpha_1 = \alpha_2 = \alpha$.

Since $m_1 \sim m_2$ we have $m_{1d} \sim m_{2d}$. Suppose $m_{1d} \neq m_{2d}$, so that $m_{id}(\{dm_{2d}/dm_{1d} \neq 1\}) > 0$, $i=1,2$, hence

$$m_{1d}(\{0 < dm_{2d}/dm_{1d} < 1\}) > 0 \quad \text{or} \quad m_{1d}(\{dm_{2d}/dm_{1d} > 1\}) > 0.$$

Assume $m_{1d}(\{dm_{2d}/dm_{1d} > 1\}) > 0$. Then there exists $\delta > 1$ such that $m_{1d}(\{dm_{2d}/dm_{1d} > \delta\}) > 0$. Since m_{1d} is nonatomic, we can find a sequence $(B_n; n \in \mathbb{N})$ of disjoint subsets of $\{dm_{1d}/dm_{1d} > \delta\}$ such that $m_{1d}(B_n) > 0$. Let $\phi: \mathbb{F}^{\mathcal{G}} \rightarrow \mathbb{F}^{\mathbb{N}}$ be the map defined by $\phi(x) = (\phi_n(x) = x(A^c \cap B_n)/m_{1d}(B_n))^{1/\alpha}; n \in \mathbb{N}$. Under μ_1 , ϕ is an i.i.d. sequence of standard SoS r.v.'s, and under μ_2 , ϕ is an independent sequence of SoS r.v.'s with $\|\phi_n\|_{\alpha}^{\alpha} = m_{2d}(B_n)/m_{1d}(B_n)$. It follows from Corollary 2.4 that $\mu_1 \phi^{-1}$ and $\mu_2 \phi^{-1}$ are either equivalent or singular, and they are singular if and only if

$$(3.1) \quad \sum_n \{ 1 - [m_{2d}(B_n)/m_{1d}(B_n)]^{1/\alpha} \}^2 = \infty.$$

Now by construction

$$m_{2d}(B_n) = \int_{B_n} \frac{dm_{2d}}{dm_{1d}} dm_{1d} > \delta m_{1d}(B_n).$$

Hence $1 < \delta < m_{2d}(B_n)/m_{1d}(B_n)$, so that (3.1) holds. Thus $\mu_1 \phi^{-1} \perp \mu_2 \phi^{-1}$ which implies $\mu_1 \perp \mu_2$.

If $m_{1d}(\{dm_{2d}/dm_{1d} > 1\}) = 0$ we have $m_{1d}(\{dm_{1d}/dm_{2d} > 1\}) > 0$ and an identical argument applies. Therefore $m_1 \sim m_2$ and $m_{1d} \neq m_{2d}$ implies $\mu_1 \perp \mu_2$.

Now assume $m_{1d} = m_{2d}$. Since $m_1 \sim m_2$, they have the same set of atoms $A = \{a_n\}$. Suppose $\mu_2 < \mu_1$ and let $\Xi: \mathbb{F}^{\mathcal{G}} \rightarrow \mathbb{F}^{\mathbb{N}}$ be defined by $\Xi(x) = (\Xi_n(x) = x(\{a_n\})/m_1(\{a_n\}))^{1/\alpha}; n \in \mathbb{N}$. Thus $\mu_2 \Xi^{-1} < \mu_1 \Xi^{-1}$ and Ξ is an i.i.d. sequence of standard SoS r.v.'s under μ_1 and under μ_2 an independent sequence of SoS r.v.'s with $\|\Xi_n\|_{\alpha}^{\alpha} = m_2(\{a_n\})/m_1(\{a_n\})$. Hence by Corollary 2.4,

$$(3.2) \quad \sum_n \{ 1 - [m_2(\{a_n\})/m_1(\{a_n\})]^{1/\alpha} \}^2 < \infty.$$

Also, if (3.2) does not hold, again Corollary 2.4 implies $\mu_1 \Xi^{-1} \perp \mu_2 \Xi^{-1}$ so that $\mu_1 \perp \mu_2$.

Note that (3.1) and (3.2) are symmetric in m_1 and m_2 and independent of α as for $q \neq 0$, $\sum_n (1-u_n)^2 < \infty$ if and only if $\sum_n (1-u_n^q)^2 < \infty$. Hence (3.2) can be replaced by iii).

Conversely, suppose that i), ii) and iii) hold. Since $m_{1d} = m_{2d}$ we have

$$Z_i \stackrel{L}{=} Z_{ia} + Z_d, \quad i=1,2,$$

where Z_{ia} and Z_d are independent, independently scattered SoS measures with control measures m_{ia} and $m_d = m_{1d} = m_{2d}$ respectively, and $\stackrel{L}{=}$ denotes equality in law. Let $\phi: \mathbb{F}^N \rightarrow \mathbb{F}^{\mathcal{G}}$ be defined by

$$[\phi(y)](B) = \phi(y, B) = \sum_{n=1}^{\infty} 1_B(a_n) m_1(\{a_n\})^{1/\alpha} y_n, \quad y=(y_n) \in \mathbb{Z}^N.$$

Thus $(\phi \circ \Xi)(Z_i) \stackrel{L}{=} Z_{ia}$, so that $\mu_{ia} = (\mu_i \Xi^{-1}) \phi^{-1}$, $i=1,2$. Now by Corollary 2.4, iii) implies $\mu_1 \Xi^{-1} \sim \mu_2 \Xi^{-1}$, hence $\mu_{1a} \sim \mu_{2a}$. Therefore, since $\mu_i = \mu_{i,a} * \mu_d$, $i=1,2$, it follows that $\mu_1 \sim \mu_2$. □

The results in Proposition 3.1 can be extended to certain symmetric (dependent) stable processes. Let Z be an independently scattered SoS measure with control measure m . For any function $f \in L_{\alpha}(I, \sigma(\mathcal{F}), m) = L_{\alpha}(m)$ the stochastic integral $\int_I f dZ$ can be defined in the usual way and is a SoS variable with $\|\int_I f dZ\|_{\alpha} = \|f\|_{L_{\alpha}(m)}$. The map $f \rightarrow \int_I f dZ$ from $L_{\alpha}(m)$ into $\mathcal{L}(Z)$ is an isometry and

$$(3.3) \quad \mathcal{L}(Z) = \left\{ \int_I f dZ; f \in L_{\alpha}(m) \right\}.$$

The stochastic integral allows for the construction of SoS processes with generally dependent values by means of the spectral representation

$$(3.4) \quad X(t) = \int_I f(t, u) Z(du), \quad t \in T,$$

where $\{f(t, \cdot); t \in T\} \subset L_{\alpha}(m)$. In fact every SoS process X has such a spectral

representation in law, in the sense that for some family $\{f(t, \cdot), t \in T\}$ in some $L_\alpha(m)$.

$$(3.5) \quad (X(t); t \in T) \stackrel{L}{=} (\int_I f(t, u) Z(du); t \in T)$$

(see e.g. [22] and [15]). Some examples of SaS processes will be considered at the end of the section.

Let $X = (X(t); t \in T)$ be a SaS process with spectral representation as in (3.4). It follows from the continuity of the stochastic integral map $f \rightarrow \int f dZ$ that the representing functions $\{(f(t, \cdot); t \in T)\}$ are linearly dense in $L_\alpha(m)$, $\overline{\text{sp}}\{f(t, \cdot); t \in T\} = L_\alpha(m)$, if and only if $\mathcal{V}(X) = \mathcal{V}(Z)$. Processes satisfying this condition will be said to have an invertible spectral representation or more simply to be invertible. Gaussian processes are invertible [1]. For non-Gaussian SaS processes this is not generally true [5]. Conditions for invertible representation are given in [3] and [5]. SaS processes with invertible representation in $L_2([0,1])$ are considered in [30].

Let $X_i = (X_i(t); t \in T)$, $i=1,2$, be two invertible Sa_iS processes with spectral representations $X_i(t) = \int_I f(t, u) Z_i(du)$, where Z_i are independently scattered Sa_iS measures with control measures m_i and $f(\cdot, t) \in L_{\alpha_1}(m_1) \cap L_{\alpha_2}(m_2)$, $t \in T$. X_1 and X_2 will be called *simultaneously invertible* if for each $B \in \mathcal{F}$ there exist $N_n(B)$, $a_{n1}(B), \dots, a_{nN_n(B)}(B)$, $t_{n1}(B), \dots, t_{nN_n(B)}(B)$ such that

$$\sum_{k=1}^{N_n(B)} a_{nk} f(t_{nk}(B), \cdot) \rightarrow 1_B(\cdot) \quad \text{as } n \rightarrow \infty,$$

in $L_{\alpha_1}(m_1)$ for both $i=1,2$. E.g., X_1 and X_2 are simultaneously invertible if they are invertible, and either $\alpha_1 = \alpha_2$ and dm_1/dm_2 is bounded above or below, or their associated random measures Z_1 and Z_2 are equivalent (cf. Proposition 3.1). The simultaneous invertibility of X_1 and X_2 allows for the study of the equivalence and singularity of μ_1, μ_2 in terms of that of Z_1, Z_2 . Indeed

$X_1(t) = \int f(t,u)Z_1(du)$ is, roughly speaking, $X_1 = L(Z_1)$, where L is a linear map from $\mathcal{L}(Z_1)$ into $\mathcal{L}(X_1)$. Simultaneous invertibility is like having $Z_1 = L^{-1}(X_1)$, so the singularity of Z_1, Z_2 should imply the singularity on equivalence of X_1 and X_2 , and vice-versa for equivalence. The next proposition makes this precise.

PROPOSITION 3.2. Let $X_i = (X_i(t); t \in T)$ be two simultaneously invertible $S\alpha_1 S$ processes with $\alpha_i \in (0, 2]$ and spectral representations $X_i(t) = \int_I f(t,u)Z_i(du)$, where Z_i are independently scattered $S\alpha_1 S$ measures with control measures m_i which are not purely discrete with a finite number of atoms. Then μ_{X_1} and μ_{X_2} are either equivalent or singular, and

$$\mu_{X_1} \sim \mu_{X_2} \quad \bullet \quad \mu_{Z_1} \sim \mu_{Z_2}, \quad \mu_{X_1} \perp \mu_{X_2} \quad \bullet \quad \mu_{Z_1} \perp \mu_{Z_2},$$

i.e. $\mu_{X_1} \sim \mu_{X_2}$ if and only if conditions i), ii) and iii) of Proposition 3.1 are satisfied, and otherwise $\mu_{X_1} \perp \mu_{X_2}$.

Proof: For $B \in \mathcal{F}$ we can define

$$\phi_n(B, x) = \sum_{k=1}^{N_n(B)} a_{nk}(B) x(t_{nk}(B)), \quad x \in \mathbb{F}^T,$$

so that $\phi_n(B, X_i(\cdot, \omega)) \rightarrow Z_i(B, \omega)$, in probability as $n \rightarrow \infty$, $i=1,2$. Let $(\phi_{n_k}(B, \cdot); k \in \mathbb{N})$ be a subsequence converging a.s. (μ_i) , $i=1,2$, and put

$$\tilde{Z}(B) = \tilde{Z}(B, \cdot) = \liminf_{k \rightarrow \infty} \phi_{n_k}(B, \cdot) 1_{\{x; \phi_{n_k}(x) \text{ converges}\}}(\cdot).$$

Hence $\tilde{Z}(B, X_i(\cdot, \omega)) = Z_i(B, \omega)$ a.s., $i=1,2$. The stochastic process $\tilde{Z} = (\tilde{Z}(B))$, $B \in \mathcal{F}$ defined on $(\mathbb{F}^T, \mathcal{G})$ is an independently scattered $S\alpha_1 S$ measure with control measure m_i under μ_{X_i} . If we also denote by \tilde{Z} the map $x \rightarrow \tilde{Z}(\cdot, x)$ then

$$\mu_{X_1} \sim \mu_{X_2} \quad \Rightarrow \quad \mu_{X_1} \tilde{Z}^{-1} \sim \mu_{X_2} \tilde{Z}^{-1} \quad (\text{i.e. } \mu_{Z_1} \sim \mu_{Z_2}) \quad \text{and} \quad \mu_{Z_1} \perp \mu_{Z_2} \quad (\text{i.e.}$$

$$\mu_{X_1}^{\tilde{Z}^{-1}} \perp \mu_{X_2}^{\tilde{Z}^{-1}} \Rightarrow \mu_{X_1} \perp \mu_{X_2}.$$

On the other hand if $\mu_{Z_1} \sim \mu_{Z_2}$, i.e. $\mu_{X_1}^{\tilde{Z}^{-1}} \sim \mu_{X_2}^{\tilde{Z}^{-1}}$, it follows that

i)-iii) of Proposition 3.1 hold. Thus, we can construct independent processes \tilde{X}_d and \tilde{X}_{ia} on $(\mathbb{F}^{\mathcal{G}}, \mathcal{C}(\mathbb{F}^{\mathcal{G}}), \mu_{Z_i})$ such that

$$X_i \stackrel{L}{=} \tilde{X}_d + \tilde{X}_{ia}, \quad i=1,2.$$

with $\mu_{X_{1a}} \sim \mu_{X_{2a}}$. Since $\mu_{X_1} = \mu_{X_d} * \mu_{X_{1a}}$, we have $\mu_{X_1} \sim \mu_{X_2}$.

Now if μ_{X_1} and μ_{X_2} are not equivalent it follows that $\mu_{Z_1} \perp \mu_{Z_2}$ (since otherwise $\mu_{Z_1} \sim \mu_{Z_2}$, which implies $\mu_{X_1} \sim \mu_{X_2}$, i.e. a contradiction) and this was shown to imply $\mu_{X_1} \perp \mu_{X_2}$. □

It follows from Proposition 3.2 that simultaneously invertible processes are singular whenever their indexes of stability are different. This is not generally true for symmetric stable processes with different indexes of stability. Indeed, let $G = \{G(t); t \in T\}$ be a Gaussian process, and for $i=1,2$, let A_i be a standard positive $(\alpha_i/2)$ -stable random variable where $\alpha_1 \neq \alpha_2$, and consider the sub-Gaussian $S\alpha_1 S$ processes $X_i = (X_i(t) = A_i^{1/2} G(t); t \in T)$. We have that $\mu_{X_i}(B) = \int_{\mathbb{R}^+} \mu_{XG}(B) \mu_{A_i}(dx)$. Since the distribution μ_{A_i} of A_i has positive density in \mathbb{R}^+ we have $\mu_{A_1} \sim \mu_{A_2}$, so that by the Corollary of Theorem 18.1 in [26], $\mu_{X_1} \sim \mu_{X_2}$. Since the linear space of a sub-Gaussian process does not contain (nondegenerate) independent random variables (see [5]), sub-Gaussian processes are not invertible (nor simultaneously invertible). Further examples of symmetric stable processes with different indexes of stability which are equivalent are $X_i = (X_i(t) = \sum_{n=1}^N A_{in}^{1/2} G_n(t); t \in T)$ where for each $i=1,2$, the vector (A_{i1}, \dots, A_{iN}) is positive $(\alpha_i/2)$ -stable, independent of the mutually

independent Gaussian processes $G_n = (G_n(t); t \in T)$, $n = 1, \dots, N$.

As a consequence of Proposition 3.2, harmonizable processes are either equivalent or singular and necessary and sufficient conditions for the two alternatives are provided.

COROLLARY 3.3. Let $X_k = (X_k(t); t \in T)$, $k=1,2$, be two harmonizable $S\alpha_k S$ processes, with $\alpha_k \in (0,2]$, i.e.

$$X_k(t) = \int_I \exp(i\langle t, u \rangle) Z_k(du), \quad t \in T,$$

where $I = \mathbb{R}^d$, respectively $[-\pi, \pi]^d$, for $T = \mathbb{R}^d$, respectively \mathbb{Z}^d , and Z_k are independently scattered $S\alpha_k S$ measures with finite spectral measures m_k which are not purely discrete with a finite number of atoms. Then μ_{X_1} and μ_{X_2} are equivalent if and only if i), ii) and iii) of Proposition 3.1 are satisfied, and they are singular otherwise.

Proof: Clearly X_1 and X_2 are simultaneously invertible, since indicator functions can be approximated uniformly, and hence in $L_{\alpha_k}(m_k)$, by linear combinations of the functions $f(t, u) = \exp(i\langle t, u \rangle)$. Hence the result follows from Proposition 3.2 □

As a special case, let S and N be harmonizable $S\alpha S$ signal and noise processes as in Corollary 3.3, that are independent of each other. Then μ_{S+N} and μ_S are equivalent if and only if

$$m_{S,d} = 0, \quad \text{the atoms of } m_S \text{ are atoms of } m_N, \text{ and}$$

$$\sum_n \left[\frac{m_S(\{a_n\})}{m_S(\{a_n\}) + m_N(\{a_n\})} \right]^2 < \infty.$$

Otherwise μ_{S+N} and μ_N are singular, and the presence of the random signal S in the additive noise N can be detected with probability one (at least in principle). In particular, μ_{S+N} and μ_N are singular when the signal has

continuous spectrum or the noise has no atomic spectrum. (Similar results hold when the signal and noise processes have simultaneously invertible representations as in Proposition 3.2).

The results in Propositions 3.1 and 3.2 and Corollary 3.3 are identical in the non-Gaussian stable case and in the Gaussian case [6]. However in the case of Corollary 3.3 much more is known for Gaussian processes. Namely, for stationary Gaussian processes ($d=1$) restricted over a finite interval, the equivalence-singularity dichotomy prevails and necessary and sufficient conditions for the two alternatives are known (see e.g. [17]). Both of these important questions remain open in the non-Gaussian stable case.

Another consequence of Proposition 3.2 is the singularity of multiples of invertible processes.

COROLLARY 3.4. *Let $X=(X(t); t \in T)$ be an invertible SaS process with $\alpha \in (0,2]$ and control measure m which is not purely atomic with a finite number of atoms. Then X and bX are singular wherever $|b| \neq 1$.*

Proof. If $X(t) = \int f(t,u)Z(du)$, where Z has control measure m , then $bX(t) = \int f(t,u)Z_b(du)$ where $Z_b = bZ$ has control measure $|b|^\alpha m$. Clearly X and bX are simultaneously invertible and the result follows from Proposition 3.2. \square

The result in Corollary 3.4 is known to hold for every Gaussian process with infinite dimensional linear space. Here again the class of SaS sub-Gaussian processes provides an example to show that the result is not true for all infinite dimensional SaS processes. In fact, if $X = (A^{1/2}G(t); t \in T)$, as before, we have for each $b > 0$, $\mu_{bX}(B) = \int_{\mathbb{R}^+} \mu_{XG}(B) \mu_{bA}(dx)$. The distributions μ_A and μ_{bA} are equivalent for all $b > 0$ so that $\mu_X \sim \mu_{bX}$.

In the Gaussian case the multiple b in Corollary 3.4 is allowed to be a function $b(t)$, but this problem remains open in the non-Gaussian stable case.

Corollary 3.4 is relevant to the detection of a constant signal in multiplicative noise. (See [2])

4. REMARKS ON SINGULARITY AND ABSOLUTE CONTINUITY OF p^{th} -ORDER AND SoS PROCESSES

For two Gaussian processes, the setwise equality of their RKHS's is a necessary condition for equivalence. For two second order processes a necessary condition for absolute continuity and a sufficient condition for singularity in terms of their RKHS's are proved in [12]. We show that these results remain true for SoS processes and for p^{th} order processes with $1 < p < 2$ respectively, with the RKHS replaced by an appropriate function space \mathcal{F} specified in the sequel.

The function space of a p^{th} order process $X = (X(t); t \in T)$ is defined in [24] by

$$\mathcal{F} = \left\{ s: T \rightarrow \mathbb{F}; \|s\|_{\mathcal{F}} \triangleq \sup_{\substack{N, a_1, \dots, a_N \\ t_1, \dots, t_N}} \frac{|\sum_{n=1}^N a_n s(t_n)|}{\|\sum_{n=1}^N a_n X_1(t_n)\|_{L_p(P)}^{1/p}} < \infty \right\}.$$

Note that when $p = 2$, $\mathcal{F} = \text{RKHS}$. If $X_i = (X_i(t); t \in T)$, $i=1,2$, are two p^{th} order processes, we say that X_1 dominates X_2 if there exists $0 < K < \infty$ such that for all $N \in \mathbb{N}$, $a_1, \dots, a_N \in \mathbb{R}^1$ and $t_1, \dots, t_N \in T$,

$$\|\sum_{n=1}^N a_n X_2(t_n)\|_{L_p(P)} \leq K \|\sum_{n=1}^N a_n X_1(t_n)\|_{L_p(P)}.$$

The relationship between domination and the function spaces is clarified in the following

PROPOSITION 4.1. Let $X_i = (X_i(t); t \in T)$ be a p^{th} order processes with function space $\mathcal{F}_i, i=1,2$.

- i) If X_1 dominates X_2 , then $\mathcal{F}_2 \subset \mathcal{F}_1$.
- ii) X_1 dominates X_2 if and only if there exists a bounded linear transformation $V: \mathcal{L}(X_1) \rightarrow \mathcal{L}(X_2)$, satisfying $V(X_1(t)) = X_2(t), t \in T$.

Consequently, if X_1 dominates X_2 and vice versa, then $\mathfrak{F}_1 = \mathfrak{F}_2$ (setwise), $\|\cdot\|_{\mathfrak{F}_1}$ and $\|\cdot\|_{\mathfrak{F}_2}$ are equivalent, and the transformation V has bounded inverse.

Proof. i) If X_1 dominates X_2 , it follows that for all functions s ,

$$\|s\|_{\mathfrak{F}_2} \leq K \|s\|_{\mathfrak{F}_1}, \text{ and thus } \mathfrak{F}_2 \subset \mathfrak{F}_1.$$

ii) Let $V: \mathcal{L}(X_1) \rightarrow \mathcal{L}(X_2)$ be defined by $V(\sum_{n=1}^N a_n X_1(t_n)) = \sum_{n=1}^N a_n X_2(t_n)$. It is clear that V is a well defined bounded linear transformation and as such it can be extended to $\mathcal{L}(X_1)$ if and only if X_1 dominates X_2 . \square

For SoS, the next Proposition shows that mutual domination is a necessary condition for absolute continuity, i.e. non-domination is a sufficient condition for singularity. This Proposition is a stochastic process version of Proposition 7 in [30].

PROPOSITION 4.2. Let $X_i = (X_i(t); t \in T)$, $i = 1, 2$, be two SoS processes. If μ_1 and μ_2 are not singular, then X_1 dominates X_2 , X_2 dominates X_1 , and $\mathfrak{F}_1 = \mathfrak{F}_2$. Equivalently if $\mathfrak{F}_1 \neq \mathfrak{F}_2$ then either X_1 does not dominate X_2 or X_2 does not dominate X_1 and $\mu_1 \perp \mu_2$.

Proof: Since for $Y \in \mathcal{L}(X_i)$, $\|Y\|_{L_p(P)} = C_{p, \alpha_i} \|Y\|_{\alpha_i}$ (see [4]), X_1 dominates X_2 if and only if

$$\|\sum_{n=1}^N a_n X_2(t_n)\|_{\alpha_2} \leq K \|\sum_{n=1}^N a_n X_1(t_n)\|_{\alpha_1}.$$

Assume X_1 does not dominate X_2 . Then for any positive sequence $K_n \rightarrow \infty$, as

$n \rightarrow \infty$, there exist $Y_n^{(i)} = \sum_{k=1}^N a_{n,k} X_i(t_{n,k})$, $i=1, 2$, such that

$\|Y_n^{(2)}\|_{\alpha_2} \geq K_n \|Y_n^{(1)}\|_{\alpha_1}$, $n=1, 2, \dots$. Without loss of generality we can assume

$\|Y_n^{(1)}\|_{\alpha_2} = 1$ for all n . Thus

$$\|Y_n^{(1)}\|_{\alpha_2} \leq 1/K_n \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Now consider the sequence of random variables $(Y_n; n \in \mathbb{N})$ defined on $(\mathbb{F}^T, \mathcal{G})$ by $Y_n(x) = \sum_{k=1}^n a_{n,k} x(t_{n,k})$, $x \in \mathbb{F}^T$. It follows that

$$\int_{\mathbb{F}^T} \exp(iuY_n) d\mu_1 = \exp(-\|Y_n^{(1)}\|_{\alpha_1}^{\alpha_1} |u|^{\alpha_1}) \rightarrow 1 \quad \text{as } n \rightarrow \infty.$$

Hence a subsequence $(Y_{n_k}; k \in \mathbb{N})$ can be chosen such that if $C_0 = \{x; Y_{n_k}(x) \rightarrow 0 \text{ as } k \rightarrow \infty\}$, then $\mu_{X_1}(C_0) = 1$. Clearly C_0 is a measurable linear subspace of \mathbb{F}^T and, since μ_2 is a $S\alpha_2S$ measure on $(\mathbb{F}^T, \mathcal{G})$, it follows by the zero-one law for stable measures [8] that $\mu_2(C_0) = 0$ or 1. On the other hand,

$$\int_{\mathbb{F}^T} \exp(iuY_{n_k}) d\mu_2 = \exp(-\|Y_{n_k}^{(1)}\|_{\alpha_2}^{\alpha_2} |u|^{\alpha_2}) = \exp(-|u|^{\alpha_2})$$

which implies that $\mu_2(C_0) = 0$ and thus $\mu_1 \perp \mu_2$. □

The crucial result used in the proof of Proposition 4.2 is the zero-one law, which is not available for general p^{th} order processes. However the proposition has some partial analogs for certain p^{th} order processes.

As in [12] we call a p^{th} order process $X = (X(t); t \in T)$ non-reduced if there exists some $\epsilon \in (0, 1]$ such that for all countable subsets T_0 of T , $P(\{\omega; X(t, \omega) = 0, t \in T_0\}) \geq \epsilon$; otherwise X is called reduced. Nontrivial $S\alpha S$ processes are reduced. When X is separable and T an interval of the real line it is shown in [12] that X is reduced if and only if $P(\{X(t) = 0, t \in T\}) = 0$, and nonreduced if and only if $P(\{X(t) = 0, t \in T\}) \geq \epsilon$ for some $\epsilon \in (0, 1]$.

Next we generalize to p^{th} order processes with $1 < p < 2$ the results in [12], Théorèmes (3.2) and (3.3.2). The proof is essentially identical to Fortet's and is presented in a shorter form.

PROPOSITION 4.3. Let $X_i = (X_i(t); t \in T)$, be a p^{th} order process with $1 < p < 2$ and function space \mathfrak{F}_i , $i=1,2$.

- i) If $\mu_2 \ll \mu_1$ then $\mathfrak{F}_1 \cap \mathfrak{F}_2$ is dense in \mathfrak{F}_2 .
- ii) If either X_1 or X_2 is reduced, and $\mathfrak{F}_1 \cap \mathfrak{F}_2 = \{0\}$, then $\mu_1 \perp \mu_2$.

Proof. i) Fix $s \in \mathfrak{F}_2$. By Proposition 1 in [24] we have

$$s(t) = E(X_2(t)Y^{\langle p-1 \rangle}) = \int_{\mathbb{F}^T} x(t)a(x)^{\langle p-1 \rangle} \mu_{X_2}(dx)$$

where $z^{\langle q \rangle} = |z|^{q-1}z$, $Y \in \mathcal{L}(X_2)$ and $a(x)$ is a representation of Y in $L_p(\mu_1) - \overline{\text{sp}}\{x(t); t \in T\} \subset \mathbb{F}^T$, $Y(\omega) = a(X(\cdot, \omega))$. Let

$$\mu_2(E) = \int_E g d\mu_1 + \mu_2(E \cap N)$$

be the Lebesgue decomposition of μ_2 with respect to μ_1 . Define

$$E_n = \{x: 0 < g(x) \leq n\} \cap N^c \text{ and}$$

$$s_n(t) = \int_{\mathbb{F}^T} x(t)a(x)^{\langle p-1 \rangle} 1_{E_n}(x) \mu_2(dx) = \int_{\mathbb{F}^T} x(t)a(x)^{\langle p-1 \rangle} g(x) 1_{E_n}(x) \mu_1(dx).$$

Since $a^{\langle p-1 \rangle} 1_{E_n} \in L_{p^*}(\mu_2)$ and $a^{\langle p-1 \rangle} g 1_{E_n} \in L_{p^*}(\mu_1)$, we have $s_n \in \mathfrak{F}_1 \cap \mathfrak{F}_2$. Also

$$|\sum_{k=1}^K c_k (s - s_n)(t_k)| \leq [\int_{\mathbb{F}^T} |\sum_{k=1}^K c_k x(t_k)|^p \mu_2]^{1/p} [\int_{\mathbb{F}^T} |a^{\langle p-1 \rangle}|^{p^*} 1_{E_n^c} d\mu_2]^{1/p^*}.$$

Thus

$$\|s - s_n\|_{\mathfrak{F}_2}^{p^*} \leq \int_{E_n^c} |a^{\langle p-1 \rangle}|^{p^*} d\mu_2 = \int_{\{g \geq n\}} |a|^p g d\mu_1 \rightarrow 0$$

as $n \rightarrow \infty$, i.e. $\mathfrak{F}_1 \cap \mathfrak{F}_2$ is dense in \mathfrak{F}_2 .

ii) For a fixed $t_0 \in T$, let $a_0(x) = x(t_0)$ and define

$$s_0(t) = \int_{\mathbb{F}^T} x(t)a_0(x)^{\langle p-1 \rangle} \mu_2(dx).$$

By Proposition 1 in [24], $s_0 \in \mathfrak{F}_2$, since $a_0(x) \in L_{p^*}(\mu_2)$. Let

$$s_{0n}(t) = \int_{\mathbb{F}^T} x(t) a_0(x)^{\langle p-1 \rangle} 1_{E_n}(x) \mu_2(dx) = \int_{\mathbb{F}^T} x(t) a_0(x)^{\langle p-1 \rangle} g(x) 1_{E_n}(x) \mu_1(dx)$$

so that $s_{0n} \in \mathfrak{F}_1 \cap \mathfrak{F}_2$. Since $\mathfrak{F}_1 \cap \mathfrak{F}_2 = \{0\}$, $s_{0n} \equiv 0$, i.e. $s_{0n}(t) = 0$ for all $t \in T$. In particular

$$s_{0n}(t_0) = \int_{\{0 < g < n\}} |x(t_0)|^p g(x) \mu_1(dx) = 0 \quad \text{for } n = 1, 2, \dots,$$

and hence

$$\int_{\{0 < g < \infty\}} |x(t_0)|^p g(x) \mu_1(dx) = 0.$$

Consequently, since $t_0 \in T$ is arbitrary, we have $x(t) = 0$ a.e. (μ_1) on $\{0 < g < \infty\}$ for each $t \in T$. But this implies that X_1 is non-reduced if

$$\mu_1(\{x: x(t) = 0, t \in T\}) \geq \mu_1(\{x; 0 < g(x) < \infty\}).$$

On the other hand if $\mu_1(\{x; 0 < g(x) < \infty\}) > 0$ then $x(t) = 0$ a.e. ($g\mu_1$) for each t and $\int_{\{0 < g < \infty\}} g d\mu_1 > 0$. Hence

$$\mu_2(\{x; x(t) = 0, t \in T_0\}) \geq \int_{\{0 < g < \infty\}} g d\mu_1 > 0,$$

i.e. X_2 is nonreduced. Since either X_1 or X_2 is reduced we must have

$$\mu_1(\{x; 0 < g(x) < \infty\}) = 0, \text{ i.e. } \mu_1 \perp \mu_2. \quad \square$$

BIBLIOGRAPHY

- [1] S. Cambanis, The measurability of a stochastic process of second order and its linear space, *Proc. Amer. Math. Soc.* 47 (1975), p. 467-475.
- [2] S. Cambanis, The relationship between the detection of sure and stochastic signals in noise, *SIAM J. Appl. Math.* 31 (1976), p. 558-568.
- [3] S. Cambanis, Complex symmetric stable variables and processes. In "Contributions to Statistics: Essays in Honour of Norman L. Johnson", P.K. Sen, Ed., North-Holland, New York (1982) p. 63-79.
- [4] S. Cambanis and A.G. Miamee, On prediction of harmonizable stable processes, *Sankhyā* (1989), to appear.

- [5] S. Cambanis and A.R. Soltani, Prediction of stable processes: Spectral and moving average representations, *Z. Wahrsch. verw. Geb.* 66 (1982), p. 593-612.
- [6] S. Chatterji and V. Mandrekar, Equivalence and singularity of Gaussian measures and applications. In "Probabilistic Analysis and Related Topics 1". A.T. Bharucha-Reid, Ed., Academic Press, New York 1978, p. 169-197.
- [7] S. Chatterji, and S. Ramaswamy, Mesures Gaussiennes et mesures produits, *Lecture Notes in Mathematics No. 920*, Springer-Verlag, Berlin, (1982), p. 570-580.
- [8] R.M. Dudley and M. Kanter. Zero-one laws for stable measures, *Proc. Amer. Math. Soc.* 45 (1974), p. 245-252.
- [9] W.H. DuMouchel, On the asymptomatic normality of the maximum likelihood estimate when sampling from a stable distribution, *Ann. Statist.* 1 (1973), p. 948-957.
- [10] J. Feldman, Equivalence and perpendicularity of Gaussian processes, *Pacific J. Math.* 9 (1958), 699-708.
- [11] W. Feller, *An Introduction to Probability Theory and Its Applications*, Vol. 2, Wiley, New York 1966.
- [12] R. Fortet, Espaces à noyau reproduisant et lois de probabilités des fonctions aléatoires, *Ann. Inst. H. Poincaré, Ser. B*, IX (1973), p. 41-48.
- [13] I. Gihman, and A. Skorohod, A., On densities of probability measures in function spaces, *Russ. Math. Surveys* 21 (1966), p. 83-156.
- [14] J. Hájek, A property of normal distribution of any stochastic process, *Czech. Math. J.* 8 (1958), p. 610-617.
- [15] C.D. Hardin, On the spectral representation of symmetric stable processes, *J. Multivariate Anal.* 12 (1982), p. 385-401.
- [16] I.A. Ibragimov and R.Z. Hasminskii, *Statistical Estimation, Asymptotic Theory*, Springer Verlag, New York 1981.
- [17] I.A. Ibragimov and Y.A. Rozanov, *Gaussian Random Processes*, Springer-Verlag, New York 1978.
- [18] S. Kakutani, On equivalence of infinite product measures, *Ann. Math.* 49 (1948), p. 239-247.
- [19] M. Kanter, Stable densities under change of scale and total variation inequalities, *Ann. Probab.* 3 (1975), p. 697-707.
- [20] M. Kanter, Equivalence singularity dichotomies for a class of ergodic measures. *Math. Proc. Cambridge Philos. Soc.* 81 (1977), p. 249-252.
- [21] J.F.C. Kingman and S.J. Taylor, *Introduction to Measure and Probability*, Cambridge University Press, London 1973.
- [22] J. Kuelbs, A representation theorem for symmetric stable processes and stable measures on H , *Z. Wahrsch. verw. Geb.* 26 (1973), p. 259-271.

- [23] L. LeCam, On the assumption used to prove asymptotic normality of maximum likelihood estimates, *Ann. Math. Statist.* 41 (1970), p. 802-828.
- [24] M. Marques and S. Cambanis, Admissible and singular translates of stable processes. In *Probability Theory on Vector Spaces IV*, S. Cambanis and A. Weron, eds., *Lecture Notes in Mathematics* 1392, Springer-Verlag, Berlin, 1989, p. 239-257.
- [25] L.A. Shepp, Distinguishing a sequence of random variables from a translate of itself, *Ann. Math. Statist.* 36 (1965), p. 1107-1112.
- [26] A. Skorohod, *Integration in Hilbert Spaces*, Springer-Verlag, Berlin 1974.
- [27] J.M. Steele, Fisher information and detection of a Euclidean perturbation of an independent stationary process, *Ann. Probab.* 14 (1986), p. 326-335.
- [28] H. Strasser, *Mathematical Theory of Statistics*, de Gruyter, Berlin 1985.
- [29] D. Thang and N. Tien, On symmetric stable measures with discrete spectral measure on Banach spaces. In "*Probability Theory and Vector Spaces II*", *Lecture Notes in Mathematics*, 828, A. Weron, Ed., Springer-Verlag, Berlin 1979, p. 286-301.
- [30] J. Zinn, Admissible translates of stable measures, *Studia Math.* 54 (1975), p. 245-257.

256. E. Mayer-Wolf, A central limit theorem in nonlinear filtering, Apr. 89.
257. C. Houdré, Factorization algorithms and non-stationary Wiener filtering, Apr. 89, Stochastics, to appear.
258. C. Houdré, Linear Fourier and stochastic analysis, Apr. 89.
259. C. Kallianpur, A line grid method in areal sampling and its connection with some early work of H. Robbins, Apr. 89, Amer. J. Math. Statist. Sci., 1989, to appear.
260. C. Kallianpur, A. G. Miaso and H. Niemi, On the prediction theory of two-parameter stationary random fields, Apr. 89, J. Multivariate Anal., 32, 1990, 120-149.
261. I. Herbst and L. Pitt, Diffusion equation techniques in stochastic monotonicity and positive correlations, Apr. 89.
262. R. Selubar, On estimation of Hilbert space valued parameters, Apr. 89, (Dissertation)
263. E. Mayer-Wolf, The noncontinuity of the inverse Radon transform with an application to probability laws, Apr. 89.
264. D. Monrad and W. Philipp, Approximation theorems for weakly dependent random vectors and Hilbert space valued martingales, Apr. 89.
265. K. Benberni and S. Cambanis, Sampling designs for estimating integrals of stochastic processes, Apr. 89.
266. S. Evans, Association and random measures, May 89.
267. H.L. Hurd, Correlation theory of almost periodically correlated processes, June 89.
268. O. Kallenberg, Random time change and an integral representation for marked stopping times, June 89, Probab. Th. Rel. Fields, accepted.
269. O. Kallenberg, Some uses of point processes in multiple stochastic integration, Aug. 89.
270. W. Wu and S. Cambanis, Conditional variance of symmetric stable variables, Sept. 89.
271. J. Mijbheer, U-statistics and double stable integrals, Sept. 89.
272. O. Kallenberg, On an independence criterion for multiple Wiener integrals, Sept. 89.
273. C. Kallianpur, Infinite dimensional stochastic differential equations with applications, Sept. 89.
274. C.W. Johnson and C. Kallianpur, Homogeneous chaos, p-forms, scaling and the Feynman integral, Sept. 89.
275. T. Hida, A white noise theory of infinite dimensional calculus, Oct. 89.
276. K. Benberni, Sample designs for estimating integrals of stochastic processes, Oct. 89, (Dissertation)
277. I. Rychlik, The two-barrier problem for continuously differentiable processes, Oct. 89.
278. C. Kallianpur and R. Selubar, Estimation of Hilbert space valued parameters by the method of sieves, Oct. 89.
279. C. Kallianpur and R. Selubar, Parameter estimation in linear filtering, Oct. 89.
280. P. Bloomfield and H.L. Hurd, Periodic correlation in stratospheric ozone time series, Oct. 89.
281. J.M. Anderson, J. Horowitz and L.D. Pitt, On the existence of local times: a geometric study, Jan. 90.
282. C. Lindgren and I. Rychlik, Slepian models and regression approximations in crossing and extreme value theory, Jan. 90.
283. H.L. Koul, M-estimators in linear models with long range dependent errors, Feb. 90.
284. H.L. Hurd, Almost periodically unitary stochastic processes, Feb. 90.
285. M.R. Leadbetter, On a basis for 'Peaks over Threshold' modeling, Mar. 90.
286. S. Cambanis and E. Masry, Trapezoidal stratified Monte Carlo integration, Mar. 90.
287. M. Marques and S. Cambanis, Dichotomies for certain product measures and stable processes, Mar. 90.
288. M. Maejima and Y. Morita, Trimmed sums of mixing triangular arrays with stationary rows, Mar. 90.
289. S. Cambanis and M. Maejima, Characterizations of one-sided linear fractional Lévy motions, Mar. 90.
290. N. Kono and M. Maejima, Hölder continuity of sample paths of some self-similar stable processes, Mar. 90.