		N/ DA	NTIN		(IGRA Approved
KEPUKI D		IN PAGE		ILL (0118 No. 0704-0184
a. REPORT SECURITY CLASSIFICATION		15. RESTRICTIVE MARKINGS			
UNCLASSIFIED					
SETURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION / AVAILABILITY OF REPORT			
b. DECLASSIFICATION / DOWNGRADING SCHEDU	.E	- Approved for Public Release; (\mathcal{R})			
N/A		Distribution Unlimited			
. PERFORMING ORGANIZATION REPORT NUMBER	R(S)	5. MONITORING ORGANIZATION REPORT NUMBER(S)			
Technical Report No. 298		AFOSR-TR. 90 0804			
a. NAME OF PERFORMING ORGANIZATION	6b. OFFICE SYMBOL	7a. NAME OF MONITORING ORGANIZATION			
Center for Stochastic Processes	,	AFOSR/NM			
C ADDRESS (City State and 710 Code)		Th ADDRESS (P)	W State and TH	اطمم)	
Statistics Department CB #3260, Phillips Hall		Bldg, 410			
		Bolling	Air Force	Base. Dr	20332-6448
Chapel Hill, NC 27599-3260				.,	
a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR NM		9. PROCUREMEN	T INSTRUMENT I	DENTIFICATIO	ON NUMBER
		FA 96 20 0	35C 0144		
		1.43070 \$	JU VI44	•	
c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF F	UNDING NUMBE	RS	
Bldg. 410		PROGRAM ELEMENT NO	PROJECT	TASK NO	WORK UNIT
Bolling AFB, DC 20332-6448		6.1102F	2304	<u> </u>	-CCESSION P
1. TITLE (Include Convinu Chapilinaian)		1	L	T H2	
Dynamic include security classifications	McKoon 1/1 - com	oustion in d	uale of mu-	lean and	202
riopayation of thats and ten t	nunedii=¥1850¥ t	quarion in a	uurs ur NUC	vicai shg	1663
2. PERSONAL AUTHOR(S)					
Chiang, T.S., Kallianpu	r, G. and Sunda	r, P.			
3a. TYPE OF REPORT 13b. TIME CO	VERED				
_wonwint _	VERED	14. DATE OF REPO	RT (Year, Month	, Day) 15.	PAGE COUNT
preprint FROM	TO	14. DATE OF REPO 1990, Maj	RT (Year, Month Y	, Day) 15. 1	PAGE COUNT 43
preprint FROM	TO	14. DATE OF REPO 1990, Maj	RT(Year, Month y	i, Day) 15. 1	PAGE COUNT 43
preprint FROM 6. SUPPLEMENTARY NOTATION None.	TO	14. DATE OF REPO 1990, Maj	RT (Year, Month Y	i, Day) 15. i	PAGE COUNT 43
preprint FROM		14. DATE OF REPO 1990, Ma	RT (Year, Month Y	, Day) 15. 1	PAGE COUNT
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP	18. SUBJECT TERMS	Continue on revers	RT (Year, Month y e if necessary an hastic diff	, Day) 15. 1 Id identify b	PAGE COUNT 43 y block number) equation
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP	18. SUBJECT TERMS Nuclear space McKean-Vlasov	Continue on revers valued stoc	RT (Year, Month y e if necessary an hastic diff ropagation	, Day) 15. Ind identify b ferential of chaos	PAGE COUNT 43 y block number) equation, 5.
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS Nuclear space McKean-Vlasov	14. DATE OF REPO 1990, Ma Continue on revers valued stoc equation, p	RT (Year, Month Y e if necessary an hastic diff ropagation	, Day) 15. Ind identify b ferential of chaos	PAGE COUNT 43 y block number) equation, 5.
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS Nuclear space McKean-Vlasov	Continue on reverse valued stoc equation, p	RT (Year, Month y e if necessary an hastic diff ropagation	<i>d identify b</i> ferential of chaos	PAGE COUNT 43 y block number) equation, 3.
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS (Nuclear space McKean-Vlasov	14. DATE OF REPO 1990, Ma Continue on reverse valued stoc equation, p	RT (Year, Month y e <i>if necessary an</i> hastic diff ropagation	<i>d identify b</i> ferential of chaos	PAGE COUNT 43 y block number) equation, 3.
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP SUB-GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Indidentify by block r	14. DATE OF REPO 1990, Ma Continue on reverse valued stoc vequation, p	RT (Year, Month Y e if necessary an hastic diff ropagation	<i>d identify b</i> ferential of chaos	PAGE COUNT 43 y block number) equation, 5.
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXX 9. ABSTRACT (Continue on reverse if necessary a DTIC	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block r Please see ov	14. DATE OF REPO 1990, Ma Continue on revers valued stoc equation, p number)	RT (Year, Month y e if necessary an hastic diff ropagation	<i>d identify</i> by erential of chaos	PAGE COUNT 43 y block number) equation,
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block r Please see ov	14. DATE OF REPO 1990, Ma Continue on revers valued stoc equation, p humber)	RT (Year, Month y e <i>if necessary an</i> hastic diff ropagation	<i>d identify b</i> erential of chaos	PAGE COUNT 43 y block number) equation, ;.
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXX 9. ABSTRACT (Continue on reverse if necessary a DTIC ELECTE	18. SUBJECT TERMS (Nuclear space McKean-Vlasov and identify by block r Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stoc vequation, p number)	RT (Year, Month y e <i>if necessary an</i> hastic diff ropagation	<i>d identify b</i> erential of chaos	PAGE COUNT 43 <i>y block number)</i> equation,
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS (Nuclear space McKean-Vlasov and identify by block r Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse Valued stoc vequation, p	RT (Year, Month y e if necessary an hastic diff ropagation	, Day) 15. Id identify by erential of chaos	PAGE COUNT 43 y block number) equation,
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS (Nuclear space McKean-Vlasov and identify by block r Please see ov	14. DATE OF REPO 1990, Ma Continue on revers valued stoc vequation, p	RT (Year, Month y e if necessary an hastic diff ropagation	<i>d identify b</i> erential of chaos	PAGE COUNT 43 / block number) equation,
preprint FROM	18. SUBJECT TERMS (Nuclear space McKean-Vlasov and identify by block r Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stoc vequation, p	RT (Year, Month y e <i>if necessary an</i> hastic diff ropagation	<i>, Day)</i> 15. Id <i>identify b</i> erential of chaos	PAGE COUNT 43 <i>y block number)</i> equation,
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXX 9. ABSTRACT (Continue on reverse if necessary a DTIC JUL 2 6 1990 D COSATI CODES COSATI CODES FIELD GROUP XXXXXXXXXXXXXXX 9. ABSTRACT (Continue on reverse if necessary a DTIC JUL 2 6 1990 COSATI CODES COSATI CODES COSATI CODES FIELD GROUP XXXXXXXXXXXXXXXX 9. ABSTRACT (Continue on reverse if necessary a COSATI CODES COSATI CODES FIELD GROUP XXXXXXXXXXXXXXXXXXXXX 9. ABSTRACT (Continue on reverse if necessary a COSATI CODES COSATI CODES FIELD GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block n	14. DATE OF REPO 1990, Ma Continue on reverse valued stock vequation, p number) ver.	RT (Year, Month y e <i>if necessary an</i> hastic diff ropagation	, Day) 15. Id identify by Perential of chaos	PAGE COUNT 43 <i>y block number)</i> equation,
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXX 9. ABSTRACT (Continue on reverse if necessary a DTIC SUPPLEMENTARY NOTATION NONE.	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block of Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stock vequation, p number) ver.	RT (Year, Month y e if necessary an hastic diff ropagation	, Day) 15. Ind identify by Perential Of chaos	PAGE COUNT 43 <i>y block number)</i> equation,
preprint FROM	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block of Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stock vequation, p number) ver.	RT (Year, Month y e if necessary an hastic diff ropagation	, Day) 15. Id identify by Ferential of chaos	PAGE COUNT 43 / block number) equation,
preprint FROM	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block in Please see ov	14. DATE OF REPO 1990, Ma; Continue on reverse valued stock vequation, p number) ver.	RT (Year, Month y e <i>if necessary an</i> hastic diff ropagation	, Day) 15. Id identify by erential of chaos	PAGE COUNT 43 <i>y block number)</i> equation,
preprint FROM	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block of Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stock vequation, p number) ver.	RT (Year, Month y e <i>if necessary an</i> hastic diff ropagation	, Day) 15. Id identify by Terential of chaos	PAGE COUNT 43 / block number) equation,
preprint FROM	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block of Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stoc vequation, p wmber) ver.	RT (Year, Month y e if necessary an hastic diff ropagation	, Day) 15. Ind identify by Perential Of chaos	PAGE COUNT 43 / block number) equation,
preprint FROM	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block of Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stoc vequation, p umber) ver. 21. ABSTRACT SE UNCLASSIFI	RT (Year, Month y e if necessary an hastic diff ropagation	(, Day) 15. Id identify by Ferential of chaos	PAGE COUNT 43 / block number) equation,
preprint FROM	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block m Please see ov	 14. DATE OF REPO 1990, Ma; Continue on reverse valued stock equation, p iumber) ver. 21. ABSTRACT SEU UNCLASSIFI 22b. TELEPHONE ((202) 767-55 	RT (Year, Month y e if necessary an hastic diff ropagation CURITY CLASSIFIC ED Include Area Cod	(, Day) 15. Id identify by erential of chaos (ATION (e) 22c. OFF AEOS	PAGE COUNT 43 / block number) equation,
preprint FROM 6. SUPPLEMENTARY NOTATION None. 7. COSATI CODES FIELD GROUP SUB-GROUP XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	18. SUBJECT TERMS (Nuclear space McKean-Vlasov Ind identify by block m Please see ov	14. DATE OF REPO 1990, Ma Continue on reverse valued stock vequation, p wmber) ver. 21. ABSTRACT SEC UNCLASSIFI 225. TELEPHONE ((202) 767-50 obsolute	RT (Year, Month y e if necessary an hastic diff ropagation CURITY CLASSIFIC Include Area Cod 26	(, Day) 15. Id identify by erential of chaos (ATION (e) 22c. OFF AFOSI (c) AFOSI	PAGE COUNT 43 / block number) equation, ICE SYMBOL R/NM FIGH OF FUE

<u>Abstract</u>: In this paper, we consider a system of interacting diffusion processes taking values in the dual of nuclear spaces:

$$dX_{j}^{n}(t) = (a(t, X_{j}^{n}(t)) + \frac{1}{n} \sum_{i=1}^{n} b(t, X_{j}^{n}(t), X_{i}^{n}(t))dt + (\sigma((t, X_{j}^{n}(t)) + \frac{1}{n} \sum c(t, X_{j}^{n}(t), X_{i}^{n}(t))dW_{t}^{j} \quad j=1,...,n.$$

We prove that under suitable conditions the system has a unique solution and its empirical distributions will converge as $n \rightarrow \infty$ to the solution of the corresponding McKean-Vlasov equation. An application to a neurophysiological model is also given.

CENTER FOR STOCHASTIC PROCESSES

.

Department of Statistics University of North Carolina Chapel Hill, North Carolina



PROPAGATION OF CHAOS AND THE MCKEAN-VLASOV EQUATION

IN DUALS OF NUCLEAR SPACES

by

T.S. Chiang

G. Kallianpur

and

P. Sundar

Technical Report No. 298

May 1990

90 07 25 076

Propagation of Chaos and the McKean-Vlasov Equation

in Duals of Nuclear Spaces

T.S. ChiangG. KallianpurP. SundarInstitute of MathematicsCenter for Stochastic ProcessesDepartment of MathematicsAcademia SinicaDepartment of StatisticsLouisiana State UniversityTaipei, TaiwanUniversity of North Carolina
Chapel Hill, NC 27599-3260Baton Rouge, LA 70803

<u>Abstract</u>: In this paper, we consider a system of interacting diffusion processes taking values in the dual of nuclear spaces:

$$dX_{j}^{n}(t) = (a(t,X_{j}^{n}(t)) + \frac{1}{n} \sum_{i=1}^{n} b(t,X_{j}^{n}(t), X_{i}^{n}(t))dt + (\sigma((t,X_{j}^{n}(t)) + \frac{1}{n} \sum c(t,X_{j}^{n}(t),X_{i}^{n}(t))dW_{t}^{j} \quad j=1,...,n.$$

We prove that under suitable conditions the system has a unique solution and its empirical distributions will converge as $n \rightarrow \infty$ to the solution of the corresponding McKean-Vlasov equation. An application to a neurophysiological model is also given.

<u>Key words</u>: Nuclear space valued stochastic differential equation, McKean-Vlasov equation, propagation of chaos.

AMS Classification (1985): 60H20, 60B05, 60B10, 60J70

Research supported by the Air Force Office of Scientific Research Contract No. F49620 85C 0144.

This author would like to thank the Center for Stochastic Processes for their financial support while doing this research.

§1. INTRODUCTION

The present paper is concerned with propagation of chaos problems for systems with an infinite number of degrees of freedom such as strings or spatially extended neurons. The investigation of the asymptotic behavior of the voltage (membrane) potentials of large assemblages of interacting neurons leads to precisely such problems and provided the immediate motivation for the work. Another example to which the approach of the present paper could be applied (we believe) is the Ginsburg-Landau model in hydrodynamics, recently studied by T. Funaki [4].

1

Sections 2 and 3 are of an introductory nature. Basic properties of duals of nuclear spaces (denoted throughout by Φ^i , the strong dual of a countably Hilbertian nuclear space Φ) are briefly discussed and the results of Kallianpur et al. [8] on the existence and uniqueness of the solution to (the martingale problem posed by) a Φ^i -valued stochastic differential equation (SDE) is extended to a system of such equations. The principal results in which the infinite dimensionality of our problem call for special arguments are derived, in Sections 3, 4, and 5.

In Theorem 4.1, the weak compactness of the sequence of empirical measures $\mu_{n}(\omega, \cdot) = \frac{1}{n} \sum_{j=1}^{n} \delta_{j=1} (\cdot, \omega) \quad (\cdot) \text{ is established and it is shown in Section 6 (Theorem 6.1) that } \eta_{n}, \text{ the law of } \mu_{n}(\omega, \cdot) \text{ converges weakly to the unique solution of the McKean-Vlasov equation.}$

The infinite dimensional (nucelar space-valued) version of the McKean-Vlasov SDE is introduced in Section 5. The existence and uniqueness of solution of this equation is investigated in detail in Baldwin et al. [1]. In view of the importance of this result for the propagation of chaos, a slightly different proof (with a somewhat stronger conclusion) is given for the special

jes

Avail and/or

Special

Dist

choice of the interaction term for our problem. The main results on the propagation of chaos are given in Theorems 6.2 and 6.3. The existence of a unique solution to the martingale problem posed by the system (3.2.1) and Theorem 5.1 on the McKean-Valsov equation are the key steps that enable Sznitman's technique for finite dimensional SDE's to be used for the nuclear space valued case.

The application, alluded to above, to the voltage potentials of interacting, spatially extended neurons is considered in Section 7. For reasons of space we have limited ourselves to the mathematics of the problem and excluded any discussion of the neurophysiological implications.

In Section 8 we introduce the assumption that the initial measure of the system (3.2.1) is μ_0 -chaotic and show that the results of the previous sections hold under this more general condition. This is of importance in application since it is more reasonable to assume (as in the case of the neurons) that the random variables $X_1^n(t), \ldots, X_n^n(t)$ are exchangeable than that they are identically distributed.

It is worth remarking that our results contain the finite dimensional results as a particular case and their relationship with other available results (e.g. Sznitman [16]) is also briefly commented upon.

An outstanding problem, to which we hope to return in a later paper, is that of proving a fluctuation or central limit theorem. The difficulties that lie ahead are foreshadowed in a recent paper by Kallianpur and Mitoma [7] that establishes such a result under restrictive conditions.

<u>\$2. PRELIMINARIES ON NUCLEAR SPACES AND \$-VALUED SDE's.</u>

In this section we provide the basics on nuclear spaces and on stochastic processes and integrals taking values in duals of nuclear spaces followed by

the results of Kallianpur, Mitoma and Wolpert [8] on the existence and uniqueness of solutions of SDE's.

2.1 Nuclear spaces

Let Φ be a real linear space whose topology is given by an increasing sequence $\|\cdot\|_{r}$. r=1,2,... of Hilbertian norms. Let Φ_{r} be the completion of Φ with respect to $\|\cdot\|_{r}$. Then Φ is called a countably Hilbertian nuclear space (CHNS) if the following two conditions are satisfied:

(i)
$$\Phi = \bigcap_{r=1}^{\omega} \Phi_r$$

(ii) For each r, there exists an m>r such that the canonical embedding $\Phi_{m} \subset \Phi_{r}$ is Hilbert-Schmidt.

Let Φ ' denote the strong dual of Φ whose topology is given by the following family of semi-norms:

 $|f|_{B} = \sup_{x \in B} |f(x)|$ where $B \subseteq \Phi$ is a bounded set in Φ .

It is well known that $\Phi' = \bigcup_{r=1}^{\infty} \Phi_{-r}$ where Φ_{-r} is the dual of Φ_{r} . Besides, the strong topology on Φ' coincides with the inductive limit topology induced by the canonical embeddings $\Phi_{-r} \subset \Phi'$. Let $\|\cdot\|_{-r}$ denote the norm in Φ_{-r} . If j_{r} denotes the canonical mapping of Φ_{r} onto its dual Φ_{-r} , then for $u \in \Phi_{-r}$ and $\phi \in \Phi_{r}$.

$$u[\phi] = \langle u, j_{\mu}\phi \rangle_{\mu} = \langle j_{\mu}, u, \phi \rangle_{\mu}$$

where $\langle \cdot \rangle$ denotes the inner product in the appropriate space.

For any T>O, C_{Φ}^{T} , denotes the space of all continuous functions from [O,T] to Φ' . If $\{|\cdot|_{\alpha} : \alpha \in A\}$ is the set of semi-norms defining the strong topology of Φ' , then by defining $\max_{\alpha} = \sup_{0 \leq t \leq T} |x_t|_{\alpha}, x \in C_{\Phi}^{T}$, the space C_{Φ}^{T} , is seen as a completely regular topological space under the projective limit topology of $\{\mathbf{m} \cdot \mathbf{m}_{\alpha} : \alpha \in A\}$. C_{ϕ} , denotes the space of all continuous functions from $[0,\infty)$ to ϕ' . $C_{\phi-\varrho}^{T}$ is the Banach space with the uniform topology, consisting of all continuous functions from [0,T] to $\phi_{-\rho}$.

2.2 ϕ' -valued processes.

Let $(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P})$ be a stochastic basis satisfying the usual hypotheses.

<u>Definition</u>: An adapted Φ' -valued stochastic process $\{M_t\}_{t\geq 0}$ is called a martingale with respect to (\mathcal{F}_t) if for each $\phi \in \Phi$, $\{M_t[\phi]\}_{t\geq 0}$ is a real-valued (\mathcal{F}_t) martingale. $\{M_t\}$ is called an L²-martingale if $EM_t[\phi]^2 < \infty$ for all $t\geq 0$ and $\phi\in\Phi$.

For a detailed discussion of Φ '-valued martingales and their properties we refer the reader to [6] and [12].

<u>Definition</u>: A continuous Φ' -valued process $\{W_t\}_{t\geq 0}$ is called a Wiener process with covariance Q, if the following conditions are satisfied:

(i) ₩₀=0 a.s.

(ii) $\{W_t[\Phi]\}\$ is a one-dimensional Wiener process with variance parameter $Q(\phi,\phi)$, where $Q(\cdot,\cdot)$ is a continuous positive definite symmetric bilinear form on Φ .

A result of Mitoma [13] implies that any Φ' -valued Wiener process W has paths that lie in the Banach space C_{Φ} for some $q \langle \infty, \text{ and which are continuous} \\ -q$ in the Φ_{-q} -topology P-a.s. The choice of q depends only on the covariance form Q. Let $r \geq q$ be a fixed integer. An important property of the quadratic form Q is that it admits a unique continuous extension to a nuclear form on Φ_r and

$$Q[\phi, \psi] = (\phi, Q_{r}\psi)_{r} = (Q_{r}^{\nu}\phi, Q_{r}^{\nu}\phi)_{r} \qquad (2.2.1)$$

for a unique non-negative trace-class operator Q_r on Φ_r . The trace norm of Q on Φ_r (or, equivalently, of Q_r^{*}) is given by

$$|Q|_{-r,-r} = \sum_{j} \sum_{j} \sum_{j} [h_{j}^{r}, h_{j}^{r}] \qquad (2.2.2)$$

where $\{h_j^r\}$ is a CONS for Φ_r .

2.3 Stochastic integrals in Φ'

Let $\{W_t\}$ be a Φ' -valued Wiener process with covariance form Q(•,•) and let $L(\Phi',\Phi')$ be the space of all continuous linear operators from Φ' to Φ' .

For each T>O and $\phi \in \Phi$, let L^2_W denote the space of progressively measurable processes H: $\mathbb{R}_+ \times \Omega \longrightarrow L(\Phi', \Phi')$ for which $E \int_0^T Q[H_s^*\phi, H_s^*\phi] ds < \infty$, where H_s^* is the operator dual to H_s .

<u>Definition</u>: The stochastic integral $I_t^H := \int_0^t H_s dW_s$ ($0 \le s \le T$) is a Φ' -valued L^2 -martingale with the quadratic variation process as $\langle I^H \rangle_t [\phi, \psi] = \int_0^t Q_{H_s}[\phi, \psi] ds$ where $Q_{H_s}[\phi, \psi] = Q[H_s^*\phi, H_s^*\psi]$.

There exists an $\ell > 0$, depending on H and T such that $I_{\cdot}^{H} \in C_{\Phi_{-\ell}}^{T}$ a.s.. If $\{h_{j}^{\ell}\} \subset \Phi$ is any CONS in $\Phi_{-\ell}$.

$$I_{t}^{H}[\phi] = \int_{0}^{t} H_{s} dW_{s}[\phi] = \sum_{j=1}^{\infty} \int_{0}^{t} (H_{s}^{\star}\phi, h_{j}^{\ell})_{\ell} dW_{s}[h_{j}^{\ell}]$$
(2.3.1)

where the right hand side is an L^2 -convergent series of Itô integrals. Besides,

$$\langle \mathbf{I}^{H} \rangle_{t} [\phi, \psi] = \sum_{i, j=1}^{\infty} \int_{0}^{t} (\mathbf{H}^{*}_{s}\phi, \mathbf{h}^{\ell}_{i})_{\ell} (\mathbf{H}^{*}_{s}\psi, \mathbf{h}^{\ell}_{j}) ds \ Q[\mathbf{h}^{\ell}_{i}, \mathbf{h}^{\ell}_{j}]$$
(2.3.2)

2.4 Φ' -valued SDE's.

We give below the result of Kallianpur, Mitoma and Wolpert [8] on the existence and uniqueness of solutions of stochastic differential equations.

For a probability measure μ_0 on Φ' and a pair of functions A: $\mathbb{R}_+ \times \Phi' \longrightarrow \Phi'$ and B: $\mathbb{R}^+ \times \Phi' \longrightarrow L(\Phi':\Phi')$, consider the following SDE:

$$dX_{t} = A(t, X_{t})dt + B(t, X_{t})dW_{t}$$

$$X_{0} = X(0)$$
(2.4.1)

where X_0 is a Φ' -valued random variable with law of X_0 given by μ_0 , and W is a Φ' -valued Wiener process with covariance form Q.

Let $\mathfrak{D}_{b}^{2}(\Phi')$ consist of all functions $f: \Phi' \to \mathbb{R}$ with $f(u) = \widetilde{f}(u[\phi])$ where $\widetilde{f} \in C_{b}^{2}(\mathbb{R})$ and $\phi \in \Phi$. The operator L_{s} is defined as follows: For each $f \in \mathfrak{D}_{b}^{2}(\Phi')$,

$$L_{s}f(u) = \widetilde{f}'(u[\phi])A(s,u)[\phi] + \frac{1}{2}\widetilde{f}''(u[\phi])Q(B^{*}(s,u)\phi,B^{*}(s,u)\phi)$$

where $B^{\star}(s,u) : \Phi \longrightarrow \Phi$ is the adjoint of B(s,u), i.e., for all $v \in \Phi'$ and $\phi \in \Phi$,

$$v[B^{\bigstar}(s,u)\phi] = B(s,u)v[\phi].$$

Let $\Omega = C_{\phi}$, \mathscr{F}_{t} = Borel σ -algebra of C_{ϕ}^{t} , and $\mathscr{F} = V \mathscr{F}_{t}$ and let $\mathfrak{T}_{t} := C_{\phi}, \longrightarrow \phi'$ be the canonical process defined by $\mathfrak{T}_{t} y = y(t)$ for all $y \in C_{\phi}$. If $\mu \in \mathfrak{T}(C_{\phi_{-k}}^{T})$, then $\mu \mathfrak{T}_{t}^{-1}(A) = \mu(y \in C_{\phi_{-k}}^{T} : y_{t} \in A)$, $A \in \mathfrak{F}(\Phi_{-k})$.

<u>Definition</u>: A solution to the martingale problem posed by (2.4.1) is a probability measure μ on C_{ϕ} , such that for any $f \in \mathfrak{D}_{b}^{2}(\Phi')$, the real-valued process $\mathbb{M}_{t}^{f} = f(x_{t}) - f(x_{0}) - \int_{0}^{t} \mathbb{L}_{s}f(x_{s}) ds$ is a $(\Omega, \mathfrak{F}, (\mathfrak{F}_{t}), \mu)$ martingale with $\mu \cdot \pi_{0}^{-1} = \mu_{0}$.

The following conditions are imposed on the space Φ , measure μ_0 and the coefficients A and B.

Let $\{h_j^m\}$ be a CONS in Φ_m obtained by the Gram-Schmidt process applied to a countable dense set $\{\xi_i\}$ in Φ . For every j, we then have

$$\boldsymbol{\xi}_{j} = \sum_{k=1}^{n} \alpha_{mk} \mathbf{h}_{k}^{m} + \eta_{j} \qquad (2.4.2)$$

where n_j (depending on m and j) \leq j and $\|\eta_j\|_m = 0$. Our assumption is the following:

(A) For each m and p, $(p \ge m)$, in (2.4.2) $\|\ln_j\|_p = 0$.

Let T>O be fixed. Then, for each sufficiently large m \ge r where r is introduced in (2.2.1), there exists a number $\theta>O$ and an index p \ge m such that for all s,t \le T,

(IC) Initial Condition:
$$c_0 := \int_{\Phi} (1 + \|u\|_{-m}^2) [\ell_n(3 + \|u\|_{-m}^2)]^2 \mu_0(du) < \infty$$

(CC) Coercivity Condition: For each
$$u \in j_m \Phi$$
,
 $2A(t,u)[j_mu] + |Q_{B(t,u)}|_{-m,-m} \leq \theta(1+||u||_{-m}^2)$
where j_m denotes the canonical map from Φ_m to Φ_{-m} , with j_{-m} as its inverse.

(LG) Linear Growth Condition: If $u \in \Phi_{-m}$, then $A(t,u) \in \Phi_{-p}$ and $\|A(t,u)\|_{-p}^{2} \leq \theta(1+\|u\|_{-m}^{2})$ $|Q_{B(t,u)}|_{-m,-m} \leq \theta(1+\|u\|_{-m}^{2})$

(JC) Joint Continuity Condition: A and B are each jointly continuous. Further,

- (i) $B(s,u)(v) \in \Phi_{m}$ if $u, v \in \Phi_{m}$ and
- (ii) $Q_{B(s,u)}(\phi,\phi)$ is continuous in u on ϕ' for each $\phi \in \phi$.

The following condition will be needed in the proof of uniqueness.

(MC) Monotonicity Condition: For all $u, v \in \Phi_{-m} (\subset \Phi_{-p})$ (A(t,u) - A(t,v), u-v)_p + $|Q_{B(t,u)-B(t,v)}|_{-p,-p} \leq \theta ||u-v||_{-p}^2$ We give below the main result of Kallianpur et al. [8].

<u>Theorem 2.4.1.</u> Assuming conditions (A), (IC), (CC), (LG) and (JC), there exists a weak solution to the stochastic differential (2.4.1). Besides, it has the pathwise uniqueness property if ((MC) is satisfied.

Next, we give a moment bound, followed by a tightness result both of which are due to Baldwin et al [1].

<u>Theorem 2.4.2.</u> Let $k \ge 1$ and $E||X_0||_{-m}^{2k} \le c_k \le \infty$. Then, under all the conditions of Theorem 2.4.1,

$$\underset{0 \leq t \leq T}{\sup} \|X_{s}\|_{-m}^{2k} \leq (2C_{k} + 1) \exp((136k^{2} - 4k)\theta T) - 1$$

<u>Remark 2.4.1</u>: $E_{t} = \frac{2k}{t} \leq (2C_{k}+1)e^{4k(k-1)\theta t} - 1$ for each $0 \leq t \leq T$.

Theorem 2.4.3: Assume that the coefficients associated with the equations

$$X_{t}^{n} = X_{0}^{n} + \int_{0}^{t} A^{n}(s, X_{s}^{n}) ds + \int_{0}^{t} B^{n}(s, X_{s}^{n}) dW_{s}^{n}$$

and

$$X_{t} = X_{0} + \int_{0}^{t} A(s, X_{s}) ds + \int_{0}^{t} B(s, X_{s}) dW_{s}$$

satisfy the conditions:

1) Conditions (IC), (CC), (LG), (JC) and (MC) hold as stated where the constants and indices are independent of n.

2) $X_{o}^{n} \xrightarrow{D} X_{O}$

3) If Q_n and Q denote the covariance forms of $\{W_s^n\}$ and $\{W_s\}$ respectively, then Q_n converges to Q.

4) For each $s \in [0,T]$ and $\phi \in \Phi$, $A^n(s, \cdot)[\phi]$ converges continuously to $A(s, \cdot)[\phi]$. 5) For each $s \in [0,T]$ and $\phi \in \Phi$, $(B^n(s, \cdot))^* \phi$ converges continuously to $B^*(s, \cdot)\phi$.

Then $P \cdot (X^n)^{-1} \Rightarrow P \cdot X^{-1}$ in $C_{\Phi_{-p}}^T$

<u>Remark 2.4.2</u>: If T>O is fixed, the solution of SDE (2.4.1) namely X will have paths in $C_{\Phi-p}^{T}$ a.s. where p is the index that appears in the conditions.

<u>Remark 2.4.3.</u> Throughout the paper, the notation \xrightarrow{D} is used to denote convergence in distribution of random variables whereas the notation => is used to denote weak convergence of measures. Thus, $X_n \xrightarrow{D} X$ is equivalent to $P \cdot (X_n)^{-1} => P \cdot X^{-1}$, where PX_n^{-1} and PX^{-1} denote the law of X_n and X respectively. We adhere to this notation even when the random variables are measure-valued.

3. SYSTEMS OF ϕ' -VALUED SDE's

The aim of this section is to extend the results of the previous section to a system of stochastic differential equations which is done by first introducing the Cartesian product of nuclear spaces.

3.1 Cartesian product of nuclear spaces.

Let Φ denote the nuclear space introduced in Section 2. Consider the linear space $\Phi \times \Phi$ with coordinatewise linear operations. Let

$$\|(\phi_1 \times \phi_2)\|_r^2 = \|\phi_1\|_r^2 + \|\phi_2\|_r^2 \text{ for } r \ge 1.$$
 (3.1.1)

An increasing sequence of Hilbertian norms is thus defined on $\Phi \times \Phi$ which preserves nuclearity of $\Phi \times \Phi$. To see this, let $\Phi_{r \times r}$ be the completion of $\Phi \times \Phi$ in the product r-norm given by (3.1.1) for all $r \ge 1$. Clearly, $\Phi_{r \times r} = \Phi_r \times \Phi_r$ and $\Phi \times \Phi = \bigcap_{r \ge 1} (\Phi_r \times \Phi_r)$. Given n>0, if m>n such that the canonical injection $r \ge 1$ i: $\Phi_m \subset \Phi_n$ is Hilbert-Schmidt, then the injection $(\Phi \times \Phi)_m \subset (\Phi \times \Phi)_n$ is also Hilbert-Schmidt.

Let $(\Phi \times \Phi)'$ denote the strong dual of $\Phi \times \Phi$ so that $(\Phi \times \Phi)' = \bigcup_{r \ge 1} (\Phi_r \times \Phi_r)'$. If $\ell \in (\Phi_r \times \Phi_r)'$, we can uniquely determine two linear functionals ℓ_1 and ℓ_2 in Φ_{-r} such that

$$\ell(\phi_1,\phi_2) = \ell_1(\phi_1) + \ell_2(\phi_2) \quad \text{for all } \phi_1,\phi_2 \in \phi_r \quad (3.1.2)$$

Likewise, given ℓ_1 and ℓ_2 in Φ_{-r} , there exists a unique ℓ in $(\Phi_r \times \Phi_r)'$ such that (3.1.2) is satisfied. In short, there exists an isomorphism between $(\Phi_r \times \Phi_r)'$ and $\Phi_{-r} \times \Phi_{-r}$ which is written as $(\Phi_r \times \Phi_r)' \cong \Phi_{-r} \times \Phi_{-r}$.

The above isomorphism is not just algebraic but topological as well, if $\Phi_{-r} \times \Phi_{-r}$ is equipped with the product -r norm. i.e. $\|(u_1 \times u_2)\|_{-r}^2 = \|u_1\|_{-r}^2 + \|u_2\|_{-r}^2$. We thus get

$$(\Phi \times \Phi)' = \bigcup_{r \ge 1} (\Phi_r \times \Phi_r)' \cong \bigcup_{r \ge 1} (\Phi_{-r} \times \Phi_{-r})$$
(3.1.3)

Besides, $\bigcup (\Phi_{-r} \times \Phi_{-r}) = \Phi' \times \Phi'$ set-theoretically. To see the topological $r \ge 1$ equivalence, consider a neighbourhood of zero in $\Phi' \times \Phi'$ i.e. Let A_1 and A_2 be two bounded sets in Φ , and $\epsilon > 0$ be given. Consider the set

$$\Lambda = ((\ell_1, \ell_2): \sup_{\phi \in A_1} |\ell_1(\phi)| < \epsilon, \sup_{\psi \in A_2} |\ell_2(\psi) < \epsilon) \subseteq \phi \times \phi'.$$

For any fixed $r \ge 1$, $\exists a_r \rightarrow A_i \subseteq \{\phi: \|\phi\|_r < a_r\}$ i=1,2 so that

$$\Lambda \supseteq ((\ell_1, \ell_2): \|\ell_1\|_{-r} < \frac{\epsilon}{a_1}, \|\ell_2\|_{-r} < \frac{\epsilon}{a_1})$$

Thus

$$\Omega \supseteq ((\ell_1, \ell_2): \|\ell_1\|_{-r}^2 + \|\ell_2\|_{-r}^2 < \epsilon^2 / a_1^2)$$
(3.1.4)

On the other hand,

$$\Lambda \subseteq \{(\ell_1, \ell_2): \sup_{\substack{(\phi_1, \phi_2) \in A_1 \times A_2}} |\ell_1(\phi_1) + \ell_2(\phi_2)| < 2\epsilon\}$$
(3.1.5)

and $A_1 \times A_2$ is a bounded set in $\Phi \times \Phi$. (3.1.4) and (3.1.5) give us the topological equivalence of $\bigcup (\Phi_r \times \Phi_r)$ and $\Phi' \times \Phi'$. Therefore (3.1.3) implies $r \ge 1$ that

$$(\Phi \times \Phi)' = \bigcup_{r \ge 1} (\Phi_r \times \Phi_r)' \cong \bigcup_{r \ge 1} (\Phi_{-r} \times \Phi_{-r}) = \Phi' \times \Phi'.$$
(3.1.6)

Equation (3.1.6) carries over for any finite number of Cartesian products.

3.2 System of SDE's

Let X_j^n be Φ '-valued processes, $1 \leq j \leq n$ governed by the SDE

$$dX_{j}^{n}(t) = (a(t,X_{j}^{n}(t)) + \frac{1}{n} \sum_{i=1}^{n} b(t,X_{j}^{n}(t),X_{i}^{n}(t)))dt + (\sigma(t,X_{j}^{n}(t)) + \frac{1}{n} \sum_{i=1}^{n} c(t,X_{j}^{n}(t),X_{i}^{n}(t)))dW_{t}^{j}$$
(3.2.1)

and $X_j^n(0)$ $1 \le j \le n$ being iid Φ' -valued r.v.'s with law of $X_j^n(0)$ given by the probability measure μ_0 . $\{W^j\}$, $1 \le j \le n$, are independent copies of a Brownian motion with Q as the covariance form. Besides,

a:
$$\mathbb{R}^+ \times \Phi' \longrightarrow \Phi'$$

b: $\mathbb{R}^+ \times \Phi' \times \Phi' \longrightarrow \Phi'$
 σ : $\mathbb{R}^+ \times \Phi' \longrightarrow L(\Phi': \Phi')$
c: $\mathbb{R}^+ \times \Phi' \times \Phi' \longrightarrow L(\Phi': \Phi')$

Let $(X_1^n(t), \ldots, X_n^n(t)) \in \Phi' \times \ldots \times \Phi'$ be a solution of (3.2.1). Then the isomorphism given by (3.1.6) between $\Phi' \times \ldots \times \Phi'$ and $(\Phi \times \ldots \times \Phi)'$ we can write the system of SDE's (3.2.1) as follows:

$$dX_t^n = (\overline{a}(t, X_t^n) + \overline{b}(t, X_t^n))dt + (\overline{\sigma}(t, X_t^n) + \overline{c}(t, X_t^n))dW_t \qquad (3.2.2)$$

where initial value X_0^n is a $(\Phi \times \ldots \times \Phi)$ '-valued r.v. such that X_0^n is isomorphic to $(X_1^n(0), \ldots, X_n^n(0)) \in (\Phi' \times \ldots \times \Phi')$, and W_t is the $(\Phi \times \ldots \times \Phi)$ '-valued Wiener process described below.

$$W_{t}(\varphi) = \sum_{j=1}^{n} W_{t}^{j}(\varphi_{j}).$$

By the independence of $\{W^j\}_{1\leq j\leq n}$

$$\langle W \rangle [\varphi, \psi] = t \sum_{j=1}^{n} Q(\varphi_j, \psi_j).$$

Besides, $\{X_t^n\}$ is a $(\Phi \times \ldots \times \Phi)$ '-valued process. The coefficients appearing in (3.2.2) are given as follows: $\square \mathbb{R}^+ \times (\Phi \times \ldots \times \Phi') \longrightarrow (\Phi \times \ldots \times \Phi)'$ $\overline{b}: \mathbb{R}^+ \times (\Phi \times \ldots \times \Phi)' \longrightarrow (\Phi \times \ldots \times \Phi)'$ $\overline{\sigma}: \mathbb{R}^+ \times (\Phi \times \ldots \times \Phi)' \longrightarrow L((\Phi \times \ldots \times \Phi)': (\Phi \times \ldots \times \Phi)'))$ $\overline{c}: \mathbb{R}^+ \times (\Phi \times \ldots \times \Phi)' \longrightarrow L((\Phi \times \ldots \times \Phi)': (\Phi \times \ldots \times \Phi)')$ For $\varphi = (\varphi_1, \ldots, \varphi_n)$ and $\varphi = (\varphi_1, \ldots, \varphi_n) \in \varphi \times \ldots \times \Phi$ and $\underline{u} \cong (\underline{u}_1, \ldots, \underline{u}_n)$ and

$$\mathbf{v} \cong (\mathbf{v}_1, \dots, \mathbf{v}_n) \in (\Phi \times \dots \times \Phi)'$$
, we have

$$\overline{\mathbf{a}}(\mathbf{t},\mathbf{u})[\boldsymbol{\varphi}] = \sum_{j=1}^{n} \mathbf{a}(\mathbf{t},\mathbf{u}_{j})[\boldsymbol{\varphi}_{j}]$$

$$\overline{\mathbf{b}}(\mathbf{t},\mathbf{u})[\boldsymbol{\varphi}] = \sum_{j=1}^{n} \frac{1}{n} \sum_{i=1}^{n} \mathbf{b}(\mathbf{t},\mathbf{u}_{j},\mathbf{u}_{i})[\boldsymbol{\varphi}_{j}]$$

$$\overline{\sigma}(\mathbf{t},\mathbf{u})(\mathbf{v})[\boldsymbol{\varphi}] = \sum_{j=1}^{n} \sigma(\mathbf{t},\mathbf{u}_{j})(\mathbf{v}_{j})[\boldsymbol{\varphi}_{j}]$$

$$\overline{\mathbf{c}}(\mathbf{t},\mathbf{u})(\mathbf{v})[\boldsymbol{\varphi}] = \frac{1}{n} \sum_{j=1}^{n} \sum_{i=1}^{n} c(\mathbf{t},\mathbf{u}_{j},\mathbf{u}_{i})(\mathbf{v}_{j})[\boldsymbol{\varphi}_{j}]$$

so that

$$\int_{0}^{t} \overline{\sigma}(s, \underline{u}) dW_{s}[\varphi] = \sum_{j=1}^{n} \int_{0}^{t} \sigma(s, u_{j}) dW_{s}^{j}[\varphi_{j}]$$
$$\int_{0}^{t} \overline{c}(s, \underline{u}) dW_{s}[\varphi] = \frac{1}{n} \sum_{j=1}^{n} \sum_{i=1}^{n} \int_{0}^{t} c(s, u_{j}, u_{i}) dW_{s}^{j}[\varphi_{j}]$$

3.3 Existence and uniqueness of solutions.

Under the conditions given below, there exists a weak solution X^n of (3.2.2). Such a solution will also be shown to be pathwise unique and thus ensures a unique strong solution $\{X_t^n\} \in (\Phi \times \ldots \times \Phi)'$ to the equation (3.2.2) (see [5]).

Analogous to (3.1.6), the isomorphism $C_{(\phi \times \ldots \times \phi)}$, $\cong C_{\phi}, \times \ldots \times C_{\phi}$, is easily established. Towards this, fix T>O and consider $y \in C_{(\phi \times \ldots \times \phi)}^{T}$, so that for each $0 \leq t \leq T$, $y(t) \in (\phi \times \ldots \times \phi)^{T}$. By using (3.1.6), y(t) is isomorphic to, say, $(y_{1}(t), \ldots, y_{n}(t)) \in \phi^{T} \times \ldots \times \phi^{T}$ for $0 \leq t \leq T$ and so, $\lim_{t \to t_{0}} y(t) = y(t_{0})$ is to the set of equivalent to $\lim_{t \to t_{0}} y_{k}(t) = y_{k}(t_{0})$ for each $1 \leq k \leq n$. Let $\{| \cdot |_{\alpha} : \alpha \in A\}$ be the set of semi-norms defining the strong topology of ϕ^{T} . Set $w_{k} w_{\alpha} = \sup_{0 \leq t \leq T} |y_{k}(t)|_{\alpha}$ for $y_{k} \in C_{\phi}^{T}$, $1 \leq k \leq n$ and $\alpha \in A$. Define $w_{y} w_{\alpha}^{2} = \sum_{k=1}^{n} \sum_{m \neq k} w_{\alpha} w_{\alpha}^{2}$ for each $\alpha \in A$. By replacing $\| \cdot \|_{r}$ by $w \cdot w_{\alpha}$ in (3.1.4), the arguments used in deriving (3.1.6) hold in the present context as well and thus $C_{(\phi \times \ldots \times \phi)}^{T}$, is isomorphic to $C_{\phi}^{T}, \ldots \times C_{\phi}^{T}$, $C_{\phi}, \times \ldots \times C_{\phi}^{T}$, $T \in \mathbb{N}$ respectively, are therefore isomorphic.

Let $X_{j}^{n}(t)$, $1 \leq j \leq n$ and t > 0, be such that for each $\varphi \cong (\varphi_{1}, \ldots, \varphi_{n}) \in \Phi \times \ldots \times \Phi$, $X_{t}^{n}(\varphi) \cong \Sigma X_{j}^{n}[\varphi_{j}]$. Then $X_{j}^{n}(\cdot) \in C_{\phi}$, for $1 \leq j \leq n$ and solve (3.2.1). The conditions for the existence and uniqueness of solutions of SDE (3.2.2) are as follows: Conditions (A) and (IC) of Section 2 are assumed to hold. It is easy to note that condition (A) on Φ implies that on $\Phi \times \ldots \times \Phi$. Likewise condition (IC) of Section 2 implies

$$\int (1+\|\mathbf{u}\|_{-\mathbf{m}}^2) [\ell \mathbf{n}(3+\|\mathbf{u}\|_{-\mathbf{m}}^2)]^2 \, \overline{\mu}_0(\mathbf{d}\mathbf{u}) < \infty$$
$$(\Phi \times \ldots \times \Phi)'$$

where $u \in (\Phi \times \ldots \times \Phi)^{+}$ and $\overline{\mu_{0}} = \mathscr{L}(X_{0}^{n})$. We will call (A) and (IC) as (SA) and (SIC) where S stands for system of stochastic differential equations. The coefficients a,b,c and σ are assumed to satisfy the following conditions: For any T>O, $\exists L_{T} \ge r$ such that for each $m \ge L_{T}$, \exists a number θ and an index p (note that θ and p depend on m) such that:

(SOC) For
$$u, v \in j_m \phi$$
, and $0 \leq t \leq T$,

$$2a(t,u)[j_{m}u] + 2b(t,u,v)[j_{m}u] + |q_{\sigma(t,u)+c(t,u,v)}|_{-m,-m} \le \theta(1+||u||_{-m}^{2} + \beta(t,u,v)||v||_{-m}^{2})$$

where

$$\beta(t,u,v) = \begin{cases} 0 & \text{if } b(t,u,v) = c(t,u,v) = 0 \\ 1 & \text{otherwise} \end{cases}$$

(SLG) Let $u, v \in \Phi_{-m}$, and $0 \leq t \leq T$. Then $a(t, u) \in \Phi_{-p}$ and $b(t, u, v) \in \Phi_{-p}$. Besides,

$$\begin{aligned} \|a(t,u)\|_{-p}^{2} &\leq \theta(1+\|u\|_{-m}^{2}) \\ \|b(t,u,v)\|_{-p}^{2} &\leq \theta(1+\|u\|_{-m}^{2}+\|v\|_{-m}^{2}) \\ |Q_{\sigma(t,u)}|_{-m,-m} &\leq \theta(1+\|u\|_{-m}^{2}) \\ |Q_{c(t,u,v)}|_{-m,-m} &\leq \theta(1+\|u\|_{-m}^{2}+\|v\|_{-m}^{2}) \end{aligned}$$

(SJC) a,b,c and σ are jointly continuous functions. Further,

(i) For $u, v, w \in \Phi_{-m}$, $\sigma(t, u)(w) \in \Phi_{-m}$ $c(t, u, v)(w) \in \Phi_{-m}$ (ii) $Q_{\sigma(t, u)}(\phi, \phi)$ is continuous in u on Φ' and $Q_{c(t, u, v)}(\phi, \phi)$ is continuous in u on Φ' for each $\phi \in \Phi$.

The following condition is needed to prove the uniqueness of solutions. (SMC) For $u_1, v_1, u_2, v_2 \in \Phi_{-m} (\subset \Phi_{-p})$

$$(a(t,u_1)-a(t,v_1),u_1-v_1)_{-p} + |Q_{\sigma(t,u_1)-\sigma(t,v_1)}|_{-p,-p} \leq \theta ||u_1-v_1||_{-p}^2$$

and

$$(b(t,u_{1},u_{2})-b(t,v_{1},v_{2}),u_{1}-v_{1})_{-p} + |Q_{c(t,u_{1},u_{2})-c(t,v_{1},v_{2})}|_{-p,-p}$$

$$\leq \theta \{||u_{1}-v_{1}||_{-p}^{2} + ||u_{2}-v_{2}||_{-p}^{2}\}$$

It is easily verified that the above conditions will imply (CC), (LG), (JC) and (MC) for the coefficients $\overline{a}, \overline{b}, \overline{c}$ and $\overline{\sigma}$ for equations (3.2.2), so that by Theorem 2.4.1 we get existence and uniqueness of solutions for the SDE (3.2.2). The moment bound given in Theorem 2.4.2 becomes

$$E \sup_{\substack{0 \le t \le T}} \|X_{j}^{n}(t)\|_{-m}^{2k} \le (2C_{k}^{+1})\exp((136k^{2}-4k)\theta T)-1 \qquad (3.3.1)$$

where $C_k = E \|X_1^n(0)\|_{-m}^{2k} < \infty$ and is independent of n. In fact, the following bound can also be derived by an obvious modification of the proof of (3.3.1): Under the Condition (IC)

$$\begin{array}{l} E \quad \sup_{\substack{|t-s| < \delta}} \|X_{j}^{n}(s) - X_{j}^{n}(t)\|_{-p}^{2} \leq c\delta^{\cancel{M}} \\ 0 \leq t, s \leq T \end{array}$$

$$(3.3.2) \quad (3.3.2)$$

where c is a constant independent of n. To see this, note that if $T \ge t \ge s \ge 0$ with t-s $\le \delta$, then

$$X_{j}^{n}(t) = X_{j}^{n}(s) + \int_{s}^{t} \{(a(s), X_{j}^{n}(s)) + \frac{1}{n} \sum_{i=1}^{n} b(s, X_{j}^{n}(s), X_{i}^{n}(s))\} ds$$

+
$$\int_{s}^{t} \{\sigma(s, X_{j}^{n}(s)) + \frac{1}{n} \sum_{i=1}^{n} c(s, X_{j}^{n}(s), X_{i}^{n}(s))\} dW_{s}^{j}.$$

We get (3.3.2) by the familiar route namely, via. Doob's inequality, Jensen's inequality wherein the condition (SLG) is used crucially.

<u>Note</u>: $0 < T < \infty$ will be kept fixed till the last paragraph in Section 6. Thus X_i^n

 $1 \leq j \leq n$, the solution of 3.2.1 will each have paths in C_{Φ}^{T} a.s. where p is the index that appears in the conditions.

§4. WEAK COMPACTNESS OF EMPIRICAL MEASURES.

Let X^n_{\bullet} denote the solution of the SDE (3.2.2) so that $X^n_{\bullet} \cong (X^n_1(\bullet), \dots, X^n_n(\bullet)) \in C^T_{\Phi_{-p}} \times \dots \times C^T_{\Phi_{-p}}$ where $X^n_j(\bullet)$, $1 \le j \le n$ solve the SDE's (3.2.1).

Let $\mathfrak{A}(C_{\Phi}^{T})$ be the Borel σ -algebra of C_{Φ}^{T} . For $\omega \in \Omega$, $B \in \mathfrak{A}(C_{\Phi}^{T})$, define -p -p -p -p -p -p

$$\mu_{n}(\omega,B) = \frac{1}{n} \sum_{j=1}^{n} \delta_{j}(\cdot,\omega)$$
(B). (4.1)

For any $k \ge 1$, let $\pi(C_{\Phi_{-k}}^{T})$ be the space of all probability measures on $C_{\Phi_{-k}}^{T}$ equipped with the topology of weak convergence of measures. Likewise $\pi(C_{\Phi^{-1}}^{T})$ will be the space of all probability measures equipped with the topology of weak convergence of measures. Note that the canonical injection $i:\pi(C_{\Phi_{-k}}^{T}) \subset \pi(C_{\Phi_{-\ell}}^{T})$ is continuous if $k \le \ell$. To see this, let $\lambda_n \in \pi(C_{\Phi_{-k}}^{T})$ such that $\lambda_n \Rightarrow \lambda$ in $\pi(C_{\Phi_{-k}}^{T})$. Therefore, for all $f \in C_b(C_{\Phi_{-k}}^{T})$, $\int_{C_{\Phi_{-k}}}^{T} f(y)\lambda_n(dy) \rightarrow$ $C_{\Phi_{-k}}^{T}$ $\int_{C_{-k}}^{T} f(y)\lambda(dy)$ as $n \to \infty$. If $g \in C_b(C_{\Phi_{-\ell}}^{T})$, let $j: \Phi_{-k} \subset \Phi_{-\ell}$ be the continuous $c_{\Phi_{-k}}^{T}$ canonical injection of Φ_{-k} into $\Phi_{-\ell}$ so that the composition $g \cdot j \in C_b(C_{\Phi_{-k}}^{T})$. Besides. $\int_{C_{\Phi_{-k}}}^{T} (g',j)(y)\lambda_n(dy) = \int_{C_{\Phi_{-\ell}}}^{T} g(y)\lambda_n(dy)$ and $\int_{C_{\Phi_{-k}}}^{T} (g',j)(y)\lambda(dy) =$ $c_{\Phi_{-k}}^{T} = C_{\Phi_{-\ell}}^{T}$ as $n \to \infty$. $c_{\Phi_{-k}}^{T} = \lambda$ in $\pi(C_{\Phi_{-k}}^{T})$ as $n \to \infty$.
$$\begin{split} & \mu_n \text{ is a random measure with } \mu_n(\omega, \cdot) \in \pi(C_{\Phi_{-p}}^T) \text{ for each } \omega \in \Omega \text{ and } n \geq 1. \\ & \text{Let } \nu_n(B) = E\mu_n(B) \text{ for all } B \in \mathfrak{B}(C_{\Phi_{-p}}^T). \text{ Let } \eta_n := \mathfrak{L}(\mu_n), \text{ the law of } \mu_n, \text{ i.e.,} \\ & \text{ the probability measure on } \pi(C_{\Phi_{-p}}^T) \text{ induced by the random measure } \mu_n. \text{ Thus,} \\ & \eta_n \in \pi(\pi(C_{\Phi_{-p}}^T)) \end{split}$$

<u>Theorem 4.1</u>: Under the conditions (SIC), (SA), (SOC), (SLG), (SJC) and (SMC), we have

(a) the sequence $\{v_n\}$ is tight in $\pi(C_{\Phi_{-q}}^T)$ for some $q \ge p$. (b) the sequence $\{\eta_n\}$ is weakly compact in $\pi(\pi(C_{\Phi_{-q}}^T))$.

<u>Proof</u>: (a) For any set $B \in \mathfrak{B}(C^{T}_{\phi})$,

$$\begin{split} \nu_n(B^C) &= E\mu_n(B^C) = \frac{1}{n} \sum_{j=1}^n P(X_j^n \in B^C) = P(X_1^n \in B^C) \text{ since } X_j^n, \ 1 \leq j \leq n, \text{ are identically} \\ \text{distributed random variables. Therefore, the tightness of } \{\nu_n\} \text{ in } \pi(C_{\phi}^T) \text{ is} \\ \text{equivalent to the tightness of } \{X_1^n\}_{n \geq 1}, \text{ i.e., of the probability measures } \{P^n\} \\ \text{ on } \pi(C_{\phi}^T) \text{ where } P^n = \text{ law of } X_1^n. \end{split}$$

By a result of Mitoma [14], the tightness of $\{P^n\}$ is equivalent to the tightness of $\{P^n \pi_{\phi}^{-1}\}$ on $\pi(C_{\mathbb{R}}^T)$ for each $\phi \in \Phi$, where $\pi_{\phi} \colon C_{\Phi}^T, \longrightarrow C_{\mathbb{R}}^T$ with $\pi_{\phi}(u) = u[\phi]$. The tightness of $\{P^n \pi_{\phi}^{-1}\}$ follows by verifying the following two conditions:

(i) Given $\epsilon > 0$, $\exists a > 0$ such that

$$\sup_{n} P^{n}(y \in C_{\phi}^{T}: \sup_{0 \leq t \leq T} |y_{t}[\phi]| > a) < \epsilon$$

(ii) Given $\epsilon > 0$ and $\rho > 0$, $\exists \delta > 0$ such that

$$\sup_{n} P^{n}(y \in C_{\phi}^{T}: \sup_{|t-s| \leq \delta} |y_{t}[\phi]| - y_{s}[\phi]| \geq \epsilon) \leq \rho.$$

Condition (i) is verified by noting that E sup $\|X_1^n(t)\|_{-p}^2$ is finite and $0 \le t \le T$ independent of n. To verify (ii), note that

$$P^{n}(y: \sup_{|t-s| < \delta} |y_{t}[\phi] - y_{s}[\phi]| \ge \epsilon) \le P^{n}(y: \sup_{|t-s| < \delta} |y_{t} - y_{s}|| - p^{||\phi||}_{p} \ge \epsilon)$$

$$\le \frac{\|\phi\|_{m}^{2}}{\epsilon^{2}} E \sup_{|t-s| < \delta} \|X_{1}^{n}(t) - X_{1}^{n}(s)\|_{-p}^{2} \le \rho$$

if δ is such that $E \sup_{|t-s| < \delta} \|X_1^n(t) - X_1^n(s)\|_{-p}^2 \leq \frac{\epsilon^2 \rho}{\|\phi\|_p^2}$.

By the moment bound given in (3.3.2) we have that

so that a δ as desired does indeed exist once we are given ϵ, ρ and ϕ .

The tightness of $\{v_n\}$ is thus established on the space $\pi(C_{\Phi}^T)$. For any $\phi \in \Phi$, and any given $\epsilon > 0$,

$$P(\sup_{0 \leq t \leq T} |X_{j}^{n}(t)[\phi]| > \epsilon) \leq P(\sup_{0 \leq t \leq T} ||X_{j}^{n}(t)||_{-m} ||\phi||_{m} > \epsilon)$$

$$\leq \frac{||\phi||_{m}^{2}}{\epsilon^{2}} E \sup_{0 \leq t \leq T} ||X_{j}^{n}(t)||_{-m}^{2}$$

$$\leq \frac{||\phi||_{m}^{2}}{\epsilon^{2}} (2C_{1}+1)e^{132\theta T} \leq \rho$$

if $\|\phi\|_m^2 \leq \frac{\rho \epsilon^2}{(2C_1+1)e^{132\theta T}}$. Thus by Mitoma ([14]; Remark (R.1)), $\{v_n\}$ are uniformly m-continuous and hence, are uniformly p-continuous as well.

Therefore, there exists an index $q \ge p$ such that $\{v_n\}$ is tight in $\pi(C_{\Phi}^T)$. (b) For the second part of the theorem, let us look upon $\mu_n(\omega, \cdot)$ as $\pi(C_{\Phi}^{T})$ -valued random variables. Then $\nu_{n} \in \pi(C_{\Phi}^{T})$ and $\eta_{n} \in \pi(\pi(C_{\Phi}^{T}))$ for all $n \ge 1$.

Note that
$$v_n(B) = \int \lambda(B)\eta_n(d\lambda) \forall B \in \mathfrak{B} \pi(C_{\Phi}^T)$$
. (4.2)
 $\pi(C_{\Phi}^T)$

Using part (a) of the theorem, for each $j \ge 1$, there exists a compact set K_j in $C_{\Phi_{-q}}^{T}$ such that $v_n(K_j^{C}) = \int \lambda(K_j^{C}) d\eta_n(\lambda) \le \epsilon/j^3$ where $\epsilon > 0$ is given. Let $K=\{\lambda: \lambda(K_j) \ge 1 - \frac{1}{j} \forall j\}$ where K_j 's can, WLOG, be taken to be increasing sets. $K \subset \pi(C_{\Phi_{-q}}^{T})$ is compact since closed tight subsets of probability measures on a complete separable metric space (in our case, on $C_{\Phi_{-q}}^{T}$) are compact (see Chapter II, Theorem 6.7 in [15]).

$$\eta_{n}(K^{C}) = P(\mu_{n} \in K^{C}) \leq \sum_{j=1}^{\infty} \eta_{n}(\lambda : \lambda(K_{j}) < 1 - \frac{1}{j})$$

$$= \sum_{j=1}^{\infty} \eta_{n}(\lambda : \lambda(K_{j}) \geq \frac{1}{j})$$

$$\leq \sum_{j=1}^{\infty} \frac{\int \lambda(K_{j}^{C}) d\eta_{n}}{(1/j)}$$

$$\leq \epsilon \sum_{j=1}^{\infty} \frac{1}{j^{2}} \leq 2\epsilon.$$

Thus tightness of $\{\eta_n\}$ in $\pi(\pi(C_{\Phi}^T))$ ensues.

Note that $\pi(C_{\Phi}^{T})$ equipped with the topology of weak convergence is a -q complete separable metric space (see Ch. II, Theorems 6.2 and 6.5 in [15]). Tight subsets of probability measures on a complete separable metric space are relatively compact. The proof is thus complete.

<u>Remark 4.1.</u> Let $\{\eta_{n_k}\}$ be the subsequence given by the above theorem so that $\eta_{n_k} \Rightarrow \eta$ (say). Let μ denote the $\pi(C_{\Phi_{-q}}^T)$ -valued random variable whose law is given by η . Thus $\eta_{n_k} \Rightarrow \eta$ is equivalent to saying that $\mu_{n_k} \xrightarrow{D} \mu$.

<u>Remark 4.2.</u> By a well-known theorem of Skorohod, there exists a probability space on which are defined $\pi(C_{\Phi-q}^{T})$ -valued random variables, say $\{Z_n\}$ and Z with law of $Z_{n_k} = \eta_{n_k}$ and law of $Z = \eta$ such that $Z_{n_k} \rightarrow Z$ a.s.. Using this representation and applying Fatou's lemma, we get

$$E \sup_{\substack{0 \leq s \leq T}} \int \|y_{s}\|_{-q}^{2} Z(dy) = \int \{\sup_{\substack{\sigma \in C_{\Phi}^{-} \\ -q}} \int \|y_{s}\|_{-q}^{2} \lambda(dy) \eta(d\lambda)$$

$$\leq \underbrace{\lim_{k \to \infty}}_{k \to \infty} E \sup_{\substack{0 \leq s \leq T}} \int \|y_{s}\|_{-q}^{2} Z_{n}(dy)$$

$$= \underbrace{\lim_{k \to \infty}}_{q} \int \{\sup_{\substack{\sigma \in S \leq T}} \int \|y_{s}\|_{-q}^{2} \lambda(dy) \eta_{n}(d\lambda)$$

$$\leq \lim_{k \to \infty} \int \|y_{s}\|_{-p}^{2} \lambda(dy) \eta_{n}(d\lambda)$$

$$\leq \lim_{k \to \infty} E \sup_{\substack{\sigma \in S \leq T}} \int \int |y_{s}\|_{-p}^{n} \lambda(dy) \eta_{n}(d\lambda)$$

$$\leq \lim_{k \to \infty} E \sup_{\substack{\sigma \in S \leq T}} \frac{1}{n_{k}} \int \int |y_{s}\|_{-p}^{n} (s) \|_{-m}^{2}$$

$$\leq (2C_{1}+1) \exp(132\theta T) \qquad (4.3)$$

by using (3.3.1). The inequality (4.3) holds with C_1 and θ remaining the same if T is replaced by any t on both sides of the inequality (4.3), $0 \le t \le T$.

<u>Remark 4.3</u>: In case μ is a degenerate random variable, and Y is a C_{Φ}^{T} -valued random variable with $\mathscr{L}(Y) = \mu$, then Remark (4.2) implies that for each $0 \le t \le T$.

$$E \sup_{0 \le s \le t} \|Y_s\|_{-q}^2 \le (2C_1 + 1) \exp(132\theta t).$$

$$(4.4)$$

§5. THE MCKEAN-VLASOV EQUATION

Let $\{Y_t: 0 \le t \le T\}$ be a Φ' -valued stochastic process that solves the following SDE known as the Global McKean-Vlasov equation: For $0 \le t \le T$,

$$Y_{t} = Y_{0} + \int_{0}^{t} A(s, Y_{s}, \mathcal{L}(Y)) ds + \int_{0}^{t} B(s, Y_{s}, \mathcal{L}(Y)) dW_{s}$$
(5.1)

where

$$A(s,u,\lambda): [0,T] \times \Phi' \times \pi(C_{\Phi'}^{T}) \longrightarrow \Phi'$$
$$B(s,u,\lambda): [0,T] \times \Phi' \times \pi(C_{\Phi'}^{T}) \longrightarrow L(\Phi': \Phi')$$

and W is a Φ '-valued Wiener process. $\mathcal{L}(Y)$ denotes the law of Y, and Y₀ is a Φ '-valued random variable.

The local McKean-Vlasov equation is of the form:

$$Y_{t} = Y_{0} + \int_{0}^{t} A(s, Y_{s}, \mathcal{L}(Y_{s})) ds + \int_{0}^{t} B(s, Y_{s}, \mathcal{L}(Y_{s})) dW_{s}$$
(5.2)

for $0 \leq t \leq T$.

By uniqueness of solutions of the SDE (5.1), we mean the following: For each $\lambda \in \pi(C_{\phi}^{T})$, let, for $0 \leq t \leq T$,

$$Y_{t}^{\lambda} = Y_{0} + \int_{0}^{t} A(s, Y_{s}^{\lambda}, \lambda) ds + \int_{0}^{t} B(s, Y_{s}^{\lambda}, \lambda) dW_{s}.$$
 (5.3)

Suppose there are λ_1 and $\lambda_2 \in \pi(C_{\Phi}^T)$ such that $\lambda_1 = \mathcal{L}(Y^{\lambda_1})$ and $\lambda_2 = \mathcal{L}(Y^{\lambda_2})$. Then $\lambda_1 = \lambda_2$.

Existence and uniqueness of solutions of equation (5.1) in full generality will appear in Baldwin et al. [1]. Here, we content ourselves with the following choice of A and B: For each $\lambda \in \pi(C_{\Phi}^{T})$, let

$$A(s,u,\lambda) = a(s,u) + b(s,u,\lambda)$$

where $\tilde{b}(s,u,\lambda) = \int b(s,u,y_s)\lambda(dy)$ and C_{ϕ}^{T} , $B(s,u,\lambda) = \sigma(s,u) + \tilde{c}(s,u,\lambda)$ (5.5) where $c(s,u,\lambda) = \int c(s,u,y_s)\lambda(dy)$. Besides, we assume that there exists a C_x^T ,

sufficiently large number M(m), possibly depending on m, where m is as in subsection 3.3, such that for each u,v $\in \Phi_{-m}$, and $0 \leq s \leq T$,

$$b(s,u,v) = \begin{cases} b_{s}(u,v) & \text{if } \|u-v\|_{-m} \leq M(m) \\ \hat{b}_{s}(u,v) & \text{otherwise} \end{cases}$$
(5.6)

where $b_{s}(u,v)$ and $\hat{b}_{s}(u,v)$ are functions of u,v with $\|\hat{b}_{s}(u,v)\|_{m} \leq C(m)$ for each $s \in [0,T]$. Likewise

$$c(s,u,v) = \begin{cases} c_s(u,v) & \text{if } \|u-v\|_{-m} \leq M(m) \\ c_s(u,v) & \text{otherwise.} \end{cases}$$
(5.7)

with $c_s(u,v)$ and $\hat{c}_s(u,v)$ are functions of u,v with $|Q_n|_{-m,-m} \leq C(m)$ for $c_s(u,v)$

each s \in [0,T].

With b and c as above we first note that \widetilde{b} and \widetilde{c} exist and are finite. To see this, consider

$$\widetilde{b}(s,u,\lambda) = \int b(s,u,v_s) d\lambda(v) = \int b(s,u,v) d\lambda \pi^{-1}(v).$$

$$C_{\Phi}^{T},$$

Since $u \in \Phi'$, there exists an index k such that $u \in \Phi_{-k}$, and such a k can be chosen to be sufficiently large.

Using such an index k in the place of m in our conditions given in subsection (3.3) as well as (5.6) and (5.7), we get for each $\varphi \in \Phi$ that

$$\begin{aligned} |\tilde{\mathbf{b}}(\mathbf{s},\mathbf{u},\lambda)[\varphi]| &\leq \int_{\Phi'} |\mathbf{b}(\mathbf{s},\mathbf{u},\mathbf{v})[\varphi]| d\lambda \pi_{\mathbf{s}}^{-1}(\mathbf{v}) \\ &= \int_{A} |\mathbf{b}(\mathbf{s},\mathbf{u},\mathbf{v})[\varphi]| d\lambda \pi_{\mathbf{s}}^{-1}(\mathbf{v}) \end{aligned}$$

+
$$\int |b(s,u,v)[\varphi] | d\lambda \pi_s^{-1}(v)$$

A^c

where $A = \{v: \|u-v\|_{-k} \leq M(k)\}$. Continuing,

$$\leq \int \|\varphi\|_{\mathbf{p}_{\mathbf{k}}} (1+\|\mathbf{u}\|_{-\mathbf{k}}+\|\mathbf{v}\|_{-\mathbf{k}}) d\lambda \pi_{\mathbf{s}}^{-1}(\mathbf{v}) + C(\mathbf{k})$$

by using (SLG) with $p_k \ge k$ as the index that corresponds to k. Continuing,

$$\leq ||\varphi||_{p_{k}} (M(k) + 1 + 2||u||_{-k} + C(k) < \infty.$$

Likewise one can establish the finiteness of $\tilde{c}(s,u,\lambda)$ by showing that $|Q_{\tilde{c}(s,u,\lambda)}|_{-k,-k} < \infty$ whenever $u \in \Phi_{-k}$.

Such a choice of b and c makes physical sense in it that a pair of particles far apart interact boundedly. This choice includes in particular the case where b(s,u,v) and c(s,u,v) are both bounded in the sense that

$$\|\mathbf{b}(\mathbf{s},\mathbf{u},\mathbf{v})\|_{-\mathbf{m}} \leq C$$

$$\left\|\mathbf{Q}_{\mathbf{c}}\right\|_{-\mathbf{m},-\mathbf{m}} \leq C$$

for all $0 \leq s \leq T$, $u \in \Phi'$ and $v \in \Phi'$. We assume that the functions a,b,c and σ satisfy the conditions (SOC), (SLG), (SJC) and (SMC). It is then a routine matter to check that A and B, as defined above, satisfy (OC), (LG), (JC) and (MC) as listed in Section 2 with the same indices and with constants independent of the measures $\lambda \in \pi(C_{\Phi'}^T)$. We need and hence introduce the following additional condition:

(SJC)' (i) For each $\lambda \in \pi(C_{\Phi'}^T)$, $\tilde{b}(s,u,\lambda)$ and $\tilde{c}(s,u,\lambda)$ are jointly continuous in s and u. Q (ϕ,ϕ) is continuous in u on Φ' for each $\phi \in \Phi$ and $\tilde{c}(s,u,\lambda)$ s $\in [0,T]$. (ii) For each $\phi \in \phi$, $u \in \phi_{-m}$ and $s \in [0,T]$, $\tilde{b}(s,u,\lambda_n)[\phi]$ converges continuously to $\tilde{b}(s,u,\lambda)[\phi]$ and $\tilde{c}^{*}(s,u,\lambda_n)\phi$ converges continuously to $\tilde{c}^{*}(s,u,\lambda)\phi$ as $\lambda_n \Rightarrow \lambda$ in $\pi(C_{\phi}^T)$.

<u>Theorem 5.1:</u> Consider the SDE (5.3) with A and B specified by equations (5.4) to (5.7). Assume conditions (SA), (SIC), (SOC), (SLG), (SJC), (SMC) and (SJC)' and that $\text{EllY}_0 \|_{-m}^4$ where C is a positive constant. Then,

(i) the McKean-Vlasov equation defined by the equation (5.3) admits a solution. The solution Y^{λ} lies in $C_{\Phi_{-n}}^{T}$ a.s.

(ii) The solution is unique if the following additional condition holds (MC π): For all u, v $\in \phi_{-m}$, and $\zeta_1, \zeta_2 \in \pi(C_{\phi}^T)$, and $0 \leq s \leq T$,

$$\langle \widetilde{\mathbf{b}}(\mathbf{s},\mathbf{u},\boldsymbol{\zeta}_{1}) - \widetilde{\mathbf{b}}(\mathbf{s},\mathbf{v},\boldsymbol{\zeta}_{2}),\mathbf{u}-\mathbf{v} \rangle_{-\mathbf{p}} + \|\mathbf{Q}_{\widetilde{\mathbf{c}}}(\mathbf{s},\mathbf{u},\boldsymbol{\zeta}_{1}) - \widetilde{\mathbf{c}}(\mathbf{s},\mathbf{v},\boldsymbol{\zeta}_{2})\|_{-\mathbf{p},-\mathbf{p}}$$

$$\leq C_{T} \{ K_{\mathbf{s}}(\boldsymbol{\zeta}_{1},\boldsymbol{\zeta}_{2}) \|\mathbf{u}-\mathbf{v}\|_{-\mathbf{p}} + \|\mathbf{u}-\mathbf{v}\|_{-\mathbf{p}}^{2} \}$$

$$(5.8)$$

where C_{T} is a constant and

$$K_{s}(\zeta_{1},\zeta_{2}) = \begin{cases} \inf_{\lambda \in \mathscr{A}(\zeta_{1},\zeta_{2})} & \int_{\Phi'} \int_{\Phi'} \|u_{1}^{-u}u_{2}\|_{-p} \lambda_{s}(du_{1}du_{2}) & \text{if } \zeta_{1} \neq \zeta_{2} \\ 0 & \text{if } \zeta_{1} = \zeta_{2} \end{cases}$$

Here, $\mathscr{A}(\zeta_1,\zeta_2) = \text{the set of all probability measures } \lambda \text{ on } C_{\Phi}^T \times C_{\Phi}^T$, with the prescribed marginals ζ_1 and ζ_2 . Besides, $\lambda_t = \lambda \pi_t^{-1}$.

<u>Proof:</u> A complete and detailed proof of this result will appear in Baldwin, et al. [1]. Here we will briefly outline the basic ideas of their proof with modifications to suit our needs.

Let $A = \{\mathscr{L}(Y^{\lambda}) : \lambda \in \pi(C_{\Phi}^{T},)\}$. Then A is a tight subset of $\pi(C_{\Phi}^{T})$ by Theorem (2.4.3). Define the map $\psi:\pi(C_{\Phi}^{T}) \to A$ by $\psi(\mu) = \mathscr{L}(Y^{\mu})$. Again by theorem (2.4.3), ψ is sequentially continuous. Note that A is a subset of $\pi(C_{\Phi_{-p}}^{T})$. By using Theorem (2.4.2) for k=1 and a result of Mitoma [14], we know that there exists an index $\overline{p} \ge p$ such that A is tight as a subset of $\pi(C_{\Phi_{-p}}^{T})$. Let $c\ell(coA)$ denote the closure of the convex hull of A in $\pi(C_{\Phi_{-p}}^{T})$. Then $c\ell(coA)$ is tight in $\pi(C_{\Phi_{-p}}^{T})$ and is therefore compact in $\pi(C_{\Phi_{-p}}^{T})$. The canonical inclusion $j:\pi(C_{\Phi_{-p}}^{T}) \subset \pi(C_{\Phi}^{T})$ is continuous as can be seen from the proof of part (c) of Theorem 6.3. Therefore, $c\ell(coA)$ can be viewed as a subset of $\pi(C_{\Phi}^{T})$ and is a compact and tight subset of $\pi(C_{\Phi}^{T})$.

It can be shown that the topology of weak convergence in $\pi(C_{\phi}^{T})$ when relativized to a compact tight subset of $\pi(C_{\phi}^{T})$ is metrizable so that $c\ell(coA)$ is a Polish space under this topology.

Let $\overline{\psi}$: $c\ell(coA) \rightarrow c\ell(coA)$ be the restriction of ψ to $c\ell(coA)$.

 $c\ell(coA)$ is metrizable and so sequential continuity of $\overline{\psi}$ is equivalent to continuity of $\overline{\psi}$. An application of the Schauder-Tychonoff fixed point theorem (see [3]) gives us the existence of the McKean-Vlasov equation. If $\lambda_0 \in \pi(C_{\Phi}^T)$ such that $\lambda_0 = \mathcal{L}(\Upsilon^0)$, then, for this choice of measure λ_0 , the coefficients A and B satisfy the conditions of existence and uniqueness of solutions as listed in Section 2. Therefore $\Upsilon^{\lambda} \in C_{\Phi_{-D}}^T$, since $A \subset \pi(C_{\Phi_{-D}}^T)$.

For part (ii) of the theorem, let λ^1 and λ^2 be two measures in $\pi(C_{\phi}^T)$ such that $\lambda^1 = \mathcal{L}(\Upsilon^{\lambda^1})$ and $\lambda^2 = \mathcal{L}(\Upsilon^{\lambda^2})$. Then Theorem 2.4.2 implies that E $\sup_{0 \leq t \leq T} ||\Upsilon^{\lambda^1}_t||^2_{-p} < \infty$ since E $\sup_{0 \leq s \leq T} ||\Upsilon^{\lambda^2}_t||^2_{-p} < \infty$. Likewise, E $\sup_{0 \leq t \leq T} ||\Upsilon^{\lambda^2}_t||^2_{-p} < \infty$. Therefore if $\Upsilon_t = \Upsilon^{\lambda^1}_t - \Upsilon^{\lambda^2}_t$, E $\sup_{0 \leq s \leq T} ||\Upsilon^{\mu^2}_t||^2_{-p} < \infty$. Applying the Itô lemma to $||\Upsilon^{\mu^2}_t||^2_{-p} \cdot \exp(-2C_T t)$ where $C_T = C_T(\lambda^1, \lambda^2)$ is the constant that appears in (5.8), and then using the condition in part (ii) of this theorem, we get that

$$\mathbb{E}\|Y_t^2\|_{-p}\exp(-2C_T t) \leq \mathbb{E}\int_0^t K_s(\lambda^1, \lambda^2) \|Y_s\|_{-p}C_T e^{-C_T s} ds$$

Since $K_s(\lambda^1, \lambda^2) \leq E \|Y_s\|_{-p}$, the above inequality yields

$$\mathbb{E} \|Y_t^2\|_{-p} \exp(-2C_T) \leq C_T \int_0^t e^{-C_T s} \mathbb{E} \|Y_s\|_{-p}^2 ds.$$

Gronwall's lemma now yields $E||Y_t^2||_{-p} = 0$ for all $t \in [0,T]$. Since Y_t is sample continuous, $\sup_{0 \le t \le T} ||Y_t^{\lambda^1} - Y_t^{\lambda^2}||_{-p}^2 = 0$ a.s.

<u>Remark 5.1:</u> Since the conditions of Theorem 5.1 hold for all sufficiently large m with p being the index determined by each such m, the conclusion of the theorem holds in particular when m is replaced by a larger index. Therefore, the measure λ_0 obtained in Theorem 5.1 is the unique solution of the McKean-Vlasov equation defined by equation (5.3) even among the measures in the larger space viz. $\pi(C_{\Phi-k}^T)$ for any $k \ge p$. This fact will be used in Section 6 for the particular choice of k=q where q is the index that appears in Theorem 4.1.

§6. PROPAGATION OF CHAOS

Let P_n denote the unique probability measure on $(C_{\Phi_{-q}}^T)^{\oplus n}$ that solves the martingale problem posed by the system of equations (3.2.1) subject to the conditions listed in subsection 3.3, and conditions (SJC)' and (MC π). Besides, we assume that b and c satisfy conditions (5.6) and (5.7). $\eta_n \in \pi(\pi(C_{\Phi_{-q}}^T))$ is

given by

$$\eta_{n}(B) = P_{n}(y_{n}^{n}: \frac{1}{n} \sum_{i=1}^{n} I \in B) \quad \forall B \in \mathfrak{B}(\pi(C_{\phi}^{T}))$$

where y_{-q}^n denotes a generic point in $(C_{\Phi-q}^T)_{-q}^{\otimes n}$ so that $y_{-q}^n = (y_1^n, y_2^n, \dots, y_n^n)$ where

each component belongs to C_{Φ}^{T} . The method employed by Sznitman [12] is used -q in proving the following theorem.

<u>Theorem 6.1:</u> Under the conditions specified in Theorem 5.1, let $\lambda_0 \in \pi(C_{\Phi_{-q}}^T)$ be the unique probability measure that solves the McKean-Vlasov equation (5.3). Then, the subsequence $\{\eta_{n_k}\}$ obtained by Theorem 4.1 is such that $\eta_{n_k} \Rightarrow \delta_{\lambda_0}$ where δ refers to the Dirac δ measure provided that there exists $\alpha > 0$ such that $\text{EllX}_1^n(0) \parallel_{-m}^{4+\alpha} \leq C$, where C is independent of n.

<u>Proof:</u> Let $f \in \mathfrak{A}_b^2(\Phi')$ (see subsection 2.4) so that $f(u) = \tilde{f}(u[\varphi])$ for some $\varphi \in \Phi$ and $\tilde{f} \in C_b^2(\mathbb{R})$.

$$L_{i,j}(f,y^{n},s) = \tilde{f}'(y^{n}_{j}(s)[\varphi])\{a(s,y^{n}_{j}(s))[\varphi] + b(s,y^{n}_{j}(s),y^{n}_{i}(s))[\varphi]\} + \frac{1}{2}\tilde{f}''(y^{n}_{j}(s)[\varphi]) \cdot Q(\frac{1}{n}\sum_{k=1}^{n}(\sigma(s,y^{n}_{j}(s)) + c(s,y^{n}_{j}(s),y^{n}_{k}(s)))^{*}[\varphi], \frac{1}{n}\sum_{k=1}^{n}(\sigma(s,y^{n}_{j}(s)) + c(s,y^{n}_{j}(s),y^{n}_{k}(s)))^{*}[\varphi])$$

where $y_{1}^{n} = (y_{1}^{n}, \dots, y_{n}^{n}) \in \Phi' \times \dots \times \Phi'$. Let

$$L(f, y(s), s, \lambda) = \int \left[\tilde{f}'(y(s)[\varphi]) \\ C_{\Phi_{-q}}^{T} \\ \{a(s, y(s))[\varphi] + b(s, y(s), z(s))[\varphi]\} + \frac{1}{2}\tilde{f}''(y(s)[\varphi]) \\ Q((\int (\sigma(s, y(s)) + c(s, y(s), z_{1}(s))dz(z_{1}))^{*}[\varphi]). \\ C_{\Phi_{-q}}^{T}$$

$$(\int_{\substack{C_{\Phi}^{T} \\ \varphi_{-q}}} (\sigma(s,y(s)) + c(s,y(s),z_{2}(s))d\lambda(z_{2}))^{*}[\varphi]) \Big] d\lambda(z)$$

By the conditions listed in subsection 3.3, the existence of a unique solution to the martingale problem posed by (3.2.1) is guaranteed. We have called such a solution as P_n . Therefore

$$\widetilde{f}(y_{j}^{n}(t)[\varphi]) - \widetilde{f}(y_{j}^{n}(r)[\varphi]) - \int_{r}^{t} \{\frac{1}{n} \sum_{i=1}^{n} L_{i,j}(f, y_{j}^{n}, s) ds$$

is a P_n -martingale with $0 \le r \le t \le T$. By the conditions listed in the statement of Theorem 5.1, a unique solution of the McKean-Vlasov equation posed by (5.3) exists and is denoted by λ_0 . Therefore the following is a λ_0 -martingale:

$$\widetilde{f}(y(t)[\varphi]) - \widetilde{f}((y(r)[\varphi]) - \int_{r}^{t} L(f,y(s),s,\lambda_{o}) ds$$

where $0 \leq r \leq t \leq T$. Consider the function F: $\pi(C_{\Phi}^{T}) \longrightarrow \mathbb{R}$ defined by -q

$$F(\lambda) = \int \{\widetilde{f}(y(t)[\varphi]) - \widetilde{f}(y(r)[\varphi]) - \int_{r}^{t} L(f,y(s),s,\lambda)ds\}$$
(6.1)
$$C_{\phi}^{T} - q$$

$$g_1(y(r_1)) \dots g_p(y(r_p))d\lambda(y)$$

where $0 \leq r_1 \leq r_2 \dots \leq r_p \leq t \leq T$ and g_1, \dots, g_p are bounded functions from $\Phi_{-q} \to \mathbb{R}$. Hence $F(\lambda_0)=0$. Now we will show that $\int F^2(\lambda)\eta(d\lambda) = 0$ by direct $\pi(C_{\Phi_{-q}}^T)$

evaluation. From this, it follows that the support of η is contained in the set of solutions to the L-martingale problem. Corresponding to each solution of the L-martingale problem, we can construct a weak solution of the McKean-Vlasov equation, by the method employed by Kallianpur et al. [8]. From the previous section we know that the McKean-Vlasov equation has a unique

strong solution namely Υ^{λ_0} . Therefore the set of solutions to the L-martingale problem is the singleton set $\{\lambda_0\}$. From the fact that λ_0 is the unique solution to the martingale problem it will then follow that $\eta = \delta_{\lambda_0}$.

$$\underbrace{ \begin{array}{c} \underline{\text{Claim}} : & \lim_{k \to \infty} \int F^2(\lambda) \eta_n(d\lambda) = \int F^2(\lambda) \eta(d\lambda) \\ & \pi(C_{\phi_{-q}}^T) & \kappa & \pi(C_{\phi_{-q}}^T) \end{array} }_{-q}$$

<u>Proof of Claim</u>. Let λ_s denote $\lambda \pi_s^{-1}$. If λ is in the support of η_n recall that λ_s has support in Φ_m for each $0 \leq s \leq T$. For each $u, v \in \Phi'$, $\varphi \in \Phi$, $\lambda \in \pi(C_{\Phi-q}^T)$ and $0 \leq s \leq T$, let for $\mathbb{R} > 0$,

$$a^{R}(s,u)[\varphi] = (-RVa(s,u)[\varphi]) \wedge R$$
$$b^{R}(s,u)[\varphi] = (-RVb(s,u,v))[\varphi]) \wedge R$$
$$Q^{R}_{h(s,u,\lambda_{s})}(\varphi,\varphi) = R \wedge Q_{h(s,u,\lambda_{s})}(\varphi,\varphi)$$

where $h(s,u,\lambda_s) = \sigma(s,u) + \tilde{c}(s,u,\lambda_s)$. Replace a,b and Q by a^R, b^R and Q^R respectively in the definition of F, and call the resulting function as F_R .

$$\lim_{k \to \infty} \int_{\pi(C_{\Phi-q}^{T})} F_{R}^{2}(\lambda) \eta_{n_{k}}(d\lambda) = \int_{\pi(C_{\Phi-q}^{T})} F_{R}^{2}(\lambda) \eta(d\lambda)$$

since $F_R \in C_b(\pi(C_{\Phi_{-q}}^T))$. The claim will be proved if we show that $\int (F^2(\lambda) - F_R^2(\lambda))\eta_n(d\lambda)) \text{ and } \int (F^2(\lambda) - F_R^2(\lambda))\eta(d\lambda) \text{ can both be made}$ $\pi(C_{\Phi_{-q}}^T) \qquad \pi(C_{\Phi_{-q}}^T)$

arbitrarily small when R is sufficiently large.

Using Fubini's theorem and Jensen's inequality,

$$\int_{\substack{\mathbf{r} \in C_{\Phi}^{T} \\ \mathbf{r} = \mathbf{q}}} (F^{2}(\lambda) - F_{R}^{2}(\lambda)) \eta_{n_{\mathbf{k}}} (d\lambda) \leq 3(t-r) (\max_{1 \leq i \leq p} \|g_{i}\|_{\infty}^{2})^{p}$$

$$\times \int_{\mathbf{r}}^{\mathbf{t}} \mathbb{E}_{\eta_{\mathbf{n}_{k}}} \begin{bmatrix} \int & (\tilde{f}(y[\varphi]a(s,y)[\varphi])^{2}\lambda_{s}(dy) & (6.2) \\ & + \int & (\tilde{f}^{*}(y[\varphi])\tilde{b}(s,y,\lambda_{s})[\varphi])^{2}\lambda_{s}(dy) \\ & + \int & (\tilde{f}^{*}(y[\varphi])\tilde{b}(s,y,\lambda_{s})[\varphi])^{2}\lambda_{s}(dy) \\ & + \int & (\tilde{f}^{*}(y[\varphi])Q_{h}(s,y,\lambda_{s})^{(\varphi,\varphi)})^{2}\lambda_{s}(dy)]ds \end{bmatrix}$$

It remains to show that each of the three terms on the right side of (6.2) can be made arbitrarily small, uniformly in k, when R is large. Since the method for each term is essentially the same, we shall consider only the third term.

$$\mathbf{Q}_{h(s,y,\lambda_{s})}(\varphi,\varphi) \leq \|\varphi\|_{m}^{2} |\mathbf{Q}_{h(s,y,\lambda_{s})}|_{-m,-m} \leq \|\varphi\|_{m}^{2} [3\theta(1+4\|y\|_{-m}^{2}+M^{2}(m))+3C^{2}(m)]$$

by using (SLG) and equation (5.7). Thus

$$\{\mathbf{y}: \mathbf{Q}_{\mathbf{h}(\mathbf{s},\mathbf{y},\boldsymbol{\lambda}_{\mathbf{s}})}(\boldsymbol{\varphi},\boldsymbol{\varphi}) > \mathbf{R}\} \subseteq \{\mathbf{y}: \|\mathbf{y}\|_{-\mathbf{m}}^{2} > \mathbf{R}/\mathbf{k}\}$$

where k is a suitable positive constant and R is sufficiently large.

Therefore, the third term on the right side of (6.2) is

$$\leq K_{1} \int_{\mathbf{r}}^{t} E_{\eta_{\mathbf{n}_{\mathbf{k}}}} \int_{\{y: \|y\|_{-m}^{2} > \mathbb{R}/k\}}^{2} (1+\|y\|_{-m}^{4})\lambda_{s}(dy)ds$$

by using (SLG) again with ${\rm K}_1$ as a suitable constant independent of $\eta_{\rm k}.$ Continuing:

$$\leq K_{1} \int_{r}^{t} \frac{1}{n_{k}} \sum_{j=1}^{n_{k}} E[(1+\|x_{j}^{n_{k}}(s)\|_{-m}^{4})I]ds \\ \{\|x_{j}^{n_{k}}(s)\|_{-m}^{2} > R/k\}$$

$$\leq K_{1}2^{4+\alpha} \int_{r}^{t} \frac{1}{n_{k}} \sum_{j=1}^{n_{k}} \{E(1+\|x_{j}^{n_{k}}(s)\|_{-m}^{4+\alpha})\}^{4/(4+\alpha)} \{P(\|x_{j}^{n_{k}}(s)\|_{-m}^{2} > R/k)\}^{\alpha/(4+\alpha)} ds$$

by Hölder's inequality.

$$\leq k_3 / R^{\alpha/(4+\alpha)} \rightarrow 0$$
 as $R \rightarrow \infty$,

wherein the last inequality uses Chebyshev's inequality $2\pi \lambda^2$ the moment bound given by (3.3.1). Besides, k_3 is a constant independent of n_k . The fact that $\int (F^2(\lambda) - F_R^2(\lambda))\eta(d\lambda)$ can be made small for large R, follows along the $\pi(C_{\Phi-q}^T)_{-q}$

same lines as above and its proof is hence omitted.

Continuation of the proof of the theorem: Using the above claim,

$$\begin{aligned} \pi(C_{\Phi_{-q}}^{T})^{F(\lambda)^{2}\eta(d\lambda)} &= \lim_{k \to \infty} \int_{\pi(C_{\Phi_{-q}}^{T})} \int_{p=1}^{\infty} \int_{p=$$

Then, the right side of (6.2) is equal to

$$\int_{\pi(C_{\Phi-q}^{T})} \int_{r}^{t} \frac{1}{\frac{1}{2}} \left(\left(g_{1}(y_{j}^{n_{k}}(r_{1})) \dots g_{p}(y_{j}^{n_{k}}(r_{p})) \right)^{2} \left(\left(\widetilde{f}'(y_{j}^{n_{k}}(s)[\varphi]) \right)^{2} \left(\left(\widetilde{f}'(y_{j}^{n_{k}}(s)[\varphi]) \right)^{2} \right)^{2} \left(\left(\widetilde{f}'(y_{j}^{n_{k}}(s)[\varphi]) \right)^{2} \left(\left((\widetilde{f}'(y_{j}^{n_{k}}(s)[\varphi]) \right)^{2} \left(\left((\widetilde{f}'(y_{j}^{n_{k}}(s)[\varphi]) \right)^{2} \left((\widetilde{f}'(y_{j}^{n$$

•
$$Q((\frac{1}{n_{k}}, \sum_{i=1}^{n_{k}} \{\sigma(s, y_{j}^{n_{k}}(s)) + c(s, y_{j}^{n_{k}}(s), y_{i}^{n_{k}}(s))\})^{*}[\varphi],$$

 $(\frac{1}{n_{k}}, \sum_{i=1}^{n_{k}} \{\sigma(s, y_{j}^{n_{k}}(s)) + c(s, y_{j}^{n_{k}}(s), y_{i}^{n_{k}}(s))\})^{*}[\varphi])dsdP_{n_{k}}(y_{k}^{n_{k}}) = o(\frac{1}{n_{k}^{2}})$

by using the condition (SLG) and the bound given by (3.3.1). Thus

$$\int_{\pi(C_{\Phi_{-q}}^{T})} F(\lambda)^{2} \eta(d\lambda) = \lim_{k \to \infty} \{o(\frac{1}{n_{k}}) + \frac{n_{k}^{-1}}{n_{k}} \int_{(C_{\Phi_{-q}}^{T})} (\tilde{f}(y_{1}^{n_{k}}(t)[\varphi]) \\ - \tilde{f}(y_{1}^{n_{k}}(r)[\varphi]) - \int_{r}^{t} \frac{1}{n_{k}} \sum_{i=1}^{n_{k}} L_{i,1}^{\varphi}(f, y_{i}^{n_{k}}, s) ds) \\ \cdot (\tilde{f}(y_{2}^{n_{k}}(t)[\varphi]) - \tilde{f}(y_{2}^{n_{k}}(r)[\varphi]) - \int_{r}^{t} \frac{1}{n_{k}} \sum_{i=1}^{n_{k}} L_{i,2}^{\varphi}(f, y_{i}^{n_{k}}, s) ds) \\ g_{1}(y_{1}^{n_{k}}(r_{1})) \dots g_{p}(y_{1}^{n_{k}}(r_{p})) g_{1}(y_{2}^{n_{k}}(r_{1})) \dots g_{p}(y_{2}^{n_{k}}(r_{p})) \\ dP_{n_{k}}(y_{i}^{n_{k}})\} = 0$$

since the independence of the Wiener processes W^1 and W^2 implies that

$$\langle M_{r}^{1}(t), M_{r}^{2}(t) \rangle = 0$$

where

$$M_{r}^{1}(t) = \tilde{f}(y_{1}^{n}(t)[\varphi]) - \tilde{f}(y_{1}^{n}(r)[\varphi]) - \int_{r}^{t} \frac{1}{n_{k}} \sum_{i=1}^{n_{k}} L_{i,1}^{\varphi}(f, y_{i}^{n_{k}}, s) ds$$

which is a P_{n_k} -martingale.

Thus $\int_{-q}^{\pi} F(\lambda)^2 \eta(d\lambda) = 0$ for all F defined by (6.1) with $f \in \mathfrak{D}_b^2(\Phi')$, $\pi(C_{\Phi-q}^T)$

 $p \in \mathbb{N}$ and g_1, \ldots, g_p continuous and bounded mapping $\Phi' \longrightarrow \mathbb{R}$, and $0 \leq r_1 \leq \ldots \leq r_p \leq t \leq T$. Since λ_0 is the unique member of $\pi(C_{\Phi \to q}^T)$ such that $F(\lambda_0) = 0$, we get that $\eta = \delta_{\lambda_0}$.

<u>Remark 6.1.</u> By Theorem 6.1, the possibly random measure μ such that $\mu_{n_{k}} \xrightarrow{D} \mu \in \pi(C_{\Phi}^{T})$ has been shown to be a non-random measure and in fact, $\mu = \lambda_{0}$.

Our result on propagation of chaos is presented in the next theorem.

<u>Theorem 6.2.</u> Under the conditions (SA), (SIC), (SOC), (SLG), (SJC), (SMC), (SJC)' and (MCm) and with the coefficients b and c satisfying (5.6) and (5.7), and sup $E||X_1^n(0)||_{-m}^{4+\alpha} \leq C$ for some $\alpha > 0$ we get

- (i) $\eta_n \Rightarrow \delta_{\lambda_0} \text{ in } \pi(\pi(C_{\Phi_{-q}}^T)).$
- (ii) If $\eta_n^t = \mathscr{L}(\mu_n^t)$ where $\mu_n^t(w, X^n) = \frac{1}{n} \sum_{i=1}^n I_i I_i (t, w)$

and $\{X_{i}^{n}(\cdot), 1 \leq i \leq n\}$ solves the system (3.2.1), then $\eta_{n}^{t} \Rightarrow \delta_{\lambda_{0}} \pi_{t}^{-1}$ for $0 \leq t \leq T$. That is to say $\mu_{n}^{t} \rightarrow \lambda_{0} \pi_{t}^{-1}$ in distribution and hence, in probability as well. <u>Proof.</u>: We have shown in Theorem 6.1 that $\eta_{n_{k}} \Rightarrow \delta_{\lambda_{0}}$. In fact, for any convergent subsequence $\{\mu_{n_{j}}, \}$ of the sequence of empirical measures $\{\mu_{n}\}$, we get from Theorem 6.1 that $\eta_{n_{j}} \Rightarrow \delta_{\lambda_{0}}$. Therefore, the whole sequence η_{n} weakly converges to $\delta_{\lambda_{n}}$.

To prove (ii), note that for all real-valued continuous, bounded functions f on $\pi(C_{\Phi}^{T})$,

$$\int_{\pi(C_{\Phi-q}^{T})}^{f(\lambda)\eta_{n}(d\lambda)} \rightarrow \int_{\pi(C_{\Phi-q}^{T})}^{f(\lambda)\delta_{\lambda_{0}}(d\lambda)}.$$
(6.2)

In particular, if $f(\lambda) = \int g(y)d\lambda(y)$ where for all $y \in C_{\Phi_{-q}}^{T}$, $g(y) = g(y_{t})$ $(C_{\Phi_{-q}}^{T})$

for t fixed in [0,T], and \tilde{g} a continuous bounded function from Φ_{-q} to R, then f(λ) is indeed a real-valued, continuous, bounded function on $\pi(C_{\Phi}^{T})$. Therefore, (6.2) for this choice of f implies that $\eta_{n}^{t} \Rightarrow \delta_{\lambda_{0}} \pi_{t}^{-1}$. Thus $\mu_{n}^{t} \rightarrow \lambda_{0} \pi_{t}^{-1}$ as $n \rightarrow \infty$ in probability since the limit is non-random.

<u>Theorem 6.3</u>: (a) For each T>O, let the conditions (SA), (SIC), (SOC), (SLG), (SJC) and (SMC) hold. Then the system of SDE's (3.2.1) admits a weak solution that is pathwise unique. That is, (3.2.1) has a unique strong solution in $(C_{\bar{a}},)^{\otimes n}$.

(b) Assume the additional conditions (SJC)' and (MC π) for each T>0. If b and c are as specified by (5.6) and (5.7), the McKean-Vlasov equation posed by (5.3) has a unique strong solution in C_{ϕ} .

(c) In view of (a) and (b) above, define $\mu_n(\cdot, w) = \frac{1}{n} \sum_{i=1}^{n} \sum_{$

law $\overline{\eta}_n \in \pi(\pi(C_{\phi},))$ $n \ge 1$. Let $\overline{\lambda}_0$ be the probability measure on C_{ϕ} , that solves the McKean-Vlasov equation posed by (5.3). If $\sup_n \text{EllX}_1^n(0) = \lim_{m \to \infty} C$ for some $\alpha \ge 0$, then

$$\overline{\eta}_n \Rightarrow \delta_{\overline{\lambda}_0} \quad \text{in } \pi(\pi(C_{\phi},)).$$

<u>Proof:</u> Part (a) follows by reading off the corresponding result in Kallianpur, Mitoma and Wolpert [8].

(b) Let the conditions of Theorem 6.2 hold for each fixed T>O. Then, the results in Section 5 and 6 hold in the interval $[0,\infty)$. To see this, suppose $\lambda_0 = \mathcal{U}(Y^{\lambda_0})$ solves the McKean-Vlasov equation in the interval $[0,T_0]$, and $\lambda_1 = \mathcal{U}(\overline{Y}^{\lambda_1})$ solves the McKean-Vlasov equation in the interval [0,T], where

 $T_1 > T_0$, then, by the uniqueness of solutions to the McKean-Vlasov equation, we get that the projection of λ_1 on the interval $[0, T_0]$ must coincide with λ_0 . Thus λ_1 is an extension of λ_0 in the above sense. Such an argument shows the existence and uniqueness of solutions to the McKean-Vlasov equation (5.3) in the interval [0,T] for any T>O.

Choose $T_n = n$, and the corresponding measures $\lambda_n \in \pi(C_{\Phi^+})$. Solve the McKean-Vlasov equation (5.3). Then, the projective limit of $\{\lambda_n\}$ is a measure $\overline{\lambda_0} \in \pi(C_{\Phi^+})$ that solves (5.3) for all $t \ge 0$.

To prove part (c), we make the following observations: For any positive index k, i: $C_{\Phi-k}^{T} \subset C_{\Phi}^{T}$, is continuous. In fact, the topology on C_{Φ}^{T} , as given in Section 2 is equivalent to the weakest topology with respect to which the above canonical inclusions are continuous. Therefore, we have

Claim: If
$$\Gamma_{\alpha} \Rightarrow \Gamma$$
 in $\pi(\pi(C_{\Phi}^{T}))$, then $\Gamma_{\alpha} \Rightarrow \Gamma$ in $\pi(\pi(C_{\Phi}^{T}))$.

<u>Proof of the claim</u>: First, note that the inclusion j: $\pi(C_{\Phi}^{T}) \subset \pi(C_{\Phi'}^{T})$ defined by $j(\lambda) = \lambda i^{-1}$ is continuous. To see this, let $\{\lambda_{\alpha}\}$ be a net in $\pi(C_{\Phi-k}^{T})$ such that $\lambda_{\alpha} \Rightarrow \lambda$ in $\pi(C_{\Phi-k}^{T})$. Therefore

$$\int_{C_{\Phi-k}^{T}} f d\lambda_{\alpha} \rightarrow \int_{C_{\Phi-k}^{T}} f d\lambda \forall f \in C_{b}(C_{\Phi-k}^{T}).$$

Let $g \in C_b(C_{\Phi}^T)$. The composition $g \cdot i$ is then in $C_b(C_{\Phi}^T)$. Also

$$\int g d\lambda_{\alpha} i^{-1} = \int g \cdot i d\lambda_{\alpha} \quad \text{for all } \alpha$$

$$C_{\phi}^{T}, \qquad C_{\phi}^{T} - k$$

and

$$\int g d\lambda = \int g \cdot i d\lambda_{\alpha}$$
$$C_{\Phi}^{T}, \qquad C_{\Phi}^{T},$$

so that

$$\int g d\lambda_{\alpha} \longrightarrow \int g d\lambda \text{ for all } g \in C_{b}(C_{\phi}^{T}).$$

$$C_{\phi}^{T}, \qquad C_{\phi}^{T}.$$

Therefore j is continuous.

Now let k be the inclusion from $\pi(\pi(C_{\Phi}^T)) \subset \pi(\pi(C_{\Phi}^T))$. Continuity of k can be proved by following step by step the proof of continuity of j. The claim is thus shown.

Now part (c) is shown by observing that Theorem 6.2 part (i) implies that

$$\eta_n \Rightarrow \delta_{\lambda_0} \quad \text{in } \pi(\pi(C_{\Phi-q}^T))$$

and hence

$$\eta_n \Rightarrow \delta_{\lambda_0}$$
 in $\pi(\pi(C_{\Phi}^T))$

by the claim shown above.

$$\eta_n \Rightarrow \delta_{\lambda_0} \quad \text{in } \pi(\pi(C_{\Phi}))$$

since the inclusion $\pi(\pi(C_{\phi}^{T},)) \subset \pi(\pi(C_{\phi},))$ is also continuous. Note that η_{n} and $\delta_{\lambda_{O}}$ are the projections of $\overline{\eta}_{n}$ and $\delta_{\overline{\lambda}_{O}}$ on $\pi(\pi(C_{\phi}^{T},))$. Thus $\overline{\eta}_{n} => \delta_{\overline{\lambda}_{O}}$ in $\pi(\pi(C_{\phi}^{T},))$.

<u>Remark 6.3</u>: The unique strong solutions mentioned in parts (a) and (b) of the above theorem are in general Φ' -valued processes and cannot be guaranteed to lie in a single Hilbert space Φ_{-j} . This is so since the indices m and p vary with T in the conditions.

§7. APPLICATION TO INTERACTING SYSTEMS OF NEURONS.

The random behavior of the voltage potential of a spatially distributed

neuron has attracted considerable attention in neurophysiology and can be modeled in the following set-up:

Let H be a separable Hilbert space and T_t be a strongly continuous contraction semigroup on H with a densely defined, closed, negative-definite generator A. In practice, H is usually taken to be $L^2(\mathfrak{A},\mu)$ where \mathfrak{A} is the membrane of a neuron and μ is a suitable measure on \mathfrak{A} . If there exists $r_1>0$ so that $(I-A)^{-r_1}$ is Hilbert-Schmidt, then there exists $\{\varphi_j\}_{j=1}^{\infty}$, a CONS for H such that $-A\varphi_j = \lambda_j \varphi_j$, $j=1,2,\ldots$ with $\sum_{j=1}^{\infty} (1+\lambda_j)^{-2r_1} < \infty$. Let $\Phi = \{\varphi \in \mathbb{H}: \sum_{j=1}^{\infty} (1+\lambda_j)^{2r} (\varphi,\varphi_j)_{\mathbb{H}}^2 < \infty$ for any $r>0\}$. Define on Φ a family of increasing Hilbertian norms $\|\cdot\|_r$ with $\|\varphi\|_r^2 = \sum_{j=1}^{\infty} (1+\lambda_j)^{2r} (\varphi,\varphi_j)^2$ and let Φ_r denote the completion of Φ w.r.t. $\|\cdot\|_r$. Since $\Phi_{r+r_1} \subset \Phi_r$ is Hilbert-Schmidt, it is easy to see that Φ is a nuclear space. The semigroup $\{T_t\}_{t>0}$ can be written as follows. For any $\varphi \in \Phi$

$$T_{t}\varphi = \sum_{j=1}^{\infty} \exp(-t\lambda_{j})(\varphi,\varphi_{j})_{0}\varphi_{j} \in \Phi.$$

The voltage potential is identified as the solution of the Φ '-valued SDE.

$$dX_{+} = A'X_{+}dt + dW_{+}$$

where A' is the adjoint operator of A and W_t is a Φ' -Brownian motion with a certain covariance function $E(W_t, \varphi)(W_1, \psi) = (t \land s)Q(\varphi, \psi)$.

More generally, suppose A_t generates a strongly continuous contraction evolution operators T(s,t), $s \leq t$ on Φ . Assume the following conditions on A_t : For any T and large enough m, there exists a p>m such that $|A_t'u|_{-p} \leq K|u|_{-m}$ for all $t \leq T$, and $u \in \Phi'_m$, i.e., as continuous linear operators from Φ'_m to Φ'_p , $\{A_t\}_{t \leq T}$ are uniformly bounded.

Then the following Φ' -valued SDE modeling the voltage potential of a neuron has a unique solution:

$$dX_{t} = A_{t}X_{t}dt + dW_{t}$$
$$X_{0} = \xi_{0}.$$

Moreover, the solution can be explicitly written as:

$$X_{t} = T'(0,t)\xi_{0} + W_{t} + \int_{0}^{t} A'T'_{s,t} W_{s} ds.$$

Here, $T'_{s,t}$ denotes the adjoint operator of $T'_{s,t}$. Now, consider the system of n-interacting neurons whose voltage potentials are governed by the following SDE:

$$dX_{i}^{(n)}(t) = (A_{t}X_{i}^{(n)}(t) + \frac{1}{n}\sum_{j=1}^{n} b_{t}(X_{i}^{n}(t), X_{j}^{n}(t))dt + dW_{i}(t), i=1,2,...,n \quad (7.1)$$

 $X_i^n(0) = \xi \in \Phi'$ where $b_t: \Phi' \times \Phi' \to \Phi'$ represents the interaction between neurons and $\{W_i(t)\}_{i=1}$ are independent copies of a Φ' -valued Brownian motion.

We require that the interaction $b_t: \Phi' \times \Phi' \to \Phi'$ satisfy the conditions (SOC), (SLG), (SJC), (SMC), (SJC)' and (MC π) as in Theorem 6.2.

The existence and uniqueness theorem in Section 3 thus guarantees that system (7.1) has a unique solution. The propagation of chaos in Section 6 asserts that the empirical distribution $\frac{1}{n} \sum_{i=1}^{n} \delta_{i=1} \left(X_{i}^{(n)} \right) \in \pi(C_{\Phi-q}^{T})$ converges in probability to a deterministic probability measure $\lambda_{0} \in \pi(C_{\Phi-q}^{T})$ which is the law of the solution of the McKean-Vlasov equation corresponding to (7.1):

$$dX_{t} = (A_{t}X_{t} + b_{t}[X_{t}, \lambda_{0}^{t}])dt + dW_{t},$$

where

$$b[X,\lambda_0^t] = \int_{\Phi'} b(x,y) d\lambda_0^t(y)$$

and λ_0^t is the law of X_t .

Thus the asymptotic behavior of a large system of neurons through mean-interactions becomes asymptotically independent with the distribution governed by the McKean-Vlasov equation (7.2).

8. CHAOTIC SYSTEMS

8.1 Exchangeable systems.

Till now, the initial random variables $X_j^n(0)$, $1 \le j \le n$ have been assumed to be i.i.d. random variables. We now relax this condition and assume that $X_j^n(0)$, $1 \le j \le n$ are exchangeable random variables for each $n \ge 1$. That is, the law of $X_j^n(0)$ $1 \le j \le n$, denoted by $\mu_0^n \in \pi(\Phi^r)^{\otimes n}$ is a symmetric probability measure on $(\Phi^r)^{\otimes n}$.

We call the symmetric measures $\mu_0^n \quad \mu_0$ -chaotic if the following condition holds: For every integer k≥1 and $f_1, \ldots, f_k \in C_b(\Phi')$,

$$\lim_{n \to \infty} \int_{\Phi'}^{\infty} f_1(u_1), \dots, f_k(u_k) d\mu_0^n(u) = \prod_{i=1}^k \int_{\Phi}^{\infty} f_i(u_i) d\mu_0(u_i)$$
(8.1)

where $u_{n} = (u_{1}, \ldots, u_{n}) \in (\Phi')^{\otimes n}$, and μ_{0} is a probability measure on Φ' . We assume that the measures μ_{0}^{n} are μ_{0} -chaotic.

In the context of the neuorphysiological model described in section 7, the assumption of exchangeability of the law of $(X_1^n(0), \ldots, X_n^n(0))$ for each $n \ge 1$ is equivalent to saying that the particular order in which the neuronal membranes are taken, is immaterial. This is so since the random variables $X_1^n(t), \ldots, X_n^n(t)$ for each $t\ge 0$ and $n\ge 1$ turn out to be exchangeable random variables. The μ_0 -chaoticity assumption is needed in showing the propagation of chaos result. To see this, consider the simplest case where the drift and diffusion coefficients are identically zero so that (8.1) itself becomes the propagation of chaos statement.

The results of the previous sections hold for the exchangeable model as well if we assume μ_0 -chaoticity and that $\operatorname{EllX}_1^n(0) ||_{-m}^{4+\delta} \leq C$ where C is a constant independent of n, and m is the index that appears in the conditions listed in subsection 3.3.

8.2 Finite-dimensional systems

By setting $\Phi_n \equiv \mathbb{R}^d$ with $\|\|x\|\|_n^2 = \sum_{i=1}^d \sum_{i=1}^{q} x_i^2$ for each $n \ge 1$, $\Phi = \mathbb{R}^d = \bigcap_{n \ge 1} \Phi_n$ is seen $n \ge 1$ to be a nuclear space with its strong dual Φ' being isomorphic to \mathbb{R}^d . In this case, all the norms $\|\cdot\|_k$, $-\infty < k < \infty$ are one and the same, namely, the Euclidean norm on \mathbb{R}^d denoted by $\|\cdot\|$. Therefore, the indices m.p.q etc. in our conditions can and will be taken to be 1. The canonical maps j_m will not appear in the conditions in this case. Besides, expressions such as $|Q_{\sigma(t,u)}|_{-m,-m}$ will simply read as trace $(\sigma\sigma^*(t,u))$. Also, the condition (SA) is trivially seen to hold for the choice of $\Phi = \mathbb{R}^d$. The propagation of chaos result for the finite-dimensional exchangeable system is given in the next theorem:

<u>Theorem 8.2.1</u>. For each T > 0, let the conditions (SIC), (SCC), (SLG), (SJC) and (SMC) hold. Let $X_j^n(0) \ 1 \le j \le n$ be exchangeable random variables and let $\mu_0^n = \text{law of } (X_1^n(0), \dots, X_n^n(0))$ be μ_0 -chaotic. Then,

a) The system of SDE's (3.2.1) has a unique strong solution in $(C_{md})^{\otimes n}$.

b) In addition, assume (SJC)' and (MCm) for each T > 0. If the coefficients b and c are as specified by (5.6) and (5.7), then the McKean-Vlasov equation (5.3) has a unique strong solution in C_{md} .

c) Assume the conditions in part (b). Further, assume that there exists $\delta > 0$ such that $E||X_1^n(0)||^{4+\delta} \leq C$ where C is independent of n. Then, in the notation

of section 6

$$\overline{\eta}_{n} \stackrel{>}{\to} \underbrace{\delta}_{\overline{\lambda}_{n}} \quad \text{in } \pi(\pi(C_{n})).$$

The above result enables us to compare our results with those of Sznitman [16].

The conditions made by Sznitman are the following:

(i) The initial random variables $X_1^n(0), \ldots, X_n^n(0)$ are \mathbb{R}^d -valued exchangeable random variables and are bounded.

(ii) $\mu_0^n = \text{law of } (X_1^n(0), \dots, X_n^n(0)) \text{ on } (\mathbb{R}^d)^{\otimes n} \text{ are } \mu_0^-\text{chaotic, where } \mu_0^-\text{ is a probability measure on } \mathbb{R}^d$

(iii) The drift and diffusion coefficients are uniformly bounded and satisfy uniform Lipschitz conditions in the space and time variables.

(iv) The covariance form Q is the identity matrix.

In the next paragraph the conditions (i) through (iv) are compared with those that appear in Theorem 8.2.1.

First (SIC) and the moment condition introduced in part (c) of Theorem 8.2.1 are satisfied since (i) says that the initial variables are bounded. (SOC) is verified as follows:

For $u, v \in \mathbb{R}^d$, $0 \le t \le T$, and $h(t, u, v) = \sigma(t, u) + c(t, u, v)$,

 $|2a(t,u)\cdot u + 2b(t,u,v)\cdot u + tr(hh^{*}(t,u,v))|$

 $\leq 2 \|a\|_{m} \|u\| + 2 \|b\|_{m} \|u\| + trace(hh^{*}(t,u,v))$

 $\leq \theta(1 + \|u\|^2)$

by using the uniform boundedness of the coefficients and condition (iv).

The verifications of (SLG) and (SJC) given the Conditions (i) to (iv) are

simple and hence left to the reader. (SMC) can be verified by using Lipschitz continuity of the coefficients. (SJC) and (MC π) are obtained by continuity and boundedness of the coefficients. Thus, our set of conditions for the propagation of chaos is weaker than that imposed in the finite-dimensional set-up by Sznitman [16]. The finite-dimensional result of Leonard [11] is close in spirit to Theorem 8.2.1 and hence, a comparison of the two is left to the reader.

Bibliography

- Baldwin, D., Hardy, G. and Kallianpur, G. (1990): The McKean-Vlasov equation in duals of nuclear spaces, (submitted for publication, 1990).
- [2] Baldwin, D. (1990): Nuclear space valued stochastic differential equations with some applications, Ph.D. Thesis, University of North Carolina at Chapel Hill.
- [3] Dunford, N. and Schwartz, J.T. (1957): Linear operators, Part I, General Theory, John Wiley and Sons.
- [4] Funaki, T. (1987): Derivation of the hydrodynamical equation for one-dimensional Ginzburg-Landau model, IMA Repring Series No. 328, University of Minnesota.
- [5] Ikeda, N. and Watanabe, S. (1981): Stochastic differential equations and diffusion processes. North-Holland.
- [6] Kallianpur, G. (1986): Stochastic differential equations in duals of nuclear spaces with some applications. IMA Preprint Series No. 244, Inst. Math. Appl., 1986.
- [7] Kallianpur, G. and Mitoma, I. (1988): A Langevin-type stochastic differential equation on a space of generalized functionals. University of North Carolina Center for Stochastic Processes Technical Report No. 238, Aug. 1988.
- [8] Kallianpur, Mitoma, I. and Wolpert, R. (1990): Diffusion equations in duals of nuclear spaces. Stochastics, 29, 1990, 285-329.
- [9] Kallianpur, G. and Perez-Abreu, (1988): Weak convergence of solutions of stochastic evolution equations on nuclear spaces, University of North Carolina Center for Stochastic Processes Technical Report No. 248, Oct. 1988. Proc. Trento Conf. on Infinite Dimensional Stochastic Differential Equations, 1990, to appear.
- [10] Kallianpur, G. and Wolpert, R. (1987): Weak convergence of stochastic neuronal models. Lect. Notes in Biomath. 70, 116-145.
- [11] Leonard, C. (1986): Une loi des grands nombres pour des systèmes de diffusions avec interaction et à coefficients non bornés. Ann. Inst. Henri Poincaré, vol. 22, 237-262.

- [12] Mitoma, I. (1981): Martingales of random distributions, Mem. Fac. Sci. Kyushu Univ. Ser. A 35, 185-197.
- [13] Mitoma, I. (1983): Continuity of stochastic processes with values in the dual of a nuclear space, Z. Wahr. verw. Gebiete 63, 271-279.
- [14] Mitoma, I. (1984): Tightness of probabilities on C([0,1]; 9') and D([0,1]: 9') Ann. Prob. 11, 989-999.
- [15] Parthasarathy, K.R. (1967): Probability measures on metric spaces. Academic Press.
- [16] Sznitman, A.S. (1984): Nonlinear reflecting diffusion process and the propagation of chaos and fluctuations associated. J. Functional Anal. 56, 311-336.

Technical Reports Center for Stochastic Processes Department of Statistics University of North Carolina Chapel Hill, NC 27599-3260

- .258. C. Houdré, Linear Fourier and stochastic analysis, Apr. 89.
- 259. G. Kallianpur, A line grid method in areal sampling and its connection with some early work of H. Robbins, Apr. 89. Amer. J. Math. Manag. Sci., 1989, to appear.
- 260. G. Kallianpur, A.G. Miamee and H. Niemi, On the prediction theory of two-parameter stationary random fields, Apr. 89. J. Multivariate Anal., 32, 1990, 120-149.
- 261. I. Herbst and L. Pitt, Diffusion equation techniques in stochastic monotonicity and positive correlations, Apr. 89.
- 262. R. Selukar, On estimation of Hilbert space valued parameters, Apr. 89. (Dissertation)
- 263. E. Mayer-Wolf, The noncontinuity of the inverse Radon transform with an application to probability laws, Apr. 89.
- 264. D. Monrad and W. Philipp, Approximation theorems for weakly dependent random vectors and Hilbert space valued martingales, Apr. 89.
- 265. K. Benhenni and S. Cambanis, Sampling designs for estimating integrals of stochastic processes, Apr. 89.
- 266. S. Evans, Association and random measures, May 89.
- 267. H.L. Hurd, Correlation theory of almost periodically correlated processes, June 89.
- -268. O. Kallenberg, Random time change and an integral representation for marked stopping times, June 89. Probab. Th. Rel. Fields, accepted.
- 269. O. Kallenberg, Some uses of point processes in multiple stochastic integration, Aug. 89.
- 270. W. Wu and S. Cambanis, Conditional variance of symmetric stable variables, Sept. 89.
- 271. J. Mijnheer, U-statistics and double stable integrals, Sept. 89.
- 272. O. Kallenberg, On an independence criterion for multiple Wiener integrals, Sept. 89.
- 273. G. Kallianpur, Infinite dimensional stochastic differential equations with applications, Sept. 89.
- 274. G.W. Johnson and G. Kallianpur, Homogeneous chaos, p-forms, scaling and the Feynman integral, Sept. 89.
- 275. T. Hida, A white noise theory of infinite dimensional calculus, Oct. 89.
- 276. K. Benhenni, Sample designs for estimating integrals of stochastic processes, Oct.
 89. (Dissertation)
- 277. I. Rychlik, The two-barrier problem for continuously differentiable processes, Oct. 89.
- .278. G. Kallianpur and R. Selukar, Estimation of Hilbert space valued parameters by the method of sieves, Oct. 89.

- 278. G. Kallianpur and R. Selukar, Estimation of Hilbert space valued parameters by the method of sieves, Oct. 89.
- 279. G. Kallianpur and R. Selukar, Parameter estimation in linear filtering, Oct. 89.
- 280. P. Bloomfield and H.L. Hurd, Periodic correlation in stratospheric ozone time series, Oct. 89.
- 281. J.M. Anderson, J. Horowitz and L.D. Pitt, On the existence of local times: a geometric study, Jan. 90.
- 282. G. Lindgren and I. Rychlik, Slepian models and regression approximations in crossing and extreme value theory, Jan. 90.
- 283. H.L. Koul, M-estimators in linear models with long range dependent errors, Feb. 90.
- 284. H.L. Hurd, Almost periodically unitary stochastic processes, Feb. 90.
- 285. M.R. Leadbetter, On a basis for 'Peaks over Threshold' modeling, Mar. 90.
- 286. S. Cambanis and E. Masry, Trapezoidal stratified Monte Carlo integration, Mar. 90.
- 287. M. Marques and S. Cambanis, Dichotomies for certain product measures and stable processes, Mar. 90.
- 288. M. Maejima and Y. Morita, Trimmed sums of mixing triangular arrays with stationary rows, Mar. 90.
- 289. S. Cambanis and M. Maejima, Characterizations of one-sided linear fractional Lévy motions, Mar. 90.
- 290. N. Kono and M. Maejima, Hölder continuity of sample paths of some self-similar stable processes, Mar. 90.
- 291. M. Merkle, Multi-Hilbertian spaces and their duals, Mar. 90
- 292. H. Rootzén, M.R. Leadbetter and L. de Haan, Tail and quantile estimation for strongly mixing stationary sequences, Apr. 90.
- 293. K. Benhenni and S. Cambanis, Sampling designs for estimating integrals of stochastic processes using quadratic mean derivatives, Apl. 90.
- 294. S. Nandagopalan, On estimating the extremal index for a class of stationary sequences, Apr. 90.
- 295. M.R. Leadbetter and H. Rootzén, On central limit theory for families of strongly mixing additive set functions, May 90.
- 296. W. Wu, E. Carlstein and S. Cambanis, Bootstrapping the sample mean for data from general distribution, May 90.
- 297. S. Cambanis and C. Houdré, Stable noise: moving averages vs Fourier transforms, May 90.
- 298. T.S. Chiang, G. Kallianpur and P. Sundar, Propagation of chaos and the McKean-Vlasov equation in duals of nuclear spaces, May 90.
- 299. J.M.P. Albin, On the upper and lower classes for stationary Gaussian fields on Abelian groups with a regularly varying entropy, June 90.