

2

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of the collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 2 Nov 1990 3. REPORT TYPE AND DATES COVERED Final Report/1 Sep 84-31 Jul 90

4. TITLE AND SUBTITLE
RESONANCE ENERGIES AND WIDTHS FROM THE POLES OF THE
MULTICHANNEL T MATRIX

5. FUNDING NUMBERS
61102F/2301/A4

6. AUTHOR(S)
Deborah K. Watson

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)
Univ of Oklahoma
Dept Physics & Astronomy
Norman, OK 73019-0225

8. PERFORMING ORGANIZATION
REPORT NUMBER
AFOSR-TR- 90 1145

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)
AFOSR/NP
Bolling AFB DC 20332-6448

10. SPONSORING / MONITORING
AGENCY REPORT NUMBER
AFOSR-84-0379

DTIC
ELECTE
NOV 21 1990
S E D
E

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT
Approved for public release; distribution is unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)
A multichannel iteration technique was developed to obtain accurate results for phase shifts from a "smooth" Schwinger K matrix. Applications were made to $e + He^+$ phase shifts in one, two, and three channel approximations. A second project was started to study excited states of helium using dimensional analysis. A moment method is being used to solve the two-electron Schrodinger equation generalized to an arbitrary number of dimensions.

14. SUBJECT TERMS

15. NUMBER OF PAGES
09

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT
UNCLASSIFIED

18. SECURITY CLASSIFICATION
OF THIS PAGE
UNCLASSIFIED

19. SECURITY CLASSIFICATION
OF ABSTRACT
UNCLASSIFIED

20. LIMITATION OF ABSTRACT
UL
SAR

AD-A229 064

Final Technical Report
1 Sep 84 - 31 Jul 90

Deborah Kay Watson
Department of Physics and Astronomy
University of Oklahoma
Norman, Oklahoma

A method of determining Rydberg states and resonance parameters based on a "smooth" K matrix has been extended to multichannel cases. An energy independent projection operator formalism to ensure orthogonality has been developed and a multichannel iteration procedure has now been successfully formulated. We have obtained accurate results for the $1P$, $3P$ and $3S$ Rydberg states of helium and for phase shifts using two channels at positive energies. We are presently working on the implementation of an improved multichannel iteration formalism which we hope will converge the three channel results to values which are independent of the initial basis set.

During the last year of the grant we began an investigation of excited states of helium using dimensional perturbation theory. We have successfully obtained energies for some singly-excited states of helium and have also completed a study of some $1,3p^e$ states using inter-dimensional degeneracies.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Final Technical Report

This final report summarizes the research performed under Grant No. AFOSR-84-0379 from, September 1987, the date of the most recent Final Technical Report, to July 1990 when this grant terminated. The goal of this proposal, "Resonance Energies and Widths from the Poles of the Multichannel T Matrix" was to determine Rydberg and resonance poles using a "smooth" K matrix. Previous calculations had successfully determined Rydberg states of some alkali atoms and beryllium as well as resonance states for beryllium and H^- by searching for the poles of the complex T matrix. The present technique avoids both the use of complex arithmetic and individual searches for each pole by formulating a "smooth" K matrix which is real and free of poles. This is done by using alternative boundary conditions to define a "smooth" Green's function which is then used to determine the K matrix. Since this K matrix has no poles at bound-state energies, a smooth quantum defect curve can be defined at all negative energies. Determining bound state energies from this curve was found to be much more efficient than performing individual searches in a single-channel study of the Rydberg states of the lithium isoelectronic sequence. At positive energies a phase shift curve can be used to extract resonance parameters using a Breit-Wigner fit. The major goal of the present grant was to extend this new K matrix method to multichannel Rydberg and resonance studies. This required the development of a multichannel "smooth" Green's function and K matrix. Implementing this formalism also required the development of a projection operator formalism to ensure that the Rydberg electron is orthogonal to the core. A further goal was to develop a multichannel iteration technique to converge results after an initial basis set is chosen.

In summary, the Research Objectives as described above are:

1. Extension of the K matrix method to multichannel Rydberg and resonance states
2. Development of a numerical multichannel "smooth" Green's function
3. Development of a projection operation formalism to enforce orthogonality
4. Development of a multichannel iteration technique
5. Application to H^- and He.

We first focused on the Rydberg states of helium using two channels. We were able to obtain very good agreement with experimental energies for the $1P$, $3P$ and $3S$ states of helium. The basis set consisted of solutions from single-channel calculations, four basis vectors in the first channel and a single basis vector in the second channel. Iteration was not necessary to obtain satisfactory results. An energy-independent projection operator formalism was successfully implemented and the effect of the closed channels was included by using a reduced K matrix formalism. This work was presented at the 19th annual meeting of the Division of Atomic, Molecular and Optical Physics in Baltimore in April 1988 and is presently being prepared for publication in Phys. Rev. A.

Publications:

T. L. Goforth and D. K. Watson, "Multichannel Quantum Defect Calculations Using a Smooth Reaction Matrix", to be submitted, Phys. Rev. A.

This method was then extended to positive energies. Accurate results for phase shifts using one or two channels were easily obtained. For three channels we found that iteration in all channels became very important for accurate results. We believe we now have an improved multichannel iteration formalism which will enable us to obtain phase shifts and resonance parameters that are independent of the initial basis set.

This work was presented at the Sixteenth International Conference on the Physics of Electronic and Atomic Collisions in New York City in July 1989.

A second project was started in March of 1989 when a postdoctoral fellow, David Goodson, joined the group. We are investigating a new way of studying atomic phenomena called dimensional scaling or dimensional perturbation theory. The particular goal of this project is to study resonance states of helium using these techniques. Dimensional analysis obtains information about physical systems by generalizing the Schrodinger equation to an arbitrary number of dimensions. A moment method is being used to solve this generalized Schrodinger equation for some excited states of helium. We have chosen to formulate the problem using normal coordinates corresponding to the normal modes of vibration to take advantage of the pseudomolecular nature of the electron-nucleus-electron system. We are hopeful that this approach will provide insight into the geometry of resonance states of helium as well as provide meaningful quantum numbers for these strongly correlated states which are not well described by independent particle models. We presently have successfully obtained energies for some singly-excited states of helium and are analyzing the behavior of their asymptotic energy series. We have also completed a study of some interdimensional degeneracies between $D=3$ and $D=5$ which allows very accurate $1,3p^e$ states for $D=3$ to be determined from S states in 5 dimensions. A description of this work has been prepared for publication in Phys. Rev. A.

Publications:

D. Goodson, D. K. Watson, J. G. Loeser and D. R. Herschbach, "Energies of Doubly-Excited Two-Electron Atoms from Interdimensional Degeneracies", to be submitted, Phys. Rev. A.

List of professional personnel:

Deborah K. Watson, Associate Professor

Terry L. Goforth, Ph.D August 1990

David Z. Goodson, Postdoctoral Assistant