

2

AD-A229 398

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including reviewing existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND DATES COVERED		
		Final Report, 1 Aug 88 to 31 Jul 89		
TITLE AND SUBTITLE		5. FUNDING NUMBERS		
PULSE PROPAGATION IN TEMPORALLY DISPERSIVE MEDIA		F49620-88-C-0094 61102F 2304/A4		
AUTHOR(S)				
Gregory A. Kriegsmann				
PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER		
WAVE PRO, INC. 311 Pine Tree Road Radnor, PA 19087		AFOSR-TR- 00 1076		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448		F49620-88-C-0094		
<p style="font-size: 2em; font-weight: bold; margin: 0;">DTIC</p> <p style="font-size: 2em; font-weight: bold; margin: 0;">ELECTE</p> <p style="font-size: 2em; font-weight: bold; margin: 0;">NOV 16 1990</p> <p style="font-size: 4em; font-weight: bold; margin: 0;">S B D</p>				
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT		12b. DISTRIBUTION CODE		
Approved for public release; distribution unlimited.				
13. ABSTRACT (Maximum 200 words)				
<p>By solving the transport equations for the amplitude of the progressing wave expansion of Maxwell's equations using the Lorentz model, we find that the amplitudes do <u>not</u> decay exponentially along the ray. Thus, we are now studying more general models of the dispersive media, to determine the qualitative features for which the amplitudes do, or do not decay exponentially. Preliminary results suggest that a classification of dispersive media is obtained depending on the relative orders of the differential operators.</p>				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			5	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	
UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED	SAR	

WAVE PRO, INC.
311 Pine Tree Road
Radnor, PA 19087

Final Technical Report for the period August 1, 1988 - July 31, 1990
Project F49620-88-C-0094

Prepared for: Air Force Office of Scientific Research

G.A. Kriegsmann (pp)
Gregory A. Kriegsmann
President

September 30, 1990

90 11 15 080

For a dispersive, non-magnetic, electrically conducting medium, Maxwell's equations for the propagation of electromagnetic waves are given in terms of dimensionless variables by

$$\nabla \times \underline{E} = - \underline{H}_t \quad , \quad \nabla \times \underline{H} = \underline{D}_t \quad , \quad (1)$$

$$\underline{D} = (\underline{E} + \underline{P}) \quad . \quad (2)$$

The vectors $\underline{D}(\underline{x}, t)$, $\underline{E}(\underline{x}, t)$, $\underline{H}(\underline{x}, t)$ and $\underline{P}(\underline{x}, t)$ are proportional to the electric displacement, the electric field strength, the magnetic field strength, and the polarization vector, respectively. The constitutive properties of the medium, which are experimentally determined give the polarization vector in (2) as a linear functional of \underline{E} . A general expression is given by the differential constitutive law,

$$\underline{P}(\underline{x}, t) = \sum_{j=1}^N \underline{P}_j(\underline{x}, t) \quad , \quad (3a)$$

$$\sum_{n=0}^r \alpha_{nj} \frac{\partial^n \underline{P}}{\partial t^n} = \sum_{m=0}^s \beta_{mj} \frac{\partial^m \underline{E}}{\partial t^m} \quad , \quad j = 1, 2, \dots, N \quad . \quad (3b)$$

The specified constitutive coefficients, α_{nj} and β_{mj} , $m = 0, 1, \dots, s$, $n = 0, 1, \dots, r$ and $j = 1, 2, \dots, N$ are experimentally determined.

More generally, we express \underline{P} as a linear functional of \underline{E} by the heredity integral,

$$\underline{P}(\underline{x}, t) = \mathcal{L} \underline{E}(\underline{x}, t) \equiv \int_{-\infty}^t \underline{E}(\underline{x}, t') \epsilon(t-t') dt' \quad , \quad (3)$$

where the "memory" function, $\epsilon(\tau)$, which is experimentally determined, vanishes for $\tau < 0$.

We have first analyzed the propagation of finite jump discontinuous solutions of the system (1)-(3) using both the method of weak solutions [1,2] and progressing wave expansions [1,3] for the Debye [4] model of the



or	<input type="checkbox"/>
	<input type="checkbox"/>
	<input type="checkbox"/>

ity Codes

and/or

Special

Dist

A-1

dispersive medium. This model is believed to be a good approximation of the dispersive properties of biological materials. Thus, each \underline{P}_j satisfies a first order ordinary differential equation

$$\frac{\partial}{\partial t} \underline{P}_j + \alpha_j \underline{P}_j = \beta_j \underline{E} \quad , \quad j = 1, 2, \dots, N \quad . \quad (4)$$

The "constitutive" coefficients $\alpha_1, \alpha_2, \dots, \alpha_n$ and $\beta_1, \beta_2, \dots, \beta_n$ are to be determined experimentally, or from quantum mechanical calculations. Then,

$$\underline{P}_j = \int_{-\infty}^t \underline{E}(\underline{x}, t') \beta_j e^{-\alpha_j(t-t')} dt' \quad (5)$$

and $\underline{P} = \int_{-\infty}^t \underline{E}(\underline{x}, t) \sum_{j=1}^N \beta_j e^{-\alpha_j(t-t')} dt'$, so that the memory function for the Debye model is given by the linear combination of exponentially decaying functions,

$$\epsilon(\tau) = \sum_{j=1}^N \beta_j e^{-\alpha_j \tau} .$$

The progressing wave expansion for the solution of (1), (2) and (5) is a representation in the form

$$\begin{aligned} \underline{E} &= \sum_{m=0}^{\infty} f_m[\phi(\underline{x}, t)] \underline{E}^m(\underline{x}, t) \quad , \quad \underline{H} = \sum_{m=0}^{\infty} f_m[\phi(\underline{x}, t)] \underline{H}^m(\underline{x}, t) \quad , \\ \underline{P}_j &= \sum_{m=0}^{\infty} f_m[\phi(\underline{x}, t)] \underline{P}_j^m(\underline{x}, t) \quad , \end{aligned} \quad (6)$$

where $f_0(\phi)$ is an arbitrary function of the single variable $\phi(\underline{x}, t)$ and f_1, f_2, \dots , are required to satisfy,

$$f'_m = f_{m-1} \quad , \quad m = 1, 2, \dots \quad . \quad (7)$$

Then we insert (6) into (1), (2) and (4) and use (7) for the derivatives of f_m . By equating to zero the coefficient of each function $f_m(\phi)$, we obtain a sequence of equations to determine ϕ , \underline{E}^m , \underline{H}^m and \underline{P}_j^m . For example, if we set

$\phi(\underline{x}, t) = t - \psi(\underline{x})$, then ψ satisfies the eikonal equation of geometrical optics,

$$|\nabla\psi|^2 = 1 \quad , \quad (8)$$

and the coefficients \underline{E}^0 , \underline{E}^1 , etc., satisfy transport equations, which reduce to first order partial differential equations along the characteristics of (8). Solving the transport equations explicitly, shows that \underline{E}^0 and \underline{H}^0 decay exponentially along the rays.

The choice of the waveform $f_0(\phi)$ is determined by the type of discontinuity to be analyzed, and the form of the pulse. For example, to recover the results of our previous analysis of the propagation of jump discontinuities, we select f_0 to be the Heaviside step function i.e.

$$f_0 = H(t-\psi) \quad .$$

For the Lorentz model [4] of dispersion the polarization vector is given by (3a) where the \underline{P}_j now satisfy

$$\frac{\partial^2 \underline{P}_j}{\partial t^2} + \alpha_j \frac{\partial \underline{P}_j}{\partial t} + \gamma_j \underline{P}_j = \beta \underline{E} \quad , \quad (11)$$

and the constitutive coefficients α_j , γ_j and β are to be experimentally determined. By solving the transport equations for the amplitude of the progressing wave expansion of Maxwell's equations using the Lorentz model, we find that the amplitudes do not decay exponentially along the ray. Thus, we are now studying more general models of the dispersive media, e.g. (3), to determine the qualitative features for which the amplitudes do, or do not decay exponentially. Preliminary results suggest that a classification of dispersive media is obtained depending on the relative orders of the differential operators on the right and left sides of (3.3b).

Finally, using the progressing wave expansion, we have determined the scattering of pulses from dispersive half spaces and from dispersive targets.

A paper entitled "Progressing Wave Expansions for Temporally Dispersive Electromagnetic Waves and the Classification of Dispersive Media" is currently in preparation.

References

1. R. Courant and D. Hilbert, "Methods of Mathematical Physics," Vol. II, Interscience, NY, 1962.
2. M. Kline and I. W. Karp, "Electromagnetic Theory and Geometrical Optics," Interscience, NY, 1965.
3. R. M. Lewis, The Progressing Wave Formalism, in "Proc. of the Symp. on Quasi-Optics," Polytechnic Institute of Brooklyn, Polytechnic Press, 1964, pp. 71-103.
4. C. J. F. Bottcher, "Theory of Electric Polarization," Elsevier, Amsterdam, 1952.