





WATER OPERATIONS TECHNICAL SUPPORT PROGRAM



**TECHNICAL REPORT E-90-1** 

# EFFECTS OF ALUM TREATMENT ON PHOSPHORUS DYNAMICS AND PHYTOPLANKTON RESPONSE IN EAU GALLE RESERVOIR, WISCONSIN

by

John W. Barko, William F. James William D. Taylor, Dwilette G. McFarland

Environmental Laboratory

DEPARTMENT OF THE ARMY Waterways Experiment Station, Corps of Engineers 3909 Halls Ferry Road, Vicksburg, MS 39180-6199



September 1990 Final Report

Approved for Public Release; Distribution Unlimited

Prepared for DEPARTMENT OF THE ARMY US Army Corps of Engineers Washington, DC 20314-1000 Destroy this report when no longer needed. Do not return it to the originator.

 $\Psi_{i} = - \Psi_{i}$  ,  $\psi_{i}$ 

🖝 🖇 🦆 🐖 🗸

. 👪 🔮 🏌 🕴

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

> The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

# Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

	N PAGE OMB NO. 0704-0188			MB No. 0704-0188	
. REPORT SECURITY CLASSIFICATION	<u> </u>	1b. RESTRICTIVE MARKINGS			
nclassified					
SECURITY CLASSIFICATION AUTHORITY		3. DISTRIEUTION		JF REPORT	
5. DECLASSIFICATION / DOWNGRADING SCHED	ULE	Approved unlimited	for public 1.	release; (	listribution
PERFORMING ORGANIZATION REPORT NUM	BER(S)	5. MONITORING	ORGANIZATION	REPORT NUMBE	R(S)
Cechnical Report E-90-1	TEN OFFICE SYMPOL	Z NAME OF M			
JSAEWES	(If applicable)	7 INAME OF W	UNITORING ORG	ANIZATION	
Environmental Laboratory					
ADDRESS (City, State, and ZIP Code)		76. ADDRESS (Ci	ty, State, and ZIP	Code)	
J909 Halls Ferry Road Jicksburg MS 39180-6199					
NAME OF FUNDING / SPONSORING	8b. OFFICE SYMBOL	9 PROCUREMEN	T INSTRUMENT I	DENTIFICATION	
ORGANIZATION	(If applicable)				= =
JS Army Corps of Engineers					
ADDRESS (City, State, and ZIP Code)		10. SOURCE OF	UNDING NUMBE	RS	
Jashington DC 20314-1000		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	ACCESSION NO.
151111gcon, DC 20314-1000					
ITLE (Include Security Classification)		<u>.</u>	1		
ffects of Alum Treatment on	Phosphorus Dynam	ics and Phyte	oplankton R	esponse in	Eau Galle
eservoir, Wisconsin					
: PERSONAL AUTHOR(S) Barko John W : James Willia	m F · Tavlor ຟi	lliam D · Mai	Farland Dw	ilette G	
a. TYPE OF REPORT	COVERED	14 DATE OF REPO	RT (Year Month	Dav) 15 PAG	
inal report FROM	TO	Septem	per 1990		48
SUPPLEMENTARY NOTATION					
vailable from National Techn	ical Information	Service, 52	35 Port Roy	al Road, S	pringtield,
	18 SUBJECT TERMS	Continue on rever	a if narassary an	d identify by bl	ock oumber)
FIELD GROUP SUB-GROUP		continue on revers	e n necessary an	a laentity by bi	
	See	reverse.			
		<u></u>			
BSTRACT (Continue on reverse if necessar	y and identify by block n	umber)	the treated	with alw	inum sulfata
alum) in 1986 to reduce exce	ssive phytoplank	ton production	was treated on associat	ed with hi	rhum surrace ph
arum, in 1900 co reduce chee		con producer		00 #2011 112	b
hosphorus loading from anoxi	c, profundal sed	iments. Prie	or to treat	ment, inte	rnal total
hosphorus loading from anoxi hosphorus (TP) loading was 3	c, profundal sed to 6 times grea	iments. Prie ter than exte	or to treat ernal TP lo	ment, inte ading duri	rnal total ng summer
bhosphorus loading from anoxi bhosphorus (TP) loading was 3 stratification. Periodic inc	c, profundal sed to 6 times grea reases in epilim	iments. Prie ter than extension netic TP mas:	or to treat ernal TP lo s and chlor	ment, inte ading duri ophyll α c	rnal total ng summer oncentra-
phosphorus loading from anoxi phosphorus (TP) loading was 3 stratification. Periodic inc tions closely corresponded wi	c, profundal sed to 6 times grea reases in epilim th elevated inte	iments. Prio ter than extended netic TP mass rnal TP. Fo	or to treat ernal TP lo s and chlor r l year fo	ment, inte ading duri ophyll α c llowing tr	rnal total ng summer oncentra- eatment,
phosphorus loading from anoxi phosphorus (TP) loading was 3 stratification. Periodic inc ions closely corresponded wi nternal TP loading and conce	c, profundal sed to 6 times grea reases in epilim th elevated inte ntrations of sol	iments. Prid ter than extanetic TP mass rnal TP. Fo uble reactive	or to treat ernal TP lo s and chlor r l year fo e phosphoru	ment, inte ading duri ophyll α c llowing tr s in the h	rnal total ng summer oncentra- eatment, ypolimnion
phosphorus loading from anoxi phosphorus (TP) loading was 3 stratification. Periodic inc tions closely corresponded wi internal TP loading and conce were substantially reduced.	c, profundal sed to 6 times grea reases in epilim th elevated inte ntrations of sol	iments. Pri- ter than ext netic TP mas. rnal TP. Fo uble reactive	or to treat ernal TP lo s and chlor r l year fo e phosphoru	ment, inte ading duri ophyll a c llowing tr s in the h	rnal total ng summer oncentra- eatment, ypolimnion
phosphorus loading from anoxi phosphorus (TP) loading was 3 stratification. Periodic inc tions closely corresponded wi internal TP loading and conce were substantially reduced. Abnormally high externa	c, profundal sed to 6 times grea reases in epilim th elevated inte ntrations of sol 1 TP loading dur	iments. Prid ter than extended netic TP mass rnal TP. Fo uble reactive ing the stra	or to treat ernal TP lo s and chlor r l year fo e phosphoru tified peri	ment, inte ading duri ophyll a c llowing tr s in the h od of 1986	rnal total ng summer oncentra- eatment, ypolimnion sustained
phosphorus loading from anoxi phosphorus (TP) loading was 3 stratification. Periodic inc tions closely corresponded wi internal TP loading and conce were substantially reduced. Abnormally high externa high TP mass and chlorophyll alum treatment on internal TP	c, profundal sed to 6 times grea reases in epilim th elevated inte ntrations of sol I TP loading dur in the epilimnio loading were es	iments. Prid ter than extense netic TP mass rnal TP. Fo uble reactive ing the stra n. During t sentially ne	or to treat ernal TP lo s and chlor r l year fo e phosphoru tified peri ne summer o gated, as e	ment, inte ading duri ophyll a c llowing tr s in the h od of 1986 f 1987, ef pilimnetic	rnal total ng summer oncentra- eatment, ypolimnion sustained fects of TP mass and
phosphorus loading from anoxi phosphorus (TP) loading was 3 stratification. Periodic inc tions closely corresponded wi internal TP loading and conce were substantially reduced. Abnormally high externa high TP mass and chlorophyll alum treatment on internal TP	c, profundal sed to 6 times grea reases in epilim th elevated inte ntrations of sol I TP loading dur in the epilimnio loading were es	iments. Prid ter than ext netic TP mas rnal TP. Fo uble reactive ing the stra n. During t sentially ne	or to treat ernal TP lo s and chlor r l year fo e phosphoru tified peri he summer o gated, as e	ment, inte ading duri ophyll a c llowing tr s in the h od of 1986 f 1987, ef pilimnetic	rnal total ng summer oncentra- eatment, ypolimnion sustained fects of TP mass and (Continued)
Distribution/AVAILABILITY OF ABSTRACT	c, profundal sed to 6 times grea reases in epilim th elevated inte ntrations of sol I TP loading dur in the epilimnio loading were es	iments. Prid ter than extense netic TP mass rnal TP. For uble reactive ing the stra n. During to sentially ne 21. ABSTRACT SE Unclassi	or to treat ernal TP lo s and chlor r l year fo e phosphoru tified peri ne summer o gated, as e CURITY CLASSIFIE fied	ment, inte ading duri ophyll a c llowing tr s in the h od of 1986 f 1987, ef pilimnetic	rnal total ng summer oncentra- eatment, ypolimnion sustained fects of TP mass and (Continued)
hosphorus loading from anoxi hosphorus (TP) loading was 3 tratification. Periodic inc ions closely corresponded wi nternal TP loading and conce ere substantially reduced. Abnormally high externa igh TP mass and chlorophyll lum treatment on internal TP DISTRIBUTION/AVAILABILITY OF ABSTRACT 3 UNCLASSIFIED/UNLIMITED SAME AS NAME OF RESPONSIBLE INDIVIDUAL	c, profundal sed to 6 times grea reases in epilim th elevated inte ntrations of sol I TP loading dur in the epilimnio loading were es	iments. Prid ter than extanetic TP mas rnal TP. For uble reactive ing the stra n. During to sentially ne 21. ABSTRACT SE Unclassi 225 TELEPHONE	or to treat ernal TP lo s and chlor r l year fo e phosphoru tified peri he summer o gated, as e CURITY CLASSIFIC fied	ment, inte ading duri ophyll a c llowing tr s in the h od of 1986 f 1987, ef pilimnetic CATION	rnal total ng summer oncentra- eatment, ypolimnion sustained fects of TP mass and (Continued)

Unclassified

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

18. SUBJECT TERMS (Continued).

Alum	Phytoplankton
Hydrology	Reservoir
Loadings	Sediment
Nutrient cycling	Water quality
Phosphorus	

19. ABSTRACT (Continued).

T

chlorophyll remained unchanged from pretreatment years. Sedimentation of external TP inputs, with subsequent alum burial, is believed to have been primarily responsible for reducing the effectiveness of alum in controlling internal TP loading in this reservoir.

Following alum treatment in 1986, phytoplankton biomass was less than half that occurring during the previous year; however, it was not significantly different from other years. Subsequent to several years of codominance with blue-green algae, the dinoflagellate *Ceratium hirundinella* assumed singular dominance in 1985. With the exception of this change in relative dominance (maintained through 1988), the phytoplankton community of Eau Galle Reservoir has been remarkably stable, despite effects of a winter drawdown in 1984, alum treatment in 1986, and climatic (wet versus dry) extremes over the 8-year study period. Without a decrease in phosphorus availability, *Ceratium* and bluegreen algae are likely to persist as codominants during the summer in this system for many years.

Acces	sion For	
NTIS	GRA&I	
DTIC	TAB	
Unanz	ounced	
Justi	fication_	
By		
Distr	ibution/	
Avat	lability (	Codes
	Avail and	/01
Dist	Special	
١		
0~/ I		
r		
	*	
	I'M S	
	( ~ ° ° ° °	, <b>)</b>
	1 Fey	' <b>J</b>

Unclassified SECURITY CLASSIFICATION OF THIS PAGE

# PREFACE

The studies reported herein were sponsored by the Headquarters, US Army Corps of Engineers (HQUSACE), Directorate of Civil Works, through the US Army Corps of Engineers Environmental and Water Quality Operational Studies (EWQOS) Program, the Algae Control Demonstration (ACD) Program, the Water Operations Technical Support (WOTS) Program, and the US Army Engineer District (USAED), St. Paul, St. Paul, MN. The EWQOS and ACD Programs were managed by the US Army Engineer Waterways Experiment Station (WES), under the direction of Dr. Jerome L. Mahloch, Manager. The WOTS Program is managed by the WES under the Environmental Resources Research and Assistance Programs, Mr. J. Lewis Decell, Manager. Technical Monitors for the study were Dr. John Bushman, Mr. James Gottesman, and Mr. David Buelow, HQUSACE.

Principal investigator for this study was Dr. John W. Barko, Environmental Laboratory (EL), WES. Experimental design, data analysis and interpretation, and preparation of the report were accomplished by Drs. Barko, William F. James, and William D. Taylor and Ms. Dwilette G. McFarland. Technical reviews were provided by Drs. Robert Gaugush, Douglas Gunnison, Thomas L. Hart, and Nancy J. McCreary of the EL. Additional reviews were provided by Messrs. Daniel Wilcox and Dennis Holme of the USAED, St. Paul.

Assistance during alum application was provided by Messrs. J. Mason, J. Wagner, G. Quinn, S. Greb, and D. Lillie and Dr. P. Garrison of the Bureau of Research, Wisconsin Department of Natural Resources, and by Dr. R. H. Kennedy of the EL. Mr. D. Knauer, also of the Bureau of Research, coordinated equipment design and operational activities. Mr. P. Bradley, Mr. H. Eakin, Ms. B. Fulton, Dr. G. Godshalk, Ms. Y. Hartz, Mr. R. Kuta, Ms. B. Nelson, Ms. K. Mueller, and Mr. R. Olewinski are gratefully acknowledged for their participation in the conduct of the fieldwork and chemical analyses.

This investigation was performed under the general supervision of Dr. John Harrison, Chief, EL, and Mr. Donald L. Robey, Chief, Ecosystem Research and Simulation Division, and under the direct supervision of Dr. Thomas L. Hart, Chief, Aquatic Processes and Effects Group.

Commander and Director of WES was COL Larry B. Fulton, EN. Technical Director was Dr. Robert W. Whalin.

This report should be cited as follows:

Barko, John W., James, William F., Taylor, William D., and McFarland, Dwilette G. 1990. "Effects of Alum Treatment on Phosphorus Dynamics and Phytoplankton Response in Eau Galle Reservoir, Wisconsin," Technical Report E-90-1, US Army Engineer Waterways Experiment Station, Vicksburg, MS.

# CONTENTS

			Page
PREFA	CE		1
PART	I:	INTRODUCTION	4
	Backg Site Objec	ground Description ctives	4 5 6
PART	II:	METHODS	7
	Alum Analy Phosp Phyto	Treatment vtical Protocol bhorus Loading Determinations oplankton Determinations	7 7 8 9
PART	III:	RESULTS	11
	Phos Phyto	phorus Dynamics	11 13
PART	IV:	DISCUSSION	16
PART	V:	FUTURE STUDIES	22
REFER	RENCES		23
TABLE	ES 1-4		
FIGUE	RES 1-1	11	
APPEN	NDIX A	PHYTOPLANKTON SPECIES LIST, EAU GALLE RESERVOIR	A1

# EFFECTS OF ALUM TREATMENT ON PHOSPHORUS DYNAMICS AND PHYTOPLANKTON RESPONSE IN EAU GALLE RESERVOIR, WISCONSIN

PART I: INTRODUCTION

#### Background

1. Eutrophication is a process frequently stimulated by the transport of nutrients from watershed to lake. While this is a natural process in all lakes and reservoirs, the rate at which it occurs is often accelerated by human activities in the watershed. Symptomatic of the advancement of eutrophication are reduced water clarity, poor dissolved oxygen conditions in bottom waters, excessive algal production, and impaired recreational value.

2. The obvious best means of ameliorating these undesirable conditions is the reduction of external material inputs. Many lake restoration techniques have been directed toward the control of external phosphorus (P) loading to reduce phytoplankton production. Successful lake restoration has often been achieved by diverting stormwater, sewage, and septic tank inputs (e.g., Edmondson 1970, Cooke et al. 1978). However, internal P loading can significantly delay lake recovery following implementation of these restoration efforts (Welch 1977, Larson et al. 1979, Ryding 1981).

3. Concentrations of P usually increase with increasing depth in the hypolimnion, a pattern closely related to changes in the area-to-volume ratio with depth and the progression of anoxia (Stauffer 1985). An increasing body of evidence indicates that anoxic, profundal sediments are an important source of internal P loading and contribute significantly to patterns of increase in P mass in the hypolimnetic water column (Mortimer 1971; Cooke et al. 1977; Theis and McCabe 1978; Larson, Schultz, and Malueg 1981; Nurnberg 1984, 1987; Riley and Prepas 1984; Nurnberg et al. 1986).

4. Phosphorus released from anoxic sediments may be transported to the epilimnion via thermally induced turbulent eddy diffusion (Jassby and Powell 1975; Imboden and Emerson 1978; Robards and Ward 1978; Stefan and Hanson 1981; Wodka et al. 1983) and wind-induced metalimnetic migration (Stauffer and Lee 1973; Larson, Schultz, and Malueg 1981; Kortmann et al. 1982; Stauffer and Armstrong 1984; Stauffer 1985, 1987; Effler et al. 1986; James, Kennedy, and

Gaugush, in press). These processes stimulate phytoplankton productivity, and thereby reduce recreational value. Under such circumstances, alternative means for improving or maintaining water quality must be sought. Many of these alternative solutions involve attempts to control the availability of P within the water body. With decreased availability of P, excessive phytoplankton growth can usually be curtailed.

5. Ferric aluminum sulfate and aluminum sulfate (alum) treatments have been widely used in lakes to reduce internal P loading from the sediment, and thereby reduce phytoplankton biomass (Peterson et al. 1973; Cooke and Kennedy 1978; Cooke et al. 1978, 1982; Soltero et al. 1981; Garrison and Knauer 1983; Foy 1985). The success of alum treatment usually depends on concurrent reduction of external P loading in addition to long-term control of sediment P release (Kennedy and Cooke 1982). While most natural lakes receive the majority of their P income from internal sources during the summer, reservoirs often receive substantial external P inputs from inflowing tributaries. Thus, in reservoirs, the success of alum treatment may largely depend on the seasonality of external P inputs.

# Site Description

6. Eau Galle Reservoir (Figure 1) is a small  $(0.6-km^2)$  impoundment located in west-central Wisconsin. It is operated by the US Army Engineer District, St. Paul. The reservoir is dimictic and approximately circular in shape. It has a mean depth of 3.2 m, a maximum depth of 9 m, and receives the majority of its water (80 percent) and nutrient inputs from the Eau Galle River. Minor inflowing tributaries include French, Lousy, and Lohn Creeks.

7. External P loading cannot be controlled effectively in this reservoir because of its large size  $(166 \text{ km}^2)$  and the agricultural nature of the watershed. External loadings occur primarily in the spring and fall, during which periods the entire volume of the reservoir can be replaced within only a few days. Loadings are usually minimal during the summer months, although freshets do occur with low frequency (Kennedy 1987). Internal P loading from anoxic sediments and the accumulation of hypolimnetic P is substantial during the summer (James, Kennedy and Gaugush, in press).

8. The outlet structure has both surface and hypolimnetic discharge ports. Surface discharge is uncontrolled and occurs at pool elevations above

286.5 m, relative to mean sea level. Discharge during the summer stratified period occurs primarily through a controlled vertical slide gate located 7.2 m below the surface discharge elevation. During the winter of 1984, the reservoir was drawn down to nearly 1 m below normal pool elevation over about a 4-month period for maintenance purposes. This event exposed variably up to 53 percent of the reservoir littoral zone to freezing and desiccation, and accounted for about a 47-percent loss of pool volume.

9. Withdrawal of cooler water during the summer stratified period gradually warms the hypoliunion, promoting a decrease in the stability of the water column (Gaugush 1984) with associated increases in phytoplankton production, primarily blue-green and dinoflagellate populations (Barko et al. 1984). During periods of declining stability in the summer, migrations of the metalimnion occur in Eau Galle Reservoir as a result of wind mixing (James, Kennedy, and Gaugush, in press). Mixing events have an important influence on P availability to phytoplankton in the epilimnion of this system.

# Objectives

10. The objective of this investigation was to limit phytoplankton population densities by controlling (via alum treatment) internal P loading from profundal sediments. To our knowledge, direct control of sediment P release with alum has heretofore not been attempted in flood-control reservoirs. The studies reported herein examine the effects of a hypolimnetic alum treatment on sediment P dynamics and phytoplankton response. Results are reported for the pretreatment (1981-1985) and posttreatment (1986-1988) periods. This report provides extensive information on P cycling and contains a detailed account of phytoplankton community composition in Eau Galle Reservoir over the 8-year study period.

### PART II: METHODS

#### Alum Treatment

11. Alum (11.3 g Al/m<sup>2</sup>; 4.5 mg Al/ $\ell$ ) was administered to the reservoir on 29 May 1986. Details of dosage and application procedures are provided in Kennedy et al. (1987). Based on previous measurements of internal total phosphorus (TP) loading, the dose applied was theoretically sufficient to control sediment P release for 5 years. The treatment area was confined to depths of 3 m and greater, since sediments at these depths experience anoxia during summer stratification. Here, only anoxic sediments were assumed to contribute to the internal P load.

### Analytical Protocol

12. Water samples for chemical analyses were collected at 1- or 2-week intervals at 1-m increments throughout the summer months in 1981, 1982, 1986, 1987, and 1988 at a deep (9-m), centrally located station. The samples were collected with either an electric pump and hose or a Van Lorn water sampler. Additional samples were collected at 25-cm intervals from the reservoir surface to a 3-m depth in 1987, and from the 7-m depth to the reservoir bottom in 1982, 1986, 1987, and 1988. A small-volume Masterflex peristaltic pump and tubing (6-mm inside diameter) were used to collect these samples near the reservoir surface, and a pneumatically driven close-interval sampler (Blakar 1979) was used to collect samples between 7 m and the sediment surface at approximately 9 m.

13. River water samples for chemical analyses were collected weekly (or more frequently) at fixed stations located on the Eau Galle River, Lohn Creek, Lousy Creek, French Creek, and the reservoir tailwater. In 1987 and 1988, samples from the Eau Galle River and the tailwater were collected on nearly every working day throughout the summer. In addition, samples were collected on the Eau Galle River during 1987 at hourly intervals using an ISCO automated water sampler during all storm periods.

14. Samples for total soluble phosphorus (TSP) and soluble reactive phosphorus (SRP) were filtered through a  $0.45-\mu$  membrane filter before analysis. Samples collected under anoxic conditions were filtered in situ with

syringe filters (Nalge). Total phosphorus and TSP were determined colorimetrically on a Technicon Auto-Analyzer II following persulfate oxidation (APHA 1.85). Soluble reactive phosphorus was determined using the ascorbic acid method on either a Technicon Auto-Analyzer II or a spectrophotometer (APHA 1985). The detection limit for P analyses was 0.005 mg/l. Samples for chlorophyll a determination were filtered onto glass fiber filters and extracted in 90-percent alkaline acetone (APHA 1985). Chlorophyll a, corrected for phaeopigments, was calculated according to Lorenzen (1967). Statistical analyses were performed using the Statistical Analysis System (SAS 1985).

# Phosphorus Loading Determinations

15. The Eau Galle River and tailwater flows were gauged continuously except during 1981. Flows from French, Lousy, and Lohn Creeks were determined from continuous-recording gauges or stage height recorders during all years except 1981 and 1988. Reservoir level (for reservoir volume determination) was measured daily with a recording gauge. Daily precipitation (in centimeters) was obtained from the nearby Eau Galle Recreation Area, Resource Manager's Office.

16. A hydrological mass balance was determined for the summer stratified period for all years except 1981. The change in storage of water mass was assessed according to the following mass balance statement:

> Change in Storage = (Gauged Inflows + Precipitation) (1) - (Gauged Outflow) ± (Ungauged Inflows and Outflows)

Ungauged inflows and outflows collectively comprised ground water, overland runoff, and evaporation. These sources were not directly measured, but were determined from the above equation.

17. Daily external TP loadings, determined for 1987 and 1988, were calculated by multiplying the inflow rate by the measured TP concentration. Daily external TP loadings for 1982 and 1986 (sampled only weekly), and for unsampled days during 1987 and 1988, were estimated using regression analyses (Method 4, Walker 1985) and relationships between measured daily TP

concentration and flow determined for 1987. Internal TP loading was determined according to the following mass balance equation:

Internal TP loadings were calculated for the years 1981, 1982, 1986, 1987, and 1988. The reservoir TP mass was determined as the sum of volume-weighted TP mass of the water column. The TP mass was also calculated separately for the 0- to 3-m depths (i.e., epilimnion) and for the 3- to 9-m depths (i.e., metalimnion and hypciimnion).

# Phytoplankton Determinations

18. Phytoplankton sampling has been uninterrupted since 1981. During most years, epilimnetic water samples for phytoplankton identification and enumeration were collected at 2-week intervals between the onset of anoxia (May-June) and autumnal overturn (August-September). In 1986 these samples were collected weekly. During other seasons, sampling frequency varied between 2 and 6 weeks. In 1981, phytoplankton samples were collected from six stations. Since it was determined that the station located over the central depression was representative of the pelagic zone (Barko et al. 1984), only phytoplankton data from that station are reported here.

19. Integrated water samples were taken using a tubular collecting device, constructed from a 7.6-cm-diam plastic pipe and equipped with a one-way check valve. The check valve allowed filling of the pipe during vertical immersion and sample retention upon withdrawal from the water. Samples were taken to a depth of 3 m. Subsamples (125 ml) were preserved with acid-Lugol's solution (Vollenweider 1969) and stored in amber glass bottles at room temperature until processed. Taxonomic identification and enumeration of phytoplankton were accomplished using phase contrast inverted microscopes (Wild M40 and Leitz Diavert) in general accordance with Lund, Kipling, and LeCren (1958). Specific enumeration procedures are described in Barko et al. (1984).

20. Permanent slides of diatoms were prepared using Hyrax medium following digestion in boiling concentrated  $HNO_3$ . These slides were examined

using a Bausch and Lomb phase contrast microscope to estimate the relative abundance of different diatom taxa enumerated by inverted microscopy. The contribution of dead cells to the diatom counts was minor.

21. Cellular volume was computed from average cell dimensions determined for 25 randomly selected cells of the more common species from each sample (Munawar and Munawar 1976). Volume calculations assumed general conformity to the geometry most closely resembling that of specific algal cells (Vollenweider 1969). Cell volume was converted to fresh weight biomass assuming that the specific gravity of phytoplankton approximates 1.0 (Nauwerck 1963). Phytoplankton biomass is reported as grams per cubic meter fresh weight.

22. Species diversity (H'), after Shannon and Weaver (1963), was calculated based on fresh weight as follows:

$$H' = -\sum_{i=1}^{s} P_{i} * \ln P_{i}$$
(3)

where

 $P_i$  = proportion of the i<sup>th</sup> taxon in the sample, calculated from  $n_i/N$  $n_i$  = biomass of individuals per cubic meter of the i<sup>th</sup> taxon N = total biomass per cubic meter

Evenness (J), a measure of how closely H' approaches the maximum possible diversity, was calculated (after Pielou 1966) as

$$\frac{H'}{\ln S}$$
(4)

where

S = number of taxa present in sample
In S = maximum diversity

#### PART III: RESULTS

### Phosphorus Dynamics

23. Large between-year differences were observed in inflow, outflow, precipitation, hydraulic residence time, and flushing rate during the stratified period (Figure 2). The Eau Galle River was the major tributary input, accounting for nearly 80 percent of the total gauged inflow during all years. Mean daily gauged inflow (including the Eau Galle River and minor tributaries) and outflow were about three times higher in 1986 (the year of alum treatment) than in other years, due primarily to the large amount and frequent periods of precipitation that occurred during that year. Ungauged inflows and outflows were greater in 1986 than in other years. Also during that year, the hydraulic residence time was greatly reduced, and the flushing rate was elevated relative to all other years. Major freshets caused peaks in external TP loading from the Eau Galle River, primarily in 1986 but also during late summer of 1987 (Figure 3). Overall, external TP loading was relatively low during most of the summer of 1987. The lack of summer storm events in 1982 and 1988 (drought years) resulted in extremely low external TP loadings.

24. Eau Galle Reservoir commonly exhibited elevated concentrations of SRP in the hypolimnion before alum treatment, as in 1982 (Figure 4). Shortly after the occurrence of anoxic conditions, concentrations of SRP increased above the sediment surface, indicating P release from anoxic sediment. Steep vertical gradients in SRP were observed, and the concentration exceeded 1.00 mg P/ $\ell$  within the bottom meter from June through September of 1982. Hypolimnetic withdrawal appeared to result in lower SRP accumulation above the 7-m depth. Declines in SRP at intermediate depths in June and early August of 1982 coincided with decreases in the volume of anoxic water due to mixing.

25. Alum treatment resulted in low SRP throughout the water column from June until late July 1986 (Figure 4). Steep vertical gradients in SRP again developed below the 6-m depth during late August 1986. However, absolute SRP concentrations in the hypolimnion were much lower in 1986 than in 1982. Slight increases in SRP at intermediate depths (4 to 6 m) in mid-August 1986 coincided with high external TP loading (Figure 3). In 1987 and 1988, the patterns of SRP increase were very similar to those observed in 1982. Periods

of major external TP loading caused increases in SRP throughout the water column in September 1986, August 1987, and September 1988.

26. During the stratified periods of 1981 and 1982, internal TP loading frequently exceeded 10 mg/m<sup>2</sup>/day (Figure 5) and was 3 to 6 times greater than external TP loading (Table 1). During these years, internal TP loading was greatest when the zone of anoxia was large, and least during periods of mixing. The passage of cold fronts with sustained high winds caused mixing, which reduced the zone of anoxia (Figure 4). Total phosphorus discharge via withdrawal accounted for a large portion (20 to >100 percent) of the total TP load (i.e., internal plus external TP loading) before treatment (Figure 5).

27. Alum treatment completely negated internal TP loading from June to mid-July 1986 (Figure 5). During that period, negative internal TP loading values reflect losses through sedimentation and/or discharge. Unfortunately, separate sources (i.e., internal versus external) of TP discharged from the reservoir could not be determined. Positive internal TP loading resumed in late July and August 1986. In 1987 and 1988, internal TP loading was similar to pretreatment years (1981 and 1982) and accounted for 58 to 85 percent of the total TP load to the reservoir during summer stratification (Table 1).

28. Before alum treatment, increases in epilimnetic TP mass (Figure 6) corresponded with increased internal TP loading (Figure 5), as during mid-June and mid-July 1981 and during late July and late August 1982. Similarly, during the 1988 drought (after alum treatment), epilimnetic TP mass exhibited increases in June and July, during periods of elevated internal TP loading. In contrast, during late July and late August 1986, the year of alum treatment, external TP loading appears to have been the major contributor to TP mass increases in the epilimnion. In 1987, both internal and external TP sources appear to have contributed to increases in TP mass in the epilimnion. The TP mass in the epilimnion increased in early July of that year, corresponding to increased internal TP loading (Figure 5). However, during late July and early August 1987, increases in epilimnetic TP mass corresponded closely with periods of elevated external TP loading.

29. Before alum treatment, TP mass in the reservoir bottom waters (i.e., metalimnion and hypolimnion) generally increased during or shortly before corresponding increases in TP mass in the epilimnion (Figure 6), suggesting vertical transport of TP. The TP mass in the bottom waters was greater than in the epilimnion before alum treatment (Figure 6, 1981 and

1982). In 1986, however, TP mass in the bottom waters decreased substantially in relation to epilimnetic TP mass due to the effects of alum treatment. The epilimnion and the reservoir bottom waters exhibited similar TP masses during the early through mid-summer months of 1987 and 1988. During these same years, TP mass in the bottom waters exceeded that of the epilimnion from late July through August.

30. Peaks in epilimnetic (0 to 3 m) chlorophyll a concentration (Figure 7) corresponded both with peaks in epilimnetic TP mass (Figure 6) and with periods of elevated internal or external TP loading (Figures 3 and 5). Chlorophyll a increases in July of 1981, 1982, 1987, and 1988 were associated with periods of elevated internal TP loading. Increases in chlorophyll a during August of 1986 and 1987 coincided with periods of elevated external TP loading. In general, alum treatment had no significant effect on mean chlorophyll a concentration and mean TP mass of the epilimnion (Table 2).

# Phytoplankton Response

31. The seasonal distribution of phytoplankton biomass is presented in Figure 8 for the 8-year study period. Generally, a spring diatom (Bacillariophyceae) bloom was followed by a clear-water phase prior to the development of a large summer population dominated by dinoflagellates (Pyrrhophyta) and blue-green algae (Cyanophyta). Spring biomass maxima varied greatly in magnitude, duration, and timing. The largest occurred in April 1981 (66 g/m<sup>3</sup> fresh weight) and the smallest in March 1987 (6 g/m<sup>3</sup>). Frequently, a second diatom bloom developed during the autumn. Cryptophytes (Cryptophyta) were occasionally as important as diatoms in the spring (1984) and autumn populations (1983, 1984, 1986, and 1987).

32. Major summer biomass peaks occurred each year between about the third week in June and the end of September (Figure 8). In five of the studied years, biomass declined abruptly in October. However, in 1981 and 1982, substantial phytoplankton biomass remained beyond October, and in 1988, biomass in the form of a blue-green algae bloom (*Aphanizomenon flos-aquae*) was increasing when sampling ended in October. Maximum summer biomass ranged from 26 g/m<sup>3</sup> in late September 1984 to 89 g/m<sup>3</sup> in July and August 1985. Maximum summer biomass occurred as early as the third week of June 1988 and as late as the third week of September 1984. Average summer biomass ranged between

10 and 20 g/m<sup>3</sup> during all years except 1985, during which biomass was much greater at 51 g/m<sup>3</sup> (Figure 9). The dinoflagellates produced more biomass than the blue-green algae in all study years except 1981 and 1984 (Figure 8). Following reservoir drawdown during the winter of 1984, mean summer blue-green algae biomass was reduced by about one half in 1985. This reduction coincided with an increase in dinoflagellate biomass during the same year.

33. Between-year differences in phytoplankton succession during the summer growing season are summarized in Figure 10. Dinoflagellates and bluegreen algae have been the perennial summer dominant groups. Only did 1981 did blue-green algae establish dominance earlier in the season than did dinoflagellates. Transitions in community structure during the summer have demonstrated no consistently recurring patterns. However, between 1985 and 1987, dinoflagellates maintained dominance throughout the summer. In this reservoir, for all practical purposes, *Ceratium* biomass is synonymcus with dinoflagellate biomass.

34. A listing of the 136 phytoplankton taxa identified in the course of this 8-year study is provided as Appendix A. This is a record of the more abundant and commonly encountered species rather than an exhaustive list. Relatively few phytoplankton species accounted for most of the biomass during the year (Table 3). Throughout the study, 21 phytoplankton taxa attained biomass levels greater than 1 g/m<sup>3</sup> on at least one occasion. Of these, only the first eight species (Table 3) made consistent contributions during each study year.

35. Stephanodiscus hantzschii was by far the most important spring species, always accompanied by S. invisitatus, and periodically by Asterionella formosa and Cyclotella meneghiniana as codominants. Throughout the study, the single most successful summer (June-September) species was Ceratium hirundinella (Pyrrhophyta). Ceratium was the dominant or codominant (i.e., accounted for >10 percent of the biomass) in 88 percent of the summer samples. Overall, it occurred at concentrations >1 g/m<sup>3</sup> fresh weight in 89 of the 198 samples collected during the investigation.

36. Summer blue-green algae biomass was usually maintained through a succession of several species, i.e., Aphanizomenon flos-aquae -----> Anabaena planctonica ----> Oscillatoria agardhii. In 1987 and 1988, A. planctonica development preceded A. flos-aquae, but O. agardhii always followed the other

two. Aphanizomenon flos-aquae was the most important of the blue-green algae contributing to summer biomass. Autumn diatoms were most obviously represented by Fragilaria crotonensis, A. formosa, and occasionally by a reoccurrence of S. hantzschii.

37. Chlorophyta was the most taxonomically diverse algal group in the plankton, followed in decreasing order by the relatively less diverse groups, Bacillariophyceae and Cyanophyta (Table 4). Despite its high taxonomic diversity (Table 4), Chlorophyta was rarely an important contributor to community biomass (Table 3), and therefore had only a minor effect on the diversity index (H'). In contrast, the abundance of a single species, *C. hirundinella*, within the taxonomically nondiverse group Pyrrhophyta (Figure 4) was often so great that it controlled H' (see below).

38. During the 8-year study, summer phytoplankton diversity ranged from 0.05 (1986) to 2.3 (1984). Average summer diversity was greatest during the first 4 years but dropped sharply in 1985 during the extended and massive *Ceratium* bloom (Figure 11). Diversity tended to remain low after 1985, as *Ceratium* continued to account for more than 50 percent of the total phytoplankton biomass. *Ceratium* biomass accounted for somewhat greater variation in diversity than did total biomass. However, *Ceratium* biomass and total biomass were highly correlated throughout the study (r = 0.96, P < 0.001). Overall, there was a significant negative relationship between biomass and diversity (r = -0.60, P < 0.001).

39. Evenness (J), a measure of how closely H' approaches the theoretical maximum diversity on a scale of 0 to 1 (Pielou 1966), ranged from 0.18 to 0.73 in summer samples from all years. Mean summer evenness for each study year was always less than 0.5, i.e., 50 percent of maximum diversity (Figure 11), an indication that community diversity was usually well below its potential, further reflecting the importance of only a few species in the community.

# PART IV: DISCUSSION

40. Before alum treatment, both elevated epilimnetic TP mass and increases in chlorophyll *a* were common in Eau Galle Reservoir during periods when external TP loading was minimal, indicating that internal sources of TP were responsible for these increases. The occurrence of SRP increases above the profundal sediment surface following the onset of anoxia and the accumulation of SRP in the anoxic hypolimnion indicated that P release from sediments was an important mechanism of internal P loading. Disassociation of iron-phosphorus complexes under anoxic conditions has been suggested as an important mechanism of sediment P release in this reservoir (James, Kennedy, and Gaugush, in press). Investigations conducted in a variety of lakes have shown that P released from anoxic sediments can account for a substantial portion of the internal TP load and TP mass increase in the water column (Theis and McCabe 1978; Larson, Schultz, and Malueg 1981; Nurnberg 1984, 1987; Riley and Prepas 1984; Nurnberg et al. 1986).

41. In lakes where P concentration gradients are present in the metalimnion, P flux via turbulent eddy diffusion often constitutes an important source to the epilimnion (e.g., Imboden and Emerson 1978, Robards and Ward 1978, Stefan and Hanson 1981, Wodka et al. 1983). However, Stauffer (1985) found that this mechanism of P transport was not important in Green Lake, Wisconsin, during late summer because of the absence of P gradients in the metalimnion. The SRP and TP concentration gradients were minimal within the metalimnion (i.e., between approximately the 3- and 6-m depths, Figure 4) of Eau Galle Reservoir, due largely to withdrawal of cooler water during the stratified period. Thus, upward movement of P to the epilimnion via turbulent eddy diffusion was probably minor in this reservoir.

42. Increases in epilimnetic TP mass and chlorophyll *a* have been observed in Eau Galle Reservoir following periods of decreased thermal stability, when wind-driven mixing results in a downward depression of the thermocline (Gaugush 1984). Entrainment of P from deeper water or from metalimnetic shelf sediments is an important mechanism for P transport in Eau Galle Reservoir. Several investigations have reported the occurrence of windinduced metalimnetic migrations with accompanying increases in epilimnetic P and algal blooms (Stauffer and Lee 1973; Kortmann et al. 1982; Stauffer and Armstrong 1984; Stauffer 1985, 1987; Effler et al. 1986). Biological uptake

and movement of P to the epilimnion by vertically migrating phytoplankton (Salonen, Jones, and Arvola 1984; Osgood 1988; Taylor, Barko, and James 1988) may also be an important transport mechanism. As reported here, the vertically migrating dinoflagellate *Ceratium hirundinella* usually dominates the phytoplankton assemblage in this reservoir during the summer.

43. While alum treatment resulted in a substantial reduction in hypolimnetic TP and internal TP loading in Eau Galle Reservoir during 1986, the abnormally high frequency of major external TP loading events clearly subsidized the mass of epilimnetic TP. Epilimnetic TP mass and chlorophyll aremained essentially unchanged from other years. Whereas during most years, particularly before treatment, inflows were quite minor during the stratified period, an exceptionally high year for precipitation and inflow occurred in 1986. Elevated external TP loading clearly mitigated the effects of reduced internal TP loading (due to alum treatment) on epilimnetic TP mass in 1986. Had alum been applied during dry years (e.g., 1982 and 1988), when internal TP loading accounted for most of the TP mass in the reservoir, control over epilimnetic TP mass and therefore chlorophyll a (phytoplankton biomass) might have been more substantial than in 1986.

44. The effectiveness of alum treatment in controlling internal TP loading from profundal sediments in Eau Galle Reservoir was greatly diminished by 1987. We suggest that burial of alum by sedimentation associated with high riverine loadings in 1986 was a most important factor in shortening the effectiveness of alum treatment in this system. Eau Galle Reservoir has an average sedimentation rate of 2 to 3 cm/year in the deep basin (James and Barko, in press). The sedimentation rate during 1986 (unmeasured) was probably considerably greater than this average rate. Foy (1985) also suggested that sedimentation was responsible for the rapid recovery of hypolimnetic P concentrations to pretreatment levels following alum treatment in White Lough. In contrast, effective control of sediment P release and TP concentration in the water column of natural lakes with lower sedimentation rates has ranged from 3 to 8 years (Born 1979, Cooke et al. 1982, Garrison and Knauer 1983). As a result, these lakes have experienced reduced levels of algal production and a shift to less noxious species of algae.

45. Recent investigations indicate that littoral sediments may release substantial amounts of P under aerobic conditions, particularly at high pH (Twinch and Peters 1984, Drake and Heaney 1987). In addition, aquatic

macrophytes, prominent in Eau Galle Reservoir (Filbin and Barko 1985), can mobilize sediment P (e.g., Barko and Smart 1980, Carignan 1985, Smith and Adams 1986) that may then be transported to the open water following macrophyte decomposition (e.g., Carpenter 1980, Landers 1982). Prentki et al. (1979) suggested that movement of P from the littoral zone accounted for 59 percent of the soluble P load to the open water of Lake Wingra. Since the shallow-water sediments of the Eau Galle Reservoir were not treated in this study, it is possible that epilimnetic TP mass during low external loading periods was influenced by internal TP loading from these areas.

46. Ungauged sources of TP, particularly from ground-water discharge, may provide another internal source of P to the epilimnion of Eau Galle Reservoir. Indeed, hydraulic discharge from this reservoir has exceeded water income throughout our studies. This imbalance was particularly apparent during the relatively dry summers of 1982 and 1988 (Figure 2). Cooke et al. (1977, 1982) suggested that ground-water flow through P-rich sediments may be an important internal P-loading mechanism in West Twin and Dollar Lakes in Ohio.

47. Although internal P loading predominates during the summer stratified period in Eau Galle Reservoir, episodic freshets frequently result in substantial external P inputs to the epilimnion. Both P sources are important in sustaining high concentrations of TP and chlorophyll *a* in this reservoir. Conversely, internal P loading appears to be much more important than external P loading in natural lakes during the summer stratified period (e.g., Cooke et al. 1977; Larson, Schultz, and Malueg 1981; Stauffer 1987). In comparison with natural lakes, reservoirs receive water and material inputs from tributaries that drain a relatively larger watershed (Baxter 1977, Thornton et al. 1981). This difference results in generally greater nutrient and sediment loads to reservoirs, making them more difficult to manage than natural lakes via P control measures.

48. As with many reservoirs, Eau Galle Reservoir has a small surface area and volume relative to the size of its drainage basin. Consequently, flushing of the pool occurs regularly with the melting of snow and ice in the spring, and with storm events throughout the year. High periodic flushing rates combined with agricultural/dairy land-use practices and local geology ensure excessive nutrient loading to the system. As a consequence of these

factors, Eau Galle Reservoir remains eutrophic and supports a large phytoplankton population composed of nuisance species.

49. The seasonal succession of phytoplankton in Eau Galle Reservoir is fundamentally similar to that described for many temperate eutrophic lakes in which spring diatoms give way to blue-green or dinoflagellate summer populations, which in turn are replaced by diatoms in the autumn (Hutchinson 1967, Round 1981). The summer phytoplankton community structure in Eau Galle Reservoir is a eutrophic Pyrrhophyta/Cyanophyta association, a combination of Hutchinson's (1967) phytoplankton association categories.

50. No significant difference in phytoplankton biomass between pretreatment and posttreatment years was noted except in 1985. The high phytoplankton biomass in the summer of 1985 (90 g/m<sup>3</sup> wet weight) is remarkable from several points of view. Rarely have such high values been reported for any lake, and in no reported case has *Ceratium hirundinella* accounted for biomass of this magnitude (Nicholls, Kennedy, and Hammett 1980). Notably, the 1985 (*Ceratium*) bloom followed a year in which the late-summer phytoplankton community was totally dominated by blue-green algae.

51. The explanation for the abrupt change in dominance from blue-green algae to Ceratium in 1985 (persisting to date) may lie partially in the differences in the life histories of the dominant species involved, relative to the effects of winter drawdown in 1984. Akinetes (Aphanizomenon) and cysts (Ceratium) resting on the sediment surface provide a seasonal inoculum for these species (Wildman, Loescher, and Winger 1975; Heaney, Chapman, and Morison 1983; respectively). While the germination of cysts appears to require only oxygen and temperatures a greater than about 4° C (Heaney, Chapman, and Morison 1983; Loeblich and Loeblich 1984; Anderson, Taylor and Armbrust 1987), warmer water temperatures and a light cue, in addition to oxygen, are required for akinete germination (Wildman, Loescher, and Winger 1975). Thus, Ceratium has the potential to derive from sediments at any depth during periods of mixing. In contrast, due primarily to its requirement for light in akinete germination, Aphanizomenon is most likely to derive from sediments of the littoral zone during periods of mixing in Eau Galle Reservoir. As a result, the winter drawdown, which impacted sediments in the littoral zone but not in deeper water, probably had a much more damaging effect on Aphanizomenon than on Ceratium. In addition, since buoyant vegetative filaments of Aphanizomenon can persist well into the winter (Jones 1979; Wildman, Loescher, and Winger

1975), it is possible that direct loss of cells of this species accompanied loss (about 47 percent) of pool volume during winter drawdown.

52. The 1985 Ceratium bloom may have also been indirectly related to the winter drawdown. The drawdown had a direct effect on the submersed macrophyte community, which may in turn have affected the phytoplankton. Ceratophyllum demersum, which earlier accounted for 58 percent of the total macrophyte standing crop (Filbin and Barko 1985), was severely diminished in its distribution, standing crop, and frequency of occurrence (Godshalk and Barko 1988). This species has been shown to be particularly sensitive to cold exposure, and its growth is often at least temporarily retarded following reduction in water levels (e.g., Beard 1973, Nichols 1975). Ceratophyllum (without a true root system) and attached epiphytes derive their nutrient supply from the water column, and are therefore in direct competition for nutrients with phytoplankton.

53. In the years prior to the drawdown, the large standing stock of macrophytes and epiphytes formed a reservoir of P and other nutrients unavailable to the phytoplankton during much of the stratified period (Filbin and Barko 1985). In 1985, this nutrient sink was much reduced, perhaps providing greater access to nutrient supplies by the phytoplankton community.

54. As discussed above, abundant P is available to the phytoplankton community of Eau Galle Reservoir from multiple sources. Alum treatment clearly and significantly reduced P supply from profundal sediments in 1986 but had no effect (except perhaps relative to 1985) on phytoplankton production. Unfortunately, the greatest number of summer storm events during the entire study was recorded in 1986. Precipitation during June and July was 2 to 3 times greater than during these months in early years. As a result, high external TP loading during the stratified period of 1986 allowed the maintenance of high TP mass and chlorophyll in the epilimnion. Weatherrelated events in 1986 apparently counteracted the effects of alum treatment.

55. One of the unique features of reservoirs compared with natural lakes is their physical and chemical instability (Baxter 1977). Eau Galle Reservoir is particularly unstable in these regards, due to its relatively large watershed and its small pool volume. Superimposed upon this inherent instability were major extremes in climate (wet versus dry years) and a winter drawdown (1984) during the 8-year study period. In spite of these extremes, plus the additional effects of alum treatment (1986) on chemical

conditions/gradients, the phytoplankton community has been remarkably stable and of low species diversity. A shift in biomass toward a single species (*Ceratium*) after 1984 further decreased species diversity. *Ceratium* appears to be particularly well suited to existence in an unstable environment. The switch in relative dominance from a mixed dinoflagellate/blue-green algae community to one dominated by *Ceratium* is notable. However, through 1988 the status of these phytoplankton groups as codominants did not change. Without a decrease in P availability, *Ceratium* and blue-green algae are likely to persist as codominants in Eau Galle Reservoir, but perhaps will vary in relative abundance from year to year. 56. Since internal loadings of P in Eau Galle Reservoir probably cannot be controlled over the long term by alum application alone, other means of reducing the availability of P to phytoplankton need to be developed. For Eau Galle Reservoir and other systems with selective withdrawal features, the opportunity to release water from specific depths provides a potentially powerful tool for reconfiguring thermal structure. Particularly in shallow systems such as Eau Galle reservoir, thermal structure has an important influence on the movement of nutrients from bottom waters to the epilimnion.

57. At present, with hypolimnetic withdrawal, weak vertical gradients in water temperature lend Eau Galle reservoir to wind-driven mixing. As discussed earlier, episodic mixing events give rise to phytoplankton blooms by intermixing P-rich bottom waters with the epilimnion. We are currently investigating the effects of a change in withdrawal from bottom to surface during the summer. The rationale for this change in reservoir operation is that nutrient recycling associated with weak thermal structure and hypolimnetic warming (Gaugush 1984; James, Kennedy, and Gaugush, in press) has contributed to past phytoplankton blooms in Eau Galle reservoir (Barko et al. 1984).

58. Effects on physical conditions following change in withdrawal (scheduled for summer 1990) are anticipated to include decreased temperature of the hypolimnion, more stable thermal stratification, greater water column stability, lesser intermixing of bottom waters with the epilimnion, and presumably, reduced phytoplankton productivity. Ultimate effects on phytoplankton productivity will depend on species composition, which may also be altered by change in withdrawal.

59. During 1991 and 1992, a combined approach will be implemented in an attempt to manage phytoplankton population densities and species composition in Eau Galle reservoir. This approach will derive from results obtained thus far on independent effects of midwinter drawdown and alum treatment, and in addition, independent effects (to be determined) of change in reservoir operation from bottom to surface withdrawal. Concurrent with ongoing studies of phytoplankton dynamics, investigations of phosphorus cycling, with attention to various internal and external sources of input to the reservoir, will continue.

#### REFERENCES

Anderson, D. M., Taylor, C. D., and Armbrust, E. V. 1987. The effects of darkness and anaerobiosis on dinoflagellate cyst germination. Limnology and Oceanography 32:340-351.

APHA (American Public Health Association, American Water Works Association, and Water Pollution Control Federation). 1985. Standard methods for the examination of water and wastewater. 16th ed. Washington, DC.

Barko, J. W., and Smart, R. M. 1980. Mobilization of sediment phosphorus by submersed freshwater macrophytes. Freshwater Biology 10:229-238.

Barko, J. W., Bates, D. J., Filbin, G. J., Hennington, S. M., and McFarland, D. G. 1984. Seasonal growth and community composition of phytoplankton in a eutrophic Wisconsin impoundment. Journal of Freshwater Ecology 2:519-533.

Baxter, R. M. 1977. Environmental effects of dams and impoundments. Annual Review of Ecology and Systematics 8:255-283.

Beard, T. D. 1973. Overwinter drawdown. Technical Bulletin 61. Wisconsin Department of Natural Resources, Madison, WI.

Blakar, I. A. 1979. A close-interval sampler with minimal disturbance properties. Limnology and Oceanography 24:983-988.

Born, S. M. 1979. Lake rehabilitation: A status report. Environmental Management 3:145-153.

Carignan, R. 1985. Nutrient dynamics in a littoral sediment colonized by the submersed macrophyte *Myriophyllum spicatum*. Canadian Journal of Fisheries and Aquatic Sciences 42:1303-1311.

Carpenter, S. R. 1980. Enrichment of Lake Wingra, Wisconsin, by submersed macrophyte decay. Ecology 61:1145-1155.

Cooke, G. D., Heath, R. T., Kennedy, R. H., and McComas, M. R. 1978. Effects of diversion and alum application on two eutrophic lakes. EPA 600/3-78-033. US Environmental Protection Agency.

. 1982. Change in lake trophic state and internal phosphorus release after aluminum sulfate application. Water Resources Bulletin 18:699-705.

Cooke, G. D., and Kennedy, R. H. 1978. Effects of hypolimnetic application of aluminum sulfate to a eutrophic lake. Internationale Vereinigung fur Theoretische und Angewandte Limnologie, Verhandlungen 20:486-489.

Cooke, G. D., McComas, M. R., Waller, D. W., and Kennedy, R. H. 1977. The occurrence of internal phosphorus loading in two small, eutrophic, glacial lakes in northeastern Ohio. Hydrobiologia 56:129-135.

Drake, J. C., and Heaney, S. I. 1987. Occurrence of phosphorus and its potential remobilization in the littoral sediments of a productive English lake. Freshwater Biology 17:513-523.

Edmondson, W. T. 1970. Phosphorus, nitrogen, and algae in Lake Washington after nutrient diversion of sewage. Science 169:690-691.

Effler, S. W., Wodka, M. C., Driscoll, C. T., Brooks, C., Perkins, M., and Owens, E. M. 1986. Entrainment-based flux of phosphorus in Onondaga Lake. Journal of Environmental Engineering Division, Proceedings of the American Society of Civil Engineers 112:617-622.

Filbin, G. J., and Barko, J. W. 1985. Growth and nutrition of submersed macrophytes in a eutrophic Wisconsin impoundment. Journal of Freshwater Ecology 3:275-285.

Foy, R. H. 1985. Phosphorus inactivation in an eutrophic lake by direct addition of ferric aluminum sulfate: Impact on iron and phosphorus. Freshwater Biology 15:613-630.

Garrison, P. J., and Knauer, D. R. 1983. Lake restoration: A five-year evaluation of the Mirror and Shadow Lakes project Waupaca, Wisconsin. EPA 600/53-83-010. US Environmental Protection Agency, Corvallis, OR.

Gaugush, R. F. 1984. Mixing events in Eau Galle Lake. Pages 286-291 in Lake and Reservoir Management; Proceedings of the North American Lake Management Society, Knoxville, Tennessee, October 18-20, 1983. EPA 440/5-84-001. US Environmental Protection Agency, Washington, DC.

Godshalk, G. L., and Barko, J. W. 1988. Effects of winter drawdown on submersed aquatic plants in Eau Galle Reservoir, Wisconsin. Pages 100-111 <u>in</u> Proceedings, 22nd Annual Meeting, Aquatic Plant Control Research Program. Miscellaneous Paper A-88-5. US Army Engineer Waterways Experiment Station, Vicksburg, MS.

Heaney, S. I., Chapman, D. V., and Morison, H. R. 1983. The role of the cyst stage in the seasonal growth of the dinoflagellate *Ceratium hirundinella* within a small productive lake. British Phycological Journal 18:47-59.

Hutchinson, G. E. 1967. A Treatise on Limnology; II. Introduction to Lake Biology and the Phytoplankton. John Wiley and Sons, New York.

Imboden, D. M., and Emerson, S. 1978. Natural radon and phosphorus as limnologic tracers: Horizontal and vertical eddy diffusion in Greifensee. Limnology and Oceanography 23:77-90.

James, W. F., and Barko, J. W. Influences of submersed aquatic macrophytes on sediment accretion and composition in a north-temperate reservoir. Ecology (in press).

James, W. F., Kennedy, R. H., and Gaugush, R. H. Effects of large-scale metalimnetic migrations on phosphorus dynamics in a north-temperate reservoir. Canadian Journal of Fisheries and Aquatic Sciences (in press).

Jassby, A., and Powell, T. 1975. Vertical patterns of eddy diffusion during stratification in Castle Lake, California. Limnology and Oceanography 20:530-543.

Jones, R. I. 1979. Notes on the growth and sporulation of a natural population of Aphanizomenon flos-aquae. Hydrobiologia 62:55-58.

Kennedy, R. H. 1987. Material loadings to Eau Galle Lake. Pages 9-22 <u>in</u> Limnological Studies at Eau Galle Lake, Wisconsin; Report 2, Special Studies and Summary. R. H. Kennedy and R. C. Gunkel, eds. Technical Report E-85-2. US Army Engineer Waterways Experiment Station, Vicksburg, MS. Kennedy, R. H., and Cooke, G. D. 1982. Control of lake phosphorus with aluminum sulfate: Dose determination and application techniques. Water Resources Bulletin 18:389-395.

Kennedy, R. H., Barko, J. W., James, W. F., Taylor, W. D., and Godshalk, G. L. 1987. Aluminum sulfate treatment of a eutrophic reservoir: Rationale, application methods, and preliminary results. Lake Reservoir Management 3:85-90.

Kortmann, R. W., Henry, D. D., Kuether, A., and Kaufman, S. 1982. Epilimnetic nutrient loading by metalimnetic erosion and resultant algal responses in Lake Waramaug, Connecticut. Hydrobiologia 92:501-510.

Landers, D. H. 1982. Effects of naturally senescing aquatic macrophytes on nutrient chemistry and chlorophyll a of surrounding waters. Limnology and Oceanography 27:428-439.

Larson, D. P., Schultz, D. W., and Malueg, K. 1981. Summer internal phosphorus supplies in Shagawa Lake, Minnesota. Limnology and Oceanography 26:740-753.

Larson, D. P., Van Sickle, J., Malueg, K. W., and Smith, P. D. 1979. The effects of wastewater phosphorus removal on Shagawa Lake, Minnesota: Phosphorus supplies, lake phosphorus and chlorophyll- $\alpha$ . Water Research 13:1259-1272.

Loeblich, A. R., III, and Loeblich, L. A. 1984. Dinoflagellate cysts. Pages 443-480 in Dinoflagellates. D. L. Spector, ed. Academic Press, New York.

Lorenzen, C. J. 1967. Determination of chlorophyll and pheo-pigments: Spectrophotometric equations. Limnology and Oceanography 12:343-346.

Lund, J. W. G., Kipling, C., and LeCren, E. D. 1958. The inverted microscope method of estimating algal numbers and the statistical basis of estimations by counting. Hydrobiologia 11:143-170.

Mortimer, C. H. 1971. Chemical exchanges between sediments and water in the Great Lakes--Speculations on probable regulatory mechanisms. Limnology and Oceanography 16:387-404.

Munawar, M., and Munawar, I. F. 1976. A lakewide study of phytoplankton biomass and its species composition in Lake Erie, April-December 1970. Journal of the Fisheries Research Board of Canada 33:581-600.

Nauwerck, A. 1963. The relationships between zooplankton and phytoplankton in Lake Erken. Symbolae Botanicae Upsalienses 17:1-163.

Nicholls, K. H., Kennedy, W., and Hammett, C. 1980. A fish-kill in Heart Lake, Ontario, associated with the collapse of a massive population of *Ceratium hirundinella* (Dinophyceae). Freshwater Biology 10:553-561.

Nichols, S. A. 1975. The use of overwinter drawdown for aquatic vegetation management. Water Resources Bulletin 11:1137-1148.

Nurnberg, G. K. 1984. The prediction of internal phosphorus load in lakes with hypolimnetic anoxia. Limnology and Oceanography 29:111-124.

. 1987. A comparison of internal phosphorus loads in lakes with hypolimnetic anoxia: Laboratory incubations versus in situ hypolimnetic phosphorus accumulation. Limnology and Oceanography 32:1162-1164.

Nurnberg, G. K., Shaw, M., Dillon, P. J., and McQueen, D. J. 1986. Internal phosphorus load in an oligotrophic Precambrian Shield lake with an anoxic hypolimnion. Canadian Journal of Fisheries and Aquatic Sciences 43:574-580. Osgood, R. A. 1988. Hypothesis on the role of Aphanizomenon in translocating phosphorus. Hydrobiologia 169:69-76. Peterson, J., Wall, J. T., Wirth, T. L., and Born, S. M. 1973. Eutrophication control: Nutrient inactivation by chemical precipitation at Horseshoe Lake, Wisconsin. Technical Bulletin 62. Wisconsin Department of Natural Resources, Madison, WI. Pielou, E. C. 1966. The measurement of diversity in different types of biological conditions. Journal of Theoretical Biology 13:131-144. Prentki, R. T., Adams, M. S., Carpenter, S. R., Gasith, A., Smith, C. S., and Weiler, P. R. 1979. The role of submersed weedbeds in internal loading and interception of allochthonous materials in Lake Wingra, Wisconsin, USA. Archive fur Hydrobiologia (Supplement) 57:221-250. Riley, E. T., and Prepas, E. E. 1984. Role of internal phosphorus loading in two shallow, productive lakes in Alberta, Canada. Canadian Journal of Fisheries and Aquatic Sciences 41:845-855. Robards, R. D., and Ward, P. R. B. 1978. Vertical diffusion and nutrient transport in a tropical lake (Lake McIlwaine, Rhodesia). Hydrobiologia 59:213-221. Round, F. E. 1981. The Ecology of Algae. Cambridge University Press, Cambridge, UK. Ryding, S. O. 1981. Reversibility of man-induced eutrophication; Experiences of a lake recovery study in Sweden. Int. Rev. Gesamten. Hydrobiol. 66:449-502. Salonen, K., Jones, R. I., and Arvola, L. 1984. Hypolimnetic phosphorus retrieval by diel vertical migrations of lake phytoplankton. Freshwater Biology 14:439-442. SAS (Statistical Analysis System). 1985. Version 5 edition of the Statistical Analysis System. SAS Institute, Cary, NC. Shannon, C. E., and Weaver, W. 1963. The Mathematical Theory of Communication. University of Illinois Press, Urbana, IL. Smith, C. S., and Adams, M. S. 1986. Phosphorus transfer from sediments by Myriophyllum spicatum. Limnology and Oceanography 31:1312-1321. Soltero, R. A., Nichols, D. G., Gasperino, A. F., and Beckwith, M. A. 1981. Lake Restoration: Medical Lake, Washington. Journal of Freshwater Ecology 1:155-165. Stauffer, R. E. 1985. Nutrient internal cycling and trophic regulation of Green Lake, Wisconsin. Limnology and Oceanography 30:347-363. . 1987. Vertical nutrient transport and its effect on epilimnetic phosphorus in four calcareous lakes. Hydrobiologia 154:87-102. Stauffer, R. E., and Armstrong, D. E. 1984. Lake mixing and its relationship to epilimnetic phosphorus in Shagawa Lake, Minnesota. Canadian Journal of Fisheries and Aquatic Sciences 41:57-69.

Stauffer, R. E., and Lee, G. F. 1973. Role of thermocline migration in regulating algal blooms. Pages 73-82 in Modeling the Eutrophication Process. E. J. Middlebrooks, D. H. Falkenburg, and T. E. Maloney, eds. Ann Arbor Science Publishers, Ann Arbor, MI.

Stefan, H. G., and Hanson, M. J. 1981. Phosphorus recycling in five shallow lakes. Journal of Environmental Engineering Division, Proceedings of the American Society of Civil Engineers 107:713-730.

Taylor, W. D., Barko, J. W., and James, W. F. 1988. Contrasting diel pattorns of vertical migration in the dinoflagellate *Ceratium hirundineila* in relation to phosphorus supply in a north temperate reservoir. Canadian Journal of Fisheries and Aquatic Sciences 45:1093-1098.

Theis, T. L., and McCabe, P. J. 1978. Phosphorus dynamics in hypereutrophic lake sediments. Water Research 12:677-685.

Thornton, K. W., Kennedy, R. H., Carroll, J. H., Walker, W. W., Gunkel, R. C, and Ashby, S. 1981. Reservoir sedimentation and water quality - An heuristic model. <u>In</u> H. G. Stefan, ed., American Society of Civil Engineers, Proceedings of a Symposium on Surface Water Impoundments 1:654-661.

Twinch, A. J., and Peters, R. H. 1984. Phosphate exchange between littoral sediments and overlying water in an oligotrophic north-temperate lake. Canadian Journal of Fisheries and Aquatic Sciences 41:1609-1617.

Vollenweider, R. A., ed. 1969. A manual on methods for measuring primary production in aquatic environments. 2nd ed. IBP Handbook No. 12. Blackwell Scientific Publications, Oxford.

Walker, W. W. 1985. Empirical methods for predicting eutrophication in impoundments; Report 4, Phase 3: Applications manual. Technical Report E-81-9. US Army Engineer Waterways Experiment Station, Vicksburg, MS.

Welch, E. B. 1977. Nutrient diversion: Resulting lake trophic state and phosphorus dynamics. Ecological Research Series Report 600/3-77-003. US Environmental Protection Agency, Corvallis, OR.

Wildman, R. B., Loescher, J. H., and Winger, C. L. 1975. Development and germination of akinetes of *Aphanizomenon flos-aquae*. Journal of Phycology 11:96-104.

Wodka, M. C., Effler, S. W., Driscoll, C. T., Field, S. D., and Devan, S. P. 1983. Diffusivity-based flux of phosphorus in Onondaga Lake. Journal of Environmental Engineering Division, Proceedings of the American Society of Civil Engineers 109:1403-1430.

	to Eau Galle Res	ervoir Du	iring Midsum	mer Periods (June-	August)	
Year	Internal TP Loading mg m <sup>-2</sup> day <sup>-1</sup>	n	ANOVA*	External TP Loading mg m <sup>-2</sup> day <sup>-1</sup>	n	ANOVA
1981	9.31 (4.13)	7	a	2.76 (1.14)	8	b
1982	7.84 (2.87)	8	а	1.28 (0.01)	92	ъ
1986	-5.58 (6.21)	6	Ъ	17.01 (5.67)	85	а
1987	6.25 (2.60)	9	ab	4.89 (1.96)	65	ь
1988	8.97 (5.01)	8	а	1.29 (0.03)	85	Ъ

				Ta	able l				
Mean	(±1	Standard	Error)	Daily	Internal	and	External	TP	Loadings

Note: Alum was applied to the reservoir in 1986.

\* Analysis of variance column summarizes results of Duncan's Multiple Range Test. Different letters associated with individual variables indicate significant differences at the 5-percent level or less.

Year	Chlorophyll a mg m <sup>-3</sup>	<u>n</u>	ANOVA*	TP Mass kg	<u>n</u>	ANOVA
1981	35.99 (6.51)	7	a	126.70 (14.06)	7	а
1982	50.15 (10.90)	6	а	108.05 (13.48)	8	а
1986	28.79 (7.88)	7	а	148.27 (18.98)	8	а
1987	56.38 (12.22)	11	а	163.13 (22.00)	12	а
1988	40.22 (7.38)	8	а	110.28 (18.72)	9	а

Mean (±1 Standard Error) Chlorophyll a and TP Mass in Epilimnion (0 to 3 m) of Eau Galle Reservoir, June-August

Table 2

Note: Alum was applied to the reservoir in 1986.

ļ

\* Analysis of variance column summarizes the results of Duncan's Multiple Range Test. Different letters associated with individual variables indicate significant differences at the 5-percent level or less.

						Total	
			Biomass Cl	asses		Occurrences	
			g m_, fres	sh weight		>1 g ⊒ ,	
Species	Group*	>20	10-20	5-10	1-5	fresh weight	Season
Ceratium hirundinella	руг	21	20	19	2	89	Summer
Stephanodiscus hantzschii	bac	6	6	e	10	25	Spring, autumn
Aphanizomenon flos-aquae	cya	1	£	e	34	40	Summer
Oscillatoria agardhii	cya	;	1	1	11	13	Summer
Asterionella formosa	bac	1	I	ł	7	8	Spring, autumn
Anabaena planctonica	cya	ł	ł	12	16	28	Summer
Fragilaria crotonensis	bac	ł	1	1	17	18	Summer, autumn
Cryptomonas erosa	cry	1	ł	l	15	16	All seasons
Rhodomonas minuta	cry	1	1	ł	9	9	Spring, summer, autumn
Cyclotella meneghiniana	bac	ł	1	1	2	5	Spring
Stephanodiscus invisitatus	bac	1	1		4	4	Spring
Anabaena spiroides	cya	ł	1	1	3	3	Summer
Anabaena flos-aquae	cya	ł	ł	ł	2	2	Summer
Cryptomonas marssonii	cry	ł		ł	2	2	Spring, autumn
Melosira granulata var. angustissima	bac bac				1	2	Au tumn Summer
Rhizosolenia eriensis	bac	ł	1	ł	1	1	Autumn
Cyclotella pseudostelligera	bac	ł		ł	1	1	Spring
Chrysochromulina parva	chr	ł	!	ł	1	1	Spring
Pandorina morum	chl	1	1	1	1	1	Summer
Sphaerocystis schroeteri	chl	ł	1	1	1	1	Summer
<pre>* pry = Pyrrhophyta, bac - E</pre>	Sacillariophy	ceae, cya	= Cyanophyta,	cry = Cryp	tophyta, c	hr = Chrysophyta,	

Biomass Attained by Dominant Phytoplankton Species in

Table 3

Eau Galle Reservoir, 1981-1988

pry = Pyrrhophyta, bac - Bacillariophyceae, cya = Cyanophyta, cry = Cryptophyta, chr = Chrysophyta, chl = Chlorophyta.

Taxon	Number of Species	Number of Dominant Species*
Chlorophyta	53	2
Bacillariophyceae	34	8
Cyanophyta	27	5
Chrysophyta	9	1
Cryptophyta	7	4
Pyrrhophyta	3	1
Euglenophyta	3	0
	136	21

# Table 4 Number of Phytoplankton Species, by Major Group, Identified

\* A species was considered dominant if it occurred in at least one sample at a biomass >1 g m<sup>-3</sup> (see Table 3).

# in Eau Galle Reservoir, Wisconsin



Figure 1. Morphometric map of Eau Galle Reservoir. Contours indicate depth (in meters). Solid circle represents sampling site in the reservoir. French Creek, located upstream on the Eau Galle River, is not shown



Figure 2. Hydrologic data (±l standard error, n = 153) for Eau Galle Reservoir, May through September: (a) mean daily gauged inflow (includes Eau Galle River, Lousy, Lohn, and French Creeks, except in 1988 when the creeks were not gauged) and outflow, (b) mean daily precipitation, (c) mean daily ungauged (nonpoint) inflows and outflows, and (d) mean daily residence time and flushing rate

![](_page_36_Figure_0.jpeg)

Figure 3. Daily TP loading from the Eau Galle River, May-September 1982, 1986, 1987, and 1988

![](_page_37_Figure_0.jpeg)

Figure 4. Seasonal and depth-related variations in soluble reactive phosphorus ( $\mu g/l$ ). Measurements were made at close depth intervals (25 cm) during May-September 1982, 1986, 1987, and 1988. Shaded areas indicate zones in which dissolved oxygen was >0.5 mg/l. Black areas above panels indicate periods during which external TP loading exceeded 25 kg TP/day

![](_page_38_Figure_0.jpeg)

Figure 5. Seasonal variations in internal TP loading (upper panels) and TP discharge (lower panels) for the years 1981, 1982, 1986, 1987, and 1988. Dashed lines indicate an internal TP loading rate of zero. Black areas above panels indicate periods during which external TP loading exceeded 25 kg TP/day

![](_page_39_Figure_0.jpeg)

.

Figure 6. Seasonal variations in TP mass within the epilimnion (0 to 3 m) and bottom waters (3 to 9 m) during 1981, 1982, 1986, 1987, and 1988. Black areas above each panel indicate periods during which external TP loading exceeded 25 kg TP/day

![](_page_40_Figure_0.jpeg)

Figure 7. Seasonal variations in epilimnetic (0 to 3 m) chlorophyll a concentrations during 1981, 1982, 1986, 1987, and 1988. Black areas above each panel indicate periods during which external TP loading exceeded 25 kg TP/day

![](_page_41_Figure_0.jpeg)

![](_page_41_Figure_1.jpeg)

![](_page_42_Figure_0.jpeg)

![](_page_42_Figure_1.jpeg)

![](_page_43_Figure_0.jpeg)

![](_page_43_Figure_1.jpeg)

Figure 10. Summary of between-year differences in seasonal succession of *Ceratium* and blue-green algae

![](_page_44_Figure_0.jpeg)

![](_page_44_Figure_1.jpeg)

# APPENDIX A PHYTOPLANKTON SPECIES LIST, EAU GALLE RESERVOIR (1981-1988)

#### BACILLARIOPHYCEAE

Achnanthes minutissima Kuetzing Amphora sp. Asterionella formosa Hassall Cocconeis placentula v. lineata (Ehr.) Van Heurck Cyclotella meneghiniana Kuetzing pseudostelligera Hustedt stelligera Cleve & Grunow Cymatopleura solea (Bréb.) W. Smith Cymbella sp. Fragilaria capucina Desmazieres capucina v. mesolepta Rabenhorst crotonensis Kitton vaucheriae (Kuetz.) Petersen Gomphonema sp. Gyrosigma spencerii (W. Smith) Cleve Melosira crenulata (Ehr.) Kuetzing granulata (Ehr.) Ralfs granulata v. angustissima Mueller varians C. A. Agardh Navicula minima Grunow Nitzschia acicularis W. Smith actinastroides (Lemm.) Van Goor palea (Kuetz.) W. Smith sigmoidea (Ehr.) W. Smith vermicularis (Kuetz.) Grunow Rhizosolenia eriensis H. L. Smith Stephanodiscus hantzschii Grunow invisitatus Hohn & Hellerman

Synedra acus Kuetzing delicatissima v. angustissima Grunow rumpens Kuetzing ulna (Nitz.) Ehrenberg ulna v. danica (Kuetz.) Van Heurck ulna v. subaequalis (Grun.) Van Heurck CHLOROPHYTA Actinastrum hantzschii Lagerheim hantzschii v. fluviatile Schroeder Ankistrodesmus convolutus Corda falcatus (Corda) Ralfs falcatus v. mirabilis (W. & W.) G. S. West gelifactum (Chod.) Bourrelly nannoselene Skuja Carteria sp. Chlamydomonas snowii Printz Chlorogonium sp. Closteriopsis sp. Closteridium sp. Closterium sp. Coelastrum cambricum Archer microporum Naegeli sphaericum Naegeli Cosmarium sp. Crucigenia truncata G. M. Smith Dictyosphaerium ehrenbergianum Naegeli pulchellum Wood Elakatothrix gelatinosa Wille

# APPENDIX A (Continued)

Eudorina elegans Ehrenberg Gloeocystis planctonica (W. & W.) Lemmermann Kirchneriella sp. Micractinium pusillum Fresenius pusillum v. elegans G. M. Smith Mougeotia sp. Oedogonium sp. Oocystis sp. Pandorina morum (Muell.) Bory Pediastrum boryanum (Turp.) Meneghini duplex v. clathratum (A. Braun) Lagerheim duplex v. reticulatum Lagerheim Quadrigula chodatti (Tanner-Fullman) B. M. Smith lacustris (Chod.) G. M. Smith Scenedesmus abundans (Kirch) Chodat acuminatus (Lag.) Chodat arcuatus v. platydisca G. M. Smith bijuga (Turp.) Lagerheim dimorphus (Turp.) Kuetzing dispar Brébisson obliquus (Turp.) Kuetzing quadricauda (Turp.) Brébisson quadricauda v. longispina (Chod.) G. M. Smith quadricauda v. parvus G. M. Smith Schroederia judayi G. M. Smith setigera (Schroed.) Lemmermann Selenastrum minutum (Naeg.) Collins Sphaerocystis schroeteri Chodat Staurastrum paradoxum Meyen Tetraëdron sp. Tetrastrum staurogeniaeforme (Schroed.) Lemmermann Treubaria setigerum (Archer) G. M. Smith

#### CHRYSOPHYTA

Chrysochromulina parva lackey Chrysococcus refuscens klebs Dinobryon sp. Kephyrion sp. Mallomonas sp. Ochromonas sp. Syncrypta volvox Ehrenberg Synura uvella Ehrenberg Stenokalyx inconstans G. Schmid

### CRYPTOPHYTA

Chroomonas sp. Cryptomonas erosa Ehrenberg marssonii Skuja ovata Ehrenberg Ketablepharis ovalis Skuja Rhodomonas minuta Skuja minuta v. nannoplanctica Skuja

### CYANOPHYTA

```
Anabaena
affinis Lemmermann
flos-aquae (Lyngb.) Brébisson
planctonica Brunnthaler
spiroides Klebahn
spiroides
v. crassa Lemmermann
Anabaenopsis
raciborskii Woloszynska
```

# APPENDIX A (Concluded)

#### CYANOPHYTA (Continued)

Aphanocapsa delicatissima West & West pulchra (Kuetz.) Rabenhorst Aphanizomenon flos-aquae (L.) Ralfs Aphanothece nidulans P. Richter Chroococcus minimus (Keissl.) Lemmermann Coelosphaerium dubium Grunow naegelianium Unger pallidum Lemmermann Dactylococcopsis raphidioides Hansgirg Gomphosphaeria lacustris v. compacta Lemmermann aponina v. delicatula Virieux Lyngbya sp. Merismopedia minima Beck tenuissima Lemmermann Microcystis aeruginosa Kuetz.; emend. Elenkin incerta Lemmermann Oscillatoria angustissima West & West agardhii Gomont limnetica Lemmermann Phormidium mucicola Naumann & Huber-Pestalozzi Spirulina sp.

#### EUGLENOPHYTA

Euglena sp. Phacus pyrum (Ehr.) Stein Trachelomonas sp.

#### PYRRHOPHYTA

Ceratium hirundinella (O.F. Muell.) Schrank Glenodinium sp. Gymnodinium sp.