

12

**CHEMICAL
RESEARCH,
DEVELOPMENT &
ENGINEERING
CENTER**

CRDEC-TR-225

AD-A229 664


**COMPARATIVE NUCLEOPHILIC SUSCEPTIBILITY
OF SELECTED PHOSPHORO- AND PHOSPHONOTHIOLATES
BY SEMI-EMPIRICAL MOLECULAR ORBITAL CALCULATIONS**

**George R. Famini
RESEARCH DIRECTORATE**

**Philip W. Bartram
PHYSICAL PROTECTION DIRECTORATE**

September 1990

**DTIC
ELECTE
DEC 11 1990**
S E D
cc



**U.S. ARMY
ARMAMENT
MUNITIONS
CHEMICAL COMMAND**

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

Aberdeen Proving Ground, Maryland 21010-5423

9 11 098

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

Distribution Statement

Approved for public release; distribution is unlimited.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1990 September	3. REPORT TYPE AND DATES COVERED Final, 90 Apr - 90 Apr	
4. TITLE AND SUBTITLE Comparative Nucleophilic Susceptibility of Selected Phosphoro- and Phosphonothiolates by Semi-empirical Molecular Orbital Calculations			5. FUNDING NUMBERS PR-1C162622A553F	
6. AUTHOR(S) Famini, George R., and Bartram, Philip W.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) CDR, CRDEC, ATTN: SMCCR-RSP-C/SMCCR-PPD, APG, MD 21010-5423			8. PERFORMING ORGANIZATION REPORT NUMBER CRDEC-TR-225	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The electronic properties of several phosphorothiolates were compared against the target molecule O-ethyl-S-(2-diisopropylaminoethyl)-methylphosphonothiolate (VX) by hierarchical clustering techniques and electrostatic potential maps to determine the best match for nucleophilic attack. Semi-empirical calculations and geometry optimization were accomplished with the MNDO algorithm in MOPAC v5.0. Graphical display of structures and electrostatic potentials were produced, using the Chemo-metric and Biometric Modeling (CBM) Branch's Molecular Modeling, Analysis and Display System (MMADS). The pattern recognition program ARTHUR was used to predict the hierarchical clustering of the electronic properties using Euclidean distance criteria. Both hierarchical clustering and electrostatic potential differences contouring predict that the molecule O-ethyl S-ethyl methylphosphonothiolate most closely resembles VX in susceptibility to nucleophilic displacement.				
14. SUBJECT TERMS Molecular orbital theory. Hierarchical clustering Hydrolysis of VX. Quantum chemistry			15. NUMBER OF PAGES 25	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

BLANK

PREFACE

The work described in this report was authorized under Project No. 1C162622A553F, CB Decontamination and Contamination Avoidance. This work was started and completed in April 1990.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, U.S. Army Chemical Research, Development and Engineering Center (CRDEC), ATTN: SMCCR-SPS-T, Aberdeen Proving Ground, Maryland 21010-5423. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce the document for U.S. Government purposes.

This report has been approved for release to the public.

Acknowledgments

The authors thank Melissa Parkinson, Physical Protection Directorate, CRDEC, for her assistance in the preparation of this report.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



BLANK

CONTENTS

	Page
1. INTRODUCTION	9
1. Problem Definition	9
1.2 Potential Simulants	10
1.3 Nucleophilic Displacement In VX	10
1.4 Theoretical Chemistry Applications	10
2. COMPUTATIONAL PROCEDURE	13
3. RESULTS AND DISCUSSION	13
3.1 Hierarchical Clustering	14
3.2 Electrostatic Potentials	14
4. SUMMARY	20
LITERATURE CITED	25

LIST OF FIGURES

1	VX	9
2	Simulant 1, O-Methyl-S-methyl Aminophosphorothiolate	11
3	Simulant 2, O-Methyl-S-methyl-N-Acetyl Amidophosphorothiolate	11
4	Simulant 3, O-Ethyl-O-chloro-bromophenyl-S-propyl Phosphorothiolate	11
5	Simulant 4, O-Ethyl-S-ethyl Methylphosphonothiolate	12
6	Simulant 5, Bis(O-ethyl)-S-ethyl Phosphorothiolate	12
7	Simulant 6, Bis(O-methyl)-chlorophenyl Phosphorothiolate ...	12
8	Simulant 7, Triethyl Phosphate	12
9	Dendrogram for Seven Electronic Properties	16
10	Dendrogram for Six Electronic Properties	16
11	Dendrogram for One Electronic Property	17
12	Electrostatic Map/VX	18
13	Electrostatic Map/Simulant 1	18
14	Electrostatic Map/Simulant 2	18
15	Electrostatic Map/Simulant 3	18
16	Electrostatic Map/Simulant 4	19
17	Electrostatic Map/Simulant 5	19
18	Electrostatic Map/Simulant 6	19
19	Electrostatic Map/Simulant 7	19
20	Difference Map/VX and Simulant 1	21
21	Difference Map/VX and Simulant 2	21
22	Difference Map/VX and Simulant 3	21
23	Difference Map/VX and Simulant 4	21
24	Difference Map/VX and Simulant 5	22
25	Difference Map/VX and Simulant 6	22

26 Difference Map/VX and Simulant 7 22
27 Difference Map/Simulant 4 and Simulant 5 20

LIST OF TABLES

1 MNDO Electronic Properties 15
2 Simulant Ranking 23

BLANK

COMPARATIVE NUCLEOPHILIC SUSCEPTIBILITY
OF SELECTED PHOSPHORO- AND PHOSPHONOTHIOLATES
BY SEMI-EMPIRICAL MOLECULAR ORBITAL CALCULATIONS

1. INTRODUCTION

1.1 Problem Definition.

O-ethyl-S-(2-diisopropylaminoethyl)-methylphosphonothiolate (VX) (Figure 1) is a potent inhibitor of the enzyme acetylcholinesterase (AChE). The inhibition mechanism involves phosphorylation of the esteratic site in the enzyme. Because of the lethal toxicity of VX, severe restrictions have been placed on its use in laboratory experimentation.




Figure 1. VX

In an effort to circumvent many difficulties involving the use of chemical agents in the laboratory, the U.S. Army Chemical Research, Development and Engineering Center (CRDEC) has been the lead organization for the development and use of chemical agent simulants (conversation with William White, Research Directorate, CRDEC, May 1990). Simulants are chemical compounds that possess similar properties and/or activities as an agent but are non-toxic. In this way, simulants will behave similarly to agents in a given experiment but require none of the precautions associated with the use of toxic chemicals. To promote the use of simulants and provide a centralized location to find properties regarding simulants, CRDEC created, as well as maintains, the Chemical Agent Simulant Data Center.¹



One area with little success in developing simulants has been in simulating reactivity, either oxidative or hydrolytic. This study will present a new approach to examining this problem.

1.2 Potential Simulants.

Figures 2-8 display the potential simulants that have been examined in this study. The compounds examined with the exception of O-ethyl-S-ethyl methylphosphonothiolate (EA-5533) (Figure 5) are common pesticides.² Research Directorate, CRDEC, (conversation with Leonard Szafraniec, Research Directorate, CRDEC, May 1990), synthesized EA-5533.

1.3 Nucleophilic Displacement In VX.

VX detoxification can cause a number of different mechanisms. This study will focus only on the nucleophilic displacement of the alkylthio group by a nucleophilic attack at the phosphorus. The products obtained by this attack are a function of the strength of the nucleophile, pH, and solvent. The VX reaction with DS2 (sodium hydroxide, methoxyethanol, diethylenetriamine) involves the substitution of the 2-(bis-isopropylamino) ethylthiol group by the 2-methoxyethoxide (OMC) (equation 1). In water/alcohol solutions, competitive substitution by the alcohol can occur (equation 2).³



1.4 Theoretical Chemistry Applications.

Computational chemistry is extensively used in the examination of chemical compound properties. Although a number of computational methods have been used successfully to predict physical and biological properties, prediction of chemical reactivity has been less successful.^{4,5,6} Theoretical methods allow for the facile computation of numerous electronic properties of




Figure 2. Simulant 1, O-Methyl-S-Methyl
: Aminophosphorothiolate




Figure 3. Simulant 2, O-Methyl-S-Methyl-
N-Acetyl Amidophosphorothiolate




Figure 4. Simulant 3, O-Ethyl-O-Chloro-
Bromophenyl-S-Propyl
Phosphorothiolate




Figure 5. Simulant 4, O-Ethyl-S-Ethyl Methylphosphonothiolate




Figure 6. Simulant 5, Bis(O-Ethyl)-S-Ethyl Phosphorothiolate




Figure 7. Simulant 6, Bis(O-Methyl)-Chlorophenyl Phosphorothiolate




Figure 8. Simulant 7, Triethyl Phosphate

molecules that can be used to infer reactivity and used to further compute reaction related properties. In effect, the ground state electronic and steric attributes are used to qualitatively predict how well a reaction can occur.

One difficulty in the application of molecular orbital calculations is the sheer amount of electronic data that can be generated. Two approaches to apply molecular orbital calculations have been examined in this study. The first approach uses hierarchical clustering techniques to find the compound that best matches those electronic properties expected to influence the nucleophilic displacement. In the second approach, molecular graphics have been applied to condense the plethora of numerical data into an easier to comprehend description of the electrostatic potential. The electrostatic potential has been used successfully in this laboratory and by others to qualitatively assess the reactivity of a substrate.⁷

2. COMPUTATIONAL PROCEDURE.

All geometry optimizations and calculations of electronic properties have been performed using the MNDO algorithm as contained in MOPAC v5.0 on a Stardent Titan super workstation.^{8,9} Graphical display of the electrostatic potentials and molecular structures were done using the Molecular Modeling Analysis and Display System (MMADS) program.¹⁰ The pattern recognition program, ARTHUR, was used to perform all hierarchical clustering.¹¹

3. RESULTS AND DISCUSSION

Two methods were employed to determine if theoretical and computational methods could be useful in selecting simulants. If the important properties that determine the reactivity of VX with respect to nucleophilic displacement could be adequately matched by a potential simulant, then the reactivities would likely be similar. However, the important properties in hydrolysis were not intuitively clear. Seven theoretically determined electronic properties were calculated and used to attempt to model reactivity (described in section 3.1). Using pattern recognition methods, it is possible to determine which of the potent simulants best match all or some of the electronic properties of the agent. A second approach using the molecular electrostatic potential (MEP) was also employed. In particular, the difference between the MEP of the simulant and VX was examined. The reactivities will more likely be similar as the MEP map approach exact overlap.

*Famini, George R., "A Computational Comparison of the Electronic Properties of G Agents with Selected Reaction Simulants," In Proceedings of the 1989 Scientific Conference on Chemical Defense Research, CRDEC-SP-024, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, September 1990, unpublished data.

3.1 Hierarchical Clustering.

The electronic properties, calculated by the MNDO algorithm, expected to contribute to the electrophilic properties in VX are as follows:

- heat of formation,
- the energy of the highest occupied molecular orbital,
- the energy of the lowest unoccupied molecular orbital,
- the electronic charge of the phosphorus,
- the electronic charge of the sulfur atom,
- the electronic charge of the P - O oxygen, and
- the dipole moment.

These electronic properties, the values for VX, and the seven simulants are listed in Table 1.

Three dendrograms were produced by the pattern recognition package ARTHUR. The first dendrogram (Figure 9) compares all seven electronic properties for VX and the simulants. The dendrogram shows that Simulant 4, O-ethyl-S-ethyl methylphosphonothiolate is the best fit for VX. Simulants 3, 6, 5, and 1 are the second best choices; Simulant 2 is the third best choice. A second dendrogram (Figure 10) compares six (the heat of formation was not considered) of the Modified Neglect of Diatomic Differential Overlap (MNDO) electronic properties for VX and the simulants.

Simulants 1, 3, 4, 5, and 6 were the first choice for the best fit and Simulant 2 the second choice. A third dendrogram compares only the energy of the orbit (ELUMO), because the Frontier Molecular Orbital Theory predicts that this property is the most significant in Sn2 chemistry. For the ELUMO dendrogram (Figure 11), the ranking is Simulant 4 followed by Simulant 1 for the best fit for VX.

3.2 Electrostatic Potentials.

Electrostatic potentials have been shown to be a good indicator of relative reactivity due to attraction and repulsion of point charges. The Molecular electrostatic Potential (MEPs) were calculated using a modified coulombic expression as shown in equation 3.

$$V_{app} = \sum_A \frac{q_a}{R_A - r} \quad (3)$$

- V_{app} is the approximate electrostatic potential
- q_a is the formal charge on atom A
- R_A is the position of the atom
- r is the electron at point r

Electrostatic potentials for VX and the simulants are shown in Figures 12-19. The planes displayed are the phosphorus, the double bonded oxygen, and the leaving group. The significant region of the plot is the area

Table 1. MNDO Electronic Properties

Compound	H	Ehomo	Elumo	qP	qO	qS	DM
VX	-88.27250	-9.30036	-1.73961	.8413	-.5898	-.1487	3.606
1	-90.44274	-10.44339	-1.93622	.9973	-.6014	-.0850	3.270
2	-129.02166	-10.41193	-2.15472	1.0980	-.5520	-.1365	6.104
3	-123.78412	-9.44083	-2.30574	1.0757	-.6325	.0108	1.667
4	-103.26889	-10.10178	-1.63999	.8195	-.6320	-.0890	1.850
5	-154.70651	-10.50167	-2.03172	1.0823	-.6091	-.0437	3.005
6	-115.91889	-9.80965	-2.23556	1.0490	-.6367	.0927	1.788
7	-209.08610	-11.11833	-0.83351	1.3962	-.6400	-.4998	2.918

H = Heat of formation

Ehomo = Energy of the highest occupied molecular orbital

Elumo = Energy of the lowest unoccupied molecular orbital

qP = Charge on the phosphorus atom

qO = Charge on the oxygen atom

qS = Charge on the sulfur atom

DM = Dipole moment

Simulant 1, O-Methyl-S-methyl aminophosphorothiolate
 Simulant 2, O-Methyl-S-methyl-N-acetyl amidophosphorothiolate
 Simulant 3, O-Ethyl-O-chlorobromophenyl-S-propyl phosphorothiolate
 Simulant 4, O-Ethyl-S-ethyl methylphosphonothiolate
 Simulant 5, Bis(O-Ethyl)-S-ethyl phosphorothiolate
 Simulant 6, Bis(O-methyl)-chlorophenyl phosphorothiolate
 Simulant 7, Triethyl phosphate




Figure 9. Dendrogram for Seven Electronic Properties




Figure 10. Dendrogram for Six Electronic Properties




Figure 11. Dendrogram for One Electronic Property




Figure 12. Electrostatic Map/VX




Figure 13. Electrostatic Map/Simulant 1




Figure 14. Electrostatic Map/Simulant 2




Figure 15. Electrostatic Map/Simulant 3




Figure 16. Electrostatic Map/Simulant 4




Figure 17. Electrostatic Map/Simulant 5




Figure 18. Electrostatic Map/Simulant 6




Figure 19. Electrostatic Map/Simulant 7

180° to the leaving group. This significant region is shown as solid contour lines and represents a positive area attractive to a negative charge. Simulants that best fit the reactivity of VX will have a positive region similar in shape and size.

Computer generated electrostatic potential difference maps have been made between VX and each of the simulants, thus simplifying the comparison. The simulants that have the best electrostatic overlaps will result in difference maps with the smallest positive lobe opposition to the leaving group. The electrostatic overlap maps are shown in Figures 20-26.

Examination of the difference map (Figure 23) shows that Simulant 4, O-ethyl-S-ethyl methylphosphonothiolate, would closely resemble VX in nucleophilic reactivity. However, Simulant 1, O-methyl-S-methyl aminophosphorothiolate, and Simulant 5, O-ethyl-O-ethyl S-ethyl phosphorothiolate (Figures 20 and 24, respectively), may provide reasonable reactivity.

Simulant 4 (EA-5533) and Simulant 5 were compared (Figure 27) to determine the difference in phosphorus reactivity between a phosphonothiolate and a similar phosphorothiolate. The large positive lobe shown indicates a large difference in reactivity between the two phosphorus.




Figure 27. Difference Map/Simulant 4 and Simulant 5

4. SUMMARY

Hierarchical clustering and electrostatic potential techniques were used to predict the similarity in reactivity between several potential phosphorothiolate simulants and VX. The technique considered the S_N2 displacement of a leaving group attached to phosphorus. Two compounds, O-methyl-S-methyl aminophosphorothiolate and O-ethyl-S-ethyl methylphosphonothiolate were identified as suitable simulants. Table 2 summarizes the ranking of simulants for MEP and hierarchical clustering.




Figure 20. Difference Map/VX
and Simulant 1




Figure 21. Difference Map/VX
and Simulant 2




Figure 22. Difference Map/VX
and Simulant 3




Figure 23. Difference Map/VX
and Simulant 4




Figure 24. Difference Map/VX
and Simulant 5



Figure 25. Difference Map/VX
and Simulant 6




Figure 26. Difference Map/VX
and Simulant 7

Table 2. Simulant Ranking

<u>Heirarchical Clustering</u>				
Rank	7 Electrostatic Properties	6 Electrostatic Properties	1 Electrostatic Property	MEP
1	simulant 4	simulants 1,3,4,5,6	simulant 4	simulant 4
2	simulants 1,3,5,6	simulant 2	simulants 1,2,3,5,6	simulant 1

BLANK

LITERATURE CITED

1. Coon, P.A., Famini, G.R., White, W.E., and Thornton, R.C., Data Base User's Guide for the Chemical Agent Simulant Data Center, CRDEC-SP-88032, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, November 1988, UNCLASSIFIED Report.
2. Eto, Morifusa, Organophosphorus Pesticides: Organic and Biological Chemistry, CRC Press, Cleveland, OH, 1974.
3. Szafraniec, L., Beaudry, W., Rohrbaugh, D. and Ward, J., "Stoichiometry of Reaction of GA, GB, VX, and HD in Mixed Aqueous Solution," In Proceedings of the 1988 U.S. Army Chemical Research, Development and Engineering Center Scientific Conference on Chemical Defense Research, Vol. I, CRDEC-SP-013, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, February 1988, UNCLASSIFIED Report.
4. Kamlet, M.J., and Taft, R.W., "Linear Free Energy Relationships. Local Empirical Rules or Fundamental Laws of Chemistry," Acta. Chem. Scand. Vol. 1339, pp 611-628 (1985).
5. Hansch, C., "A Quantitative Approach to Biochemical Structure Activity Relationship," Acc. Chem. Res. Vol. 2, p 232 (1969).
6. Famini, G.R., "Using Theoretical Descriptors in Structure Activity Relationships. V. A Review of the Theoretical Parameters," CRDEC-TR-085, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, July 1989, UNCLASSIFIED Report.
7. Politzer, P., and Truhlar, D., Chemical Applications of Molecular Electrostatic Potentials, Plenum Press, New York, NY, and London, England, 1981.
8. Dewar, M.J.T., and Thiel, W., "Ground States of Molecules 38. The MNDO Method," J. Am. Chem. Soc. Vol. 99, p 4899 (1977).
9. Steward, J.J.P., MOPAC Manual (5th Edition), FJSRL-TR-88-0007, Frank J. Seiler Research Lab, U.S. Air Force Academy, Colorado Springs, CO, December 1988, UNCLASSIFIED Report.
10. Leonard, J.M., and Famini, G.R., A User's Guide to the Molecular Modeling Analysis and Display System (2nd Edition), CRDEC-TR-030, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, January 1989, UNCLASSIFIED Report.
11. ARTHUR, Informetrix, Inc., Seattle, WA, 1982.