

4

REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

tion is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this reporting burden, including this burden estimate, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Washington, DC 20540, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

AD-A230 022

2. REPORT DATE  
11/5/90

3. REPORT TYPE AND DATES COVERED  
Technical Rept. (Interim) 9/30/90

4. TITLE AND SUBTITLE

Dynamics of Solitary Waves Induced by Shock Impulses in a Linear Atomic Chain

5. FUNDING NUMBERS

C: N00014-89-WX-24146  
N00014-90-WX-24103  
PE: 0601153N  
PR: RR011-03-OK  
TA: 412n006---02  
WU: 61-3299-00

6. AUTHOR(S)

Phuoc X. Tran, D.W. Brenner and C.T. White

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Naval Research Laboratory  
Washington, DC 20375-5000  
(Code 6119, C.T. White)

8. PERFORMING ORGANIZATION REPORT NUMBER

Technical Report #10

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000  
(Code 1112AI, D.H. Liebenberg)

10. SPONSORING / MONITORING AGENCY REPORT NUMBER

Technical Report #10

SDTIC  
ELECTE  
DEC 13 1990  
S E D

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT

This document has been approved for public release and sale, distribution of this document is unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

The propagation of nonlinear waves induced by a shock impulse in a linear atomic chain of finite but large length is investigated numerically with a model interatomic potential. The shock impulse is initiated by giving the first atom in the chain an initial velocity  $v_i$  toward its neighbor. We find that there exists a velocity  $v_c$  such that for  $v_i > v_c$  a soliton is produced by this impulse with a constant energy and supersonic velocity  $v_s$ . Studies of the motion of the atom during the passage of this soliton reveal a behavior similar to that expected from the collision of hard spheres. However, for  $v_i < v_c$ , the induced pulse propagates below the speed of sound and gradually disperses through the emission of phonons. In the supersonic regime,  $v_i > v_c$  we find that there exists a velocity  $v_m$  such that if  $v_i > v_m$ , then  $v_s < v_i$ , but if  $v_i < v_m$ , then  $v_s > v_i$ . Finally, in agreement with earlier studies using other model potentials, we find that if  $v_i$  is large enough, a soliton can be generated with sufficient energy to spall an atom from the end of the chain.

14. SUBJECT TERMS

molecular dynamics, chains, shock impulse, solitary waves

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT  
UNCLASSIFIED

18. SECURITY CLASSIFICATION OF THIS PAGE  
UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT  
UNCLASSIFIED

20. LIMITATION OF ABSTRACT

OFFICE OF NAVAL RESEARCH

CONTRACT #'s:

N00014-89-WX-24146

N00014-90-WX-24103

R&T CODE 412n006---02

TECHNICAL REPORT # 10

DYNAMICS OF SOLITARY WAVES INDUCED BY SHOCK IMPULSES IN  
A LINEAR CHAIN\*


Phuoc X. Tran, D.W. Brenner, and C.T. White

Code 6119  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Accepted for publication in the

*Int. J. Quantum Chem. S24, xxxx (1990).*

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-120	



\*Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

## Dynamics of Solitary Waves Induced by Shock Impulses in a Linear Atomic Chain\*

PHUOC X. TRAN, DONALD W. BRENNER, and C. T. WHITE

*Naval Research Laboratory, Washington, DC 20375-5000*

### Abstract

The propagation of nonlinear waves induced by a shock impulse in a linear atomic chain of finite but large length is investigated numerically with a model interatomic potential. The shock impulse is initiated by giving the first atom in the chain an initial velocity  $v_i$  toward its neighbor. We find that there exists a velocity  $v_c$  such that for  $v_i > v_c$  a soliton is produced by this impulse with a constant energy and supersonic velocity  $v_s$ . Studies of the motion of the atom during the passage of this soliton reveal a behavior similar to that expected from the collision of hard spheres. However, for  $v_i < v_c$  the induced pulse propagates below the speed of sound and gradually disperses through the emission of phonons. In the supersonic regime,  $v_i > v_c$ , we find that there exists a velocity  $v_m$  such that if  $v_i > v_m$ , then  $v_s < v_i$ , but if  $v_i < v_m$ , then  $v_s > v_i$ . Finally, in agreement with earlier studies on other model potentials, we find that if  $v_i$  is large enough, a soliton can be generated with sufficient energy to spall an atom from the end of the chain.

using

### Introduction

Since J. Scott Russell's observation (1834) of solitary waves on the surface of water, many other physical systems have been found to support the propagation of solitons. Initial analytic studies by Korteweg and de Vries [1] has resulted in an equation (KdV) which became a model for nonlinear wave propagation in fluid dynamics, plasma physics, optical communications, and other discrete nonlinear lattices. Zabusky and Kruskal [2] found a particular solution to the KdV equation in the form of stable pulses that can retain their initial shapes even after colliding into one another. These pulses were termed solitons as they behave like stable particles.

Ed. ok to delete  
one "to"?

Due to their remarkable properties, solitons have been proposed as a mechanism for energy transport in biological systems as well as in solids. Certain types of nonlinear discrete lattices such as the Toda model [3] are found to be integrable and have exact solitonic solutions. A few nonlinear models for interatomic potentials to some degree of approximation can be reduced to the KdV equation and hence admit solitons.

Studies of shock compression in solids and in energetic materials have been undertaken normally within the framework of continuum mechanics where macroscopic properties such as the temperature, pressure, and velocity profiles of atoms

\* This work was supported by ONR through NRL and the ONR Physics Division.

near the shock front are obtained. The continuum approach, however, does not include the dispersion in a discrete anharmonic lattice that may lead to an unsteady shock front, a result which is not allowed in hydrodynamic equations. Discrete models for shock wave propagation can provide physical details of the shock front and of individual atoms on a microscopic scale. Discrete models can also be used to study chemically sustained shock waves [4,5]. A large quantity of theoretical and experimental works on shock compression has been published, e.g., Ref. [8]. Some previous works have suggested that solitonic behavior [6,7] of a steady shock can result in the breaking of chemical bonds [8]. Moreover, the solitons initiated without shock waves in the Toda chain were shown numerically to cause fragmentation [9]. In this work, we study the propagation of solitons generated by varying the impact velocity of an instantaneous shock upon a linear chain. Other aspects associated with shock compression such as thermal excitation and chemical reaction in a heterogeneous lattice will be left to further studies.

In addition to existing models for interatomic potentials of Lennard-Jones, Morse, and Rydberg types, we can construct a simple potential that describes a repulsive interaction at short distance, attractive interaction within a finite range and vanishing at infinite, written as

$$V_{i,j} = C[1 - K(x_i - x_j)]e^{-K(x_i - x_j)} \quad (1)$$

where  $x_i$  is the position of the  $i$ th atom and  $K$  and  $C$  are positive constants proportional to the equilibrium separation and binding energy, respectively. The order of  $i$  and  $j$  is chosen such that  $x_i > x_j$ . If we consider only nearest neighbor interactions, Hamilton's equation for a one-dimensional lattice with free ends can be reduced to

$$\frac{d^2 Y_i}{dT^2} = (Y_{i+1} - Y_i - 2)e^{-(Y_{i+1} - Y_i)} - (Y_i - Y_{i-1} - 2)e^{-(Y_i - Y_{i-1})} \quad (2)$$

where  $Y$  and  $T$ , respectively, are the atomic position and time in dimensionless units. In the next section, we show a solution of the above system of ordinary differential equations obtained by using numerical algorithms. Analytical solutions in the continuum limit will also be presented.

### Results

The dimensions used in this calculation were scaled such that each unit of velocity, distance, and time correspond to 14.29 km/s, 1.6 Å, and  $1.21 \times 10^{-14}$  s, respectively. These values were based on an atomic mass of 14 amu, lattice spacing of 3.2 Å, and binding energy of 4 eV. The number of atoms in the chain is between 150 and 500.

The shock impulse is initiated by giving the first atom in the chain an initial velocity  $v_i$ , termed the impact velocity, toward its neighbors. The shock propagates further into the chain through interaction between neighboring atoms. At any moment, the position of the shock front can be defined as that of the last atom with the highest velocity, adjacent to the remaining atoms which are not yet disturbed.

Each atom is displaced in the same direction as the shock front passes. This translation is similar to the result of a collision of hard spheres. The graph of atomic velocity versus chain length shows a sharp pulse at the shock front [Fig. 1(a)]. Below a limit value  $v_c$  of the impact velocity, the amplitude of the pulse decreases (or disperses) as the front advances further into the lattice. This dispersion vanishes when the speed of the front reaches a value  $v_s$ , which requires an impact velocity  $v_c$ . Analytical calculation (given below) in the long wavelength limit shows that  $v_s$  is the speed of sound in the lattice and also that it is a minimum velocity a soliton must have.

For  $v_i > v_c$ , the amplitude of the pulse remains constant in time as it passes through the lattice. Furthermore, the pulse travels with a constant speed and energy until it gets reflected at the end of the chain. These results suggest solitonic behavior for the shock pulse (atomic velocity). The plot of this pulse in time [Fig. 1(b)] is similar to a hyperbolic secant function which is indeed a result obtained analytically in continuum limit. Supersonic solitons that emerge from a collision initiated by impacts upon both ends of the chain retain their initial shapes (Fig. 2). Also in this supersonic regime, there exists a value  $v_m > v_c$  such that the speed of soliton (or shock front) is found to be faster than the impact velocity if  $v_i < v_m$ , and slower if  $v_i > v_m$ . This characteristic is illustrated in Figure 3.

Depending on the value of the impact velocity, the energy of the soliton can attain a critical value to spall an atom from the end bond, which is the most weakly bound atom in the chain. This amount of energy is independent of the total number of atoms in the chain. Therefore, it is possible to have a stream of shock impulses acting on a lattice to generate a sequence of solitons with sufficient energy to break




Figure 1(a). Atomic velocities vs. atomic positions.  $v_i = 1.4$ ,  $N = 250$  atoms.




Figure 1(b). Atomic velocity vs. time. The curve with lower peak is obtained by numerical method for a discrete lattice. The one with higher peak is an analytic solution, Eq. (6) in continuum limit,  $v_i = 0.6$ ,  $N = 250$  atoms.

away the atoms one by one from a free end [9]. This may suggest an intrinsic mechanism for the breaking of chemical bonds without detonation by shock waves.

We now show an approximate analytic solution. In the long wavelength limit, the continuous version of Eq. (2) can be obtained by using




Figure 2(a). Collision of two solitons initiated by the same impact velocity on both ends of the chain.  $v_i = 1.5$ ,  $N = 250$  atoms.




Figure 2(b). Two solitons emerge intact after the collision.

$$R_{i\pm 1} = R_i \pm \frac{dR_i}{di} + \frac{1}{2} \frac{d^2 R_i}{di^2} \pm \frac{1}{3!} \frac{d^3 R_i}{di^3} + \frac{1}{4!} \frac{d^4 R_i}{di^4} \pm \dots \quad (3)$$

with the atomic displacement  $R_i$  being used instead of  $Y_i$ . First, Eq. (2) is expanded about the atomic equilibrium position up to second-order terms to obtain an ap-




Figure 3. Soliton (or shock front) velocity vs. impact velocity. The supersonic soliton is generated only in the region  $v_i > v_c$ .

proximate discrete equation. The resulting equation is then transformed into the continuum limit using Eq. (3) and keeping only leading terms up to fourth order in derivatives to obtain

$$e^2 \frac{d^2 R}{dT^2} = \frac{d^2 R}{di^2} - 2 \frac{d^2 R}{di^2} \frac{dR}{di} + \frac{1}{12} \frac{d^4 R}{di^4}; \quad (4)$$

Let  $z = K(x - vt)$  be a coordinate moving with the wave; then Eq. (4) becomes

$$\frac{\partial S}{\partial \tau} + \frac{1}{2} S \frac{\partial S}{\partial z} + \frac{1}{24} \frac{\partial^3 S}{\partial z^3} = 0 \quad (5)$$

where  $S = \partial R / \partial T$  being the atomic velocity,  $\omega = T/t$ ,  $\tau = 32\alpha^{-1}T$ , and  $\alpha = (eKv/\omega)^2 - 4$ . This is the Korteweg-de Vries equation, which can be shown to admit a particular solution in the form

$$S = S_\infty + \frac{\sigma}{4} \operatorname{sech}^2\left(\frac{\sqrt{\sigma}}{2} z\right) \quad (6)$$

with  $\sigma = \frac{3}{4}\alpha(Kv/\omega)$ . The stability of this solution requires  $\alpha$  to be positive or

$(2/e)\sqrt{C}/m$  with  $m$  being the atomic mass. This minimum is equal to the speed of sound traveling in the lattice, obtained analytically by solving Eq. (2) in the long wavelength limit. The soliton therefore exists only in a supersonic regime. Figure 1(b) shows a comparison between a continuous solution (6) and a discrete solution (numerical).

#### Discussion

It has been known that nonlinear lattices are not integrable except in very few models (e.g., the Toda exponential potential). Nonlinear interactions have a wide range of applications in many physical systems. Moreover, the balance between the nonlinear interaction in the lattice and the dispersion in the medium may give rise to interesting properties such as the recurrence phenomenon [10] (Fermi-Pasta-Ulam) and nondispersive solitary waves. Most other work [6-8] on shock waves and solitons in solids considered only the case of steady shocks. Although Karo has observed single soliton generated by impacting one end of the chain [11], the dynamical behavior of the lattice in response to shocks with different impact velocities has not been investigated in detail. By considering the shock impulse, we found that the shock front (or soliton) can propagate only with a supersonic speed, and that it is not always linearly proportional to the impact velocity. The energy delivered to a crystalline lattice by shock waves, as shown in this simplified model, is transported by lattice solitons. Chemical bonds within the lattice may break when the energy carried by the shock-induced solitons attain a certain critical threshold. Not only the bonds near the end of the lattice are broken up, but also those far away from the end as we have observed in the case of a sequence of impulses or a steady shock [12].

Edi powers on  
bracket needed  
here?, ok as is

run in  
(do on  
return)



We also found a minimum value of the impact velocity for the shock to generate a soliton that exists only in a supersonic regime. The speed of the soliton varies with the impact velocity but is time-invariant and independent of the length of the chain. The shock front (determined by the position of soliton) will continue to travel with the same speed when it first impacts upon the lattice only if the impact velocity has a particular value  $v_m$ . If the impact velocity falls below  $v_m$ , the front propagates faster and otherwise if above that value. The different behavior of the shock front when the impact velocity is below and above  $v_m$  arises most likely from the nonlinearity, an intrinsic property of the lattice. For an impact velocity greater than  $v_m$ , the force exerted by a shock is able to suppress the nonlinearity and therefore results in a slower speed of the shock front. Furthermore, it has been known that nonlinearity enhances the energy transport in a disordered lattice [13].

Our finding suggests a mechanism of energy transport in a lattice by a shock impulse, and determines how fast a shock front can propagate in a material given the impact velocity. Other aspects of shock compression in a lattice are subject to future works.

#### Acknowledgments

This work was supported by ONR through the ONR Physics Division, Contract #N00014-89-WX-24146 and through NRL.

#### Bibliography

- Av. Ref. 11. City 2
- [1] D. J. Korteweg and G. de Vries, *Phil. Mag.* **39**, 422 (1895).
  - [2] N. J. Zabusky and M. D. Kruskal, *Phys. Rev. Lett.* **15**, 240 (1965).
  - [3] M. Toda, *Prog. Theor. Phys. Suppl.* **59**, 1 (1976).
  - [4] M. L. Elert, D. M. Deaven, D. W. Brenner, and C. T. White, *Phys. Rev. B* **39**, 1453 (1989).
  - [5] D. W. Brenner, C. T. White, M. L. Elert, and F. E. Walker, *Int. J. Quant. Chem.* **S33**, 333 (1989).
  - [6] G. K. Straub and B. L. Holian, *Phys. Rev. B* **19**, 4049 (1979).
  - [7] J. H. Batteh and J. D. Powell, *Phys. Rev. B* **20**, 1398 (1979).
  - [8] A. Karo and J. R. Hardy, *J. Phys. Coll. (Paris)* **C4**, 235 (1987).
  - [9] M. Toda, *Prog. Theor. Phys. Suppl.* **59**, 148 (1976).
  - [10] E. Fermi, J. R. Pasta, and S. M. Ulam, *Collected Works of E. Fermi* (Chicago University Press, Chicago, 1965), Vol. II, p. 978.
  - [11] A. Karo, in *Fast Reactions in Energetic Systems*, C. Capellos and R. F. Walker, Eds., NATO Advanced Study Institutes Series C (Reidel, 1981), p. 611.
  - [12] P. X. Tran, unpublished.
  - [13] D. N. Payton, M. Rich, and W. M. Visscher, *Phys. Rev.* **160**, 706 (1967).

Received April 18, 1990

interim/revised June 1988

REPORTS DISTRIBUTION FOR ONR PHYSICS DIVISION  
UNCLASSIFIED CONTRACTS

Director Defense Advanced Research Projects Agency Attn: Technical Library, T10 1400 Wilson Blvd. Arlington Virginia 22209-2309	1 copy
Office of Naval Research Physics Division Office (Code 1112) 800 North Quincy Street Arlington, Virginia 22217-5000	2 copies
Office of Naval Research Director, Technology (Code 20) 800 North Quincy Street Arlington, Virginia 22217-5000	1 copy
Naval Research Laboratory Department of the Navy (Code 2625) Attn: Technical Library Washington, D.C. 20375-5000	1 copy
Office of the Director of Defense Research and Engineering Information Office Library Branch The Pentagon, Rm. 3E 1006 Washington, D.C. 20301	1 copy
U.S. Army Research Office Box 12211 Research Triangle Park North Carolina 27709-2211	2 copies
Defense Technical Information Center Cameron Station Alexandria, Virginia 22314	4 X copies
Director National Bureau of Standards Research Information Center Attn: Technical Library (Admin E-01) Gaithersburg, MD 20899	1 copy
Commander U.S. Army Belvoir Research, Development and Engineering Ctr. Attn: Technical Library (STRBE-BT) Fort Belvoir, Virginia 22060-5606	1 copy

interim/revised June 1986

ODDR&E Advisory Group on Electron Devices  
201 Varick Street, 11th Floor  
New York, New York 10014-4877

1 copy

Air Force Office of Scientific Research  
Department of the Air Force  
Bolling AFB, DC 22209

1 copy

Air Force Weapons Laboratory  
Technical Library  
Kirtland Air Force Base  
Albuquerque, NM 87117

1 copy

Lawrence Livermore Laboratory  
Attn: Dr. W. F. Krupke  
University of California  
P.O. 808  
Livermore, CA 94550

1 copy

Harry Diamond Laboratories  
Technical Library  
2800 Powder Mill Road  
Adelphi, MD 20783

1 copy

Naval Weapons Center  
Technical Library (Code 753)  
China Lake, CA 93555

1 copy

Naval Underwater Systems Center  
Technical Center  
New London, CT 06320

1 copy

Commandant of the Marine Corps  
Scientific Advisor (Code RD-1)  
Washington, DC 20380

1 copy

Naval Ordnance Station  
Technical Library  
Indian Head, MD 20640

1 copy

Naval Postgraduate School  
Technical Library (Code 0212)  
Monterey, CA 93940

1 copy

Naval Missile Center  
Technical Library (Code 5632.2)  
Point Mugo, CA 93010

1 copy

Naval Ordnance Station  
Technical Library  
Louisville, KY 40214

1 copy

interim/revised June 1988

Commanding Office Naval Ocean Research & Development Activity Technical Library NSTL Station, MS 39529	1 copy
Naval Oceans Systems Center Technical Library San Diego, CA 92152	1 copy
Naval Surface Weapons Center Technical Library Silver Spring, MD 20910	1 copy
Naval Ship Research and Development Center Central Library (Codes L42 and L43) Bethesda, MD 20084	1 copy
Naval Avionics Facility Technical Library Indianapolis, IN 46218	1 copy

*a*