

12

P.G. Saffman
Professor of Applied Mathematics

CALIFORNIA INSTITUTE OF TECHNOLOGY

APPLIED MATHEMATICS 217-50
FIRESTONE LABORATORY
PASADENA, CALIFORNIA 91125

pgs@ama-1.caltech.edu
FAX (818) 449-2677

AD-A231 205

January 2 1991

Dr Edwin P. Rood
Office of Naval Research
Code 113FF
800 N. Quincy Street
Arlington, VA 22217-5000

DTIC
ELECTE
JAN 14 1991
S D

Dear Dr Rood

Research in Nonlinear Water Waves
Navy Grant No. N00014-89-J-1164


Quarterly letter progress report Oct 1, 1990 - Dec 31, 1990

Work has continued during the quarter on the dynamical effects of a thin shear layer on the structure and stability of finite amplitude water waves. One problem of interest is the possibility of an 'explosive instability' discussed extensively in the Russian literature, which occurs when there is resonance between waves of opposite 'signature'. The signature can be thought of as the energy of the wave, and the explosive instability is an interaction between waves with positive and negative energy. (The concept is familiar in Plasma Physics.) The signature is easy to calculate when the undisturbed profile is of the 'stick' type, and this is the case for the Russian studies, but is not yet unambiguously defined when the profiles are smooth. We are studying this problem with the help of Dr R. MacKay from the University of Warwick, who is an expert on dynamical system theory, and hope to clarify the problem and be able to extend the Russian results to smooth profiles which may be more realistic.

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

We have also started work on the effect of a thin, vorticity containing shear layer in the water on the Kelvin-Helmholtz and Miles instabilities of the air flow over water. We are in the process of formulating the equations and are now considering efficient methods of solution and deciding upon the relevant parts of parameter space to search. Preliminary calculations with a 'stick' profile suggest that some rather interesting results may be obtained. In order to determine the effect of a thin vortical layer, it is necessary to study first the classical case of wind over water in the absence of vorticity in the water. Our approach differs from the classical Miles-Lighthill method based on asymptotic analysis of the critical layer. We solve the Rayleigh equation numerically, and we are finding growth rates that are five times greater than those obtained by Miles. In view of the continuing interest and belief in Miles's calculations, we are endeavoring to check this result as carefully as possible and compare also with numerical solutions of the Orr-Sommerfeld equation carried out by Valenzuela and others.

Yours sincerely



P.G. Saffman

- cc: ONR Pasadena
- cc: Director, Naval Research Laboratory
- cc: Defense Technical Information Center

Accession For	
NIS	CRAC
DIR	TAB
Undetermined	
Justification	
By AD-A 224065	
Distribution	
Availability	
Date	AVS
A-1	

