AD-A232 688

, Al	D-A2	232 68	8	•			(\mathcal{D})	
			PORT DOCUM	AENTATION	PAGE			
1a. REPORT SECURITY CLASSIFICATION Unclassified				16. RESTRICTIVE MARKINGS				
28. SECURITY CLASSIFICATION AUTHORITY				3. DISTRIBUTION/AVAILABILITY OF REPORT Distribution unlimited				
26. DECLASSIFICATION / DOWNGRACING SCHEDRIG 8 1991				Approved for public release				
4. PERFORMIN None	G ORGANIZAT	ION ROOM NUMB	ier(s)	s. monitoring organization report number(s) AEOSR-TR- 9-1 0-1-2-3				
6a. NAME OF Texas Tec Departmen	PERFORMING h Univers	ORGANIZATION ity tr. Engin.	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NP (Dr. R. Barker)				
6c. ADDRESS (City, State, and	d ZIP Code)		7b. ADDRESS (City. State and ZIP Code)				
Texas Tec	h Univers	ity		Building 410				
Departmen	t of Elect	trical Engine	eering	Bolling Air Force Base				
Ba. NAME OF	<u>TX /9409</u> FUNDING/SPO	3102 INSORING	86. OFFICE SYMBOL	Washington, DC 20332-0448				
ORGANIZATION AFOSR			(If applicable) NP	Grant AFOSR 86-0156				
Sc. ADDRESS (City, State, and	I ZIP Code)	1	10. SOURCE OF	UNDING NUMBE	RS		
Building	410	-		PROGRAM	PROJECT	TASK	WORK UNIT	
Bolling A	ir Force	Base		ELEMENT NO.	NO.	NO.	ACCESSION NO.	
Washingto	n <u>, DC</u> 203. ude Security C	32-6448		611 02 F	2301	<u>A8</u>	N. A.	
THEORETICAL AND NUMERICAL STUDY OF ANOMALOUS TURBULENT TRANSPORT IN PLASMAS (U)								
12. PERSONAL AUTHOR(S) Osamu Ishihara - Principal Investigator								
13a. TYPE OF	REPORT	13b. TIME	COVERED	14. DATE OF REPO	ORT (Year, Month	, Day) 15. PA		
Final Rep	OTT	FROM 6	<u>/15/86</u> 10 <u>12/14/</u> 9	1991, Febr	uary 5	4. f. activit	L	
contract	86-0156.	"Theoretical	and Numerical St	udv of Anom	alous Turbu	lent Tran:	sport in Plasmas	
for the p	eriod of	06/15/86 to 1	2/14/90.	, 			•	
17.	COSATI	CODES	18. SUBJECT TERMS (C	Continue on revers	e if necessary an	d identify by	block number)	
FIELD	GROUP	SUB-GROUP	Plasma Physics	, Plasma Tu	rbulence, P	lasma Ins	tabilities,	
			Turbulence, An	Langmuir Iurbulence, Drift Waves, Ion-Acoustic				
19. ABSTRACT	(Continue on	reverse if necessary	y and identify by block n	umber)				
This final technical report describes research work performed under AFOSR contract 86-0156 during the period of June 15, 1986 to December 14, 1990. Theoretical and computational analysis of particle transport in a plasma turbulence was carried out. Particle behavior was investigated by test particle simulations in different types of plasma turbulence, including								
in real space around mode rational surface was confirmed in drift wave turbulence. Nonresonant interaction of waves and particles were studied in ion acoustic turbulence. Resonance broaden-								
ing in velocity space was found as a result of non-Markovian process in Langmuir turbulence.								
ities in toroidal geometry were studied as eigen value problems. Undergraduate research								
assistantship program associated with this contract is also described.								
20. DISTRIBUT	ION / AVAILAB	ULTY OF ABSTRACT		21. ABSTRACT S	ECURITY CLASSIFI	CATION		
22 AUNCLASSIFIED AUNLIMITED DIS SAME AS RPT. DTIC USERS				Unclassified				
ober	FRESPONSIBLE t J. Bark	E INDIVIDUAL		226. TELEPHONE	22b. TELEPHONE (Include Area Code) 22c. OFFICE SYMBOL			
PAN 1	473. 84 MAD	R2 /	PR edition may be used un	202-767-4	904	AFOSR/	NP	
		- C - F			SECURITY	CLASSIFICAT	ION OF THIS PAGE	

.

• ,

FINAL TECHNICAL REPORT

.

2

on

Contract AFOSR 86-0156

THEORETICAL AND NUMERICAL STUDY OF ANOMALOUS TURBULENT TRANSPORT IN PLASMAS

June 15, 1986 to December 14, 1990

Osamu Ishihara

Principal Investigator Professor Department of Electrical Engineering Texas Tech University

91 8 06 128

on

Contract AFOSR 86-0156

THEORETICAL AND NUMERICAL STUDY OF ANOMALOUS TURBULENT TRANSPORT IN PLASMAS

for the period June 15, 1986 to December 14, 1990

Submitted to

The Air Force Office of Scientific Research

by

Dr. Osamu Ishihara Plasma Laboratory Department of Electrical Engineering Texas Tech University Lubbock, TX 79409

for

Dr. Robert J. Barker Directorate of Physics Air Force Office of Scientific Research Bolling Air Force Base Washington, D.C. 20332-6448

February 5, 1991

Principal Investigator:

Osamu Ishihara

Accesion For

NTIS CTASE

Unannounded Justification

Available

cope ?

DTIC T/ G

By Dist ibution (

Di.t

H-1

Phone 806-742-3463

TABLE OF CONTENTS

		Page	
I.	Summary	2	
II.	Research Accomplishments	3	
III.	Undergraduate Fellowship Program	5	
IV.	Listing of Conference Presentations	6	
V.	Listing of Publications	8	
VI.	Listing of Theses and Dissertations	9	
VII.	Staffs	10	
APPENDIX			

Conference Abstracts Supported by Grant AFOSR 86-0156 Publication Abstracts Supported by Grant AFOSR 86-0156

I. SUMMARY

This final technical report describes research work "Theoretical and Numerical Study of Anomalous Turbulent Transport in Plasmas" performed under AFOSR contract 86-0156 during the period of June 15, 1986 to December 14, 1990.

Theoretical and computational analysis of particle transport in a plasma turbulence was carried out. Particle behavior was investigated by test particle simulations in plasma turbulences, including drift wave turbulence, ion acoustic turbulence and Langmuir turbulence. A semiclassical approach to analyze plasma turbulence was taken to understand particle transport.

The report includes a brief description of research accomplishments under the support of this grant in Sec. II. Section III describes the undergraduate fellowship program associated with this grant. Section IV lists conference presentations, while Sec. V lists papers published in professional journals. Section VI lists the titles of master theses and PhD dissertations supported by this grant. The report concludes in Sec. VII by listing staffs involved in this contract.

II. RESEARCH ACCOMPLISHMENTS

Drift wave turbulence ---- Anomalous transport of electrons in the low frequency electrostatic turbulence (drift wave turbulence) has been studied extensively. Electrons were known to diffuse across a magnetic field by E x B drift, where E is a turbulent electric field fluctuation and B is a background magnetic field. We consider the transport of electrons in the presence of shear structure in the magnetic field. We have formulated the transport of electrons in a moderately strong turbulence based on the drift kinetic equation. Electrons are in resonance with waves at a mode rational surface, which is determined by the rational numbers associated with Fourier modes of drift waves. We have shown that the resonance region broadens as the turbulence becomes stronger and that the broadening develops with time. Our new formulation of resonance broadening explains our observation of particle transport in the test particle numerical experiment. The resulting transport coefficient can describe nonlinear aspect of wave-particle interaction in the drift wave turbulence. This principal investigator was invited to give a talk on this subject at the 1989 IEEE International Conference on Plasma Science at Buffalo, New York in May 1989. The full paper is published in Physics of Fluids (1990).

Langmuir Turbulence --- Time dependence of velocity-diffusion coefficient of plasma particles in Langmuir turbulence has been studied based on the generalized Langevin equation and the derived Fokker-Planck-type equation. The time dependent nature of the diffusion coefficient is caused by the retarded effect of the turbulent collisions, which is responsible for the non-Markovian stochastic process in a Gaussian (or non-Gaussian) system. The projection operator method has been used to derive the evolution equation known as the generalized Langevin equation for the particle velocity in an electrostatic plasma turbulence. Upon completion of hundreds of realizations in a test particle simulation, we obtained numerical results in agreement with analytical results. Our results were reported by a series of papers at IEEE International Conference on Plasma Science and at the Annual Meeting of Division of Plasma Physics of the American Physical Society.

Ion Acoustic Turbulence --- Nonresonant interaction of plasma particles and plasma waves has been studied in the presence of ion acoustic turbulence. We have confirmed by test particle computer experiments that particles away from the resonance layer could indeed diffuse as was predicted by our analytical work. The results were reported in the Annual Meeting of Division of Plasma Physics of the APS (1987). Relativistic particle acceleration --- Relativistic particle acceleration by plasma turbulence has been studied in a quasilinear regime. Relativistic charged particles can be accelerated by taking energy from plasma turbulence. We have formulated the problem classically and quantum mechanically. The theory describes the interaction between charged particles and plasmons. The self-field created by the particle itself is also considered. Our equation reduces to the Balescu-Lenard equation in the classical limit. This result was presented at the APS Topical Conference on Plasma Astrophysics at Santa Fe in September, 1988.

Turbulence in toroidal geometry --- Analytical study of turbulent fluctuations in toroidal geometry has been carried out in collaboration with Professor A. Hirose of University of Saskatchewan, Canada. In a collisionless plasma we have generalized the ideal MHD mode equation to describe drift-Alfven mode and examined numerically the stability problem (published in Nuclear Fusion (1987)). In a collisional plasma, the resistivity could be responsible for the instability. We have examined the resistive MHD balloning mode by solving the second order mode equation (published in the Canadian Journal of Physics (1988) and Nuclear Fusion (1989)). The mode equation is extended to incorporate the plasma inhomogeneity like ion-temperature-gradient. The numerical study of the ion-temperature-gradient instability is published in Nuclear Fusion (1987).

Semiclassical approach to plasma turbulence --- We have applied the method of quantum electrodynamics to the analysis of plasma turbulence. The wave-particle interaction is viewed as a scattering process between plasma particles and quasiparticles(plasmons, phonons). The particle transport in plasma turbulence is then closely related to the transition probability of absorption and emission processes. Successful application of the method resulted in clarification of the nonresonant wave-particle interaction in plasma turbulence (published in Physical Review A (1987)). On the other hand, such a viewpoint is applied to the interaction between photons and plasmons, and a novel scheme of photon acceleration (or frequency up-shifting of electromagnetic wave) is proposed (accepted for publication in Physical Review A).

III. UNDERGRADUATE RESEARCH FELLOWSHIP PROGRAM

The undergraduate research fellowship program was initiated by Dr. Robert Barker of AFOSR and our plasma laboratory was involved in the program during the period of the contract. The selected undergraduate students were involved in the related research. They participated in a weekly plasma seminar with graduate students and presented their assigned topics at the seminar. The names of undergraduate students and the titles of their final reports are listed here.

From May 16 to August 31, 1987 :

- 1. Theodore Christopher Grabowski
- "Anomalous Particle Transport in Drift Wave Turbulence in a Plasma" 2. Christopher James Koath
- "Study of Diffusion and Friction in an Electrostatic Turbulent Plasma" 3. David Edward Stahlke
 - "Ion Acoustic Plasma Turbulence and Data Processing"

From June 15, 1988 to May 31, 1989 :

1. Hok Tung Wong

"Application of MAGIC Code to the Study of Plasma Turbulence"

- 2. David Stahlke
 - "A Computational Study of Various Two-Stream Instabilities"
- 3. Mark Thompson
 - "A Numerical Experiment of Relativistic Particle Diffusion in a Plasma Turbulence"

IV. LISTING OF CONFERENCE PRESENTATIONS

Conference presentations of the research results obtained under the contract AFOSR 86-0156 are listed here.

- 1. Nonresonant Wave-Particle Interaction from the Quantum Mechanical Viewpoint, O. Ishihara, Bull. Am. Phys. Soc. 31, 1560 (1986).
- 2. Numerical Study of Turbulence, E. Robinson and O. Ishihara, Bull. Am. Phys. Soc. 31, 1497 (1986).
- 3. Real Space Diffusion in Drift Wave Turbulence, E. Robinson and O. Ishihara, IEEE Int. Conf. Plasma Sci., paper 2X5 (Arlington, VA 1987).
- 4. Analytical and Numerical Study of Anomalous Friction in a Plasma Turbulence, W. Ho and O. Ishihara, IEEE Int. Conf. Plasma Sci., paper 5T3 (Arlington, VA 1987).
- 5. Resistive Modes in Tokamaks, A. Hirose, T.L. Kroeker, and O. Ishihara, Bull. Am. Phys. Soc. 32, 1773 (1987).
- Study of Velocity Diffusion of Test Particles in Ion Acoustic Turbulence, K.M. Yuen, O. Ishihara, and A. Hirose, Bull. Am. Phys. Soc. 32, 1859 (1987).
- 7. Quasilinear Effects on Turbulent Bremsstrahlung, O. Ishihara and A. Hirose, Bull. Am. Phys. Soc. 32, 1859 (1987).
- 8. Stochastic Heating of Plasma Particles in Ion Acoustic Turbulence, K.M. Yuen and O. Ishihara, Spring Meeting of the Texas Section of the American Physical Society (Austin, TX March, 1988).
- 9. Plasma Particle Diffusion in Drift Wave Turbulence, C. Grabowski and O. Ishihara, Spring Meeting of the Texas Section of the Am. Phys. Soc. (Austin, TX March, 1988).
- Stochastic Diffusion in the Absence of Resonances, O. Ishihara, W. Ho, K.M. Yuen, and A. Hirose, IEEE Int. Conf. Plasma Sci. (Seattle, WA 1988).
- 11. Resistive MHD Ballooning Mode in Tokamaks, A. Hirose and O. Ishihara, IEEE Int. Conf. Plasma Sci. (Seattle, WA 1988).
- Relativistic Quasilinear Particle Acceleration in Plasma Turbulence, O. Ishihara, APS Topical Conf. on Plasma Astrophysics (Santa Fe, NM 1988). Bull. Am. Phys. Soc. 34, 1287 (1989).
- 13. Saturation of Nonlinear Drift Mode Induced by Toroidicity, O. Ishihara and A. Hirose, Bull. Am. Phys. Soc. 33, 1941 (1988).
- Time-Dependent Diffusion Coefficient for Non-Markovian Process in Plasma Turbulence, H. Xia and O. Ishihara, Bull. Am. Thys. Soc. 33, 2020 (1988).

- 15. Study of Stochastic Heating Mechanism of Plasma Particles in the Presence of Ion Acoustic Turbulence, K.M. Yuen and O. Ishihara, Fall Meeting of the Texas Section of the American Physical Society (Lubbock, TX 1988). Bull. Am. Phys. Soc. 34, 1521(1989).
- Velocity Diffusion Coefficient as a Function of Time, H. Xia and O. Ishihara, Fall Meeting of the Texas Section of the Am. Phys. Soc. (Lubbock, TX 1988). Bull. Am. Phys. Soc. 34, 1521 (1989).
- 17. Particle Diffusion in Turbulent Fields: Transition from Quasilinear to Nonlinear Stage, O. Ishihara, IEEE Int. Conf. Plasma Sci. (Buffalo, NY 1989). (Invited Talk).
- 18. Generalized Langevin Equation for Plasma Turbulence, H. Xia and O. Ishihara, IEEE Int. Conf. Plasma Sci. (Buffalo, NY 1989).
- 19. Study of Properties of the Stochastic Processes in Plasma Turbulence, H. Xia and O. Ishihara, Bull. Am. Phys. Soc. 34, 1926 (1989).
- 20. Photon Acceleration: Three Wave Interaction, O. Ishihara, IEEE Int. Conf. Plasma Sci. (Oakland, CA 1990).
- Effect of Turbulent Collisions on Diffusion in Stationary Plasma Turbulence, H. Xia and O. Ishihara, IEEE Int Conf.Pasma Sci. (Oakland, CA 1990).
- 22. Photon Acceleration by the Plasmon-Photon-Photon Interaction, O. Ishihara, Bull. Am. Phys. Soc. 35, 2125 (1990).
- Analysis of the Non-Markovian Behavior in Velocity Diffusion Process, H. Xia and O. Ishihara, Bull. Am. Phys. Soc. 35, 1956 (1990).
- 24. Photon Acceleration by Plasma Turbulence, O. Ishihara, IEEE Int. Conf. Plasma Sci. (Williamsburg, VA 1991).
- 25. Characteristics of Distribution Function in a Time-Dependent Velocity Diffusion Process, H. Xia and O. Ishihara, IEEE Int. Conf. Plasma Sci. (Williamsburg, VA 1991).

V. LISTING OF PUBLICATIONS

Publications of the research results obtained under the contract AFOSR 86-0156 are listed here.

- 1. Nonresonant Wave-Particle Interaction in Semiclassical Quasilinear Theory, O. Ishihara, Phys. Rev. A35, 1219-1225 (1987).
- 2. Drift Alfven Eigenmode in Tokamaks, A. Hirose and O. Ishihara, Nuclear Fusion 27, 1231-1238 (1987).
- 3. Search for Ion Temperature Gradient Driven, Electrostatic Hydrodynamic Instability in Tokamaks, A. Hirose and O. Ishihara, Nuclear Fusion 27, 1439-1451 (1987).
- 4. Resistive Ballooning Mode in Tokamaks, A. Hirose, T.L. Kroeker, and O. Ishihara, Can. J. Phys. 66, 1069-1075 (1988).
- 5. Resistive MHD Ballooning Mode in Tokamaks, A. Hirose and O. Ishihara, Nuclear Fusion 29, 795-803 (1989)
- 6. Resonance Broadening in Drift Wave Turbulence, O. Ishihara, C. Grabowski, and A. Hirose, Phys. Fluids B2, 270-279 (1990).
- 7. Photon-Plasmon-Photon Interaction, O. Ishihara, Phys. Rev. A (in press).

VI. LISTING OF THESES AND DISSERTATIONS

Master theses and doctoral dissertations supported by the contract AFOSR 86-0156 are listed here.

Maser Theses

- 1. Edward Kyle Robinson, "Real Space Diffusion in Drift Wave Turbulence" (1987).
- 2. Kin-Ming Yuen, "Velocity Space Diffusion in Ion Acoustic Turbulence" (1989).

PhD Dissertations

- 1. Wai H. Ho, "Wave-Particle Interaction in Plasma Turbulence" (final version is in preparation).
- 2. Huajuan Xia, "The Non-Markovian Process in Velocity Diffusion in the Stationary Plasma Turbulence" (in preparation).

VI. STAFFS

Faculty Investigator:

Professor Osamu Ishihara of Department of Electrical Engineering and Department of Physics at Texas Tech University is a principal investigator.

Graduate Research Assistants:

During the period of the contract, two doctoral students and two master students were supported.

Undergraduate Research Assistants:

Five undergraduate students were supported under the program of undergraduate research fellowship.

Accountants:

Financial record was kept by Ms. Sandra Branch of the Department of Electrical Engineering. She was partially supported by this contract.

APPENDIX

.

Program of the Twenty-Eighth Annual Meeting of the Division of Plasma Physics Baltimore, Maryland; 3–7 November 1986

RI 9C

Monresonant Vare-Particle Interaction from the Quantum Mechanical Viewpoint,* Osemu Ishihara, Texas Tech. University -- Honresonant nature of wave-particle interaction is clarified from the viewpoint of quantum mechanics. The interaction of particles and quasiparticles (plasmons, phonons, etc.) is described by a scattering process which is charaterized by a transition probability. A proper treatment of the time-dependent interaction Hamiltonian reveals that both the resonant and nearesonant mature of particle-quasiparticle interaction contributes to transition probability. The resenant transition probability [1], known as Farmi's golden rule, is now supplemented by a nonresonant part, which guarantees the conservation of emergy and momentum in the system as is predicted by the classical quasilinear theory.

* Work supported by AFOSE under grant AFOSE-86-0156.
[1] E. G. Harris, in Advances in Plasma Physics, ed. by A. Simon and W. B. Thompson, Vol. 3, p.157 (Wiley, New York, 1969).

4T 26

<u>Hamerical Study of Test Particle Transport in Drift</u> <u>Namerical Study of Test Particle Transport in Drift</u> <u>Namerical Study</u> - In drift wave turbulence where the <u>magnetic field is expressed as $B = B_0(0, \pi/L_g, 1)$, L_g shear length, a teal space diffusion coefficient has been formulated¹ and is found to have similiar characteristics as the time dependent diffusion coefficient derived for velocity space in Langmuir turbulence.² A numerical experiment reveals the particle diffusion in real space in the x direction. Test particles are followed in their trajectories in the prescribed random electrostatic fields, $\Phi(\mathbf{X}, t) = \sum_{min} \Phi_{mentical}(\mathbf{x}_N)$ + $\mathbf{k}_R = -\omega_{min}(t)$, where Φ_{min} is the electric potential of the (m, 0) mode. A preliminary result of the observed diffusion coefficient demonstrates the time dependent structure of the diffusion coefficient as predicted by the analytical study.</u>

* Work supported by AFOSR under grant AFOSR-86-0156

A. Salat, Z. Maturforsch. Teil A <u>38</u>, 1189 (1983).
 O. Ishihara and A. Mirose, Phys. Fluids <u>28</u>, 2159 (1983).

CONFERENCE RECORD - ABSTRACTS

1987 IEEE INTERNATIONAL CONFERENCE ON PLASMA SCIENCE

June 1-3, 1987

Arlington, Virginia

2X5

Real Space Diffusion In Drift Wave Turbulence*

E. Robinson and O. tshihara

Department of Electrical Engineering Texas Tech University, Lubbock, Texas 79409

In drift wave turbulence characterized by a sheared magnetic field $\vec{B} = B_0(\vec{e}_z + x/L_z\vec{e}_y)$, where L_z is the shear

length, and random electrostatic fields, $\Phi(x,t) = \sum_{mn} \Phi_{mn} x \Theta^{i}(k_m y + k_n z - \omega_{mn})$, where Φ_{mn} is the electric potential of the (m,n) mode, a numerical experiment has been carried out to examine the particle diffusion process in real space. Two cases are examined, one with an infinite number of n-modes and the other with a single n-mode. 100 m-modes are used for both cases. For infinite n-modes, diffusion coefficients found

are in agreement with the value $D = \pi \sum_{m} \langle |m \Phi_m|^2 \rangle$ as was predicted¹ for a low turbulence level, while for a single n-mode we observe that diffusion greatly depends on the choice of m and n.

An analytical expression for the space diffusion in the x-direction is given by

$$D(X,t) = \frac{1}{2} \frac{d}{dT} \sum_{mn} \left< \left| m \tilde{\Phi}_{mn} \right|^2 \right> \int_{0}^{1} \int_{0}$$

where X is the initial position of a test particle, exhibits analogous characteristics to the quasilinear velocity space diffusion coefficient.² For low turbulence levels, Eq. (1) shows quasilinear-like behavior,

$$D(X) = \pi \sum_{mr} \left\langle \left| m \Phi_{mn} \right|^2 \right\rangle \delta(mX + n).$$
 (2)

at the resonant locations, X = -(n/m). From further examination of the single n-mode at a rather high turbulence level, we have observed deviation of the diffusion coefficient from that of Eq. (2), i.e. high turbulence levels cause diffusion in the space surrounding the resonant locations as is predicted by Eq. (1). A comparative study between the real space diffusion in drift wave turbulence and the velocity space diffusion in Langmuir turbulence, $\omega \propto \omega_{pe}^2$ at a rather strong turbulence level demonstrates the time dependent nature of the diffusion coefficient as was described by Eq. (1).

* Work supported by AFOSR under grant AFOSR-86-0156.

- 1. A. Salat, Z. Naturiorsch. Tell a 38, 1189 (1983).
- 2. O. Ishihara and A. Hirose, Phys. Fluids 28, 2159 (1985).

Analytical and Numerical Study of Anomolous Friction in a Plasma Turbulance.•

Wai Ho and Osamu Ishihara, Texas Tech University, Labbock, TX 79409-4439.---The anomolous dragging effect on plasma particles caused by the electrostatic plasma turbulence in the absence of magnetic field is studied numerically and analytically. It has been known that the frictional effect together with the diffusion effect appears in the presence of succhastic electric fields. In the surbulent electric fields, the distortion of particle trajectories can be given by

$$\Delta \hat{x}(t) = \int_{0}^{t} \Delta \hat{v}(t) dt', \text{ where } \Delta \hat{v}(t) = \frac{q}{m} \int_{0}^{t} \hat{E}[\hat{x}(t), t] dt'.$$

We thus define the friction coefficient as $F(v,t) = d<\Delta V(t)/dt$. It is straightforward to show that the friction coefficient is given by

$$\vec{R}(\vec{v},t) = \left(\frac{q}{m}\right)^2 \frac{1}{t} \int_0^t d\tau \int_0^t ds \frac{\partial}{\partial \vec{v}} \cdot s^{-1} \left(\vec{v},s\right) \exp(i\vec{k} \cdot \Delta \vec{x}(t-s))$$
(1)

where $\hat{S}(\vec{vs},s) = \sum_{k} \int \frac{d\omega}{2\pi} \hat{s}(\vec{k},\omega) \exp[i(\omega - \vec{k} \cdot \vec{v})s]$

and
$$\int (\hat{\mathbf{k}}, \omega) = \sum_{\omega} \int \frac{d\omega}{2\pi} \langle \hat{\mathbf{k}}(\hat{\mathbf{k}}, \omega) \hat{\mathbf{k}}(\hat{\mathbf{k}}, \omega) \rangle$$
.

Alternatively, we can write F(v,t) as

$$\vec{F}(\vec{v},t) = \left(\frac{q}{m}\right)^{2} \frac{1}{t} \frac{\partial}{\partial \vec{v}} \circ \int_{0}^{t} \frac{d}{dt} \int_{t-\tau^{1}}^{t} \frac{d}{ds} \int_{0}^{\infty} \frac{d}{ds} \left(\frac{d}{ds}\right)^{2} \left(\frac{d}$$

A numerical experiment has been carried out, in which one dimensional random electric fields are modeled for Langmuir turbulence and are described by $\frac{1}{2}$

$$\langle E(x,t)E(0,0)\rangle = -\sigma^2 \left(\frac{m\omega_p^2}{qk_0^2}\right) \frac{\partial^2}{\partial x^2} \left[\exp(-\frac{1}{25} k_0^2 x^2) \cos(k_0 x - \omega_p t) \right]$$

where up is the plasma frequency, k_0 is the center wavenumber and σ is the normalized rms value of the turbulent potential. In the limiting case of weak urbulence ($\sigma \ll 0.01$), we have observed a well-defined friction coefficient as given by the limit $\Delta \vec{x} \rightarrow 0$ of Eq. (1);

$$\mathbf{F}(\vec{v}) = \frac{1}{2} \begin{pmatrix} \mathbf{q} \\ \mathbf{m} \end{pmatrix} \sum_{\mathbf{k}} d\omega \left[\frac{\partial}{\partial \omega} \vec{k} \cdot \vec{s} \right] \left[\vec{k} \cdot \omega \right] \mathbf{s} \left(\omega \cdot \vec{k} \cdot \vec{v} \right) \quad (3)$$

In the moderate turbulence level ($\tau \sim 0.01$), the friction coefficient is observed to deviate from the value given by Eq. (3), and is observed to show some broadening effect as represented by $<\Delta T >$ term in Eq. (2).

The formulation of the friction coefficient is also studied by a renormalized method with Green's function based on the Vlasov equation.² With the help of diagrammatic technique, we formulate the turbulent friction by taking secular terms in the Green's function which agrees with Eq. (1) in some limiting situation.

* Work supported by AFOSR under grant AFOSR-86-0156.

1. O. Ishihara and A. Hirose, Phys. Fluids 23, 2159(1985). 2. M. Kono and Y. H. Ichikawa, Prog. Theor. Phys. 49,754(1973).

5T3

. . +

Program of the Twenty-Ninth Annual Meeting of the Division of Plasma Physics San Diego, California; 2–6 November 1987

3R 18 <u>Resistive Modes in Tokenska</u>⁴ A. HIROSE, T.L. KROEKER, <u>Univ. of Sask.</u>, O. ISHIRARA, <u>Texas Tech</u> <u>Univ.</u> —The resistive g and rippling modes in tokensk geometry have been reszenized in terms of the generalized moments' free from any ordering energy frequencies, ω , ω_n , and $\omega_D(n) = 2\xi_1\omega_n$ (cost + of simi) (the megnetic drift frequency in the ballousing space). Such generalization is necessary for resistive modes because of strong radial localization about a rational surface and corresponding large mode width in u-space, $|\omega_D(n)| >>$ $|\omega| \ge \omega_n$. For typical tokensk edge parameters, a memerical search for unstable, bounded eigenfunctions of resistive modes has failed. Only the stable (stabilized by shear and toroidicity) drift mode (predicted analyti-

*Sponsored by MSERC (Canada) and USAF-OSR

cally) has been found.

¹A. Hirose, Nucl. Fusion (in press). Also, Coum. Plasme Phys. Controlled Fusion <u>10</u>, 229 (1987)

6T 11

Study of Velocity Diffusion of Test Particles in Ion Acoustic Turbulence.* K.M. Yuen and O. Ishihara, Texas Tech Univ., and A. Hirose, Univ. of Saskatchewan .- in order to study velocity diffusion of ions in ion acoustic turbulence, we set the test particles in prescribed turbulent electric fields. The numerical experiments are carried out with particles placed randomly in the system, or alternatively we followed the test particles with the same initial position for different realization of electric fields. The ion acoustic turbulence is modeled by the dispersion relation which is characterized by either nondispersive or dispersive nature. The results of the numerical experiments indicate that in both nondispersive and dispersive cases, ions which do not satisfy the resonant condition do diffuse in velocity space. The analytical study also shows a possibility of diffusion of nonresonant particles if the turbulent spectrum has a finite correlation time. The numerical results show fair agreement with theoretical predictions provided that the initial velocities are set within critical values, which are located well above and well below the phase velocity of the waves. It has also been observed that there is no stochastic heating of particles if their initial velocities are set outside the critical range

*Supported by AFOSR under Grant AFOSR-86-0156A.

1. A. Hirose & O. Ishihara, Bull. Am. Phys. Soc. 29, 1238 (1984). 6T 12

.

Quasilinear Effects on Turbulent Bremsstrahlung.⁴ O. Ishihara, <u>Texas Tech Univ.</u>, and A. Hirose, <u>Univ. of Sas-katchewan</u>—The upconversion of low frequency ion acoustic turbulence into high frequency Langmuir waves is re-examined in the context of weak turbulence theory. Con-

trary to the earlier observations^{1,2} that there exists turbulent bremsstrahlung of Langmuir waves with a background of ion-acoustic turbulence, it is shown that quasifinear effects on the electron distribution function cause stabilization of Langmuir waves. The quasilinear effects are found tion of Langmuir waves. The quasilinear effects are sound to cancel the destabilizing terms originated from the direct turbulent corrections to the linear response function. To-gether with the earlier findings^{2,3} that polarization effects do not cause destabilization of Langmuir waves, it is now concluded that there should be no upconversion of ion acoustic turbulence into Langmuir turbulence, which agrees with the conclusion derived by the symmetry argument.

*Supported by AFOSR under grant AFOSR-86-0156A (O.L.) and NSERC, Canada (A.H.).

and reserve, Canada (A.F.).
V.N. Tsytovich, L. Stenflo & H. Wilhelmsson, Physica Scripta 11, 251 (1975).
A. Hirose, Com. Plasma Phys. Contr. Fus. 5, 117 (1984).
D.F. Duflois & D. Pesme, Phys. Fluids 27, 218 (1984).
J. Kuijpers & D.B. Melrose, Astrophys. J. 294, 28 (1985).

PROGRAM OF THE JOINT SPRING MEETING of the Texas Section of the American Physical Society Texas Section of the American Association of Physics Teachers Society of Physics Students, Zone 10 The University of Texas Austin, Texas March 4 & 5, 1988

9:12

AD2

Stochastic Heating of Plasma Particles in Ion Acoustic Jurbulence.* K.M. YUEN and O. ISHIHARA, <u>Texas Tech</u> Univ. Stochastic process of plasma ions placed in the background ion acoustic turbulence is studied analytically and numerically. The dispersion relation of the ion acoustic wave is given by $\omega - kC_g$, where C_g is the ion acoustic velocity. The random nature of the turbulence is characterized by the correlation function of electric fields. We model the turbulence by discrete Fourier modes and the envelope of its correlation function forms a Gaussian spectrum. By controlling the width of the spectrum, the model is applicable to a single mode as a limit and to the wide-band turbulence. It is shown that when the spectrum width becomes wider the heating rate of plasma particles becomes larger. Stochastic heating is found to take place for ions in resonance and also not in resonance with ion acoustic waves.

*Supported by AFOSR under grant AFOSR-86-0156A.

11:12

AD12

Plasma Particle Diffusion in Drift Wave Turbulence. C. GRABOUSKI and O. ISHIHARA, Texas Tech U.* Notions of plasma particles have been studied numerically and analytically when the particles are subjected to both a background magnetic field and electric fields created by drift wave curbulence. For these studies, test particles are placed in a field model in slab geometry. Both the random electric fields and the background magnetic field, with increasing shear in the x direction, are set to be confined in the y-z plane. Humerical and analytical results indicate that particle motions increase slong the x axis as the magnitude of the electric fields increases and as the density of the mode rational surfaces increases along the x axis, where the mode ratiomal surface is defined by x = -n/n with n and m being mode numbers of the electric fields.

*Supported by AFOSR under grant AFOSR-86-0156A.

CONFERENCE RECORD - ABSTRACTS

1988 IEEE INTERNATIONAL CONFERENCE ON PLASMA SCIENCE

June 6-8, 1988

Seattle, Washington

4C6

Stochastic Diffusion in the Absence of Resonances*

O. Ishihara, W. Ho and K.M. Yuen Department of Electrical Engineering Texas Tech University, Lubbock, Texas and A. Hirose Department of Physics University of Saskatchewan, Saskatoon, Canada

The nonresonant nature of charged particle motion in the electrostatic plasma turbulence is studied numerically and analytically by artificially suppressing the resonant conditions of wave-particle interaction. Nonresonant structure in the phase space is realized in the test particle numerical experiments by two different methods. One approach is to make use of discreteness of the spectrum of plasma turbulence. Test particles placed in velocity space may not reach any of two adjacent modes in the extremely weak turbulence. This situation is modeled for electrons placed in the Langmuir turbulence.¹ Another approach is to place test particles completely out of the resonant region in velocity space. This situation is modeled for jons placed in the ion acoustic turbulence.² Nu-

merical experiments are carried out for test particles placed in prescribed random electric fields, whose second order correlation functions are characterized by sinusoidal structure with the exponential damping in real space, described by the summation of Fourier modes as

$E(x,t) = \sum_{j} f(k_j) \cos[w(k_j)t - k_j x + \beta_j],$

where $f(k_j)$ is a shaping factor with Gaussian form. β_j is a random phase. and $\omega(k_j) = 1$ for Langmuir waves and $\omega(k_j) = k_j/(1+k_j^2)^{1/2}$ for ion acoustic waves. Test particles, ranging from 20 to 2000 in number, are followed in their trajectories in the discrete fluctuation spectrum with mode numbers 20 to 400. Diffusion coefficients are measured in velocity space by taking mean square variation as a function of time and compared with the analytic predictions evaluated by

$$D(v) = \begin{pmatrix} g \\ g \end{pmatrix}^2 \int_0^{\infty} dt \int_{-\infty}^{\infty} dx \ \delta(x \cdot vt) \langle E(x', t')E(x' + x, t' + t) \rangle,$$

where e/m is a charge/mass ratio and set to be 1. Stochastic diffusion observed numerically both in Langmuir turbulence and in ion acoustic turbulence is in good agreement with the theory in the scale of Kolmogorov time defined by $\tau_{\rm K} \sim (k^2 \ D)^{-1/3}$. In the Langmuir turbulence, diffusion is observed even well below the critical turbulence level given by the criterion for the resonance overlapping. Our observation suggests the possible stochastic acceleration of charged particles in the nonresonant nature in the plasma turbulence.

- 1. O. Ishihara and A. Hirose, Phys. Fluids 28. 2159 (1985).
- 2. A. Hirose and O. Ishihara, Bull. Am. Phys. Soc. 29. 1238 (1984).

Work supported by AFOSR under Grant AFOSR-86-0156A (0.1., W.H., K.Y.) and by NSERC of Canada (A.H.). RESISTIVE HHD BALLOONING HODE IN TOKANAKS*

A. Hirose Dept. of Physics. Univ. of Saskatchewan Saskatoon, Saskatchewan, S7N 0MO, Canada

O. Ishihara Dept. of Elect. Eng., Texas Tech Univ. Lubbock, TX 79409

A mode equation for the resistive NKD ballooning mode in low 8 tokamak magnetic geometry has been derived and analyzed both analytically and numerically. Any resistive modes in a high temperature tokamak require strong radial localization about a mode rational surface in order to satisfy $v_g \gg k_g(r)v_{T_0}$. Here, v_g is the electron-ion collision frequency and $k_g(r) = k_g r/L_g$ is the parallel gradient with r being the radial distance from the rational surface, and L_g the shoar length. Strong radial localization makes the magnetic drift frequency operator

$$u_{D} = 2c_{n}u_{n}\left(\cos\theta - \frac{1\sin\theta}{k_{0}}\frac{\partial}{\partial r}\right)$$

secular. and the conventional ordering

totally breaks down. The formulation in the present study is free from any ordering among the frequencies

w, w_g and w_D and thus offers a generalization of the conventional mode equation for arbitrarily strong radial localization. The mode equation, based on electrostatic ion response and fully electromagnetic electron response, is given in Fourier space by

$$\left[\frac{d}{d\kappa} \frac{1+\kappa^2}{A(\kappa)} \frac{d}{d\kappa} + \left(\frac{L_{\theta} k_{De}}{ck_{\theta}}\right)^2 \frac{G(\kappa)}{B(\kappa)}\right] \phi(\kappa) = 0$$

where

۰.

$$A = u - u_{ee} + u_{De}(\kappa) + i \left(\frac{ck_0}{u_{pe}}\right)^2 v_e^{(1+\kappa^2)}$$

$$B = u - u_{ee} + u_{De}(\kappa)(1+\tau^{-1}) + \tau(u + u_{e1})(1 - \Gamma_0)$$

$$G = [u_{ee}^{-u}u_{De}(\kappa)](1+\tau^{-1})u_{De}(\kappa)+\tau[u-u_{De}(\kappa)](u+u_{e1})(1-\Gamma_0)$$

$$\Gamma_0 = e^{-b}I_0(b) , b = (k_0\rho_1)^2(1+\kappa^2) , \tau = T_0/T_1$$

No unstable modes have been predicted analytically and found numerically for the mode equation. The only bounded solution analytically predicted is the stable ion diamagnetic mode. The damping factor is proportional to the magnetic shear and ϵ_n , the toroidicity. This has been fully confirmed by numerical integration with parameters relevant to present-day tokamaks.

*Sponsored by NSERC, Canada (A.H.) and AFOSR (0.1.)

4C7

Minutes of the American Physical Society Topical Conference on Plasma Astrophysics Santa Fe, New Mexico; 19–23 September 1988

P112 Relativistic Ouasilinear Particle Acceleration in Plasma Turbulence.* O. ISHTHARA, Texas Tech U.--Relativistic quasilinear kinetic theory is developed to describe particle acceleration in the presence of plasma turbulence. Scattering between charged particles and plasmons is considered. It is assumed that each time a particle makes a transition the energy it gains (or loses) comes from (or goes into) the energy of turbulent fluctuations. The theory includes nonresonant interactions between charged particles and plasmons since plasmons are known to carry not only electric field energy but also oscillating energy of particles.1 The formulation takes into account the self fields created by the particle itself, which can be incorporated as the effect of renormalization of the particle mass.² The derived equation reduces to the Balescu-Lenard equation in the classical limit. A possible mechanism of cosmic ray acceleration is discussed based on this theory.

*Supported by US AFOSR Grant AFOSR-86-0156. 10. Ishihara, Phys. Rev. A. <u>35</u>, 1219 (1987). ²V.N. Tsytovich, Sov. Phys. JETP <u>52</u>, 3 (1980).

Program of the Thirtieth Annual Meeting of the Division of Plasma Physics Hollywood, Florida; 31 October-4 November 1988

Drift 11 23 Saturation 10 Nonlinear Induced by Toroidicity. * O. ISHIHARA, Texas Inch. University, and A. HIROSE, University of Sakatchevan-Tokamaks are known to be unstable with respect to a drift mode because of their toroidicity. The turbulence level at their toroidicity. The turbulence level at saturation of the drift mode is studied with the assumption of cold ions and the electron the assumption of coid ions and the electron temperature gradient as a principal nonlinear-ity. A positive nonlinear frequency shift is given by $\Delta \Omega = (q_0^2/12k^2)|\phi|^2$, where q_0 is the electron temperature gradient relative to a density gradient, $k = k_{d/2}(k_d)$ is a poloi-dal asimuthal wavenumber, ρ_0 is an ion Larmor module with an electron temperature), and ϕ is radius with an electron temperature), and ϕ is the peak boundary value of the potential. The nonlinearity sensitively modifies the radial profile of the drift mode, which in turn induces nonlinear wave damping. The resulting turbulence level at saturation falls in be-tween the strong turbulence limit due to ion mode coupling and the weak turbulence limit due to electron trapping. *Supported by NSERC, Canada (A.H.) and AFOSR under grant 86-0156 (0.I.).

Time-Dependent Diffusion Coefficient 6S 2 for Non-Markovian Process in Plasma Turbulence.* H. XIA and O. ISHIHARA, Texas Tech University -- Recent study of resonance broadening theory showed the time dependent nature of a diffusion coefficient of plasma particles in velocity space, 1, 2 where a particle orbit modification is properly treated. Such a time dependence of diffusion coefficient is studied based on the generalized Langevin equation, where a frictional force exerted by the effective collisions due to plasma turbulence is included with its retarded effect. This retarded effect is responsible for the non-Markovian process in a Gaussian system. It is shown that a derived Pokker-Planck-type equation has a diffusion coefficient which depends on time and reduces to the conventional quasilinear diffusion coefficient in a short time limit of much less than the effective collision time.

*Supported by APOSR under grant \$6-0156.

- 1. O. Ishihara and A. Hirose, Phys. Fluids 28, 2159 (1985).
- 2. A. Salat, Phys. Fluids 31, 1499 (1988).

Minutes of the Seventh Annual Fall Meeting of the Texas Section Lubbock. Texas: 4-5 November 1988

CF 11 Study of Stochastic Heating Mechanism of Plasma Particles in the Presence of Ion Acoustic Turbulence⁶. K.M. YUEN and O. ISHIHARA, Texas Tech University. The velocity diffusion of ions in ion acoustic turbulence is studied numerically and analytically. Turbulence is modeled by random electric fields with discrete Fourier modes and the correlation function of the fields is assumed to be Gaussian. Test ions are placed in a background of ion acoustic turbulence and the statistics of particle motions are studied. We have observed that stochastic heating can take place for ions both in resonance and not in resonance with ion acoustic waves. Resonant particles keep absorbing energy from the background turbulence while the non-resonant particles undergo alternate heating and cooling periods with a net heating effect. For the ions which are heating effect. which are located in velocity space closer to the phase velocity of the ion acoustic waves are observed to be heated more effectively than those which are located far from the phase velocity of the waves. The numerical results fairly agree with the analytical prediction.

* Supported by AFOSR under grant 86-0156 1. K.M. Yuen and O. Ishihara, Butl. Am. Phys. Soc. 32, 1859 (1987).

CF 12 <u>Velocity Diffusion Coefficient as a Function</u> of Time⁵. H. XIA, and O. ISHIHARA, <u>Terms Tech</u> <u>University</u>—The time-dependent nature of the velocity diffusion coefficient in the plasma turbulence was presented by Weinstock¹ and was developed as extended resonance broadening theories^{2,3}. The theories were developed on the assumption of Gaussian and Markovian stochastic process. Based on the developed on the assumption of Gaussian and Markovian stochastic process. Based on the Mori-Kubo generalized Langevin equation, the exact generalized Fokker-Planck equation corresponding to the Gaussian but otherwise arbitrary stochastic force is studied. It agrees asymptotically to the phenomenological Fokker-Planck equation for systems exhibiting short-time tail decay. And it shows a velocity diffusion coefficient as a function of time. Those theories are examined by the numerical experients where the velocity diffusion coefficients are measured in plasma turbulence. * Supported by AFOSR under grant 86-0156 1. J. Weinstock, Phys. Fluids 12, 1045 (1969). 2. O. Ishihara and A. Hirose, Phys. Fluids 28, 2159 (1985).

2159 (1985). 3. A. Salat, Phys. Fluids 31, 1499 (1988).

Vol. 34, No. 6 (1989)

CONFERENCE RECORD - ABSTRACTS

1989 IEEE INTERNATIONAL CONFERENCE ON PLASMA SCIENCE

May 22 - 24, 1989

Buffalo, New York

3A3-4-INVITED PAPER

Particle Diffusion in Turbulent Fields: Transition from Quasilinear to Nonlinear Stage*

Osamu Ishihara Department of Electrical Engineering Texas Tech University, Lubbock, Texas

Charged particles in random electric fields are known to diffuse over phase velocity resonant layer in velocity space [1,2], while charged particles placed in a sheared magnetic field diffuse over the mode rational surfaces due to ExB random fluctuations [3]. We study in detail the transition of particle diffusion in turbulent fields from quasilinear (constant diffusion rate) to nonlinear regime (time dependent rate) by test particle numerical experiments.

We consider the equation of motion: $d\xi/dt = \zeta$, $d\zeta/dt = R(\xi,t)$, where $R(\xi,t)$ is a random $(\xi, \zeta, R) = (x, v, qE/m)$ for electric field, particle motion in random fields, and $(\xi, \zeta, R) = (y, w \times /L_s. cw E_y / L_s B_0)$ for guiding center motion in a sheared magnetic field, where Ls is the shear length, W is the particle velocity along the magnetic field, and c is the speed of light. Random fields are prescribed to have Gaussian structure in Fourier modes with the spectrum width δk . Trajectories of test particles are followed and statistics of ζ variance is examined. We define the diffusion coefficient by $D_{\zeta} = \frac{1}{2} d\langle \Delta \zeta \rangle / dt$, where the time slope of the ζ variance is to be taken at $t_{sc} \langle t \langle t_{c}, t_{sc} \rangle$ is the autocorrelation time, $1/(\delta k\zeta)$, and t_c is the time at which particles reach the edge of the resonance layer. We have observed that particle diffusion rate is substantially deviated from the quasilinear value and depends on time when the amplitude of random field becomes larger. Such a transition from quasilinear to nonlinear diffusion is observed without particle loss from the resonance region. Time dependent diffusion coefficient may be explained by --the effect of retarded friction caused by turbulent fluctuations [4].

- 1. O. Ishihara and A. Hirose, Phys. Fluids <u>28</u>, 2159 (1985).
- A. Salat, Phys. Fluids 31, 1499 (1988).
 S. P. Hirshman and K. Molvig, Phys. Rev.
- S. P. Hirshman and K. Molvig, Phys. Rev. Lett. <u>42</u>, 648 (1979).
- 4. H. Xia and O. Ishihara, Bull. Am. Phys. Soc. <u>33</u>, 2020 (1988).

*Supported by the AFOSR under Grant 86-0156.

Generalized Langevin Equation for Plasma Turbulence*

Huajuan Xia and Osamu Ishihara Department of Electrical Engineering Texas Tech University, Lubbock, Texas

The generalized Langevin equation, originally derived for the study of Brownian motion by Mori(1) and Kubo(2), has been applied to the study of plasma turbulence. The evolution equation for the particle velocity in an electrostatic plasma turbulence is derived by the projection operator method. The time integral appearing in the equation indicates that the wave-particle interaction in a plasma is not a Markovian process. Thus the wave-particle interaction cannot be described as instantaneous collisions between particles and wave quanta (quasi-particles) as is usually done in a conventional weak turbulence theory.

We consider plasma particles placed in an electrostatic plasma turbulence characterized by random electric fields (Gaussian process). Particles placed initially at (x,v) move in the random fields to (x_t, v_t) in phase space at time t. Then the particle velocity can be expressed as

$v_t = e^{-iJt}v$

where $-i\Sigma$ is the Liouville operator and is given by $-i\Sigma = v \partial/\partial x + (e/m)E(x)\partial/\partial v$. The evolution equation (the generalized Langevin equation) for v_t can be derived by the method of the projection operator as

 $\frac{d}{dt} v_t = \frac{e}{m} e^{-i\Omega t} E(x) - \int_0^t \gamma_{t-\tau}(x,v) v_{\tau}(x,v) d\tau$

where $-i\Omega = -i(1-P)\mathcal{I}$, \mathcal{P} is a projection operator, and $\mathcal{I}_{t-\tau}$ is a friction function. The time integral of the right hand side demonstrates the memory effect in the wave-particle interaction. The cumulant expansion of the characteristic function of v_t , $\langle \exp(i\xi v_t) \rangle$, shows the time-dependent nature of the velocity variance. We find that the memory effect or the influence of the past on particle trajectories in phase space limits the particle diffusion in velocity space, in agreement with the earlier observations (3,4).

1.H. Mori, Prog. Theor. Phys. 33, 423 (1965).

2.R. Kubo, Prog.Theor.Phys.29, Part 1, 255 (1966).

3.0. Ishihara and A. Hirose, Phys. Fluids 28, 2159 (1985).

4.A. Salat, Phys. Fluids 31, 1499 (1988).

*Mork supported by AFOSR under Grant 86-0156.

1P3

CONFERENCE RECORD - ABSTRACTS

1990 IEEE INTERNATIONAL CONFERENCE ON PLASMA SCIENCE

May 21-23, 1990

Oakland, California

2P2-16

PHOTON ACCELERATION: THREE-WAVE INTERACTION

O. Ishihara Department of Electrical Engineering Texas Tech University, Labbock, Texas

Recently a frequency upshifting of short pulses of laser light was studied by Dawson and others[1]. Photons, which propagate in a plasma with effective mass of $\hbar \omega_{po}/c^2$, were shown to be accelerated (upshifted in 60) by propagating down the density gradient of a longitudinal plasma wave.

We study the possibility of photon acceleration from the viewpoint of three-wave interaction[2]. We consider that a photon with momentum $\hbar q_{phi}$ and energy $\hbar \omega_{phi}$ will combine a plasmon with momentum $\hbar q_{phi}$ and energy $\hbar \omega_{phi}$ to give a photon with momentum $\hbar q_{phi}$ and energy $\hbar \omega_{phi}$. The waves will interact one another through the wave-particle interaction as shown below:



where the straight line indicates a plasma particle (charge e, mass m and momentum p). The interaction vertex function for the plasmon-particle can be obtained by the use of the interaction Hamiltonian between particles and longitudinal quasiparticles

$$H_{L} = \int d^{3}x \Psi^{\dagger}(x) e\phi(x) \Psi(x) .$$

where $\Psi(\mathbf{x})$ is a wave function of plasma particles, while the interaction vertex function for the photon-particle can be obtained from the interaction Hamiltonian between particles and transverse quasiparticles

$$H_{\rm r} = -\frac{e}{mc} \int d^3x \Psi^{\dagger}(x) \left(p - \frac{e}{c} A_{\rm e}\right) \cdot A_{\rm i}(x) \Psi(x) ,$$

where A_0 and A_1 are the zero-order and the first-order vector potential, respectively. Our calculation of the vertex function for this three-wave interaction process reveals that there will be no transition to occur if the plasma is cold and stationary. We have found that the transition probability becomes maximum when the plasma particles move in the direction of longitudinal plasma oscillations (plasmons) along with the direction of polarization of vector potential A, resulting in the upshifting of frequency of the outgoing photon fields.

*Supported by AFOSR under Grant 86-0156.

- [1] S.C. Wilks, J.M. Dawson, W.B. Mori, T. Katsouleas, and M.E. Jones, Phys. Rev. Letters 62, 2600 (1989).
- [2] E.G. Harris, in Advances in Plasma Physics, ed. by A. Simon and W.B. Thompson (Interscience, New york, 1969), Vol. 3, p. 157.

EFFECT OF TURBULENT COLLISIONS ON DIFFUSION IN STATIONARY PLASMA TURBULENCE*

H. Xia and O. Ishihara Department of Electrical Engineering Texas Tech University, Lubbock, Texas

Recently the velocity diffusion process was studied by the generalized Langevin equation derived by the projection operator method[1]. The further study shows that the retarded frictional function, $\gamma(t)$, plays an important role in suppressing particle diffusion in the velocity space in stronger turbulence as much as the resonance broadening effect. The retarded frictional effect, produced by the effective collisions due to the plasma turbulence, is described as

$$\gamma(t-\tau) = \frac{q^2}{m^2(v,v)} (E_{\tau}(x,v), E_{t}(x,v)),$$

where (.,.) indicates an average over real (x) and velocity (v) spaces and random fluctuation field $E_{\tau}(x,v)$ is assumed to be a Gaussian, but non-Markovian and non-wide-sense stationary process. The velocity diffusion coefficient in a stationary plasma turbulence in the absence of magnetic field is given as

$$D(\mathbf{v},t) = \frac{q^2}{2m^2} \frac{\partial}{\partial t} \int_0^t d\tau_1 \int_0^t d\tau_2 \,\chi(t-\tau_1) \,\chi(t-\tau_2) \,\beta(\mathbf{v},\tau_1,\tau_2) \,,$$

where

$$\beta(\mathbf{v}, \tau_1, \tau_2) = \langle \mathbf{E}_{\tau_1}(\mathbf{x}, \mathbf{v}), \mathbf{E}_{\tau_2}(\mathbf{x}, \mathbf{v}) \rangle$$
$$= \sum_{\mathbf{k}} \mathbf{E}_{\mathbf{k}} \mathbf{E}_{\mathbf{k}}^* \mathbf{e}^i \left(\mathbf{k} \mathbf{v} \cdot \mathbf{\omega}_{\mathbf{k}} \right) \left(\tau_1 \cdot \tau_2 \right) \left(\mathbf{e}^{-i \mathbf{k} \left(\Delta \mathbf{x}_{\tau_1} \cdot \Delta \mathbf{x}_{\cdot 2} \right)} \right)$$

with < >, a spatial average, is the resonance broadening term . discussed in the reference 2. Here

$$\chi(t) = -\frac{1}{2\pi} \int d\omega \frac{e^{-i\omega t}}{-i\omega + \gamma(\omega)}$$

introduces the effect of turbulent collisions on the velocity space diffusion ($\gamma(\omega)$ is the Fourier transform of $\gamma(t)$). The diffusion rate becomes time-dependent because of the non-Markovian nature of the random process involved. The relation between the proposed formulation and the extended resonance broadening theory[2] is discussed. We also carry out test particle numerical experiment for Langmuir turbulence to test the theories. In a stronger turbulence a deviation of the diffusion rate from the one predicted by both the quasilinear and the extended resonance theories has been observed and is explained qualitatively by the present formulation.

* Supported by AFOSR under grant 86-0156.

[1] H. Xia and O. Ishihara, Bull. Am. Phys. Soc. 34, 1926 (1989).

[2] O. Ishihara and A. Hirose, Phys. Fluids 28, 2159 (1985).

1B-7

Program of the Thirty-Second Annual Meeting of the Division of Plasma Physics Cincinnati, Ohio; 12–16 November 1990

ST21 Photon Acceleration by the Plasmon-Photon Interaction.* O. ISHIHARA, Texas Tech Univer-sity - Photon acceleration in a plasma will be observed as a frequency upshifting in the electromagnetic wave propagating in a plasma. We study photon acceleration from the viewpoint of photon emission by the viewpoint of photon emission by the interaction between plasmon and photon. Three waves, photon, plasmon and photon, interact one another through the interaction between the wave and plasma particles. The relativistic interaction Hamiltonian between particles of species s and photons are given by $H=-e_1^3/d^3x\Psi^+a_A\Psi$, where a is a 4X4 matrix expressed in terms of Pauli matrices and A is a vector potential, while the interaction Hamiltonian between particles and plasmons is $H=e_{a}d^{3}x\Psi^{4}\phi\Psi$, where ϕ is a scalar potential. The frequency upshifting is analyzed in terms of the energy gain of photon as a result of three wave interaction.

*Supported by AFOSR under grant 86-0156.

2R 2 Analysis of the Non-Markovian Behavior <u>_ 1 n</u> Velocity Diffusion Process.* H. XIA and O. ISHIHARA, <u>Texas Tech University</u> -- The non-Markovian diffusion in velocity space was observed in the test particle computer simulations of the plasma turbulence[1]. The intrinsic non-Markovian effects such 25 resonance broadening and retarded turbulent collisions were found to be responsible for such behavior[2]. The numerical experiment in the moderately strong Langmuir turbulence, where the resonance region is bounded by the lower boundary, shows that particles diffuse toward the upper resonance layer after they hit the lower boundary. It is found that the non-Markovian behavior due to the boundary effect further deviates the time-dependent diffusion coefficient predicted by the intrinsic non-Markovian effects.

*Supported by AFOSR under grant 86-0156. 1. O.Ishihara and A.Hirose, Phys.Fluids <u>28</u>, 2159 (1985); A.Salat, Phys.Fluids <u>31</u>, 1499 (1988). 2. H.Xia and O.Ishihara, IEEE Int. Conf. on Plasma Science (Oakland, CA, 1990).

Abstract Submitted for the 18th 1991 IEEE International Conference on Plasma Science

PHOTON ACCELERATION BY PLASMA TURBULENCE*

O. Ishihara Department of Electrical Engineering Texas Tech University, Lubbock, Texas

Recently, we proposed a novel method of photon acceleration¹ in which electromagnetic waves injected into a current carrying plasma are scattered by the plasma turbulence and the frequency of the electromagnetic waves are upshifted by absorbing energy of the plasma turbulence,

We examine the conditions of effective photon acceleration for the above mentioned scheme for the case either plasma particles are non-relativistic or relativistic. We start with Hamiltonian for a particle of species s in an electromagnetic field:

 $\mathcal{H}_{s}^{\text{nonrol}} = \frac{1}{2m_{s}} \left| p - \frac{c_{s}}{c} A(x) \right|^{2} + c_{s} \Phi(x)$ $\mathcal{H}_{s}^{\text{rel}} = c \Omega \cdot \left(p - \frac{c_{s}}{c} A(x) \right) + \beta m_{s} c^{2} + c_{s} \Phi(x)$

where A(x) and $\Phi(x)$ are the vector and scalar potentials of the field. α and β are the dimensionless Dirac 4 x 4 matrices described as 2 x 2 matrices. The three-wave interaction takes place among photon (Ω_1 , q_1), plasmon (Ω_2 , q_2) and photon (Ω_3 , q_3). The scattered photon (Ω_3 , q_3) will have the frequency of $\Omega_3 = \Omega_1 + \Omega_2$ governed by the conservation law. The vertex function for the photon-plasmon-photon interaction is given by²

$$M = \sum_{i} \int d^{3}v \quad f(v) \frac{\pi \hbar e_{z}^{2}}{\gamma^{2} m_{z}^{2}} \left(\frac{2\pi \hbar e_{z}^{2} \Omega_{2}}{V q_{z}^{2} \Omega_{1} \Omega_{3}} \right)^{1/2}$$

$$\times \quad v \cdot \left\{ \frac{2u_{q_{1}\sigma} q_{1} \cdot u_{q_{2}\sigma} - q_{1} u_{q_{1}\sigma'} u_{q_{2}\sigma}}{(\Omega_{1} - v \cdot q_{1}) (\Omega_{2} - v \cdot q_{2})} \left(\frac{-q_{z}^{2}}{\Omega_{1} - v \cdot q_{1}} + \frac{q_{z}^{2}}{\Omega_{z} - v \cdot q_{2}} \right) + \frac{2u_{q_{1}\sigma} q_{z} u_{q_{1}\sigma} - q_{3} u_{q_{1}\sigma'} u_{q_{1}\sigma}}{(\Omega_{3} - v \cdot q_{2}) (\Omega_{3} - v \cdot q_{3})} \left(\frac{q_{z}^{2}}{\Omega_{z} - v \cdot q_{3}} + \frac{q_{z}^{2}}{\Omega_{z} - v \cdot q_{3}} \right) \right\}$$

where $u_{q_1\sigma}$ and $u_{q_3\sigma}$ are unit polarization vectors of photon fields. The terms with $u_{q_1\sigma} u_{q_3\sigma}$ are resulted from relativistic effect. This vertex function suggests that the frequency upshifting becomes maximum when the wave vector of the emitted photons is in the direction of electron streams in the plasma unbulence.

*Supported by AFOSR under Grant 86-0156.

1. O. Ishihara, 1990 IEEE International Conference on Plasma Science, paper 2P2-16 (Oakland, CA).

2. O. Ishihara, Phys. Rev. A (in press).

2. Plasma Waves and

Instabilities

Jshihas

Osamu Ishihara

Texas Tech University Box 4439-MS3102

Lubbock, TX 79409

USA

X

806 742-3463

Alexandre Alexandre Alexandre Obsilogenerations Alexandre Alexandr

Abstract Submitted for the 18th 1991 IEEE International Conference on Plasma Science

Characteristics of Distribution Function in a Time-Dependent Velocity Diffusion Process*

Huajuan Xia and Osamu Ishihara

Department of Electrical Engineering Texas Tech University, Lubbock, Texas

The time-dependency of the particle velocity diffusion in plasma turbulence has been discussed by various theoretical models and shown in the test-particle computer simulations [1]-[4]. The non-Markovian effects are responsible for the time-dependent nature of the diffusion coefficient that deviates far from the quasilinear value at a moderately strong turbulence level in Langmuir turbulence [5].

Under the assumption of Gaussian turbulent fields, the non-Markovian effects were discussed by treating the particle Non-Markovian effects were discussed by freading the particle velocity as a Wiener process. The memory property appears in the spatial increment of particle trajectory as well as in the term of retarded effective turbulent friction in the generalized Langevin equation [5]. We further study such a non-Markovian behavior in velocity diffusion by introducing the non-Gaussian nature in turbulent fields. This modification leads to a deviation of nethability distribution forget of particle subscitic forget the of probability distribution function of particle velocity from the Gaussian distribution. The effect of non-Gaussian turbulent fields on the particle diffusion coefficient is discussed in detail.

To examine the long time behavior of the diffusion : process, we perform test-particle numerical experiments by a process, we perform test-particle numerical experiments by a newly installed MasPar massively parallel computer at the Texas Tech University. In the test-particle simulation, the k-spectrum of stationary Langmuir turbulence is modeled by two different shapes; one is Gaussian-type, and the other is square-like. When particles hit the boundary of resonance region, a fictitious non-Markovian effect on the diffusion coefficient is observed. By varying the spectral shape, this fictitious non-Markovian behavior could be suppressed. The settles of cost-markovian behavior could be suppressed. The results of test-particle numerical experiments are presented and compared with the theory based on the assumption of non-Gaussian turbulent fields.

* Supported by AFOSR under grant 86-0156.

- [1] J. Weinstock, Phys. Fluids 12, 1045 (1969).

- F. Dovell and D. Gresillon, Phys. Fluids 25, 1396 (1982).
 O. Ishihara and A. Hirose, Phys. Fluids 28, 2159 (1985).
 A. Salat, Phys. Fluids 31, 1499 (1988).
 H. Xia and O. Ishihara, 1990 IEEE International Conference on Plasma Science, paper 1B-7 (Oakland, CA).

2. Plasma Waves and

Instabilities

x

Grene Ishiha

Osamu Ishihara Texas Tech University Box 4439-MS3102

Lubbock, TX 79409

USA

. ~

. . .

. . .

806 742-3463

....

PHYSICAL REVIEW A

VOLUME 35, NUMBER 3

Nonresonant wave-particle interaction in semiclassical quasilinear theory

Osamu Ishihara

Department of Electrical Engineering, Texas Tech University, P.O. Box 4439, Lubbock, Texas 79409-4439 (Received 20 August 1986)

A nonresonant nature of wave-particle interaction is clarified from the viewpoint of quantum mechanics. The interaction of particles and quasiparticles can be described by the use of transition probability which is found to have both resonant and nonresonant contributions. The resonant transition probability is known as Fermi's golden rule, which is now supplemented by the nonresonant contribution, resulting in the proper conservation of energy and momentum in the particle-quasiparticle system.

NUCLEAR FUSION, Vol.27, No.8 (1987)

DRIFT ALFVÉN EIGENMODE IN TOKAMAKS

A. HIROSE

Department of Physics, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

O: ISHIHARA Department of Electrical Engineering, Texas Technical University, Lubbock, Texas, United States of America

ABSTRACT. A drift Alfvén eigenmode having a phase velocity significantly larger than the electron diamagnetic velocity is shown to exist in the tokamak magnetic geometry. For typical ohmical heated tokamak discharges the approximate dispersion relation is 40 H 36000. Such a last drift mode has recently been observed in the TEXT tokamak.

1231

ł

SEARCH FOR ION TEMPERATURE GRADIENT DRIVEN ELECTROSTATIC HYDRODYNAMIC INSTABILITY IN TOKAMAKS

A. HIROSE Department of Physics, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

O. ISHIHARA Department of Electrical Engineering, Texas Tech University, Lubbock, Texas, United States of America

ABSTRACT. The ion temperature gradient driven electrostatic instability in tokamak magnetic geometry has been investigated numerically in terms of a generalized ion density moment. For typical tokamak parameters (ratio between the density gradient scale length and the major radius, $\epsilon_a = L_a/R$, about 0 (0.1) and $T_i \cong T_e$) the search for rapidly growing instabilities of a hydrodynamic nature has been unsuccessful, even with the ion temperature gradient relative to the density gradient, $\eta_i = d(\ln T_i)/d(\ln n_o)$, as large as six.

CAN. J. PHYS. VOL. 66, 1988

Resistive ballooning mode in tokamaks

1069

A. HIROSE AND T. L. KROEKER¹ Department of Physics, University of Saskatchewan, Saskatoon, Sask., Canada S7N OWO

> AND O. Ishihara

Department of Electrical Engineering, Texas Tech University, Lubbock, TX 79409, U.S.A. Received May 3, 1988²

The resistive ballooning mode in tokamak geometry has been re-investigated without imposing any ordering among characteristic frequencies, $|\omega|$ (growth rate), ω_0 (diamagnetic frequency), and $\omega_D(\eta)$ (magnetic-drift frequency in the ballooning space). An extensive numerical search for unstable eigenfunctions has failed. Only a stable-ion diamagnetic mode (analytically predicted) has been found. NUCLEAR FUSION, Vol.29, No.5 (1989)

RESISTIVE MHD BALLOONING MODE IN TOKAMAKS

A. HIROSE Department of Physics, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

O. ISHIHARA Department of Electrical Engineering, Texas Tech University, Lubbock, Texas, United States of America

ABSTRACT. A mode equation for the resistive MHD ballooning mode in low beta tokamak magnetic geometry has been derived and analysed without imposing any ordering among the frequencies ω , $\omega_{\rm e}$ (diamagnetic frequency) and $\omega_{\rm D}$ (magnetic drift frequency operator). Only a stable, ion diamagnetic mode has been found. The resistive ballooning mode is strongly stabilized by magnetic shear and toroidal effects.

Phys. Fluids B 2 (2), February 1990

0899-8221/90/020270-10\$02.00

© 1990 American Institute of Physics

270

Resonance broadening in drift wave turbulence

O. Ishihara and C. Grabowski^{a)}

Department of Electrical Engineering, Texas Tech University, Lubbock, Texas 79409

A. Hirose

Department of Physics, University of Saskatchewan, Saskatoon, Saskatchewan, S7N 0W0, Canada

(Received 30 June 1989; accepted 4 October 1989)

Turbulent diffusion of electrons in drift modes in a sheared magnetic field is studied by using a test particle numerical experiment. Electrons diffuse across the magnetic field over mode rational surfaces, where electrons interact with waves in resonance. A spatial diffusion coefficient, which describes resonance broadening, is found to be time dependent and departs from quasilinear predictions in stronger turbulence even well before the time when particles hit the resonance boundary of the rational surfaces.

Accepted for publication

in Physical Review A (1991).

Photon-Plasmon-Photon Interaction

Osamu Ishihara

Department of Electrical Engineering and Department of Physics Texas Tech University, Lubbock, Texas 79409-3102

ABSTRACT

A theory of photon-plasmon-photon three-wave interaction is presented. An electromagnetic wave launched into the plasma turbulence interacts each other through the wave-particle interaction resulting in the emission of an electromagnetic wave with frequency upshifted. The theory is based on the Dirac relativistic wave equation, where the interaction of plasma particles with electromagnetic waves are described. It is shown that the frequency upshifting of the electromagnetic wave can take place in a current carrying plasma.

PACS numbers: 52.40.Db, 52.35.Ra, 52.40.Nk