

DTIC FILE COPY

MISCELLANEOUS PAPER EL-91-5

2



**US Army Corps
of Engineers**

MACROINVERTEBRATES OF LUXAPALILA CREEK, MISSISSIPPI AND ALABAMA, 1987-89

by

Barry S. Payne, Andrew C. Miller, Christina Miller-Way, C. Rex Bingham

Environmental Laboratory

DEPARTMENT OF THE ARMY

Waterways Experiment Station, Corps of Engineers
3909 Halls Ferry Road, Vicksburg, Mississippi 39180-6199

**DTIC
ELECTE
MAR 21 1991
S B D**

AD-A232 870



February 1991

Final Report

Approved For Public Release; Distribution Unlimited

Prepared for US Army Engineer District, Mobile
Mobile, Alabama 36628-0001

91 3 18 124



Destroy this report when no longer needed. Do not return
it to the originator.

The findings in this report are not to be construed as an official
Department of the Army position unless so designated
by other authorized documents.

The contents of this report are not to be used for
advertising, publication, or promotional purposes.
Citation of trade names does not constitute an
official endorsement or approval of the use of
such commercial products.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE: February 1991 3. REPORT TYPE AND DATES COVERED: Final report

4. TITLE AND SUBTITLE: Macroinvertebrates of Luxapalila Creek, Mississippi and Alabama, 1987-89 5. FUNDING NUMBERS:

6. AUTHOR(S): Barry S. Payne, Andrew C. Miller, Christina Miller-Way, C. Rex Bingham

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES): USAE Waterways Experiment Station, Environmental Laboratory, 3909 Halls Ferry Road, Vicksburg, MS 39180-6199 8. PERFORMING ORGANIZATION REPORT NUMBER: Miscellaneous Paper EL-91-5

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES): US Army Engineer District, Mobile, AL 36628-0001 10. SPONSORING/MONITORING AGENCY REPORT NUMBER:

11. SUPPLEMENTARY NOTES: Available from National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161

12a. DISTRIBUTION/AVAILABILITY STATEMENT: Approved for public release; distribution unlimited 12b. DISTRIBUTION CODE:

13. ABSTRACT (Maximum 200 words): A survey of the macroinvertebrate community of four pools and four riffles in Luxapalila Creek, Mississippi and Alabama, was conducted in 1987-89. The purpose was to collect preconstruction data before the creek is impacted by selective clearing and snagging, bank protection, construction of notched sills, and a fabric dam. Luxapalila Creek can be characterized as exhibiting high macroinvertebrate species richness, diversity, and equitable distribution of chironomid (midge) larvae and oligochaete (worm) species. These two groups dominated the fauna and comprised 77 and 62 percent of the macroinvertebrates in pools and riffles, respectively. Total macroinvertebrate density was lower although more stable in pools (6,433-7,229 individuals/sq m) than in riffles where it was higher and more variable (9,662-28,820 individuals/sq m). Species richness of chironomids appeared to be greater in pools than riffles probably because of drift out of riffles and into pools. High winter discharge led to greatly reduced macroinvertebrate densities in all riffles and in two pools

(Continued)

14. SUBJECT TERMS: Invertebrates, Stream ecology 15. NUMBER OF PAGES: 89 16. PRICE CODE:

17. SECURITY CLASSIFICATION OF REPORT: UNCLASSIFIED 18. SECURITY CLASSIFICATION OF THIS PAGE: UNCLASSIFIED 19. SECURITY CLASSIFICATION OF ABSTRACT: UNCLASSIFIED 20. LIMITATION OF ABSTRACT:

13. (Concluded).

during the spring of 1989. Post-construction biological monitoring will determine if community composition and total density is affected by proposed channel modifications.

PREFACE

In September 1987, the US Army Engineer Waterways Experiment Station (WES) initiated invertebrate studies on Luxapalila Creek, Mississippi and Alabama, for the US Army Corps of Engineers, Mobile District (CESAM). The purpose was to collect baseline information on invertebrates to evaluate the effects of proposed channel modifications.

This report was prepared by Dr. Barry S. Payne, Dr. Andrew C. Miller, Ms. Christina Miller-Way, and Mr. C. Rex Bingham, all of the Aquatic Habitat Group (AHG), WES. Mr. Edwin Theriot was Chief, AHG, Dr. Conrad J. Kirby was Chief, Environmental Resources Division, and Dr. John Harrison was Chief, Environmental Laboratory, WES, during preparation of this report. Mr. Brian Peck, CESAM, monitored the study and reviewed an early draft of the report. The report was edited by Ms. Janean C. Shirley of the WES Information Products Division, Information Technology Laboratory.

Commander and Director of WES was COL Larry B. Fulton, EN. Technical Director was Dr. Robert W. Whalin.

This report should be cited as follows:

Payne, B. S., Miller, A. C., Miller-Way, C., and Bingham, C. R. 1991. "Macroinvertebrates of Luxapalila Creek, Mississippi and Alabama, 1987-89," Miscellaneous Paper EL-91-5 , US Army Engineer Waterways Experiment Station, Vicksburg, MS.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

CONTENTS

	<u>Page</u>
PREFACE	1
CONVERSION FACTORS, NON-SI TO SI (METRIC) UNITS OF MEASUREMENT	3
PART I: INTRODUCTION	4
Background	4
Purpose and Scope	4
PART II: STUDY AREA AND METHODS	5
Study Area	5
Methods	7
PART III: RESULTS	9
Physicochemical Conditions	9
Biological Conditions	14
PART IV: DISCUSSION	27
Major Findings	27
Recommendations	28
REFERENCES	30
APPENDIX A: COUNTS OF MAJOR TAXA PER CORE SAMPLE, FALL 1987	A1
APPENDIX B: SPECIES COMPOSITION IN BENTHIC SAMPLES COLLECTED IN THE SPRING OF 1988	B1
APPENDIX C: COUNTS OF MAJOR TAXA PER CORE SAMPLE, FALL, 1988	C1
APPENDIX D: SPECIES COMPOSITION IN SPRING, 1989 SAMPLES	D1
APPENDIX E: COUNTS OF MAJOR TAXA, PER CORE SAMPLE, COLLECTED FROM SITE 4 IN THE FALL OF 1989	E1

CONVERSION FACTORS, NON-SI TO SI (METRIC)
UNITS OF MEASUREMENT

Non-SI units of measurement used in this report can be converted to SI
(metric) units as follows:

<u>Multiply</u>	<u>By</u>	<u>To Obtain</u>
cubic feet per second	0.02831685	cubic metres per second
feet	0.3048	metres
gallons (US liquid)	3.785412	cubic decimetres
inches	2.54	centimetres
miles (US statute)	1.609347	kilometres
square miles	2.589998	square kilometres

MACROINVERTEBRATES OF LUXAPALILA CREEK,
MISSISSIPPI AND ALABAMA, 1987-89

PART I: INTRODUCTION

Background

1. The US Army Corps of Engineers, Mobile District (CESAM) is preparing a flood control plan for Luxapalila Creek, Mississippi and Alabama. The project area includes the lower reach of the creek, from its mouth to river mile (RM) 25.5. The current plan calls for selective clearing and snagging, bank protection, and construction of notched sills, excavation of pools, and a fabric dam. Proposed channel modifications would protect rural areas upstream of Columbus, MS, against floods expected to occur every 1.5 years, and urban areas within Columbus would be protected against floods expected every 5 years.

2. Many aquatic habitats are altered by channel diversion, modification, or construction of dams (Standford and Ward 1979). This demand on lotic habitats has brought about an interest in habitat improvement or development to mitigate losses caused by man's activities. For example, the fabric dam and other modifications proposed for Luxapalila Creek would help to provide access to the upper watershed by walleye for 45 percent of their critical spawning migration period (US Fish and Wildlife Service 1987). A wide array of measures, including construction of sills, artificial riffles, ponds, and planting of riparian vegetation, have been used to improve habitat for aquatic biota in streams throughout the United States (Shields 1983; Woods and Griswold 1981; US Soil Conservation Service 1971a, 1971b; King, Miller, and Glover 1982; Miller 1987).

Purpose and Scope

3. The purpose of this study was to obtain baseline data on sediment characteristics and benthic invertebrates at Luxapalila Creek, Mississippi and Alabama. This information will be used by CESAM to evaluate the effects of proposed channel alterations and mitigation measures on selected reaches of the creek.

PART II: STUDY AREA AND METHODS

Study Area

4. Luxapalila Creek is located in northeast Mississippi and northwest Alabama (Figure 1). The creek originates in southern Marion County near Winfield, AL and flows in a southerly and then a southwestern direction for about 75 miles* before joining the Tombigbee River at RM 362.35. The drainage basin includes 794.5 square miles of mainly forested and agricultural land that is relatively unaffected by urban development. Riparian vegetation consists of bald cypress (*Taxodium distichum*), river birch (*Betula nigrum*), maples (*Acer* spp.) and oaks (*Quercus* spp.).

5. The study area included approximately 30 miles of Luxapalila Creek in Lowndes County, Mississippi, and Lamar County, Alabama. In this reach the creek consists of riffles, runs, and short pools. Most pools are narrow, comparatively shallow, and appear to function more like runs than pools. Substrate in the riffles consists of gravelly sand with very little cobble or bedrock. A 26-mile segment of the creek between Winfield, AL and the Alabama-Mississippi State line was channelized in 1922. In 1967 the lower 2.1 miles of Luxapalila Creek was channelized (Arner et al. 1976; US Army Corps of Engineers 1986; US Fish and Wildlife Service 1987).

6. Luxapalila Creek is near the northern edge of the Gulf Coastal Plains Physiographic Province. The topography is hilly and ranges from low, smoothly rounded hills of 40-50 ft relief within broad intervening valleys to hills and ridges up to 200 ft high separated by narrow valleys with steeply sloping sides. The basin has a temperate climate characterized by long, warm summers, and short, usually mild winters. The basin receives abundant rainfall that is fairly well distributed throughout the year. Annual rainfall at Columbus from 1951 to 1980 was 56.75 in.

7. Four sampling sites, each consisting of a single pool and a nearby riffle, were located on Luxapalila Creek. Sites 1-3 were within the reach to be modified by proposed channel alterations. Site 4, which is in a previously channelized reach, was upstream of proposed channel modifications. The following is a brief description of the four sites surveyed. More detailed data

* A table of factors for converting non-SI to SI (metric) units is presented on page 3.

on sediments and water chemistry at each site can be found in Part III of this report.

Site 1

8. At the site closest to the mouth of the creek (Figure 1) the channel was braided and approximately 80 ft wide. Substrate in the riffle consisted of firm gravel over hardpan clay. There was no submersed vegetation present, although the emergent macrophyte water willow (*Justicia americana*) had colonized gravelly shoals along the left descending bank.




Figure 1. Study sites on Luxapalila Creek, Mississippi and Alabama

9. The pool at site 1 was located approximately 500 ft upriver of the riffle. Approximately 20 percent of the substrate in the lower reach of the pool, where samples were taken, was stabilized by logs or other large snags. Substrate consisted of sand overlain by a 2-in. layer of pea gravel. Canopy coverage was approximately 20 percent and there was no submersed or emergent vegetation in the pool.

Site 2

10. The riffle at site 2 was approximately 50 ft wide and was bordered by extensive stands of *J. americana*. The gravel was colonized by scattered clusters of *Sparganium*: sp. covered by an unidentified algae. The riparian canopy was open.

11. The pool, which was located approximately 100 ft downstream, was approximately 100 ft wide. Substrate consisted primarily of sand and silt overlain by a layer of leaf litter and twigs. Snags were present in the pool, but to a lesser extent than at site 1. No emergent or submersed vegetation

was observed. The riparian cover, which consisted of *T. distichum* and water oak (*Quercus nigra*), covered approximately 50 percent of the channel. Banks were low, 3-4 ft high, and stable.

Site 3

12. The riffle at site 3 was about 50 ft wide, and exposed gravel bars were colonized by *J. americana*. Substrate consisted mainly of gravel that appeared to be slightly less consolidated than in the riffle at site 2 but more consolidated than gravel at site 1. There were no snags in the channel and no riparian cover.

13. The pool was located approximately 1,500 ft downstream of the riffle and 500 ft upstream of a US Geological Survey (USGS) discharge gage (No. 02443500 near Columbus, MS). No submersed vegetation was present and the riparian vegetation did not form a noticeable canopy. *Justicia americana* was present along the border of the pool, but to a lesser extent than in the riffle.

Site 4

14. This site, which was located in Alabama, was approximately 20 miles upstream of site 3; this river reach experienced considerably lower average discharge than the lower reach. This reach was straight (the result of channel straightening in the 1920's, Figure 1), less than 50 ft wide, with a nearly closed canopy consisting of oaks (*Quercus* spp.), maples (*Acer* spp.), sweetgum (*Liquidambar styraciflua*), and sycamore (*Platanus occidentalis*). River banks were 8-10 ft high and many trees along the bank were undercut by erosion.

15. Meanders had begun to form within the channelized reach at site 4. The riffle was less than 2 ft deep where samples were taken. There were no exposed gravel bars or emergent vegetation. The pool was located approximately 500 ft downstream of the riffle and had been created by woody snags.

Methods

16. Macroinvertebrates were collected in September, 1987, June and September 1988, and June and October, 1989. Sites 1-3 were sampled from the beginning of the survey through June 1989; site 4 was first sampled in June 1988 and the final sample was taken in October, 1989. Therefore, four consecutive samples were taken at each site; two in the spring and two in the fall. A decision was made by CESAM to add site 4 after the main sampling program had

begun. Counts of major taxa and species composition for each of the sampling periods are presented in Appendixes A-E.

17. At each site, 13 sediment samples were collected from each pool and each riffle. Ten samples were taken for macroinvertebrates, one for total organic content, and two (that were later combined) for grain size analysis. All samples were obtained with a hand-held coring device (Miller and Bingham 1987) that penetrated to a depth of 10 cm and sampled 0.0079 sq m. Organic content was determined by first drying the sample at 65° C, weighing, then heating in a muffle furnace to 550° C. The weight after firing at this latter temperature is termed "ash-free dry weight" and is the loss due to volatilization of organic material. Grain size was determined by sieving the sediments through a standard set of USGS sieves, and weighing each fraction. Sediment samples obtained for macroinvertebrates were preserved in the field with buffered 10-percent Formalin that was stained with rose bengal to facilitate removal of organisms.

18. In the laboratory benthic invertebrates were removed from sediments by an elutriation process. Sediment samples were agitated (swirled in a 3-gal bucket and poured through a 500- μ mesh sieve. Lighter material (detritus and invertebrates) was poured out of the bucket, sand and gravel remained on the screen, and fine silt passed through the screen. Each sample was elutriated five times; lighter material was combined and sand and gravel were discarded. Tests have indicated that this process retrieves 90-100 percent of the invertebrates on sand and gravel substrate.

19. Invertebrates were picked from the elutriated sample with the aid of a binocular microscope. Organisms were first sorted to major group (chironomids, oligochaetes, ephemeropterans, etc.) and counted. Following this initial analysis, chironomids, oligochaetes, and other invertebrates were identified to the lowest possible taxon with appropriate keys. Voucher specimens have been retained at the US Army Engineer Waterways Experiment Station (WES). A complete listing of invertebrates found at each sampling site appears in Appendix A.

PART III: RESULTS

Physicochemical Conditions


Discharge

20. The mean daily discharge for Luxapalila Creek at Columbus, MS (16 years of record; Tharpe et al. 1987) is 1,130 cfs but varies greatly from winter and spring maxima to summer and fall minima (Figure 2a). Mean daily discharge was usually greater than 1,000 cfs in winter and spring and often ranged up to 6,000 cfs. During the 16-year period of record the maximum daily discharge was 40,400 cfs on 14 April, 1979. Mean daily discharge was low (< 500 cfs and often 50 to 200 cfs) between June and August; the minimum discharge reported by Tharpe et al. 1987) occurred during the present study and was 20 cfs on 19 August, 1988.


21. Mean daily discharge during the winter (1986-87) prior to this study was near average (Figure 2b). Winter and spring discharge in 1987-88 was low; values less than 1,000 cfs were typical and no values greater than 3,000 cfs occurred. In contrast, the winter and spring of 1988-89 were characterized by higher than average daily discharge; during this period a maximum value of nearly 17,000 cfs was recorded (i.e., approximately three times higher than the average maximum value; compare Figures 2a and 2b). The mean daily discharges for dates sampled during this survey are given below:

<u>Sampling Date</u>	<u>Mean, cfs</u>
16 September 1987	102
27 June 1988	40
28 June 1988	37
10 October 1988	216
11 October 1988	177
31 May 1989	239
1 June 1989	217
2 November 1989	194

22. The present study included samples affected by a wide range of hydrologic conditions. The samples collected in the fall of 1987 were taken during a low-water period after a winter and spring characterized by high water. Samples collected in 1988 were taken during a sustained drought after extremely low water during the previous winter and spring. The June 1989 samples were collected after high discharge that occurred during the winter and spring of 1988-89.



a. Average daily discharge \pm standard error, 1974-88



b. Average daily discharge during the study period (sampling periods are noted with an asterisk)

Figure 2. Average daily discharge at Luxapalila Creek, Mississippi

Water chemistry


23. The water in Luxapalila Creek was moderately low in both calcium hardness (averaging 7 to 8 mg/l) and alkalinity (averaging 7-10 mg/l), was slightly acidic, and was clear to moderately turbid (Figures 3a-3d). Dissolved oxygen was high (6.6-8.2 mg/l) and was always greater than 90-percent air saturation on dates when macroinvertebrate and sediment samples were taken. These aspects of water chemistry are related to land use and soil conditions in the basin. The low hardness and alkalinity reflect moderate to low deposits of limestone. Clear to moderate turbidity is a consequence of lack of heavily farmed agricultural land.

24. Community composition and density of benthic invertebrates are primarily affected by substrate composition and water velocity (Hynes 1970). It should be noted that lakes and rivers low in carbonates are generally not as productive (i.e., producing large amounts of biomass per unit of time) as hard-water habitats (Russell-Hunter 1970). Despite their generally lower productivity, soft-water habitats that are relatively unaffected by industrial, agricultural, or residential development (such as Luxapalila Creek) provide valuable habitat for a diverse fauna.


Sediment characteristics

25. The substrate of Luxapalila Creek can be characterized as poorly sorted gravel and coarse sand (Figures 4a and 4b). At all sites the silt-clay fraction (< 0.02 mm), and cobble fraction (< 100 mm) each constituted less than 5 percent of the total sediment weight. Median grain sizes for all samples ranged from 4.11 to 18.13 mm; sorting coefficients ranged from 0.90 to 2.72. Pools typically had slightly smaller median particle diameters than did riffles (7.79, ± 2.32 (\pm standard deviation (SD)) versus 10.23 ± 4.56 , respectively). However, these differences are nonsignificant and should not obscure the fact that with respect to substrate characteristics pools and riffles in Luxapalila Creek are relatively similar. Differences between pools and riffles were affected by season. During the summer when discharge events capable of redistributing gravel became less frequent, differences between pools and riffles became even less apparent.


26. The average substrate organic content at all sites and habitat types was low and equaled 1.18 percent (± 0.89). There was no significant difference in percentage organic carbon between pools (1.05, ± 0.41) and riffles (1.31, ± 1.19). In addition, there were no significant differences among stations (pools and riffles combined) for all dates. Average organic content




a. Calcium



b. Total alkalinity




c. pH




d. Turbidity

Figure 3. Five-day averages for selected water quality parameters at Columbus, Luxapalila Creek, Mississippi. Data courtesy of the Columbus Water Department



a. Pools



b. Riffles

Figure 4. Sediment characteristics in pools and riffles in Luxapalila Creek, Mississippi and Alabama during the study period


for all dates was 1.2 (± 0.63), 1.1 (± 0.47), 0.93, (± 0.33), and 0.88 (± 0.41) for sites 1, 2, 3, and 4, respectively. There was no significant difference in organic content with respect to season, although organic contents of sediments were slightly less during the spring (0.91, ± 0.45) than in the fall ($1.5 \leq (1.5, \pm 1.15)$).

Biological Conditions


Macroinvertebrate density

27. The density of macroinvertebrates, averaged for all pools and dates, equaled 7,364 individuals per sq m. Variation in average macroinvertebrate density in pools was not great among sites or dates (Figure 5a). The lowest average density was observed at pool 4 in October 1988 and equaled 5,427 individuals per sq m; the highest average density of 12,561 individuals per sq m was observed at pool 2 in September 1987. Neither spatial nor seasonal density patterns were evident among sites and dates. The ratio of the total range of variation in average density (7,134 individuals per sq m) to the overall average density (7,364 individuals per sq m) equaled 1.0. This low value indicated lack of extreme variation in macroinvertebrate density data among pools and sampling dates.


28. In contrast to pools, average macroinvertebrate density in riffles varied greatly among sites and dates (Figure 5b). Averaged for all sites and dates, macroinvertebrate density in riffles equaled 15,964 individuals per sq m and was slightly greater than twice the overall average density in pools. At sites 1, 2, and 3 (sampled from fall 1987 through spring 1989), average density appeared to follow a seasonal pattern. At each of these sites, the average density in spring was substantially less than in the preceding fall. At site 4, the average density of macroinvertebrates declined between spring 1988 and fall 1989; however, density at this site declined between October 1988 and June 1989 as was observed at the other three riffles. Site- and date-specific average densities in riffles varied from 2,184 (riffle 4, November 1989) to 40,276 (riffle 2, September 1987) individuals per sq m. The ratio of this range (38,092 individuals per sq m) to the overall average density (15,964 individuals per sq m) equaled 2.4, indicating greater than twice as much variation in total macroinvertebrates in riffles than in pools.



a. Density ($\bar{x} \pm SE$) of all macroinvertebrates in pools



b. Density ($\bar{x} \pm SE$) of all macroinvertebrates in riffles



lux1
c. Percentage composition of major macroinvertebrates in pools and riffles

Figure 5. Characteristics of macroinvertebrates in Luxapalila Creek, Mississippi and Alabama, 1987-89

29. A summary of mean macroinvertebrate density (and standard error, SE) by sampling date and habitat type, appears below:

<u>Date</u>	<u>Pools</u>		<u>Riffles</u>	
	<u>Mean</u>	<u>SE</u>	<u>Mean</u>	<u>SE</u>
Sep 87	8,587	4,426	28,820	20,469
Jun 88	7,229	2,421	15,248	3,985
Oct 88	7,025	2,221	18,730	8,553
Jun 89	6,433	1,701	9,662	12,614

The greater stability of biological conditions in pools than riffles is indicated by the more consistent mean and mean-to-SE ratio of macroinvertebrate density in the former habitat.

30. Analysis of variance of total macroinvertebrate density was performed for spring and fall 1988 and spring 1989 samples (i.e., those dates when all four pools and riffles were sampled). Significant variation in total macroinvertebrate density was evident among sites in both the spring and fall of 1988 (Table 1). Paired comparisons of sites confirmed the intersite patterns in Figures 5a and 5b; namely, density was higher in riffles than pools, intersite variation among riffles was high, and intersite variation among pools was low (Table 2). In both the spring and fall of 1988, 17 of 28 possible pairwise combinations between sites revealed significant differences. In the spring of 1988, 14 of these significant differences were for pool-versus-riffle comparisons, and all such comparisons indicated lower densities in pools than riffles. The remaining differences observed in spring involved inter-riffle comparisons, with densities at riffle 2 being significantly greater than densities at the other three riffles. In the fall, 12 of the 17 significant differences among sites involved riffle-versus-pool comparisons, and, as in the spring, riffles always had higher densities than did pools. The remaining five significantly different pair comparisons involved intersite differences among riffles. Riffle 4 had significantly lower macroinvertebrate density than all three other riffles, and riffle 2 had higher density than riffles 1 and 3. No significant intersite differences among pools were evident in either the spring or fall of 1988.

31. Lack of significant intersite differences (i.e., among pool, pool versus riffle, or among riffle) in total macroinvertebrate densities in the spring of 1989 (Table 1) was principally the result of the reduction in density in riffles during that sampling period (Figure 5b). These low densities in riffles in the spring of 1989 eliminated pool-versus-riffle differences

Table 1

Analysis of Variance of Total Macroinvertebrate Density at all Sites
in Luxapalila Creek, Spring and Fall, 1988 and Spring, 1989

<u>Date</u>	<u>Source</u>	<u>DF</u>	<u>Sum Squares</u>	<u>Mean Square</u>	<u>F</u>	<u>p</u>
Spring 1988	Between sites	7	53,180	7,597	29.1	0.0001
	Within sites	32	8,346	261		
	Total	39	61,526			
Fall 1988	Between sites	7	167,338	23,905	89.0	0.0001
	Within sites	32	8,592	269		
	Total	39	175,930			
Spring 1989	Between sites	7	58,217	8,317	1.93	0.097*
	Within sites	32	137,881	4,309		
	Total	39	196,098			

* Not significant at the 0.05 level.

that were evident in 1988. Furthermore, within-riffle density in the spring of 1989 was high and obscured the between-riffle differences that were evident in the spring of 1989.

32. Chironomids and oligochaetes were the numerically dominant macroinvertebrates in both pools and riffles (Figure 5c). These two taxa comprised an average of 77 and 62 percent of all macroinvertebrates in Luxapalila Creek pools and riffles, respectively. *Corbicula fluminea*, trichopterans (caddisflies), acarines (mites), and nematodes (unsegmented worms) were less abundant than chironomids and oligochaetes but also comprised a substantial proportion of the total macroinvertebrate community. The bivalve community was dominated by the introduced Asiatic clam, *Corbicula fluminea*. Dominant trichopterans were *Hydroptila* and *Chimarra*. Bivalves, trichopterans, and acarines were more abundant in riffles than pools, but nematodes were more abundant in pools than riffles. *Corbicula*, *Hydroptila*, and *Chimarra* typically are found in lotic habitats. Neither acarines nor nematodes were identified to a lower taxonomic level.

33. The average density of chironomids, the dominant major taxon, was less variable in pools (Figure 6a) than in riffles (Figure 6b). Maximum density of this group was higher and minimum density was lower in riffles than pools. Averaged for all sites and dates, chironomid density in pools and riffles was 3,981 and 7,419 individuals per sq m, respectively. Site- and

Table 2

Paired Comparisons of Site-Specific Density of Total Macroinvertebrates
in Luxapalila Creek, Spring and Fall 1988

		Spring 1988		Fall 1988	
		Mean Difference*	Scheffe's F-test	Mean Difference*	Scheffe's F-test
Interpool comparisons	P1 vs P2	-15.4	0.325	3.2	0.014
	P1 vs P3	8.4	0.097	-27.0	0.970
	P1 vs P4	-5.8	0.046	8.8	0.103
	P2 vs P3	23.8	0.776	-30.2	1.213
	P2 vs P4	9.6	0.126	5.6	0.042
	P3 vs P4	-14.2	0.276	35.8	1.705
	Pool-to- riffle comparisons	P1 vs R1	-62.0	5.264**	-92.0
P1 vs R2		-106.8	15.620**	-188.8	47.413**
P1 vs R3		-51.2	3.590**	-90.4	10.870**
P1 vs R4		-47.0	3.025**	-11.4	0.173
P2 vs R1		-46.6	2.974**	-95.2	12.055**
P2 vs R2		-91.4	11.440**	-192.0	49.034**
P2 vs R3		-35.8	1.755	-93.6	11.653**
P2 vs R4		-31.6	1.367	-14.6	0.284
P3 vs R1		-70.4	6.787**	-65.0	5.620**
P3 vs R2		-115.2	18.174**	-161.8	34.822**
P3 vs R3		-59.6	4.864**	-63.4	5.347**
P3 vs R4		-55.4	4.203**	15.6	0.324
P4 vs R1		-56.2	4.325**	-100.8	13.515*
P4 vs R2	-101.0	13.969**	-197.6	51.936**	
P4 vs R3	-45.4	2.823	-99.2	13.089**	
P4 vs R4	-41.2	2.324**	-20.2	0.543	
Inter-riffle comparisons	R1 vs R2	-44.8	2.748**	-96.8	12.464**
	R1 vs R3	10.8	0.160	1.6	0.003
	R1 vs R4	15.0	0.308	80.6	8.641**
	R2 vs R3	55.6	4.233**	98.4	12.879**
	R2 vs R4	59.8	4.897**	177.4	41.861**
	R3 vs R4	4.2	0.024	79.0	8.301**

* Negative values indicate lower mean density at first site listed for each paired comparison.

** Probability of $F < 0.05$.


date-specific average density ranged from 1,953 to 7,036 individuals per sq m in pools and from 709 to 21,139 individuals per sq m in riffles. The ratios of these ranges to the overall average densities in pools and riffles were 1.3 and 2.9, respectively. In pools, variation of chironomid density did not follow a discernible intersite or seasonal pattern. In riffles, chironomid density was generally lowest in riffle 4 and highest in riffle 2. The seasonal pattern for total macroinvertebrates of lower density in spring than in the preceding fall was evident for chironomids in riffles 1 and 2. Riffle 1 was especially depauperate of chironomids in June 1989 compared to the three previous dates, and all four riffles had relatively low chironomid densities in June 1989.

34. The average density of oligochaetes was similar in pools and riffles and varied greatly among sites and dates (Figures 6c and 6d). Averaged for all sites and dates, the densities of oligochaetes in pools and riffles were 1,697 and 2,861 individuals per sq m, respectively. Site- and date-specific average densities of oligochaetes ranged from 178 to 3,620 and from 51 to 9,709 individuals per sq m in pools and riffles, respectively. Exceptionally low densities were observed during June 1989 (pools 2 and 4 and riffle 4) and in November 1989 in both the pool and riffle of site 4 (the only site samples in November 1989).


35. *Corbicula fluminea*, which was moderately abundant in riffles, had especially low densities at all riffles in June 1989 (Figure 7). Less than 500 individuals per sq m were found at riffle 1 in the spring of 1989, although densities of this clam had averaged 2,000 individuals per sq m during 1988. In riffles 2 and 3, 1988 densities of *C. fluminea* averaged 1,000 and 2,750 individuals per sq m, respectively, but no Asiatic clams were collected from these two riffles in June 1989. No *C. fluminea* were found in riffle 4 in June 1989, although high density populations were never noted at this site.

Community composition


36. In Luxapalila Creek the abundant chironomid and oligochaete communities were rich in species. In both pools and riffles, cumulative species of chironomids and oligochaetes was a linear function of the cumulative number of individuals (Figures 8a and 8b). The rate of acquisition of new species was clearly higher in pools than riffles. In pools, 63 species of chironomids were represented among 1,503 individuals identified to the species level. In riffles, 50 species of chironomids were represented among 1,910 individuals.




a. Chironomids in pools



b. Chironomids in riffles



c. Oligochaetes in pools



d. Oligochaetes in riffles

Figure 6. Total density of chironomids and oligochaetes in pools and riffles, Luxapalila Creek, Mississippi and Alabama, 1987-89






Figure 7. Total density of *Corbicula fluminea* in riffles in Luxapalila Creek, Mississippi and Alabama, 1987-89

37. The distribution of individuals among species of both chironomids and oligochaetes was highly equitable (i.e., evenly distributed) based on Simpson's index of equitability (Simpson 1949), and equitability was slightly higher in pools than riffles (Tables 3 and 4 for chironomids and oligochaetes, respectively). The value of this index equals one minus the sum of the proportional abundance of each species in a community. The index is relatively insensitive to underestimation of species richness and ranges from theoretical minima and maxima approaching 0.0 and 1.0, respectively. Values between 0.2 and 0.8 are observed in most samples of naturally occurring communities (e.g., Whittaker 1965). Simpson's index of equitability was slightly less than 1.0 for both taxonomic groups in both habitats (Tables 3 and 4 for chironomids and oligochaetes, respectively). The theoretical maximum equitability of chironomids in pools would have been observed if there was equal distribution of the 1,503 individuals among all 63 species identified from pool samples (Simpson's



a. Naidids and tubificids



b. Chironomids

Figure 8. Species richness as a function of cumulative individuals for oligochaetes and chironomids, Luxapalila Creek, Mississippi and Alabama, 1987-89

Table 3
Distribution of Individuals Among Species of Chironomids
in Luxapalila Creek, Mississippi*

Species	Pools		Riffles	
	p	n	p	n
Chironomidae				
Chironominae				
Chironomini				
<i>Chironomus</i> sp.	0.0160	24	0.0005	1
<i>Cladopelma</i> sp.	0.0013	2	0.0000	0
<i>Cryptochironomus fulvus</i>	0.0213	32	0.0011	2
<i>Cryptochironomus</i> sp.	0.0027	4	0.0011	2
<i>Dicrotendipes neomodestus</i>	0.0140	21	0.0661	124
<i>Dicrotendipes nervosus</i> Type I	0.0319	48	0.0016	3
<i>Dicrotendipes nervosus</i> Type II	0.0013	2	0.0000	0
<i>Dicrotendipes</i> sp. I	0.0120	18	0.0080	15
<i>Endochironomus</i> sp.	0.0093	14	0.0005	1
<i>Glyptotendipes</i> sp.	0.0033	5	0.0005	1
<i>Harnischia</i> sp.	0.0040	6	0.0005	1
<i>Microtendipes</i> sp.	0.0007	1	0.0000	0
<i>Nilothauma babiysi</i>	0.0106	16	0.0016	3
<i>Parachironomus abortivus</i>	0.0020	3	0.0005	1
<i>Paracladopelma undine</i>	0.1240	186	0.0016	3
<i>Paralauterborniella nigrohalteralis</i>	0.0313	47	0.0016	3
<i>Paratendipes albimanus</i>	0.0013	2	0.0000	0
<i>Paratendipes nudisquama</i>	0.0007	1	0.0000	0
<i>Phaenopsectra dyari</i>	0.0838	126	0.0000	0
<i>Phaenopsectra flavipes</i>	0.0186	28	0.0000	0
<i>Polypedilum convictum</i>	0.0067	10	0.0762	143
<i>Polypedilum fallax</i>	0.0013	2	0.0000	0
<i>Polypedilum illinoense</i>	0.0126	19	0.0053	10
<i>Polypedilum nr. scaloneum</i>	0.1190	179	0.0410	77
<i>Pseudochironomus</i> sp.	0.0146	22	0.0016	3
<i>Robackia</i> sp.	0.0033	5	0.0624	117
<i>Stenochironomus</i> sp.	0.0007	1	0.0000	0
<i>Stictochironomus</i> sp.	0.0020	3	0.0005	1
<i>Tribelos</i> sp.	0.0000	0	0.0005	1
<i>Xenochironomus</i> sp.	0.0053	8	0.0213	40
Unidentified chironomini		4		5
Tanytarsini				
<i>Cladotanytarsus</i> sp.	0.0146	22	0.0320	10
<i>Lauterborniella</i> sp.	0.0013	2	0.0000	0
<i>Micropsectra</i> sp.	0.0033	5	0.0000	0
<i>Rheotanytarsus</i> sp.	0.0140	21	0.0736	138
<i>Stempellina</i> sp.	0.0013	2	0.0000	0

(Continued)

* Note: p = relative abundance; n = number present.

Table 3 (Concluded)

Species	Pools		Riffles	
	p	n	p	n
<i>Stempellinella</i> sp.	0.0020	3	0.0000	0
<i>Tanytarsus coffmani</i>	0.0000	0	0.0016	3
<i>Tanytarsus glabrescens</i>	0.0319	48	0.0474	89
<i>Tanytarsus querlus</i>	0.1530	230	0.0245	46
Unidentified tanytarsini		0		4
Orthoclaadiinae				
<i>Brillia</i> sp.	0.0007	1	0.0000	0
<i>Coryoneura celeripes</i>	0.0013	2	0.0144	27
<i>Coryoneura taris</i>	0.0013	2	0.0197	37
<i>Coryoneura</i> sp.	0.0007	1	0.0016	3
<i>Cricotopus bicinctus</i>	0.0120	18	0.0938	176
<i>Cricotopus trifascia</i>	0.0000	0	0.0011	2
<i>Cricotopus</i> sp.	0.0013	2	0.0016	3
<i>Eukiefferiella</i> sp.	0.0027	4	0.0032	6
<i>Nanocladius crassicornus</i>	0.0027	4	0.0037	7
<i>Nanocladius distinctus</i>	0.0027	4	0.0059	11
<i>Nanocladius rectinervis</i>	0.0000	0	0.0027	5
<i>Nanocladius minimus</i>	0.0013	2	0.0000	0
<i>Nanocladius</i> sp.	0.0067	10	0.0006	1
<i>Parakiefferiella</i> sp.	0.0619	93	0.0448	84
<i>Rheocricotopus</i> sp.	0.0000	0	0.0053	10
<i>Thienemanniella</i> nr. <i>fusca</i>	0.0047	7	0.2569	482
<i>Thienemanniella xena</i>	0.0007	1	0.0219	41
Unidentified orthoclaadiinae		5		15
Tanypodinae				
<i>Ablabesymia mallochi</i>	0.0033	5	0.0011	2
<i>Ablabesymia parajanta</i>	0.0446	67	0.0139	26
<i>Ablabesymia tarella</i>	0.0020	3	0.0016	3
<i>Clinotanypus</i> sp.	0.0007	1	0.0000	0
<i>Labrundinia pilosella</i>	0.0007	1	0.0059	11
<i>Macropelopia</i> sp.	0.0013	2	0.0005	1
<i>Natarsia</i> sp.	0.0013	2	0.0005	1
<i>Nilotanypus</i> sp.	0.0007	1	0.0219	41
<i>Pentaneura</i> sp.	0.0067	10	0.0000	0
<i>Procladius</i> sp.	0.0599	90	0.0021	4
<i>Thienemannimyia</i> sp.	0.0013	2	0.0005	1
Unidentified tanypodinae		34		10
Diamesinae				
<i>Potthasia</i> sp.	0.0007	1	0.0000	0
Unidentified Chironomidae		67		85
Total number of species		63		85
Total number of individuals identified		1,504		1,876
Simpson's index of equitability		0.93		0.90

Table 4

Distribution of Individuals Among Species of Naidid and Tubificid Oligochaetes
in Pool Versus Riffles in Luxapalila Creek, Mississippi*

Taxon	Pools		Riffles	
	n	p	n	p
Naididae				
<i>Amphichaeta leydigi</i>	6	0.0160	1	0.0016
<i>Bratislavia bilongata</i>	0	0.0000	2	0.0033
<i>Bratislavia unidentata</i>	0	0.0000	4	0.0065
<i>Chaetogaster diaphanus</i>	3	0.0080	51	0.0831
<i>Dero digitata</i>	3	0.0080	10	0.0163
<i>Dero furcata</i>	13	0.0347	0	0.0000
<i>Dero nivea</i>	8	0.0213	78	0.1270
<i>Dero obtusa</i>	11	0.0293	76	0.1238
<i>Dero trifida</i>	3	0.0080	9	0.0147
<i>Dero sp.</i>	5	0.0133	3	0.0049
<i>Homochaeta naidina</i>	1	0.0027	0	0.0000
<i>Nais behningi</i>	0	0.0000	3	0.0049
<i>Nais bretscheri</i>	0	0.0000	3	0.0049
<i>Nais communis</i>	0	0.0000	3	0.0049
<i>Nais elinquis</i>	1	0.0027	0	0.0000
<i>Nais pardalis</i>	12	0.0320	90	0.1466
<i>Nais pseudobtusa</i>	0	0.0000	10	0.0163
<i>Nais simplex</i>	0	0.0000	3	0.0049
<i>Nais variabilis</i>	2	0.0053	53	0.0863
<i>Piquetiella michiganensis</i>	5	0.0133	29	0.0472
<i>Pristina aquiseta</i>	4	0.0107	28	0.0456
<i>Pristina leidyi</i>	6	0.0160	33	0.0537
<i>Pristina synclites</i>	29	0.0773	1	0.0016
<i>Pristina sp.</i>	0	0.0000	1	0.0016
<i>Pristinella jenkiniae</i>	2	0.0053	1	0.0016
<i>Pristinella longidentata</i>	3	0.0080	2	0.0033
<i>Pristinella longisoma</i>	0	0.0000	9	0.0147
<i>Pristinella osborni</i>	8	0.0213	52	0.0847
<i>Pristinella sima</i>	0	0.0000	2	0.0033
<i>Slavina appendiculata</i>	2	0.0053	21	0.0342
<i>Specaria josinae</i>	52	0.1387	1	0.0016
<i>Stevensoniana trivandrama</i>	1	0.0027	26	0.0423
Total identified naidids	180	0.4800	602	0.9805
Total unidentified naidids	0		8	
Total naidids	180		610	
Tubificidae				
<i>Aulodrilus limnobius</i>	19	0.0507	0	0.0000
<i>Aulodrilus piqueti</i>	127	0.3387	4	0.0065

(Continued)

* Note: n = number present; p = relative abundance.

Table 4 (Concluded)


Taxon	Pools		Riffles	
	n	p	n	p
<i>Aulodrilus pluriset</i>	6	0.0160	0	0.0000
<i>Branchiura sowerby</i>	31	0.0827	7	0.0114
<i>Limnodrilus hoffmestri</i>	11	0.0293	0	0.0000
<i>Limnodrilus rubripenis</i>	0	0.0000	1	0.0016
Total identified tubificids	195	0.5200	12	0.0195
Total unidentified tubificids	78		28	
Total tubificids	275		40	
Tubificid-to-Naidid Ratio	1.53		0.07	
Simpson's Index of Equitability	0.84		0.91	

index value equal to 0.98). In contrast, the most inequitable possible distribution would have been observed if there was a single individual of each of 62 species and 1,441 individuals of one extremely dominant species (Simpson's index equal to 0.08). The observed value of 0.92 is near the theoretical maximum of 0.98. Similarly high values were observed for chironomids in riffles and oligochaetes in both pools and riffles.


38. The unusually high equitability of chironomids and oligochaetes in pools and riffles was especially evident in plots of species-specific percentage abundance as a function of species rank (Figures 9a-9d). It is not unusual for the two or three most abundant species in a community of benthic macroinvertebrates to comprise 75-90 percent of the entire community. In Luxapalila Creek, no individual species comprised greater than 30 percent and the three most abundant species represented approximately 50 percent of the total community. Species relative abundance in these highly equitable communities spanned only 2.0 to 2.5 orders of magnitude in each community sample, although total species richness was high, ranging from 27-63.

39. Distinct differences were apparent in the most abundant species of chironomids in pools versus riffles, although at least a few individuals of most species could be found in either habitat type (Table 3). None of the six most abundant species in pools were among the six most abundant species in riffles. The six dominant chironomids in pools were *Tanytarsus querlus* (15.3 percent), *Paracladopelma undine* (12.4 percent), *Polypedilum* nr. *scalaenum* (11 percent) *Phaenopsectra dyari* (8.4 percent), *Parakiefferiella* sp. (6.2 percent), and *Procladius* (6.0 percent). All but one of these species (*P. dyari*) were obtained in riffles as well as pools, although both *P. undine* and *Procladius* sp. were very uncommon in riffles. *Tanytarsus querlus*, *P.* nr. *scalaenum*, and *Parakiefferiella* sp. were moderately abundant in riffles; these species comprised 2.5, 4.1, and 4.5 percent, respectively, of the chironomids in riffles. Considered in total, the six most abundant species in pools comprised 60.2 percent of the pool assemblage of chironomids, versus 11.5 percent of the riffle assemblage of chironomids.


40. The six most abundant species of chironomids in riffles were *Thienemanniella* nr. *fusca* (25.7 percent), *Cricotopus bicinctus* (9.4 percent), *Polypedilum convictum* (7.6 percent), *Rheotanytarsus* sp. (7.4 percent), *Dicrotendipes neomodestus* (6.6 percent), and *Robackia* sp. (6.2 percent). The combined abundance of these six species in riffles and pools was 62.9 and 6.3 percent, respectively. None of the dominant species in riffles was found




a. Chironomids in pools



b. Chironomids in riffles



c. Naidids and tubificids in pools



d. Naidids and tubificids in riffles

Figure 9. Species relative abundance in relation to dominance for chironomids and oligochaetes in pools and riffles, Luxapalila Creek, Mississippi and Alabama, 1987-89

in greater than 1.6 percent abundance among pool-dwelling chironomids, although some individuals of all six species were obtained from the more lentic habitat.

41. Differences were also evident between pool and riffle chironomid communities among uncommon species. Eighteen uncommon species of chironomids were found in pools but not in riffles, but only four such species were found only in the riffles (Table 3). Community structure in terms of uncommon species must be evaluated cautiously, because a high degree of uncertainty is associated with presence or absence data for uncommon species. Nonetheless, the magnitude of pool-versus-riffle difference in the number of species unique to each habitat type (i.e., a ratio of 18 to 4) indicates that more species of this group probably occurred in pools than in riffles. Drift of chironomid larvae is a major means of dispersal in rivers and streams (Simpson and Bode 1980). Settlement of drifting larvae in pools is probably more likely than in riffles given the more depositional nature of the former habitat. Thus regardless of the origin of drifting chironomids, successful immigration via drift into pools is probably higher than into riffles and may account for the slightly greater richness observed in pools versus riffles.

42. As with chironomids, the pool community of oligochaetes was dominated by different species than the riffle community (Table 4). The six most abundant species in pools included three tubificids (*Aulodrilus piqueti* (33.9 percent), *Branchiura sowerbyi* (8.3 percent), and *A. limnobioides* (3.5 percent)) and three naidids (*Specaria josinae* (13.9 percent), *Pristina synclites* (7.7 percent), and *Dero furcata* (3.5 percent)). Two of these species (*A. limnobioides* and *D. furcata*) were not obtained in riffles, and the other four dominant species in pools were uncommon in riffles. The combined abundance of these six species was 72.4 percent in pools and only 2.2 percent in riffles. The six most abundant species in riffles were all naidids, and included *Nais pardalis* (14.7 percent), *Dero nivea* (12.7 percent), *D. obtusa* (12.4 percent), *N. variables* (8.6 percent), *Pristinella osborni* (8.5 percent), and *Chaetogaster diaphanus* (8.3 percent). All six of these species were also obtained from pools, four occurred in pools in moderate abundance (2.0 percent), but none individually comprised more than 3.2 percent of the oligochaete community in the more lentic habitat type. The combined abundance of these six species was 65.2 percent in riffles and 11.6 percent in pools.

43. As reflected in the species composition of dominant oligochaetes in pools and riffles, the pool community was a mixed assemblage of tubificids and

naidids, but the riffle community was almost entirely comprised of naidids (Table 4). Tubificids are lentic species, collected frequently in ponds and lakes, whereas naidids are found in riffles. The ratio of tubificids to naidids equaled 1.52 and 0.07 in pools and riffles, respectively. Among uncommon species, this inter-habitat difference was also notable. Ten species of naidids were obtained from riffles but not pools, whereas three species of naidids were found in pools but not riffles. Three species of tubificids were found in pools but not riffles, and only one species of tubificid (*Limnodrilus rubripennis*) was found only in riffles.

44. The particular combination of species occurring at a given site (especially on a particular date) was generally consistent with, but not identical to, the pool or riffle communities indicated by the composite data summarized in Table 5. Intersite comparisons of chironomid and oligochaete species composition were made using Jaccard's similarity index of beta diversity. Beta diversity is essentially a measure of how different sites are in terms of the variety of species found in them (Magurran 1988). Jaccard's index of beta diversity is equal to $j/(a+b-j)$; where a and b equal the number of species in sites a and b , respectively, and j equals the number of species found in both sites. Intersite similarity was low, as exemplified by comparisons among sites in the fall of 1987 (Table 5). The highest observed value was only 0.65 for chironomids at the site 2 riffle versus the site 3. These results indicate that species composition varies among pools and riffles, although the degree of dissimilarity is probably overestimated due to the low abundance of most species (Figures 9a-9d) and the uncertainty of even the presence or absence of such species based on anything less than extremely extensive sampling.

45. Community composition at the species level also varied greatly among sampling dates. For example, although *Nais pardalis* was the most abundant oligochaete in riffles based on all data combined, this species did not occur in the fall 1988, the spring 1989, or the fall 1989 samples.

46. Only by combining data for all pools and all riffles for all sampling dates are characteristic interhabitat differences evident (Table 5). For example, rheophilic (flow-loving) chironomids that dominated the combined data set for riffles (*Thienemanniella* nr. *fusca*, *Cricotopus bicinctus*, *Poly-pedilum convictum*, and *Rheotanytarsus* sp. as shown in Table 3 did not occur in the same relative abundance on each riffle on a particular date or in similar abundance in a particular riffle on different dates. In addition, although

Table 5
Community Comparisons (Jaccard's Index) for Chironomids and
Oligochaetes in Luxapalila Creek, Mississippi, 1987-89

<u>Taxonomic Group</u>	<u>Habitat</u>	<u>Comparison Between Sites</u>		
		<u>1 vs 2</u>	<u>2 vs 3</u>	<u>1 vs 3</u>
Chironomidae	Pool	0.55	0.63	0.41
	Riffle	0.59	0.65	0.61
Oligochaeta	Pool	0.31	0.23	0.43
	Riffle	0.35	0.35	0.36

naidids generally dominated the oligochaete community in riffles, species composition of this group exhibited great variation among sites and dates. In general, dynamic changes in species composition even among dominants were generally so great that they masked seasonal patterns of density change that were evident at the family level of description.

47. In June 1988, 20 specimens of the oligochaete *Piquetiella michiganensis* were found in four of five samples of the riffle at site 4. Densities were estimated at 314.3 (\pm 258.2) individuals/sq m. In a nearby pool two of five core samples yielded one specimen each with an estimated density of 31.4 (\pm 38.5) individuals per sq m. Similar sampling at two pools and two riffles in Luxapalila Creek near its confluence with the Tombigbee River yielded no *P. michiganensis*. In the fall of 1988, a single *P. michiganensis* was collected in a riffle in the upper section of the lower reach.

48. This species has been reported in north central North America as far south as Virginia (Brinkhurst 1986). It has been collected in the Great Lakes and upper Mississippi River east to the Susquehanna and Chemung Rivers in New York (Hiltunen and Klemm 1980; Klemm 1985), and south to the Wabash River in southern Indiana.* More complete macroinvertebrate surveys in the central United States would establish whether the apparent disjunct distribution of this species is the result of incomplete data or specific habitat requirements that are occasionally met at the periphery of its range.

* Personal Communication, Feb 1989, Dr. Michael S. Loden, Jefferson Parish Environmental Department, Jefferson, LA.

PART IV: DISCUSSION

Major Findings

49. In this macroinvertebrate study in Luxapalila Creek, samples were taken after exceptionally high (June and November 1989) and low (June and September 1988) water. Floods and droughts are physical factors of potentially major significance to the structure and abundance of stream invertebrate communities. Intersite and interdate variability in community structure were probably intensified by the extreme range of physical conditions that occurred during this study. This factor was mainly responsible for the high biological diversity in Luxapalila Creek. In addition, the high equitability of species within chironomid and oligochaete communities reflected effects of a wide range of extreme physical conditions that occurred between September 1987 and November 1989.

50. Especially low densities of chironomids, oligochaetes, and total invertebrates were prominent characteristics of Luxapalila Creek in June 1989, although each site did not exhibit reduced density of all three groups. It is noteworthy that the highest mean daily discharge (14,000 cfs) recorded during the course of this study occurred in January 1989. The prominence of especially low densities in June 1989 is evidence that scouring floods affected standing crops (although to different degrees for particular taxa at particular sites). By November 1989, oligochaetes at site 4 (the only site sampled in the fall of 1989) had not recovered in the pool or riffle, but chironomids in pool 4 were at the highest average density observed for this group of invertebrates in Luxapalila Creek pools (Figure 6a). Midges are notable for their ability to rapidly recolonize after decimation, with their aerial adult stage and the aquatic drift of larvae contributing to this opportunistic characteristic (e.g., Simpson and Bode 1980). Oligochaetes are of course fully aquatic; naidids can enter the drift although tubificids rarely do (Milbrink 1973). During early biological colonization of a manmade gravel bar in the Tennessee-Tombigbee Waterway, chironomids were among the first colonists while oligochaetes appeared later (Bingham and Miller 1989).

51. In addition to scouring high flows during the winter and spring of 1989, this study included a period of sustained and exceptionally low water during the summer and fall of 1988. The lowest recorded discharge (25 cfs) in the 16-year history of records was measured during the summer of 1988. The

abundance of *Paracladopelma undine*, a dominant chironomid in pools, declined greatly in the fall of 1988, from initially high densities in the fall of 1987 and spring of 1988. *Paracladopelma undine* is a member of a genus that is apparently restricted to cool water (Wierderholm 1983). It is possible that the decline of this species in the fall of 1988 may have been related to physical stress (such as reduced dissolved oxygen and increased water temperature) associated with extremely low stream discharge for a sustained period. Similarly, it is noteworthy that the highest measured ratio (5.0) of tubificid to naidid oligochaetes on any date was measured for the October 1988 pool samples. Tubificids are recognized to be tolerant of pool water quality, including low dissolved oxygen and high temperature (Brinkhurst and Cook 1974).

52. Ordinarily, cumulative species is a linear function of the *logarithm* of the cumulative number of individuals (e.g., McNaughton and Wolf 1973). The lack of a semilogarithmic relationship between cumulative species and cumulative individuals (Figure 8) during this study was primarily because even more species of chironomids and oligochaetes are likely to occur in Luxapalila Creek than were identified. Nonetheless, the extreme physical conditions of both low and high flow allowed more species per individuals identified to be accounted for than if more stable discharge conditions had prevailed.

53. The slopes of dominance-diversity plots of chironomid and oligochaete communities (Figures 9a-9d) were extraordinarily low and indicate the high equitability of species in Luxapalila Creek. Percentage abundance of species changed only two orders of magnitude in community samples of 27-63 species. Generally, a range in species abundance of 3 to 5 orders of magnitude would be associated with rich assemblages of species (McNaughton and Wolf 1973; Whittaker 1965). In comparison, species abundances of 15-30 species in samples of riverine mussel communities typically span the same range as that observed among 27-63 species of chironomids or oligochaetes in Luxapalila Creek. The high equitability among species of chironomids and oligochaetes in Luxapalila Creek pools and riffles observed during the present study may have been enhanced by the extreme range of hydraulic conditions.

Recommendations

54. Choice of sites within the project area for a post-construction macroinvertebrate study should include at least one pool-riffle sequence not directly affected and one pool-riffle sequence directly affected by the project. Pools directly affected by construction may become more depositional in nature than pools not directly affected by construction. Thus, pool-versus-riffle comparisons at locations directly affected by the project may show clearer differences in macroinvertebrate community structure than pool-versus-riffle comparisons at locations not directly affected by the project. As in the present study, characterizations of the macroinvertebrate community should focus on density and species relative abundance. Macroinvertebrates should be studied at the pool and riffles of site 4, because this site is upstream of the project area and is not likely to show direct or indirect effects of project construction.

REFERENCES

- Arner, D. H., Robinette, H. R., Frasier, J. E., and Gray, M. H. 1976. "Effects of Channelization on the Luxapalila River on Fish, Aquatic Invertebrates, Water Quality, and Furbearers," Contract Report No. 14-16-0008-739, Department of Wildlife and Fisheries, Mississippi State University.
- Bingham, C. R., and Miller, A. C. 1989. "Colonization of a Man-Made Gravel Bar by Oligochaete," Hydrobiologia, Vol 180, pp 229-234.
- Brinkhurst, R. O. 1986. "Guide to the Freshwater Aquatic Microdrile Oligochaetes of North America," Canadian Special Publication of Aquatic Sciences 84, Department of Fisheries and Oceans, Ottawa, Canada.
- Brinkhurst, R. O., and Cook, D. G. 1974. "Aquatic Earthworms (Annelida: Oligochaeta)," Pollution Ecology of Freshwater Invertebrates, C. W. Hart, Jr., and S. L. H. Fuller, eds., Academic Press, New York and London.
- Hiltunen, J. K., and Klemm, D. J. 1980. "A Guide to the Naididae (Annelida: Clitellata: Oligochaeta) of North America," EPA-6000/4-80-031, Environmental Monitoring and Support Laboratory, Cincinnati, OH.
- Hynes, H. B. N. 1970. "The Ecology of Running Waters," University of Toronto Press, Toronto, Canada.
- King, R. H., Miller, A. C., and Glover, J. E. 1982. "Proposed Riffle Construction in an Old River Channel," Journal of the Mississippi Academy of Sciences, Vol 27, pp 151-161.
- Klemm, D. J. 1985. "A Guide to the Freshwater Annelida (Polychaeta, Naidid and Tubificid Oligochaeta, and Hirudinea) of North America, Kendall/Hunt Publishing Company, Dubuque, IA.
- Magurran, A. E. 1988. "Ecological Diversity and its Measurement," Princeton University Press, Princeton, NJ.
- McNaughton, S. J., and Wolf, L. L. 1973. General Ecology, Holt, Rinehart, and Winston, Inc., New York.
- Milbrink, G. 1973. "On the Vertical Distribution of Oligochaetes in Lake Sediments," Report No. 53, Institute of Freshwater Research, Drottingholm.
- Miller, A. C., and Bingham, C. R. 1987. "A Hand-held Benthic Core Sampler," Journal of Freshwater Ecology, Vol 4, pp 77-81.
- Miller, A. C. 1987. "Habitat Development in Navigable Waterways Using Dredged Material," Proceedings of the Twenty-first Annual Dredging Seminar, John B. Herbich, ed., Texas Engineering Experiment Station, Texas A&M University, College Station, TX, pp 2-12.
- Russell-Hunter, W. D. 1970. "Aquatic Productivity: An Introduction to Some Basic Aspects of Biological Oceanography and Limnology," MacMillan Publishing Co., New York.
- Shields, F. 1983. "Design of Habitat Structures for Open Channels," Journal of Water Resources Planning Management, Vol 109, pp 331-344.
- Simpson, E. H. 1949. "Measurement of Diversity," Nature, Vol 163, p 688.
- Simpson, K. W., and Bode, R. W. 1980. "Common Larvae of Chironomidae (Diptera) from New York State Streams and Rivers," Bulletin No. 439, New York State Museum, New York.

- Standford, J. A., and Ward, J. V. 1979. "Stream Regulation in North America," The Ecology of Regulated Streams, J. V. Ward and J. A. Standford, eds., Plenum Press, New York, pp 215-236.
- Tharpe, E. J., Plunkett, M. L., Morris, F., and Oakley, W. T. 1987. "Water Resources Data--Mississippi Water Year 1987," US Geological Survey Water-Data Report MS-87-1, Jackson, MS.
- US Army Corps of Engineers. 1986. "Environmental Description of Luxapalila Creek, Mississippi and Alabama," US Army Engineer District, Mobile, AL.
- US Fish and Wildlife Service. 1987. "Supplementary Report to the Final Fish and Wildlife Coordination Act Report on Luxapalila Creek," US Fish and Wildlife Service, Daphne, AL.
- US Soil Conservation Service. 1971a. "Guidelines for Planning and Review of Channel Improvements," US Soil Conservation Service Watershed Memorandum 108.
- _____. 1971b. "Planning and Design of Open Channels," US Soil Conservation Service Technical Report 25.
- Whittaker, R. H. 1965. "Dominance and Diversity in Land Plant Communities," Science, Vol 147, pp 250-260.
- Wiederholm, T., ed. 1983. "Chironomidae of the Holarctic Region. Keys and Diagnoses; Part I. Larvae," Entomological Scandinavica, Supplement 19.
- Woods, L. C., III, and Griswold, B. L. 1981. "Channelization and Mitigation: Their Effects on Macroinvertebrate Communities in the Olentangyi River, Columbus, Ohio," Warmwater Streams Symposium of the American Fisheries Society, L. A. Krunholz, ed., Lawrence, KS, pp 113-118.

APPENDIX A
COUNTS OF MAJOR TAXA PER CORE SAMPLE, FALL 1987

Lumpkin Creek, Mississippi Sampling Date: 9/16/87 LX987C		Counts of major taxa: raw data (for those species identified to species counts are summed here, all others (*) are raw data)		P1P3		P1P4		P1P5		P1P6		P1P7		P1P8		P1P9		P1P10		P1P11		P1P12		P1P13		P1P14		P1P15		P1P16		P1P17		P1P18					
Turbellaria	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0					
Oligochaeta	14	9	7	18	6	14	59	31	8	31	8	14	16	11	27	30	18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Chironomidae	46	22	40	34	17	21	22	58	35	61	12	31	23	9	26	8	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Ceratopogonid	0	0	6	2	3	0	0	5	1	3	0	0	1	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Coleoptera	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Ephemeroptera	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Odonata	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Plecoptera	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Trichoptera	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Amphipoda	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Isopoda	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bivalvia	2	5	1	1	0	13	10	28	17	21	2	3	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Gastropoda	2	0	0	0	0	1	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mollusca	4	5	1	1	0	14	10	29	18	22	2	3	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Acarina	5	0	7	0	1	2	1	11	0	1	0	0	0	0	2	7	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Polychaeta	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Nirudinea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Epididae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Collembola	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Nematoda	20	8	17	10	3	2	2	8	4	9	3	1	19	10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Nemertes	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Aeolosomatidae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
TOTAL	172	45	77	69	30	53	98	145	66	131	25	50	64	53	87	41	28	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Luxapalila Creek, Mississippi		SITE		P117		P118		P119		P1110		P1213		P1216		P1218		P1219		P12110	
Sampling Date: 9/16/87		GENUS SPECIES		P113		P117		P118		P119		P1110		P1213		P1216		P1218		P1219	
Raw species data: LY987RAM		GENUS SPECIES		P113		P117		P118		P119		P1110		P1213		P1216		P1218		P1219	
TAXA	GENUS SPECIES	P113	P117	P118	P119	P1110	P1213	P1216	P1218	P1219	P12110										
PLATYHELMINTHES																					
TURBELLARIA																					
	Dugesia tigrina																				
	Planaria sp.							1													
	Unidentified																				
ANNELIDA																					
POLYCHAETA																					
OLIGOCHAETA																					
	Sparganophilus																				
	Haplotaxidae		1																		
	Lumbriculidae		2																		
	Naidae																				
	Amphichaeta leydigi		1																		
	Bratislavia bilongata																				
	Bratislavia unidentata																				
	Cheetogaster diaphanus																				
	Dero digitata																				
	Dero furcata				1																
	Dero nivea																				
	Dero obtusa																				
	Dero trifida		1																		
	Neis behningi																				
	Neis bretscheri																				
	Neis communis																				
	Neis pardalis		1																		
	Neis pseudobtusa																				
	Neis variabilis																				
	Pristina equiseti																				
	Pristina leidy		1																		
	Pristina syncletes																				
	Pristinella jenkinsae																				
	Pristinella longidentata																				
	Pristinella longisoma																				
	Pristinella osborni																				
	Pristinella sima																				
	Slavina appendiculata																				
	Specaria josinae		1																		
	Stevensoniana trivandrama																				
	Unidentified naididae																				
	Tubificidae																				
	Aulodrilus limobius		1		2																
	Aulodrilus piqueti		6		5																
	Branchiura sowerbyi																				
	Limnodrilus hoffmeisteri																				
	Potamothenis vejovskyi																				
	Unidentified tubificidae		2		1																
	Unidentified																				
	Helobdella elongata																				
	Unidentified																				
HIRUDINEA																					
AELOSOMATIDAE																					

TAXA	GENUS SPECIES	P113	P117	P118	P119	P1110	P1213	P1216	P1218	P1219	P12110
INSECTA											
COLEOPTERA											
	<i>Bidessus</i> sp.										
	<i>Stenelmis</i> sp.										
	<i>Malpūs</i> sp.				1			1			
	<i>Hydrochus</i> sp.										
COLLEMBOLA											
	<i>Isotomurus palustris</i>	1									
	<i>Sminthurides</i> sp.										
DIPTERA											
	CHIRONOMID										
	Chironominae										
	<i>Chironomus</i> sp.	2									
	<i>Cryptochironomus fulvus</i>	1			2		3	1		2	
	<i>Dicrotendipes nemodestus</i>	4	1		2		2			6	
	<i>Dicrotendipes nervosus</i>					1				1	
	<i>Dicrotendipes</i> sp. 1										
	<i>Endochironomus</i> sp.										
	<i>Milothauma babiyi</i>	2									
	<i>Parachironomus abortivus</i>	2									
	<i>Paracleodopelma uridine</i>	4	2	17	16	5	2	6	10	11	9
	<i>Paralauterborniella</i> sp.			1							
	<i>Polypedilum convictum</i>		1								
	<i>Polypedilum illinoense</i>	2		2	3			1		2	
	<i>Polypedilum</i> nr. <i>scaeloenum</i>		2								
	<i>Pseudochironomus</i> sp.	6						1		1	
	<i>Robackia</i> sp.										
	<i>Stenochironomus</i> sp.				1						1
	<i>Xenochironomus</i> sp.								4		
	Unidentified chironomini										
	Tanytarsini										
	<i>Tanytarsus</i> sp.	2	3			6			2		5
	<i>Cladotanytarsus</i> sp.										
	<i>Microspectra</i> sp.								2		1
	<i>Rheotanytarsus</i> sp.		1								
	<i>Stempellina</i> sp.										
	<i>Stempellinella</i> sp.										
	<i>Tanytarsus glabrescens</i>	2	3	2	3		2			2	3
	<i>Tanytarsus querlus</i>	4	1	13				2	4	5	16
	Orthocladinae										
	<i>Corynoneura celeripes</i>	2									
	<i>Cricotopus bicinctus</i>				1			3			1
	<i>Eukiefferiella</i> sp.										
	<i>Menocladus crassicornus</i>										
	<i>Menocladus distinctus</i>										
	<i>Menocladus rectinervis</i>										
	<i>Menocladus</i> sp.										
	<i>Parakiefferiella</i> sp.	6	6		2	2	4	5	8	13	5
	<i>Rheocricotopus</i> sp.										
	<i>Thienemanniella</i> nr. <i>fusca</i>										
	<i>Thienemanniella xena</i>										
	Unknown sp.										
	Tanytarsinae										
	<i>Ablabesmyia parajanta</i>	4			1						1
	<i>Ablabesmyia tarella</i>										
	<i>Labrundinia pitosella</i>										

TAXA	GENUS SPECIES	P113	P117	P116	P119	P1110	P1213	P1216	P1218	P1219	P1210
	Macropetopia sp.	2					1				
	Milotenopus sp.			1							
	Pentaneura sp.						3	1	4		2
	Procladius sp.	2		1	1		2	1	2		9
	Unidentified tenypodinae						1		6		
	Unidentified										
	Ceratopogonidae										
	Alluaudomyia sp.			4	2	3			1		
	Bezzia sp.			2					4	1	3
	Empididae										
	Hemerodromia										
	HEMEROPTERA										
	Ceania sp.										
	Ephemerella sp.										
	Tricorythodes sp.										
	Clypeum subaequalis										
	Spinedia wallace										
	Unknown sp. A (squatly bodies)										
	Unknown sp. B (slim guys)										
	Unknown sp. C										
	COGNATA										
	Argia sp.										
	Macromia sp.								1		
	Archilestes										
	Immatures		1								
	PLECOPTERA										
	Perlina ephyre										
	Immatures										
	TRICHOPTERA										
	Macronema zebratum										
	Unknown hydroptychid sp.										
	Hydroptila sp.								1		
	Oecetis sp.	1									
	Chimarra sp.									1	
	Polycentropus sp.								1		2
	Type diversa										
	Unidentified										
	AMPHIPODA										
	Synurella sp.	1									
	ISOPODA										
	Asellus sp.		3								
	ACARINA										
	MOLLUSCA										
	Corbicula fluminea	5		7		1	2	1	11		1
	Ferrissia rivularis										
	PELECYPODA	2		1	1		13	10	28	17	21
	GASTROPODA	2								1	1
	Ammicula sp.										
	Gyraulus sp.										
	Unidentified						1				
	OTHER										
	NEMATODA										
	NEHERTEA										
	Prostoma graecense										
	TOTALS	92	48	79	69	30	53	98	145	66	131
	SPECIES NUMBER	27	23	17	23	13	20	22	32	15	31

Luxapalila Creek, Mississippi

Sampling Date: 9/16/87

Raw Species data: LX987RAW

TAXA	P331	P334	P337	P338	P3310	R111	R112	R113	R114	R118	R123	R124
GENUS SPECIES												
PLATYHELMINTHES												
TURBELLARIA												
<i>Dugesia tigrina</i>						1	2		1			
<i>Planaria</i> sp.				1						1		
Unidentified							4	1				
ANNELIDA												
POLYCHAETA												
OLIGOCOAETA												
<i>Sparganophilus</i>												1
<i>Haptaxidae</i>						6	35	5		3	4	3
<i>Lumbriculidae</i>												
<i>Naididae</i>												
<i>Amphichaeta leydigi</i>												
<i>Bratistavia bilongata</i>						2						
<i>Bratistavia unidentata</i>												
<i>Chaetogaster diaphanus</i>												7
<i>Dero digitata</i>	3											
<i>Dero furcata</i>				7	2.5							
<i>Dero nivea</i>				1		1		1	4			
<i>Dero obtusa</i>		4	1		2.5					1	1	
<i>Dero trifida</i>			1									
<i>Nais behningi</i>												
<i>Nais bretscheri</i>												
<i>Nais communis</i>						2						
<i>Nais pardalis</i>		2	2	2	2	1					8	1
<i>Nais pseudobtusa</i>												14
<i>Nais variabilis</i>				2								
<i>Pristina e quiseta</i>			1								2	9
<i>Pristina leidy</i>		1		2								
<i>Pristina synclites</i>			2									
<i>Pristinella jenkiniae</i>												
<i>Pristinella longidentata</i>										1		
<i>Pristinella longisoma</i>										1		
<i>Pristinella osborni</i>			1	2	1	2			2			
<i>Pristinella sima</i>												
<i>Slavina appendiculata</i>				1								
<i>Specaria josinae</i>				1								
<i>Stevensomiana trivandana</i>												
Unidentified naididae								4				
TUBIFICIDAE												
<i>Aulodrilus limnobius</i>				1								
<i>Aulodrilus piqueti</i>	2	5										
<i>Branchiura sowerbyi</i>		2										
<i>Limnodrilus hoffmeisteri</i>	1	1										
<i>Potamothenis vejovskyi</i>												1
Unidentified tubificidae		1	2	3	6	1						
Unidentified					4							
<i>Helobdella elongata</i>				1								
Unidentified								8	6	16	2	
AEOLOSOMATIDAE												

TAXA	GENUS SPECIES													R1214
	P1311	P1314	P1317	P1318	P1310	R1111	R1112	R1113	R1114	R1116	R1118	R1213	R1214	
INSECTA														
COLEOPTERA														
	Bidessus sp.													
	Stenelmis sp.	1												
	Halipus sp.							1						
	Hydrochus sp.							2						
COLLEMBOLA														
	Isotomurus palustris							1					2	
	Sminthurides sp.							4						
DIPTERA														
CHIRONOMID														
	Chironominae													
	Chironomus sp.													
	Cryptochironomus fulvus													
	Dicrotendipes neomolestus													
	Dicrotendipes nervosus													
	Dicrotendipes sp. I													
	Endochironomus sp.													
	Miltotholma babiyi													
	Parachironomus abortivus													
	Paracladopelma undine													
	Paraleuterborniella sp.	7												
	Polypedilum convictum													
	Polypedilum filinoense													
	Polypedilum nr. scalloenum													
	Pseudochironomus sp.													
	Robackia sp.													
	Stenochironomus sp.													
	Xenochironomus sp.													
	Unidentified chironomini													
	Tanytarsini													
	Cladotanytarsus sp.													
	Microsectra sp.													
	Rheotanytarsus sp.													
	Stempellina sp.													
	Stempellinella sp.													
	Tanytarsus glabrescens													
	Tanytarsus querlus													
	Orthocleidiinae													
	Corynoneura ceteripes													
	Cricotopus bicornis													
	Eukiefferiella sp.													
	Eukiefferiella sp.													
	Manocledius crassicornus													
	Manocledius distinctus													
	Manocledius rectinervis													
	Manocledius sp.													
	Parakiefferiella sp.													
	Rheocricotopus sp.													
	Thienemannella nr. fusca													
	Thienemannella xena													
	Unknown sp.													
	Tanyptodinae													
	Ablabesmyia parajanka													
	Ablabesmyia taralla													
	Labrundinia pilosella													

TAXA	GENUS SPECIES	P1311	P1316	P1317	P1318	P13110	R1111	R1112	R1113	R1114	R1118	R1213	R1214
	Macropelopia sp.						21	4		1	5.2		
	Nilotanytus sp.												
	Pentaneura sp.												
	Procladius sp.	1											
	Unidentified tanypodinae												
	Unidentified						8	2					3.5
	Ceratopogonidae												
	Alluaudomyia sp.												
	Bezzia sp.	1											3.5
	Empididae												
	Hemerodromia												
EPHEMEROPTERA													
	Caenis sp.												
	Ephemerella sp.												
	Tricorythodes sp.											2	
	Cinygmula subaequalis												
	Spinadisa wallacei												
	Unknown sp. A (squatly bodies)							1		1			1
	Unknown sp. B (slim guys)												
	Unknown sp. C												
ODONATA													
	Argia sp.												
	Macromia sp.												
	Archilestes												
	Immatures												
PLECOPTERA													
	Perlunella ephyre												
	Immatures												
TRICHOPTERA													
	Macronema zebratum												
	Unknown hydrogrychid sp.												
	Hydroptila sp.								4	4			4
	Deceit sp.								1	1			3
	Chimarra sp.						68	35	15	5			4
	Polycentropus sp.												
	Lype diversa												
	Unidentified												
AMPHIPODA													
	Synurella sp.												2
ISOPODA													
	Asellus sp.												
ACARINA													
							2	7	38	48	39	13	17
MOLLUSCA													40
PELECYPODA													
	Corbicula fluminea	3											
GASTROPODA													
	Ferrissia rivularis												
	Amnicola sp.												
	Gyraulus sp.												
	Unidentified												
OTHER													
NEMATODA													
NEMERTEA													
	Prostoma graecense												
TOTALS		23	64	38	87	28	267.2	232	111	67	189	233	299
SPECIES NUMBER		12	20	10	31	12	26	26	20	16	23	25	25

Lunapilla Creek, Mississippi																						
Sampling Date: 9/16/87																						
Counts of major taxa: raw data																						
(for those samples identified to species, counts are summed here, all others (*) are raw data)																						
MAJOR TAXA	R111	R112	R113	R114	R118	R123	R124	R126	R128	R129	R1210	R131	R1312*	R133	R136	R135	R1316*	R1317*	R1318*	R1319*	R1310	
Turbellaria	1	6	1	1	1	0	1	1	1	4	0	12		17	0	14	4	2			7	0
Oligochaeta	15	53	6	9	15	9	37	11	17	16	16	46	72	320	76	69	27	42	32	46	37	
Chironomidae	131	77	24	6	129	111	173	50	154	109	142	15	36	51	48	122	5	12	3	4	4	
Ceratopogonid	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	1		1	
Coleoptera	1	2	1	0	0	0	0	1	1	0	0	0	0	0	0	0	1	0			0	
Ephemeroptera	0	1	1	1	0	3	2	1	11	25	4	0	13	6	10	5	0	1			0	
Odonata	0	0	0	0	1	0	0	0	1	0	0	1	1	1	3	1	1	1			2	
Plecoptera	0	0	1	0	0	0	0	0	0	0	0	0	0	2	0	2	0	0			0	
Trichoptera	68	35	20	10	11	13	13	13	15	21	20	8	3	7	11	7	3	1	1	1	1	
Amphipoda	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0	
Isopoda	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0	
Bivalvia	4	1	5	11	8	17	22	9	22	23	30	39	35	132	55	11	7	7	2	15	25	
Gastropoda	0	0	1	0	0	13	5	0	13	56	15	4	8	0	2	8	0	0			1	
MINOR GROUPS																						
Acarina	38	48	39	13	17	60	40	47	104	220	152	5	6	36	16	31	6	12	13	13	16	
Polychaeta	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0	
Hirudinea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0	
Empididae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			0	
Collembola	5	1	5	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0			0	
Nematoda	4	0	1	0	4	2	3	1	3	11	4	3	5	14	20	6	2	2	1	2	3	
Nemertea	0	0	0	0	1	5	0	0	1	0	2	1	7	9	0	4	2	4			3	
Aeolosomatidae	0	8	6	16	2	0	0	0	9	5	0	6	6	98	44	21					3	
TOTAL	267	232	111	67	189	235	299	135	353	490	385	141	185	693	285	302	53	85	54	94	96	

Luxapalila Creek, Mississippi											
Sampling Date: 9/16/87											
Raw species data: LX987RAU											
TAXA	GENUS SPECIES	R1216	R1218	R1219	R1210	R1311	R1313	R1314	R1315	R1310	
PLATHHELMINTHES											
TURBELLARIA											
	Dugesia tigrina	1		1		8	6		10		
	Planaria sp.						4				
	Unidentified		1	3		4	7		4		
ANNELIDA											
POLYCHAETA											
OLIGOCHAETA											
	Sparganophilus										
	Haplotaxidae										
	Lumbriculidae					2					
	Naididae										
	Amphichaeta leydigi			2							
	Bratislavia bilongata							4			
	Chaetogaster diaphanus	3	4	12	3			5	12.5	7	
	Dero digitata						10.7	1.9		3.3	
	Dero furcata										
	Dero nivea					18	85.4	14.9	12.5	1.7	
	Dero obtusa					12	10.7	7.5	6.7	6.7	
	Dero trifida							3.7		3.3	
	Nais behningi								1.8		
	Nais bretscheri							1	3.5		
	Nais communis					2					
	Nais pardalis	3	8		1		44.5	7	30.2		
	Nais pseudobutusa				1		26.7		1.8		
	Nais variabilis				8			2			
	Pristina aquiseta	2	1								
	Pristina leidy		4		3		44.5	8	3.5		
	Pristina synclites										
	Pristinella jenkinsae									1	
	Pristinella longidentata										
	Pristinella longisoma										
	Pristinella osborni						53.4	5		14	
	Pristinella sima							2			
	Slavina appendiculata					4	26.7	6	3.5		
	Specaria josinae										
	Stevensoniana trivandana						8.9	2			
	Unidentified naididae							5			
	Tubificidae										
	Aulodrilus limnobius										
	Autodrilus piqueti									1	
	Branchiura sowerbyi										
	Limnodrilus hoffmeisteri										
	Potamothenis vejovskyi										
	Unidentified tubificidae	2				2		1			
	Unidentified	1		2		4	8.9				
HIRUDINEA											
	Melobella elongata					1					
	Unidentified		9	5		6	97.9	44	21.3	3	
AELOSOMATIDAE											

TAXA	GENUS SPECIES	R126	R128	R129	R1210	R131	R133	R134	R135	R1310
INSECTA										
COLEOPTERA										
	<i>Bidessus</i> sp.									
	<i>Stenelmis</i> sp.									
	<i>Halipus</i> sp.	1								
	<i>Hydrochus</i> sp.								1	
COLLEMBOLA										
	<i>Isotomurus pallustris</i>									
	<i>Sminthurides</i> sp.									
DIPTERA										
CHIRONOMID										
	Chironominae									
	<i>Chironomus</i> sp.									
	<i>Cryptochironomus fulvus</i>				2.8		4.1			
	<i>Dicrotendipes neomolestus</i>	6	21.6	8.7	36.2			2	2.4	
	<i>Dicrotendipes nervosus</i>									
	<i>Dicrotendipes</i> sp. 1									
	<i>Endochironomus</i> sp.									
	<i>Nilothauma babiyi</i>				2.8					
	<i>Parachironomus abortivus</i>									
	<i>Paracladopetma undine</i>			2.2						
	<i>Paralauterborniella</i> sp.									
	<i>Polypedium convictum</i>	12	33.8	10.9	16.8	1		1	4.8	
	<i>Polypedium illinoense</i>	2	9.2							
	<i>Polypedium nr. scaloenum</i>						4.1			
	<i>Pseudochironomus</i> sp.			2.2						
	<i>Robackia</i> sp.									1
	<i>Stenochironomus</i> sp.									
	<i>Xenochironomus</i> sp.		6.2			1				
	Unidentified chironomini									
	Tanytarsini									
	<i>Cladotanytarsus</i> sp.	4	3.1		13.9			2		
	<i>Micropectra</i> sp.							1		
	<i>Rheotanytarsus</i> sp.	2	21.6	8.7	5.5			2		
	<i>Stempellina</i> sp.									
	<i>Stempellinella</i> sp.									1
	<i>Tanytarsus glabrescens</i>	2	6.2	13.1	11.1		4.1			
	<i>Tanytarsus querlus</i>		3.1				4.1			
	Orthocleidiinae									
	<i>Corynoneura ceteripes</i>									
	<i>Cricotopus bicinctus</i>	2	12.2	2.2	2.8	4		11	7.4	
	<i>Eukiefferiella</i> sp.								4.8	
	<i>Nanocleadius crassicornus</i>							1		
	<i>Nanocleadius distinctus</i>							4		
	<i>Nanocleadius rectinervis</i>							5		
	<i>Nanocleadius</i> sp.									
	<i>Parakiefferiella</i> sp.	4	6.2	26	8.4	1	28.6		7.2	2
	<i>Rheocricotopus</i> sp.									
	<i>Thienemanniella nr. fusca</i>	16	27.7	13	33.4			7	26.4	
	<i>Thienemanniella xena</i>									
	Unknown sp.									1
	Tanyptodinae									
	<i>Ablabesmyia parajanta</i>			4.4	2.8	5	4.1	10		
	<i>Ablabesmyia terella</i>							1		
	<i>Labrundinia pitoselin</i>			2.2		1				

TAXA	GENUS SPECIES	R1216	R1218	R1219	R12110	R1311	R1313	R1314	R1315	R13110
	Macropelopia sp.									
	Nilotanytus sp.		6.5		5.5					1
	Pentaneura sp.									
	Procladius sp.									
	Unidentified tenypodinae					1				
	Unidentified					1				
	Ceratopogonidae		3.1	10.9						2.4
	Alluaudomyia sp.									
	Bezzia sp.									
	Empididae									
	Hemerodromia		1							1
EPHEMEROPTERA										
	Caenis sp.									
	Ephemerella sp.		1							2
	Tricorythodes sp.		1	8						4
	Cinygmula subaequalis									4
	Spinadid wallace									4
	Unknown sp. A (squatly bodies)			3						
	Unknown sp. B (slim guys)		2	4	4		1			2
	Unknown sp. C									3
ODONATA										
	Argia sp.					1				3
	Macromia sp.									1
	Archilestes		1							
	Immatures						1			
PLECOPTERA										
	Perlunella ephyre									1
	Immatures						2			1
TRICHOPTERA										
	Macronema zebrastrum				1					
	Unknown hydropsychid sp.									
	Hydroptila sp.		2	12	6	8	1	3		3
	Oecetis sp.		2	1	9	3	1	4		3
	Chimarra sp.		9	2	6	2	7	2		4
	Polycentropus sp.									
	Lype diversa									5
	Unidentified									1
AMPHIPODA										
	Synurella sp.									
ISOPODA										
	Asellus sp.		47	104	220	152	5	36	16	31
ACARINA										
MOLLUSCA										
PELECYPODA										
	Corbicula fluminea		9	22	23	30	39	132	55	11
GASTROPODA										
	Ferussia rivularis		13	56	15	4		2		8
	Amnicola sp.									
	Gyraulus sp.									
	Unidentified									
OTHER										
HEMATODA										
HEMERTEA										
	Prostoma graecense		1	1	2	1	9			4
TOTALS		135	553	690	385	141	693.4	285	302.6	96
SPECIES NUMBER		24	32	30	28	28	32	42	31	19

APPENDIX B
SPECIES COMPOSITION IN BENTHIC SAMPLES COLLECTED IN THE SPRING OF 1988

Luxapallila Creek, Mississippi		Counts of major taxa: raw data																				
Sampling Date: 6/26-27/1968		(for those samples identified to species, counts are summed here, all others (*) are raw data)																				
LX68CNT																						
Counts of major taxa: raw data																						
(for those samples identified to species, counts are summed here, all others (*) are raw data)																						
TAXA	31E	P112*	P113*	P117*	P119*	P1110	P122	P124	P126	P128	P1210	P131*	P132*	P133*	P136*	P137*	P146	P147	P148	P149	P1510	
Turbellaria				1						1			4									1
Oligochaeta	16		7	12	18	3	9	5	18	27	10	11	14	17	21	21	17	18	15	17		
Chironomidae	34		19	27	31	23	35	41	60	63	21	27	17	22	19	25	27	30	40	49		
Ceratopogonid			3	2					4													
Coleoptera						1		1							1							
Ephemeroptera																						
Odonata	1																					
Plecoptera						1																
Trichoptera						1		2		1	2		1	2		2	2	1				
Amphipoda						1																
Isopoda	1																					
Bivalvia	2		4	1		2	4	13	7	6	2	3	2	1								
Gastropoda	1			2	1		1			1	1	4	2									2
MINOR GROUPS																						
Acarina	3		2			1	1	2	4	2					1							2
Polychaeta																						
Hirudinea							1															
Empididae																						
Collembola																						
Nematoda	5		3	6	1	5	5	3	2	4	1	4	2	4	1	2	1	1	2	4		
Nemertea																						
Aeolosomatidae																						
TOTAL	62		35	40	45	59	34	56	77	91	110	37	53	38	47	44	49	49	53	62	74	

Luxapallila Creek, MISSISSIPPI
 Sampling Date: 6/27-28/86

9a. Species data

TAXA	SITE									
	GENUS SPECIES	P1212	P1214	P1216	P1218	P1210	P1416	P1417	P1418	P1419
PLATHHELMINTHES										
TURBELLARIA										
	Dugesia tigrina					1				1
	Planaria sp.									
	Unidentified									
ANNELIDA										
POLYCHAETA										
OLIGOCHAETA										
	Sparganophilus									
	Replotaxidae									
	Lumbriculidae									1
	Naididae									
	Amphichaeta leydigi		1							
	Bratislavia bilongata									
	Bratislavia unidentata									
	Chaetogaster diaphanus			1						
	Dero digitata									
	Dero furcata									
	Dero nivea				1					5
	Dero obtusa						1			1
	Dero trifida									
	Homocheata naidina							1		
	Nais behningi									
	Nais bretscheri									
	Nais communis									
	Nais perdalis									
	Nais pseudobtusa									
	Nais variabilis									
	Piquetiella michiganensis									1
	Pristina equiseti					1				
	Pristina leioly							1		
	Pristina synclites									
	Pristinella jenkinsae							1		
	Pristinella longidentata									
	Pristinella longisoma									
	Pristinella osborni									3
	Pristinella sine									
	Stavina appendiculata									
	Specaria josinae				2					
	Stevensoniana trivandrami					3				
	Unidentified naididae									
	Tubificidae									
	Autodrilus limobius								1	
	Autodrilus piqueti				1				4	
	Autodrilus pluriseti					2			8	
	Branchiura sowerbyi									2
	Limodrilus hoffmeisteri									1
	Potamothenis vejvodskyi									2
	Unidentified tubificidae					3			1	
	Unidentified									3
										1
										2

TAXA	GENUS SPECIES	P122	P124	P126	P128	P1210	P126	P127	P128	P129
HIRUDINEA	<i>Melobella elongata</i>		1							
	Unidentified									
AEOLOSOMATIDAE										
ARTHROPODA										
INSECTA										
COLEOPTERA	<i>Bidesus</i> sp.									
	<i>Stenelmis</i> sp.									
	<i>Halipus</i> sp.	1				1				
	<i>Hydrochus</i> sp.									
COLLEMBOLA	<i>Isotomurus palustris</i>									
	<i>Sminthurides</i> sp.									
DIPTERA										
CHIRONOMID										
	Chironominae									
	<i>Chironomus</i> sp.	3	3	2	5	3	2	1	5	
	<i>Cryptochironomus fulvus</i>									
	<i>Dicrotendipes nemodestus</i>	1	2	3						
	<i>Dicrotendipes</i> sp. 1	1	1	1	4					
	<i>Endochironomus</i> sp.	1	1	1	1					
	<i>Glyptotendipes</i>									
	<i>Harnischia</i> sp.									
	<i>Mitothuma babiyi</i>									
	<i>Parachironomus abortivus</i>									
	<i>Paracladopelma undine</i>	6	8	3	7	17	3	8	8	12
	<i>Paralauterborniella</i> sp.									
	<i>Paratendipes albinus</i>									
	<i>Phaenopsectra dyair</i>									
	<i>Polypedilum convictum</i>									
	<i>Polypedilum illinoense</i>									
	<i>Polypedilum nr. scalloenum</i>	1	4	4	7	3	2	11	3	1
	<i>Pseudochironomus</i> sp.			2	1	1				
	<i>Robackia</i> sp.									
	<i>Stenochironomus</i> sp.									
	<i>Stictochironomus</i> sp.									
	<i>Xenochironomus</i> sp.									
	Unidentified chironomini									
	<i>Tanytarsini</i>									
	<i>Cladotanytarsus</i> sp.									
	<i>Microsectra</i> sp.									
	<i>Rheotanytarsus</i> sp.									
	<i>Stempellina</i> sp.									
	<i>Stempellina</i> sp.									
	<i>Tanytarsus glabrescens</i>	2	5	1	2	2				
	<i>Tanytarsus querlus</i>	2	5	11	13	3	2	8	12	7
	Orthocladinae									
	<i>Corynoneura celeripes</i>									
	<i>Corynoneura teris</i>									
	<i>Cricotopus bicornis</i>									
	<i>Eukiefferiella</i> sp.									
	<i>Manocladus crassicornis</i>									
	<i>Manocladus distinctus</i>									

TAKA	GENUS SPECIES	P1212	P1214	P1216	P1218	P1210	P1416	P1417	P1418	P1419
	Menocladus rectinervis									
	Menocladus sp.					1				
	Parakiefferiella sp.		7	9	4	7		1		
	Rheocricotopus sp.									
	Thienemanniella nr. fusca				1					
	Thienemanniella xena									
	Unknown sp.									
	Tanypodinae									
	Ablabesmyia parajanta		1		4	1				
	Ablabesmyia tarella		1		1	1				
	Labrundinia pilosella									
	Macropetopia sp.									
	Nilotanybus sp.									
	Pentaneura sp.									
	Procladius sp.	6	1	1	3	8	2	4	2	6
	Unidentified tanypodinae				2		6		2	
	Unidentified	2		2	3	3	2			2
	Ceratopogonidae									
	Alluaudomyia sp.			1		2				
	Bezzia sp.					2		1		1
	Empididae									
	Hemerodromia									
	EPHEMEROPTERA									
	Caenis sp.									
	Ephemerella sp.									
	Tricorythodes sp.									
	Cinygmula subaequalis									
	Spinadis wallace									
	Unknown sp. A (squatly bodies)									
	Unknown sp. B (slim guys)									
	Unknown sp. C									
	COONATA									
	Argia sp.									
	Macromia sp.									
	Archilestes								1	
	Immatures									
	PLECOPTERA									
	Perlina ephyre									
	Immatures	1								1
	TRICHOPTERA									
	Macronema zebratum									
	Unknown hydroptychid sp.									
	Hydroptila sp.	1		2				1		1
	Cecetis sp.									
	Chimarra sp.									1
	Polycentropus sp.									
	Type diverse									
	Unidentified					1				
	AMPHIPODA									
	Synurella sp.									
	ISOPODA									
	ASCELLUS									
	ACARINA									
		1	1	2	4	2				2

TAXA	GENUS SPECIES	P1212	P1214	P1216	P1218	P12110	P1416	P1417	P1418	P1419
MOLLUSCA										
PELECYPODA	Corbicula fluminea	2	4	13	7	6		2		1
GASTROPODA	ferrissia rivularis	1	1			1				1
	Amnicola sp.									
	Gyraulus sp.									
	Unidentified									
OTHER										
NEMATODA			5	3	2	4	2	1	1	2
NEMERTEA										
	Prostoma graecense									
TOTAL NUMBER		33	55	71	81	83	44	59	46	58
NUMBER OF SPECIES		16	20	22	25	28	20	22	17	23

Luxapalita Creek, Mississippi																									
Sampling Date: 6/26-27/1988																									
LX68CNT																									
Counts of major taxa: raw data																									
(for those samples identified to species, counts are summed here, all others (*) are raw data)																									
TAXA	R111*	R112*	R113*	R114*	R115*	R116*	R117*	R118*	R119*	R120*	R121*	R122*	R123*	R124*	R125*	R131*	R132*	R133*	R134*	R135*	R136*	R137*	R138*	R139*	R140*
Turbellaria	6	7	6	2	4	4	1	2	5	5	2	1	4	3	1	1	18								
Oligochaeta	26	24	33	17	32	30	27	21	17	15	12	20	21	17	24	21	47	20	15						
Chironomidae	54	46	57	40	103	108	117	94	86	49	52	55	50	57	47	49	43	51							
Ceratopogonid																									
Collembola	1																								
Ephemeroptera		2								4						3									
Odonata					1					1															
Plecoptera																									
Trichoptera	10	6	2	7	3	20	9	12	13	6	5	5	2	3	13	5	8	4							
Amphipoda																									
Isopoda																									
Bivalvia	21	7	16	21	12	2	4	2	11	8	31	14	19	28	11	4	7	2	5						
Gastropoda	2	2	3																						
MINOR GROUPS																									
Acarina	11	9	5	14	8	16	11	20	14	17	1	1	5	2	7	6	8	14	10	11					
Polychaeta																									
Hirudines																									
Empididae																									
Collembola																									
Nematoda	1	2	2					2	1	1	1	1	1	1	1	1	1	2	1	1					
Nemertea	2	1								2	2														
Acolosomatidae																									
TOTAL	129	102	116	141	83	178	167	182	157	146	109	90	107	110	101	97	101	101	97	101	97	101	97	101	97

Luxapalila Creek, Mississippi
 Sampling Date: 6/27-28/88

Raw species data

TAXA	GENUS SPECIES	P1410	R1211	R1212	R1213	R1214	R1215	R1216	R1217	R1218	R1219	R1220
PLATYHELMINTHES												
TURBELLARIA												
	Dugesia tigrina		4	2	1	2	1					
	Planaria sp.			2								2
	Unidentified											
ANNELIDA												
POLYCHAETA												
OLIGOCHAETA												
	Sparganophilus											
	Kaplotaxidae											
	Lumbriculidae		5									1
	Maillidae											
	Amphichaeta leydigi											
	Bratiavia bilongata											
	Bratiavia unidentata											
	Chaetogaster diaphanus										1	
	Dero digitata		2								4	
	Dero furcata											
	Dero nivea		1	7	6		3					
	Dero obtusa			4		25	1	9				
	Dero trifida			1								1
	Homochaeta naidina											
	Maie behningi											
	Maie bretscheri											
	Maie communis											
	Maie pardalis		2	1	3	1	3	1				1
	Maie pseudobutusa		1									1
	Maie variabilis		4	1	1	5	1	4				4
	Piquetiella michiganensis		1								1	9
	Pristina equisea					3					3	1
	Pristina leioly			1	2	2	3					1
	Pristina syncyites		1									
	Pristinella jenkiniae											
	Pristinella longidentata											
	Pristinella longisoma					1						
	Pristinella osborni		1	2	1	1	1					1
	Pristinella sime											
	Stevina appendiculata		3		1							
	Specaria josineae											
	Stevensoniensia trivandrama											
	Unidentified naididae			2								
	Tubificidae											
	Autodrilus limobius											
	Autodrilus piqueti											1
	Autodrilus pluriseti											
	Branchiura sowerbyi											
	Limnodrilus hoffmeisteri				3		2					
	Potamothenis vejovskyyi											
	Unidentified tubificidae		3			3						1
	Unidentified											

TAXA	GENUS SPECIES	R1410	R1211	R1212	R1213	R1214	R1215	R1415	R1417	R1418	R1419	R1410
MIRIDIINA	<i>Helobdella elongata</i>				1							
	Unidentified											
AELOSOMATIDAE			10		3	1	2	4				
ARTHROPODA												
INSECTA												
COLEOPTERA												
	<i>Bidessus</i> sp.											
	<i>Stenelmis</i> sp.			1					1			
	<i>Halipus</i> sp.											
	<i>Hydrochus</i> sp.											
COLLEMBOLA												
	<i>Isotomurus palustris</i>											2
	<i>Sminthurides</i> sp.											
DIPTERA												
	CHIRONOMID											
	Chironominae											
	<i>Chironomus</i> sp.	1	1									
	<i>Cryptochironomus fulvus</i>											
	<i>Dicoretendipes neomodestus</i>		11	11	15	7	7		2	2		
	<i>Dicoretendipes nervosus</i>											
	<i>Dicoretendipes</i> sp. 1	2	6	2	2	4	1					
	<i>Endochironomus</i> sp.	4				1						
	<i>Glyptotendipes</i>											
	<i>Harnischia</i> sp.											
	<i>Milothauma babiyi</i>											2
	<i>Parachironomus abortivus</i>	1										
	<i>Paracledopelma undine</i>	8		1	1							
	<i>Paralauterborniella</i> sp.											
	<i>Paratendipes albimanus</i>	1										
	<i>Phaenopsectra dyair</i>	1										
	<i>Polypedium convictum</i>	2	2	2	1	3	2	24	6	6	6	4
	<i>Polypedium illinoense</i>	1										
	<i>Polypedium nr. scaloenum</i>	6							2	6	2	
	<i>Pseudochironomus</i> sp.					1	1					
	<i>Robackia</i> sp.							10		1	2	14
	<i>Stenochironomus</i> sp.											
	<i>Stictochironomus</i> sp.					1						
	<i>Xenochironomus</i> sp.	1										
	Unidentified chironomini											
	Tenytarsini											
	<i>Cladotanytarsus</i> sp.	1				3			6	4	4	18
	<i>Micropectra</i> sp.											
	<i>Rheotanytarsus</i> sp.	2			1		2	6			6	8
	<i>Stempellina</i> sp.											
	<i>Stempellinella</i> sp.											
	<i>Tanytarsus glabrescens</i>	1	2	7	3	2	5	4	12	2		
	<i>Tanytarsus querlus</i>	10	2			1	1		10	10	2	
	Orthoclelineae											
	<i>Corynoneura celeripes</i>											
	<i>Corynoneura taris</i>											
	<i>Cricotopus bicornis</i>	1		6	23	12	7	4	4	4	2	
	<i>Eukiefferiella</i> sp.											
	<i>Manocleadius crassicornis</i>	1										
	<i>Manocleadius distinctus</i>	1			1	1	1					

TAXA	GENUS SPECIES	P1610	R121	R122	R123	R124	R125	R145	R147	R148	R149	R1410
	<i>Nanocladus rectinervis</i>											
	<i>Nanocladus</i> sp.											
	<i>Parakiefferiella</i> sp.		13	10	3	1						
	<i>Rheocricotopus</i> sp.		1							2	2	
	<i>Thienemannella</i> nr. <i>fusca</i>		38	64	57	51	36					2
	<i>Thienemannella</i> <i>xena</i>											
	Unknown sp.		1								2	
	Tanypodinae											
	<i>Ablabesmyia parajanta</i>		1									
	<i>Ablabesmyia tarella</i>		1		1							
	<i>Lebrundinia pilosella</i>									4		
	<i>Macropalopia</i> sp.								4	2		
	<i>Milotanypus</i> sp.											
	<i>Pentaneura</i> sp.											
	<i>Procladius</i> sp.		8		1							
	Unidentified tanypodinae		4			2	1					2
	Unidentified		2	3	1		2			4	6	
	Ceratopogonidae											
	<i>Alluaudomyia</i> sp.											
	<i>Bezzia</i> sp.											
	Epididae											
	Hemerodromia											
	EPHEMEROPTERA											
	<i>Caenis</i> sp.											
	<i>Ephemerella</i> sp.											
	<i>Tricorythodes</i> sp.											
	<i>Cinygmula subaequalis</i>											
	<i>Spinradis wellace</i>											
	Unknown sp. A (squatly bodies)											
	Unknown sp. B (slim guys)											
	Unknown sp. C											
	ODONATA											
	<i>Argia</i> sp.							1				
	<i>Macromia</i> sp.											
	<i>Archilestes</i>									1		
	Immatures		1									
	PLECOPTERA											
	<i>Perlina</i> ephyre											
	Immatures							14		8	2	2
	TRICHOPTERA											
	<i>Macronema zebretum</i>											
	Unknown hydroptychid sp.											
	<i>Hydroptila</i> sp.		1	6	8	7	10	2		2	6	1
	<i>Oecetis</i> sp.		3	1		1	1		13	3		3
	<i>Chimarra</i> sp.		5	2	4	5	2	1			2	
	<i>Polycentropus</i> sp.											
	Type diverse											
	Unidentified											
	AMPHIPODA											
	<i>Synurella</i> sp.											
	ISOPODA											
	<i>Asellus</i> sp.											
	ACARINA		1	16	20	14	17	6	8	14	10	11

TAXA	GENUS SPECIES	P1410	R1211	R1212	R1213	R1214	R1215	R1216	R1217	R1218	R1219	R1220
MOLLUSCA												
PELECYPODA	Corbicula fluminea		2	4	2	11	8	4	4	7	2	5
GASTROPODA	Ferrissia rivularis	2										
	Amiticola sp.											
	Gyraulus sp.											
	Unidentified											
OTHER												
NEMATODA		4		2		1	1		1	2	1	1
NEMENEA	Prostoma graecense			1					1			
TOTAL NUMBER		71	158	153	168	179	121	104	81	87	75	85
NUMBER OF SPECIES		24	36	25	28	32	26	21	21	25	25	25

APPENDIX C
COUNTS OF MAJOR TAXA PER CORE SAMPLE, FALL, 1988

Lupatilla Creek, Mississippi		Counts of major taxa: raw data																				
Sampling Date: 10/1988		(for those samples identified to species, counts are summed here, all others (*) are raw data)																				
LX1088C																						
Counts of major taxa: raw data																						
(for those samples identified to species, counts are summed here, all others (*) are raw data)																						
SITE																						
TAXA	P/1/1	P/1/2	P/1/6	P/1/7	P/1/9	P/2/1	P/2/2	P/2/4	P/2/5	P/2/7	P/3/1	P/3/2	P/3/3	P/3/4	P/3/9	P/4/2	P/4/3	P/4/5	P/4/6	P/4/9		
Turbellaria	2	2			3	2	3					2									2	
Oligochaeta	15	12	17	10	12	11	11	10	8	8	32	19	17	27	27	5	12	10	8	12		
Chironomidae	31	25	27	32	19	34	21	30	28	23	39	36	30	31	33	18	14	36	10	0		
Ceratopogonid							2		3					4	5							
Coleoptera		1						2			1											
Ephemeroptera									2		2											
Odonata		1					1								3						2	
Plecoptera				1																		2
Trichoptera	3		2	1	3			3	2	4	5	6	2	7	5	5	2	10	6		3	
Amphipoda																						
Isopoda																						
Bivalvia	2	3	4	2	5	2	2	2	3	3	7	14	8	7	11							4
Gastropoda			1								1	1										
MINOR GROUPS																						
Acarina	1		2	3	6	1	1	4	5	3	2	1	1	1	1	1	3	5	5			
Polychaeta																						
Hirudinea							1															
Epididae										1												
Collembola					2																	
Nematoda	3		3		1		2	2	3													
Nemertea							1	1														
Aeolosomatidae																						
Tanyderidae							1															
TOTAL	57	44	56	50	50	51	43	50	53	44	90	80	59	80	83	43	42	71	33	24		

TAXA		SITE										TOTAL		
GENUS SPECIES		P121	P122	P124	P125	P127	FREQ	P142	P144	P145	P146	P149	FREQ	TOTAL
		P/Z										P/Z		
FLATHELMINTHES														
TURBELLARIA														
	Dugesia tigrina	2	3				2		1			2	2	4
	Planaria sp.													
	Unidentified													
ANNELIDA														
POLYCHAETA														
OLIGOCOAETA	Sparganophilus													
	Heptotaxidae													
	Lumbriculidae		1				1		2	3			2	3
	Naididae													
	Amphichaeta leydigi			1	1		2							2
	Bratislavia bilongata													
	Bratislavia unidentata													
	Cheetogaster diaphanus													
	Dero digitata													
	Dero furcata													
	Dero nivea													
	Dero obtusa													
	Dero trifida													
	Dero sp.		2		2	1	3							3
	Homochaeta naidina													
	Naie behningi													
	Naie bretscheri													
	Naie communis													
	Naie pardalis													
	Naie pseudobtusa													
	Naie simplex													
	Naie variabilis													
	Piquetiella michiganensis											3	1	1
	Pristina aquiseta													
	Pristina leidy											1		
	Pristina synclytes													
	Pristina sp.													
	Pristinella jenkinsae													
	Pristinella longidentata													
	Pristinella longisoma													
	Pristinella osborni													
	Pristinella sima													
	Slavina appendiculata													
	Specaria josinae			1			1							1
	Stevensoniana trivandana													
	Unidentified naididae													
	Tubificidae													
	Autodrilus limnobius		1				1							1
	Autodrilus piqueti													
	Autodrilus pluriseti	4			2	4	3		2	3		4	4	7
	Branchiura sowerbyi													
	Limnodrilus hoffmeisteri	3		4	2	2	4							4
	Limnodrilus rubripennis	1	1			1	3		1	1			3	6
	Potamothenis vejovskyi													

TAXA	GENUS SPECIES	P121	P122	P126	P125	P127	AVG	P142	P144	P145	P146	P149	AVG	TOTAL
	Unidentified tubificidae	3	6	4			3		8	3	4	4	4	7
	Enchytraeidae													
	Barbdrillus paucisetus				1		1							1
	Unidentified													1
MIRUDIINEA	Melobdella elongata		1				1			1			1	1
	Actinobdella sp.													1
	Unidentified	1					1							1
AELOSOMATIDAE														
ARTHROPODA														
INSECTA														
COLEOPTERA	Bidessus sp.													
	Sternelmis sp.													
	Melipus sp.			2			1		1				1	2
	Hydrochus sp.													
COLLEMBOLA	Isotomurus palustris							2						1
	Sminthurides sp.													
	Unidentified													
DIPTERA														
CHIRONOMID														
	Chironominae													
	Chironomus sp.													
	Cladopelma	2					1							1
	Cryptochironomus fulvus													
	Cryptochironomus sp.		1				2	1	1				2	4
	Dicrotendipes neodestus			1			1							1
	Dicrotendipes nervosus Type 1													
	Dicrotendipes sp. 1													
	Endochironomus sp.													
	Glyptotendipes													
	Glyptotendipes		1				2						1	1
	Harnischia sp.													
	Milothauma babiyi	1					2	2		1			2	4
	Parachironomus abortivus													
	Paracadopelma undine													
	Paralauterborniella nigrohalt	2	1	5	2	1	5	1	1	1			2	7
	Paratendipes albinus													
	Paratendipes nudisquama													
	Phaenopsectra dyair													
	Phaenopsectra flavipes													
	Polypedium convictum			2			1						1	1
	Polypedium illinoense	1					1							2
	Polypedium nr. scalorum	7	5	9	1	2	5	1	2	1	3	4	4	9
	Pseudochironomus sp.	1	2		2	2	4			1	1		2	6
	Robeckia sp.													
	Stenochironomus sp.													
	Stictochironomus sp.													
	Xenochironomus sp.													
	Unidentified chironomini													
	Tanytarsini													
	Cladotanytarsus sp.			1		1	2							2
	Micropectra sp.													
	Rheotanytarsus sp.			1	2		2			10			1	3
	Stempellina sp.	1					1							1

TAXA	P1211	P1212	P1214	P1215	P1217	AVG	P1412	P1414	P1415	P1416	P1419	AVG	TOTAL
GENUS SPECIES													
Stempellinella sp													
Tanytarsus coffmani													
Tanytarsus glabrescens	2			6	5	3							3
Tanytarsus querlus	6			2		2			2				3
Unidentified tanytarsini													
Orthocladiinae													
Corynoneura celeripes													
Corynoneura taris													
Corynoneura sp													
Cricotopus bicinctus	2		2	3		3	1						4
Cricotopus sp		1											1
Eukiefferiella sp.	3					1							1
Manocladus crassicornus													
Manocladus distinctus			1			1			3				2
Manocladus rectinervis													
Manocladus minimus	1					1							1
Manocladus sp.				3	2	2	2	1					6
Parakiefferiella sp.				1		1							1
Rheocricotopus sp.													
Thienemannimyia										1			1
Thienemannielle nr. fusca				1		1							1
Thienemannielle xena									1				1
Unidentified orthocladiinae									2				1
Tanypodinae													
Ablebsmyia mallochi				1		1							1
Ablebsmyia parajanta	5	3	1	7	4	4			7	1			7
Ablebsmyia terella													
Clinotanytus										1			1
Labrundinia pilosella					1	1							1
Macropelopia sp.													
Mataria sp													
Milotanytus sp.													
Pentaneura sp.													
Procladius sp.	4	5	2	2	1	5	3	9	1	3			9
Unidentified tanypodinae	2					1	2						2
Diaesinae													
Potthesia	1					1							1
Unidentified Chironomidae	1			1		2	1			1			4
Ceratopogonidae													
Alluaudomyia sp.										1			2
Bezzia sp.													
Unidentified ceratopogonid													
Empididae													
Memorodromia						1							1
Tanyderidae	1					1							1
Simuliidae													
Simulium sp.													
EPHEMEROPTERA													
Baetisca sp													
Caenis sp.													
Ephemerella sp.													
Tricorythodes sp.													
Cinygmula subaequalis													
Stenonema sp.													

TAXA	GENUS SPECIES	P/2/1	P/2/2	P/2/4	P/2/5	P/2/7	AVG	P/4/2	P/4/4	P/4/5	P/4/6	P/4/9	AVG	TOTAL
	Spinedis wallace				2						2	2		3
	Unknown sp. A (squatly bodies)						1							
	Unknown sp. B (slim guys)													
	Unknown sp. C													
	Unidentified							1						1
ODONATA														
	Argia sp.													
	Macromia sp.													
	Archilestes													
	Dromogomphus sp.		1				1							1
	Immatures													
PLECOPTERA														
	Perlinella ephyre													
	Immatures													
	Unidentified									1				1
TRICHOPTERA														
	Ceraclea													
	Macronema zebratum													
	Hydroptila sp.					1	1	1	2	2	1	1	5	6
	Oecetis sp.			1			1				2		1	2
	Chimarra sp.			2	2	3	3	4	7	3	2		4	7
	Polycentropus sp.									1			1	1
	Type diverse													
	Unidentified													
AMPHIPODA														
	Synurella sp.													
ISOPODA														
	Asellus sp.													
ACARINA														
MOLLUSCA														
	Asellus sp.	1		1	4	5	4	1	3	5	5		4	8
PELECYPODA														
	Corbicula fluminea	2		2	3	3	4	1	2	3	1	4	4	8
GASTROPODA														
	Ferrissia rivularis													
	Ammicola sp.													
	Syraulus sp.													
	Unidentified													
OTHER														
	Unidentified													
HEMATODA														
	Unidentified													
MEMERTEA														
	Unidentified	1	2	2	3		3	5			1	1	3	6
	Unidentified		1	1			2	1					1	3

Lukopatia Creek, Mississippi												TOTAL		
Sampling Date: 10/88												FREQ		
LX10889												FREQ		
TAXA	GENUS SPECIES	R121	R122	R123	R125	R129	R/2	R144	R145	R148	R149	R150	FREQ	FREQ
PLATYHELMINTHES														
TURBELLARIA														
	<i>Dugesia tigrina</i>	3		3	5	1	4				1		1	5
	<i>Planaria</i> sp.													
	Unidentified													
ANNELIDA														
POLYCHAETA														
OLIGochaETA	<i>Sparganophilus</i>													
	Heptotaxidae					2	3	6	6	1	20	4	5	6
	Lumbriculidae	4	3											
	Naididae													
	<i>Amphichaeta leydigi</i>													
	<i>Bratislavia bilongata</i>													
	<i>Bratislavia unidentata</i>													
	<i>Chaetogaster diaphanus</i>											1	1	1
	<i>Dero digitata</i>													
	<i>Dero furcata</i>													
	<i>Dero nivea</i>	5	2	1	3	3	5							5
	<i>Dero obtusa</i>		2	1	8	3	4							4
	<i>Dero trifida</i>		2				1				3		1	1
	<i>Dero</i> sp.													
	<i>Homochaeta naidina</i>													
	<i>Nais behningi</i>													
	<i>Nais bretscheri</i>													
	<i>Nais communis</i>													
	<i>Nais pardalis</i>	3	4	1	1	2	5	1		1			2	7
	<i>Nais pseudobtusa</i>	1					1			1			1	2
	<i>Nais simplex</i>	2					1			1			1	2
	<i>Nais variabilis</i>	5	1			8	3						3	3
	<i>Piquetiella michiganensis</i>									9			1	1
	<i>Pristina equiseti</i>									1			2	2
	<i>Pristina leidy</i>													
	<i>Pristina synclites</i>													
	<i>Pristina</i> sp.												1	1
	<i>Pristinella jenkinsae</i>													
	<i>Pristinella longidentata</i>													
	<i>Pristinella longisoma</i>													
	<i>Pristinella osborni</i>	1	1				2				1		1	3
	<i>Pristinella sima</i>													
	<i>Stelvina appendiculata</i>													
	<i>Specaria josinae</i>		1				1						1	1
	<i>Stevensoniana trivandiana</i>										1		1	1
	Unidentified naididae	1					1						1	1
	Tubificidae													
	<i>Aulodrilus limnobius</i>													
	<i>Aulodrilus piqueti</i>												1	1
	<i>Aulodrilus pluriseti</i>													
	<i>Branchiura sowerbyi</i>		1	1			2							2
	<i>Limnodrilus hoffmeisteri</i>													
	<i>Limnodrilus rubripennis</i>	1					1						1	1
	<i>Potamothenis vejdovalkyi</i>													

TAXA	GENUS SPECIES										AVG	R1410	R1418	R1419	R1420	AVG	TOTAL	
	Unidentified tubificidae																	
	Barbdrilus paucisetus																	
	Unidentified																	
HIRUDINEA	Helobdella elongata																	
	Actinobdella sp.																	
	Unidentified																	
AELOSOMATIDAE																		
ARTHROPODA																		
INSECTA																		
COLEOPTERA																		
	Bidessus sp.																	
	Stenelmis sp.																	
	Halipus sp.																	
	Hydrochus sp.																	
COLLEMBOLA																		
	Isotomurus palustris																	
	Sminthurides sp.																	
	Unidentified																	
DIPTERA																		
CHIRONOMID																		
	Chironominae																	
	Chironomus sp.																	
	Cladopelma																	
	Cryptochironomus fulvus																	
	Cryptochironomus sp.																	
	Dicrotendipes neomodestus																	
	Dicrotendipes nervosus Type 1																	
	Dicrotendipes sp. 1																	
	Endochironomus sp.																	
	Glyptotendipes																	
	Harnischia sp.																	
	Miltothuma babiyi																	
	Parachironomus abortivus																	
	Paracledopelma undine																	
	Paratuberborniella nigrohalteralis																	
	Paratendipes albimanus																	
	Paratendipes nudisquama																	
	Phaenopspectra dyar																	
	Phaenopspectra flavipes																	
	Polypedilum convictum																	
	Polypedilum illinoense																	
	Polypedilum nr. scatoenum																	
	Pseudochironomus sp.																	
	Robackia sp.																	
	Stenochironomus sp.																	
	Stictochironomus sp.																	
	Xenochironomus sp.																	
	Unidentified chironomini																	
	Tanytarsini																	
	Cledotanytarsus sp.																	
	Micropectra sp.																	
	Rheotanytarsus sp.																	
	Stempellina sp.																	

TAXA	GENUS SPECIES	R1211	R1212	R1213	R1215	R1219	AVG	R1414	R1415	R1418	R1419	R14110	AVG	TOTAL
	Stempellinella sp													
	Tanytarsus coffmani							2						1
	Tanytarsus glabrescens	7.92		3.67	3.51	19.76	4	1	3					2
	Tanytarsus querlus	3.96					1		13					6
	Unidentified tanytarsini				3.51		1		1					2
	Orthocladinae													
	Corynoneura celeripes													
	Corynoneura taris	3.96					1							1
	Corynoneura							1	1					2
	Cricotopus binctus	19.8	75.4	58.66	17.55	23.06	5			15				2
	Cricotopus								2					6
	Eukiefferiella sp.								2					1
	Manocladus crassicornus	7.92					1							1
	Manocladus distinctus													
	Manocladus rectinervis		5.2		3.29		2							2
	Manocladus minimus													
	Manocladus sp.													
	Parakiefferiella sp.													
	Rheocricotopus sp.													
	Thienemanniella								1					1
	Thienemanniella nr. fusca	63.36	13	60.67	87.75	85.65	5		4					6
	Thienemanniella zena	23.76					1							1
	Unidentified orthocladinae		5.2	7.33	7.02		3							3
	Tanypodinae													
	Ablabesmyia mallochi		2.6				1							2
	Ablabesmyia parajanta	3.96	5.2	3.67			3							4
	Ablabesmyia tarella													
	Clinotanypus													
	Labrundinia pilosella								2					2
	Macropelopia sp.													
	Nataria sp							1						1
	Nilotanypus sp.													
	Pentaneura sp.													
	Procladius sp.								1					2
	Unidentified tanypodinae								1					1
	Potthasia													
	Unidentified	3.96	13	7.33	21.06	9.88	5	3	9	1	1	4		9
	Ceratopogonidae													
	Alluaudomyia sp.													
	Bezzia sp.													
	Unidentified								4					1
	Empididae													
	Memorodromia													
	Tanyderidae								2	1				2
	Simuliidae													
	Simulium sp.									2				1
	Beetisca sp													
	Caenis sp.													
	Ephemerella sp.													
	Tricorythodes sp.	1	1				2		3					3
	Cinygmula subaequalis													
	Stenomema sp.		1	1			2		7		4			4

TAXA	GENUS SPECIES	R/21	R/22	R/23	R/25	R/29	AVG	R/44	R/45	R/48	R/49	R/50	AVG	TOTAL
	Spinedis wallace													
	Unknown sp. A (squatly bodies)													
	Unknown sp. B (slim guys)													
	Unknown sp. C													
	Unidentified													
OOONATA														
	Argia sp.													
	Macromia sp.													
	Archilestes													
	Dromogomphus sp.													
	Immatures													
PLECOPTERA														
	Perlunella ephyre													
	Immatures													
	Unidentified	1				2	2	2	2	3	1		3	5
TRICHOPTERA														
	Ceraclea sp													1
	Macronema zebraatum													
	Hydroptila sp.	6		6	4	16	4	4	4	4			1	5
	Deetis sp.	1		4	2	1	4	4	4				1	5
	Chimarra sp.			1	1		2	2	2	4	4		3	5
	Polycentropus sp.													
	Type diverse													
	Unidentified													
AMPHIPODA														
	Synurella sp.													
ISOPODA														
	Asellus sp.													
ACARINA		30	33	24	27	28	5	2	9	12	10	7	5	10
MOLLUSCA														
PELECYPODA	Corbicula fluminea	9	12	17	10	5	5	6	3	4	4	4	4	9
GASTROPODA	Ferrissia rivularis													
	Amnicola sp.													
	Gyraulus sp.													
	Unidentified							2						1
OTHER														
NEMATODA		1	8		1		3	7	2			2	4	7
NEMERTEA		4	1	2	2		4	3	1		1		3	7

APPENDIX D
SPECIES COMPOSITION IN SPRING, 1989 SAMPLES

Luxapalila Creek, Mississippi

Sampling Date: 6/1989
LX689cnt

Counts of major taxa: raw data
(for those samples identified to species, counts are summed here, all others are raw data)

TAXA	P/1/2	P/1/3	P/1/6	P/1/8	P/1/9	P/2/3	P/2/4	P/2/5	P/2/6	P/2/7	P/3/3	P/3/6	P/3/7	P/3/9	P/3/10	P/4/3	P/4/5	P/4/7	P/4/8	P/4/10
Turbellaria	1			1			3	2	1			1								
Oligochaeta	19	11	14	20	32		1	3	1	6	17	13	15	11	16	9	0	0	1	1
Chironomidae	20	30	27	25	35		32	34	10	60	32	16	22	29	37	35	42	38	47	43
Ceratopogonid					2															
Coleoptera					1					1						1				2
Ephemeroptera																				1
Odonata														3	5					1
Plecoptera											1				2					2
Trichoptera					1		1		3						1					2
Amphipoda								2	1					1						
Isopoda																				
Bivalvia			2	4	2		1	2	3	3										
Gastropoda			1							2										
MINOR GROUPS																				
Acarina							1				3		4		1		1	1	2	4
Heteroptera																				
Polychaeta																				
Mirudiina																				
Empididae																				
Collembola		3		1																1
Nematoda	4	7	5	5	2			3	1	6	1					11	6	2	9	2
Nemertea																				
Simuliidae																				
Aeolosomatidae																				
Tanyderidae																				
TOTAL	45	51	50	56	75	36	51	45	21	76	57	30	41	45	63	56	31	44	62	55

TAXA		SITE										
		P123	P124	P125	P126	P127	P133	P135	P137	P138	P1410	
GENUS SPECIES												
FLATHELMINTHES												
TURBELLARIA												
	Dugesia tigrina		3	2	1							
	Planaria sp.											
	Unidentified											
ANNELIDA												
POLYCHAETA												
OLIGOCHAETA												
	Sparganophilus											
	Haptotaxidae											
	Lumbriculidae			1								
	Naididae											
	Amphichaeta leydigi											
	Bratislavia bilongata											
	Bratislavia unidentata											
	Chaetogaster diaphanus											
	Dero digitata											
	Dero furcata											
	Dero nivea											
	Dero obtusa											
	Dero trifida											
	Dero sp.											
	Homochaeta naidina											
	Neis behningi											
	Neis bretscheri											
	Neis communis											
	Neis elinguis											
	Neis pardalis											
	Neis pseudobtusa											
	Neis simplex											
	Neis variabilis											
	Piquetiella michiganensis											
	Pristina aquiseta											
	Pristina leidy					1						
	Pristina synclites											
	Pristina sp.											
	Pristinella jenkiniae											
	Pristinella longidentata											
	Pristinella longisoma											
	Pristinella osborni											
	Pristinella sine											
	Stevina appendiculata											
	Specaria josinae											
	Stevensoniene trivandrama		1	1								
	Unidentified naididae											
	Tubificidae											
	Aulodrilus limobius											
	Aulodrilus piqueti											
	Aulodrilus pluriset											
	Branchiura sowerbyi											
	Limnodrilus hoffmeisteri											
	Limnodrilus rubripennis											

TAXA	GENUS SPECIES	P1213	P1214	P1215	P1216	P1217	P1413	P1415	P1417	P1418	P1410
	Potamothis vejovskyi			1							
	Unidentified tubificidae					2					
	Enchytraididae					1				1	
	Barbidrilus paucisetus						8				
	Unidentified oligochaeta						1				
HIRUDINEA	Helobdella elongata										
	Actinobdella sp.										
	Unidentified										
AEOLOSOMATIDAE											
ARTHROPODA											
INSECTA											
COLEOPTERA											
	Bidessus sp.										
	Cyphon										
	Malipus sp.			1							1
	Hydrochus sp.										
	Stenelmis sp.										
COLLEMBOLA											
	Isotomurus palustris										
	Sminthurides sp.										
	Unidentified										
DIPTERA											
CHIRONOMID											
	Chironominae										
	Chironomus sp.		1								1
	Cladopelma										
	Cryptochironomus fulvus					1	2				
	Cryptochironomus sp.										
	Dicrotendipes neomodestus		1								1
	Dicrotendipes nervosus Type I		3				5	9	8	4	10
	Dicrotendipes nervosus Type II										
	Endochironomus sp.										
	Glyptotendipes		2								
	Harnischia sp.			1							1
	Microtendipes										2
	Miltothema bebiyi										
	Parachironomus abortivus					1					
	Paracledopelma undine										
	Paraleuterbornielle nigrohalteralis		5	1	2	1	5	2	7	5	5
	Paratendipes albinus										
	Paratendipes nudisquama										
	Phaenopsectra dyair									3	1
	Phaenopsectra flavipes									1	
	Polypedilum convictum										
	Polypedilum fallax		1								
	Polypedilum illinoense					1	1				2
	Polypedilum nr. scalloenum		5	4	6	2	8	6	2	10	2
	Pseudochironomus sp.										
	Robackia sp.										
	Stenochironomus sp.										
	Stictochironomus sp.				1						
	Tribelos sp.		12	3							1
	Xenochironomus sp.										
	Unidentified chironomini										2

TAXA	GENUS SPECIES	P1213	P1214	P1215	P1216	P1217	P1413	P1415	P1417	P1418	P1410
	Tanytarsini										
	Cladotanytarsus sp.	1				1					
	Leuterborniella		2								
	Micropectra sp.		5								
	Rheotanytarsus sp.		3							1	
	Stempellinella sp.										
	Tanytarsus coffmani										
	Tanytarsus glabrescens		1								
	Tanytarsus querlus		10	11		24	7	9	10	11	8
	Unidentified tanytarsini	5									
	Orthocladinae										
	Brillia										
	Corynoneura celeripes										1
	Corynoneura taris										1
	Corynoneura sp.						1				
	Cricotopus bicinctus										
	Cricotopus trifascia										
	Cricotopus sp.										1
	Eukiefferiella sp.										1
	Nanocladus crassicornus										
	Nanocladus distinctus										2
	Nanocladus rectinervis										
	Nanocladus minimus										1
	Nanocladus sp.			1							
	Parakiefferiella sp.		2								
	Rheocricotopus sp.										
	Thienemannella nr. fusca										
	Thienemannella xena								1		
	Unidentified orthocladinae										
	Tanypodinae										
	Ablabesmyia mallochi								1	1	1
	Ablabesmyia parajanta				1		1	2	3	1	4
	Ablabesmyia tarella										
	Clinotanypus										
	Lebrundinia pilosella								1		
	Macropelopia sp.										
	Metarsia sp.										2
	Nilotanypus sp.										
	Pentaneura sp.										
	Procladius sp.			3	1	1				1	
	Thienemannimyia										
	Unidentified tanypodinae								2		2
	Diamesinae										
	Potthastia										
	Unidentified Chironomidae		6	6	1	6	2	1	1	3	3
	Ceratopogonidae										
	Alluaudomyia sp.										
	Bezzia sp.		1								2
	Unidentified ceratopogonid										
	Empididae										
	Hemerodromia										1
	Tanyderidae										
	Simuliidae										
	Simulium sp.										

TAXA	GENUS SPECIES	P12/3	P12/4	P12/5	P12/6	P12/7	P14/3	P14/5	P14/7	P14/8	P14/10
EPHEMEROPTERA	Beetisce sp										
	Cennis sp.										
	Ephemerebella sp.										
	Isonychia										
	Tricorythodes sp.										
	Cinygmula subaequalis										
	Stenocron sp.		1								
	Stenonema sp.										1
	Spinadis wallace										
	Unknown sp. A (squatly bodies)										
	Unknown sp. B (slim guys)										
	Unknown sp. C										
Unidentified											
METEROPTERA	Mesovelia										
COGNATA	Argia sp.										
	Macromia sp.										
	Archilestes										
	Dromogomphus sp.								1		2
PLECOPTERA	Immatures										
	Perlinelle ephyre										
TRICHOPTERA	Immatures										
	Unidentified										
	Ceraclea										
AMPHIPODA	Macronema zebratum										
	Hydroptila sp.	1									
	Oecetis sp.		1								
	Chimarra sp.										2
	Polycentropus sp.										
	Type diverse										
	Unidentified								1	1	
	Synurella sp.										
	Gammarus sp.							2			
	Asellus sp.										
ISOPODA											
ACARINA											
MOLLUSCA											
PELECYPODA	Corbicula fluminea										
	Ferrissia rivularis	1	2	3	3	2					
GASTROPODA	Amnicole sp.										
	Gyraulus sp.										
OTHER	Unidentified										
NEMATODA											
NEMERTEA											
TOTAL NUMBER PER CORE		36	51	45	21	76	56	31	44	68	35
NUMBER OF SPECIES		11	20	17	16	22	15	18	16	27	20

Luxapalila Creek, Mississippi		Site																				
Sampling Date: 6/1989		R/1/1	R/1/2	R/1/4	R/1/5	R/1/10	R/2/1	R/2/5	R/2/6	R/2/7	R/2/8	R/3/1	R/3/2	R/3/4	R/3/5	R/3/7	R/4/1	R/4/2	R/4/8	R/4/9	R/4/10	
Counts of major taxa: raw data (for those samples identified to species, counts are summed here, all others are raw data)																						
Turbellaria	4																					
Oligochaeta	10	36	11	18	3	21	5	13	21	14	44	20	13	34	46	2	0	0	0	1	1	
Chironomidae	4	10	6	4	4	56	6	37	75	20	34	38	6	358	39	17	14	8	12	23	23	
Ceratopogonid																						
Coleoptera	1																					
Ephemeroptera																						
Odonata																						
Plecoptera																						
Trichoptera	5	2				12	21	81	6	6	6	6	1	22	22	3	1	2	1	4	4	
Amphipoda	2																					
Isopoda																						
Bivalvia	4	11																				
Gastropoda	2																					
MINOR GROUPS																						
Acarina	2	1																				
Heteroptera																						
Polychaeta																						
Hirudinea																						
Empididae																						
Collembola																						
Nematoda	4	4	2	4	4	4	2	4	2	4	2	4	6	13	3	7	4	7	4	1	1	
Nemertea																						
Simuliidae																						
Aeolosomatidae																						
Tanyderidae																						
TOTAL	31	70	19	28	8	113	15	94	233	43	93	68	29	430	118	32	20	18	20	35	35	

Luxapalila Creek, Mississippi
 Sampling Date: 6/89
 Raw species data: LX689RAW

TAXA	GENUS SPECIES	R/2/1	R/2/5	R/2/6	R/2/7	R/2/8	R/4/1	R/4/2	R/4/8	R/4/9	R/4/10
PLATYHELMINTHES											
TURBELLARIA											
	Dugesia tigrina										2
	Planaria sp.										
	Unidentified										
ANNELIDA											
POLYCHAETA											
	Sparganophilus										
	Haplotaenidae										
	Lumbriculidae	4		1	3	1					
	Naididae										
	Amphichaeta leydigi			1	2						
	Bretislevia bilongata										
	Bretislevia unidentata										
	Cheetogaster diaphanus										
	Dero digitata										
	Dero furcata										
	Dero nivea										
	Dero obtusa										
	Dero trifida										
	Dero sp.										
	Homochaeta naidina										
	Mais behningi				1	1					
	Mais bretscheri										
	Mais communis										
	Mais elinguis										
	Mais pardalis										
	Mais pseudobtusa										
	Mais simplex										
	Mais variabilis										
	Piquetella michigenesis										
	Pristina equisetata	3	3	1	3	6					
	Pristina leidy										
	Pristina synclites										
	Pristina sp.										
	Pristinella jenkinsae										
	Pristinella longidentata										
	Pristinella longisoma										
	Pristinella osborni										
	Pristinella sina	2		4							
	Stelvinia appendiculata										
	Specaria josinae										
	Stevensoniana trivandrama	4	1	1	5	5					
	Unidentified naididae										
	Tubificidae										
	Aulodrilus limobius										
	Aulodrilus plqueti										
	Aulodrilus pluriset										
	Branchiura sowerbyi										
	Limodrilus hoffmeisteri										
	Limodrilus rubripennis										

TAXA	GEMUS SPECIES	R1211	R1215	R1216	R1217	R1218	R1411	R1412	R1418	R1419	R14110
	Potamothenis vejovskyi	1	1	2		1	1				
	Unidentified tubificidae										
	Enchytraeidae										
	Barbidrilus paucisetus									1	
	Unidentified oligochaeta	7		3	9		1				
HIRUDINEA	Melobella elongata										
	Actinobdella sp.										
	Unidentified										
	AEOLOSOMATIDAE										
ARTHROPODA											
INSECTA											
	COLEOPTERA										
	Bidessus sp.				1						
	Cyphon										
	Halipus sp.										1
	Hydrochus sp.										
	Stenelmis sp.										
	COLLEMBOLA										
	Isotomurus palustris										
	Sminthurides sp.										
	Unidentified										
DIPTERA											
	CHIRONOMID										
	Chironominae										
	Chironomus sp.										
	Cladopelma										
	Cryptochironomus fulvus										
	Cryptochironomus sp.										
	Dicrotendipes neomolestus										
	Dicrotendipes nervosus Type 1										
	Dicrotendipes nervosus Type II										
	Endochironomus sp.										
	Glyptotendipes										
	Harnischia sp.			1							
	Microtendipes										1
	Miltothama babyi										
	Parachironomus abortivus										
	Parachironomus undine										
	Paralauterborniella nigrohalteralis									1	
	Paratendipes albinus										
	Paratendipes nudisquama										
	Phaenopsectra dyar										
	Phaenopsectra flavipes										
	Polypedilum convictum										1
	Polypedilum fallax										
	Polypedilum illinoense										1
	Polypedilum nr. scaloenum			2	1	2	1	3			
	Pseudochironomus sp.										
	Robackia sp.									5	
	Stenochironomus sp.										
	Stictochironomus sp.										
	Tribelos sp.										
	Xenochironomus sp.										
	Unidentified chironomini										2

TAXA	R1211	R1215	R1216	R1217	R1218	R1411	R1412	R1418	R1419	R14110
Tanytarsini										
Cladotanytarsus sp.		1	2	2	1					
Leuterborniella										
Micropectra sp.										
Rheotanytarsus sp.	2	2	4	22	1				1	5
Stempellinella sp.										
Tanytarsus coffmani	1									
Tanytarsus glabrescens										
Tanytarsus querlus	5			1						
Unidentified tanytarsini										
Orthocleidiinae										
Brillia										
Corynoneura celeripes	9		1	3				1		
Corynoneura taris	16		1	11	3					
Corynoneura sp.							1			
Cricotopus bicinctus		1		2	1					
Cricotopus trifascia								2		
Cricotopus sp.										1
Eukiefferiella sp.										
Manocleadius crassicornus	3									
Manocleadius distinctus										
Manocleadius rectinervis										
Manocleadius minimus										
Manocleadius sp.			1							1
Parakiefferiella sp.										4
Rheocricotopus sp.				1						1
Thienemanniella nr. fusca	3		6	9	4					1
Thienemanniella xena	4		15	17	4			1		
Unidentified orthocleidiinae				2	3					2
Tanypodinae										
Ablabesmyia mallochi										
Ablabesmyia parajanta	1									
Ablabesmyia taralle										
Clinotanypus										
Lebrundinia pilosella										1
Macropelopia sp.										
Natarsia sp.										
Milotanypus sp.	5		2	4	3					
Pentaneura sp.										
Procladius sp.										
Thienemanniya										
Unidentified tanypodinae	1									
Diamasinae										
Potthastia										
Unidentified Chironomidae	3	2	1	2	1	1	3	1	1	2
Ceratopogonidae										
Alluaudomyia sp.										
Bezzia sp.										
Unidentified ceratopogonid										
Empididae										
Hemerodromia										
Tanyderidae									9	3
Simuliidae										
Simulium sp.			4		1					
				33						

TAXA	GENUS SPECIES	R1211	R1215	R1216	R1217	R1218	R1411	R1412	R1418	R1419	R14110
EPHEMEROPTERA	Baetisca sp.										
	Caenis sp.										
	Ephemereilla sp.				2	1					
	Isonychia										
	Tricorythodes sp.										
	Cinygmula subaequalis										
	Stenonema sp.	24		7	6						
	Spinadisa wallace			3	3						
	Unknown sp. A (squatly bodies)										
	Unknown sp. B (slim guys)										
	Unknown sp. C			1	4						
	Unidentified										
	ODONATA	Mesovelia		1							
Argia sp.											
PLECOPTERA	Macromia sp.										
	Archilestes										
	Dromogomphus sp.										
	Immatures										
TRICHOPTERA	Perlinella ephyre										
	Immatures						3	1	2	1	4
	Unidentified										
MOLLUSCA	Ceraclea										
	Macronema zebrastrum										
	Hydroptila sp.				73						
	Oecetis sp.										
	Chimarra sp.	12		19	8	6					
	Polycentropus sp.										
	Type diversa										
	Unidentified			2							
	Synurella sp.										
	Gammarus sp.										
ISOPODA	Asellus sp.										
				5	7	1	3	1	1	1	2
PELECYPODA	Corbicula fluminea										
	Ferrissia rivularis										
	Ammicula sp.										
	Gyraulus sp.										
	Unidentified										
OTHER			8	2			7	4	7	4	1
NEMATODA											
TOTAL NUMBER		115	19	94	233	43	32	20	18	20	35
	NUMBER OF SPECIES	24	8	24	26	18	11	10	8	11	17

APPENDIX E
COUNTS OF MAJOR TAXA, PER CORE SAMPLE, COLLECTED FROM SITE 4
IN THE FALL OF 1989

Table E1. Counts of Major Taxa, per Core Sample, Collected From Site 4
in the Fall of 1989

Major Taxa	Pool Samples					Riffle Samples				
	1	3	4	6	9	2	3	5	7	9
Oligochaeta	1	4	1	3			1	1		
Chironomidae	15	59	78	100	42	30	19	18	21	13
Ceratopogonidae				1	3					
Coleoptera									1	1
Ephemeroptera		1								
Acarina					2	1				
Nematoda	2		2	2	1					
Nemertea					2					

Table E2. Species Composition of Oligochaetes and Chironomids
Collected From Site 4 in Fall 1989

Taxa	Pool Samples					Riffle Samples				
	1	3	4	6	9	2	3	5	7	9
Oligochaeta										
Naididae										
<i>Dero furcata</i>			1		2					
<i>Pristina leidy</i>									1	
<i>Specaria josinae</i>			2							
Tubificidae										
Unidentified				1					1	
Chironomidae										
Chironominae										
<i>Chironomus</i> sp.				1						
<i>Dicrotendipes nervosus</i> II				1						
<i>Harnischia</i> sp.										
<i>Phaenopsectra dyari</i>		3	27	38	33	17				
<i>Phaenopsectra flavipes</i>		1	5	10	3	6				
<i>Polypedilum fallax</i>				1						
<i>Polypedilum illinoense</i>			1			1				
<i>Polypedilum</i> nr. <i>scaloneum</i>	11	4	20	8	7	8	8	4	6	7
<i>Robackia</i> sp.							21	9	12	14
Tanytarsini										
<i>Tanytarsus querlus</i>				2	2	1				
Tanypodinae										
<i>Ablabesmyia parajanta</i>			5		1	3				
<i>Procladius</i> sp.				1						
Orthocladinae										
<i>Coryoneura celeripes</i>								1	1	1
<i>Coryoneura taris</i>							1		1	1
<i>Thienemanimyia</i> sp.				1						
<i>Thienemanniella</i> nr. <i>fusca</i>						1				