

AD-A238 222



A mixed limit theorem for stable random fields.


By

T.V. Kurien¹

and

J. Sethuraman²

Department of Statistics
Florida State University



September 1990
FSU Technical Report Number M 832
U.S. Army Technical Report Number D 114
A: OSR Technical Report Number 90-251

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Key Words: Mixed limit theorem, stable random field, pattern analysis.

¹ Research supported by the Air Force Office of Scientific Research under Grant AFOSR88-0040

² Research supported by Army Research Office Grant DAA03-90-G.



DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

91-05202



A random image can therefore be modelled by a collection of random variables U_α located at the vertices α of a graph $G = (V, e)$. Gibbs distributions have been used to specify the distribution of random images because the graph structure G can be tailored to take into account the dependencies intrinsic to the structure of the images arising in practice.

The general pattern analysis problem can be broadly described as follows. There is a true image I which cannot be observed. However, a deformed version I^D of I can be observed. The physical process deforming the true image I is due to the process of observation itself. This physical process could be mathematically modelled as an additive noise or by some other process. The pattern analysis problem is to reconstruct the true image I based on the observed image I^D by using a statistical procedure that is optimal in some sense.

Bayesian Image analysis first formulates a prior distribution for I which incorporates what is known about the structure of the true image I . Gibbs distributions have been used as prior distributions in Bayesian Image analysis [5]. The next step in Bayesian Image analysis is to update the prior notion of I using I^D . This gives rise to a posterior distribution on the set of images. Under fairly general conditions, the posterior distribution is a Gibbs distribution. The posterior mean or the posterior mode is usually used as an estimate of the true image I .

Bayesian Image analysis has had a good deal of success in dealing with pattern analytic problems. The HANDS project at Brown University [8] is an application of Bayesian Image analysis to recognizing biological shapes like human hands. For Bayesian Image analysis, it is necessary to generate an observation from the posterior distribution, and this is commonly done using the Gibbs Sampler [5], which in turn uses the Metropolis algorithm [9]. In fact, we can simulate observations from the prior distribution using the same Gibbs Sampler, to evaluate its suitability for the problem at hand. This is called pattern synthesis. However these methods are not very fast and require a good deal of computer time.

A natural question that now arises is whether we can replace simulations with direct mathematical approximations of such distributions when the number of vertices (sites) of the graph is very large. The answer to this question is in the affirmative, for special graphs, provided that the number of vertices increases to infinity and the scale of gray levels present at each site increases at a certain rate. Results that describe this kind of behavior are called mixed limit theorems and have been studied previously by Grenander and Sethuraman [7] and also by Chow [4].

A graph of interest in pattern analysis is the linear connection graph G_1 defined below. G_1 has vertex set $\{0, \dots, n\}$ and edge set $\{(i, i+1) : 0 \leq i < n\}$.

We will introduce some notation in order to state the mixed limit theorem for the graph G_1 . Let $A(\cdot)$ be a symmetric density and $Q(\cdot)$ be a density function on \mathbb{R} . Let (U_0, \dots, U_n) which reside at the vertices of G_1 have a joint density proportional to

$$\prod_{i=0}^{n-1} A(\sqrt{n}(u_{i+1} - u_i)) \prod_{i=0}^n Q(u_i).$$

The function A is called the acceptor function in the pattern analysis literature. The acceptor function models the local dependence between the U_i 's. The situation when the local

dependence specified by the pattern analytic problem at hand should make neighboring U_i alike can be modelled by taking the acceptor function A to be a symmetric density which is decreasing on \mathfrak{R}^+ . The situation when the local dependence should make neighboring U_i 's different can be modelled by taking A to be a symmetric density which has bounded support and is increasing on \mathfrak{R}^+ . In the equation above, the factor of \sqrt{n} represents an increase in the scale of gray levels present at each site as alluded to earlier.

Define

$$\begin{aligned} X_i &= \sqrt{n}U_i \\ \text{and } Y_i &= X_i - X_0, i = 0, \dots, n. \end{aligned} \tag{1.1}$$

Note that $Y_0 = 0$ and that the distributions of the X_i 's do depend on n . We shall not explicitly indicate this dependence for the rest of this paper. The case when $A(\cdot)$ has a second moment and $Q(x)$ is essentially of the form $\exp(-x^2)$ was the first situation to be considered. Mixed limit theorems for the distribution in $C[0, 1]$ of the process formed by linearly interpolating the X_i 's have been derived by Chow [4]. The limiting process in this case is a Gaussian process with a known covariance function. This corresponds to the situation in which the acceptor function A has thin tails. The case where the acceptor function A has thick tails is the situation that we will study in this paper. Let $A(\cdot)$ be a stable density with index α , $0 < \alpha < 2$. Then A has thicker than normal tails. Theorem 2.2 is a mixed limit theorem for the distribution of a family of stochastic processes on $[0, 1]$ based on the X_i 's. The processes that we will study are related to the stable processes (see Breiman [10]). Since stable processes are defined on the space $D[0, 1]$, this is the natural topological space to prove distributional limit theorems for the processes under study.

Section 2.

Construct the stochastic processes $G_n(t)$ and $X_n(t)$ on $[0, 1]$ as follows:

$$\begin{aligned} X_n(t) &= n^{-1/\alpha}X_{[nt]}, \\ G_n(t) &= n^{-1/\alpha}Y_{[nt]}. \end{aligned} \tag{2.1}$$

The random variables X_i and Y_i are as defined in equation (1.1). The processes $X_n(\cdot)$ and $G_n(\cdot)$ are random variables in $D[0, 1]$. A theorem that establishes the weak convergence in $D[0, 1]$ of the stochastic processes $G_n(t)$ and $X_n(t)$ will be referred to as a mixed limit theorem.

For the rest of this paper we shall assume that $Q(x) = \exp(-x^2)$ and that $A(\cdot)$ is a symmetric stable density of index α , $0 < \alpha < 2$.

Let $f_\alpha(\cdot)$ denote the symmetric stable density with index α . Let T_0 have density $f_\alpha(\cdot)$. Then $f_\alpha(\cdot)$ is a bounded density and $P\{T_0 > x\} \sim x^\alpha$. We will make use of these facts in the proof of Lemma 2.1.

Now (X_0, Y_1, \dots, Y_n) has a joint density proportional to

$$\prod_{i=0}^{n-1} f_\alpha(y_{i+1} - y_i) \exp(-n^{-1} \sum_{i=1}^n (y_i + x_0)^2 - x_0^2/n). \tag{2.2}$$

In Theorem 2.1 we will show that $G_n(\cdot)$ converges weakly to a process $G(\cdot)$ in $D[0,1]$. Corollary 2.1 of this theorem will show that $X_n(t)$ will also converge weakly in $D[0,1]$ to the same process $G(\cdot)$.

The method of proof of Theorem 2.1 is as follows. We will show that the distribution of $G_n(\cdot)$ conditional on $X_0 = x_0$ converges weakly in $D[0,1]$. We then show that the distribution of X_0 converges strongly on \mathfrak{R} . A result of Sethuraman [3] which is stated as Lemma 2.2 in the proof of Theorem 2.1 will establish the weak convergence of $G_n(\cdot)$. Lemma 2.1 is a weak convergence result for stable laws which will be used in the proof of Theorem 2.2.

Let X and Y be random variables. Then $X \stackrel{d}{=} Y$ denotes that X and Y are equal in distribution. Let $\{X_n, n = 1, \dots\}$ be a sequence of random variables on a complete separable metric space S and let X and X_0 be random variables in S . Then $X_n \xrightarrow{w} X$ denotes that $\mathcal{L}(X_n)$ converges weakly to $\mathcal{L}(X)$ in S . $\mathcal{L}(X_n|X_0 = x_0)$ denotes the conditional distribution of $(X_n$ given that $X_0 = x_0)$. Let $R(t)$ be a stationary and independent increment process in $D[0,1]$ with $R(t) \stackrel{d}{=} t^{1/\alpha} T_0$. Lemma 2.1 below explicitly constructs such a process $R(\cdot)$ in $D[0,1]$.

Lemma 2.1

Let $0 < \alpha < 2$ and let T_0, T_1, \dots be i. i. d. with density f_α . Consider the partial sums $S_i = \sum_{j=1}^i T_j, i = 1, 2, \dots, S_0 = 0$ and let the process $(R_n(t), 0 \leq t \leq 1)$ be defined by

$$R_n(t) = n^{-1/\alpha} S_{[nt]}.$$

Then: $R_n(\cdot) \xrightarrow{w} R(\cdot)$ in $D[0,1]$.

Proof

The method of proof is the usual one, i.e., we shall first prove finite-dimensional convergence and then verify the appropriate tightness conditions.

Let us calculate the characteristic function $\phi_n(\cdot, t)$ of $R_n(t), t \in [0,1]$.

$$\begin{aligned} \phi_n(u, t) &= E(\exp(iuR(t))) \\ &= E(\exp(iun^{-1/\alpha} \sum_{1 \leq j \leq [nt]} T_j)) \\ &= \exp(-|u|^\alpha d_n(t)) \end{aligned} \tag{2.3}$$

where $d_n(t) = (\frac{[nt]}{n})^{1/\alpha} \rightarrow t^{1/\alpha}$ as $n \rightarrow \infty$ for $t \in [0,1]$.

Let $t_0 = 0 < t_1 < \dots < t_k < t_{k+1} = 1$, and let $f_\alpha(\cdot)$ be the symmetric stable density with index α .

Let

$$X_{i,n} = R_n(t_i) - R_n(t_{i-1}), \quad 1 \leq i \leq k+1, \quad n = 1, 2, \dots$$

Note that $\{X_{i,n}, i = 1, \dots, k+1\}$ are mutually independent random variables. Let $f_{i,n}$ be the density of $X_{i,n}$. Then by (2.3) it is clear that $f_{i,n}$ is given by

$$f_{i,n}(x) = f_\alpha\left(\frac{x}{d_n(t_j - t_{j-1})}\right) / d_n(t_j - t_{j-1}).$$

Since $X_{i,n}$ are mutually independent and $R_n(t_i) = \sum_{j=0}^i X_{j,n}$, $i = 1, \dots, k+1$, the density of $(R_n(t_1), \dots, R_n(t_k))$ is given by

$$\prod_{i=1}^k f_{i,n}(y_i - y_{i-1}) \text{ where } y_0 = 0.$$

This density converges to the density of $(R(t_1), \dots, R(t_k))$. Hence the finite dimensional distributions converge. We will verify tightness next.

Let $0 = t_0 < t_1 < t_2 < t_3 < t_4 = 1$, $\epsilon > 0$ and let U, V be stable symmetric random variables with index α . Then by the mutual independence of the increments $X_{i,n}$, and the facts about the tail probabilities of stable symmetric random variables, it follows that

$$\begin{aligned} & P\{|R_n(t_3) - R_n(t_2)| \geq \epsilon, |R_n(t_2) - R_n(t_1)| \geq \epsilon\} \\ &= P\{|R_n(t_3) - R_n(t_2)| \geq \epsilon\} \cdot P\{|R_n(t_2) - R_n(t_1)| \geq \epsilon\} \\ &= P\{|U| \geq \frac{\epsilon}{d_n(t_2 - t_1)}\} \cdot P\{|V| \geq \frac{\epsilon}{d_n(t_3 - t_2)}\} \\ &\leq \frac{M \cdot (t_3 - t_1)^2}{\epsilon^{2\alpha}}. \end{aligned}$$

Using Theorem (15.6) in Billingsley [1], tightness of the distributions is immediate. This proves the lemma. \square

Define $\nu(x_0, x)$, $f(u)$ and $\mu(u)$ as follows

$$\begin{aligned} \nu(x_0, x) &= \exp\left(-\int_0^1 (x(t) + x_0)^2 dt\right) \left(\int_{D[0,1]} \exp\left(-\int_0^1 (x(t) + x_0)^2 dt\right) dP(x)\right)^{-1} \\ f(u) &= \int_{D[0,1]} \exp\left(-\int_0^1 (x(t) + u)^2 dt\right) dP(x) \\ \mu(u) &= f(u) / \int_{\mathbb{R}} f(y) dy. \end{aligned}$$

Note The finiteness of these integrals becomes clear in the proof of Theorem 2.1 below.

In order to state the mixed limit theorem for G_n we introduce the following notation.

Let $Q_n = \mathcal{L}(G_n)$, $P_n = \mathcal{L}(R_n)$, $P = \mathcal{L}(R)$. For x_0 in \mathbb{R} let $Q_{n,x_0} = \mathcal{L}(G_n | X_0 = x_0)$. These are probability measures on $D[0, 1]$.

Theorem 2.1

Let $A(\cdot)$ be a stable density of index α , $0 < \alpha < 2$. Then Q_n converges to Q weakly in $D[0, 1]$ where the density of Q is given by

$$dQ(x) = \frac{\exp(-\int_0^1 (x(t) - \int_0^1 x(u)du)^2 dt)}{\int_{D[0,1]} \exp(-\int_0^1 (x(t) - \int_0^1 x(u)du)^2 dt) dP(x)} dP(x).$$

Proof

If we omit the exponential term in (2.2), then $G_n(\cdot)$ and $R_n(\cdot)$ have the same distribution. To study Q_n , we shall first show that Q_{n,x_0} converges weakly in $D[0, 1]$. We will then show that the densities of X_0 converge. This proves strong convergence for the densities of X_0 via Scheffe's theorem [2]. This will prove a weak convergence result for the distribution of $Q_n(\cdot)$ on $D[0, 1]$, via a result of Sethuraman [3] stated below.

Lemma 2.2

Let Λ_n be a sequence of probability measures on $V \times W$ where V and W are topological spaces. Let μ_n be the marginal distribution of Λ_n on V and $\nu_n(v, \cdot)$ be the conditional p. m. of Λ_n on W . Suppose that $\mu_n(A) \rightarrow \mu(A) \forall A \subseteq V$ and $\nu_n(v, \cdot) \rightarrow \nu(v, \cdot)$, weakly for almost all v (w. r. t μ). Then, $\Lambda_n \rightarrow \Lambda$ weakly where

$$\Lambda(A \times B) = \int_A \nu(v, B) d\mu(v)$$

for each measurable rectangle $A \times B \in V \times W$.

We shall apply this theorem where $W = D[0, 1]$, $V = \mathfrak{R}$, μ_n is the distribution of X_0 , and $\nu_n(x_0, \cdot) = \mathcal{L}(Q_n(\cdot) | X_0 = x_0)$.

We will now show the weak convergence of $\mathcal{L}(Q_n(\cdot) | X_0 = x_0)$ by considering it's Radon-Nikodym derivative with respect to P_n and using the fact that $P_n \xrightarrow{w} P$. We shall denote a function $x(\cdot)$ in $D[0, 1]$ by x from now on. Let $\frac{dQ_{n,x_0}}{dP_n}(x)$ denote the Radon-Nikodym derivative of Q_{n,x_0} with respect to $P_n(\cdot)$ at x for x in $D[0, 1]$. For x in $D[0, 1]$ let

$$k_n(x) = \exp(-n^{-1} \sum_{i=1}^n (x(i/n) + x_0)^2)$$

$$\text{and } k(x) = \exp(-\int_0^1 (x(t) + x_0)^2 dt).$$

Then

$$\frac{dQ_{n,x_0}}{dP_n}(x) = k_n(x) / \int_{D[0,1]} k_n(y) dP_n(y).$$

Let f be a bounded continuous function from $D[0, 1]$ to \mathfrak{R} . Then

$$\int_{D[0,1]} f(x) dQ_{n,x_0} = \int_{D[0,1]} f(x) k_n(x) dP_n / \int_{D[0,1]} k_n(x) dP_n. \quad (2.4)$$

The weak convergence of Q_{n,x_0} will be proved by showing that

$$\begin{aligned} \int_{D[0,1]} f(x) k_n(x) dP_n(x) &\rightarrow \int_{D[0,1]} f(x) k(x) dP(x) \text{ and} \\ \int_{D[0,1]} k_n(x) dP_n(x) &\rightarrow \int_{D[0,1]} k(x) dP(x). \end{aligned} \quad (2.5)$$

This will prove that $Q_{n,x_0} \xrightarrow{w} Q_{x_0}$ and that

$$\frac{dQ_{x_0}}{dP}(x) = k(x) / \int_{D[0,1]} k(y) dP(y).$$

Lemma 2.3 will help us to verify this.

Lemma 2.3

Let $f : \mathfrak{R} \rightarrow \mathfrak{R}$ be continuous and $x_n \rightarrow x \in D[0, 1]$. Then,

$$n^{-1} \sum_{i=1}^n f(x_n(i/n)) \rightarrow \int_0^1 f(x(t)) dt.$$

Proof

Without loss of generality, we can assume that $f(x) = x$ since the map $x \rightarrow f(x)$ induced on $D[0, 1]$ by f is a continuous map from $D[0, 1] \rightarrow D[0, 1]$. Hence, it suffices to show that

$$n^{-1} \sum_{i=1}^n x_n(i/n) \rightarrow \int_0^1 x(t) dt.$$

Let us define

$$y_n(t) = \begin{cases} x_n(i/n) & \text{if } (i-1)/n < t \leq i/n \\ x_n(0) & \text{if } t = 0. \end{cases}$$

Then, $n^{-1} \sum_{i=1}^n x_n(i/n) = \int_0^1 y_n(t) dt$. Since $x_n \rightarrow x$ in $D[0, 1]$, there exist λ_n which are strictly increasing, continuous and surjective mappings of $[0, 1] \rightarrow [0, 1]$ such that

$$\begin{aligned} x_n(\lambda_n(t)) &\rightarrow x(t) \text{ uniformly in } t \\ \text{and } \lambda_n(t) &\rightarrow t \text{ uniformly in } t. \end{aligned} \quad (2.6)$$

Let $s \in [0, 1]$ be a continuity point of $x(\cdot)$. The following argument shows that $y_n(s) \rightarrow x(s)$.

There exists i_n such that $(i_n - 1)/n < s \leq i_n/n$. Let $s_n^* = \lambda_n^{-1}(i_n/n)$. Then $y_n(s) = x_n(\lambda_n s_n^*)$. Hence

$$\begin{aligned} |y_n(s) - x(s)| &\leq |x_n(\lambda_n s_n^*) - x(s_n^*)| + |x(s_n^*) - x(s)| \\ &\leq \sup_{t \in [0, 1]} |x_n(\lambda_n t) - x(t)| + |x(s_n^*) - x(s)|. \end{aligned} \quad (2.7)$$

Now $s_n^* \rightarrow s$ and since s is a continuity point of $x(\cdot)$, $|x(s_n^*) - x(s)| \rightarrow 0$ as $n \rightarrow \infty$. The first term in (2.7) converges to 0 by (2.6). Hence $y_n(s) \rightarrow x(s)$, as asserted. Since the number of discontinuity points of $x(\cdot)$ is countable, $y_n(s) \rightarrow x(s)$ a. s. (Lebesgue). By the Dominated Convergence Theorem, and because x_n and hence y_n are uniformly bounded,

$$\int_0^1 y_n(t) dt \rightarrow \int_0^1 x(t) dt.$$

This proves Lemma 2.3. \square

Hence it follows that

$$\begin{aligned} n^{-1} \sum x_n^2(i/n) &\rightarrow \int_0^1 x^2(t) dt \text{ and} \\ n^{-1} \sum x_n(i/n) &\rightarrow \int_0^1 x(t) dt. \end{aligned} \quad (2.8)$$

Thus $\exp(-n^{-1} \sum_{i=1}^n (x(i/n) + x_0)^2) \rightarrow \exp(-\int_0^1 (x(t) + x_0)^2 dt)$ By (2.8) since $\exp(-t)$ is a bounded continuous function on $[0, \infty]$, $k_n(x_n) \rightarrow k(x)$ and $f(x_n)k_n(x_n) \rightarrow f(x)k(x)$ as $x_n \rightarrow x$ in $D[0, 1]$. Also note that $k_n(x)$ and $k(x)$ are bounded.

Then by using Theorem 5.5 of Billingsley [1] it follows that both terms in (2.5) converge as stated.

Thus $Q_{n, x_0} \xrightarrow{w} Q_{x_0}$ as $n \rightarrow \infty$ and

$$\begin{aligned} \frac{dQ_{x_0}}{dP} &= \exp\left(-\int_0^1 (x(t) + x_0)^2 dt\right) \left(\int_{D[0, 1]} \exp\left(-\int_0^1 (x(t) + x_0)^2 dt\right) dP(x)\right)^{-1} \\ &= \nu(x_0, x). \end{aligned} \quad (2.9)$$

Now, we will show that the distribution of X_0 converges strongly by using Scheffe's theorem [2].

Define

$$f_n(x_0) = \exp(-x_0^2/n) \int_{D[0, 1]} \exp\left(-n^{-1} \sum_{i=1}^n (x(i/n) + x_0)^2\right) dP_n(x).$$

Then, the density of X_0 is given by $f_n(x_0) / \int_{-\infty}^{\infty} f_n(x_0) dx_0$. By (2.5) we know that $f_n(x_0) \rightarrow f(x_0)$ as $n \rightarrow \infty$ where

$$f(x_0) = \int_{D[0, 1]} \exp\left(-\int_0^1 (x(t) + x_0)^2 dt\right) dP(x). \quad (2.10)$$

In order to successfully apply Scheffe's theorem, we need to check that $\int_{\mathfrak{R}} f_n(x_0) dx_0 \rightarrow \int_{\mathfrak{R}} f(x_0) dx_0$.

By Fubini's Theorem and elementary algebra we know that

$$\begin{aligned} \int_{\mathfrak{R}} f_n(x_0) dx_0 &= \int_{\mathfrak{R}} \exp(-x_0^2/n) \int_{D[0,1]} \exp(-n^{-1} \sum_{i=1}^n (x(i/n) + x_0)^2) dP_n(x) dx_0 \\ &= \int_{D[0,1]} \exp\left(-n^{-1} \sum_{i=1}^n x^2(i/n) + ([n(n+1)]^{-1} \sum_{i=1}^n x(i/n))^2\right) \\ &\quad \times \int_{\mathfrak{R}} \exp\left(-n/(n+1)(x_0 - [n(n+1)]^{-1} \sum_{i=1}^n x(i/n))^2\right) dx_0 dP_n(x) \\ &\rightarrow \sqrt{\pi} \int_{D[0,1]} \exp\left(-\int_0^1 x^2(t) dt + \left(\int_0^1 x(t) dt\right)^2\right) dP(x). \end{aligned}$$

This limit is equal to $\int_{\mathfrak{R}} f(x_0) dx_0$, because

$$\begin{aligned} \int_{\mathfrak{R}} f(x_0) dx_0 &= \int_{\mathfrak{R}} \int_{D[0,1]} \exp\left(-\int_0^1 (x(t) + x_0)^2 dt\right) dP(x) dx_0 \\ &= \sqrt{\pi} \int_{D[0,1]} \exp\left(-\int_0^1 x^2(t) dt + \left(\int_0^1 x(t) dt\right)^2\right) dP(x). \end{aligned} \tag{2.11}$$

Thus the distribution of X_0 converges strongly to a distribution with density function given by

$$d\mu(x_0) = f(x_0) / \int_{\mathfrak{R}} f(x_0) dx_0. \tag{2.12}$$

Hence, by Lemma 2.2 it follows that Q_n converges weakly to a distribution Q on $D[0,1]$ with density given by

$$dQ(x) = \left(\int_{\mathfrak{R}} \nu(x, x_0) d\mu(x_0) \right) dP(x). \tag{2.13}$$

The integral in (2.13) can be further simplified by using (2.10) and (2.11).

$$\begin{aligned} \int_{\mathfrak{R}} \nu(x_0, x) d\mu(x_0) &= \frac{\int_{\mathfrak{R}} \exp\left(-\int_0^1 (x(t) + x_0)^2 dt\right) dx_0}{\int_{\mathfrak{R}} f(x_0) dx_0} \\ &= \frac{\exp\left(-\int_0^1 (x(t) - \int_0^1 x(u) du)^2 dt\right)}{\int_{D[0,1]} \exp\left(-\int_0^1 (x(t) - \int_0^1 x(u) du)^2 dt\right) dP(x)}. \end{aligned}$$

This completes the proof of Theorem 2.1 \square

The following corollary is a mixed limit theorem for $X_n(t)$. This shows that mixed limit theorems exist for the stable case and the limits are Gibbs distributions.

Corollary 2.1

$X_n(t)$ converges weakly in $D[0, 1]$ to the process $G(\cdot)$ which has distribution Q .

Proof

Since $X_n(t) = n^{-1/\alpha+1/2}X_0 + G_n(t)$, $0 < \alpha < 2$, and the distributions of X_0 converge strongly as $n \rightarrow \infty$ it follows from Slutsky's Theorem that the distributions of $X_n(\cdot)$ converges weakly in $D[0, 1]$ to $Q(\cdot)$. \square

Remark. The next problem that has to be addressed is that of explicitly evaluating the integral $\int_{D[0,1]} \exp(-\int_0^1 (x(t) - \int_0^1 x(u)du)^2 dt) dP(x)$. This will be necessary in order to simulate observations from the limiting distribution Q .

References

- [1] Billingsley, P. (1968). Convergence of Probability Measures. *John Wiley, New York.*
- [2] Billingsley, P. (1986). Probability and Measure. *John Wiley, New York.*
- [3] Sethuraman, J. (1961). Some limit theorems for joint distributions. *Sankhya A, 23, 379-386.*
- [4] Chow, Y. (1989). A limit theorem for Pattern synthesis in Image processing. *Preprint.*
- [5] Geman, D. and Geman, S. (1984). Stochastic relaxation, Gibbs distributions and the Bayesian restoration of images. *IEEE-PAMI 6, 721-741.*
- [6] Chow, Y. and Grenander, U. (1985). A singular perturbation problem. *J. of Integral Eqns. 9, 63-73.*
- [7] Grenander, U. and Sethuraman, J. (1986). Mixed limit theorems for pattern synthesis. *Unpublished.*
- [8] Grenander, U. and Keenan, D. M. (1988). HANDS. A pattern theoretic approach to processing biological images. *Unpublished.*
- [9] Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. N. and Teller, E. (1953). Equations of state calculations by fast computing machines. *J. of Chem. Phys. 21, 1087-1091.*
- [10] Breiman, L. (1968). Probability. *Addison Wesley, Massachusetts.*

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) FSU Tech Report No. M832		5. MONITORING ORGANIZATION REPORT NUMBER(S) USARO Tech Report No. D-114	
6a. NAME OF PERFORMING ORGANIZATION Florida State University	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION U. S. Army Research Office	
6c. ADDRESS (City, State, and ZIP Code) Department of Statistics Florida State University Tallahassee, Florida 32306		7b. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION U. S. Army Research Office	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAAL03-90-G-0103	
8c. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO. DAAGL03
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) A mixed limit theorem for stable random fields			
12. PERSONAL AUTHOR(S) T. V. Kurien and J. Sethuraman			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) September, 1990	15. PAGE COUNT 11
16. SUPPLEMENTARY NOTATION The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
A mixed distributional limit theorem for a stable random field of index $0 < \alpha < 2$ is derived. These random fields are of special interest in pattern analysis, in particular, in pattern synthesis. This paper considers the case when the underlying graph that the random field is defined on is linear. This result is encouraging insofar as it shows that the mixed limit theorems do exist in the stable case. The final limiting distribution can be written in terms of the stable process of index α in $D[0, 1]$.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL