Marine Physical Laboratory



1



TEST OF NEWTONIAN GRAVITY USING DSV SEA CLIFF

Final Report to Office of Naval Research Grant N00014-90-J-1063 For the Period 10-01-88 - 12-31-90 Principal Investigator(s): John A. Hildebrand and Mark A. Zumberge

	Acous	Alva Bor	···· /·		
10	BTT: BTT: Uncom Jasti	GMARI IND COMERC Floaticn			
	Sy Distribution, Aveilabtion				
2	Dist	Special	07		
	A-1				

MPL-U-48/91 June 1991

Approved for public release; distribution unlimited.



University of California, San Diego Scripps Institution of Oceanography



91 7 15 070

Best Available Copy

SECURITY CLASSIFICATION OF THIS PAGE

			REPORT DOCUM	MENTATION I	PAGE					
1. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			16 RESTRICTIVE MARKINGS							
28. SECURITY	CLASSIFICATION	AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT						
26. DECLASSIFICATION / DOWNGRADING SCHEDULE			Approved for public release; distribution unlimited.							
4. PERFORMIN	G ORGANIZATI	ON REPORT NUMBER	R(S)	5. MONITORING ORGANIZATION REPORT NUMBER(S)						
MPL-U-48/91										
68. NAME OF PERFORMING ORGANIZATION			66. OFFICE SYMBOL	78. NAME OF MONITORING ORGANIZATION						
Marine Physical Laboratory			(If applicable) MPL	Office of Naval Research Department of the Navy						
6c. ADDRESS (City, State, and ZIP Code)			7b. ADDRESS (City, State, and ZIP Code)							
University of California, San Diego			800 North Quincy Street							
\cdot San Die	rgo, CA 9	2152	graphy	Allington	, VA 22217	-5000				
8a. NAME OF	a, NAME OF FUNDING / SPONSORING 8b. OFFI			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER						
Office	Office of Naval Research ONR			N00014-90-J-1063						
BC. ADDRESS (City, State, and	ZIP Code)	┨╌╼╕┄╌╡╾╼╼╍┽╼╸ _{╢┺} ┉┍┉╸┽┶╾═╺┑	10. SOURCE OF	UNDING NUMBE	RS				
Departn	ent of the	e Navy Stroot		PROGRAM	PROJECT	TASK		WORK UNIT		
Arlingt	con, VA = 22	2217-5000		ELEMENT NO.	NO.					
11. TITLE (Incl	ude Security Cl	lassification)		-I	<i>I</i>			L		
TEST OF NEWTONIAN GRAVITY USING DSV SEA CLIFF										
12. PERSONAL	LAUTHOR(S)	and Mark A	7umaorgo	<u></u>	<u></u>					
John A. Hildebrand and Mark A. Zuligerge				14 DATE OF REPO	14 DATE OF REPORT I Year Month Davi 15 PAGE COUNT					
Final R	eport	FROM	to	June 19	91		3			
16. SUPPLEM	ENTARY NOTAT	ION								
17.	COSATI	CODES	18. SUBJECT TERMS	(Continue on rever	se if necessary ar	nd identify b	y bloc	k number)		
FIELD	GROUP	SUB-GROUP	geophysical	measurement	s, gravitat	ional co	nsta	nt,		
			Newton's in	nverse square	e law					
19 ABSTRAC	T (Continue on	reverse if necessary	and identify by block	number)						
	•		, ,							
] .	Saveral reco	nt geophysical mag	suramants have sourch	ad for deviations	from Newton's i	nverco caua	re law	hv		
testing for range dependence of the gravitational constant G . One group of these experiments observes the vertical										
gravity gradient above the earth using measurements on tall towers (Eckhardt et al. 1988; Thomas et al. 1989;										
Speake et al. 1990) and tests for deviations from Newtonian gravity but provides no direct estimate of G . Another										
group or geophysical experiments observes gravity from material stabs of known density by measuring within mines and horeholes (Stacey et al. 1987; Zumberge et al. 1990) or above fluctuating lakes (Moore et al. 1988; Muller et al.										
1990) and yield estimates of $G(r)$. Collectively, these experiments offer little evidence for deviations from										
Newtonian gravity in the scale range 10 m to 1200 m.										
20 DISTRIBU	TION / AVAILAB	ILITY OF ABSTRACT		21, ABSTRACT S	ECURITY CLASSIF					
	55IFIED/UNLIMI1	TED UX SAME AS	RPT. 🗌 DTIC USERS	I INCLAS	SIFIFD					

TEST OF NEWTONIAN GRAVITY USING DSV SEA CLIFF

Principal Investigator(s) John A. Hildebrand Mark A. Zumberge

Final Report to Office of Naval Research Grant N00014-90-J-1063 for the Period 10-01-88 - 12-31-90

Scripps Institution of Oceanography University of California at San Diego La Jolla, CA 92093-0205 (619) 534-4069, 534-3533

Scientific Objectives

Several recent geophysical measurements have searched for deviations from Newton's inverse square law by testing for range dependence of the gravitational constant G. One group of these experiments observes the vertical gravity gradient above the earth using measurements on tall towers (Eckhardt *et al.* 1988; Thomas *et al.* 1989; Speake *et al.* 1990) and tests for deviations from Newtonian gravity but provides no direct estimate of G. Another group of geophysical experiments observes gravity from material slabs of known density by measuring within mines and borcholes (Stacey *et al.* 1987; Zumberge *et al.* 1990) or above fluctuating lakes (Moore *et al.* 1988; Muller *et al.* 1990) and yield estimates of G(r). Collectively, these experiments offer little evidence for deviations from Newtonian gravity in the scale range 10 m to 1200 m.

Project Objectives

Our project was designed to measure the gravitational constant within the ocean using the submersible *Sea Cliff* over a scale length of 5000 m. A number of observables are needed to calculate the gravitational constant in an oceanic setting including: gravitational acceleration, depth, oceanic density, scalloor density, and scalloor topography. The strengths of our experiment are that gravity measurements were made over three-dimensions, that oceanic density was accurately characterized over three-dimensions, that scalloor density and topography were accurately mapped, and that the scale-length was large. This project was the first time a submersible was used as a platform for water-column gravity measurements.

Present Status and Progress During the Project Duration

Our experimental site was in the northeast Pacific ocean ($35^{\circ}13$ 'N $132^{\circ}00$ 'W), approximately 600 km west of Monterey, California. This site was chosen to minimize gravity perturbations from the ocean-continent boundary, from oceanic fracture zones, and from oceanic currents and fronts. A 7000 km² area surrounding the site was mapped by multi-beam echo-sounding to characterize the topographic relief and seismic reflection profiling to measure the sediment thickness. Water density was measured with conductivity, temperature and depth (CTD) profiles, which determine density to better than 1 part in 10^4 using the seawater equation of state.

We measured gravitational acceleration within the water-column using a Bell Aerospace BGM-3 gravity meter on a gyro-stabilized platform. The gravimeter sensor is a pendulous accelerometer, which has a high-gain servo loop to mechanically null a proofmass using a magnetic force. The sensor and associated electronics are housed in an oven to minimize thermal drift. The gravity sensor has a time constant of 4.5 seconds. A stabilized platform provides a vertical reference for the gravity sensor, using two gyro-stabilized feedback loops. It is capable of stabilization over an angular range of $\pm 30^{\circ}$ for pitch and $\pm 45^{\circ}$ for roll and maintains verticality to better than 1 milliradian.

The gravimeter was placed in the U. S. Navy submersible *Sea Cliff*, which is capable of descending to depths of 6100 m (20000 ft). A submersible pilot and an equipment operator accompanied the gravity meter to the scafloor on a total of four dives. The dives were, on average, 11 hours long and reached depths of more than 5000 m. At the beginning of each dive the submersible descended at a central position. At the scafloor, the submersible ballast was trimmed to be nearly neutrally buoyant, and gravity was read at a consistent location, allowing a check for instrumental drift. The submersible depth was monitored by two quartz pressure gauges (Paroscientific Model 410KT), which are accurate to better than one-meter when corrected for water density. The depth measurements at the central bottom site were repeatable to 0.2 m and the gravity measurements were repeatable to 200 μ Gal, with an instrumental drift of 100 μ Gal/day. During three of the dives, the submersible was driven along the bottom approximately 1.5 km and an additional gravity reading was made before ascent to the surface. The resulting pattern of dive ascent locations is an equilateral triangle with a central point.

During the submersible deployments, gravity and depth were recorded every 12 seconds. To ascend, a scries of weights were dropped giving the submersible positive buoyancy. We maintained the submersible at a uniform ascent rate, between 10 m/min and 25 m/min, however, weight drop events produced detectable vertical accelerations. To correct for vertical accelerations, we subtracted the second-derivative of the depth from the gravity data. These corrections substantially removed from the gravity the effect of weight drops. Figure 1 shows the gravity corrected for depth only, the corrections from the second-derivative of depth, and the corrected gravity for one of the four dives.

To account for the attraction of local terrain we used a digital representation of the multi-beam echo-sounding map. The terrain from a region 30 km by 30 km surrounding the dive sites was included at a 0.2 km grid spacing. The average scafloor density in this region is 2690 kg/m³, derived from the vertical gradient of an on-bottom gravity survey. We assumed the terrain to be uniformly 2690 kg/m³, and calculated its attraction using the average scafloor depth (5104 m) as a reference plane. The terrain attraction was calculated for locations along the dive ascents and is shown as a dashed line in the lower panel of Figure 1.

In Figure 1, the depth correction to the data accounts for the attraction of the sea water using the laboratory value of G. Because there is little or no residual slope in the corrected gravity profile, we see that the laboratory value of the gravitational constant does an excellent job of matching the observed vertical gravity gradient, especially for the oceanic slab that is more than one km above the bottom (where the terrain effect is negligible). This leads to a preliminary conclusion that the gravitational constant at scale lengths of one to four km equals the laboratory value. We expect that after completing the analyses of the other *Sea Cliff* dives, and incorporating these data with preveiously obtained observations with the submarine *Dolphin*, on-bottom gravity measurements, and sea-surface gravity surveys, this experiment will provide the best constraints on deviations from Newton's law to date.

Publications for FY90

....

Stevenson, J. M., Zumberge, M. A., and Hildebrand J. A., 1990. Deep-Ocean Gravity Measurements Aboard the Submersible Seacliff, EoS, 71, 1277.



Figure 1. a) Raw gravity and depth data for a submersible dive. b) Gravity data corrected for depth using the laboratory value of $G = 6.672 \times 10^{-11}$ MKS (top), vertical acceleration correction from second differences of the pressure depth data (middle), and the sum of gravity and vertical acceleration data (bottom). c) Ascent gravity data as a function of depth (solid line) and terrain correction (dashed line).