

AD-A243 085



**EXCITED STATE KINETICS OF MERCURY AND
CADMIUM HALIDES AND NITROGEN FLUORIDE**

2

D. W. Setser

Kansas State University
Department of Chemistry
Manhattan, KS 66506

DTIC
ELECTE
OCT 16 1991
S D D

September 1991

Final Report

APPROVED FOR PUBLIC RELEASE DISTRIBUTION UNLIMITED.

91-13367



**PHILLIPS LABORATORY
Directorate of Lasers and Imaging
AIR FORCE SYSTEMS COMMAND
KIRTLAND AIR FORCE BASE, NM 87117-6008**



REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE September 1991	3. REPORT TYPE AND DATES COVERED Final 15 Mar 84 - 14 Apr 85	
4. TITLE AND SUBTITLE Excited State Kinetics of Mercury and Cadmium Halides and Nitrogen Fluoride		5. FUNDING NUMBERS G: F29601-84-K-0030 PE: 61102F PR: 2303 TA: 03 WU: 01	
6. AUTHOR(S) D. W. Setser		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Kansas State University Department of Chemistry Manhattan, KS 66506		10. SPONSORING / MONITORING AGENCY REPORT NUMBER PL-TN--91-1062	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Phillips Laboratory Kirtland AFB, NM 87117-6008		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release. Distribution unlimited.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Twelve progress reports on the investigation of the excited state kinetics of mercury and cadmium halides and nitrogen fluoride conducted by the Department of Chemistry, Kansas State University, Manhattan, Kansas, between 15 March 1984 and 1 April 1985.			
14. SUBJECT TERMS Excited State Kinetics, Hg Halides, Cd Halides, Nitrogen Fluorides, NF(b) Afterglow, NF(b) Quenching Rate Constants			15. NUMBER OF PAGES 32
17. SECURITY CLASSIFICATION OF REPORT Unclassified			16. PRICE CODE
18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT SAR	

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U. S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES AND
NITROGEN FLUORIDE

Contract No: F29601-84-0030



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, KS 66506

Scientific Program Officer: Dr. Vernon Schlie
Contract Duration: March 15, 1984 - March 14, 1985
Monthly Progress Report (First Report)
Report Period: March 15 - April 15 (1984)

Progress Report (1)

The contract award date was March 15, 1984. However, the paperwork giving permission to start formal spending on the contract was not received and approved by Kansas State University until ~April 2. Thus, the efforts of one postdoctoral fellow and one student were transferred to the Air Force Contract as of April 1.

Two small scale flowing-afterglow systems were in operation at the time the contract was awarded. Therefore, experimental work could be started immediately. The first projects were to measure some quenching rate constants for NF(b) and to demonstrate the general utility of a flowing-afterglow source for NF(b) for both afterglow systems. The NF(b) source is the reaction of Ar(³P_{0,2}) metastable atoms with NF₂. The following total quenching rate constants have been measured for NF(b, v=0) with Air Force support.

<u>Reagent</u>	<u>Rate Constants</u> (10 ⁻¹⁴ cm ³ molec ⁻¹ sec ⁻¹)
O ₂	4.2 ± 0.1
Cl ₂	7200 ± 1200
HF	967 ± 15 96.7 ± 15
NF ₂	0.6 ± 0.4
N ₂ F ₄	1.4 ± 0.5

In the absence of reagent and with ~ 2 torr of Ar carrier gas pressure, the decay of NF(b) in our system is largely determined by the purely radiative decay rate, i.e., the quenching rate of NF(b) at the Pyrex glass walls is slow (but observable).

During the next month kinetic studies of NF(b) will continue and we will modify the larger flowing-afterglow apparatus for a study of vibrational-to-electronic energy transfer between HF(v) and NF(a).

D. W. Setser

*PS. On my final contract I noted that cadmium had evolved to calcium in the title of the contract. Please check to see that your listing of our contract is correct. I do not want calcium kinetics showing on your computer print-out of our contract title.

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U. S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES AND
NITROGEN FLUORIDE

Contract No: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, KS 66506

Scientific Program Officer: Dr. Vernon Schlie

Contract Duration: March 15, 1984 - March 14, 1985

Monthly Progress Report (Second Report)

Report Period: April 15 - March 15 (1984).

Progress Report 2

This monthly period was one of minor experimental activity because the student, Daimay Lin, was writing her Master's Thesis and because the postdoctoral fellow, Dr. Singh, accepted a permanent position and left KSU. The new postdoctoral person arrived (and started work) on May 14. Thus, work has begun on rebuilding the $O_2(a)$, $O_2(b)$ apparatus for the $NF(a) + HF(v)$ experiment. Daimay did the final flow calibrations on her $NF(b)$ apparatus this month and has refined the rate constants for $NF(b)$ quenching. Next month's report will contain a listing of all those rate constants. In general there seems to be a close parallel between the trends in $NF(b)$ quenching rate constants and those for $O_2(b)$.

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES AND
NITROGEN FLUORIDE

Contract No: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, KS 66506

Scientific Program Officer: Dr. Vernon Schlie

Contract Duration: March 15, 1984 - March 14, 1985

Monthly Progress Report (~~Second~~ ^{Third} Report)

Report Period: May 15 - June 15 (1984).

Progress Report 3 (May 15 - June 15)

A new student, Andy Slagle, and a new postdoctoral fellow, Dr. Habadas, arrived and started work on the contract. Mr. Slagle will continue the HF(b) quenching measurements and Dr. Habadas will do the NF(a) + HF(v) experiments. Mr. Slagle has learned the techniques and added a few rate constants to our listing (attached) of NF(b) quenching, rate constants. During the next two months he will study NF(b) with halogens. Dr. Habadas rebuilt the O₂(a,b) apparatus so that NF(a) + HF(v) could be studied. The building phase is nearly completed and preliminary experiments will be done during late June. Since this work is difficult and since Dr. Habadas is learning techniques, approximately two more months are expected before reliable data can be obtained.

During the June 15 - Sept. 15 period, Setser will be in Grenoble France studying N₂(A) + CdX₂ (X = Br,I) excitation transfer reactions. Dr. Y. C. Yu will be at KSU during this period and he will be handling phone calls, reports etc. Dr. Yu also will supervise Mr. Slagle in the NF(b) + halogen quenching studies. If any formal administrative action on this contract is required, Dr. Copeland, assistant Dept. Head, should be contacted.

Comparison of Rate Constants at 300K (in units of $10^{-14} \text{ cm}^3 \text{ molec}^{-1} \text{ sec}^{-1}$)
 for the Quenching of $\text{NF}(b^1\Sigma^+)$ and $\text{O}_2(b^1\Sigma_g^+)$

Q	$k_Q^{\text{NF}(b)}$	$k_Q^{\text{O}_2(b)}$
HCl	$30.2 \pm 1.9^{(a)}$	$4 \pm 2^{(d)}$ $7.3 \pm 0.2^{(e)}$ $6.7 \pm 3.5^{(f)}$
HF	$96.7 \pm 15^{(a)}$ $100^{(b)}$	$110 \pm 20^{(d)}$
CH ₄	$22.1 \pm 1^{(a)}$	$7.5^{(f)}$
CH ₃ F	$20.7 \pm 0.3^{(a)}$	
CH ₃ Cl	$23.6 \pm 2.3^{(a)}$	$80 \pm 40^{(d)}$
CH ₃ Br	$17.9 \pm 0.8^{(a)}$	
CH ₃ I	$35.9 \pm 4.2^{(a)}$	
CH ₂ Br ₂	$25.9^{(a)}$	
CHCl ₃	$6.4 \pm 1.7^{(a)}$	
CHF ₃	$63.5^{(a)}$	
CF ₃ NO	$11.6 \pm 1.2^{(a)}$	
CF ₃ Br	$1.20 \pm 0.18^{(a)}$	
NH ₃	$8.91 \pm 0.18^{(a)}$	$200^{(f)}$ $130 \pm 20^{(d)}$
CO ₂	$0.53 \pm 0.41^{(a)}$	$30 \pm 10^{(d)}$
O ₂	$3.8 \pm 0.1^{(a)}$	$0.01^{(g)}$
CO	$1.50 \pm 1.23^{(a)}$	$0.33^{(g)}$
Cl ₂	$7020 \pm 1130^{(a)}$	
Br ₂	$11830 \pm 839^{(a)}$	
NF ₂	$0.5 \pm 0.4^{(a)}$	
N ₂ F ₄	$0.4 \pm 0.1^{(a)}$	
NF ₃	$180 \pm 70^{(c)}$	
Ar	$<0.001^{(c)}$	
H	$300^{(b)}$	

- a. This work.
- b. M. A. Kwok, J.M. Herbelin and N. Cohen, *Electronic Transition Lasers*, p. 8, 2nd Ed., The MIT Press, 1975.
- c. P. H. Tennyson, A. Fintijn and M. A. A. Clyne, *Chem. Phys.*, 62, 171, 1981.
- d. J. P. Singh and D. W. Setser, *J. Phys. Chem.*, to be published.
- e. R. G. O. Thomas and B. A. Thrush, *Proc. Roy. soc. A*, 287, 350, 1977.
- f. M. J. E. Gunthjer and D. R. Snelling, *J. Photochem.*, 4, 27, 1975.
- g. F. Stuhl and K. H. Welge, *Can. J. Chem.*, 47, 1870, 1969.

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U. S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE

Contract No.: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, KS 66506

Scientific Program Officer: Dr. Vernon Schlie

Contract Duration: March 15, 1984 - March 14, 1985

Monthly Progress Report (Fourth Report)


Report Period: June 15 - July 15 (1984)

Progress Report 4


The student, Daimay Lin, who was working on this project, finished her M.S. degree and a new student, Andy Slgale, is able to run her apparatus after a couple weeks' training. Andy is currently extending Daimay's measurements on NF(b) quenching to other interesting molecules.

The initial testing on the NF(a) + HF(v) experiment has been successful. Qualitative work by Dr. Jan Habdas shows that $\text{HF}(v) + \text{NF}(a) \rightarrow \text{NF}(b)$ is observed. An emission spectrum of NF(a), NF(b) and possibly HF(v) is attached to this report. The monochromator calibration is in progress. This is essential before doing any quantitative measurements. Also computer simulations of the kinetic equations are being carried out.


→ Telephone conversation w Dr Habdas 25 JUL 84
 spectrum attached is that produced
 by the reaction



Next step is direct production of HF by




APR 15 1950
MK 0 20
50 A
2.5"




0 655

11

- 5100



NF(b) afterglow



MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE

Contract No.: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Scientific Program Officer: Dr. Vernon Schlie

Contract Duration: March 15, 1984 - March 14, 1985

Monthly Progress Report (Fifth Report)

Report Period: July 15 - August 15 (1984)

Progress Report 5

In collaborative work at Grenoble Dr. Setser has demonstrated that the $\text{CdI}_2 + \text{N}_2(\text{A})$ reaction gives $\text{CdI}(\text{B}) + \text{I}$ with a branching fraction of ~ 0.1 and a total quenching rate constant of $-4.3 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

V-E transfer from $\text{HF}(\text{v})$ to $\text{NF}(\text{a})$ was observed. The preliminary experiments indicated about 30% increase in $\text{NF}(\text{b})/\text{NF}(\text{a})$ ratio upon the addition of $\text{HF}(\text{v})$ to the $\text{H} + \text{NF}_2$ system. Quantitative measurements are in progress.

The spectrum attached to last report was obtained under these conditions:

Total Ar flow	$2.8 \times 10^{-3} \text{ mole/s}$
Total reactor pressure	0.76 torr
Pumping speed	- 82 m/s
H_2 flow	$2.4 \times 10^{-6} \text{ mole/s}$
NF_2 flow	$1.7 \times 10^{-6} \text{ mole/s}$
Slits	400 μm
Scan rate	50 A/min

Experiments on the E-E transfer between $\text{NF}(\text{b})$ and halogens or interhalogens are also being carried out by Mr. Slagle and Dr. Y. C. Yu. A pre-reactor has been constructed to generate $\text{IF}(\text{X})$, etc.

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE

Contract No.: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Scientific Program Officer: Dr. Vernon Schlie



Contract Duration: March 15, 1984 - March 14, 1985

Monthly Progress Report (Sixth Report)


Report Period: August 15 - September 15 (1984)


Progress Report 6

The pre-reactor for in situ synthesis of unstable interhalogens works well for IF(X). E-E transfer from both NF(b) ^{and} N₂(A) to IF(X) has been observed by Dr. Y. C. Yu. IF(B-X) emission spectrum from NF(b) + IF(X) → NF(X) + IF(B) and N₂(A) + IF(X) → N₂(X) + IF(B) are attached. Other E-E transfer reactions observed are summarized below:



No emission can be found for NF(b) + Cl₂, Br₂, ICl, IBr and N₂(A) + Cl₂, Br₂, ICl. For N₂(A) + BrF(X) reaction, definite conclusion can not be made since it is not sure that BrF(X) is generated by the pre-reactor from F + Br₂.





N₂ (G) + IF (G)

10K

450A

W

78
77
76
75
74
73
72
71
70
69
68
67
66
65
64
63
62
61
60
59
58
57
56
55
54
53
52
51
50
49
48
47
46

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE

Contract No.: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Scientific Program Office: Dr. Vernon Schlie



Contract Duration: March 15, 1984 - March 14, 1985

Monthly Progress Report (Seventh Report)

Report Period: Sept. 15 - Oct. 15 (1984)

Progress Report 7

Professor Setser returned from France Sept. 20 and Dr. Yu returned to Taiwan the next day. During this report period, no significant experiments have been done because we are waiting on arrival of N_2F_4 (we have been promised some tanks from Aerospace and a small gift from Hercules). This should be enough for the critical experiments. During this period, we have designed and built second generation flow reactors which should enable the critical experiments to be done for characterizing the following two reactions.



Dr. Yu's last report (#6) was a bit too brief. Indeed, transfer from $NF(b)$ to IF was observed; however, the $IF(B)$ intensity was very low and there may have been traces of $N_2(A)$ in the $NF(b)$ flow system. Thus, we have no firm information about the $NF(b) + IF$ reaction. With good fortune, we may have conclusive results in another month.

During Setser's stay in Grenoble, the $N_2(A)$ reactions with all the Cd and Zn halides were examined. The best system seems to be CdI_2 . The CdI_2 total quenching rate constant was $4.3 \pm 1.5 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and the branching fraction for $CdI(B)$ formation was 0.08. All the other compounds had similar total quenching rate constants, but even smaller branching fractions for excited state product formation. Thus, dissociative excitation transfer from $N_2(A)$ with the Hg, Cd, and Zn halides do not have sufficiently large branching fractions for $HgX(B)$, $CdX(B)$ or $ZnX(B)$ to be very interesting as laser systems. We plan to do some spectroscopic interpretations of the $ZnX(B-X)$ and $CdX(B-X)$ spectra from $N_2(A)$ excitation and write a paper in collaboration with the Grenoble laboratory. However, this is not a high priority item and work will not begin until summer of 1985.

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE


Contract No.: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Scientific Program Office: Dr. Vernon Schlie
Contract Duration: March 15, 1984 - March 14, 1985
Monthly Progress Report (Eighth Report)
Report Period: October 15 - November 15 (1984)



Progress Report #8

During this report period, we have installed the improved flow reactors that will allow the two following reactions to be characterized.



We have received a N_2F_4 tank from Aerospace Corporation and we are expecting additional tanks of N_2F_4 from Hercules Corporation and from the Air Force Weapons Laboratory. We have received CF_3I from the Air Force Laboratory and we have found a commercial source of $\text{C}_2\text{F}_5\text{I}$. These are used to prepare IF from F atom reactions. Thus, all chemicals finally are available for the experiments.

During this report period, Dr. Habadas explored the $\text{F} + \text{H-N}_3$ reaction as an alternative source for NF(a) for study of reaction(i).



This NF(b) source, if fact, appears to be superior to the $\text{H} + \text{NF}_2$ reaction for our purposes. In particular, there appears to be less complication from secondary reactions because excess F with NF(a) causes fewer difficulties than H with NF(a). Based upon our preliminary work, we plan to use reactions (iii) to generate NF(b) for study of reaction (i) and we will not need to use N_2F_4 ($\text{NF}_2 + \text{H}$) to generate NF(a). From our qualitative observation, we believe that reaction (iii-b) must have a rate constant larger than the $2 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ value quoted in the literature. After finishing with reaction (i) we would like to study (iii).

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE

Contract No.: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Scientific Program Office: Dr. Vernon Schlie


Contract Duration: March 15, 1984 - March 14, 1985

Monthly Progress Report (Tenth Report)

Report Period: December 15, 1984 - January 15, 1985

Progress Report # 10

We have continued our experiments to characterize the following two reactions.



Some progress has been made in each case.

The disk-drive for the data system of the FT-IR was returned and new calibrations for the FT-IR system were done last week. Thus, we are ready to observe HF(v) and HF(v-2) via infrared emission from (i). The kinetic modeling studies mentioned in the last report were continued, and we confirmed our suspicion that HF(v=0) from the NF(a) source reaction (either $\text{F} + \text{HN}_3$ or $\text{H} + \text{NF}_2$) was causing kinetic complications because of the quenching of NF(b) by HF(0). Thus, we are trying to develop a NF(a) source that is free of HF(0). We are exploring the $\text{F} + \text{Cl-N}_3$ reaction. The preparation line for ClN_3 has been built and will soon be tested. If this reaction proves to be a source for NF(a), we believe that the assignment of k_v from (i) will be more reliable because the reverse reaction will be less important.

We began the study of (ii) by training the graduate student operator. Stable halogen molecules were selected for investigation. To my surprise the work proved not to be routine and a mistake was discovered in our earlier quenching measurements (by Ms. Lin) for Cl_2 and Br_2 . Those rate constant values, which were included in previous reports and in a preprint, are too large by factors of 2-3. The correct values are

$$k_{\text{Cl}_2} = 1.8 \pm 0.3 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

$$k_{\text{Br}_2} = 6.0 \pm 1.0 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

$$k_{\text{ICl}} = 1.0 \pm 0.3 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

As reported by Dr. Yu in the work done in the summer, these reactions do not give observable Cl_2^* , Br_2^* , or ICl^* emission. These rate constants have been measured several times and the results are reproducible. Apparently Ms. Lin made a mistake when she prepared the Cl_2 and Br_2 mixtures with Ar in her preliminary investigation of Cl_2 and Br_2 . The rate constants for the other reagents reported in the preprint were measured with pure gases and those rate constants are satisfactory. We now are ready to test our method for generating IF and to measure the room temperature quenching rate constant and branching fraction for IF(B) formation. We are conserving N_2F_4 to ensure that we can do the IF experiments. However, we also are proceeding in a conservative step by step fashion to make sure that errors are avoided (or at least identified) in handling the halogens.

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab
Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE


Contract No.: F29601-84-0030

Principal Investigator: D. W. Setser
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Scientific Program Office: Dr. Vernon Schlie
Contract Duration: March 15, 1984 - March 14, 1985
Monthly Progress Report (Eleventh Report)
Report Period: January 15, 1985 - February 15, 1985

Progress Report # 11

We have continued our experiments to characterize the following two reactions:



Some progress has been made in each case.

The attempt to develop a NF(a) source that was free of HF(0) for study of reaction (i) failed. The $\text{F} + \text{Cl-N}_3$ reaction does not cleanly give FCl and N_3 , rather the first step seems to be addition to Cl-N_3 with $\text{NFC1} + \text{N}_2$ being the products. Thus, we have returned to $\text{F} + \text{NH}_3$ as the NF(a) source reaction. We now are collecting data (NF(b) and HF(v) emission intensities) and we will be using these data to assign k_v^b . This project is in the terminal phase and soon will be completed.

We have been testing our method of generating IF for the study of reaction (ii). Preliminary experiments have been done; however, some kinetic difficulties have been discovered. For example, we have found that the microwave discharged Ar/CF₄ mixture gives some quenching of both $\text{N}_2(\text{A})$ and NF(b), although pure CF₄ gives no measurable quenching of either. Therefore, we are exploring different sources for F atoms, since the $\text{F} + \text{CF}_3\text{I}$ reaction is used to generate IF. We have observed the rather weak IF(B-X) emission excited by the NF(b) excitation-transfer reaction. Comparison of the NF(b) and IF(B) emission intensities for a known [IF] suggests an excitation rate constant of $\sim 2-5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. *tentative* Definitive studies can be done to establish this value. Measuring the total quenching rate constant will be more difficult. During the next 4 week period, efforts will be made to obtain quantitative data for the total quenching of (ii), so that a branching reaction for IF(B) formation can be assigned.

me

MONTHLY REPORTS FROM KANSAS STATE UNIVERSITY
TO U.S. AIR FORCE

Air Force Weapons Lab

Kirtland Air Force Base, NM

EXCITED STATE KINETICS OF MERCURY AND CADMIUM HALIDES
AND NITROGEN FLUORIDE

Contract No.: F29601-84-0030


Principal Investigator: D.W. Setser
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Scientific Program Office: Dr. Vernon Schlie
Contract Duration: March 15, 1984 - March 14, 1985 (extended to April 1,
1985)

Monthly Progress Report (Twelfth Report)
Report Period: February 15, 1985 - April 1, 1985

Progress Report # 12

We have continued our experiments to characterize the following two reactions:



Some progress has been made for each case.

We have collected data, NF(b) and HF(v) emission intensities, using F + C₂H₆ as the HF(v) source and F + HN₃ as the NF(a) source for reaction (i). The major experimental effort this month has been development of a method to measure [F] and to more fully characterize the NF(a) source reaction. We have used these data to assign $k_{v=2}^b$ as $\sim 0.5 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. We plan to choose another fuel, probably SiH₄, and generate mainly HF(v=3) so that $k_{v=3}^b$ can be assigned. This project is in the terminal phase and soon will be completed.

The IF(B) excitation and the NF(b) total quenching rate constants have been measured at room temperature using the F + CF₃I reaction to generate IF(X). Comparison of the NF(b) and IF(B) emission intensities for a range of known [IF] given an excitation rate constant of $1.5 \pm 0.5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. The total quenching rate constant for an equal molar mixture of IF(X) and CF₃ was measured by observing the decay of NF(b) in the flow reactor as a function time and [IF]. The total quenching rate constant was $1.1 \pm 0.4 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. This rate constant pertains to an equal concentration of IF and CF₃, but IF is expected to dominate the quenching.

We now are in the process of preparing the final report on this contract, which terminated on April 1, 1985. The two week extension from March 15 to April 1 was granted by the Chicago Monitoring Office for administrative reasons.