

AD-A244 811



AD _____

✓
①

GRANT NO: DAMD17-91-Z-1023

TITLE: ADJUVANT ACTION OF LIPOPOLYSACCHARIDES AND DERIVATIVES

PRINCIPAL INVESTIGATOR: Jon A. Rudbach, Ph.D.

CONTRACTING ORGANIZATION: Ribi Immunochem Research, Inc.
P.O. Box 1409
Hamilton, MT 59840

REPORT DATE: September 30, 1991

DTIC
SELECTE
JAN 17 1992
S B D

TYPE OF REPORT: Final Proceedings

PREPARED FOR: U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

1991 September 30 Final Proceedings (4/1/91 - 9/30/91)

ADJUVANT ACTION OF LIPOPOLYSACCHARIDES AND DERIVATIVES

Subtitle: Newer Aspects of the Adjuvant Action of Lipid
A and Its Analogs

Grant No.
DAMD17-91-Z-1023

Jon A. Rudbach, Ph.D.

Ribi Immunochem Research, Inc.
P.O. Box 1409
Hamilton, MT 59840

U.S. Army Medical Research and Development Command
Fort Detrick
Frederick, Maryland 21702-5012

Published proceedings of the Conference which took place April 6-9, 1991,
in Airlie, Virginia

Approved for public release; distribution unlimited


Conference, RA I

Unclassified

Unclassified

N/A

N/A



Newer Aspects
of the
Adjuvant Action
of
Lipid A
and
Its Analogs

Airlie Conference Center • Airlie, Virginia • April 6-9, 1991


92-01510



92 1 16 080

Sponsored By:

*Vaccine Research and Development Branch, Division of AIDS
National Institute of Allergy and Infectious Diseases, NIH,
and
Walter Reed Army Institute of Research,
U.S. Army Medical Research and Development Command*



Newer Aspects
of the
Adjuvant Action
of
Lipid A
and
Its Analogs

Airlie Conference Center • Airlie, Virginia • April 6-9, 1991

Sponsored By:

*Vaccine Research and Development Branch, Division of AIDS
National Institute of Allergy and Infectious Diseases, NIH,
and
Walter Reed Army Institute of Research,
U.S. Army Medical Research and Development Command*



Contents

Agenda.....	1
Keynote Address	9
Abstracts	
Session I.....	11
Session II.....	21
Evening Workshop.....	31
Session III.....	33
Session IV.....	45
Evening Lecture.....	59
Session V	63
List of Participants	73
Notes.....	93

Program Committee

Dr. Carl R. Alving	Walter Reed Army Institute of Research
Dr. Phillip J. Baker	National Institute of Allergy and Infectious Diseases
Dr. Barbara Detrick	National Institute of Allergy and Infectious Diseases
Dr. Arthur G. Johnson	University of Minnesota
Dr. Jon A. Rudbach	RIBI ImmunoChem Research, Inc.
Dr. Stefanie N. Vogel	Uniformed Services University of Health Sciences

Note: These abstracts are considered privileged communications and should not be used for citation.



Agenda

**NEWER ASPECTS OF THE ADJUVANT ACTION
OF LIPID A AND ITS ANALOGS**

**Airlie Conference Center
Airlie, Virginia
April 6-9, 1991**

AGENDA

Saturday, April 6

12:00 noon - 6:00 p.m.	Registration	
7:25 p.m. - 7:30 p.m.	Welcome Notes	Dr. Arthur G. Johnson
7:30 p.m. - 8:15 p.m.	Keynote Address <i>Current Developments in AIDS Vaccine Adjuvant Research</i>	Dr. Wayne C. Koff
8:30 p.m.	Reception	

Sunday, April 7

8:30 a.m. - 12:10 p.m.	Session I New Advances in the Structural and Functional Relationships of Lipid A and Its Analogs (Part I)	<i>Chairperson:</i> Dr. Ernst Rietschel
8:30 - 8:50	<i>Overview</i>	Dr. Ernst Rietschel
8:50 - 9:35	<i>Lipid A Analogs Aiming at Prevention of the Deleterious Effects of Endotoxin: Structure- Activity Relationships</i>	Dr. Peter Stuetz
9:35 - 10:20	<i>Biological Activities of Nonpyrogenic Synthetic Lipid A Analogues with Monosaccharide Backbone</i>	Dr. Motohiro Matsuura
10:20 a.m. - 10:40 a.m.	Coffee Break	
10:40 - 11:25	<i>Current Status of the Structure- Function Relationships of Lipid A</i>	Dr. Kuni Takayama
11:25 - 12:10	<i>Regulation of Nitric Oxide Synthesis by Lipid A Derivatives</i>	Dr. Gary L. Gustafson
12:15 p.m. - 1:30 p.m.	Lunch	

Sunday, April 7 (continued)

1:30 p.m. - 4:50 p.m.	Session II New Advances in the Structural and Functional Relationships of Lipid A and Its Analogs (Part II)	<i>Chairperson:</i> Dr. Alois Nowotny
1:30 - 2:15	<i>Over Expression of KDO Transferase Activity</i>	Dr. Christian R.H. Raetz
2:15 - 3:00	<i>Modulation of Antibody Isotype by LPS and Nonionic Block Copolymers</i>	Dr. Robert L. Hunter
3:00 p.m. - 3:20 p.m.	Coffee Break	
3:20 - 4:05	<i>Lipid A-Like Molecules Antagonize the Effects of Endotoxins on Human Monocytes and Neutrophils</i>	Dr. Douglas Golenbock
4:05 - 4:50	<i>Host Responses to LPS and Bacteria</i>	Dr. Christopher Galanos
7:30 p.m.	Evening Workshop Implications and Consequences of Lipid A Administration	<i>Chairperson:</i> Dr. Carl R. Alving

Monday, April 8

8:30 a.m. - 12:10 p.m.	Session III Cellular Mechanisms and Cytokines	<i>Chairperson:</i> Dr. Phillip J. Baker
8:30 - 9:10	<i>Inactivation of Suppressor T Cell Activity with MPL and Its Analogs</i>	Dr. Phillip J. Baker
9:10 - 9:50	<i>Immunoregulation by Lipid A Analogs</i>	Dr. Arthur G. Johnson
9:50 - 10:30	<i>Early Endotoxin Tolerance: A Model for Differential Cytokine Induction and Gene Expression by LPS and MPL</i>	Dr. Stefanie N. Vogel
10:30 a.m. - 10:50 a.m.	Coffee Break	
10:50 - 11:30	<i>Induction and Modulation of IL-1, IL-6, and TNF by LPS and Its Analogs</i>	Dr. Hans-Dieter Flad
11:30 - 12:10	<i>LPS Processing by Phagocytes: An Update</i>	Dr. Robert S. Munford
12:10 p.m. - 1:15 p.m.	Lunch	
1:30 p.m. - 5:10 p.m.	Session IV Non-Specific Increases in Resistance	<i>Chairperson:</i> Dr. Richard A. Proctor

Monday, April 8 (continued)

1:30 - 2:10	<i>Role of Purinoreceptor in Endotoxin Activation of Macrophages</i>	Dr. Richard A. Proctor
2:10 - 2:50	<i>Identification and Characterization of LPS Binding Proteins on Mammalian Cells</i>	Dr. Mei-Guey Lei
2:50 - 3:30	<i>Protective Effects of MPL in Models of Sepsis and Endotoxemia</i>	Dr. J. Terry Ulrich
3:30 p.m. - 3:50 p.m.	Coffee Break	
3:50 - 4:30	<i>MPL as Prophylaxis for Surgical Wound Infection: A Phase I-II Study</i>	Dr. David Beatty
4:30 - 5:10	<i>Induction of Endotoxin Tolerance by Non-Toxic Derivatives of Lipid A</i>	Dr. Mark E. Astiz
7:30 p.m.	Evening Lecture <i>Molecular Mechanisms of Intracellular Signalling</i>	Dr. Masayasu Nakano

Tuesday, April 9

8:30 a.m. - 12:10 a.m.	Session V Current Vaccine Strategies Using Lipid A Derivatives	<i>Chairperson:</i> Dr. Barbara Detrick
8:30 - 8:50	<i>Overview</i>	Dr. Barbara Detrick
8:50 - 9:35	<i>Development of Synthetic Lipid A Analogue Vaccine</i>	Dr. Alan S. Cross
9:35 - 10:20	<i>Adjuvant Effects of Liposomal Lipid A: Human Vaccine Induces Antibodies to Both Malaria Antigen and Lipid A</i>	Dr. Carl R. Alving
10:20 a.m. - 10:40 a.m.	Coffee Break	
10:40 - 11:25	<i>An Effective Vaccine for Syphilis: The Importance of Adjuvant Action of MPL in Reversing Immunosuppression</i>	Dr. Thomas Fitzgerald
11:25 - 12:10	<i>Evaluation of MPL as Adjuvant for Human Immunization: Enhanced Antibody Responses with Concurrent Injection of MPL and Polysaccharide Conjugates</i>	Dr. Sarvamangala Devi



Keynote Address

Current Developments in AIDS Vaccine Adjuvant Research

Saturday, April 6 • 7:30 p.m. - 8:00 p.m.

Dr. Wayne C. Koff

National Institute of Allergy and Infectious Diseases, NIH • Bethesda, MD



Session I

New Advances in the Structural and Functional Relationships of Lipid A and Its Analogs (Part I)

Sunday, April 7 • 8:30 a.m. - 12:00 noon

Speakers' Summaries (in order of presentation)

LIPID A ANALOGS AIMING AT PREVENTION OF THE DELETERIOUS EFFECTS OF ENDOTOXIN: STRUCTURE-ACTIVITY RELATIONSHIPS.

P. Stuetz, SANDOZ Forschungsinstitut, Vienna, Austria.

Despite the advent of potent new antibiotics with a broad spectrum of activity, overall morbidity and lethality in patients with Gram-negative bacteremia have not changed over the years. At present, two different approaches are being pursued with lipid A analogs to prevent the deleterious effects of endotoxin whose dominant role in septic shock has been well established.

Highly lipophilic lipid A analogs, i.e. monosaccharidic or disaccharidic glucosamine-phosphates that contain at least 3 N,O-bound long-chain fatty acids per glucosamine-phosphate backbone have the ability to enhance non-specific resistance to bacterial and fungal infections, presumably by stimulating the secretion of colony stimulating factors, and by priming for oxidative burst and bacterial killing of macrophages and neutrophils. These events are paralleled by the development of so-called early-phase tolerance to endotoxin, a reversible state of hyporesponsiveness against otherwise lethal doses of LPS. Paradoxically, repeated administration of this class of compounds potentiates resistance to infections, but also tolerance to endotoxin even further and for a longer period of time. From the limited information available it appears that this class of compounds, like LPS or lipid A are also good adjuvants. Possible modes of action underlying these effects will be discussed.

The second class represent much more polar monosaccharidic or disaccharidic lipid A analogs with a maximum of 2.5 N,O-bound long-chain fatty acid residues per glucosamine-phosphate which can directly block LPS-effector mechanisms, presumably by competing with LPS-binding sites. These compounds also protect against a lethal LPS-dose in rodents. Their adjuvant effects have not been investigated so far, but in the prolymphoid 70/32 cell line, prototype compounds are able to effectively block the LPS-induced differentiation into mature B cells.

Synthetic lipid A analogs with monosaccharide backbone,
which are nonpyrogenic with various biological activities

Motohiro Matsuura¹, Makoto Kiso² and Akira Hasegawa²

¹Jichi Medical School, Tochigi, Japan and ²Gifu University, Gifu, Japan.

A typical lipid A (Escherichia coli type) is composed of glucosamine disaccharide backbone with 1,4'-bisphosphate and hexaacyl substituents at the 2, 2', 3 and 3' positions. We have synthesized lipid A analogs with monosaccharide backbone carrying phosphate and acyl substituents, and their biological activities examined. Analogs possessing monophosphate and triacyl substituents to monoglucosamine backbone were generally active. A representative analog (GLA-60), which exhibits various immunomodulating activities such as induction of mediators, activation of B cells or macrophages, enhancement of non-specific resistance and so on, has a 3-hydroxytetradecanoyl ($C_{14}-OH$), a 3-tetradecanoyloxytetradecanoyl ($C_{14}-O-(C_{14})$) and a phosphate groups at the 2, 3 and 4 positions of a glucosamine backbone, respectively. An analog GLA-63 with a $C_{14}-O-(C_{12})$ group instead of a $C_{14}-O-(C_{14})$ group in GLA-60 exhibited as high activities as GLA-60, but the activities of GLA-64 with a $C_{14}-O-(C_{16})$ group diminished markedly. Moreover, analogs with a tetradecanoyl (C_{14}) and a $C_{14}-O-(C_{16})$ groups as acyl substituents at the 2 and 3 positions (GLA-69 and GLA-58) were nearly nonactive. Tolerance inducing activity against lethal toxicity of LPS was, however, observed even in such scarcely active analogs. The most interesting point in biological activities of the synthetic monosaccharide analogs is that neither pyrogenicity nor Shwartzman reactivity is exhibited even by the highly active analogs such as GLA-60 and GLA-63.

CURRENT STATUS OF THE STRUCTURE-FUNCTION RELATIONSHIP OF LIPID A

Kuni Takayama

Mycobacteriology Research Laboratory
William S. Middleton Memorial Veterans Hospital
Madison, Wisconsin 53705

and
Department of Bacteriology, CALS
University of Wisconsin
Madison, Wisconsin 53706

The endotoxic moiety of the lipopolysaccharide (LPS) of Gram-negative bacteria is the lipid A. Since its "correct" structure was elucidated in 1983, many investigators have studied the structural basis for the numerous biological activities of lipid A. This included chemical synthesis of analogs. From all these studies, one can reach a simple conclusion that the most endotoxic lipid A structure is the hexaacyl diphosphoryl lipid A found in the LPS of the *Salmonella* strains and *Escherichia coli* [$\beta(1\rightarrow6)$ glucosamine disaccharide containing phosphate groups at the 1- and 4'-positions, HOC_{14} fatty acids at the 2- and 3-positions, $\text{C}_{12}\text{OC}_{14}$ at the 2'-position, and $\text{C}_{14}\text{OC}_{14}$ at the 3'-position]. We call this the model "toxic" lipid A. A departure from this basic structure leads to lower biological activities. Nature modulates the biological activities of this toxic lipid A by controlling (i) the number of phosphate groups in the disaccharide, (ii) the number, kind, and size of the hydroxy and normal fatty acids in the disaccharide, and (iii) the polar substituents to be added to the basic structure. Another variation that nature has devised is to use 2,3-diamino-2,3-dideoxyglucose in the disaccharide backbone.

A model "nontoxic" lipid A can be obtained from the LPS of *Rhodopseudomonas sphaeroides*. It is closely related structurally to the model toxic lipid A. We will show that by merely reducing the fatty acid content from six to five and lowering the size of the hydroxy fatty acids at the 3- and 3'-positions from C_{14} to C_{10} , the toxic lipid A is converted to nontoxic lipid A. These results emphasize the strong influence of specific hydroxy fatty acids in the lipid A moiety of LPS in determining its endotoxic property.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

REGULATION OF NITRIC OXIDE SYNTHESIS BY LIPID A DERIVATIVES

G.L. Gustafson and M.J. Rhodes
Ribi ImmunoChem Research, Inc., Hamilton, MT 59840

Recent studies in several laboratories have shown that LPS can induce the expression of nitric oxide (NO) synthetase in macrophages as well as in tissue cells of animals. Expression of this enzyme in macrophages is a protective response since macrophage production of NO enhances antimicrobial and antitumor activities. However, expression in tissue cells is detrimental since tissue NO is a mediator of endotoxin shock. In the present work the potentials of LPS and lipid A derivatives to regulate NO synthesis were studied 1) in cultured murine peritoneal macrophages (macrophage model) and 2) in intact mice (systemic model). NO production was quantified indirectly by measuring the accumulations of nitrite and nitrate in cell culture media and in blood samples. Lipid A derivatives were compared with regard to their activities to stimulate NO production and to block LPS-stimulated NO production. The regulation of NO synthesis in the systemic model differed in many ways from the regulation of macrophage NO synthesis. These included 1) differences in structure/activity relationships for lipid A derivatives between the two models, 2) differences in the kinetics of NO production, and 3) differences in the patterns of blocking of NO production by lipid A derivatives. Of special interest, it was found that monophosphoryl lipid A (MPL) did not enhance systemic production of NO, but that it did induce NO synthesis in cultured macrophages. Based on all of the differences observed, it was hypothesized that tissue cells rather than macrophages might be the major producers of NO in mice challenged systemically with LPS. The demonstrations that pretreatment of mice with MPL attenuated the ability of LPS to induce systemic NO production but did not itself induce NO production supported the view that MPL might have applications as a prophylactic drug for endotoxic shock.



Session II

New Advances in the Structural and Functional Relationships of Lipid A and Its Analogs (Part II)

Sunday, April 7 • 1:20 p.m. - 4:00 p.m.

Speakers' Summaries (in order of presentation)

OVEREXPRESSION OF KDO TRANSFERASE ACTIVITY. C. R. H. Raetz. Merck Sharp and Dohme, Rahway, NJ.

An autoradiographic assay applicable to colonies immobilized on filter paper was developed for obtaining temperature sensitive mutants of *Escherichia coli* defective in the transfer of 3-deoxy-D-manno-octulosonic acid (KDO) from CMP-KDO to a tetraacyldisaccharide-1,4'-bisphosphate precursor of lipid A, designated lipid IV_A. Cell-free extracts from two mutants found in a population of 30,000 mutagen treated cells showed normal KDO transferase activity when assayed at 30°C, but almost no activity at 42°C. The mutation was mapped by mating one of the mutants with different Hfr strains and analyzing genetic linkage of KDO transferase activity to selectable markers. The lesion was located to a position between 80-84 min on the *E. coli* chromosome. A plasmid from the Clarke and Carbon collection, pLC17-24, known to contain genes from the *rfa* region (81 min), was shown to overexpress KDO transferase activity 4-5 times and to correct the mutation when the plasmid was conjugated into the mutant strains. The KDO transferase gene, designated *kdtA*, was subcloned from pLC17-24 into a multicopy vector. The resulting plasmid, pCL3, overproduced transferase activity -100-fold. The *kdtA* gene was shown to code for a 43 kDa polypeptide, as judged by radiolabeling of minicells. Its DNA sequence was determined. The results demonstrate that overexpression of this single gene product greatly stimulates the incorporation of two stereochemically distinct KDO residues during LPS biosynthesis in extracts of *E. coli*.

Modulation of Antibody Isotype by Lipopolysaccharides in Combination With Nonionic Block Copolymers

Robert L. Hunter
Department of Pathology,
Emory University,
Atlanta, GA 30322

Kuni Takayama
Mycobacteriology Laboratory,
William S. Middleton Memorial Veterans Hospital,
Madison, WI 53705;

Nonionic block copolymers and lipopolysaccharides are both effective immunologic adjuvants which are thought to act via distinct mechanisms. We hypothesized that they might produce synergistic effects when used together. We prepared a series of LPS preparations ranging from the smallest precursor, lipid X through complete LPS with O-polysaccharide chains. Three preparations with reduced toxicity, monophosphoryl lipid A, partially hydrolyzed Ra-LPS and LPS of *R. sphaeroides* were also utilized. All LPS preparations except the smallest were effective adjuvants for inducing early antibody responses to TNP-HEA when injected in squalane-in-water emulsions with copolymer L141. Only the larger LPS preparations induced sustained antibody responses. By itself, emulsions of copolymer L141 induced a predominant IgG1 antibody isotype response with lesser amounts of IgG2a and IgG2b. Surprisingly, all of the LPS preparations tested increased the proportion of IgG2 isotypes even though some had little effect on overall titers. The detoxified Ra-LPS (Ra-detox) was the most effective preparation for both increasing antibody titers and inducing the desirable IgG2a and IgG2b isotypes. These results demonstrate that the combination of LPS and block polymer adjuvants can produce synergistic effects without unacceptable toxicities.

Lipid A-like Molecules Antagonize the Effects of Endotoxins on Human Monocytes and Neutrophils. Douglas Golenbock, M.D., Boston City Hospital, Boston MA. Lipopolysaccharide (LPS) is implicated as the bacterial product responsible for the clinical syndrome of gram-negative bacterial septicemia. Although the lipid A domain of LPS appears to be responsible for the toxicity of endotoxin, lipid A from the photosynthetic bacterium *Rhodobacter sphaeroides* (RSLA), two disaccharide precursors of lipid A from enteric bacteria termed lipid IV_A and KDO₂-lipid IV_A, and the monosaccharide lipid A precursor lipid X have little stimulatory activity on human cells. Using human mononuclear and polymorphonuclear phagocytes, we have shown that these molecules are specific antagonists of LPS. Complete, apparently competitive, inhibition of LPS activity of both monocytes and neutrophils was possible at a 10-fold excess of the disaccharide antagonists, as judged by measuring the release of cytokines and PGE₂ from monocytes or the surface expression of CD11b/CD18 (CR3). The LPS inhibitors had no effect on the stimulation of monocytes by purified protein derivative from *Mycobacterium tuberculosis* or heat-killed *Staphylococcus aureus* nor on neutrophil CR3 upregulation by TNF α , formyl-met-leu-phe or phorbol myristate acetate. Lipid X had no activity when tested against monocytes (up to a concentration of 1 μ g/ml) and did not inhibit LPS-induction of TNF α or PGE₂ release. At high concentrations (10 μ g/ml), lipid X acted as an antagonist when tested against LPS-induced PMN responses. The effect of the disaccharide inhibitors could not be removed when treated monocytes or neutrophils were extensively washed and then exposed to LPS. In contrast, the effects of lipid X could easily be washed away from treated PMN suggesting either that lipid X rapidly dissociates from neutrophil LPS receptors or inhibits LPS through non-competitive means (such as sequestration of LPS). The lipid A-based inhibitors may be useful in the design of novel therapeutic agents for gram-negative bacterial septicemia.

Host-Responses to LPS and Bacteria.

C.Galanos and M.A.Freudenberg

Max-Planck-Institut für Immunbiologie, 7800 Freiburg, FRG

The lethal toxicity of LPS is mediated by macrophages through tumor necrosis factor α (TNF α) which is probably the primary mediator of the lethal activity of LPS. LPS lethality may be completely inhibited by anti-TNF α antibodies. Smooth and rough (Ra-Re) LPS as well as free enterobacterial lipid A induce lethal effects in D-galactosamine treated mice. Synthetic hexaacylated lipid A (506) and monophosphorylated tetraacylated lipid A precursor Ia (406) were also active. The ability of synthetic compounds 506 and 406 to stimulate macrophages to produce TNF α in vitro was determined and compared to that of LPS. The experiments revealed that all 3 agents exhibited similar effects in cultures of mouse macrophages. However only LPS, but not compound 506 or 406, stimulated TNF α in cultures of human macrophages. It indicates that different structural prerequisites may be required in different animal species and/or different macrophage populations may respond differently to different LPS substructures. Like lethality also the tolerance-inducing properties of LPS are mediated by macrophages through TNF α . Pretreatment of mice with minute amounts of LPS or TNF α will protect them to a subsequent challenge with lethal amounts of either agent in D-GalN sensitized mice. TNF α is the mediator of the lethal activity of gram-negative and gram-positive bacteria. While several mg of purified LPS will not kill D-GalN treated LPS-resistant mice, a few μ g of the same bacteria do so very effectively. This is evidence for the presence of shock inducing components, other than LPS, in gram-negative bacteria. Killed gram-positive bacteria also exhibit lethal effects in D-GalN sensitized LPS-sensitive and -resistant mice. The lethal activity of gram-negative and -positive bacteria may be inhibited by anti-TNF α antibodies. This is evidence that TNF α is a common mediator of the lethal activity of gram-negative and -positive bacteria. Minute amounts of both types of bacteria when used to pretreat mice, induce non-specific resistance to LPS and to their own lethal effects, and vice versa.



Evening Workshop

Implications and Consequences of Lipid A Administration

Sunday, April 7 • 7:30 p.m.



Session III

Cellular Mechanisms and Cytokines

Monday, April 8 • 8:30 a.m. - 12:00 noon

Speakers' Summaries (in order of presentation)

Inactivation of Suppressor T Cell (Ts) Activity by Monophosphoryl Lipid A (MPL) and its Structural Analogs (P.J. Baker)

Previous studies have shown that treatment with MPL abrogates the expression of Ts activity without adversely effecting other T cell functions (e.g., helper and amplifier T cell activity). LPS-responsive (LPS^r) and LPS-defective (LPS^d) strains of C3H mice do not differ in their capacity to make an antibody response to Type III pneumococcal polysaccharide (SSS-III) or in the degree of Ts activity generated after exposure to this antigen; however, treatment with MPL abolishes the expression of Ts function in LPS^r -- but not in LPS^d -- mice. This suggests that these mice may differ with respect to the presence of (a) a cell-surface receptor needed for the specific binding of MPL to activated Ts, and/or (b) a unique biochemical pathway in Ts that is sensitive to MPL. Several synthetic analogs of lipid A were tested to obtain relevant information on this issue. The results obtained suggests that the minimal structural unit required to produce the effects described is a diglucoseamine to which must be attached at least 5 fatty acids with perhaps one of them being an acyloxyacyl fatty acid at the 2' or 3' position. The ability of MPL or its analogs to inactivate Ts function appears to be independent of its ability to polyclonally activate B cells and the number of phosphate groups present.

IMMUNOREGULATION BY LIPID A ANALOGS. Arthur G. Johnson, Marilyn Odean, and Akira Hasegawa. University of Minnesota, Duluth, MN and Gifu University, Gifu, Japan.

LPS can either enhance or suppress the immune response depending on its time of administration relative to antigen. Our current research is directed towards (a) determining the minimal structural requirements responsible for enhanced antibody titers and/or their down regulation, and (b) identifying the target cell(s) and mediator(s) initiating both of these regulatory phenomena. In earlier studies with monophosphoryl lipid A (Ribi) we have implicated a Th₁ cell as the initial target cell and secretion of interferon gamma as vital to the enhancing action. In addition, suppression of antibody was also induced by MPL when given before antigen. Thus, neither KDO nor the reducing sugar phosphate is required for these actions.

Recently, a number of glycolipids representing the non-reducing moiety of lipid A have been synthesized and characterized by Hasegawa, et al. We have initiated testing of three of these analogs for their ability to suppress PFC formation when given before antigen. Our findings to date indicate that 25 µg of GLA 27 and GLA 60 injected ip effectively suppressed the murine PFC response. However compound 47, was ineffective, even at twice the dosage. Examination of their structural differences revealed that addition of a C₁₄-O-acyl group to the C₁₄ fatty acid at the R'3 site to GLA 27 (which then becomes GLA 47) inactivated the capacity to suppress PFC formation. However, keeping the O-acyl linkage on R'3 but removing it from the C₁₄ fatty acid on R'2 (now GLA 60) once again permitted suppression to be expressed. No significant pattern emerged from the suppression induced by three other analogs.

EARLY ENDOTOXIN TOLERANCE: A MODEL FOR DIFFERENTIAL
CYTOKINE INDUCTION AND GENE EXPRESSION BY LPS AND MPL.

Stefanie N. Vogel and Beth E. Henricson, Uniformed Services
University of the Health Sciences, Bethesda, MD 20814 USA.

Previous studies have shown that both smooth lipopolysaccharide (LPS) and the non-toxic, lipid A derivative, monophosphoryl lipid A (MPL), are capable of inducing a state of "early endotoxin tolerance" (EET) in mice; however, one needs significantly more MPL than LPS to achieve a level of desensitization comparable to that induced by LPS. Since cytokines have been implicated in the induction of EET, we examined levels of cytokines induced by LPS vs. MPL and found that although comparable levels of CSF were induced by tolerizing doses of each, the levels of other cytokines (e.g., TNF, IL-6, and IFN) are markedly reduced in animals injected with MPL. An in vitro model of LPS desensitization was established to dissect the problem further. Briefly, the macrophage cell line, RAW264.7, or thioglycollate-elicited peritoneal exudate macrophages were exposed first to medium, LPS, or MPL for 20 hr and then "challenged" with LPS for an additional 20 hr and the level of TNF produced in the supernatants measured. As was observed in vivo, significantly more MPL was required to induce the same level of desensitization induced by LPS. Finally, induction of transcription of "early" endotoxin-induced genes was compared in macrophages treated with LPS vs. MPL. For seven genes examined, the concentration of MPL required to induce transcription of LPS-induced "early" gene expression ranged from 20- to 2500-fold that of LPS, depending on the individual RNA species analyzed. This approach should provide significant insights into the mechanisms by which LPS and MPL transmit their respective signals and result in toxic vs. beneficial effects in the host. (Supported by USUHS R07338 and NMRDC 63706.0095.001).

Induction and modulation of IL-1, IL-6, and TNF production by LPS and its partial structures

H.-D.Flad, Department of Immunology and Cell Biology, Forschungsinstitut Borstel, D-2061 Borstel, FRG

Bacterial lipopolysaccharides (LPS) consist of a hydrophilic heteropolysaccharide component and a covalently linked hydrophobic lipid portion termed lipid A. In studying the structure-function relationship of LPS and lipid A partial structures it could be established that S- and R-form LPS are very potent inducers of IL-1 and TNF in human monocytes in vitro, while synthetic *S.minnesota* heptaacyl lipid A (compound 516) and the monophosphorylated partial structures (compounds 504 and 505) were less active. Evidence is presented that the synthetic tetraacyl partial structure of *E.coli* lipid A (precursor Ia, compound 406) and the monosaccharide structure (lipid X, compound 401) do not induce IL-1, IL-6, and TNF, but modulate LPS-induced cytokine production. These compounds are able to suppress and/or delay LPS-induced TNF and IL-1 β in a concentration-dependent fashion. Compound 406 is not able to inhibit phorbol-12-myristate-13-acetate (PMA)-induced TNF and IL-1 release, which suggests that its modulating effect is LPS-specific. Evidence is presented which indicates that the modulating effect of compound 406 on LPS-induced cytokine production is due to a competition of binding.

LPS PROCESSING BY PHAGOCYTES -- AN UPDATE. R. S. Munford. University of Texas Southwestern Medical Center, Dallas, TX.

Phagocytes process LPS principally by hydrolyzing the fatty acyl chains and phosphates from lipid A, leaving the saccharide chain (O-antigen) largely intact. Of the catalytic enzymes known to take part in LPS degradation by animal phagocytes, acyloxyacyl hydrolase (AOAH) has been most intensively studied. This enzyme removes the secondary acyl chains from LPS, producing a lipid A moiety that resembles the LPS precursor IVa and synthetic analog 406. In keeping with the reported bioactivities of these lipid A partial structures, enzymatically deacylated S. typhimurium, E. coli, P. aeruginosa, and H. influenzae LPSs are non-toxic (e.g., unable to stimulate human vascular endothelial cells) yet retain other biological activities, including murine spleen cell mitogenicity. Interestingly, Neisseria LPS is unable to elicit murine spleen cell mitogenesis and will inhibit the stimulatory ability of other LPSs in this assay system. It therefore appears that the contribution of the lipid A secondary acyl chains to the bioactivity of a given LPS cannot be predicted from the bioactivities of the existing lipid A analogs; possible explanations for this observation will be discussed.



Session IV

Non-Specific Increases in Resistance

Monday, April 8 • 1:30 p.m. - 4:45 p.m.

Speakers' Summaries (in order of presentation)



"Role of the Purinoreceptor in Endotoxin
Activation of RAW264.7 Macrophages".

T. TANKE, H. RHIM, J-W. VAN DE LOO, P.S.

LEVENTHAL, R.A. PROCTOR,* and P.J. BERTICS. Univ. of
Wisc. Medical School, Madison, WI.

Studies of LPS-induced release of toxic mediators from macrophages have implicated G-proteins on the basis of pertussis toxin (PT) inhibition of IL-1 release. However, direct studies of G-protein activity have not been reported. Hence, we isolated membranes from RAW264.7 macrophages and studied GTPase activity as a measure of G-protein activation. Kinetic analyses showed maximal LPS-induced increase (3-fold) of velocity within 15 min that is enhanced by ATP, ADP, AMPPNP, ammonium sulfate, but is PT insensitive, even though PT was able to ribosylate a 40kDa protein. However, ADP doses that inhibit ATPase activity stimulated LPS-induced GTPase activity. Lipid X, which reduces LPS-lethality, dose-dependently blocked LPS-activation of the GTPase in the presence of ATP. These effects appear specific as other lipids, detergents, and glycosides neither stimulated nor inhibited LPS-induced GTPase activity. Thus, LPS may activate macrophages via coupling a G-protein to a membrane receptor, and this interaction may be modulated by ATP. As ATP infusion causes shock and death, LPS might cause toxicity by sensitizing cells to ambient ATP levels.

IDENTIFICATION AND CHARACTERIZATION OF LPS BINDING PROTEINS ON MAMMALIAN CELLS. M.-G. Lei, T.-Y. Chen, J. Halling, S. Field, L. Flebbe and D. C. Morrison. Univ. of Kansas Medical Center, Kansas City, KS 66103.

Studies from our laboratory have focused upon the identification and characterization of endotoxic lipopolysaccharide receptors expressed on mammalian lymphoreticular cells. These experiments have employed a photoactivatable disulfide reducible radioiodinated derivative of LPS synthesized by our laboratory (Wollenweber and Morrison, J. Biol. Chem., 260, 15068, 1985). Using S-LPS from *E. coli* 0111:B4 we identified a membrane glycoprotein of approximately 80 kDa on mouse lymphocytes and macrophages which manifest a number of properties consistent with its functioning as a specific LPS receptor. Our studies suggested that binding specificity for this 80 kDa protein was for the lipid A region. We have recently confirmed this hypothesis by the synthesis of photoderivatized LPS from the deep rough *S. minnesota* Re chemotype LPS, where binding to an 80 kDa protein can readily be demonstrated. Extension of these studies to selected subpopulations of human peripheral blood cells has confirmed the existence of similar 80 kDa proteins on polymorphonuclear leukocytes, monocytes, B and T lymphocytes and platelets. These proteins are not detectable on human erythrocytes. Competitive inhibition studies readily demonstrate inhibition of binding to the 80 kDa protein with a variety of LPS and lipid A, including that from R.1 spheroides but not with a variety of peptidoglycans. We have recently begun characterization of a second LPS binding protein of approximate molecular mass 38 kDa. Of interest, photoderivatized binding of LPS to this protein is inhibited by both S and R chemotype LPS but not by purified Lipid A. The functional role of this 38 kDa protein remains to be defined (Supported by AI 23447).

Protective Effects of Monophosphoryl Lipid A in Models of Sepsis and Endotoxemia

J.T. Ulrich, Y.M. Hudson, and J.R. Ward
Ribi ImmunoChem Research, Inc., Hamilton MT 59840

Monophosphoryl lipid A (MPL) and 3-O deacylated MPL (3D-MPL) are prepared by partial degradation of Re LPS from Salmonella minnesota R595. These materials are greatly attenuated with respect to their toxic activities. A major focus in our laboratory has been to evaluate the ability of these substances to enhance non-specific resistance to bacterial challenge and to protect against lethal endotoxemia. Pretreatment of mice (I.P.) with 25-50 µg MPL or 3D-MPL results in maximal protection (80-100% survival) to Escherichia coli (5×10^8 CFU) given 48 hr. later. Increased non-specific resistance in this model was temporary. Less than 10% survival was observed if the challenge was given 5 days after MPL or 3D-MPL treatment. Experiments in which mice received multiple treatments with 3D-MPL prior to challenge indicated that a minimum of 6 days was required before re-treatment could elicit maximum protection. D-galactosamine (D-gal) treatment profoundly sensitizes mice to endotoxin/LPS. Using the D-gal model we have found that 3D-MPL pretreatment (I.V.) produces a hyporesponsive state to LPS that occurs within 6 hrs. and which persists for 48 hrs., with protection rapidly declining by 72 hrs. Multiple dosing studies with 3D-MPL given daily or every other day demonstrated that this hyporesponsive state to LPS challenge could be extended for up to 7 days. Tumor necrosis factor (TNF α) production by macrophages in response to LPS stimulation has been shown to mediate lethality in D-gal compromised mice. Changes in TNF α concentrations in mouse sera were measured following D-gal/LPS challenge. Mice pretreated with 3D-MPL produced only 30% as much serum TNF α as control mice when challenged with LPS. D-gal treatment of mice has also been shown to increase their sensitivity to the direct toxic/lethal effects of human recombinant TNF α (hrTNF α). We found that 3D-MPL pretreatment abrogates the toxicity of injected hrTNF α . We have conducted a series of studies in rabbits designed to evaluate the effect of 3D-MPL pretreatment on a subsequent lethal challenge (35 µg/kg) or pyrogenic challenge (40 ng/kg) of Salmonella abortus equi LPS. Pretreatment of rabbits I.V. with 3D-MPL over a non-pyrogenic dose range of 2.5 µg/kg to 12.5 µg/kg provided significant protection (80-100%) to a lethal challenge of LPS given 24 hrs. later. However, pretreatment with non-pyrogenic doses of 3D-MPL (≤ 12.5 µg/kg) did not significantly alter the biphasic fever response to a pyrogenic (40 ng/kg) challenge of LPS. Of interest was that rabbits treated with doses of 3D-MPL in excess of 50 µg/kg exhibited a monophasic fever (peak at 1.0 hr.), yet when challenged 24 hrs. later with 40 ng/kg LPS they appeared to be refractile with respect to the second fever peak typically seen at 2.5-3.0 hr. in control rabbits. These studies taken collectively have suggested to us that 3D-MPL may have prophylactic value in ameliorating the problem of sepsis and septic shock in clinical medicine.

Monophosphoryl Lipid A as Prophylaxis for Surgical Wound Infection
A Phase I/II Study

J. David Beatty¹, Jon A. Rudbach², Merle Smith¹, Terry Ulrich¹, Angeles Mison¹,
Kenneth B. Von Eschen², Barbara G. Beatty¹, Jose Esteban¹

¹ Division of Surgery, City of Hope National Medical Center, Duarte CA

² Ribic Immunochem Inc. Hamilton MI

³ Division of Pathology, City of Hope National Medical Center, Duarte CA

This study was undertaken to examine the effect of monophosphoryl lipid A (MPL), an attenuated analogue of toxic lipid A, as a nonspecific pre-operative immune stimulant in patients undergoing complex surgical resections for head and neck cancer. MPL was given 24-48h prior to the surgical procedure. All patients received prophylactic cephoxitin and gentamicin for 3 days beginning 8 hours prior to surgery. Patients were observed for toxicity, wound complications and systemic complications. Hematological, biochemical and immunological blood tests were performed pre-MPL infusion; 2h and 24h post-MPL; pre-surgery; and 3, 7 and 28 days after surgery.

Escalating doses of MPL were given to 19 patients: 10 $\mu\text{g}/\text{m}^2$ (2 pt), 35 $\mu\text{g}/\text{m}^2$ (2 pt) and 100 $\mu\text{g}/\text{m}^2$ (15 pt). At 100 $\mu\text{g}/\text{m}^2$, 60% of patients (9/15) had MPL related toxicity including chills and rigor (8/15), fever (7/15), headache (4/15), muscle pain (2/15), lethargy (1/15) and weakness (1/15). There were 14 grade 1, 9 grade 2, and 1 grade 3 toxicities. To decrease toxicity during the study, the dose of MPL was limited to the greater of 200 μg or 2.5 $\mu\text{g}/\text{kg}$ ideal body weight in patients with body surface area over 2m^2 .

At 100 $\mu\text{g}/\text{m}^2$ MPL, 7/15 patients had surgical wound complications including decreased flap viability (6/15), erythema (6/15), induration (4/15), infection (2/15), fistula (1/15), skin contamination (1/15) and dehiscence (1/15). Postoperatively, all patients (15/15) had fever and one patient had respiratory tract infection. There were no cardiac or renal complications. No association was noted between MPL toxicity, and local or systemic complications.

Standard hematological and biochemical tests were unaffected by MPL except for neutrophil count. Juvenile neutrophils (bands) were increased from <1% to 6% at 2h post-MPL and mature neutrophils from 66% to 78%. By 24h they had returned to normal. The surgical procedure resulted in a rise in neutrophils, platelets, fibrinogen and liver function tests.

Levels of immunoglobulin (IgG, IgA, IgM), β_2 microglobulin, complement (C_{3c} , C_4), lymphocyte markers (CD2, CD4, CD8, CD20, CD25, DR, NK), monocyte/macrophage adherence and phagocytosis, monocyte superoxide dysmutase, tetanus toxoid recall and the cytokines IL-2, GM-CSF and γ -INF were not affected by MPL infusion. The levels of serum CRP were elevated by both MPL and surgery. Serum IL-6 and TNF- α were normal (<10ng/ml) pre-MPL and 24h post-MPL. At 2h post-MPL IL-6 was elevated in 11/15 patients (358 ± 109 $\mu\text{g}/\text{ml}$) and TNF- α was elevated in 8/15 patients (32.3 ± 9.3 $\mu\text{g}/\text{ml}$). Levels returned to normal by 24h post-MPL. Correlation was noted between IL-6 and TNF- α levels, and between these cytokines and toxicity. Neopterin levels were increased at 24h post-MPL in 13/15 patients.

In conclusion, prophylactic MPL was safely administered prior to major surgical procedures and resulted in non-specific indications of an immune response and increased levels of cytokines associated with macrophage activation. Confirmation that prophylactic MPL significantly decreases the incidence of local wound sepsis would require a large multi-center trial.

**Induction of Endotoxin Tolerance by Nontoxic Derivatives of
Lipid A**

Mark E. Astiz, M.D.
Associate Professor of Medicine
St. Vincent's Hospital and Medical Center of New York
New York, New York

Several nontoxic derivatives of endotoxin (LPS) and lipid A have been recently isolated. These substances are significantly less toxic than the parent molecule yet maintain the beneficial immunomodulatory activities of endotoxin. We have been particularly interested in the ability of these substances to modify the systemic toxicity of endotoxin and to induce endotoxin tolerance.

Monophosphoryl lipid A (MPL) is a nontoxic derivative of lipid A which lacks a phosphate group at the reducing end of the molecule. The ability of MPL to modulate endotoxin activity was examined in several studies.

In initial studies, rats were pretreated with *S. minnesota* monophosphoryl lipid A fifteen minutes prior to endotoxin infusion. A significant attenuation of the decrease in arterial pressure and decrease in cardiac output was observed in MPL treated animals. Mean arterial pressure (MAP) decreased from 134 mmHg to 90 mmHg and cardiac output (CO) decreased from 356 ml/kg/min to 229 ml/kg/min in LPS animals while the animals pretreated with MPL showed no significant changes in MAP or CO.

In a subsequent study effect of MPL on LPS induced priming of human neutrophil superoxide radical production was examined. In contrast to LPS, MPL exhibited no neutrophil priming activity in concentrations ranging from 1 ng/ml to 10^3 ng/ml. Incubation of neutrophils with MPL prior to LPS resulted in dose-dependent inhibition of LPS-inducing neutrophil priming.

The induction of MPL was also examined in large and small animals models. In rats, exposure to MPL 48 hours prior to lethal endotoxin infusion produced 100%.

Seventy-two hour survival as compared to 100% mortality in animals pretreated with diluent control. Significant attenuation of decrease in arterial pressure and cardiac output were also observed. A similar study in a porcine model demonstrated that a pretreatment with MPL 48 hours prior to lethal endotoxemia produced 20% mortality as compared to 100% mortality in diluent controls. Hemodynamic changes following endotoxin infusion were significantly attenuated and tumor necrosis factor release was significantly reduced in the MPL treated animals.

The toxicity and ability of *R. sphaeroides* lipid A (RS) to induce tolerance was also studied. LPS and RS were infused in equal doses in rats. LPS resulted in significant decreases in cardiac output and arterial pressure whereas infusion of RS and was not associated with significant hemodynamic changes. Exposure of rats to RS forty-eight hours prior to lethal endotoxin infusion significantly decreased mortality from 100% in control animals to 20% of the tested animals.

These studies suggest that nontoxic derivatives of lipid A may be able to modify the systemic response to endotoxin by inducing endotoxin tolerance.



Evening Lecture

Molecular Mechanisms of Intracellular Signalling

Monday, April 8 • 7:15 p.m.

Dr. Masayasu Nakano

Jichi Medical School • Tochigi-Ken, Japan

MOLECULAR MECHANISMS OF INTRACELLULAR SIGNALLING IN MURINE
MACROPHAGES BY LPS OR LIPID A ANALOGS

Masayasu Nakano, Department of Microbiology, Jichi Medical
School, Tochigi-ken 329-04, Japan

Intracellular events in the peritoneal exudate macrophages of LPS-responsive C3H/He and LPS-low responsive C3H/HeJ mice after the stimulation with LPS or lipid A analogs were studied. LPS-induced IL-1 production by C3H/He macrophages seemed to be dependent on calmodulin-dependent pathway rather than the pathways dependent on protein-kinase C, A, or G. LPS induces phosphorylation of a distinct set of substrate proteins in the C3H/He, but not C3H/HeJ, macrophages prior to final LPS-induced cellular functions such as IL-1 or TNF production. The phosphorylated proteins (pp) were located in either a cytosol fraction (pp65 & 28) or a membrane fraction (pp58, 43, 33, & 19) or in both fractions (pp105 & 75). Among the pp, pp65 was the most heavily phosphorylated cytosolic soluble protein, and was a substrate for serine kinase. When the macrophages were stimulated with biologically active lipid A analogs, the patterns of phosphorylation were similar to that of the macrophages stimulated with LPS. However, when the macrophages were stimulated with biologically non-active lipid A analogs, some of the pp were not observed.



Session V

Current Vaccine Strategies Using Lipid A Derivatives

Tuesday, April 9 • 8:30 a.m. - 11:25 a.m.

Speakers' Summaries (in order of presentation)

Abstract

Development of a Synthetic Lipid A Analogue Vaccine. Bhattacharjee AK, Drabick J, Wright CD, Sadoff JC, Cross AS. Department of Bacterial Diseases, WRAIR, Washington DC and Univax Biologics, Inc, Rockville, MD

The mortality from sepsis due to Gram-negative bacteria remains high despite the availability of potent antimicrobial agents. The recently reported success of a human MAb to lipid A, HA-1A, derived from a patient immunized with a J5 vaccine, suggests that immunotherapy directed at lipid A may be an important supplement to the conventional care of septic patients. Bieber et al. demonstrated that the binding of anti-J5 MAb to lipid A is inhibited by gentiobiose octaacetate. Consequently we constructed a synthetic lipid A analogue conjugate vaccine by coupling gentiobiose heptaacetate (GHA) to a protein carrier, succinylated diphtheria toxoid, through spacer compounds. This vaccine bound to the anti-lipid A MAb in ELISA assay, and induced antibody to GHA in rabbits and mice (>2500 ELISA antibody units [U]), but less antibody to J5 LPS (127 U v. 15 U in unimmunized rabbits) or to lipid A (100 U v. 25 U in unimmunized rabbits). Human antibody to lipid A derived from a volunteer immunized with monophosphoryl lipid A in a liposome, also bound to GHA (>675 units). A functionally-active lapine anti-J5 Ab bound to J5 LPS and lipid A (>2900 units) but not to GHA (16 units). Functional activity of anti-GHA Ab was demonstrated in a Limulus lysate neutralization test, and in protection against active and passive dermal Shwartzman reactions. Phase I testing with this vaccine will be initiated this year. These data also suggest there might be multiple epitopes to which functionally active Ab to lipid A may be directed.

ADJUVANT EFFECTS OF LIPOSOMAL LIPID A: HUMAN VACCINE INDUCES ANTIBODIES TO BOTH MALARIA ANTIGEN AND LIPID A

CARL R. ALVING, ROBERTA L. RICHARDS, JAMES E. EGAN, CLYDE SCHULTZ, DANIEL M. GORDON, AND LOUIS F. FRIES

Depts of Membrane Biochemistry, Immunology, and Biologics Research, Walter Reed Army Inst of Research, Washington, DC 20307-5100, and Ctr. for Immuniz. Res., Johns Hopkins Univ., Baltimore, MD 21205 USA

Liposomes containing lipid A served as carriers of a synthetic antigen for immunizing mice, rabbits, monkeys, and humans against sporozoites of *Plasmodium falciparum*. Liposomal native lipid A and liposomal monophosphoryl lipid A (MPL) had potent dose-dependent adjuvant properties. Although large amounts of lipid A were required for optimal activity, even very high doses of liposomal lipid A were nonpyrogenic in rabbits. An experimental vaccine was developed consisting of liposomes containing both MPL and a recombinant antigen (R32NS1₈₁) having epitopes from the repeat region of the circumsporozoite protein of *P. falciparum*. The liposomes were adsorbed with alum. A phase I safety trial of the vaccine in humans revealed no significant acute toxic effects and the vaccine was nonpyrogenic in humans even at an MPL dose of approximately 2.2 mg. High titers of IgG antibodies both to repeat region epitopes of the malaria antigen and to the MPL itself developed within two weeks after a single injection of the vaccine in humans. The highest dose of liposomal MPL induced the highest titers of antibodies to lipid A. Liposomes effectively blocked potential toxic effects but retained the adjuvant effects of lipid A. The vaccine consisting of alum-adsorbed liposomes containing high doses of MPL and encapsulated antigen appears to be safe and efficacious for inducing humoral immune responses in humans.

AN EFFECTIVE VACCINE FOR SYPHILIS: THE IMPORTANCE OF THE ADJUVANT ACTION OF
MPL IN REVERSING IMMUNOSUPPRESSION. Dr. Tom Fitzgerald, University of Minnesota
Duluth, MN. 55812

Abstract

Our research involves mechanisms of immunoregulation in chronic diseases, using syphilitic infection as the model. Following rabbit inoculation with Treponema pallidum, a number of immune functions are stimulated within the initial 10 days of infection. At this time, down-regulation of these functions becomes detectable, and levels of IL-1, IL-2, γ -interferon, and class II antigens are diminished. Macrophages and T-lymphocytes participate in this suppression through their secretion of PGE₂ and transforming growth factor (TGF). We have proposed that this down-regulation is premature and that residual treponemes then establish chronic infection. We have been able to reverse this down-regulation by using indomethacin to block macrophage suppression and monophosphoryl lipid A (MPL) to block T-suppression. Soluble factors including TGF are produced by activated lymphocytes. These factors inhibit Con A induced IL-2 synthesis and T-cell proliferation. MPL reverses both of these 2 suppressive effects. In addition, treponemes directly inhibit IL-2 generation and MPL reverses this defect. We have extended these findings to vaccine development. Our premise was that past vaccine failures were attributed to the ability of treponemes to induce down-regulation, thereby preventing immune stimulation. We have now succeeded in developing an effective vaccine by counter-acting these suppressive tendencies. Our protocol involves: cyclophosphamide day -2 (neutralizes natural suppressors and T-suppressors), inactivated T. pallidum plus Ribi adjuvant (MPL and trehelose dimycolate) day 0, MPL boost day +2 (neutralizes elicited T-suppressors), challenge with viable organisms day 29, and indomethacin day 29 to 36 (neutralizes macrophage suppressors). This unique vaccine approach of preventing suppression may have applications to other chronic diseases that involve macrophage suppression and/or T-cell suppression.

Dr. Tom Fitzgerald
University of Minnesota-Duluth

EVALUATION OF MONOPHOSPHORYL LIPID A (MPL) AS AN ADJUVANT FOR HUMAN IMMUNIZATION: ENHANCEMENT OF THE SERUM ANTIBODY RESPONSE IN YOUNG MICE TO POLYSACCHARIDE-PROTEIN CONJUGATES BY CONCURRENT INJECTION WITH MPL

Rachel Schneerson,¹ Ali Fattom,¹ Shousun C. Szu,¹ Dolores Bryla,² Ulrich,³ Jon A. Rudbach,³ Gerald Schiffman,⁴ and John B. Robbins¹

¹Laboratory of Developmental and Molecular Immunity, ²Biometry and Mathematical Statistics Branch, National Institute of Child Health and Human Development, NIH, Bethesda, MD 20892; ³RIBI Immunochemicals, Inc., P.O. Box 1409, Hamilton, MT 59840; and ⁴Department of Microbiology and Immunology, Downstate Medical Center SUNY, 480 Clarkson Avenue, Brooklyn, NY

Concurrent injection of monophosphoryl lipid A (MPL) in saline or as an oil-in-water emulsion (o/w) enhanced both the primary and secondary serum antibody responses to the capsular polysaccharide (CP) components of seven conjugates: the enhanced responses were Ag-specific. In contrast, MPL did not enhance the serum antibody response to 5 of the 6 unconjugated CPs. MPL and trehalose dimycolate (TDM) injected concurrently with the unconjugated Vi CP of *Salmonella typhi* (Vi) enhanced the serum antibody response to that Ag. MPL further enhanced the Vi antibody levels when injected with conjugates of this CP. The serum antibody responses to *Pseudomonas aeruginosa* exotoxin A (ETA), used as the carrier protein for the *Staphylococcus aureus* types 5 and 8 conjugates, were also enhanced by MPL. MPL in o/w was generally more effective than when administered in saline.



List of Participants

Newer Aspects of the Adjuvant Action of Lipid-A
and its Analogs

April 6-9, 1991

List of Speakers/Participants

Dr. Carl R. Alving
Chief
Dept. of Membrane Biochemistry
Walter Reed Army Institute of Research
Washington, DC 20307-5100
(202)-576-3248

Dr. Mark E. Astiz
St. Vincent's Hosp. & Med. Ctr. of NY
153 West 11th Street
New York, NY 10011

Dr. Phillip Baker
NIAID\NIH
Twinbrook II
12241 Parklawn Drive
Rockville, MD 20852

Dr. Michael Balady
Biologics
USAMMDA
Fort Detrick
Frederick, MD 21702
(301)-663-7661

Dr. W. Ripley Ballou
Dept. of Immunology
Walter Reed Army Institute of Research
Washington, DC 20037
(202)-576-1257

Dr. J. David Beatty
Division of Surgery
Research and Education
City of Hope National Medical Center
1500 East Duarte Road
Duarte, CA 91010

Dr. Apurba K. Bhattacharjee
Bacterial Diseases
Walter Reed Army Institute of Research
14th and Dahlia
Washington, DC 20037
(202)-576-3338

Dr. Donald S. Burke
Division of Retrovirology
Walter Reed Army Institute of Research
13 Taft Court, Suite 201
Rockville, MD 20850
(301)-295-6414

Dr. Robert Burrell
Microbiology & Immunology
WVU Health Sciences Center
2082 HSN
Morgantown, WV 26506
(304)-295-1177

Dr. Yupin Charoenvit
Malaria Program
Naval Medical Research Institute
12300 Washington Avenue
Rockville, MD 20852
(301)-295-1177

Ms. Carole Cole
Biologist
Resources and Centers Branch/BRDP
Division of AIDS
NIAID/NIH
6003 Executive Boulevard
Bethesda, MD 20892
(301)-402-0755

Mr. James Cradock
Acting Chief
Developmental Therapeutics Branch/BRDP
Division of AIDS
NIAID/NIH
6003 Executive Boulevard
Bethesda, MD 20892
(301)-496-0636

Dr. Alan Cross
Bacterial Diseases
Walter Reed Army Institute of Research
Washington, DC 20307
(202)-576-3326

Dr. Jacques C. Darbord
Microbiology
University
7 Rue Fer A Aoulin
75221 Paris Cedex 05,
FRANCE

Dr. Michael J. Darsley
Hyrridoma
IGEN, Inc.
1530 E. Jefferson Street
Rockville, MD 20852
(301)-984-8000

Dr. Barbara Detrick
Microbiologist
Vaccine Res. & Devel. Branch\BRDP
Division of AIDS
NIAID/NIH
6003 Executive Boulevard
Bethesda, MD 20892
(301)-496-8200

Dr. Sarmangala Devi
Laboratory of Develop. & Mole. Immunity
NICHD\NIH
Building 6, Room 145
Bethesda, MD 20892

Dr. Roman Dziarski
Northwest Center for Medical Education
Indiana University School of Medicine
3400 Broadway
Gary, IN 46408
(219)-980-6551

Dr. Emilio Emini
Virus and Cell Biology Division
Merck, Sharp & Dohme Research Labs.
Building 16, Room 100
West Point, PA 19486
(215)-661-7859

Dr. Martin E. Evans
Infectious Diseases\Dept. of Medicine
Uniformed Services
University of Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20814
(301)-295-3618

Dr. Patricia Fast
Medical Officer
Vaccine Research & Devel. Branch\BRDP
Division of AIDS
NIAID/NIH
6003 Executive Boulevard
Bethesda, MD 20892
(301)-496-8200

Dr. Thomas Fitzgerald
Microbiology
UMD Medical School
10 University Drive
Duluth, MN 55812
(218)-726-7278

Dr. H.D. Flad
Immunology and Cell Biology
Forschungsinstitut Borstel
D-2061 Borstel,
GERMANY

Dr. Chris Galanos
Max Planck Instit. fur Immunobiologie
Stubeweg 51
Freiburg, D-7800
GERMANY

Dr. Nathalie Garcon
Vaccine Formulation Science
Smith Kline Beecham Biologicals
Rue de L'institut, 89
Rixensart, 1330
BELGIUM

Ms. Marta W. Glass
Biologist
Vaccine Research & Develop. Branch\BRDP
Division of AIDS
NIAID/NIH
6003 Executive Boulevard
Bethesda, MD 20892
(301)-496-8200

Dr. Douglas Golenbock
Boston City Hospital
Boston, MA

Dr. Gary L. Gustafson
Ribi ImmunoChem Research, Inc.
P.O. Box 1409
Hamilton, MT 59840
(406)-363-6214

Ms. Beth E. Henricson
Dept. of Microbiology
Unif. Serv. Univ. Health Sci./Vogel Lab.
4301 Jones Bridge Road
Bethesda, MD 20814
(301)-295-3447

Dr. Stephen L. Hoffman
Malaria Program
Naval Medical Research Institute
12300 Washington Avenue
Rockville, MD 20852
(301)-295-2000

Dr. John J. Hooks
Chief
Virology and Immunology Section
NEI/NIH
Building 10, Room 6N228
9000 Rockville Pike
Bethesda, MD 20892
(301)-496-6578

Dr. Tomas Hraba
Laboratory of Immunogenetics
NIAID/NIH
Twinbrook II Research Facility
12441 Parklawn Drive
Rockville, MD 20852
(301)-496-1220

Dr. Robert L. Hunter
Professor
Dept. of Pathology/Lab. Medicine
Emory University School of Medicine
762 Woodruff Memorial Building
Atlanta, GA 30322

Dr. Arthur G. Johnson
Head
Dept. of Medical Microbiology/Immunology
Univ. of Minnesota School of Medicine
Duluth, MN 55804
(218)-726-7561

Dr. Trevor R. Jones
Malaria Program
Naval Medical Research Institute
12300 Washington Avenue
Rockville, MD 20852
(301)-295-0178

Dr. Charlotte R. Kensil
Section Head
Natural Products Chemistry
Cambridge BioScience Corporation
365 Plantation Street
Worcester, MA 01605
(508)-797-5777

Dr. Srisin Khusmith
Malaria Program
Naval Medical Research Institute
12300 Washington Avenue
Rockville, MD 20852
(301)-295-0171

Ms. Janet Kirkley
Dept. of Biochemistry
The George Washington University
Medical Center, Ross Hall
2300 Eye Street, NW, Room 537
Washington, DC 20037
(202)-994-2312

Dr. Scott Koenig
MedImmune, Inc.
19 Firstfield Road
Gaithersburg, MD 20878
(301)-590-2621

Dr. Wayne C. Koff
Chief
Vaccine Research and Devel. Branch\BRDP
Division of AIDS
NIAID/NIH
6003 Executive Boulevard
Bethesda, MD 20892
(301)-496-8200

Dr. Dale N. Lawrence
Chief
International Studies, BRDP
Division of AIDS
NIAID/NIH
6003 Executive Blvd.
Rockville, MD 20892
(301)-496-8200

Mr. Nicholas J. Legakis
Dept. of Microbiology
University of Athens
Athens Tower, Building B
Athens 115 27
Mesogeion 2-4,
GREECE

Dr. Mei-Guey Lei
Dept. of Microb., Molec. Gen. & Immun.
University of Kansas Medical Center
39th and Rainbow Boulevard
Kansas City, KS 66103
(913)-588-7061

Dr. Anita Malik
Malaria Program
Naval Medical Research Institute
12300 Washington Avenue
Rockville, MD 20852
(301)-295-2000

Dr. Motohiro Matsuura
Dept. of Microbiology
Jichi Medical School
3311-1, Yakushiji, Minami
Kawachi-Machi
Tochigi, 329-04
JAPAN

Dr. Edward McSweegan
BMB
NIAID/NIH
5333 Westbard Avenue
Westwood Building, Room 738
Bethesda, MD 20892
(301)-496-7728

Dr. Sylvia Mellouk
Malaria Program
Naval Medical Research Institute
12300 Washington Avenue
Rockville, MD 20852
(301)-295-6173

Dr. Donald Milton
Environmental Health
Harvard School of Public Health
665 Huntington Avenue
Boston, MA 02115
(617)-432-3324

Dr. Robert S. Munford
Internal Medicine and Microbiology
University of Texas SW Medical Center
5323 Harry Hines Blvd.
Dallas, TX 75235

Dr. Kent R. Myers
Research and Development
Ribi ImmunoChem Res., Inc.
P.O. Box 1409
Hamilton, MT 59840
(406)-363-6214

Dr. Masayasu Nakano
Dept. of Microbiology
Jichi Medical School
Tochigi-ken 329-04,
JAPAN

Dr. Paul Naylor
Associate Professor
Dept. of Biochemistry and Molecular Bio.
George Washington Univ. Medical Center
2300 Eye Street, NW
Washington, DC 20037
(202)-994-1742

Dr. Alois H. Nowotny
Dept. of Immunology
University of Pennsylvania
L. Levy Center for Oral Health Research
Bldg. A-2, 4010 Locust Street
Philadelphia, PA 19104

Ms. Marilyn Odean
Dept. of Med. Microbiology/Immunology
UMD School of Medicine
10 University Drive
Duluth, MN 55812
(218)-726-7940

Dr. I. Outschoorn
Instituto Nac. Microbiology
Majadahonda
Madrid,

Dr. Matthew Pollack
Division of Infectious Diseases
Dept. of Medicine
Uniformed Services Univ. of Heal. Sci.
4301 Jones Bridge Road
Bethesda, MD 20814
(301)-295-3618

Dr. Richard A. Proctor
Dept. of Med. Medical Micro. & Immunol.
Univ. of Wisconsin - Madison Med. Sch.
436 Service Memorial Institutes
1300 University Avenue
Madison, WI 53706

Dr. Nilofer Qureshi
Dept. of Mycobacteriology
William S. Middleton Mem. Vet. Hospital
2500 Overlook Terrace
Madison, WI 53705
(608)-256-1901

Dr. Christian R.H. Raetz
Division of Merck & Co., Inc.
Merck, Sharp and Dohme Research Labs.
P.O. Box 2000
Rahway, NJ 07065

Dr. Mangala Rao
Membrane Biochemistry
Walter Reed Army Institute of Research
Washington, DC 20307
(202)-576-3545

Dr. Elmer Reist
Bio-Organic Chemistry Laboratory
SRI International
333 Ravenswood Avenue
Menlo Park, CA 94025
(415)-859-3543

Dr. Robert M. Rice
Biological Systems, USAMMDA
Fort Detrick
Frederick, MD 20702
(301)-663-7661

Dr. Roberta L. Richards
Membrane Biochemistry
Walter Reed Army Institute of Research
Washington, DC 20307
(202)-576-3940

Dr. Ernst Th. Rietschel
Professor
Institut für Exper. Biologie und Medizin
Forschungsinstitut Borstel,
Parkalle 1-40 D 2061,
GERMANY

Dr. Einar Rosenqvist
Vaccine Department
National Inst. of Public Health
Geitmyrsveien 75
0462 Oslo 4,
NORWAY

Dr. Jon A. Rudbach
Vice President
Research and Development
Ribi ImmunoChem Research, Inc.
P.O. Box 1409
Hamilton, MT 59840
(406)-363-6214

Dr. John L. Ryan
Infectious Diseases - Clinical Research
Merck, Sharp & Dohme Research Labs.
Sumneytown Pike (BL 3-4)
West Point, PA 19486
(215)-834-2557

Dr. Jerald C. Sadoff
Walter Reed Army Institute of Research
Washington, DC 20307

Dr. Martha Sedegah
Malaria Program
Naval Medical Research Institute
12300 Washington Avenue
Rockville, MD 20852
(301)-295-2000

Dr. M. Slaoui
Molecular and Cellular Biology
Smith Kline Beecham Biologicals
89 Rue de L'institut 330
Rixemart,
BELGIUM

Dr. H. Snippe
Medical Microbiology
Eijkman-Winkler Lab. of Med. Micro.
Medical School AZU G 04.614
Heidelberglaan 100
3584 CX Utrecht,
NETHERLANDS

Dr. P. Stuetz
Sandoz Forschungsinstitut Gesellschaft
Brunner Strasse 59
A-1235 Wien,
AUSTRIA

Dr. Kuni Takayama
Mycobacteriology Research Lab.
Williams S. Middleton Mem. Veter. Hosp.
2500 Overlook Terrace
Madison, WI 53705

Dr. Nobumasa (Nobi) Tanaka
US-Japan Science Consulting Serv., Inc.
72 Paxwood Road
Delmar, NY 12054
(518)-439-8334

Dr. Chao-Ming Tsai
Bacterial Products
Center for Biologics
Food and Drug Administration
8800 Rockville Pike
Bethesda, MD 20892
(301)-496-2154

Dr. J. Terry Ulrich
Infectious Disease Research
Ribi ImmunoChem Research, Inc.
P.O. Box 1409
Hamilton, MT 59840
(406)-363-6214

Dr. Frederick R. Vogel
Lederle Biologicals
Lederle Laboratories
Lederle Lab. Bldg. 114/100
Pearl River, NY 10965
(914)-732-4481

Dr. Stephanie Vogel
Microbiology
Uniformed Svcs. Univ. of the Hlth. Sci.
4301 Jones Bridge Road
Bethesda, MD 20814
(301)-295-3446

Dr. Shaw Warren
Infectious Diseases
Massachusetts General Hospital
Gray 5, Fruit Street
Boston, MA 02114
(617)-726-5772

Dr. Nabila M. Wassef
Membrane Biochemistry
Walter Reed Army Institute of Research
Washington, DC 20307
(202)-576-3528

Dr. Richard O. Williams
Research and Development
IGEN, Inc.
1530 E. Jefferson Street
Rockville, MD 20852
(301)-984-8000

Dr. Scott Winston
Univax Biologics, Inc.
12111 Parklawn Drive
Rockville, MD 20852
(301)-770-3099

Dr. James F. Young
MedImmune, Inc.
19 Firstfield Road
Gaithersburg, MD 20878
(301)-590-2642

REGISTRANTS: Airlie Conference Center, April 6-9.

- Dr. Carl R. Alving,
Membrane Biochemistry, Walter Reed Army Institute of Research,
Washington, D.C. 20307, 202-576-3248.
- Dr. Shimon Amselem,
Nova Pharmaceutical Corporation, 6200 Freeport Center,
Baltimore, MD 21224, 202-576-0777.
- Dr. Mark E. Astiz,
St. Vincent's Hospital and Medical Center of New York, 153 West
11th Street, New York, N.Y. 10011.
- Dr. Phillip Baker,
NIAID-NIH, Twinbrook II, 12241 Parklawn Drive, Rockville, MD
20852.
- Dr. Michael Balady,
Biologics, USAMMDA, Fort Detrich, Frederick, MD 21702, 301-
663-7661.
- Dr. W. Ripley Ballou,
Dept. of Immunology, Walter Reed Army Institute of Research,
Washington, D.C. 20037, 202-576-1257.
- Dr. J. David Beatty,
Research and Education, Division of Surgery, City of Hope
National Medical Center, 1500 East Duarte Road, Duarte, CA
91010.
- Dr. Apurba K. Bhattacharjee,
Bacterial Diseases, Walter Reed Army Inst. of Research, 14th and
Dahlia, Washington, D.C. 20307, 202-576-3338.
- Dr. Donald S. Burke,
Division of Retrovirology, Walter Reed Army Institute of
Research, 13 Taft Court, Suite 201, Rockville, MD 20850, 301-
295-6414 (ext. 15/16).
- Dr. Robert Burrell,
Microbiology and Immunology, WVU Health Sciences Center, 2082
HSN, Morgantown, WV 26506, 304-293-3559.
- Dr. Yupin Charoenvit,
Malaria Program, Naval Medical Research Institute, 12300
Washington Ave., Rockville, MD 20852, 301-295-1177.
- Carole Cole,

M.S., Resources and Centers Branch, BRDP, Division of AIDS,
National Institute of Allergy and Infectious Diseases, NIH, 6003
Executive Blvd., Bethesda, MD 20892, 301-402-0755.

James Craddock,

M.S., Developmental Therapeutics Branch, BRDP, Division of AIDS,
National Institute of Allergy and Infectious Diseases, NIH, 6003
Executive Blvd., Bethesda, MD 20892, 301-496-0636.

Dr. Robert Crawford,

Cellular Immunology, Walter Reed Army Institute of Research,
18321 Queen Elizabeth Drive, Olney, MD 20832.

Dr. Alan Cross,

Bacterial Diseases, Walter Reed Army Institute of Research,
Washington, D.C. 20307, 202-576-3326.

Dr. Bernard Danve,

Microthique/Serologi, Pasteur Merieux Serums and Vaccines,
65280 Nancy l'Etoile, France.

Dr. Jacques C. Darbord,

Microbiology, University, 7 Rue Fer A Aoulin 75221 Paris Cedex
05, Paris, France.

Dr. Michael J. Darsley,

Hyrridoma, IGEN, Inc., 1530 E. Jefferson St., Rockville, MD 20852,
301-984-8000.

Dr. Barbara Detrick,

Vaccine Research and Development Branch, BRDP, Division of
AIDS, National Institute of Allergy and Infectious Diseases, NIH,
6003 Executive Blvd., Bethesda, MD 20892, 301-496-8200.

Dr. Sarmangala Devi,

Laboratory of Development and Molecular Immunity, NICHD, NIH,
Building 6, Room 145, Bethesda, MD 20892.

Dr. Roman Dziarski,

Northwest Center for Medical Education, Indiana University
School of Medicine, 3400 Broadway, Gary, IN 46408, 219-980-
6551.

Dr. Karen L. Elkins,

Dept. Cellular Immunology, Walter Reed Army Institute of
Research, Rockville, MD 20850, 301-295-6953.

Dr. Emilio A. Emini,

Virus and Cell Biology, Merck, Sharp and Dohme Research
Laboratories, Building 16, Room 100, West Point, PA 19486, 215-

661-7859.

Dr. Martin E. Evans,

Infectious Disease Dept. of Medicine, Uniformed Services
University of the Health Sciences, 4301 Jones Bridge Road,
Bethesda, MD 20814, 301-295-3618.

Dr. Patricia Fast,

Vaccine Research and Development Branch, BRDP, Division of
AIDS, National Institute of Allergy and Infectious Diseases, NIH,
6003 Executive Blvd., Bethesda, MD 20892, 301-496-8200.

Dr. Thomas Fitzgerald,

Microbiology, UMD Medical School, 10 University Drive, Duluth,
MN 55812, 218-726-7278.

Dr. H. D. Flad,

Immunology and Cell Biology, Forschungsinstitut Borstel, D-2061
Borstel, Germany.

Dr. Bruce D. Forrest,

Medical Research Council, UK, CDSC, PHLS 61 Colindale Ave.,
London, UK. NW9 5EQ.

Dr. Marina Freudenberg,

Max Planck Institut fur Immunobiologie, Stubeweg 51, Freiburg,
Germany.

Dr. Chris Galanos,

Max Planck Institut fur Immunobiologie, Stubeweg 51, Freiburg,
D-7800, Germany.

Dr. Nathalie Garcon,

Vaccine Formulation science, Smith Kline Beecham Biologicals,
Rue de l'institut, 89, Rixensart 1330 Belgium.

Marta W. Glass,

M.S., Vaccine Research and Development Branch, BRDP, Division of
AIDS, National Institute of Allergy and Infectious Diseases, NIH,
6003 Executive Blvd., Bethesda, MD 20892, 301-496-8200.

Dr. Douglas Golenbock,

Boston City Hospital, Boston MA.

Dr. Shawn Green,

Cellular Immunology, Walter Reed Army Institute of
Research, 9620 Medical Center Drive, Rockville, MD 20850, 301-
295-6956.

Dr. Gary L. Gustafson,

Ribi ImmunoChem Research, Inc., P.O. Box 1409, Hamilton, MT
59840, 406-363-6214.

- Dr. Scott W. Harkonen,
Medical and Regulatory Affairs Dept., Univax Biologics, In. 12280
Wilkins Ave., Rockville, MD 20852, 301-770-3099.
- Beth E. Henricson,
Microbiology, Uniformed Services University of Health
Science/Vogel Laboratory, 4301 Jones Bridge Road, Bethesda, MD
20814, 301-295-3447.
- Dr. Stephen L. Hoffman,
Malaria Program, Naval Medical Research Institute, 12300
Washington Ave., Rockville, MD 20852, 301-295-2000.
- Dr. John J. Hooks,
Virology and Immunology Section, National Eye Institution, NIH,
Bldg. 10 Rm 6N228, Bethesda, MD 20892, 301-496-6578.
- Dr. Tomas Hraba,
Laboratory of Immunogenetics, National Institute of Allergy and
Infectious Diseases, NIH, Twinbrook II Research Facility, 12441
Parklawn Dr., Rockville, MD 20852, 301-496-1220.
- Dr. Robert Hunter,
Emory University School of Medicine, Dept. of Pathology and
Laboratory Medicine, Atlanta, GA 30322.
- Dr. Arthur G. Johnson,
Medical Microbiology and Immunology, University of Minnesota,
School of Medicine, Duluth, MN 55804, 218-726-7561.
- Dr. Margaret Johnston,
DAIDS.
- Dr. Trevor R. Jones,
Malaria Program, Naval Medical Research Institute, 12300
Washington Ave., Rockville, MD 20852, 301-295-0178.
- Dr. Charlotte R. Kensil,
Natural Products Chemistry, Cambridge Biotech Corporation, 365
Plantation St., Worcester, MA 01605, 508-797-5777.
- Dr. Srisin Khusmith,
Malaria Program, Naval Medical Research Institute, 12300
Washington Ave., Rockville, MD 20852, 301-295-0171.
- Janet Kirkley,
Dept. of Biochemistry, George Washington University Medical
Center, 2300 Eye St., NW, Ross Hall Rm. 537, Washington, D.C.
20037, 202-994-2312.
- Dr. Scott Koenig,

MedImmune, Inc., 19 Firstfield Road, Gaithersburg, MD 20878,
301-590-2621.

Dr. Wayne C. Koff,

Vaccine Research and Development Branch, BRDP, Division of
AIDS, National Institute of Allergy and Infectious Diseases, NIH,
6003 Executive Blvd., Bethesda, MD 20892, 301-496-8200.

Dr. Dale N. Lawrence,

International Studies, Office of the Chief Vaccine Research and
Development Branch, BRDP, Division of AIDS, National Institute of
Allergy and Infectious Diseases, NIH, 6003 Executive Blvd.,
Bethesda, MD 20892, 301-496-8200.

Nicholas J. Legakis,

Microbiology, University of Athens, Athens Tower, Building B,
Athens 115 27, Mesogeion 2-4.

Dr. Mei-Guey Lei,

Microbiology, Molec. Genetics and Immunology, University of
Kansas Medical Center, 39th and Rainbow Blvd., Kansas City, KS
66103, 913-588-7061.

Dr. Heinz Leibl,

Immuno Ag, Immunological Research, Industriestrasse 131, A-
1220 Vienna, Austria (01143) 2300, Ext. 2721.

Dr. Jeffrey L. Jorde,

Biology Division, Office of Naval Research, 800
N. Quincy St., Arlington, VA 22217, 703-696-4055.

Dr. Anita Malik,

Malaria Program, Naval Medical Research Institute, 12300
Washington Ave., Rockville, MD 20852, 301-295-2000.

Dr. Josef W. Mannhalter,

Immunological Research, Immuno Ag, Industriestrasse 131, A-
1220 Vienna, Austria (01143) 2300, Ext. 2721.

Dr. Motohiro Matsuura,

Microbiology, Jichi Medical School, 3311-1, Yakushiji, Minami-
Kawachi-Machi, Tochigi, 329-04, Japan.

Dr. Edward McSweegan,

NIAID/BMB, NIH, Westwood Bldg. Room 738, Bethesda, MD 20892,
301-496-7728.

Dr. Sylvia Mellouk,

Malaria Program, Naval Medical Research Institute, 12300
Washington Ave., Rockville, MD 20852, 301-295-6173.

Dr. Monte Meltzer,

Dept. Cellular Immunology, Walter Reed Army Institute of Research, 9620 Medical Center Drive, Suite 200, Rockville, MD 20850, 301-295-6949.

Dr. Donald Milton,

Environmental Health, Harvard School of Public Health, 665 Huntington Ave., Boston, MA 02115, 617-432-3324.

Dr. Robert S. Munford,

Internal Medicine and Microbiology, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, TX 75235.

Dr. Kent R. Myers,

Research and Development, Ribic ImmunoChem Res., Inc., P.O. Box 1409, Hamilton, MT 59840, 406-363-6214.

Dr. Masayasu Nakano,

Dept. of Microbiology, Jichi Medical School, Tochigi-ken 329-04, Japan.

Dr. Paul Naylor,

Biochemistry and Molecular Biology, George Washington University Medical Center, 2300 Eye St., NW, Washington, D.C. 20037, 202-994-1742.

Dr. Alois H. Nowotny,

Immunology, University of Pennsylvania, L. Levy Ctr. for Oral Health Research, Bldg. A-2, 4010 Locust St., Philadelphia, PA 19104.

Marilyn Odean M.S.,

Medical Microbiology and Immunology, UMD School of Medicine, 10 University Drive, Duluth, MN 55812, 218-726-7940.

Dr. I. Outschoorn,

Instituto Nac. Microbiology, Majadahonda, Madrid, Spain.

Dr. Matthew Pollack,

Infectious Diseases/Dept. of Medicine, Uniformed Services University of the Health Sciences, 4301 Jones Bridge Road, Bethesda, MD 20814, 301-295-3618.

Dr. Richard A. Proctor,

Medicine and Medical Microbiology and Immunology, University of Wisconsin-Madison Medical School, 436 Service Memorial Institutes, 1300 University Ave., Madison, WI 53706.

Dr. Nilofer Qureshi,

Mycobacteriology, William S. Middleton Memorial Veterans'

Hospital, 2500 Overlook Terrace, Madison, WI 53705, 608-256-1901.

Dr. Christian R. H. Raetz,

Merck, Sharp and Dohme Research Laboratories, Division of Merck and Co., Inc., P.O. Box 2000, Rahway, NJ 07065.

Dr. Mangala Rao,

Membrane Biochemistry, Walter Reed Army Institute of Research, Washington, D.C. 20307, 202-576-3545.

Dr. Elmer Reist,

Bio-Organic Chemistry Laboratory, SRI International, 333 Ravenswood Ave., Menlo Park, CA 94025, 415-859-3543.

Dr. Robert M. Rice,

Biological Systems, USAMMDA, Fort Detrick, Frederick, MD 20702, 301-663-7661.

Dr. Roberta L. Richards,

Membrane Biochemistry, Walter Reed Army Institute of Research, Washington, D.C. 20307, 202-576-3940.

Prof. Dr. Ernst Th. Rietschel,

Institut für Experimentelle Biologie und Medizin, Forschungsinstitut Borstel, Parkallee 1-40, D 2061, Germany.

Dr. Einar Rosenqvist,

Vaccine Dep., National Inst. of Public Health, Geitmyrsveien 75 0462 Oslo 4-Norway.

Dr. Jon A. Rudbach,

Ribi ImmunoChem Research, Inc., P.O. Box 1409, Hamilton, MT 59840, 406-363-6214.

Dr. Jerald C. Sadoff,

Walter Reed Army Institute of Research, Washington, D.C. 20307.

Dr. Martha Sedegah,

Malaria Program, Naval Medical Research Institute, 12300 Washington Ave., Rockville, MD 20852, 301-295-2000.

Dr. M. Slaoui,

Molecular and Cellular Biology, Smith Kline Beecham Biologicals, 89 Rue de l'Institut 330, Rixemart, Belgium.

Dr. H. Snippe,

Medical Microbiology, Eijkman-Winkler Laboratory of Medical Microbiology, Medical School AZU G 04.614, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands.

Dr. P. Stuetz,

Sandoz Forschungsinstitut Gesellschaft M. B. H., Brunner Strasse

- 59, A-1235 Wien, Austria.
- Dr. Kuni Takayama,
Mycobacteriology Research Lab., William S. Middleton Memorial
Veterans Hospital, 2500 Overlook Terrace, Madison, WI 53705.
- Dr. Nobumasa (Nobi) Tanaka,
US-Japan Science Consulting Services, Inc., 72 Paxwood Road,
Delmar, NY 12054, 518-439-8334.
- Dr. Chao-Ming Tsai,
Bacterial products, Center for Biologics, FDA, 8800 Rockville
Pike, Bethesda, MD 20892, 301-496-2154.
- Dr. J. Terry Ulrich,
Infectious Disease Research, Ribic ImmunoChem Research, Inc.,
P.O. Box 1409, Hamilton, MT 59840, 406-363-6214.
- Dr. Stephanie Vogel,
Microbiology, Uniformed Services University of the Health
Sciences, 4301 Jones Bridge Road, Bethesda, MD 20814, 301-
295-3446/7.
- Dr. Frederick R. Vogel,
Lederle Biologicals, Lederle Laboratories, Lederle Laboratories
Bldg. 114/100, Pearl River, NY 10965, 914-732-4481.
- Dr. Shaw Warren,
Infectious Diseases, Massachusetts General Hospital, Gray 5,
Mass. General Hospital, Fruit St., Boston, MA 02114, 617-726-
5772.
- Dr. Nabila M. Wassef,
Membrane Biochemistry, Walter Reed Army Institute of Research,
Washington, D.C. 20307, 202-576-3528.
- Dr. Richard O. Williams,
Research and Development, IGEN, Inc., 1530 E. Jefferson St.,
Rockville, MD 20852, 301-984-8000.
- Dr. Scott Winston,
Univax Biologics Inc., 12111 Parklawn Drive, Rockville, MD
20852, 301-770-3099.
- Dr. James F. Young,
MedImmune, Inc., 19 Firstfield Road, Gaithersburg, MD 20878,
301-590-2642.

THIS DOCUMENT HAS NO MISSING PAGES. REPORT IS MISNUMBERED