Woods Hole Oceanographic Institution



SELECTE FEB 28 1992

Characterizing Acoustic Features of Marine Animal Sounds

by

Kurt M. Fristrup and William A. Watkins

January 1992

Technical Report

Funding was provided by NAVSEA under Contract No. N00140-90-D-1979 and a series of contracts and grants by ONR including Grant N00014-91-J-1445 with supplemental support by NOARL and ORINCON/DARPA.

Approved for public release; distribution unlimited.

92-04682

WHOI-92-04

Characterizing Acoustic Features of Marine Animal Sounds

bу

Kurt M. Fristrup and William A. Watkins

Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543

January 1992

Technical Report

Funding was provided by NAVSEA under Contract No. N00140-90-D-1979 and a series of contracts and grants by ONR including Grant N00014-91-J-1445 with supplemental support by NOARL and ORINCON/DARPA.

Reproduction in whole or in part is permitted for any purpose of the United States Government. This report should be cited as Woods Hole Oceanog. Inst. Tech. Rept., WHOI-92-04

Approved for public release; distribution unlimited.

Approved for Distribution:

Peter H. Wiebe, Chairman Department of Biology

Accesion For

NTIS CRA&I
DTIC TAB
Unannounced
Justification

By
Dist ibution/
Availability Codes

Dist Avail and for Special

Abstract

Software tools were designed to characterize the acoustic features of marine animal sounds. These have resulted in a set of calculated measurements that summarize particular aspects of sound sequences. The specificity of these measurements was enhanced by adjusting calculations to compensate for ambient noise. The sound measures included statistics for Aggregate Bandwidth, Intensity, Duration, Amplitude Modulation, Frequency Modulation, Short-term Bandwidth, Center Frequency, and Amplitude Frequency Interaction. The efficacy of noise compensation was tested for each statistic. Then, the sound measures were tested on a subset of 200 sequences of marine animal sounds, including sequences from 20 species: six baleen whales, 13 toothed species, and one seal. The statistics were reviewed for each species and a graphical comparison of all species was generated using principal components analysis. Preliminary results confirm that such sounds can be classified by means of relatively simple statistical algorithms, and we are encouraged to continue toward a system for automatic classification of marine animal sounds.

1 Introduction

Marine animals produce a remarkable variety of sounds (Watkins and Wartzok 1985). A primary goal of the bioacoustic program at the Woods Hole Oceanographic Institution (WHOI) has been to parse this variation into sensible classes of signals. Marine mammal sounds in particular contain distinctive features associated with species (op. cit.), individual identity (Caldwell, Caldwell and Tyack 1990), and certain behaviors. These features have never been examined in a broad context, comparing the sounds of a wide variety of species. Do the differences in these sound features remain distinctive as the scope of comparison widens? With our own ears, we can often distinguish acoustic features that appear to be species-specific, and sometimes features unique to individual animals; can we specify numerical algorithms that objectively recognize these distinctions?

The logistic requirements for addressing these questions have been formidable. To quantify the interspecific and intraspecific variability in marine animal sounds, a large number of sounds must be analyzed for each individual or species to be differentiated. Many biological and environmental attributes potentially explain acoustic variability. Therefore, these numeric results had to be referenced: species, population, group, social context, behavior, activity, individual identity, sex, reproductive situation, age, season, geographic location, water depth, and sound propagation. Thus, a necessary resource for such acoustic distinctions has been a system for integrating the sound sequences with associated biological and environmental data.

The SOUND database system organized for marine animal sounds (Watkins, Fristrup, and Daher 1991) has provided this resource. The databases and associated files contain thousands of digitized sound segments spanning more than seventy species recorded from all the world's oceans. The database describes the time, geographic location, recording conditions, identity of the animal(s) producing the sounds, the behavioral observations

associated with sound production, etc. These SOUND databases represent years of work by several people, and the analyses reported here depend on the availability of such. In turn, these analyses complemented and extended the capabilities of the database. New relational database structures have been implemented to permit flexible and convenient integration of these statistical results with the biological and environmental information about the sounds.

The quantification of time-frequency characters of the animal sounds for these analytic distinctions has had no precedent on this scale. No prior work has dealt with so many species and such a variety of repertoires from individual animals. The WHOI studies of marine animal acoustics, which have continued since William E. Schevill's work in the late 1940's, have provided the heuristic basis for these statistical decisions. We have learned to utilize many different acoustic features to describe and diagnose sounds. As a first step toward the development of an automatic, non-subjective system for separating the different animal sound sequences, we have devised statistical measures to recognize familiar acoustic features.

This report describes the numerical procedures that have been used, and it demonstrates their effectiveness with a trial set of 200 digitized sequences of marine mammal sounds. These preliminary results suggest that the gross acoustic features we analyzed can be useful indicators of species identity, and that with refinement they might provide the basis for finer distinctions.

2 The Statistics

Our statistical estimation techniques were based on our experience with the marine animal sounds in our tape library. We were guided by the following criteria:

- Each statistic was designed to emphasize particular parameters of animal sounds that we recognized as important for distinguishing species.
- Each statistic had to be insensitive to sound artifacts introduced by propagation in the ocean (multipath, fading, frequency-dependent attenuation, etc.).
- Most statistics needed to be relatively insensitive to noise and assumed a minimum
 of 15 dB signal/noise.
- Most statistics had to be insusceptible to the shape (relative frequency emphases) of the ambient noise power spectra.
- Most statistics needed to be related to obvious features in the time-frequency analysis
 displays of these sounds (duration, frequency range, etc.) so we could recognize the
 effectiveness of the statistics in making the discriminations.

These criteria reflected our interest in discriminating among the animal sounds rather than making selections that were largely controlled by differences in the ambient backgrounds. The choice of criteria did not take into consideration changes to the sounds contributed by the orientation and movements of sound sources. A number of other effects also have not been addressed, including means for dealing with frequency-dependent attenuation. Statistics that obviously would be sensitive to distortion of phase information have been avoided in these analyses.

The basic unit of data used for our feature extraction programs was one FFT (Fast Fourier Transform) block. For most files, this was 256 sample points, but for very short files

(low sampling rates) the FFT size was decreased to obtain no fewer than 16 blocks. Adjacent blocks did not overlap. A constant was subtracted from each sample point such that the mean for each block became zero, and the data were then tapered with a Hamming window. These choices eliminated explicit correlation between adjacent blocks and smoothed the resulting power spectra; the cost of this was a reduction in the degrees of freedom for our analyses.

The noise compensation technique begins by estimating the power spectrum of sounds that were present throughout the sound cut. To identify blocks of data containing only noise energy, intensity measures were computed for up to 600 blocks of data distributed evenly through a sound cut. Blocks were sorted by intensity, and the blocks between the fifth and tenth percentiles in level were used to form a noise power spectrum. We eliminated the bottom five percent to avoid using atypically quiet sections (tape dropout, etc.). During subsequent processing of these data, a multiple (currently 6.67x) of this noise spectrum was subtracted from each block's power spectrum (negative values set to zero). All spectral statistics were computed from this reduced power spectrum. To obtain the amplitude estimate for the block, the adjusted spectrum values were summed; this indirect method of computing amplitude, which exploits Parseval's relation (Oppenheim and Schafer 1989, p. 574), prevents loud noise components from dominating the amplitude statistics.

2.1 Abbreviations in Statistical Formulae

- t_i , time in seconds: the interval from the beginning of the sound cut to the beginning of the i^{th} FFT block.
- ss_i , relative intensity in arbitrary units: sum of the adjusted power spectrum values for the i^{th} block.
- $mm_i = min(ss_{i-1}, ss_i)$: the smaller of two adjacent ss values.

- \bullet F_{50} in Hertz: the frequency that bisects the area under the power spectrum density.
- N₅₀ in Hertz: the minimum number of frequency bins required to accumulate fifty percent of the total signal energy.
- F_{75} in Hertz: the highest frequency encountered when calculating N_{50} .
- F_{25} in Hertz: the lowest frequency encountered when calculating N_{50} .

Considerable use of symbols could not be avoided, but wherever possible we have used descriptive terms for ease of interpretation.

Table 1: Summary of Statistics

Duration: Results in Table 2 Total Duration 2	age			
Duration: Results in Table 2 Total Duration 2				
County Consendanting	8			
Sound Concentration 3	8			
Amplitude Modulation: Results in Table 3				
Amplitude Mean 4	9			
Amplitude Standard Deviation 5	9			
Attack Fraction 6	9			
Attack Proportion 7	10			
Amplitude Skewness 8	10			
Frequency Modulation: Results in Table 4				
Upsweep Mean 9	10			
Upsweep Fraction 10	11			
Upsweep Proportion 11	11			
Time Frequency Correlation 12	11			
Time Upsweep Correlation 13	12			
Short-term Bandwidth: Results in Table 5				
Short-term Bandwidth Mean 14	12			
Short-term Spectral Concentration 15	13			
Short-term Spectral Asymmetry 16	13			
Aggregate Bandwidth: Results in Table 6				
Total Upper Frequency - F_{75} 5	, 13			
Total Lower Frequency - F_{25} 5	, 13			
Total Spectrum Concentration - N_{50} 5	, 13			
Modal Upper Frequency - F_{75} 5	, 13			
Modal Lower Frequency - F_{25} 5	, 13			
Modal Spectrum Concentration - N_{50} 5	, 13			
Center Frequency: Results in Table 7				
Median Frequency Mean 17	14			
Total Spectrum Median Frequency				
Modal Spectrum Median Frequency				
Amplitude Frequency Interaction: Results in table 8				
Amplitude Frequency Correlation 18	14			
Amplitude Upsweep Correlation 19	14			

Intensity measures were computed for each block of data. These values were used as weights to find the "center" of the sound, defined as the weighted average of the time values (eq. 1). This statistic is only used as a reference point for subsequent calculations of Sound Duration (eq. 2) and Amplitude Skewness (eq. 8).

• Signal Center: weighted mean of t, ss as weights.

$$\bar{t} = \frac{\sum_{i=0}^{N} s s_i t_i}{\sum_{i=0}^{N} s s_i}$$
 (1)

2.2 Sound Duration

The Sound Duration was computed by Equation 2. It yielded a gross estimate of total duration, including any intervals of silence between sound elements.

• Sound Duration: weighted standard deviation of t, ss as weights.

$$4\sigma_t = 4\sqrt{\frac{\sum_{i=0}^{N} ss_i t_i^2}{\sum_{i=0}^{N} ss_i} - \bar{t}^2}$$
 (2)

The Sound Concentration was computed by Equation 3. It yielded an estimate of duration that would result if the sound were "packed" such that all silent sections were removed. It responds only to the relative amplitudes of different blocks, and it is insensitive to their ordering in the sound.

• Sound Concentration: equivalent statistical bandwidth of the amplitude values.

$$\frac{\left(\sum_{i=0}^{N} s s_{i}\right)^{2}}{\sum_{i=0}^{N} s s_{i}^{2}} \tag{3}$$

The ratio of these two duration estimates can be used to measure duty cycle.

2.3 Amplitude Modulation

A reference value was computed for the average level of the sound: the Amplitude Mean was computed by Equation 4.

• Amplitude Mean: average ss value.

$$\overline{ss} = \frac{\sum_{i=0}^{N} ss_i}{N} \tag{4}$$

The Amplitude Standard Deviation (Eq. 5) can be used to measure the average magnitude of amplitude modulation. However, Amplitude Mean and Amplitude Standard Deviation are scaled arbitrarily by idiosyncrasies of the digitizing process. To form a useful diagnostic, they must be used together to form a scale-independent measure like the coefficient of variation (Eq. 5/ Eq. 4).

• Amplitude Standard Deviation: standard deviation of ss values.

$$\sigma_{ss} = \sqrt{\frac{\sum_{i=0}^{N} ss_{i}^{2}}{N} - \left(\frac{\sum_{i=0}^{N} ss_{i}}{N}\right)^{2}}$$
 (5)

The occurrence and magnitude of sections of sound with increasing and decreasing amplitudes was measured by assessing the fraction of blocks in which a subsequent block had a larger (or lower) amplitude than the current block. This Attack Fraction was computed by Equation 6.

• Attack Fraction: the fraction of blocks in which subsequent block has a larger ss value than the current block.

$$\frac{\sum_{ss_{i+1}>ss_i}1}{N} \tag{6}$$

The proportion of average changes in amplitude to the sum of the average increases and decreases in amplitude values was calculated and called the Attack Proportion. These statistics (Eq. 6 and Eq. 7) were similar in function to the "attack" and "decay" terms used to refer to the initial and terminal amplitude modulation of individual notes in usic. The attack proportion was computed by Equation 7.

 Attack Proportion: the proportion of average increase in ss values relative to the sum of average increases and decreases in ss values.

$$\frac{\sum_{\Delta s s_{i} > 0} \Delta s s_{i}}{\sum_{\Delta s s_{i} > 0} \Delta s s_{i} - \sum_{\Delta s s_{i} < 0} \Delta s s_{i}}$$

$$(7)$$

Gross asymmetry in the amplitude modulation relative to the "center" of the sound was weighted by a function of time and amplitude. This Amplitude Skewness was computed by Equation 8.

• Amplitude Skewness: weighted skewness of t, ss as the weights.

$$\frac{\sum_{i=0}^{N} ss_{i}t_{i}^{3}}{\sum_{i=0}^{N} ss_{i}} - 3\bar{t}\frac{\sum_{i=0}^{N} ss_{i}t_{i}^{2}}{\sum_{i=0}^{N} ss_{i}} + 2\bar{t}^{3}}{\sigma_{i}^{3}}$$
(8)

2.4 Frequency Modulation

The frequency modulation of a sound was expressed by differences between median frequencies of adjacent power spectrum estimates. The average upsweep trends (downsweep is a negative upsweep) for entire sound cuts were calculated by comparing weighted averages of the change in frequency and amplitude values, giving the Upsweep Mean using Equation 9.

• Upsweep Mean: weighted average of the change in F_{50} values, mm as weights.

$$\frac{\sum_{i=1}^{N} m m_i \Delta F_{50i}}{\sum_{i=1}^{N} m m_i} \tag{9}$$

Estimates of the relative occurrence of frequency modulation were calculated by determining the fraction of total energy that coincided with increases in frequency. This Upsweep Fraction was computed by Equation 10.

• Upsweep Fraction: fraction of summed mm values that coincide with increases in F_{50} .

$$\frac{\sum_{\Delta F_{50i} > 0} mm_i}{\sum_{i=1}^{N} mm_i} \tag{10}$$

The relative magnitudes of frequency upsweeps and downsweeps were calculated as the Upsweep Proportion, using Equation 11.

• Upsweep Proportion: proportion of average weighted increase in F_{50} to the sum of the weighted average increases and decreases in F_{50} .

$$\frac{\sum_{\Delta F_{50i}>0} m m_i \Delta F_{50i}}{\sum_{\Delta F_{50i}>0} m m_i} \frac{\sum_{\Delta F_{50i}>0} m m_i}{\sum_{\Delta F_{50i}>0} m m_i \Delta F_{50i}} \frac{\sum_{\Delta F_{50i}<0} m m_i \Delta F_{50i}}{\sum_{\Delta F_{50i}>0} m m_i} \frac{\sum_{\Delta F_{50i}<0} m m_i}{\sum_{\Delta F_{50i}<0} m m_i}$$
(11)

The magnitudes of linear relationships between time and median frequency were estimated by calculating correlation coefficients. This correlation used intensity values as weights to focus on portions of the sound cuts with loud signals. This was computed as a Time Frequency Correlation by Equation 12.

• Time Frequency Correlation: weighted correlation between F_{50} and t, ss as weights.

$$\frac{\sum_{i=0}^{N} \frac{ss_{i}t_{i}F_{50i}}{\sum_{i=0}^{N} \frac{ss_{i}}{i}} - \overline{tF_{50}}}{G_{4}G_{5}}$$
(12)

where

$$\sigma_F = \sqrt{\frac{\sum_{i=0}^{N} ss_i F_{50i}^2}{\sum_{i=0}^{N} ss_i} - \overline{F_{50}}^2}$$

The magnitudes of linear relationships between time and frequency upsweep were also estimated by a weighted correlation coefficient. This Time Upsweep Correlation was computed by Equation 13.

• Time Upsweep Correlation: weighted correlation between ΔF_{50} and t, mm as weights.

$$\frac{\sum_{i=1}^{N} \frac{mm_{i}t_{i}\Delta F_{50i}}{\sum_{i=1}^{N} \frac{mm_{i}}{\tau_{t}\tau_{F}}} - \hat{t}\hat{F}_{50}}{\tau_{t}\tau_{F}}$$
(13)

Where

$$\hat{t} = \frac{\sum_{i=1}^{N} m m_i t_i}{\sum_{i=1}^{N} m m_i}, \ \hat{F_{50}} = \frac{\sum_{i=1}^{N} m m_i \Delta F_{50i}}{\sum_{i=1}^{N} m m_i}$$

$$\tau_t = \sqrt{\frac{\sum_{i=1}^{N} m m_i t_i^2}{\sum_{i=1}^{N} m m_i} - \hat{t}^2}, \ \tau_F = \sqrt{\frac{\sum_{i=1}^{N} m m_i \Delta F_{50i}^2}{\sum_{i=1}^{N} m m_i} - \hat{F}_{50}^2}$$

2.5 Short-term Bandwidth

Two measures of bandwidth were computed for each block. The gross spread in power spectral values (frequency) was calculated to give the Short-term Bandwidth Mean by Equation 14.

• Short-term Bandwidth Mean: weighted average of $F_{75} - F_{25}$, ss as weights.

$$\frac{\sum_{i=0}^{N} ss_i(F_{75i} - F_{25i})}{\sum_{i=0}^{N} ss_i}$$
 (14)

The effective number of frequency bins in each block containing significant power levels (ignoring gaps in the sound) also was computed to give the Short-Term Spectral Concentration by Equation 15.

• Short-term Spectral Concentration: weighted average of N₅₀, ss as weights.

$$\frac{\sum_{i=0}^{N} ss_i N_{50i}}{\sum_{i=0}^{N} ss_i} \tag{15}$$

Then, the relative emphasis of sideband energy on either side of the dominant frequency was estimated by calculating the Short-term Spectral Asymmetry using Equation 16.

• Short-term Spectral Asymmetry: weighted average spectral asymmetry, with ss as weights.

$$\frac{\sum_{i=0}^{N} ss_{i} \frac{F_{50i} - F_{25i}}{F_{75i} - F_{25i}}}{\sum_{i=0}^{N} ss_{i}}$$
 (16)

2.6 Aggregate Bandwidth

Two aggregate power spectra were computed for each sound cut: the Total Spectrum was the average of all FFT power spectra for the sound, and the Modal Spectrum accumulated only the power spectrum magnitudes for the frequency bin with the largest value in each FFT block. Both aggregate spectra were processed to extract three statistics related to bandwidth. The Upper (F_{75}) and Lower (F_{25}) Frequencies estimated the bounds of the aggregate spectra. These could be used to compute a bandwidth spanning any gaps in the spectral density. The Spectrum Concentration (N_{50}) provided an estimate of a "packed" or gap-free bandwidth.

2.7 Center Frequency

Three statistics were used to estimate the aggregate center frequency, or "average" frequency, of the entire sound cut. They were the F_{50} values computed for the Total and Modal Spectra, and a weighted average of the instantaneous F_{50} values computed by Equation 17. These statistic produced very similar values in most instances.

• Median Frequency Mean: weighted average of F₅₀, ss as weights.

$$\overline{F_{50}} = \frac{\sum_{i=0}^{N} ss_i F_{50i}}{\sum_{i=0}^{N} ss_i}$$
 (17)

2.8 Amplitude-Frequency Interaction

The strength of a linear trend between amplitude and center frequency was computed as the Amplitude-Frequency Correlation by Equation 18.

• Amplitude Frequency Correlation: correlation between ss and F₅₀.

$$\frac{\sum_{i=0}^{N} ss_{i} F_{50i} - \overline{ss} \sum_{i=0}^{N} F_{50i}}{\sigma_{ss} \sqrt{N \sum_{i=0}^{N} F_{50i}^{2} - \left(\sum_{i=0}^{N} F_{50i}\right)^{2}}}$$
(18)

The magnitude of a linear trend between amplitude and frequency modulation was calculated as the Amplitude Upsweep Correlation, using Equation 19.

• Amplitude Upsweep Correlation: correlation between mm and ΔF_{50} .

$$\frac{\sum_{i=1}^{N} m m_i \Delta F_{50i} - \overline{m} \overline{m} \sum_{i=1}^{N} \Delta F_{50i}}{\sigma_{mm} \sqrt{N \sum_{i=1}^{N} \Delta F_{50i}^2 - \left(\sum_{i=1}^{N} \Delta F_{50i}\right)^2}}$$
(19)

where

$$\overline{mm} = \frac{\sum_{i=1}^{N} mm_i}{N}, \ \sigma_{mm} = \sqrt{\frac{\sum_{i=1}^{N} mm_i^2}{N} - \overline{mm}^2}$$

3 Noise Performance

To explore the effectiveness of our system of noise compensation, test sounds were copied and contaminated with increasing levels of noise. These test files were processed to generate plots of each statistic relative to signal/noise ratios. The differences between adjacent points along each curve reflected estimation error, because adjacent points represented different noise sequences with similar signal/noise ratios. Any overall trend in these plots reflected imperfect noise compensation.

Against Gaussian white noise, these noise tests were successful. However, the density of this type of noise was favorable for other noise compensation algorithms as well, and we were reminded that noise backgrounds in the ocean were rarely "white" (equal energy at all frequencies). Therefore, noise was generated synthetically to resemble the ambient background for many of our recordings at sea using MATLAB software (The Math Works, Inc.). Parametric spectral estimation procedures extracted parameters for a sixth order autoregressive (AR) process from an ambient noise sample. Noise sequences were generated by filtering white noise with a finite impulse response (FIR) filter constructed from the AR estimates.

The sound statistics were tested against noise and plotted in Figures 1 through 27 to provide an indication of performance for each statistic. In these figures, the horizontal axis represented signal/noise ratios, proceeding from low to relatively high values. Note that most of our marine mammal sound sequences exceeded 15 dB signal/noise. The vertical axes represented the estimated numeric level for this statistic. The variance may be seen in the relative amplitude fluctuations on the vertical axes, and the trend for this statistic may be seen in the relative changes in the progression from lower to higher signal/noise along horizontal axes.

The performance of each of the statistics relative to noise was summarized in the table

below, with the plots subjectively classified as follows.

Table of Noise Performance for the Statistics

	Low Variance	High Variance
	Signal Center (Fig. 1)	Attack Fraction (Fig. 6)
	Median Frequency Mean (Fig. 25)	Attack Proportion (Fig. 7)
1	Upsweep Mean (Fig. 9)	Upsweep Proportion (Fig. 11)
	Time Frequency Correlation (Fig. 12)	Short-term Spectral Asymmmetry (Fig. 16)
į.	Time Upsweep Correlation (Fig. 13)	Amplitude Skewness (Fig. 8)
small	Upsweep Fraction (Fig. 10)	
trend	Total Spectrum Median (Fig. 23)	
	Modal Lower Frequency (Fig. 21)	
	Modal Spectrum Median (Fig. 24)	
	Modal Upper Frequency (Fig. 20)	
	Sound Duration (Fig. 2)	Amplitude Frequency Correlation (Fig. 26)
	Sound Concentration (Fig. 3	Amplitude Upsweep Correlation (Fig. 27)
large	Amplitude Mean (Fig. 4)]
trend	Amplitude Standard Deviation (Fig. 5)	j
	Short-term Bandwidth Mean (Fig. 14)	Ì
	Short-term Spectral Concentration (Fig. 15)	
ll i	Total Upper Frequency (Fig. 17)	
1	Total Lower Frequency (Fig. 18)	
	Total Spectrum Concentration (Fig. 19)	
	Modal Spectrum Concentration (Fig. 22)	

Generally, the statistics performed well in noise. Higher order statistics (standard deviations, correlations) were less consistent, and the least useful were those statistics measuring frequency-amplitude relations and gross asymmetry in the sound waveform envelope.

4 Preliminary Analysis of Marine Animal Sounds

A subset of approximately 200 sequences of marine mammal sounds were used to test these statistics. Sounds were selected without attention to any particular acoustic features. They included sequences from 20 species: six baleen whales, 13 toothed whales and dolphins, and one seal. A wide variety of sound types was included in this subset; we also included sounds from pairs of species that were difficult to distinguish aurally.

4.1 Statistical Interdependence

The redundancy in these statistics, for this data set, was examined by a stepwise multiple regression procedure. This analysis treated each sound equally, ignoring the identity of the sequence. At each stage, the algorithm identified the statistic that had the highest linear correlation with other statistics for these data. This statistic was removed, and the analysis repeated. When the correlation coefficients and scatter plots indicated relatively poor fits, the analysis was terminated. We anticipated some redundancy in our statistics; we intended to test alternative sound measures. A more conclusive analysis of redundancy and performance awaits analysis of larger data sets.

For the comparisons described here, the multiple regression functions explained more than 80% of the variance for the first nine statistical estimators. The statistics that were successively eliminated were (with percent explained variance): Amplitude Standard Deviation (0.979, fig. 28), Median Frequency Mean (0.974, fig. 29), Modal Spectrum Median (0.974, fig. 31), Total Upper Frequency (0.944, fig. 31), Total Spectrum Concentration (0.900, fig. 32), Modal Lower Frequency (0.896, fig. 33), Modal Upper Frequency (0.885, fig. 34), Short-term Spectral Concentration (0.878, fig. 15), and Total Spectrum Median (0.839, fig. 36). In these figures, the horizontal axes represented the predicted value, the vertical axes represented the observed value, and the dark line represented the regression

line. Eight of the nine variables appear to scale with center frequency, and we suspect that re-expressing these in relation to center frequency would remove much of this redundancy. Amplitude Standard Deviation and other amplitude variables sould be re-expressed relative to Amplitude Mean for similar reasons. It was not resolved whether simple division by these scaling factors would be appropriate.

4.2 Acoustical Analyses and Biological Information

The ability to select and analyze acoustic measurements based on related biological or environmental observations was crucial for these data. This could have been done by segregating data files for different species, activities, locations, etc. and independently processing each batch. However, it would have been cumbersome and difficult to manage such sorting and data segregation for each new query, especially as the selections became more complicated. A more powerful technique was to link the numerical analyses directly to the text databases. All sound cuts were processed in one batch, and these extensive computations proceeded automatically, unattended. Interactive exploration of relationships among statistics and biological or environmental factors followed, with all of the flexibility and convenience of database queries and reports.

The SOUND text databases for the recordings and the digital sound sequences (Watkins, Fristrup, and Daher 1991) could have accommodated new numeric data from the statistical analyses, but with INMAGIC software this required restructuring the entire database each time the number of numeric fields changed. This was not feasible: the analyses required many iterations and modifications. Therefore, PARADOX software (supports relational database models, with visual, query-by-example interface) was used to provide more flexible linkage between biological and acoustic information. The text information from the SOUND databases remained unmodified as a single table, and additional tables were created to handle the numeric results. Fields were used in these numeric tables that identified

the related SOUND text records. Then, subsets of the statistical results were obtained by selecting particular fields in SOUND and reporting the linked numerical information. Note that these queries could be reversed to select pertinent biological or environmental information based on acoustic criteria.

4.3 Preliminary Species Summaries

We generated summaries of the numerical results for each of the species in the trial data set (see Tables 2 through 8). For each species, the mean value of the statistic was listed with the maximum as a superscript and the minimum as a subscript. The number of sounds analyzed (Count) for each species was indicated in Table 1, but not repeated in the other tables. Tables were divided in three sections: baleen whales, toothed species, one seal and a transient sequence (hammer simulating clicks of *Physeter catodon*).

Two aspects of the summaries of sound duration (Table 2) merit comment. Species that are represented by more than ten sounds showed a dramatic variation between upper and lower bounds on both statistics. This may have reflected different selection criteria for the sound cuts. These sound cuts may have been a mix of isolated sounds, long sound sequences with intervals of silence, or continuous choruses from many individuals. All of these are valid data, but we need to differentiate among these classes of recordings in future analyses. Also, the sound "duty cycle" could be calculated by dividing Sound Concentration by Sound Duration. Table 2 indicates that baleen whales could largely be distinguished from toothed species by comparison of sound duty cycles.

Table 2						
Species	Count	Sound	Sound			
		Duration s.	Concentration s.			
B. mysticetus	5	1.62 2.50	.403 .782			
C. marginata	3	.86 .83	$.297_{-217}^{+000}$			
E. glacialis	5	.83 1.27	.265 :639 :066			
E. australis	16	$.82_{.18}^{2.39}$.443 ^{1.319}			
B. acutorostrata	5	$.51^{.70}_{.27}$.255 .403			
B. physalus	15	$1.07_{0.00}^{4.80}$.332 ^{.829} _{.100}			
P. catodon	28	4.15 69.97	.057 :172			
D. leucas	1	$1.17_{1.17}^{1.17}$	$.517^{:517}_{:517}$			
S. longirostris	25	$1.39_{.10}^{2.63}$.089 ^{.256}			
S. $long. + P. cat.$	4	$1.29^{1.88}_{.76}$	$.079^{ \cdot 127}_{ \cdot 023}$			
S. bredanensis	7	$1.72_{.98}^{\:2.41}$.230 :384 :095			
C. commersonii	2	$1.46_{.61}^{2.31}$	$.102^{.142}_{.061}$			
D. delphis	13	$1.68_{.66}^{4.97}$	$.182_{.020}^{1.147}$			
G. griseus	23	$4.06_{-17}^{8.32}$	$.414_{.013}^{1.295}$			
G. macrorhynchus	11	$1.68_{.62}^{4.47}$.403			
G. melaena	11	$1.37_{.14}^{8.10}$	$.253_{.039}^{1.584}$			
O. orca	7	.98 ^{1.83}	$.393^{+921}_{-040}$			
P. crassidens	6	$5.94_{3.23}^{9.79}$	$.470_{.077}^{1.098}$			
P. phocoena	8	$1.54^{3.48}_{.47}$	$.066^{ +125}_{ +030}$			
I. geoffrensis	2	$1.31_{1.02}^{1.60}$.121 :132			
A. phillipi	4	4.43 7.26 1.99	$.670_{.235}^{1.019}$			
Hammer on metal	1	$2.41_{\ 2.41}^{\ 2.41}$.079 :079			

Table 3 displayed statistics related to amplitude modulation. The first two of these statistics were not useful diagnostics in themselves because the absolute value of each was inherently tied to equipment gain settings during any stages of processing. However, the proportion of Amplitude Standard Deviation/Amplitude Mean (standard error) was a useful indication of amplitude modulation, with larger standard errors indicating more modulation.

Attack Fraction and Attack Proportion were negatively correlated because sound cuts were edited so that initial and terminal noise levels were approximately the same. One of these probably would have been sufficient.

Amplitude Skewness appeared to be a less robust statistic. However, some gross differences agreed with our experience. Among baleen whales, *Eubalaena glacialis* tended to start loudly and taper to silence (negative skew), while *B. acutorostrata* often started softly and swelled in level.

Table 3

	Table 3			
A mplitude	Amplitude	Attack	Attack	Amplitude
Mean	StdDev	Fraction	Proportion	Skewness
.095 :265	.336 .889	.51 .57	.49 .51	.439 1.643
$.161^{:331}_{:028}$	$.291^{.548}_{.062}$.44 :56 .38	.56 .63	$140rac{.016}{289}$
$.195^{.513}_{.011}$	$.501_{.057}^{1.584}$.51 .57	.49 .55	$-1.127 \substack{.486 \\ -5.160}$
$1.254_{.034}^{5.651}$	$2.263_{.125}^{7.928}$.50 ^{.69}	$.50_{-31}^{.60}$	$232_{-1.391}^{2.658}$
$19.177_{\ 4.948}^{\ 38.858}$	35.734 64.134 12.505	.45 .56	.55 . ⁷⁵	$1.123_{.339}^{1.648}$
$1.612_{.000}^{7.413}$	$3.383_{.001}^{12.925}$	$.50^{.75}_{.29}$.56 .38	$.444_{-2.211}^{3.828}$
.077 :310	.442 ^{1.565}	.46 .57	.54 .68	$226_{-2.052}^{.885}$
$.010^{.010}_{.010}$	$.021^{.021}_{.021}$	$.52_{-52}^{+52}$.48 .48	.678 :678
$.143_{.000}^{1.387}$	$.427_{.000}^{3.700}$.50 .56 .45	.50 .55	$.131_{-3.049}^{4.023}$
$.010^{.016}_{.004}$	$.046_{.032}^{.058}$	$.51^{.53}_{.48}$.49 . 5 2	$392^{2.632}_{-5.964}$
$.113^{.383}_{.014}$	$.298_{.048}^{1.023}$	$.49_{-48}^{+51}$	$.51^{.52}_{.49}$	$.369 {}^{3.189}_{575}$
$.008^{+012}_{-004}$	$.043^{.056}_{.030}$	$.58_{-54}^{-62}$.42 :46 :38	$-5.200{}^{840}_{-9.560}$
$.126^{+430}_{-004}$	$.484_{.022}^{1.560}$.50 . 54	.49 : ₃₉	$041_{-4.224}^{4.272}$
$.060_{.001}^{.299}$	$.163^{ +622}_{ -007}$.49 .53	.51 .55	$.038_{-3.067}^{2.064}$
.381 ^{1.079}	$.952_{.090}^{2.785}$.50 : ⁵⁷	.50 . 54	$.017_{-1.606}^{2.366}$
$.726_{.153}^{1.757}$	$1.653^{3.603}_{.377}$	$.50^{.57}_{.42}$.50 : ⁵⁸	$.192_{-2.607}^{2.065}$
$.228^{ +451}_{ .011}$	$.384_{.048}^{+610}$	$.50^{.56}_{.46}$.50 . 54	$.280_{360}^{ 1.348}$
$.084^{:331}_{:001}$	$.231^{.682}_{.006}$.49 ^{.53}	$.51^{.54}_{.47}$.030 .984 840
$.171^{.933}_{.000}$	$.944_{001}^{5.185}$	$.48^{.56}_{.39}$	$.42^{.56}_{.09}$	$106{}^{3.919}_{-5.415}$
$.217^{:244}_{:190}$	$1.007^{1.140}_{.874}$.43 :44	.57 ^{.58}	$992{}^{819}_{-1.166}$
$.566_{.161}^{1.138}$	$1.359_{.585}^{2.517}$.51 :54	.49 .52	.081301
$1.489_{1.489}^{1.489}$	$10.214 \substack{10.214 \\ 10.214}$.40 :40	.60 .60 .60	$192{}^{192}_{192}$
	Mean .095.265 .013 .161.331 .161.328 .195.513 .195.513 1.254.5651 1.254.5651 19.177 38.858 1.612.7413 .000 .077.310 .077.310 .010.010 .143 1.387 .000 .010.016 .010.000 .010.016 .010.000 .010.016 .010.000 .010.000 .010.00000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.00000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.0000 .010.00000 .010.00000 .010.00000 .010.00000 .010.00000 .010.00000 .010.000000 .010.000000 .010.00000000	Amplitude Amplitude Mean StdDev .095 .265 .263 .336 .889 .336 .889 .291 .548 .291 .062 .195 .513 .291 .062 .195 .513 .501 1.584 .263 .125 .263 .125 .263 .125 .263 .125 .263 .125 .263 .125 .125 .263 .125 .263 .125 .263 .125 .263 .125 .263 .125 .263 .125 .263 .125 .263 .263 .263 .263 .263 .263 .263 .263	Amplitude Amplitude Attack Mean StdDev Fraction .095 .265 .013 .336 .889 .01 .51 .57 .47 .57 .47 .51 .57 .42 .51 .51 .54 .51 .51 .54 .51 .51 .51 .54 .51 .51 .51 .51 .51 .51 .51 .51 .51 .51	Amplitude Amplitude Attack Attack Attack Mean StdDev Fraction Proportion .095 .265 .336 .889 .51 .57 .49 .51 .161 .331 .291 .548 .44 .56 .56 .63 .195 .513 .501 .057 .51 .57 .49 .55 .1254 .6341 2.263 .125 .50 .69 .50 .60 .19.177 38.858 35 .734 64.136 .45 .56 .55 .75 .1612 .7003 3.383 .001 .50 .75 .56 .92 .077 .310 .442 .1565 .46 .37 .54 .42 .010 .010 .021 .021 .52 .52 .48 .48 .143 .000 .427 .000 .50 .56 .50 .55 .010 .016 .046 .058 .51 .53 .49 .57 .113 .383 .298 .048 .49 .51 .51 .52 .008 .001 .004 .032 .51 .53 .42 .46 .008 .001 .043 .036 .58 .54 .42 .46 .060 .299 .163 .622 .50 .54 .49 .38 .726 .1.757

Table 4 displays statistics relating to frequency modulation. Baleen whales and toothed species clearly had different magnitudes of Upsweep Mean, but much of this could have resulted from higher center frequencies of most toothed whale sounds.

Upsweep Fraction and Upsweep Proportion were not always negatively correlated, because the sounds from marine mammals could start or end at very different frequencies. The numbers for *E. glacialis* were instructive. Upsweep Mean was negative for this species, indicating that the sounds had lower frequencies toward the end, and Upsweep Fraction was nearly half indicating that about half of the block-to-block changes in frequency were positive. Thus, most of these downsweeps were greater in magnitude than the upsweeps, and indeed the Upsweep Proportion was less than one-half.

The Time Frequency and Time Upsweep Correlations showed considerable variation within larger samples. We know that the features they target are useful diagnostics, so we must seek better means of measuring them. Alternatives include non-parametric measures of correlation.

Table 4

Species	Upsweep	Upsweep	Upsweep	Time-Freq.	Time-Upsweep
•	Mean Hz/s.	Fraction	Proportion	Correlation	Correlation
B. mysticetus	-395^{18}_{-1149}	.47 .58	.50 : ₃₈	24 ^{.42}	.02 -24
C. marginata	-144^{-107}_{-191}	$.00^{+01}_{-00}$.68 :88	$27_{45}^{.01}$	$.18_{22}^{.50}$
E. glacialis	-1320_{-3879}^{145}	.54 .99	.34 :56	$17^{.35}_{64}$	$.10^{.32}_{03}$
E. australis	102_{-92}^{553}	.69 ^{1.00}	$.57_{.20}^{1.00}$	$.18_{98}^{.75}$	$.16_{15}^{.62}$
B. acutorostrata	0 98	.72 :97	.35 :50	$08_{63}^{.73}$	$04^{.33}_{41}$
B. physalus	-36_{-134}^{12}	$.17^{.94}_{0.00}$	$.32_{0.00}^{.91}$	$10_{-1.0}^{1.00}$	$12^{.86}_{98}$
P. catodon	45628 799608 -962976	.58 1.00	$.52_{0.00}^{1.00}$	$13^{.74}_{76}$	$03^{.72}_{63}$
D. leucas	-680^{-680}_{-680}	.18 .18	.69 :69	$82{}^{82}_{82}$	$02{}^{02}_{02}$
S. longirostris	$172907 {}^{3788800}_{-1082692}$	$.70_{.09}^{1.00}$	$.59_{-16}^{1.00}$	$00^{.89}_{89}$.0432
S. $long. + P. cat.$	$81780_{-115619}^{477717}$.54 .83	.29 :41	$40{}^{02}_{90}$	$03^{.09}_{13}$
S. bredanensis	$41495 {}^{331392}_{-41335}$.56 .63	$.45^{:51}_{.36}$	$04_{16}^{.09}$	$02_{13}^{.05}$
C. commersonii	-6998^{-790}_{-13207}	$.17^{ 26}_{ 08}$	$.37^{:50}_{:25}$	$.10^{ .42}_{22}$	$41{}^{30}_{51}$
D. delphis	$-70028{}^{81760}_{-788441}$	$.58_{.05}^{1.00}$	$.58_{.22}^{1.00}$.00 ^{.66}	$.00_{38}^{ .23}$
G. griseus	4432 52320 -9104	.55 :73	$.49 \substack{.80 \\ .22}$	$.04 \frac{.87}{45}$	$02^{.16}_{27}$
G. macrorhynchus	10434_{-7471}^{57709}	.51 :74	$.51^{.79}_{.28}$	$04_{66}^{.70}$	$01_{35}^{.11}$
G. melaena	2431_{-11684}^{29658}	.51 :88	.49 : ⁷⁸	.01 ^{.96}	$08^{.17}_{47}$
O. orca	$-129211_{-896083}^{2504}$	$.48^{.71}_{.09}$.43 :68 :25	$05_{40}^{.41}$	$16_{90}^{.05}$
P. crassidens	-161_{-2131}^{1082}	.55 .65	.38 :62	.45 : ⁷⁹	$00^{.04}_{11}$
P. phocoena	8550_{-2429}^{46219}	$.68_{0.00}^{1.00}$	$.70_{0.00}^{1.00}$	$.02_{99}^{1.00}$	$09^{.16}_{36}$
I. geoffrensis	$-43940{}^{-6242}_{-81638}$.35 .41	.54 .55	$15{}^{09}_{21}$.0402
A. phillipi	-1633^{-6}_{-4367}	.23 .44	.45 :70	.0602	.0202
Hammer on metal	$-503277 {}^{-503277}_{-503277}$.01 :01	.20 :20	36 ³⁶	.48 .48

The short-term bandwidth statistics in Table 5, the aggregate bandwidth statistics in Table 6, and the center frequency statistics of Table 7 were the most diagnostic for this set of sound sequences. They apparently separated the sounds of different species. Bandwidths appeared to scale with frequency, suggesting that they could be expressed best in proportion to center frequency.

Table 5						
Species	Short-term	Short-term	Short-term			
•	Bandwidth	Spectral	Spectral			
	Mean Hz/s.	Concentration Hz/s.	Asymmetry			
B. mysticetus	34502 39597 32023	3016 3756 2592	.11 .22			
C. marginata	881 949 810	106_{99}^{118}	$.17^{+25}_{-08}$			
E. glacialis	7444 19157 3200	335_{200}^{572}	$15^{.36}_{0.00}$			
E. australis	4378 6773 3269	417 ⁶⁸⁰ ₂₆₅	$.22_{0.00}^{40}$			
B. acutorostrata	907 1094	102_{74}^{158}	$.20^{.54}_{0.00}$			
B. physalus	464 ⁹⁴⁵	40 ⁹⁶	.06 :29			
P. catodon	1898178 6104578 244488	135440 339051	.41 :76			
D. leucas	52254 52254	4652 4652	.13 :13			
S. longirostris	$1189241 {}^{2556271}_{821087}$	$102772{}^{316991}_{51781}$	$.12^{.31}_{.00}$			
S. $long. + P. cat.$	3387894 6005715	$221269 {}^{403662}_{54720}$.22 :42			
S. bredanensis	3454478 7664267 1336456	$166331{}^{312275}_{106172}$	$.22^{.31}_{.15}$			
C. commersonii	30991 33563 28419	2471_{2176}^{2765}	.04 ^{.05}			
D. delphis	$1266464 {}^{3159867}_{132364}$	$102935{}^{336716}_{13007}$	$.13^{.32}_{.01}$			
G. griseus	508874 2462444 57730	$26081{}^{67239}_{5404}$.24 ^{.64} .08			
G. macrorhynchus	721235 1932495 129749	38673_{11581}^{102974}	.25 :49			
G. melaena	327154 653042 220740	27724 48173 17660	.18 :41			
O. orca	214199 322328 14074	17242_{1113}^{27372}	.19 :46			
P. crassidens	200492 430008 113113	20372 47612 9063	.14 :23			
P. phocoena	39577_{12800}^{126412}	3815_{800}^{10007}	$.17^{.47}_{0.00}$			
I. geoffrensis	1102209 1576500 627918	47188 56594 37782	.49 .52			
A. phillipi	137311 164278	12491 14300	.18 :20			
Hammer on metal	3383956 3383956 3383956	119674_{119674}^{119674}	.36 .36			

Table 6

		Total Spectrum			Modal Spectrum		
Species	Upper	Lower	Spectrum	Upper	Lower	Spectrum	
	Frequency	Frequency	Concentration	Frequency	Frequency	Concentration	
	Hz	Hz	Hz	Hz	Hz	Hz	
B. mysticetus	821 250	545 1438	149 302	903 2578	558 1438	114 32	
C. marginata	93 ⁹⁹	79 8 0	15 19 11	91 94 90	76 80	10 ½7	
E. glacialis	361_{200}^{717}	156 300	48 119	314 40°	158 310	27 55	
E. australis	174 ²⁴⁰	124 208	38 ⁶⁹	164 ²⁴⁰	132_{20}^{219}	22 44 10	
B. acutorostrata	89 130	70 92 28	15 ²⁴	88 130	72 100 30	9 15	
B. physalus	74 ¹⁴⁰	58 129 20	6 35	74 140	67 130	4 10	
P. catodon	6463 17100	1411 8480	1385 4044	3093 10240	1919 9734	215 610	
D. leucas	2563 2563	2266_{2266}^{2266}	298 ²⁹⁸	2517 2517 2517	2266 2266	251 251 251	
S. longirostris	11329_{640}^{27200}	6987 35280	1009_{174}^{2569}	11050_{640}^{26243}	7166_{320}^{25920}	757 2209	
S. long. + P. cat.	20036 30080	8640 10560 4480	2105 3778	$11258{}^{13120}_{10207}$	8640 10560 4480	674 1256	
S. bredanensis	15250_{7045}^{22400}	4118_{320}^{7360}	2237 5306 751	8849 15040	5517 8202	798 1315	
C. commersonii	3574 3750 3398	2785 3683	77 ¹⁰⁹	3574 3750	3503 3685	47 57 36	
D. delphis	16484_{5120}^{37153}	9127_{1753}^{15040}	1284 3212	13598 27200	9097 15306	807 ²⁰⁰³	
G. griseus	8682 13945	4185_{117}^{12344}	903 2221	7815 13594	4440 12344	525 1279	
G. macrorhynchus	5941 11840	2186 3447	659 1373	5064 11680	2586 6553	356 979 109	
G. melaena	$5707 {}^{8715}_{2880}$	4586 8451 1992	556 1037	5599 8640 2880	4729 8442 1992	317 542	
O. orca	1936_{800}^{3394}	949 2064	336 557	1635 2488	1067 ²⁰⁶²	189 389	
P. crassidens	6891_{234}^{13359}	5184 7266 117	473 698	6706 13359	5219 7284 117	313 579	
P. phocoena	1522 3080	610_{0}^{2480}	$110^{\frac{231}{25}}$	660 2560	619_{0}^{2520}	28 ³⁹ 23	
I. geoffrensis	5291 7734 2847	$1875{}^{1992}_{1758}$	1143 1548 738	3379 3946 2813	2184 2493 1875	278 301 256	
A. phillipi	234 234	117 117	100 113	234 234	117 117	67 76	
Hammer on metal	1440_{1440}^{1440}	640 640	800 800 800	1111 1111	800 800 800	210 210	

Table 7						
Species	Median	Total	Modal			
	Frequency	Spectrum	Spectrum			
	Mean Hz	Median Hz	Median Hz			
B. mysticetus	698 1723	642 1586 187	634 1591			
C. marginata	83 86 81	82 84	82 86 80			
E. glacialis	34 5 593	317 622 139	226 ³³⁰			
E. australis	$157\frac{235}{39}$	145^{218}_{27}	141_{25}^{219}			
B. acutorostrata	78 110	75 ¹⁰³	73 300			
B. physalus	66_{25}^{129}	66_{25}^{129}	67_{25}^{130}			
P. catodon	4906 11283	4340 11578	2606 10069			
D. leucas	2058 2058 2058	$2279{}^{2279}_{2279}$	$2312{}^{2312}_{2312}$			
S. longirostris	9451_{349}^{26124}	9702_{320}^{26160}	9664 321			
S. $long. + P. cat.$	$16107{}^{22928}_{10992}$	16848 ²⁶⁷⁷²	10874_{10013}^{12547}			
S. bredanensis	11038_{7180}^{16942}	8837 17320 3849	7230 ⁹⁰¹³ 2832			
C. commersonii	3236 3664 2809	3473 3683 3262	3503 3685 3322			
D. delphis	13538_{3285}^{26735}	13571_{1753}^{30097}	$11917{}^{19455}_{2286}$			
G. griseus	6518_{939}^{11305}	6763 12346 869	6408 12397 286			
G. macrorhynchus	4339 9384 2630	3859 8766 1573	3398 8568 198			
G. melaena	5195_{2530}^{7203}	$5198 \frac{8451}{2534}$	5136_{2518}^{8442}			
O. orca	1850_{938}^{3357}	$1480_{\ 622}^{\ 2064}$	1290_{583}^{2062}			
P. crassidens	6558 9636 2091	5623 ⁷⁹⁶⁴	5459_{148}^{7955}			
P. phocoena	$1423_{ 536}^{ 2537}$	1346_{14}^{3017}	625_{9}^{2520}			
I. geoffrensis	3622 4250 2994	$3387 {}^{4072}_{2702}$	$2840{}^{3129}_{2551}$			
A. phillipi	263 393 185	149 161	125 135			
Hammer on metal	$2718{}^{2718}_{2718}$	$1217{}^{1217}_{1217}$	840 840 840			

Table 8 displays statistics relating to amplitude-frequency interactions. Although relatively less robust than many of the previous statistics, these measures also appeared to be useful. For example, the statistics for *C. marginata* had positive Amplitude Frequency Correlation suggesting that the higher frequency sections were louder than low frequency ones, and the negative Amplitude Correlation indicated that sections with downsweeps (or smaller upsweeps) tended to be the loudest.

Note that all of these biological sounds except A. phillipi tended to have positive Amplitude Frequency Correlations (higher frequency sections were louder).

Table 8						
Species	A mplitude	Amplitude				
	Frequency	Upsweep				
	Correlation	Correlation				
B. mysticetus	.27 :40	01 ^{.01}				
C. marginata	.51 ^{.91}	$28{}^{17}_{40}$				
E. glacialis	$.11 \frac{.37}{04}$	$12_{20}^{.00}$				
E. australis	$.24_{15}^{ .92}$	$.04 \stackrel{.21}{06}$				
B. acutorostrata	$.13^{.74}_{40}$	$.01_{05}^{.12}$				
B. physalus	$.65_{22}^{1.00}$	$37^{.08}_{98}$				
P. catodon	$.22_{02}^{ .65}$	01 ^{.43} ₉₈				
D. leucas	.69 .69	$10{}^{10}_{10}$				
S. longirostris	$.34_{02}^{.90}$	$.09_{24}^{1.00}$				
S. long. $+ P.$ cat.	$.51^{.57}_{.42}$	$00^{.11}_{07}$				
S. bredanensis	.28 :47	$.00_{02}^{.06}$				
C. commersonii	.62 .63	$26{}^{22}_{31}$				
D. delphis	.37: ⁷⁷	$02_{37}^{ .13}$				
G. griseus	.44 : ⁷⁸	$.03_{05}^{.15}$				
G. macrorhynchus	.19 :48	$.01 rac{.08}{04}$				
G. melaena	.28 :43	$.01 \stackrel{.07}{05}$				
O. orca	.40 : ⁷²	$09_{63}^{.02}$				
P. crassidens	.40 :68 :03	$00_{02}^{.00}$				
P. phocoena	.32 :95	$02_{-1.00}^{.56}$				
I. geoffrensis	$.24_{-24}^{+25}$	$09{}^{01}_{16}$				
A. phillipi	04 ⁰²	01 ^{.01} ₀₂				
Hammer on metal	.47 :47	63 ⁶³				

4.4 Principal Component Analysis

To obtain a better perspective on the overall distribution of sounds as measured by our statistics, we performed a principal components analysis on the numerical results. The first two principal components provided axes for scatter plots that expressed about half of the total variability in our statistics. As discussed earlier, segregation of the principal

PARADOX structure. The baleen whales and toothed species were plotted separately in Figures 38 and 39. These two groups generally could be separated by frequency alone, and the separate plots focus attention on the variation within these groups. Each sound cut is represented by a colored symbol on the plot; color and symbol type redundantly code species identity.

In both Figure 38 and Figure 39, the data for most species tended to cluster in a relatively discrete portion of the plot. Some regions were shared by a few neighboring species with similar sound types. These results appeared to confirm that acoustic features could be analyzed so as to compute sound statistics that would be useful to classify an unknown biological sound as one of a few potential candidates.

This result was particularly remarkable because the sound data used for the tests included both specific sounds produced by individuals and sounds produced by many animals, with temporally overlapping sounds by two or more animals. If we distinguish between the individual and group recordings, our acoustic classifier will perform much better.

A goal for such analyses has been the development of a system for automatic diagnosis of marine animal sounds based on acoustic criteria. The statistical problem for classification of an unknown relative to known groups is usually addressed by forming estimates of the distance between the unknown and the known, with the most likely classification being the one that minimizes this distance. The distance measure that is usually employed is the Mahalanobis distance (e. g. Morrison 1976, p. 241). Alternatively, we may find that methods which make fewer assumptions (Efron and Tibshirani 1991) will be better suited to developing the classifier. Non-acoustic criteria also could be incorporated from the text databases for additional refinement of these judgements.

5 Discussion and Summary

Software tools have been developed and tested for statistical analysis of marine animal sounds. The preliminary results suggest that such sounds can be classified by means of relatively simple statistical algorithms. Three areas of this research are planned for particular attention.

The SOUNDC database will need to be modified to identify additional information about the sound cuts. These additions will include identification of choruses of overlapping sounds and notes regarding the sequential organization of discrete sound elements. Sequences also needed to be identified relative to the usual pattern of sound production for that species or behavior, and noted if atypical.

Continued evolution of our sound statistics is inevitable. All frequency statistics should be re-scaled, such that the other information is not swamped by gross differences in center frequency, for example. Some correlations may need to be replaced with more robust (perhaps non-parametric) alternatives. Methods for expressing some of these statistics may need to be investigated to increase their independence from each other.

A comprehensive analysis of noise sensitivity and compensation techniques is also important. The specificity of our acoustic measurements is improved by removing the influence of ambient noise, but our ability to classify could be critically impaired if we erroneously discard portions of the signal.

Acknowledgements

This report covers one portion of a larger project addressing many characteristics of marine animal sounds, with all in our laboratory contributing. We particularly thank Mary Ann Daher, Terrance Howald and Peter Tyack for their contributions. We also thank Paul Boutin, Jesse Stanbrough (WHOI), John Coakley and Carla Lundquist (TRICCSMA), who were instrumental in identifying research objectives of Navy interest and framing this project. Funding for the statistical analyses has been through TRICCSMA (NUSC, Newport) from NAVSEA, Contract Number N00140-90-D-1979. The compilation of SOUND and related databases for marine animal sounds is supported by the Office of Naval Research, Ocean Acoustics Program through a series of contracts and grants, including N00014-91-J-1445. Supplemental support for specific enhancements of the marine animal sound data have been provided by NOARL (Stennis Space Center, MI), and ORINCON/DARPA (NOSC, San Diego).

6 LITERATURE CITED

Caldwell, M. C., D. K. Caldwell and P. L. Tyack 1990, The Bottlenose Dolphin: recent progress in research, Academic Press, San Diego.

Efron, B. and R. Tibshirani 1991, Statistical data analysis in the computer age, SCIENCE 253:390-395.

Morrison, D. F. 1976, Multivariate Statistical Analysis, 2nd ed., McGraw-Hill Inc., New York.

Watkins, William A., Kurt Fristrup, and Mary Ann Daher 1991. Marine animal SOUND database. Technical Report WHOI-91-21, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, 51 pp.

Watkins, William A., and Douglas Wartzok 1985. Sensory biophysics of marine mammals.

Marine Mammal Science 1:219-260.




Figure 1. Noise performance of Signal Center (eq. 1). The vertical axis scores Signal Center in seconds; the horizontal axis scores signal/noise. Note the small range of values on the vertical axis. We subjectively label this as low variance, small trend.




Figure 2. Noise performance of Signal Duration (eq. 2). The vertical axis scores Signal Duration in seconds; the horizontal axis scores signal/noise. We subjectively label this as low variance, large trend.




Figure 3. Noise performance of Signal Concentration (eq. 3). The vertical axis scores Signal Concentration in seconds; the horizontal axis scores signal/noise. We subjectively label this as low variance, large trend.




Figure 4. Noise performance of Amplitude Mean (eq. 4). The vertical axis scores Amplitude Mean in arbitrary units; the horizontal axis scores signal/noise. We subjectively label this as low variance, large trend.




Figure 5. Noise performance of Amplitude Standard Deviation (eq. 5). The vertical axis scores Amplitude Standard Deviation, in arbitrary units; the horizontal axis scores signal/noise. We subjectively label this as low variance, large trend.




Figure 6. Noise performance of Attack Fraction (eq. 6). The vertical axis scores Attack Fraction (values of $0 \leftrightarrow 1$ are possible); the horizontal axis scores signal/noise. We subjectively label this as high variance, small trend.




Figure 7. Noise performances of Attack Proportion (eq. 7). The vertical axis scores Attack Proportion (values of $0 \leftrightarrow 1$ are possible); the horizontal axis scores signal/noise. We subjectively label this as high variance, small trend.




Figure 8. Noise performance of Amplitude Skewness (eq. 8). The horizontal axis scores Amplitude Skewness (scale independent); the horizontal axis scores signal/noise. We subjectively label this as high variance, small trend.




Figure 9. Noise performance of Upsweep Mean (eq. 9). The vertical axis scores Upsweep Mean in Hz/s.; the horizontal axis scores signal/noise. We subjectively label this as low variance, small trend.




Figure 10. Noise performance of Upsweep Fraction (eq. 10). The vertical axis scores Upsweep Fraction (values of $0 \leftrightarrow 1$ are possible); the horizontal axis scores signal/noise. We subjectively label this as low variance, small trend.




Figure 11. Noise performance of Upsweep Proportion (eq. 11). The vertical axis scores Upsweep Proportion (values of $0 \leftrightarrow 1$ are possible); the horizontal axis scores signal/noise. We subjectively label this as high variance, small trend.




Figure 12. Noise performance of Time Frequency Correlation (eq. 12). The vertical axis scores Time Frequency Correlation coefficients (values $-1 \leftrightarrow 1$ possible); the horizontal axis scores signal/noise. We subjectively label this as low variance, small trend.




Figure 13. Noise performances of Time Upsweep Correlation (eq. 13). The vertical axis scores Time Upsweep Correlation coefficients (values of $-1 \leftrightarrow 1$ possible); the horizontal axis scores signal/noise. We subjectively score this as low variance, small trend.




Figure 14. Noise performance of Short-term Bandwidth Mean (eq. 14). The vertical axis is Short-term Bandwidth Mean in Hz/s.; the horizontal axis scores signal/noise. We subjectively label this as low variance, large trend.




Figure 15. Noise performance of Short-term Spectral Concentration. The vertical axis scores Short-term Spectral Concentration in Hz/s.; the horizontal axis scores signal/noise. We subjectively label this as low variance, large trend.




Figure 16. Noise performance of Short-term Spectral Asymmetry (eq. 16). The vertical axis scores Short-term Spectral Asymmetry (values of $0 \leftrightarrow 1$ are possible); the horizontal axis scores signal/noise. We subjectively label this as high variance, small trend.




Figure 17:. Noise performance of Total Upper Frequency (see p. 5). The vertical axis scores Total Upper Frequency in Hz; the horizontal axis scores signal/noise. We subjectively latel this as low variance, large trend.




Figure 18. Noise performance of Total Lower Frequency (see p. 5). The vertical axis score Total Lower Frequency in Hz; the horizontal axis scores signal/noise. We subjectively label this as low variance, large trend.




Figure 19. Noise performance of Total Spectrum Concentration (see p. 5). The vertical axis scores Total Spectrum Concentration in Hz; the horizontal axis scores signal/noise. We subjectively label this low variance, large trend.




Figure 20. Noise performance of Modal Upper Frequency (see p. 5). The vertical axis scores Modal Upper Frequency in Hz; the horizontal axis scores signal/noise. We subjectively label this low variance, small trend.




Figure 21. Noise performance of Modal Lower Frequency (see p. 5). The vertical axis scores Modal Lower Frequency in Hz; the horizontal axis scores signal/noise. We subjectively label this low variance, small trend.




Figure 22. Noise performance of Modal Spectrum Concentration (see p. 5). The vertical axis scores Modal Spectrum Concentration in Hz; the horizontal axis scores signal/noise. We subjectively label this low variance, small trend.




Figure 23. Noise performance of Total Spectrum Median (see p. 5). The vertical axis scales Total Spectrum Median in Hz; the horizontal axis scales signal/noise. We subjectively label this low variance, small trend.




Figure 24. Noise performance of Modal Spectrum Median (see p. 5). The vertical axis scales Modal Spectrum Median in Hz; the horizontal axis scales signal/noise. We subjectively label this low variance, small trend.




Figure 25. Noise performance of Median Frequency Mean (eq. 17). The vertical axis scales Median Frequency Mean in Hz; the horizontal axis scales signal/noise. We subjectively label this as low variance, small trend.




Figure 26. Noise performance of Amplitude Frequency Correlation (eq. 18). The vertical axis scales Amplitude Frequency Correlation coefficients (values of $-1 \leftrightarrow 1$ are possible); the horizontal axis scales signal/noise. We subjectively label this as high variance, large trend.




Figure 27. Noise performance of Amplitude Upsweep Correlation (eq. 19). The vertical axis scales Amplitude Upsweep Correlation coefficients (values of $-1 \leftrightarrow 1$ are possible); the horizontal axis scales signal/noise. We subjectively label this as high variance, large trend.




Figure 28. Linear prediction of Amplitude Standard Deviation based on 25 other variables (Signal Center not used for any regressions). The axes are in arbitrary units. The percent of variance explained by the linear regression is displayed in the title.




Figure 29. Linear prediction of Median Frequency Mean based on 24 other variables (Amplitude Standard Deviation already removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 30. Linear prediction of Modal Spectrum Median based on 23 other variables (two variables previously removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 31. Linear prediction of Total Upper Frequency based on 22 other variables (three variables previously removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 32. Linear prediction of Total Spectrum Concentration based on 21 other variables (four variables previously removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 33. Linear prediction of Modal Lower Frequency based on 20 other variables (five variables previously removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 34. Linear prediction of Modal Upper Frequency based on 19 other variables (six variables previously removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 35. Linear prediction of Short-term Spectral Concentration based on 18 other variables (seven variables previously removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 36. Linear prediction of Total Spectrum Median based on 17 other variables (eight variables previously removed). The axes are in Hz, but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.




Figure 37. Linear prediction of Amplitude Upsweep Correlation based on 16 other variables (nine variables previously removed). The axes are correlation coefficients (values $-1 \leftrightarrow 1$ possible), but the data are shifted such that the means are zero. The percent of variance explained by the linear regression is displayed in the title.

Figure 38. Plot of sound samples from six baleen whale species. The horizontal axis scales each sample's score on the first principal component (which basically reflects center frequency). The vertical axis scales each sample's score on the second principal component (not easily interpreted).





Figure 39. Plot of sound samples from thirteen toothed whale species. The horizontal axis scales each sample's score on the first principal component (which basically reflects center frequency). The vertical axis scales each sample's score on the second principal component (not easily interpreted).



DOCUMENT LIBRARY

March 11, 1991

Distribution List for Technical Report Exchange

Attn: Stella Sanchez-Wade Documents Section Scripps Institution of Oceanography Library, Mail Code C-075C La Jolla, CA 92093

Hancock Library of Biology & Oceanography
Alan Hancock Laboratory
University of Southern California
University Park
Los Angeles, CA 90089-0371

Gifts & Exchanges Library Bedford Institute of Oceanography P.O. Box 1006 Dartmouth, NS, B2Y 4A2, CANADA

Office of the International Ice Patrol c/o Coast Guard R & D Center Avery Point Groton, CT 06340

NOAA/EDIS Miami Library Center 4301 Rickenbacker Causeway Miami, FL 33149

Library Skidaway Institute of Oceanography P.O. Box 13687 Savannah, GA 31416

Institute of Geophysics University of Hawaii Library Room 252 2525 Correa Road Honolulu, HI 96822

Marine Resources Information Center Building E38-320 MIT Cambridge, MA 02139

Library
Lamont-Doherty Geological
Observatory
Columbia University
Palisades, NY 10964

Library Serials Department Oregon State University Corvallis, OR 97331 Pell Marine Science Library University of Rhode Island Narragansett Bay Campus Narragansett, RI 02882

Working Collection Texas A&M University Dept. of Oceanography College Station, TX 77843

Library
Virginia Institute of Marine Science
Gloucester Point, VA 23062

Fisheries-Oceanography Library 151 Oceanography Teaching Bldg. University of Washington Seattle, WA 98195

Library R.S.M.A.S. University of Miami 4600 Rickenbacker Causeway Miami, FL 33149

Maury Oceanographic Library Naval Oceanographic Office Stennis Space Center NSTL, MS 39522-5001

Marine Sciences Collection Mayaguez Campus Library University of Puerto Rico Mayaguez, Puerto Rico 00708

Library
Institute of Oceanographic Sciences
Deacon Laboratory
Wormley, Godalming
Surrey GU8 5UB
UNITED KINGDOM

The Librarian CSIRO Marine Laboratories G.P.O. Box 1538 Hobart, Tasmania AUSTRALIA 7001

Library
Proudman Oceanographic Laboratory
Bidston Observatory
Birkenhead
Merseyside L43 7 RA
UNITED KINGDOM

0272-101				
REPORT DOCUMENTATION PAGE	1. REPORT NO. WHOI-92-04	2.	·	's Accession No.
I. Title and Subtitle			5. Report D	
Characterizing Acoust	ic Features of Marine Anim	nal Sounds		uary 1992
			6.	
7. Author(s) Kurt M. Fristrup and William A. Watkins			1	ng Organization Rept. No.
				IOI-92-04
9. Performing Organization Name and	Address		10. Project/	Task/Work Unit No.
Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543			11. Contrac	t(C) or Grant(G) No.
			(C) N(00	140-90-D-1979
, , , , , , , , , , , , , , , , , , , ,			3,700	014-91-J-1445
			(G) NUU	01-713-14-3
i2. Sponsoring Organization Name an	xd Address		13. Type of	Report & Period Covered
NAVSEA, Office of Naval Research, NOARL and ORINCON/DARPA			Tec	hnical Report
			14.	***************************************
This report should be cited a	as: Woods Hole Oceanog. Inst. Tech.	Rept., WHOI-92-04.		
enhanced by adjusting calcul Bandwidth, Intensity, Durati Amplitude Frequency Interactested on a subset of 200 seq species, and one seal. The staprincipal components analys	at summarize particular aspects of soundations to compensate for ambient noise on, Amplitude Modulation, Frequency oction. The efficacy of noise compensate quences of marine animal sounds, inclusive were reviewed for each species sis. Preliminary results confirm that sue are encouraged to continue toward a	se. The sound measures Modulation, Short-tention was tested for each uding sequences from 2 and a graphical compatch sounds can be class	included statistic in Bandwidth, Ce statistic. Then, t 0 species: six ba- rison of all speci- ified by means o	cs for Aggregate enter Frequency, and he sound measures were leen whales, 13 toothed es was generated using f relatively simple
17. Document Analysis a. Descriptor sound database marine animal sounds automatic characterization of b. Identifiers/Open-Ended Terms				
c. COSATI Field/Group		19. Security Class (This Report)	21. No. of Pages
Approved for public rel	lease; distribution unlimited.	UNCLASS 20 Security Class (IFIED	82 22 Price

20. Security Class (This Page)

22. Price