

2

AD-A248 026

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE



Form Approved
OMB No. 0704-0188

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION Unclassified		1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release Distribution unlimited	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE		4 PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 7	
4 PERFORMING ORGANIZATION REPORT NUMBER(S)		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Brigham Young University	6b. OFFICE SYMBOL (If applicable) BYU	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Provo, UT 84602		7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, VA 22217-5000	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable) ONR	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-91-J-1710	
8c. ADDRESS (City, State, and ZIP Code) 800 No. Quincy Street Arlington, VA 22217-5000		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO	PROJECT NO
		TASK NO	WORK UNIT ACCESSION NO.

11. TITLE (Include Security Classification) Recognition of the Enantiomers of Chiral Organic Ammonium Salts By Chiral Dibenzyl- and Diphenyl-Substituted Diamido-or Dithionoamidopyridino-18-crown-6 Ligands

12. PERSONAL AUTHOR(S) P. Huszthy, J.S. Bradshaw, C.Y. Zhu, T. Wang, and R.M. Izatt

13a. TYPE OF REPORT Interim	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1992, March 24	15. PAGE COUNT
--------------------------------	--	---	----------------

16 SUPPLEMENTARY NOTATION

17 COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)
FIELD	GROUP	SUB-GROUP	

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

New chiral pyridino-18-crown-6 ligands containing two amide, thionoamide or amine units and two benzyl or phenyl substituents on chiral macroring positions have been synthesized (see Figure 1). Some of these chiral ligands exhibit only moderate or no recognition for the enantiomers of [α -(1-naphthyl)ethyl]ammonium perchlorate as determined by ¹H NMR spectral techniques.

CONTINUED ON BACK


20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. Harold Guard		22b TELEPHONE (Include Area Code) (202) 696-4409	22c OFFICE SYMBOL ONR

92-07983



92 3 20 930

Figure 1. Chiral Diamido-, Dithionoamido-, Diaza- and Azapyridino-18-crown-6 Ligands




- | | |
|---|-----------------------------------|
| 1, X = NH; Y = O; R = benzyl (S,S) | |
| 2, X = NH; Y = S; R = benzyl (S,S) | |
| 3, X = NH; Y = H ₂ ; R = benzyl (S,S) | |
| 4, X = NH; Y = O; R = phenyl (S,S) | |
| 5, X = NH; Y = S; R = phenyl (S,S) | |
| 6, X = NH; Y = H ₂ ; R = phenyl (S,S) | |
| 7, X = NCH ₃ ; Y = O; R = phenyl (S,S) | |
| 8, X = NCH ₃ ; Y = S; R = phenyl (S,S) | |
| 9, X = NCH ₃ ; Y = H ₂ ; R = phenyl (S,S) | |
| | 10, R = H (S,S) |
| | 11, R = C(O)CH ₃ (S,S) |

Complexation of the enantiomeric forms of NapEt by some of the new ligands has been studied by the temperature-dependent ¹H NMR technique¹⁻⁵, and by determining the log K values for the association of the chiral ligand and the enantiomer salt by a direct ¹H NMR spectral method.^{5,9} Table I contains the free energy of activation (ΔG_c^\ddagger) and log K values for the interaction of some of these new chiral ligands with the enantiomers of [α -(1-naphthyl)ethyl]ammonium perchlorate(NapEt). The data show that, in general, these chiral ligands exhibit only moderate or no recognition for the enantiomers of NapEt. The ΔG_c^\ddagger data show that ligands 7, 8, and 9 form stable

Accession For	
Yield (Gram)	<input checked="" type="checkbox"/>
Yield (%)	<input type="checkbox"/>
Unpurified	<input type="checkbox"/>
Purification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Special	

A-1



complexes in CD_2Cl_2 with the (R) and (S) forms of NapEt but the resulting complexes each have about the same kinetic stability. This is in sharp contrast to complexation by the chiral diphenyl-substituted pyridino-18-crown-6 ($Y = \text{H}_2$, $X = \text{O}$, $R = \text{phenyl}$) and diesterpyridino-18-crown-6 ($Y = \text{O}$, $X = \text{O}$, $R = \text{phenyl}$) ligands (see Figure 1) which exhibited considerable recognition of one form of NapEt over the other form in CD_2Cl_2 as determined by the temperature dependent ^1H NMR method.⁵ Log K values in Table I do show that some of these ligands exhibit enantiomeric recognition in a mixture of 50% CDCl_3 - 50% CD_3OD or in CD_3NO_2 . Dithionoamide (S,S)-5 favored (R)-NapEt over (S)-NapEt by 0.37 log K units. This recognition compares favorably with that obtained by the similar diphenyl-substituted crown ($X = \text{O}$, $Y = \text{O}$) ligand.⁵ Ligand (S,S)-8 also exhibited moderate recognition for (R)-NapEt in the 50% CDCl_3 - 50% CD_3OD mixture. Ligands (S,S)-1 and (S,S)-7 exhibit some recognition for one form of NapEt in CD_3NO_2 . Thus, in some cases, these new chiral ligands show moderate enantiomeric recognition but they have inferior recognition properties as compared to the diphenyl- and di-t-butyl-substituted pyridino- and diesterpyridino-18-crown-6 ligands (all have $X = \text{O}$).⁵

The data in Table I also show that log K values for macrocycle-ammonium salt interactions increase as the polarity of the solvent decreases. This is an expected trend for solvent effects in these types of interactions. It is also instructive to note that there is a significant decrease in log K values when substituting an amide group for an ester in these systems. This is best observed by comparing the log K value of 0.7 (Table 1) for the interaction of (S,S)-4 with (R)-

NapEt in a mixture of 50% CDCl₃ - 50% CD₃OD with a log K value of 2.15 for the same interaction of the comparable ester (X and Y = O, R = phenyl) in a mixture of 30% CDCl₃ - 70% CD₃OD.⁵ The two solvent systems are slightly different but the latter solvent is more polar which should decrease the log K value. This significant decrease in complexing ability for the diamido-crowns is probably a result of distorted macrorings that will be mentioned later. The diamino-crowns, on the other hand, form stronger complexes as shown by the log K value of 3.2 for the interaction of (S,S)-9 with (R)-NapEt.

Table I. Free Energies of Activation (ΔG_c^\ddagger)^a and log K Values^b for the Interaction of Some Chiral Macrocyclic Ligands with the R and S Forms of [α -(1-Naphthyl)ethyl]ammonium Perchlorate (NapEt)

Ligand	ΔG_c^\ddagger values (Kcal/mol) ^c		log K values at 20 °C		
	(<u>R</u>)-NapEt	(<u>S</u>)-NapEt	(<u>R</u>)-NapEt ^d	(<u>S</u>)-NapEt ^d	(<u>S</u>)-NapEt ^e
(<u>S</u> , <u>S</u>)-1			g	g	0.8
(<u>S</u> , <u>S</u>)-4	f	f	0.7	g	
(<u>S</u> , <u>S</u>)-5			1.39	1.02	
(<u>S</u> , <u>S</u>)-7	11.3	11.2	g	g	2.75
(<u>S</u> , <u>S</u>)-8	>14.0	>13.9	0.7	0.5	
(<u>S</u> , <u>S</u>)-9	12.0	12.1	3.2	3.3	h

^a A Varian VXR-500 spectrometer was used to record all ¹H NMR Spectra for determination of both ΔG_c^\ddagger and log K values. The procedure for the determination of ΔG_c^\ddagger values was as described in ref. 1-5.

^b Log K values were determined by a direct ¹H NMR procedure as described in ref. 9.

^c CD₂Cl₂ was used as a solvent.

^d 50%/50% (v/v) CDCl₃-CD₃OD mixture was used as a solvent.

^e CD₃NO₂ was used as a solvent.

^f No splitting was observed in the ¹H NMR spectra.

^g Log K value was so low that accurate measurements could not be made.

^h Log K value was so high that accurate calculations could not be made.

References

- (1) Davidson, R. B.; Bradshaw, J. S.; Jones, B. A.; Dalley, N. K.; Christensen, J. J.; Izatt, R. M.; Morin, F. G.; Grant, D. M. J. Org. Chem. 1984, 49, 353.
- (2) Bradshaw, J. S.; Thompson, P. K.; Izatt, R. M.; Morin, F. G.; Grant, D. M. J. Heterocyclic Chem. 1984, 21, 897.
- (3) Bradshaw, J. S.; Colter, M. L.; Nakatsuji, Y.; Spencer, N. O.; Brown, M. F.; Izatt, R. M.; Arena, G.; Tse, P.-K.; Wilson, B. E.; Lamb, J. D.; Dalley, N. K.; Morin, F. G.; Grant, D. M. J. Org. Chem. 1985, 50, 4865.
- (4) Bradshaw, J. S.; Huszthy, P.; McDaniel, C. W.; Zhu, C. Y.; Dalley, N. K.; Izatt, R. M.; Lifson, S. J. Org. Chem. 1990, 55, 3129.
- (5) Huszthy, P.; Bradshaw, J. S.; Zhu, C. Y.; Izatt, R. M.; Lifson, S. J. Org. Chem. 1991, 56, 3330.
- (6) Kyba, E. B.; Koga, K.; Sousa, L. R.; Siegel, M. G.; Cram, D. S. J. Am. Chem. Soc. 1973, 95, 2692.
- (7) Li, Y.; Echegoyen, L.; Martinez-Diaz, M. V.; Mendoza, J.; Torres, T. J. Org. Chem. 1991, 56, 4193.
- (8) Izatt, R. M.; Zhu, C.-Y.; Huszthy, P.; Bradshaw, J. S.
"Enantiomeric Recognition in Macrocycle-Primary Ammonium Cation Systems," in "Crown Ethers: Toward Future Applications," Cooper, S. R. Ed., VCH Press, New York, in press.
- (9) Zhu, C. Y.; Bradshaw, J. S.; Oscarson, S. L.; Izatt, R. M. J. Incl. Phenom. in press.