

AD-A248 359

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188



Estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including instructions, to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

2 REPORT DATE March 1992		3 REPORT TYPE AND DATES COVERED professional paper	
4 TITLE AND SUBTITLE NARROW-BAND SONAR SIGNALS OF SMALL CETACEANS		5 FUNDING NUMBERS PR: MMB2 PE: 0602435N WU: DN688674	
6 AUTHOR(S) W. W. L. Au		8 PERFORMING ORGANIZATION REPORT NUMBER	
7 PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Command, Control, and Ocean Surveillance Center (NCCOSC), Research, Development, Test and Evaluation Division (NRaD) San Diego, CA 92152-5000		10 SPONSORING/MONITORING AGENCY REPORT NUMBER	
9 SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Ocean Systems Center (now NRaD) Block Programs San Diego, CA 92152-5000		11 SUPPLEMENTARY NOTES	
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE	
13 ABSTRACT (Maximum 200 words) The sonar signals of some of the smallest cetaceans are very similar to each other and very different than that of mid-sized cetaceans. Signals of <i>Cephalorhynchus hectori</i> , <i>Cephalorhynchus commersonii</i> (genus <i>Cephalorhynchus</i>), <i>Phocoena phocoena</i> , <i>Neophocoena phocoena</i> and <i>Phocoenoides dalli</i> (family phocoenidae) are compared with those of some larger dolphins. Signals of <i>Tursiops truncatus</i> , <i>Delphinapterus leucas</i> and <i>Pseudorca crassidens</i> , housed in Kaneohe Bay, Oahu, Hawaii, have peak frequencies between 100-200 kHz, with high amplitudes (210-225 dB re 1 μ Pa), short durations (50-70 μ s), and wide bandwidths (30-40 kHz). Some of the smaller cetaceans emit signals having peak frequencies between 120 and 140 kHz, with low amplitudes (< 170 dB re 1 μ Pa), long durations (170-430 μ s) and narrow bandwidths (7-11 kHz). Double pulses are also emitted regularly by some of the smaller dolphins and very infrequently by the larger dolphins. Signals used by the smaller animals may reflect constraints associated with their small size and differences in generation mechanisms. For a given peak acoustic pressure, there is 3 to 4 times more energy in the signals of the smaller cetaceans. However, because of the narrower bandwidths, the distance resolution capability of the small cetacean signals is between 2 to 3.5 times inferior to that of the larger animals. Furthermore, the narrow bandwidth signals do not possess any doppler resolution properties. Published in <i>Proceedings of 9th Biennial Conference on the Biology of Marine Mammals</i> , December, 1991.			
14 SUBJECT TERMS marine biosystems marine biology		15 NUMBER OF PAGES	
17 SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		16 PRICE CODE	
18 SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19 SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	
		20 LIMITATION OF ABSTRACT SAME AS REPORT	

DTIC
APR 7 1992

1



UNCLASSIFIED

21a. NAME OF RESPONSIBLE INDIVIDUAL W. W. L. Au	21b. TELEPHONE (include Area Code) (808) 257-1647	21c. OFFICE SYMBOL Code 512

Accession For
NTIS Grant
DTIC TAB
Unannounced
Justification

Published in Program of Ninth Biennial Conference on the Biology of Marine Mammals, Chicago, Illinois, Dec. 5-9, 1991.

NARROW-BAND SONAR SIGNALS OF SMALL CETACEANS

Whitlow W. L. Au
Naval Ocean Systems Center
P.O. Box 997, Kailua, Hi 96734

Distribution
Availability Code
Dist Sp. 101
A-1 20

ABSTRACT

The sonar signals of some of the smallest cetaceans are very similar to each other and very different than that of mid-sized cetaceans. Signals of *Cephalorhynchus hectori*, *Cephalorhynchus commersonii*, (genus *Cephalorhynchus*), *Phocoena phocoena*, *Neophocoena phocoena* and *Phocoenoides dalli* (family phocoenidae) are compared with those of some larger dolphins. Signals of *Tursiops truncatus*, *Delphinapterus leucas* and *Pseudorca crassidens*, housed in Kaneohe Bay, Oahu, Hawaii, have peak frequencies between 100-120 kHz, with high amplitudes (210-225 dB re 1 μ Pa), short durations (50-70 μ s), and wide bandwidths (30-40 kHz). Some of the smaller cetaceans emit signals having peak frequencies between 120 and 140 kHz, with low amplitudes (< 170 dB re 1 μ Pa), long durations (170-430 μ s) and narrow bandwidths (7-11 kHz). Double pulses are also emitted regularly by some of the smaller dolphins and very infrequently by the larger dolphins. Signals used by the smaller animals may reflect constraints associated with their small size and differences in generation mechanisms. For a given peak acoustic pressure, there is 3 to 4 times more energy in the signals of the smaller cetaceans. However, because of the narrower bandwidths, the distance resolution capability of the small cetacean signals is between 2 to 3.5 times inferior to that of the larger animals. Furthermore, the narrow bandwidth signals do not possess any doppler resolution properties.