

AD-A250 055

DOCUMENTATION PAGE

2

1a. REPC
Uncl

1b. RESTRICTIVE MARKINGS
None

2a. SECL

3. DISTRIBUTION/AVAILABILITY OF REPORT
OPEN

2b. DECLASSIFICATION/DOWNGRADING SCHEDULE
SELECTE
MAY 14 1992

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

5. MONITORING ORGANIZATION REPORT NUMBER(S)
AFOSR-TR- 92 0310

6a. NAME OF PERFORMING ORGANIZATION
University of Rochester

6b. OFFICE SYMBOL (if applicable)

7a. NAME OF MONITORING ORGANIZATION
AFOSR/NL

6c. ADDRESS (City, State and ZIP Code)
River Station
Rochester, New York 14642

7b. ADDRESS (City, State and ZIP Code)
Building 410
Bolling AFB, DC 20332-6448

8a. NAME OF FUNDING/SPONSORING ORGANIZATION
US Air Force

8b. OFFICE SYMBOL (if applicable)
AFOSR, NL

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER
AFOSR-89-C041

10. SOURCE OF FUNDING NOS.

PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.
60102F	2313	A8	

11. TITLE (Include Security Classification)
Function of P and M Pathways in Primates

12. PERSONAL AUTHOR(S)
William Merigan

13a. TYPE OF REPORT
Annual Technical Rept.

13b. TIME COVERED
FROM 10/1/89 TO 9/30/90

14. DATE OF REPORT (Yr., Mo., Day)
13 April 1992

15. PAGE COUNT
3

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD	GROUP	SUB-GR

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

ABSTRACT (Continue on reverse if necessary and identify by block number)

Psychophysical threshold measures were used in combination with precisely located lesions of the sub-cortical visual pathway to examine segregation of function between P and M pathways. Sceleral search coils monitored fixation locus in the tested monkeys, to insure that test stimuli were presented in visual field regions corresponding to the lesion location. Results of these studies indicated that the P pathway (or color-opponent pathway) is the major contributor to visual acuity, color vision, and luminance contrast sensitivity. On the other hand, the M pathway (or broad contours), as well as sensitivity to rapidly shifting visual stimuli.

This document has been approved for public release and sale; its distribution is unlimited.

19. DISTRIBUTION/AVAILABILITY OF ABSTRACT
CLASSIFIED/UNLIMITED SAME AS RPT DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION
Unclassified

NAME OF RESPONSIBLE INDIVIDUAL
Dr. John F. Taniguchi

22b. TELEPHONE NUMBER (Include Area Code)
301-761-5031

22c. OFFICE SYMBOL
AFOSR/NL

1. Abstract

Psychophysical threshold measures were used in combination with precisely located lesions of the sub-cortical visual pathway to examine segregation of function between P and M pathways. Scleral search coils monitored fixation locus in the tested monkeys, to insure that test stimuli were presented in visual field regions corresponding to the lesion location. Results of these studies indicated that the P pathway (or color-opponent pathway) is the major contributor to visual acuity, color vision, and luminance contrast sensitivity. On the other hand, the M pathway (or broadband pathway) appeared crucial for the visibility of low spatial frequencies (broad contours), as well as sensitivity to rapidly drifting visual stimuli.

2. Research objectives

The goal of this project was to determine if there is a segregation of function between the sub-cortical P and M visual pathways of the macaque. Lesions were made by the injection of ibotenic acid (a glutamate agonist) into carefully selected regions of the parvocellular and magnocellular layers of the lateral geniculate nucleus. Visual function was assessed by the measurement of psychophysical thresholds in portions of the visual field corresponding to the lesions.

3. Status of the research

As part of our assessment of the pre-lesion vision of two monkeys, we determined spatial resolution across the horizontal meridian of the visual field. As was expected from preliminary results in humans, acuity declined monotonically with distance from the fovea. The rate of this decline, and the absolute acuity at each eccentric location were in close agreement with the resolution limits imposed by sampling density of retinal ganglion cells of the P pathway. Later results (below) on the effects of lesions of the P pathway on visual acuity confirmed that the P pathway limits visual acuity.

In initial studies, lesions were made in the parvocellular layers of the lateral geniculate of two monkeys. Psychophysical testing was followed by physiological and anatomical reconstruction of the lesion. Fortunately, both lesions fully destroyed the parvocellular and spared the magnocellular layers in the region corresponding to the tested locus in the visual field. Results of these studies showed clearly that the P pathway plays a dominant role in spatial resolution and chromatic contrast sensitivity as well as luminance contrast sensitivity over a broad range of spatial and temporal frequencies.

Determining the effect of M pathway lesions on visual function has proven to be somewhat more complex. In our first studies, we found that temporal resolution was not reduced by lesions of the M pathway. This result was somewhat counter-intuitive, since visual system physiologists have generally considered the M pathway to be the basis of temporal resolution. Effects on contrast sensitivity were consistent with the findings for temporal resolution. Low spatial and higher temporal frequency vision was devastated by M lesions. However, at moderate or higher spatial frequencies, lesions of the M pathway had no measurable effect on contrast sensitivity.

92-12621



92 5 11 093

4. Publications

Merigan, W.H. and Katz, L.M. (1990) Spatial resolution across the macaque retina. *Vision Res.*, 7, 985-991.

Merigan W.H., Maunsell, J.H.R. (1990) Macaque vision after magnocellular lateral geniculate lesions. *Vis. Neurosci.*, 5, 347-352.

5. Participating personnel

William H. Merigan Associate Professor Ophthalmology Univ. Roch. Med. Ctr.

John H. R. Maunsell Associate Professor Physiology Univ. Roch. Med. Ctr.

Thomas A. Eskin Associate Professor Pathology Univ. Roch. Med. Ctr.

6. Oral presentations and conferences

Merigan, W.H., Katz, L. M., and Maunsell, J. H. R. (1989) Contribution of the primate parvocellular pathway to acuity and contrast sensitivity. *Invest. Ophthamol. Vis. Sci.* 30,

7. Discoveries and inventions

None

8. Significance of accomplishments

Our earlier studies, proposed in the original application, and supported by AFOSR to completion, showed that subcortical visual pathways in the macaque, and possibly in the human, are quite independent, and appear specialized primarily for the transmission of different portions of the spatio-temporal visual spectrum. In addition, the P pathway appears to be the sole conduit for color vision. Future work will examine specific suggestions that the M pathway may be the substrate for motion perception.

<input checked="" type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>

Availability Codes	
Dist	Avail and/or Special
A-1	

