

AD-A250 745

DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

This is estimated to average 1 hour per response including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collection of information, sending comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Boulevard, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

2. REPORT DATE 1992		3. REPORT TYPE AND DATES COVERED Open literature publication	
4. TITLE AND SUBTITLE (U)Evidence of NK1 and NK2 tachykinin receptors and their involvement in histamine release in a murine mast cell line		5. FUNDING NUMBERS 61102A 3M161102BS11 AA	
6. AUTHOR(S) Krumins, SA and Broomfield, CA		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: SGRD-UV- PB Aberdeen Proving Ground, MD 21010-5425		10. SPONSORING/MONITORING AGENCY REPORT NUMBER USAMRICD-P91-017	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) USAMRICD ATTN: SGRD-UV-RC APG, MD 21010-5425		11. SUPPLEMENTARY NOTES Appeared in Neuropeptides, vol 21, 65-72, 1992	
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Abstract—Binding of [³ H]substance P (SP) and histamine release were examined using a cloned mouse mast cell line. SP binding was saturable and specific. In the presence of 30mM Na ₂ SO ₄ /50mM Tris buffer, SP interacted with two types of binding sites with K _d values of 0.3 and 40nM. High-affinity SP binding was blocked by the inclusion of 0.5uM of the NK1 receptor selective ligand septide in the binding mixture. Neurokinin A (NKA) evoked concentration dependent histamine release. At concentrations in the nanomolar range, the NK1 preferring agonists SP, SP methylester and physalaemin evoked $\leq 5\%$ net release of histamine, which was substantially less than the maximum effect of NKA (+37%) in the micromolar range. Pretreatment of the cells with the NK2 antagonist peptide A reduced NKA-induced histamine release. [D-Arg ¹ ,D-Phe ² ,D-Trp ³ ,Leu ¹¹]-substance P, a putative SP antagonist, also elicited histamine release in the micromolar range, apparently acting as an agonist at the NK2 site. Compound 48/80, N-terminal SP fragments, neurokinin B and the two selective NK2 receptor antagonists cyclo[Gln-Trp Phe (R)-[ANC-2]Leu-Met] (peptide A) and cyclo[Gln-Trp-Phe-Gly Leu Met] (peptide B) were ineffective. Although the results suggest the coexistence of functional NK1 and NK2 receptors, it appears that in this mast cell line neurokinin-induced histamine release is primarily mediated by the NK2 receptor, characterized biochemically as a low affinity binding site with a K _d value of 40nM for SP.			
14. SUBJECT TERMS histamine release, substance P, tachykinin receptors		15. NUMBER OF PAGES 8	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT None

Evidence of NK1 and NK2 Tachykinin Receptors and their Involvement in Histamine Release in a Murine Mast Cell Line

S. A. KRUMINS and C. A. BROOMFIELD

Biochemical Pharmacology Branch, United States Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, Maryland 21010, USA. (Reprint request to CAB)

Abstract—Binding of [3 H]substance P (SP) and histamine release were examined using a cloned mouse mast cell line. SP binding was saturable and specific. In the presence of 30 mM Na_2SO_4 /50 mM Tris buffer, SP interacted with two types of binding sites with K_d values of 0.3 and 40 nM. High-affinity SP binding was blocked by the inclusion of 0.5 μM of the NK1 receptor selective ligand septide in the binding mixture. Neurokinin A (NKA) evoked concentration-dependent histamine release. At concentrations in the nanomolar range, the NK1 preferring agonists SP, SP methylester and physalaemin evoked $\leq 5\%$ net release of histamine, which was substantially less than the maximum effect of NKA (+37%) in the micromolar range. Pretreatment of the cells with the NK2 antagonist peptide A reduced NKA-induced histamine release. [D-Arg¹, D-Phe⁵, D-Trp^{7,9}, Leu¹¹]-substance P, a putative SP antagonist, also elicited histamine release in the micromolar range, apparently acting as an agonist at the NK2 site. Compound 48/80, N-terminal SP fragments, neurokinin B and the two selective NK2 receptor antagonists cyclo[Gln-Trp-Phe-(R)-[A¹C-2]Leu-Met] (peptide A) and cyclo[Gln-Trp-Phe-Gly-Leu-Met] (peptide B) were ineffective. Although the results suggest the coexistence of functional NK1 and NK2 receptors, it appears that in this mast cell line neurokinin-induced histamine release is primarily mediated by the NK2 receptor, characterized biochemically as a low affinity binding site with a K_d value of 40 nM for SP.

Introduction

The undecapeptide substance P (SP), considered to be a neurotransmitter involved in sensory processes, is now thought to cause a wide range of peripheral effects, e.g., vasodilation, plasma extravasation and immunomodulation (1). SP

belongs to the peptide family known as tachykinins, characterized by a common C-terminal sequence, Phe-X-Gly-Leu-Met-NH₂ (Table 1). In addition to SP, two other mammalian tachykinins, named neurokinins, are known, the decapeptides neurokinin A (NKA) and neurokinin B (NKB) (2). The biological effects of SP, NKA and NKB are mediated through their respective tachykinin receptors, NK1, NK2 and NK3 (2-4). Among the

Date received 28 May 1991
Date accepted 3 October 1991

Table 1 Amino acid sequences of the mammalian tachykinins (neurokinins).

Substance P (SP)	Arg-Pro-Lys-Pro-Gln-Gln-Phe-Phe-Gly-Leu-Met-NH ₂
Neurokinin A (NKA)	His-Lys-Thr-Asp-Ser-Phe-Val-Gly-Leu-Met-NH ₂
Neurokinin B (NKB)	Asp-Met-His-Asp-Phe-Val-Gly-Leu-Met-NH ₂

three neurokinins, only SP is known to induce histamine release from some types of mast cells, such as rat peritoneal (5, 6) and human skin mast cells (7, 8). SP action on mast cells is not fully understood. Whereas most of the biological effects of neurokinins depend on the interaction of the common C-terminal sequence (4), the histamine releasing activity of SP, in contrast, requires the basic N-terminal domain of the SP molecule (9, 10); i.e., Arg-Pro-Lys-Pro. However, interactions of the C-terminal portion are needed for maximum histamine release (11). Because receptors on mast cells recognizing the N-terminal domain of the SP molecule have not yet been identified (12), mediation of SP action via such receptors is presently being questioned. Other mechanisms for SP action on mast cells have been suggested. For example, a receptor-independent pathway involving GTP-binding regulatory proteins (G proteins) has been suggested as a common mechanism for the histamine releasing action of SP and the poly-basic molecules mastoparan and compound 48/80 (12-14). Mastoparan, a peptide toxin from wasp venom, and compound 48/80 are potent histamine releasers in rat peritoneal mast cells (11, 13) but not in guinea pig mast cells (15), human pulmonary mast cells (16) or human basophilic leukocytes (17, 18). SP displays similar variation in evoking histamine release from mast cells of different tissues and species (5-11, 13, 19). These variations in response remain inexplicable.

In this report, the action of neurokinins on a cloned mouse mast cell line, MC/9, derived from fetal liver cells from a (B6 X A/J)F₁ mouse (20), is described. The aim of this study was to examine biochemically SP binding, using a homogeneous cell population, and correlate the binding pattern with selective neurokinin-induced histamine release to 1) identify the tachykinin receptors present on mast cells and 2) assess their role in neurokinin-induced histamine release. Biochemical studies of SP binding to mast cells are lacking. Most studies

of SP-induced histamine release have been performed on heterogeneous but enriched populations of rat peritoneal mast cells or blood basophils, not well suited for biochemical assays. Autoradiographic studies of SP binding have failed to show specific binding of SP to rat peritoneal mast cells (21).

In pilot experiments, NKA was found to release histamine from MC/9 cells, while SP, N-terminal SP fragments, compound 48/80 and the mast cell degranulating peptide (MCD), another potent histamine releasing compound isolated from wasp venom, caused little or no histamine release. Moreover, specific binding of SP to MC/9 cells was observed. These experiments were expanded and are described in detail here. A part of this investigation has previously been presented in an abstract (22).

Materials and Methods

Materials

Radioligand used was [2-prolyl-3,4-³H]substance P ([³H]SP; 36.3 and 42.6 Ci/mmol, New England Nuclear, MA). Unlabeled peptides were purchased from Peninsula Laboratories, CA, or Cambridge Research Biochemicals, NY, and mast cell degranulating peptide (MCD) was from Bachem, CA. Bovine albumin, Fraction V, bacitracin, bestatin, compound 48/80 and histamine diphosphate salt were purchased from Sigma Chemical Company, MO. Fluoraldehyde *o*-Phthalaldehyde reagent solution was purchased from Pierce Chemical Company, IL. The MC/9 mast cell line was purchased from American Type Culture Collection (ATCC, CRL 8306), MD.

Methods

The MC/9 cells were incubated as suspension cultures in Dulbecco's modified Eagle's medium

(DME) containing L-arginine HCl (116 mg/l), L-asparagine (36 mg/l), folic acid (6 mg/l), non-essential amino acids (0.1 mM), sodium pyruvate (1.0 mM) and fetal bovine serum (10%), supplied by ATCC. Actively growing cultures, seeded at 10^5 cells/ml in Corning 25 cm² polystyrene tissue culture flasks and supplemented with fresh culture medium every 2-3 days, were allowed to multiply at 37°C in a 7% CO₂-air mixture for 5 to 6 days. The cultures were harvested at a cell density of 1.5×10^6 cells/ml. Aliquots of the cultured cells were prepared for histamine release studies, while remaining cells were pelleted and stored at -70°C or in liquid nitrogen. Cell viability was assessed by the Trypan blue exclusion test. Cultured cells of viability $\geq 75\%$ were used in the experiments.

Binding studies. A modification of the method described by Maruyama (23) was used. The frozen cell pellets were thawed and diluted with 50 volumes of ice-cold 50 mM Tris-HCl buffer (pH 7.4), vortexed vigorously, then centrifuged at 40000g for 15 min at 4°C and washed one more time. The resulting pellet was resuspended in Tris buffer (600 mg protein/ml), now containing bacitracin (final concentration 40 µg/ml), and distributed in 0.25 ml aliquots to a set of polypropylene tubes containing 40 µM bestatin, 0.4 mg bovine albumin and with or without 30 mM Na₂SO₄ (total volume 1 ml). For binding saturation analysis, increasing concentrations of [³H]SP were added in the presence and absence of 1 µM of unlabeled SP. In competitive inhibition studies, increasing concentrations (0.1 nM-10 µM) of unlabeled SP were added in the presence of a fixed concentration of [³H]SP (0.2-2 nM). After 20 min incubation on ice, the labeled membranes were collected by filtration over Whatman GF/B filters under vacuum, followed by three rapid washes with ice-cold Tris buffer using the M-24R Cell-Harvester (Brandel Instruments). Both the tubes and the filters were soaked with 0.1% polyethylenimine for 24 h before use. The binding of [³H]SP to crude MC/9 membranes was determined from the difference between parallel experiments with membranes or the vehicle for each combined concentration of labeled and unlabeled SP to rule out contributions of [³H]SP binding to non-biological materials.

Binding data were analyzed using the LIGAND computer program (24) to obtain estimates for the apparent equilibrium dissociation constant, K_d, and maximum binding capacity, B_{max}. Models involving one or two independent classes of binding sites were evaluated for each experiment. The most appropriate model was shown on the basis of the 'extra sum-of-squares' F-test. The protein concentration was determined by Peterson's modification of the micro-Lowry method (Sigma Protein Assay Kit) using bovine serum albumin as a standard.

Histamine release studies. Cultured cells were pelleted, washed twice and resuspended in Tyrode solution of the following composition (mM): NaCl 137, KCl 2.7, NaH₂PO₄ 0.4, MgCl₂ 1, glucose 5.6, HEPES (4-(2-hydroxyethyl)-1-piperazine-ethanesulphonic acid) 10, and with or without CaCl₂ 1.8 mM (pH 7.4, 20°C). Aliquots (0.9 ml) of cell suspensions (10^6 cells/ml), subsequent to temperature equilibration (5 min, 37°C), were incubated with the indicated concentrations of a possible releasing agent or the vehicle (spontaneous release) in a volume of 0.1 ml. The reaction was stopped after 15 min incubation by chilling the tubes in an ice bath. The cells were separated from the medium by centrifugation for 10 min at 300g, 4°C. Released histamine was measured in the supernatants. Residual histamine in the cells was measured after lysing the cells by addition of distilled water to the pellets followed by protein precipitation with 20% trichloroacetic acid and centrifugation at 1000g. Histamine was assayed fluorimetrically by the method of Shore et al. (25) using an excitation wavelength of 348 nm and emission wavelength of 439 nm. Histamine standard ranging from 25 ng/ml to 250 µg/ml were used for calibration of the assay. Histamine released in the supernatant was expressed as a percentage of total histamine present in the cells plus supernatant. Spontaneous release in the absence of a releasing agent was subtracted from the stimulated release to estimate net induced release. The releasing agents were examined for possible interference with histamine determination, and corrections were made when required.




Fig. 1 Concentration dependence of the specific binding of [^3H]SP to mast cell membranes in 50 mM Tris buffer, 4°C. All values are the mean of quadruple samples. The difference between samples were less than 18%.

Results

Binding of [^3H]substance P to mast cell membranes. The binding of [^3H]SP to mast cell membranes was rapid. Time course studies of 0.2 nM [^3H]SP binding revealed that maximum specific binding was reached after 20 min incubation at 4°C (data not shown). After 1 h incubation, 87% of maximum binding remained bound, while after 3 h incubation, 72% remained bound. Hence, 20 min incubation was routinely used. Figure 1 demonstrates that specific binding of [^3H]SP is saturable. Artfactual non-receptor interactions were presumably absent since background counts were subtracted from each assay point (see Methods). The specific binding, obtained by deducting non-specific binding in the presence of 1 μM unlabeled SP from that of total binding without SP, was only 10 to 15% of total binding. For example, the addition of 18 nM [^3H]SP (797300 cpm) resulted in total [^3H]SP binding of 11154 cpm and specific [^3H]SP binding of 1267 cpm (2 mL total volume of binding mixture) after binding to nonbiological substances had been accounted for.

The shape of the saturation curve is indicative of cooperative binding. For positive cooperativity, the ascending portion of the curve is almost parallel with the y-axis (26), here indicated by [^3H]SP bound. Facilitated or cooperative binding

results in concave downward Scatchard plots (not shown), in which K_d varies with amount bound. Therefore, the parameters for the binding of [^3H]SP were not calculated based on these binding data.

Competitive inhibition studies of [^3H]SP binding by increasing concentrations of unlabeled SP were performed in high ionic strength solutions by the inclusion of Na_2SO_4 in the binding mixtures, which according to Bahouth and Masacchio (27) acts mainly by reducing nonspecific SP binding. The lowering of nonspecific SP binding was confirmed. The addition of 30 mM Na_2SO_4 to the binding mixture produced nonspecific binding as low as 35% of total binding and was routinely used in the competition experiments. The binding parameters were estimated from the displacement of 2 nM [^3H]SP binding (88000 cpm) by increasing concentrations of unlabeled SP. The radioligand was used at the lowest practical concentration for detection of binding to sites of disparate characteristics. This concentration provided about 1200 cpm of total binding and about 800 cpm of specific binding after corrections for [^3H]SP binding to non-biological material. The displacement curve with the 95% confidence limit curves (Fig. 2) represents a conglomerate plot of binding data from four separate experiments. A two-site model was found to be a best fit for [^3H]SP binding to mast cell membranes. The estimated binding parameters are shown in Table 2.

Blocking studies with the highly selective NK1 receptor ligand [pGlu⁶,Pro⁹]substance P (septide) (28) were performed to further examine the heterogeneity of SP binding sites. Fig. 3 shows the effect of 0.5 μM septide on the displacement curve in the

Table 2 Parameters for substance P binding to mast cell (MC/9) membranes in the presence of 30 mM Na_2SO_4

	Apparent K_d (nM)	B_{max} fmole/mg protein
High-affinity site	0.3	67
Low-affinity site	40.0	2700

Binding parameters were derived from the combined analysis of four displacement experiments using the LIGAND program (24). Data are shown in Figure 2.




Fig. 2 Displacements of the binding of 2nM [3 H]SP by unlabeled SP from mast cell membranes in the presence of 30mM Na_2SO_4 . The displacement curve with 95% confidence limit curves is a conglomerate plot based on four competition experiments, indicating binding to two sites (Table 2). B/T = bound divided by total radioactivity; $\log(T)$ = logarithm of the total ligand concentration.

presence of 30mM Na_2SO_4 . The absence of high-affinity displacement of [3 H]SP binding, shown by an initial plateau in the plot, is indicative of septide blocking of NK1 sites.

Release of histamine from mast cells. The amount of histamine in the MC/9 cells varied with number of days in culture. A maximum amount up to $0.608 \mu\text{g}/10^6$ cells was measured, which was more than 10-fold that of mast cells derived from murine bone marrow cultures (29), $0.053 \mu\text{g}/10^6$ cells, but much lower than that of either human pulmonary, $4.23 \mu\text{g}/10^6$ cells, or rat peritoneal mast cells, $30 \mu\text{g}/10^6$ cells, (16).




Fig. 3 Representative curve of the displacements of the binding of 2nM [3 H]SP by unlabeled SP from mast cell membranes in the presence of both 30mM Na_2SO_4 and 0.5uM septide. Analysis of binding data was based on duplicate measurements, and the experiment was repeated once with similar results.

Table 3 Histamine release from mast cells (MC/9) induced by tachykinin receptor ligands

Ligand	Conc ($M \times 10^{-6}$)	(n)	Histamine release (% of total)
Substance P (SP) ^a	0.010	(3)	5 ± 1
..	0.100	(1)	4
..	1.0	(1)	3
..	10.0^b	(1)	3
..	100.0	(2)	4 ± 0.7
SP methylester	0.010	(3)	3 ± 1
..	0.100	(2)	4 ± 0.7
Physalaemin ^a	0.010	(5)	4 ± 0.7
Neurokinin A (NKA)	0.010	(1)	0
..	0.1	(2)	2 ± 0.7
..	1	(10)	4 ± 0.6
..	10	(5)	7 ± 1
..	20	(2)	10 ± 0.7
..	50	(3)	13 ± 0.7
..	100	(5)	37 ± 5
[D-Arg ¹ , D-Phe ⁵]	0.01	(1)	0
..	0.10	(1)	0
..	1	(1)	3
D-Trp ⁷ , Leu ¹¹]	10	(2)	4 ± 2
substance P (peptide X) ^b	50	(2)	13 ± 0.7
	100	(2)	41 ± 0

The values, expressed as Mean \pm SEM ($n \geq 3$) and Mean \pm SD ($n = 2$), represent induced release, which is ligand stimulated release minus spontaneous release.

^aDetectable in the absence of extracellular CaCl_2 . ^bPotent bombesin antagonist (30) and putative SP antagonist (31)

Histamine was released from suspensions of MC/9 cells by micromolar concentrations of the NK2 preferring agonist, NKA, and only to a minor degree by nanomolar concentrations of the NK1 selective agonists, SP, SP methylester and the amphibian tachykinin, physalaemin, (Table 3). Increasing concentrations of the NK1 preferring agonists up to 100uM did not produce an increase in histamine release. Septide appeared to have no histamine releasing capability. To detect induced release for SP and physalaemin, it was necessary to omit extra-cellular CaCl_2 from the medium. This is a condition the MC/9 cells share with rat peritoneal mast cells (10). Omitting CaCl_2 extra-cellularly yielded an overall slightly lower value for the spontaneous release, $16 \pm 1\%$ ($n = 16$), as

compared to $18 \pm 1\%$ ($n = 32$) in the presence of 1.8mM CaCl_2 , but had little or no effect on induced histamine release except when SP or physalaemin were the releasers. Extending the incubation period with the releasing agents from 15 min up to 60 min did not change the values of the spontaneous release. Induced histamine release reached a maximum at 7.5 min incubation and remained constant up to 60 min, the longest period examined. [D-Arg¹,D-Phe⁵,D-Trp^{7,9},Leu¹¹] substance P (peptide X), a potent bombesin antagonist (30) and putative SP antagonist (31) induced the release of histamine in a similar concentration range as did NKA, while two selective antagonists for the NK2 receptor (32), cyclo[Gln-Trp-Phe-(R)Gly-[ANC-2]Leu-Met] (peptide A) and cyclo[Gln-Trp-Phe-Gly-Leu-Met] (peptide B), were without effect up to 1000uM . Compound 48/80, examined from 0.001 to 1000ug/ml , a range known to cause release in rat peritoneal mast cells (10, 11, 13), MCD at 1uM , the N-terminal SP fragments, SP(1-4) (0.01 - 100uM) and SP(1-6), SP(1-7) or SP(1-9) at 100uM , as well as NK3, a preferential NK3 receptor agonist, in the range 0.01 - 35uM , evoked little or no histamine release (data not shown). The two NK2 selective antagonists, peptide A and B (see above), were examined for their ability to inhibit NKA-induced histamine release. Peptide A appeared to be most effective. Preincubation of the cells with either 10 or 100uM of peptide A for 10min at 37°C prior to the exposure to equimolar concentrations of NKA reduced the histamine release by 25 and 57% , respectively. Preincubation with the SP antagonist (33), [D-Arg¹,D-Trp^{7,9},Leu¹¹] substance P (spantide), on the other hand, had no reducing effect.

The MC/9 cells, like rat peritoneal mast cells (34), released histamine with concanavalin A (con A), an IgE-type secretagogue. Incubation for 15min with 2.5 and 5ug/ml of con A without the potentiator phosphatidylerine evoked 11% and 9% induced release, respectively.

Discussion

The results demonstrate that the MC/9 mast cell line possesses specific binding sites for SP. Moreover, analysis of the binding data has revealed that

SP binds to two types of sites of vastly different binding properties. As SP interacts with all three tachykinin receptors, but preferentially with the NK1 type, the high-affinity binding is compatible with that of SP binding to NK1 receptors. While blocking of high-affinity SP binding using the selective NK1 ligand septide (Fig. 3) confirms the presence of NK1 receptors, the functional studies support the presence of NK2 receptors.

We have demonstrated here that none of the well-known histamine releasers, compound 48/80, MCD and N-terminal SP fragments evoked histamine release from the MC/9 cells. On the other hand, the results have revealed the novel observation that NKA acts as a histamine secretagogue in this cell type. Moreover, the reduction in NKA-induced histamine release observed after pretreatment with a NK2 antagonist but not with a NK1 antagonist (spantide) suggests the involvement of NK2 receptors in this process and thereby the existence of NK2 receptors on this cell line. The weak activity of the selective NK1 agonists suggests the presence of a minor population of functional NK1 sites. However, the involvement of NK1 receptors in NKA-induced histamine release was dismissed on the following grounds: (1) NKA is shown to interact preferentially with NK2 receptors; (2) pretreatment of the cells with the SP (NK1) antagonist spantide led to no loss in NKA-induced histamine release; while (3) pretreatment with a NK2 antagonist did. A lack of effect by the NK3-preferring ligand NKB precluded the possible involvement of NK3 receptors in NKA-induced histamine release.

Recently, pharmacological and biochemical evidence has suggested that there are multiple subtypes of NK2 receptors (35). A series of linear and cyclic NK2 antagonists were found to have varying potency in different NK2 containing tissues. The finding that the selective NK2 receptor antagonist, peptide A, inhibited only partially NKA-induced histamine release might indicate the presence of a heterogeneous pool of NK2 sites on MC/9 cells.

The present studies have provided evidence for the existence of NK1 and NK2 receptors on the MC/9 cell line. The coexistence of physiologically relevant NK1 and NK2 receptors has recently been described for different functions in other types of

tissues. For example, both SP and NKA were found to stimulate the release of 5-hydroxytryptamine via NK1 and NK2 receptors in the rat cerebral cortex (33). Both receptor types have also been implicated in SP-induced superoxide anion production in guinea-pig alveolar macrophages (36).

It has been argued that, because some SP antagonists, such as [D-Arg¹, D-Pro², D-Trp⁷⁻⁹, Leu¹¹] substance P, elicit histamine release (37), the induction of histamine release by SP is not tachykinin receptor-mediated. Because several antagonists of SP with two or more D-Trp residues of limited selectivity for all three tachykinin receptors have considerable agonist properties for these sites (38, 39), it is possible that we deal with a class of compounds with mixed agonist-antagonist properties. Peptide X (Table 3) is apparently such a compound. This peptide described initially as a SP antagonist that blocks the hyoscine-resistant opiate withdrawal contracture in guinea pig ileum (32), was later found to be a potent antagonist of the amphibian tetradecapeptide bombesin and structurally related mammalian peptides, including gastrin-releasing peptide (GRP), (31). Peptide X has also been found to inhibit the action of the neurohypophysial hormone vasopressin (40). Since vasopressin and the peptides of the bombesin family interact with distinct and specific high-affinity receptors, it appears that peptide X interacts as a blocking agent with several independent receptors. However, present studies indicate that peptide X, which has minimal structural homology with NKA (only Leu¹⁰), elicits a biological response. Thus, peptide X functions apparently as a mixed agonist-antagonist, expressing agonist characteristics at the NK2 receptor and antagonist characteristics at various independent receptors, including the NK1 tachykinin receptor. In conclusion, this study has demonstrated the presence of both NK1 and NK2 receptors on a murine liver derived mast cell line and is the first to show that histamine is released by NKA via NK2 tachykinin receptor activation. Whereas septide appears to have no histamine releasing activity in this cell type, other NK1 preferring agonists, e.g., SP methylester, SP and physalaemin, exert a minor stimulating effect on such release, suggesting the possible involvement of NK1 receptors. We have

shown here that in a mast cell type, unresponsive to known basic secretagogues such as compound 48/80 and N-terminal SP fragments, a NK2 receptor-dependent pathway might exist for the release of histamine.

Acknowledgements

This work was done while the author, Dr S. A. Krumm, held a National Research Council (USAMRICD) Senior Research Associateship. The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Army or the Department of Defense.

References

- 1 Lembeck, F. (1988). The 1988 Ulf von Euler Lecture. Substance P. From extract to excitement. *Acta Physiol. Scand.* 133: 435-454.
- 2 Regoli, D., Drapeau, G., Dion, S. and D'Orleans-Juste, P. (1987). Mini-review. Pharmacological receptors for substance P and neurokinins. *Life Sci.* 40: 109-117.
- 3 Helke, C. J., Krause, J. E., Mantyh, P. W., Couture, R. and Bannon, M. J. (1990). Diversity in mammalian tachykinin peptidergic neurons: multiple peptides, receptors, and regulatory mechanisms. *FASEB J.* 4: 1606-1615.
- 4 Regoli, D., Drapeau, G., Dion, S. and Couture, R. (1988). New selective agonists for neurokinin receptors: pharmacological tools for receptor characterization. *Trends Pharmacol. Sci.* 9: 290-295.
- 5 Johnson, A. R. and Erdos, E. G. (1973). Release of histamine from mast cells by vasoactive peptides. *Proc. Soc. Exp. Biol. Med.* 142: 1252-1256.
- 6 Erjavec, F., Lembeck, F., Horjanc-Irman, T., Skofitsch, G., Donnerer, J., Saria, A. and Holzer, P. (1981). Release of histamine by substance P. *Arch. Pharmacol.* 317: 67-70.
- 7 Devillier, P., Regoli, D., Aseraf, A., Descours, B., Marsac, J. and Renoux, M. (1986). Histamine release and local responses of rat and human skin to substance P and other mammalian tachykinins. *Pharmacology* 32: 340-347.
- 8 Church, M.K., Benyon, R. C., Lowman, M. A., Hutson, P. A. and Holgate, S. T. (1989). Allergy or inflammation? From neuro-peptide stimulation of human skin mast cells to studies on the mechanism of the late asthmatic response. *Agents and Actions* 26: 22-30.
- 9 Devillier, P., Drapeau, G., Renoux, M. and Regoli, D. (1989). Role of the N-terminal arginine in the histamine-releasing activity of substance P, bradykinin and related peptides. *Eur. J. Pharmacol.* 168: 53-60.
- 10 Fewtrell, C. M. S., Foreman, J. C., Jordan, C. C., Oehme, P., Renner, H. and Stewart, J. M. (1982). The effects of substance P on histamine and 5-hydroxytryptamine release in the rat. *J. Physiol.* 330: 393-411.
- 11 Mazurek, N., Pecht, L., Feichberg, V. I. and Blumberg, S. (1981). The role of the N-terminal tetrapeptide in the

- histamine releasing action of substance P. *Neuropharmacology* 20: 1025-1027.
12. Moush, M., Bueh, J.-L., Bronner, C., Rouot, B. and Landry, Y. (1990). G protein: a receptor-independent mode of action for cationic amphiphilic neuropeptides and venom peptides. *Trends Pharmacol. Sci.* 11: 358-362.
 13. Moush, M., Bronner, C., Landry, Y., Bockaert, J. and Rouot, B. (1990). Direct activation of GTP-binding regulatory proteins (G-proteins) by substance P and compound 48/80. *FEBS Letters* 259: 260-262.
 14. Repke, H. and Bienert, M. (1987). Mast cell activation - a receptor-independent mode of substance P action. *FEBS Letters* 221: 236-240.
 15. Feldberg, W. and Mongar, J. L. (1954). Comparison of histamine release by compound 48/80 and octylamine in perfused tissues. *Br. J. Pharmacol.* 9: 197-201.
 16. Ennis, M. (1982). Histamine release from human pulmonary mast cells. *Agents and Actions* 12: 60-63.
 17. Pearce, F. L. (1986). On the heterogeneity of mast cells. *Pharmacology* 32: 61-71.
 18. Foreman, J. C. and Lichtenstein, L. M. (1980). Induction of histamine secretion by polycations. *Biochim. Biophys. Acta* 629: 587-603.
 19. Ali, H., Leung, K. B. P., Pearce, F. L., Hayes, N. A. and Foreman, J. C. (1986). Comparison of the histamine-releasing action of substance P on mast cells and basophils from different species and tissues. *Int. Arch. Allergy App. Immunol.* 79: 413-418.
 20. Nabel, G., Fresno, M., Chessman, A. and Cantor, H. (1981). Use of cloned populations of mouse lymphocytes to analyze cellular differentiation. *Cell* 23: 19-28.
 21. Burcher, E., Müssap, C. J., Geraghty, D. P. and McClure-Sharp, J. M. (1990). Concepts in characterization of tachykinin receptors. Abstract and oral presentation at the Substances P and Related Peptides: Cellular and Molecular Physiology, New York Academy of Sciences Conference, Worcester, Mass. 18 to 21 July 1990.
 22. Krumins, S. A. (1990). Binding of [³H]-substance P to mast cells. Abstract and poster presentation at *ibid.*
 23. Maruyama, M. (1986). Selective solubilization of physalamin-type substance P binding sites from rat brain membranes by glycodeoxycholate and NaCl. *Brain Res.* 370: 186-190.
 24. Munson, P. J. and Rodbard, D. (1980). LIGAND: A versatile computerized approach for characterization of ligand-binding systems. *Anal. Biochem.* 45: 530-556.
 25. Shore, P. A., Burkhalter, A. and Cohn, V. H. (1959). A method for the fluorometric assay of histamine in tissues. *J. Pharmacol. Exp. Ther.* 127: 182-186.
 26. De Lean, A. and Rodbard, D. (1979). Kinetics of cooperative binding. In: *The Receptors*, Vol. 1, (O'Brien, R. D. ed.) Plenum Publishing Corporation, p. 143-192.
 27. Bahouth, S. W. and Musacchio, J. M. (1985). Specific binding of [³H]-substance P to the rat submaxillary gland. The effects of ions and guanine nucleotides. *J. Pharmacol. Exp. Ther.* 234: 326-336.
 28. Wormser, U., Laufer, R., Hart, Y., Chorev, M., Gilon, C. and Selinger, Z. (1986). Highly selective agonists for substance P receptor subtypes. *EMBO J.* 5: 2805-2808.
 29. Wodnar-Filipowicz, A., Heusser, C. H. and Moroni, C. (1989). Production of the haemopoietic growth factors GM-CSF and interleukin-3 by mast cells in response to IgE receptor-mediated activation. *Nature* 339: 150-152.
 30. Woll, P. J. and Rozenzurg, E. (1988). [D-Arg¹, D-Phe⁴, D-Trp^{7,9}, Leu¹¹] substance P, a potent bombesin antagonist in murine Swiss 3T3 cells, inhibits the growth of human small cell lung cancer cells in vitro. *Proc. Natl. Acad. Sci. USA* 85: 1859-1863.
 31. Tsou, K., Wu, S.-X., Lu, Y.-A. and Way, E. L. (1985). Block of the hyoscine-resistant opiate withdrawal contraction of ileum by a new substance P antagonist [D-Arg¹, D-Phe⁴, D-Trp^{7,9}, Leu¹¹] substance P. *Eur. J. Pharmacol.* 110: 155-156.
 32. Williams, B. J., Curtis, N. R., McKnight, A. T., Maguire, J., Foster, A. and Tridgett, R. (1988). Development of NK-2 selective antagonists. *Reg. Pep.* 22: 189.
 33. Iverfeldt, K., Solii, M. and Barfai, T. Y. (1990). Substance P and Neurokinin A, two coexisting tachykinins stimulating the release of [³H]-5-HT from rat cerebral cortical slices. *Brain Res.* 506: 335-338.
 34. Carnuccio, R., Di Rosa, M., Ialenti, A., Iuvone, T. and Sautebin, L. (1989). Selective inhibition by vasocortin of histamine release induced by dextran and concanavalin-A from rat peritoneal cells. *Br. J. Pharmacol.* 98: 32-36.
 35. Buck, S. H. and van Giersbergen, P. L. M. (1990). Pharmacological and biochemical evidence for multiple types of tachykinin NK₂ receptors. Abstract and oral presentation at the Substance P and Related Peptides: Cellular and Molecular Physiology, New York Academy of Sciences Conference, Worcester, Mass. 18 to 21 July 1990.
 36. Brunelleschi, S., Vanni, L., Ledda, F., Giotti, A., Maggi, C. A. and Fantozzi, R. (1990). Tachykinins activate guinea-pig alveolar macrophages: involvement of NK₂ and NK₁ receptors. *Br. J. Pharmacol.* 100: 417-420.
 37. Hakanson, R., Horig, J. and Leander, S. (1982). The mechanism of action of a substance P antagonist (D-Pro², D-Trp^{7,9})-SP. *Br. J. Pharmacol.* 77: 697-700.
 38. Regoli, D., Mizrahi, J., D'Orleans-Juste, P., Dion, S., Drapeau, G. and Escher, E. (1985). Substance P antagonists showing some selectivity for different receptor types. *Eur. J. Pharmacol.* 109: 121-125.
 39. Buck, S. H. and Shatzer, S. A. (1988). Agonist and antagonist binding to tachykinin peptide NK-2 receptors. *Life Sci.* 42: 2701-2708.
 40. Zachary, I. and Rozenzurg, E. (1986). A substance P antagonist also inhibits specific binding and mitogenic effects of vasopressin and bombesin-related peptides in Swiss 313 cells. *Biochem. Biophys. Res. Commun.* 137: 135-141.



1

Jada F
CASH
RAB
100000
JANUARI

By	
Distributer	
Availability	
Dist	Avail
A-1	20