· · ·					(		
SECURITY CL AD-A253		MENTATION	PAGE				
Ta. REPORT :		TID. RESTRICTIVE					
TO, THE STATE INTO A STATE AND A ST		NA					
2a. SECURITY CLASSIFICATION UP ORTY	3. DISTRIBUTION / AVAILABILITY OF REPORT						
25. DECLASSIFICATION / DOWNGN, ING SCHEDU	<u>6 1992 </u>	4					
NA		Distribution Unlimited					
4. PERFORMING ORGANIZATION REPORT NUMIER(S)		S. MONIFORING ORGANIZATION REPORT NUMBER(S)					
NA	NA						
6a. NAME OF PERFORMING ORGANIZATION	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION					
University of Tennessee	NA	Office of Naval Research					
6c. ADDRESS (City, State, and ZIP Code)	7b. AUDRESS (City, State, and ZIP Code)						
Center for Environemntal Biot		800 N. Quincy Street					
10515 Research Drive Suite 30 Knoxville, TN 37932	0	Arlington, VA 22217-5000					
8a. NAME OF FUNDING/SPONSORING	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER						
ORGANIZATION	8b. OFFICE SYMBOL (If applicable)						
Office of Naval Research	ONR	N00014-88-K-0489					
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street		10. SOURCE OF FUNDING NUMBERS PROGRAM PROJECT TASK WORK UNIT					
Arlington, VA 22217-5000		ELEMENT NO.	NO.	NO.	ACCESSION NO		
11. TITLE (Include Security Classification) Ecol		<u> </u>	1123B	422s001	l		
Final FROM 5/ 16. SUPPLEMENTARY NOTATION Ecology of Microfouling Community Biosig					tial		
17. COSATI CODES	Continue on reverse if necessary and identify by block number)						
FIELD GROUP SUB-GROUP	ia - Initial Microfouling Community, Test						
	Antifouling	lest	•				
19. ABSTRACT (Continue on reverse if necessary The research program represent examination of the lipid meta 2) the analysis of the initia biomarker techniques, and 3) microbial biofouling on paint Significant progress was made from <u>in situ</u> samples and in the chemical noise were developed system for detecting biofilm the initial microfouling organ	nted a consolida abolism of the A al microfouling the development ed surfaces for in detecting a pioreactors. Ne for assessing formation and s	tion of thre rchaebacteri community us of an asses the Antifou nd monitorin w methodolog the initial ublethal tox	a from the h ing ultrasen sment progra ling Coating g the hydrot ies for sign microfouling	ydrothermal sitive sign m for the d s Developem hermal vent ificantly r film and a	vents, ature lipid etection of ent Program. microbiota educing test		
	RPT. DTIC USERS Redition may be used un All other editions are o	(U) 22b. TELEPHONE ( (202) 696– ntil exhausted.					
<b>92–18660</b>	92		i	2	•		

······································						
(U)						
SECURITY CLASSIFICATION OF THIS PAGE				·		
1a. REPORT SECURITY CLASSIFICATION	REPORT DOCU	TIL RESTRICTIVE				
		NA				
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION / AVAILABILITY OF REPORT				
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE		Distribution Unlimited				
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)				
NA		NA				
6a. NAME OF PERFORMING ORGANIZATION	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION				<u>.</u>
University of Tennessee	NA	Office of Naval Research				
5c. ADDRESS (City, State, and ZIP Code)	7b. AUDRESS (Ci	ty, State, and ZIF	Code)			
Center for Environemntal Bio 10515 Research Drive Suite 3	800 N. Quincy Street Arlington, VA +22217-5000					
Knoxville, TN 37932						
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMEN	IT INSTRUMENT I	DENTIFIC	ATION NU	IMBER ·
Office of Naval Research	ONR	N00014-88-K-0489 10. SOURCE OF FUNDING NUMBERS				
8c. ADDRESS (City, State, and ZIP Code) . 800 N. Quincy Street		PROGRAM	PROJECT	TASK	·····	WORK UNIT
Arlington, VA 22217-5000		ELEMENT NO.	NO. 1123B	NO.	s001	ACCESSION NO
1. TITLE (Include Security Classification) Eco	logy of Archaeab	acteria for				d the
6. SUPPLEMENTARY NOTATION Ecology Microfouling Community Biosi	gnatures - Antif	ouling Coati	ing Developm	ent P	rogram	
17. COSATI CODES FIELD GROUP SUB-GROUP	18. SUBJECT TERMS ( Archaebacter					
	Antifouling				···	,
19. ABSTRACT (Continue on reverse if necessar The research program represe examination of the lipid met 2) the analysis of the initi biomarker techniques, and 3) microbial biofouling on pair Significant progress was mad from <u>in situ</u> samples and in chemical noise were develope system for detecting biofilm the initial microfouling org	ented a consolidate abolism of the A al microfouling the development ated surfaces for le in detecting a bioreactors. Ne ad for assessing a formation and s	tion of three archaebacters community us of an asses the Antifou and monitorin w methodolog the initial sublethal to	a from the sing ultrase sement progra ling Coatin ng the hydro gies for sig microfoulin dicity on the	hydro nsiti am fo gs De therm nific g fil	thermal ve sign or the d velopen al vent antly r .m and a	vents, nature lipic letection of nent Program microbiota reducing test
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT						
UNCLASSIFIED/UNLIMITED SAME AS 22a. NAME OF RESPONSIBLE INDIVIDUAL	RPT. DTIC USERS	22b. TELEPHONE	(Include Area Cod	e) 22c.	OFFICE SY	MBOL
Dr. Bernard J. Zahuranec	APR edition may be used ur	(202) 696- ntil exhausted.	·		ONR	
	All other editions are o		SECURITY	•	FICATION ( U)	OF THIS PAGE
						:

#### **Contract #:** N0014-88-K-0489

Principal Investigators: David C. White, James B. Guckert

Institute: Institute for Applied Microbiology, University of Tennessee

**Contract Title:** Ecology of Archaebacteria, for Extreme Environments and the Initial Microfouling Community by Signature Biomarker Techniques

**Period of Performance:** 05-01-88 through 04-03-91

#### **Objectives:**

**\***` '

1. To develop and analyze the ether lipid components of the Archaebacteria from the hydrothermal vents.

2. To develop sensitive methods for analysis of signature biomarkers for the initial microfouling community.

3. To develop an assay system for the formation and toxicology of the microbial fouling community on the surfaces of developed in the antifouling coatings program.

### Accomplishments:

¥

1. We were able to develop a chromatographic method for the separation and quantitative analysis of the ether lipids from Archaebacteria. These lipids which consist of diphytanyl- and bidiphytanyl-glycerol ether derivatives are too high molecular weight to be made volatile at the temperatures that are possible for gas chromatography. D. B. Hedrick developed a method based on supercritical fluid chromatography (SFC) with supercritical carbon dioxide with a pressure gradient where he was able t separate the large molecular weight alcohols formed after the acid methnolysis of the lipids. These were then separated by capillary chromatography and detected by flame ionization detection. The analysis allowed us to show that there are diphytanyl glycerol ethers (DPE) and bi-diphytanyl glycerol ethers (BGE) in the neutral lipid, glycolipid, and polar lipid We were able to account for low levels in some fraction. Archaebacteria by showing that in these organisms the lipids are not extractible until after acid methanolysis (1). Using this technique it was possible to develop a rapid method for distinguishing the ether lipids utilizing FT/IR (2).

2. The principal problem with the ultrasensitive analysis of polar lipid biomarkers for very sparse microbial biofilms such as the initial microfouling community is the contamination of the solvent used for extraction with fatty acids. This leads to chemical noise. To decrease this problem we developed a microtechnique where we utilize a very small volume 5 ml total of

extract and microcolumns for extraction. This was then further improved by a "reverse serial extraction technique" in which the same small volume of extract is repeatedly exposed to biofilm samples. The development of a o-ring device based on a joint that is clamped to the surface for recovery of the contents inside the extractor allowed the recovery of multiple samples of biofilm from the same coupon. With this technique and the utilization of single ion extraction software in the gas chromatograph/mass spectrometer (GC/MS) it was possible to detect sub picomolar quantities of microbes in biofilms. This technique has been published in relationship to the recovery of microbes from the deep subsurface (3). With this technique and new developments it proved possible to show absolute correspondence between the phylogenetic relationships based on 16S RNA sequences and the cluster analyses of the phospholipid ester-linked fatty acids (4).

3. To evaluate the microfouling potential of newly developed antifouling coatings we developed a marine bacterial challenge assay. This assay was designed to test both the microfouling potential as well as the toxicological effects with respect to the bacterial physiological stress associated with exposure to any leached materials. This challenge assay can be expanded to include a marine diatom to provide an algal/bacterial challenge assay. The system developed involved an oligotrophic, continuous-flow (approximately 5 ml/min) flow of ASTM seawater with very dilute nutrients containing an inoculum of Pseudomonas atlantica, a gram-negative bacteria like those in the initial microfouling community. With this system the U.S. Navy standard F121 copper based paint was used and compared to stainless steel surfaces. Although the F121 paint gave off between 4 -8 mg Cu/L the bulk phase microbial densities remained at 3 x  $10^6$  and the biofilms developed to densities of  $10^9$  cells/sq. cm. The test system also allows the detection of sublethal toxicity as the oring extraction system allows recovery of the organisms which are then extracted and tested for the proportions of cyclopropane and trans monoenoic phospholipid ester-linked fatty acids. Both these accumulate in <u>P. atlantica</u> that are exposed to toxic stimuli. This test system in the presence of toxins such as mercury show these changes.

5

## Significance:

ŧ

Y

In this research program the methods and techniques necessary to study the ecology of the hydrothermal vent extreme thermophilic Archaebacteria were developed. The test system using the <u>P. atlantica</u> challenge assay system was developed and shown to be effective in testing the anti-fouling coatings developed in the ONR program.

Future Programs:

Research based on the data and techniques developed in this test system have led to the development of an on-line test system utilizing the bioluminescent bacteria.

# Publications:

t

1. Hedrick, D. B., J. B. Guckert, and D. C. White. 1991a. Archaebacterial ether lipid diversity analyzed by supercritical fluid chromatography: Integration with bacterial lipid protocol. J. Lipid Research <u>32</u>: 659-666.

2. Hedrick, D. B., D. F. Nivens, C. Stafford, and D. C. White. 1991b. Rapid differentiation of Archaebacteria from eubacteria by diffuse reflectance Fourier-transform infrared spectoscopic analysis of lipid preparations. J. Microbiol. Methods <u>13</u>: 67-73.

3. White, D. C., D. B. Ringelberg, J. B. Guckert, and T. J. Phelps. 1991. Biochemical markers for <u>in situ</u> microbial community structure. <u>In</u> Proceedings of the First International Symposium on Microbiology of the Deep Subsurface, (C. B. Fliermans and T. C. Hazen, eds.). January 15-19, 1990, Orlando, FL. WSRC Information Services, Aiken, SC. pp. 4-45 to 4-56.

4. Guckert, J. B., D. B. Ringelberg, D. C. White, R. S. Henson, B. J. Bratina. 1991. Membrane fatty acids as phenotypic approach to taxonomy of methylotrophs within the proteobacteria. J. Gen. Microbiol. <u>137</u>: in press.

Accession For NTTO GRAAI WALL TAB Unestablineed  $\Box$ Justification. Dv\_ Distribution/ Availability Codes Avail and/or Special Dist