	_			
A)-A254 0	94 ENTATION P	PAGE	OMB NC 0704 3188
		mategina (Jarket) in Urio, previewing the Dilard on S Sourcen - Mashingtine e Driver - Mashingtine e Driver - Mashenent (r	Prospecter on using the time thread end in othermation rend comments regard on nadquarters her uses unredonate for of to sudget Paperwork Reduction Project Component Typer Apple.	
1		July 1992	Final 1 Ap	r 89-30 Apr 92
4. TITL New	E AND SUBTITLE Developments in .	Atom Interferometry	DTIC 5.	FUNDING NUMBERS
6. AUT	THOR(S) David E. Pritc	hard	AUG 1 7 1992	DAAL03-89-K-0082
7. PER	FORMING ORGANIZATION M Massachusetts In Cambridge, MA	NAME(S) AND ADDRESS(ES)	у у	PERFORMING ORGANIZATION REPORT NUMBER
9. SPO U.	S. Army Research	ENCY NAME(S) AND ADDRESS(E Office	5) 10	D. SPONSORING / MONITORING AGENCY REPORT NUMBER
P. Res	0. Box 12211 search Triangle Pa	rk, NC 27709-2211		ARO 26566.5-PH
au po	thor(s) and should sition, policy, or	nd/or findings contained as decision, unless so	ined in this report an official Depart designated by othe	are those of the ment of the Army r documentation.
au po 12a D Ap	thor(s) and should sition, policy, or ISTRIBUTION / AVAILABILITY proved for public	release; distribution	ined in this report an official Depart designated by othe 12 n unlimited.	are those of the ment of the Army r documentation. 2b. DISTRIBUTION CODE
au <u>po</u> 12a D Ap 13. AB	thor(s) and should sition, policy, or ISTRIBUTION / AVAILABILITY proved for public	release; distribution	ined in this report an official Departs designated by othe n unlimited.	are those of the ment of the Army r documentation. 2b. DISTRIBUTION CODE
au po 12a D Ap 13. AB T a W T i	thor(s) and should sition, policy, or ISTRIBUTION / AVAILABILITY proved for public STRACT (Maximum 200 wor the first true ator s optical elements as constructed whi his demonstration interometers which	and/or findings contain i not be construed as <u>c decision, unless so</u> STATEMENT release; distribution (as) m interferometer was a s for atom deBroglie ich physically separation was closely followed used laser light as	ined in this report an official Departs designated by othe n unlimited.	are those of the ment of the Army r documentation. The DISTRIBUTION CODE g transmission gratin ing atom interferomet re recombining them. ons of atom
au po 12a D Ap 13. AB T a W T i A i o s	thor(s) and should sition, policy, or ISTRIBUTION / AVAILABILITY proved for public STRACT (Maximum 200 wor The first true ator s optical elements vas constructed whi this demonstration nterometers which tom interferometer nvolving inertial f basic quantum pl tructures using at	and/or findings contain interferometer was a sofor atom deBroglie was closely followed used laser light as for studies of a number of a sofor and may ultimation con holograms.	ined in this report an official Departs designated by othe n unlimited.	are those of the ment of the Army r documentation. The DISTRIBUTION CODE g transmission gratin ing atom interferomet re recombining them. ons of atom types of experiments r properties, tests to make ultra-small
au po 12a O Ap 13. AB T a w T i A i o s	thor(s) and should sition, policy, or ISTRIBUTION / AVAILABILITY proved for public STRACT (Maximum 200 wor the first true ator is optical elements as constructed which this demonstration interometers which tom interferometer nvolving inertial f basic quantum pl tructures using at	and/or findings contain in not be construed as <u>decision, unless so</u> STATEMENT release; distribution (ds) in interferometer was s for atom deBroglie was closely followed used laser light as rs will make possible effects, studies of a hysics, and may ultimation holograms.	ined in this report an official Departs designated by othe n unlimited.	are those of the ment of the Army <u>r documentation.</u> The DISTRIBUTION CODE g transmission gratin ing atom interferomet re recombining them. ons of atom types of experiments r properties, tests to make ultra-small
au po 12a D Ap 13. AB T a W T i i A i o s	thor(s) and should sition, policy, or istribution / AVAILABILITY proved for public STRACT (Maximum 200 wor the first true ator s optical elements as constructed which this demonstration nterometers which tom interferometer nvolving inertial f basic quantum pl tructures using at	and/or findings contain in not be construed as <u>decision, unless so</u> STATEMENT release; distribution ds) in interferometer was s for atom deBroglie ich physically separat was closely followed used laser light as rs will make possible effects, studies of a hysics, and may ultimation holograms.	ined in this report an official Departs designated by othe n unlimited.	are those of the ment of the Army r documentation. b. DISTRIBUTION CODE g transmission gratin ing atom interferomet re recombining them. ons of atom types of experiments r properties, tests to make ultra-small 15. NUMBER OF PAGES
au po 12a D Ap 13. AB T a w T i A i o s	thor(s) and should sition, policy, or istribution, AVAILABILITY proved for public STRACT (Maximum 200 wor the first true ator is optical elements tas constructed which this demonstration interometers which tom interferometer involving inertial f basic quantum pl tructures using at tructures using at tructures using at tructures using at	and/or findings contain in not be construed as <u>decision, unless so</u> STATEMENT release; distribution (ds) in interferometer was s for atom deBroglie is for atom deBroglie is was closely followed used laser light as rs will make possible effects, studies of a hysics, and may ultimation holograms.	ined in this report an official Departs designated by othe n unlimited.	are those of the ment of the Army r documentation. b. DISTRIBUTION CODE g transmission grating ing atom interferometer re recombining them. ons of atom types of experiments r properties, tests to make ultra-small 15. NUMBER OF PAGES 4 15. PRICE CODE
au po 12a D Ap 13. AB T a W T 13. AB T a W T 14. SU	thor(s) and should sition, policy, or istribution / AVAILABILITY proved for public STRACT (Maximum 200 wor the first true ator s optical elements vas constructed whi this demonstration nterometers which tom interferometer nvolving inertial f basic quantum pl tructures using at tructures using at	and/or findings conta. I not be construed as <u>decision, unless so</u> STATEMENT release; distribution (ds) In interferometer was of s for atom deBroglie was ich physically separative was closely followed used laser light as rs will make possible effects, studies of a hysics, and may ultimation tom holograms. 18. SECURITY CLASSIFICATION OF THIS PAGE	ined in this report an official Departs designated by othe n unlimited. demonstrated. Usin waves, a three grat tes atom waves befo by two demonstrati the beam splitters. qualitatively new atomic and molecular ately open the way tings, Atom Optics,	are those of the ment of the Army r documentation. 2b. DISTRIBUTION CODE g transmission grating ing atom interferometer re recombining them. ons of atom types of experiments r properties, tests to make ultra-small 15. NUMBER OF PAGES 4 15. PRICE CODE TION 20. LIMITATION OF ABSTR

Prescribed by ANSI Std. 239/15 298/102

New Developments in Atom Interferometry: Final Report DAAL03-89-K-0082 (ARO proposal number 26566-PH) David E. Pritchard Massachusetts Institute of Technology Department of Physics/Research Laboratory of Electronics

During this grant period we demonstrated the first true atom interferometer. Using transmission gratings which we fabricated ourselves as optical elements for atom deBroglie waves, we constructed a three grating atom interferometer which physically separates atom waves before recombining them. Our demonstration was closely followed by two demonstrations of atom interferometers which used laser light as the beam splitters.

Atom interferometers will make possible qualitatively new types of experiments involving inertial effects, studies of atomic and molecular properties, tests of basic quantum physics, and may ultimately open the way to make ultra-small structures using atom holograms.

The relatively large mass and low velocity of atoms makes atom interferometers especially sensitive to inertial effects such as rotation, acceleration, and gravity. Sagnac rotation in accord with theoretical predictions has been observed (4) and sensitivity to gravitational acceleration at the 3×10^{-6} level demonstrated (5). Atom interferometers may become the best absolute accelerometers and gravimeters in the next few years.

Atom interferometers can be applied to a number of experiments in fundamental physics: tests of quantum mechanics such as the Aharonov-Casher effect (6), measurement of the equality of proton and electron charges, and a precise measurement of the momentum of a photon. This latter measurement should produce a new high precision value for the fundamental constants $N_A\hbar$.

Interferometers for atoms and molecules, will offer more accurate ways to measure intrinsic properties of these particles, like their polarizability. They will also open up new areas of study, such as measurements of the "index of refraction" of a gas for a particle beam which passes through it.

The key component of our interferometer is the set of three matched transmision diffraction gratings which we constructed at the National Nanofabrication Facility (NNF) at Cornell University. Our interferometer consists of three 400 nm period gratings, mounted 0.66m apart on separate translation stages inside the vacuum envelope. During operation, the 0th and 1st order beams from the first grating strike the middle grating (which is 140 μ m wide) where they are diffracted in the 1st and -1st orders so that they converge at the third grating. At the second (middle) grating the beams have widths of 30 μ m (FWHM) and are separated by 27 μ m. The first two gratings form an interference pattern in the plane of the third grating, which acts as a mask to sample this pattern. The detector, located 0.30 m beyond the third grating, records the flux transmitted by the third grating.

The data necessary to determine the interferometer phase contrast are acquired by modulating the position of one grating relative to the other two and simultaneously recording the signal from the atom counting electronics as well as the signal from an optical interferometer used to measure the relative position of the gratings. After removing data obscured by noise spikes from the hot wire, the atom count rate data are averaged into bins according to relative grating position, resulting in a fringe pattern.

The peak to peak amplitude of our interference signal is 70 Hz, which enables us to determine the interferometer phase to a precision of 0.1 rad in 1 min. The excellent long-term stability of our position stabilization system provides measured atom-interferometer phase drift of less than 0.1 rad over 10 min.

In addition to our progress with the atom interferometer, we also devoted effort to the development of new processes to fabricate atom optics as well as their fabrication. Our students made two trips to Cornell University and the NNF to develop and build gratings. The first trip resulted in the development of a new process for fabricating atom optics. The process allows fabrication of precisely positioned openings in thin silicon nitride membranes mounted in silicon frames. The pattern created in the membrane is determined by an electron beam writer, making the process quite versitile. This process was used to create the diffraction gratings used in the interferometer. In addition, several zone plates (atom lenses) were also built, and were later successfully demonstrated. During our second trip we devised ways to reduce the electron beam writing time. This decreased thermal drift during the writing period assuring higher overall accuracy, and also increased our overall productivity. We made a wide variety of diffraction gratings with various heights and periods between 100 and 300 nm, as well as an assortment of single and double slits.

Since the development of the interferometer, we have added a second year graduate student and a visiting postdoctoral fellow to our research team, which now has two graduate students who have finished the two written parts of their qualifiers (one has finished his oral as well). This has enabled us to deal with the experiment on several fronts: we improved the detector and the source, and improved the vacuum system, particularly in the main chamber where the beam used to suffer considerable attenuation. We have also built a septum to place between the two legs of our interferometer. This consists of a mount with a tightly stretched aluminum foil in the middle which extends along the beam axis for 10 cm. It is positioned by a three degree of freedom rotator/translator and we have been able to position it so that its shadow is only 35 microns wide as seen by the atomic beam.

The scientific future of atom interferometers looks bright: atom beam sources are inexpensive and intense relative to other particle beams/sources (eg. neutrons, electrons), several techniques have now been demonstrated to make interferometers for them, and the atoms which may be used in them come with a wide range of parameters such as polarizability, mass, and magnetic moment. This assures the applicability of these instruments to a wide range of measurements of both fundamental and practical interest.

We wrote several papers which were accepted or published, and most science magazines have now written up our pioneering work in atom interferometry.

PUBLICATIONS SUPPORTED BY ARO

An Interferometer for Atoms, D. W. Keith, C. R. Ekstrom, Q. A. Turchette, D. E. Pritchard: Phys. Rev. Lett. 66, 2693 (1991).

Free-standing gratings and lenses for atom optics, D. W. Keith, M. J. Rooks: J. Vac. Sci. Technol. B 9, 2846 (Nov/Dec 1991)

Numerical Model of a Multiple Grating Interferometer, Q. A. Turchette, D. E. Pritchard, D. W. Keith: Submitted to JOSA B, (1991).

"Atom Interferometers", D.E. Pritchard, Opt. Photonics News, p. 19 (November 1991).

"Atom Optics", D.E. Pritchard, In *Atomic Physics* 12, Ed. J.C. Zorn, R.R. Lewis and M. Weiss, New York, American Journal of Physics, 1991, pp. 165-170.

THESES

. .

David W. Keith, "An Interferometer for Atoms" Ph.D. Department of Physics, MIT 1991.

Quentin A. Turchette, "Numerical Model of a Three Grating Interferometer for Atoms", S.B., Department of Physics, 1991.

Acces	sion For	
NTIS DTIO Undum Jugat	- 684 &1 743 9900-00 <u>1</u> 1-0001-011	
By Lines Lines	11 - 11 - 14 - 10 - 19 - 19 - 19 - 19 - 19 - 19 - 19	······································