

2

OFFICE OF NAVAL RESEARCH

AD-A254 599



Contract N00014-85-K-0896

Technical Report No. 47

Luminescent Nitro Derivatives of 5,11-Dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole

by

Q. Lu and J. H. Boyer

Published in

Heteroatom Chem., 1992, accepted.

DTIC  
S ELECTE D  
AUG 24 1992  
A

Department of Chemistry  
University of New Orleans  
New Orleans, LA 70148-2820

August 7, 1992

Reproduction in whole or in part is permitted for any purpose  
of the United States Government

This document has been approved for public release and sale;  
its distribution is unlimited

92-23488



40909/ 16pgs

92 8 21 161

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION <b>Unclassified</b>		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE <b>Unclassified</b>			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report # 47 Contract N00014-85-K-0896		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION University of New Orleans Department of Chemistry	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) University of New Orleans New Orleans, LA 70148		7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-85-K-0896	
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Luminescent Nitro Derivatives of 5,11-Dehydro-5H,11H-benzotriazol[2,1-a]benzotriazole			
12. PERSONAL AUTHOR(S) Q. Lu and J. H. Boyer			
13a. TYPE OF REPORT Technical Report	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) August 7, 1992	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  See Attached Report			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> OTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Joseph H. Boyer		22b. TELEPHONE (Include Area Code) (504) 286-7225	22c. OFFICE SYMBOL

## Luminescent Nitro Derivatives of Benzotriazolo[2,1-a]benzotriazole

Qi Lu and Joseph H. Boyer\*

Department of Chemistry, University of New Orleans, New Orleans, LA 70148

Received ...

## ABSTRACT

Fluorescence was enhanced and laser activity introduced by substitution in 5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole 6 to give 2-nitro, 2,8-dinitro, 2,4,8-trinitro, and 2,4,8,10-tetranitro derivatives 9a-d. Luminescence for compounds 6, 9a-d, and the 2,8-dinitro-3,9-dimethyl and 2,3,8,9-tetramethyl-4,10-dinitro derivatives 11a,b was erratically solvent dependent when examined in ethyl acetate, acetonitrile, and acetone, and was most efficient in the 2,8-dinitro derivative 9b, [ $\lambda_f$  479 nm (ethyl acetate)  $\Phi$  0.98,  $\lambda_f$  501 nm (acetonitrile)  $\Phi$  0.58, and  $\lambda_f$  494 nm (acetone)  $\Phi$  0.61] and in the tetranitro derivative 9d [ $\lambda_f$  509 nm (acetonitrile)  $\Phi$  0.81 and  $\lambda_f$  511 nm (acetone)  $\Phi$  0.66]. With laser activity at 560-590 nm (acetonitrile) the dye 9b was 30 percent as efficient as rhodamine 6G (ethanol) in power output. Luminescence was quenched by the reduction of nitro groups to give 2-amino and 2,8-diamino derivatives 9e,f and by the conversion of the tetranitro compound 9d to an unassigned diazido dinitro derivative 9g. Luminescence was not detected in 2,5-dimethyl-3,6-dinitro-1,3a,4,6a-tetraazapentalene 14 and ethyl 2,5-dimethyl-1,3a,4,6a-tetraazapentalene-3,6-dicarboxylate 15. Azidoazobenzenes were obtained from 4-methyl- and 4,5-dimethyl-


DTIC QUALITY INSPECTED 5

A-1




1,2-phenylene diamines via oxidation with lead dioxide to aminoazobenzene derivatives followed by treatment of the diazotized amines with sodium azide and thermolysis of azido intermediates to give 3,9-dimethyl and 2,3,8,9-tetramethyl derivatives 10 a,b of the triazolotriazole 6. Nitration converted the triazole 6 to the 2,4,8-trinitro derivative 9c and the alkyltriazoles to their dinitro derivatives 11 a,b.

### INTRODUCTION


Rarely has the C-nitro substituent, a "killer of luminescence" [1], been contained in a fluorescent or laser dye. The known laser dyes with nitro substituents included 4-dimethylamino-4'-nitrostilbene 1 [2-4], 1,1'-diethyl-11-nitro-4,4'-quinodicyanone tetrafluoroborate 2 [5], 2-amino-7-nitrofluorene 3 [5], methylene green 4 [5], and 1,3,5,7,8-pentamethyl-2,6-dinitropyrromethene-BF<sub>2</sub> complex 5 [6]. To accommodate fluorescence and laser activity in the stilbene 1 and the absence of luminescence in 4-nitrostilbene a contribution from electronic interaction between the nitro and amino substituents was proposed [2-4].




A reexamination of the qualitative observation of strong fluorescence in solution for a dinitro derivative of 5,11-dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole 6 [7] was undertaken. The unsubstituted dibenzotetraazapentalene 6, a hybrid of mesoionic structures, was a yellow crystalline solid with a very weak yellow-green fluorescence under ultraviolet light and absorption in the electronic spectrum at  $\lambda_{\max}$  (ethanol) 402 nm,  $\log \epsilon$  4.58 [8]. Its chromophore and luminophor properties were comparable to those reported for isomeric and isosteric dipyrido [1,2-*a*:1',2'-*e*]1,3,4,6-tetraazapentalene 7 with absorption  $\lambda_{\max}$  (methanol) 396 nm,  $\log \epsilon$  4.00, and  $\lambda_f$  (ethanol) 433 nm,  $\Phi$  0.27 [9].



- |    |   |     |  |
|----|---|-----|--|
| 6  | W = X = Y = Z = R = H                   | 9f  | W = Y = NH <sub>2</sub> , X = Z = R = H                  |
| 9a | W = NO <sub>2</sub> , X = Y = Z = R = H | 9g  | unassigned diazidodinitro derivative                     |
| 9b | W = Y = NO <sub>2</sub> , X = Z = R = H | 10a | W = X = Y = Z = H, R = CH <sub>3</sub>                   |
| 9c | W = X = Y = NO <sub>2</sub> , Z = R = H | 10b | W = R = Y = CH <sub>3</sub> , X = Z = H                  |
| 9d | W = X = Y = Z = NO <sub>2</sub> , R = H | 11a | W = Y = NO <sub>2</sub> , X = Z = H, R = CH <sub>3</sub> |
| 9e | W = NH <sub>2</sub> , X = Y = Z = R = H | 11b | W = Y = R = CH <sub>3</sub> , X = Z = NO <sub>2</sub>    |




- 8 A = B = H  
 12 A = CH<sub>3</sub>, B = H  
 13 A = B = CH<sub>3</sub>



- 14 V = NO<sub>2</sub>  
 15 V = CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>

A preparation of the tetraazapentalene 6 from *o*-phenylene diamine 8, eq (1); its nitration to the mono, di, and tetranitro derivatives 9a,b,d; reduction of the mono and dinitro compounds to the corresponding mono and diamino derivatives 9e,f; and replacement of two nitro groups in compound 9d with two azido groups to give an unassigned diazido dinitro derivative 9g were previously reported [7].



## RESULTS AND DISCUSSION

A suitable variation in the nitration of compound 6 [7,8] gave 2,4,8-trinitro-5,11-dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole 9c. The assignment was complementary to the structures of the other nitro derivatives 9a,b,d and supported by <sup>1</sup>H NMR analysis. In straightforward extensions the preparations of 2,8-dinitro-3,9-dimethyl-5,11-dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole 11a and 4,10-dinitro-2,3,8,9-tetramethyl-5,11-dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole 11b were obtained from 3,4-diaminotoluene 12 and 4,5-dimethyl-1,2-phenylenediamine 13 via the inter-

mediate dimethyl and tetramethyldibenzotetraazapentalenes **10a,b** respectively.

Longest wavelength absorption in the electronic spectrum, fluorescence emission, and laser activity data were obtained for twelve examples of tetraazapentalene derivatives **6**, **9a-g**, **11a,b**, **14** and **15**. Limited solubilities afforded measurements on each tetraazapentalene in ethyl acetate, acetonitrile, and acetone. Absorption maxima and extinction coefficients were virtually solvent independent and bathochromic shifts progressed from 403 nm for the unsubstituted compound **6** in increments of about 35, 55, 80, and 90 nm as the number of nitro substituents increased from 1 to 4 in compounds **9a-d** (Table). The effect of dinitro substitution was independent of position assignment insofar as the 2,8-dinitro derivatives **9b** and **11a** and the 4,10-dinitro derivative **11b** absorbed at 450 nm ( $\pm 7$ ),  $\log \epsilon$   $4.43 \pm 0.09$ .

In comparison with the weak luminophor in dibenzotetraazapentalene **6** fluorescence intensification was brought about by nitro substitution in the derivatives **9a-d** and **11a,b** (Table). Although the quantum yields were erratically solvent dependent dibenzotetraazapentalene **6** and its nitro derivatives **9a-c** and **11a,b** gave highest values in ethyl acetate. The tetranitro derivative **9d** was exceptional with weak fluorescence in ethyl acetate and strong fluorescence in acetonitrile and in acetone (Table). The highest fluorescent quantum yield on record for a nitro compound was observed for 2,8-dinitro-5,11-dehydro-5*H*,11*H*-benzotriazol[2,1-*a*]benzotriazole **9b**,  $\lambda_f$  (ethyl acetate) 479 nm,  $\Phi$  0.98. This high value was not sustained by either the structurally similar 2,8-dinitro-3,9-dimethyl derivative **11a**,  $\Phi$  0.46 (ethyl acetate), the 2,3,8,9-tetramethyl-4,10-dinitro derivative **11b**,  $\Phi$  0.15 (ethyl acetate), or the unassigned diazido-dinitro derivative **9g** in which fluorescence was not detected. Reduction of nitro


groups also led to the disappearance of fluorescence in the monoamino and diamino derivatives **9e**, **9f**. Fluorescence was not detected in either 2,5-dimethyl-3,6-dinitro-1,3a,4,6a-tetraazapentalene **14** or ethyl 2,5-dimethyl-1,3a,4,6a-tetraazapentalene-3,6-dicarboxylate **15**. The simpler tetraazapentalene chromophore in compounds **14** and **15** was shifted to  $\lambda_{\text{max}}$  near 380 and 310 nm.

Laser activity was shown by four nitro derivatives **9a–d** of the dibenzotetraazapentalene **6** but was not detected in the dinitro derivatives **11a,b**. In general the relative efficiency (RE) in power output of a laser dye depended on several factors including a high extinction coefficient of absorption, a high fluorescence quantum yield and minimal overlap of absorption (singlet-singlet and triplet-triplet) in the fluorescence spectral region. These were presumably operative conditions for the nitro derivatives **9a–d** to show laser activity over the range 530 to 590 nm (Table). Enigmatic results showed the dinitro derivative **9b** with  $\Phi$  0.98 in ethyl acetate to be laser inactive and with  $\Phi$  0.58 in acetonitrile or  $\Phi$  0.61 in acetone to be laser active. A low relative efficiency, RE 30 (on a scale where RE 100 was arbitrarily assigned to laser dye rhodamine 6G), was determined for the dye **9b** in acetonitrile [10]. Photoinstability accounted for its decomposition under flash lamp excitation and precluded an accurate measurement of triplet-triplet absorption [10].

Electronic interaction between conjugated nitro and amino groups was a common feature of laser dyes **1–5** and presumably diminished the role of the nitro group as a quencher of luminescence. By extension the laser activity in each dye **9a–d** can tentatively be partially attributed to a charge separated luminophor brought about by electronic interaction between a nitro group in conjugation with an electron rich meso-



ionic nitrogen atom as illustrated for the 2,8-dinitro compound **9b**, eq (2). A similar charge separated luminophor failed to bring about laser activity in the 2,8- and 4,10-dinitrodibenzotetraazapentalenes **11a,b** and in 2,5-dimethyl-3,6-dinitro-1,3a,4,6a-tetraazapentalene **14**.



### EXPERIMENTAL

Instruments for spectroscopic measurements included: Perkin-Elmer 1600 FTIR, Varian Gemini 300 NMR, Hewlett-Packard 5985 (70 eV) GC-MS, Cary 17 (UV), Perkin-Elmer LS-5B Luminescence spectrometer, and a Phase-R DL-1100 dye laser with a DL-5Y coaxial flashlamp. Literature procedures were followed to prepare 5,11-dehydro-5*H*,11*H*-benzo-triazolo[2,1-*a*]benzotriazole **6**, its mononitro, dinitro, and tetra-nitro derivatives **9a,b,d**, its 2-amino- and 2,8-diamino derivatives **9e,f** [7,8], 2,5-dimethyl-3,6-dinitro-1,3a,4,6a-tetraazapentalene **14**, and ethyl 2,5-dimethyl 1,3a,4,6a-tetraazapentalene-3,6-dicarboxylate **15** [11,12]. Light absorption, luminescence, and laser activity for the dyes **6**, **9a-d**, and **11a,b**, are described in the Table.

Each recorded UV absorption was restricted to the highest wavelength.  $^1\text{H}$  NMR spectra were run in  $\text{CDCl}_3$  with tetramethylsilane as an internal standard. Fluor-

rescence quantum yields of the dyes were determined for solutions in ethyl acetate, acetonitrile, and acetone with excitation at 450 and 460 nm by reference to acridine orange,  $\Phi$  0.46 [13]. Melting points were determined on a Thomas Hoover melting point apparatus and were uncorrected. Elemental analyses were obtained from Midwest Micro Lab, Indianapolis, Ind., and Galbraith Laboratories, Inc., Knoxville, Tenn.

*2,4,8-Trinitro-5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole 9c*


The compound 5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole **6** (450 mg, 1.5 mmol) was added in small portions to concentrated nitric acid (15 ml) at 5 °C with stirring. After stirring was continued for 2 h the mixture was poured into ice-water.

A precipitate was isolated and recrystallized from dimethylformamide to give 2,4,8-trinitro-5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole **9c** as a yellow crystalline solid (460 mg, 62%), mp 282–283 °C;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  8.24 (d,  $J_{10,9}$  9.6 Hz, H 10), 8.66 (dd,  $J_{9,10}$  9.3 Hz,  $J_{9,7}$  2.1 Hz, H 9), 8.77 (s, H 1), 9.44 (s, H 3), 9.54 (d,  $J_{7,9}$  2.1 Hz, H 7); IR (KBr):  $\nu$  3094, 1617, 1523, 1351, 1325, 1285, 1159, 1108, 819; EI-MS ( $m/z$ ) (%): 343 (30, M), 75 (31), 30 (100). Anal. calcd for  $\text{C}_{12}\text{H}_5\text{N}_7\text{O}_6$ : C, 41.96; H, 1.46; N, 28.55; found: C, 41.73; H, 1.61; N, 28.21.

*General procedure for the preparation of tetraazapentalenes 10a,b*

A straightforward adaptation of the preparation of the dibenzotetraazapentalene **6** from *o*-phenylenediamine **8** was followed. Lead dioxide (1.1 mol) was added to a stirred solution of an *o*-phenylenediamine (0.5 mol) in benzene (1.5 L). After 1 h at 25 °C the mixture was kept at 80 °C for 3 h. After insoluble lead salts were removed and the deep red mixture was filtered through silica gel (300 g, 230–400 mesh, 60 Å, di-

chloromethane) and concentrated, recrystallization of the residue from toluene gave a diaminoazobenzene. The 4,4'-dimethyl-2,2'-diaminoazobenzene **18** (37%) was obtained as an orange crystalline solid, mp 142–143 °C (lit. [14], 120 °C); IR, MS, and  $^1\text{H}$  NMR data were in agreement with reported values [14]. Anal. calcd for  $\text{C}_{14}\text{H}_{16}\text{N}_4$ : C, 69.79; H, 6.68; N, 22.90; found: C, 69.97; H, 6.17; N, 23.31. The 4,4',5,5'-tetramethyl-2,2'-diaminoazobenzene **19** (34%) was obtained as red leaf crystals, mp 203–204 °C (lit. [14], 205 °C); IR, MS, and  $^1\text{H}$  NMR data were in agreement with reported values [14].



**18** A =  $\text{CH}_3$ , B = H, D =  $\text{NH}_2$

**20** A =  $\text{CH}_3$ , B = H, D =  $\text{N}_3$

**19** A = B =  $\text{CH}_3$ , D =  $\text{NH}_2$

**21** A = B =  $\text{CH}_3$ , D =  $\text{N}_3$

An extension of the conversion of the diamine **16** to the diazide **17** afforded the diazides **20** and **21** from the diamines **18** and **19** respectively. The 4,4'-dimethyl-2,2'-diazidoazobenzene **20** (89%) was obtained as orange needles (toluene), mp 122 °C (explosive dec.);  $^1\text{H}$  NMR:  $\delta$  2.39 (s, 2  $\text{CH}_3$ ), 6.96 (s, H 3 / H 3'), 6.99, 7.63 (d,  $J$  8.4 Hz, H 5, H 6 / H 5', H 6'); IR (KBr):  $\nu$  2117 ( $\text{N}_3$ ), 1603, 1284; EI-MS ( $m/z$ ) (%): 292 (4, M), 236 (78), 192 (19), 39 (100). Anal. calcd for  $\text{C}_{14}\text{H}_{12}\text{N}_8$ : C, 57.53; H, 4.11; N, 38.36; found: C, 57.68; H 4.03; N, 38.11. The 4,4',5,5'-tetramethyl-2,2'-diazidoazobenzene **21** (63%) was obtained as orange needles (toluene) mp 118 °C (explosive dec.);  $^1\text{H}$  NMR:  $\delta$  2.25 (s, 2  $\text{CH}_3$ ), 2.29 (s, 2  $\text{CH}_3$ ), 6.92 (s, 2 H), 7.51 (s, 2 H); IR (KBr):  $\nu$  2108 ( $\text{N}_3$ ), 1443, 1378, 1212, 991, 838. Anal. calcd for  $\text{C}_{16}\text{H}_{16}\text{N}_8$ : C, 59.99; H, 5.03; N, 34.98; found: C, 59.89; H, 4.92; N, 34.77.

In an extension of the conversion of the diazide **17** to dibenzotetraazapentalene

6 the diazides **20** and **21** gave the tetraazapentalene derivatives **10a,b**. The 3,9-dimethyl-5,11-dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole **10a** (99%) was obtained as yellow needles (toluene) mp 197–198 °C; <sup>1</sup>H NMR: δ 2.60 (s, CH<sub>3</sub>), 2.61 (s, CH<sub>3</sub>), 7.21 (dd, *J*<sub>8,7</sub> 8.7 Hz, *J*<sub>8,10</sub> 4.5 Hz, H 8), 7.44 (d, *J*<sub>7,8</sub> 9.0 Hz, H 7), 7.66 (bs, H 10), 7.80 (d, *J*<sub>1,2</sub> 8.4 Hz, H 1), 7.93 (s, H 4), 8.03 (d, *J*<sub>2,1</sub> 8.4 Hz, H 2); IR (KBr): ν 1616, 1496, 1370, 1340, 1102, 798; UV (CH<sub>3</sub>CN) λ<sub>max</sub> 405 (4.54), 387 (4.35), 258 (4.86); EI-MS (*m/z*) (%): 236 (100, M). Anal. calcd for C<sub>14</sub>H<sub>12</sub>N<sub>4</sub>: C, 71.17; H, 5.12; N, 23.71; found: C, 71.17; H, 5.06; N, 23.52. The 2,3,8,9-tetramethyl-5,11-dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole **10b** (77%) was obtained as a yellow powder (toluene), mp 280 °C (dec); <sup>1</sup>H NMR: δ 2.28 (s, 2 CH<sub>3</sub>), 2.46 (s, 2 CH<sub>3</sub>), 7.69 (s, 2 H); IR (KBr): ν 1624, 1595, 1449, 1354, 1271, 999; EI-MS (*m/z*) (%) 264 (47, M). Anal. calcd for C<sub>16</sub>H<sub>16</sub>N<sub>4</sub>: C, 72.73; H, 6.06; N, 21.21; found: C, 72.61; H, 6.04; N, 20.94.

*2,8-dinitro-3,9-dimethyl-5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole 11a*

Nitric Acid (25%, 50 ml) at 0–5 °C was added to 3,9-dimethyl-5,11-dehydro-5*H*,11*H*-benzotriazolo[2,1-*a*]benzotriazole **10a** and after 1.5 h, the temperature was allowed to rise to room temperature and stored for 4 h. An orange-red solid was isolated, washed with water, and dried. Purification by column chromatography (silica gel, dichloromethane/light petroleum ether, 1:1) and recrystallization from dimethylformamide and toluene afforded the dinitro derivative **11a** (*R<sub>f</sub>* 0.42) as an orange crystalline solid, mp 260 °C (dec); <sup>1</sup>H NMR: δ 2.90 (s, CH<sub>3</sub>), 7.91 (s, H 4 and H 10), 9.04 (s, H 1 and H 7); IR (KBr): ν 1613, 1530, 1331, 1116, 846, 777, 752; EI-MS (*m/z*) (%): 326 (100,

M). Anal. calcd for  $C_{14}H_{10}N_6O_4$ : C, 51.53; H, 3.07; N, 25.77; found: C, 51.48; H, 3.15; N, 25.53.

*4,10-Dinitro-2,3,8,9-tetramethyl-5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole 11b*

To a stirred solution of 2,3,8,9-tetramethyl-5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole **10b** (528 mg, 2.0 mmol) in acetonitrile (10 ml) nitronium tetrafluoroborate (531 mg, 4.0 mmol) in acetonitrile (15 ml) was slowly added at 0 °C. The mixture was held at 0 °C for 1 h. A yellow precipitate was isolated and recrystallized from toluene to give the dinitro derivative **11b** as a yellow crystalline solid (524 mg, 74%), mp 310 °C (dec);  $^1H$  NMR:  $\delta$  2.59 (s,  $CH_3$ ), 2.60 (s,  $CH_3$ ), 8.22 (s, H 1 and H 7); IR (KBr):  $\nu$  1625, 1523, 1359, 1321, 1094, 1027, 865, 813, 769; EI-MS ( $m/z$ ) (%): 354 (100, M). Anal. calcd for  $C_{16}H_{14}N_6O_4$ : C, 54.24; H, 3.95; N, 23.73; found: C, 54.04; H, 3.81; N, 23.60.

Spectroscopic data were obtained for 2-amino- and 2,8-diamino-5,11-dehydro-5H,11H-benzotriazolo[2,1-a]benzotriazole **9e,f**, 2,5-dimethyl-3,6-dinitro-1,3a,4,6a-tetraazapentalene **14**, and ethyl 2,5-dimethyl-1,3a,4,6a-tetraazapentalene-3,6-dicarboxylate **15**: UV absorption [compound, (solvent),  $\lambda_{max}$  nm, log  $\epsilon$ ]: **9e** ( $CH_3CO_2C_2H_5$ ,  $CH_3CN$ , or  $CH_3COCH_3$ ), 436, 4.29; **9f** ( $CH_3CN$  or  $CH_3COCH_3$ ), 477, 4.43; **14** ( $CH_3CO_2C_2H_5$ ), 382, 4.37; **14** ( $CH_3CN$ ), 384, 4.50; **14** ( $CH_3COCH_3$ ), 384, 4.61; **15** ( $CH_3CO_2C_2H_5$ ), 310, 4.34; **15** ( $CH_3CN$ ), 310, 4.61; **15**, 310, 4.35. The diamine **9f** was insoluble in ethyl acetate. Luminescence was not detected in solutions of compounds **9e,f**, **14** and **15**.

### ACKNOWLEDGMENTS

Financial support was received from ONR, ARO, and the Louisiana Board of Regents (LEQSF-RD-B-06 and RD-B-15).

### REFERENCES

- [1] B. M. Krasovitskii and B. M. Bolotin, "Organic Luminescent Materials," Engl. transl. by V. G. Vopian, VCH, Weinheim, 1988, p 17.
- [2] (a) P. P. Shorygin and T. M. Ivanova, *Doklady Akad. Nauk S.S.S.R.*, 121, 1958, 70; *Chem. Abstr.*, 54, 1960, 23785c. (b) Ref 1, pp 44, 215.
- [3] Fluorescence for the stilbene 1,  $\lambda_f$  590 nm,  $\Phi$  0.7 (benzene) [2] did not agree with the more recently reported  $\lambda_f$  600 nm,  $\Phi$  0.085 (benzene);  $\lambda_f$  502 nm,  $\Phi$  0.46 (cyclohexane);  $\lambda_f$  678 nm,  $\Phi$  0.38 (tetrahydrofuran);  $\lambda_f$  772 nm,  $\Phi$  0.0006 (acetonitrile) [4].
- [4] D. M. Shin and D. G. Whitten, *J. Phys. Chem.*, 92, 1988, 2945.
- [5] M. Maeda, "Laser Dyes," Academic Press Inc., New York, 1984, pp 37, 38, 123, 124, 144, 148.
- [6] M. Shah, K. Thangaraj, M.-L. Soong, L. T. Wolford, J. H. Boyer, I. R. Politzer, and T. G. Pavlopoulos, *Het. Chem.*, 1, 1990, 389.
- [7] R. A. Carboni, J. C. Kauer, W. R. Hatchard, and R. J. Harder, *J. Amer. Chem. Soc.*, 89, 1967, 2626.
- [8] R. A. Carboni, J. C. Kauer, J. E. Castle, and H. E. Simmons, *J. Amer. Chem. Soc.*, 89, 1967, 2618.
- [9] M. P. Groziak, S. R. Wilson, G. L. Clauson, and N. J. Leonard, *J. Amer. Chem.*

*Soc.*, 108, 1986, 8002.

- [10] T. G. Pavlopoulos, personal communication.
- [11] R. Pflieger, E. Garthe, and K. Rauer, *Chem. Ber.*, 96, 1963, 1827.
- [12] We are indebted to Dr. A. Natesh for the preparation of compounds 14 and 15.
- [13] A. W. Johnson, I. T. Kay, E. Markham, R. Price, and K. B. Shaw, *J. Chem. Soc.*, 1959, 3416.
- [14] G. Crank and M. I. H. Makin, *Aust. J. Chem.*, 37, 1984, 845.

**Table**  
**Absorption and Luminescence in Dibenzotetraazapentalenes**

No.	Solvent <sup>a</sup>	$\lambda_{\max}^b$	$\log \epsilon$	$\lambda_f^b$	$\Phi_f$	$\lambda_{\text{las}}^c$
<b>6</b>	EA	403	4.68	452	0.02	—
	AN	403	4.68	—	—	—
	A	403	4.85	—	—	—
<b>9a</b>	EA	438	4.40	498	0.53	560
	AN	440	4.42	—	—	—
	A	440	4.42	527	0.18	—
<b>9b</b>	EA	454	4.45	479	0.98	—
	AN	457	4.52	501	0.58	575
	A	457	4.52	494	0.61	562
<b>9c</b>	EA	476	4.50	509	0.61	575
	AN	482	4.39	516	0.27	—
	A	483	4.44	516	0.36	—
<b>9d</b>	EA	497	4.56	512	0.08	—
	AN	493	4.60	509	0.81	560
	A	493	4.71	511	0.66	—
<b>9g</b>	EA	463	4.32	—	—	—
	AN	464	4.32	—	—	—
	A	464	4.37	—	—	—
<b>11a</b>	EA	447	4.34	480	0.46	—
	AN	446	4.36	—	—	—
	A	448	4.42	495	0.10	—
<b>11b</b>	EA	443	4.47	517	0.15	—
	AN	447	4.30	—	—	—
	A	441	4.49	—	—	—

<sup>a</sup> Ethyl acetate (EA); acetonitrile (AN); acetone (A). <sup>b</sup>  $6 \times 10^{-6}$  M solutions.

<sup>c</sup>  $2 \times 10^{-4}$  M solutions.