

AD-A254 737



2

TECHNICAL REPORT BRL-TR-3394

**BRL**

FIXED POINTS OF  
EXPANSIVE ANALYTIC MAPS (II)

WALTER O. EGERLAND  
CHARLES E. HANSEN

SEPTEMBER 1992

DTIC  
SELECTE  
SEP 02 1992  
S B D

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.

U.S. ARMY LABORATORY COMMAND

BALLISTIC RESEARCH LABORATORY  
ABERDEEN PROVING GROUND, MARYLAND

92 8 31 058

92-24148



050750

23p4

## NOTICES

Destroy this report when it is no longer needed. DO NOT return it to the originator.

Additional copies of this report may be obtained from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The use of trade names or manufacturers' names in this report does not constitute indorsement of any commercial product.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> September 1992	<b>3. REPORT TYPE AND DATES COVERED</b> Final, October 1990-June 1992	
<b>4. TITLE AND SUBTITLE</b> Fixed Points of Expansive Analytic Maps (II)			<b>5. FUNDING NUMBERS</b> PR: 1L162618AH43	
<b>6. AUTHOR(S)</b> Walter O. Egerland and Charles E. Hansen				
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Ballistic Research Laboratory ATTN: SLCBR-DD-T Aberdeen Proving Ground, MD 21005-5066			<b>10. SPONSORING/MONITORING AGENCY REPORT NUMBER</b>  BRL-TR-3394	
<b>11. SUPPLEMENTARY NOTES</b>				
<b>12a. DISTRIBUTION/AVAILABILITY STATEMENT</b> Approved for public release; distribution is unlimited			<b>12b. DISTRIBUTION CODE</b>	
<b>13. ABSTRACT (Maximum 200 words)</b>  The main result of this note, the "Encircling Theorem," states conditions that assure that an analytic function is expansive in the closed unit disk $D$ and has a fixed point in $D$ . A corollary describes in detail the case of a conformal map. From a new covering lemma for polynomials further sufficient conditions are deduced that guarantee that a polynomial of degree $n$ , $n = 1, 2, \dots$ , is expansive and has a fixed point in $D$ . On the other hand, an important example shows that for each $n \geq 3$ polynomials of degree $n$ exist that cover $D$ but do not have a fixed point in $D$ . Finally, the distribution of the fixed points of any finite Blaschke - product is established.				
<b>14. SUBJECT TERMS</b> Fixed points; expansive analytic maps; conformal maps; polynomials; Blaschke - products; conformal mapping			<b>15. NUMBER OF PAGES</b> 25	
			<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> UNCLASSIFIED	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> UNCLASSIFIED	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> UNCLASSIFIED	<b>20. LIMITATION OF ABSTRACT</b> UNCLASSIFIED	

**INTENTIONALLY LEFT BLANK.**

**TABLE OF CONTENTS**

	<u>Page</u>
LIST OF FIGURES .....	v
1. INTRODUCTION .....	1
2. FEJÉR'S REMARK .....	3
3. THE "ENCIRCLING" THEOREM .....	4
4. A COVERING LEMMA .....	6
REFERENCES .....	10
APPENDIX 1 .....	11
APPENDIX 2 .....	13
DISTRIBUTION LIST .....	15

**DTIC QUALITY INSPECTED 3**

<b>Accession For</b>	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution/ _____	
<b>Availability Codes</b>	
<b>Dist</b>	<b>Avail and/or Special</b>
A-1	

**INTENTIONALLY LEFT BLANK.**

## LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1.	$p$ covers $D$ .....	1
2.	$q$ covers $K$ .....	2
3.	$z + (z + 1.1i)^3$ .....	5
4.	The location of the zeros of $G_n$ .....	8

**INTENTIONALLY LEFT BLANK.**



## 1. INTRODUCTION.

In BRL-TR-3063, Fixed Points Of Expansive Analytic Maps, Nov. 1989 [7], the notion of an “expansive analytic map” was introduced. Let  $D$  be the closed unit disk in the complex plane and  $f$  an analytic function in  $D$ . By definition,  $f$  is “expansive” in  $D$  if the image  $f(D)$  satisfies the condition  $f(D) \supseteq D$ . Equivalently, we say that  $f$  “covers”  $D$  whenever  $f(D) \supseteq D$  holds. The principal result in BRL-TR-3063 states that a quadratic complex polynomial that covers  $D$  has a fixed point in  $D$ . More generally, it was shown that the result holds for any closed disk  $K$ , i.e.,  $f(K) \supseteq K$  implies that there exists  $z_0 \in K$  such that  $f(z_0) = z_0$ . Figure 1. and Figure 2. illustrate the principal result for  $D$  and  $K: |z-1| \leq 1$ . The quadratic polynomials chosen are  $p(z) = z + ie^{-i\frac{3}{8}\pi}(z - e^{i\frac{\pi}{4}})(z + 1)$  and  $q(z) = z^2 - \frac{3}{2}z + \frac{3}{2}$ , respectively.




Figure 1.  $p$  covers  $D$

The purpose of this note is to give further results on expansive maps. The main result, given in Section 3., is the “Encircling” Theorem. It states that if for an analytic function  $f$  in  $D$  (not necessarily a polynomial) it is known that it maps an interior point of  $D$  into the open unit disk and the image of the unit circle “encircles” the unit circle (“ $f$  pushes the unit circle out”), then  $f$  is expansive and has a fixed point in  $D$ . For example, the function of Figure 1. has this “encircling” property, whereas the function of Figure 2. does not have this property. As a corollary to the “encircling” theorem we

obtain a proof of an important theorem for univalent expansive maps that is independent of Brouwer's fixed point theorem for the plane and more informative than a weaker version obtainable from this theorem. The proof of the "encircling" theorem uses at key junctures a pertinent but not widely publicized remark of Fejér regarding Rouché's theorem. In Section 2. we state Fejér's addendum to Rouché's theorem. It is a consequence of Fejér's remark that a finite Blaschke-Product has a fixed point on the unit circle. We proved that if the number of factors is  $n$ , then at least  $n - 1$  fixed points are on the unit circle. This result was proposed to the problem section of the American Mathematical Monthly and appeared as Problem 6654 on page 273 of the March 1991 issue (Vol. 98, Number 3) of this journal. Our solution, as submitted to the editors, is reproduced in Appendix 1. In Section 4. we prove a covering lemma for polynomials and derive a sufficient condition for a polynomial of degree  $n$  to cover  $D$  and have a fixed point in  $D$ . We also define for each  $n \geq 3$  a polynomial  $g_n$  that covers  $D$  but does not have a fixed point in  $D$ . This example shows that the principal result of BRL-TR-3063 cannot be generalized to polynomials of degree  $n \geq 3$  without additional assumptions. Finally, in Appendix 2, we give a proof of Lemma F that was left to the reader in BRL-TR-3063. This particular proof keeps the necessary algebraic manipulations to a minimum.




Figure 2.  $q$  covers  $K$

## 2. FEJÉR'S REMARK.

What we call "Fejér's Remark" is explicitly attributed to him by Lipka [1,p.143]. It is also recorded in [2,p.5,Exercise 7]. If we introduce the notation  $C = \{ z \mid |z| = 1 \}$  for the unit circle and  $U = \{ z \mid |z| < 1 \}$  for the open unit disk, "Fejér's Remark" is the following statement.

**Fejér's Remark.** Let  $F$  and  $G$  be analytic in  $D$ . If for  $z \in C$

- (i)  $F(z) \neq 0$
- (ii)  $F(z) + G(z) \neq 0$
- (iii)  $|F(z)| \geq |G(z)|$ ,

then  $F$  and  $F + G$  have the same number of zeros in  $U$ .

For the proof, see [1,p.143].

As a first application, we observe that a "Blaschke-factor"

$$B_n(z) = e^{i\omega} \prod_{k=1}^n \frac{z - a_k}{1 - \bar{a}_k z}, \quad |a_k| < 1, \quad k = 1, 2, \dots, n, \quad n \geq 2$$

has a fixed point on the unit circle  $|z| = 1$ .

Suppose this were false. Since a Blaschke-factor is expansive as well as non-expansive, the inequality  $|B_n(z)| \geq 1 = |-z|$  for  $z \in C$  would imply that  $B_n$  has at least two fixed points in  $U$ , while the inequality  $|z| \geq 1 = |-B_n(z)|$  for  $z \in C$  would imply that  $B_n$  has exactly one fixed point in  $U$ . This contradiction proves the assertion. A more detailed analysis shows that  $B_n$  has at least  $n - 1$  fixed points on the unit circle (see Problem 6654 in the March 1991 issue of the American Mathematical Monthly and Appendix 1. of this note).

As a second application, we remark that for analytic functions the following stronger form of Brouwer's fixed point theorem for the disk holds: If  $f$  is analytic in  $D$  and  $f(D) \subseteq D$ , then  $f$  has a fixed point in  $D$ . If  $f$  does not have a fixed point on the unit circle, then  $f$  has a unique fixed point in the open unit disk. For the proof, we apply Fejér's remark to the functions  $F(z) = -z$  and  $G(z) = f(z)$ .

### 3. THE "ENCIRCLING" THEOREM.

**Theorem ("encircling" theorem).** Let  $f$  be analytic in  $D$  and  $|f(a)| < 1$  for some  $|a| < 1$ . If  $|f(C)| \geq 1$ , where  $|f(C)| := |f(e^{i\phi})|$ ,  $0 \leq \phi < 2\pi$ , then

- (i)  $f(D) \supseteq D$ .
- (ii)  $f$  has a fixed point in  $D$ .
- (iii) If  $f$  does not have a fixed point on  $C$ ,  $f$  has as many fixed points in  $U$  as  $f$  has zeros in  $U$ .

**Proof.** Since  $|f(C)| \geq 1$ , the inequality  $|f(z)| \geq 1 > |-f(a)|$  holds for  $z \in C$ . Therefore, by Rouché's theorem,  $f$  has a zero in  $U$ . If  $u_0 \in U$ , it follows from  $|f(z)| \geq 1 > |-u_0|$  for  $z \in C$ , again by Rouché's theorem, that  $f$  covers  $U$  since  $u_0$  is arbitrary. If  $u_0 \in C$ , either  $f(e^{i\phi_0}) = u_0$  for some  $\phi_0$  or, by Fejér's remark,  $f(z_0) = u_0$  for some  $z_0 \in U$ . This proves (i). Suppose (ii) were false, i.e., assume it were possible that  $f(z) \neq z$  for  $z \in D$ . Then, in particular, we would have  $f(z) - z \neq 0$  for  $z \in C$  and the inequality  $|f(z)| \geq 1 = |-z|$  would imply, by Fejér's remark (set  $F(z) = f(z)$ ,  $G(z) = -z$ ), that  $f(z)$  and  $f(z) - z$  have the same number of zeros in  $U$ . Since  $f$  has at least one zero in  $U$  by (i),  $f(z) - z = 0$  for some  $z_0 \in U$ , i.e.,  $f$  has at least one fixed point in  $U$ . This contradiction proves (ii) and (iii) is immediate. This completes the proof.

**Corollary.** Let  $f$  be analytic and univalent in  $D$ . If  $f$  is not a conformal automorphism of  $U$  and  $f(D) \supseteq D$ , then

- (i)  $f$  has a fixed point in  $D$ .
- (ii) If  $f$  does not have a fixed point on the unit circle  $C$ ,  $f$  has a unique fixed point  $z_0 \in U$  such that  $|f'(z_0)| > 1$ , i.e.,  $z_0$  is a repeller.
- (iii) If  $f(a) = 0$ ,  $a \neq 0$ , and (ii) occurs, the repeller  $z_0$  lies in the "annular sector"  $A$ , defined by

$$A = \left\{ z \left| \begin{array}{l} |a| \left( 1 + \sqrt{1 - |a|^2} \right)^{-1} < |z| < 1 \\ \cos(\arg(z/a)) > |a| \end{array} \right. \right\}.$$

**Proof.** Clearly  $|f(C)| \geq 1$ . Since  $f$  covers  $D$  and is univalent, there exists a unique  $a \in U$  such that  $f(a) = 0$ . Therefore, (i) follows by the “encircling” theorem. If  $f$  does not have a fixed point on  $C$ , it follows from (iii) of the “encircling” theorem that  $f$  has a unique fixed point  $z_0 \in U$  because it has the unique zero  $a \in U$ . Since  $f(D) \supseteq D$  implies  $f^{-1}(D) \subseteq D$ , the inverse  $g = f^{-1}$  of  $f$  is bounded by 1 and  $g(z_0) = z_0$ . The Schwarz-Pick lemma [4,5] applies to  $g$  and  $g'$  satisfies the strict inequality

$$(1 - |z|^2)|g'(z)| < 1 - |g(z)|^2 \quad (z \in U)$$

because  $g$  is not a conformal automorphism of  $U$ . Setting  $z = z_0$ ,  $|g'(z_0)| < 1$ , and, hence,  $|f'(z_0)| > 1$  follows. This proves (ii). To prove (iii), we note that by Schwarz’s lemma  $z_0$  is subject to the inequality  $|z_0 - a| < |z_0||1 - \bar{a}z_0|$ . Hence,  $z_0 \in A_s$ . This completes the proof of the Corollary.

**Remarks.**

1. If an analytic function in  $D$  covers  $D$  and does not have a fixed point in  $D$ , then necessarily  $f(C) \cap U \neq \emptyset$ . This intersection property may be observed for the function of Figure 3., where the closed curve shown is the image of  $C$  under  $f$ .
2.  $p(z) = z + (1 + z)^3$  illustrates (i) and  $p(-z)$  illustrates (ii) and (iii) of the Corollary.
3. The proof of the Corollary applies verbatim to analytic  $f$  (not necessarily univalent) for which  $f(D) \subseteq D$  with an attractor and  $f(0)$  taking the place of the repeller and  $a$ . This strengthens, in particular, the perennial problem 197[6,p.142].




Figure 3.  $z + (z + 1.1i)^3$

#### 4. A COVERING LEMMA.

The following covering lemma for polynomials is motivated by the elementary observation that every polynomial  $p(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0$  such that  $|a_n| \geq 1 + |a_0|$  covers  $D$  and has a fixed point in  $D$ .

**Lemma.** Let  $p_n(z) = a_1 z + \dots + a_{n-1} z^{n-1} + a_n z^n$ ,  $a_n \neq 0$ , be a polynomial of degree  $n \geq 3$  and such that  $|a_1| + |a_{n-1}| > 0$ . Then, for some  $R \geq R_0 \geq |a_n|$ ,  $p_n(D) \supseteq D_R = \{z \mid |z| \leq R\}$ . The radius  $R_0$ , depending only on  $|a_1|$ ,  $|a_{n-1}|$ , and  $|a_n|$ , is given by

$$R_0 = \begin{cases} \frac{1}{2} \left( |a_{n-1}| + \sqrt{(|a_{n-1}| - 2|a_n|)^2 + 4(|a_{n-1}| - |a_1|)|a_n|} \right) & (1) \\ \quad \text{if } |a_{n-1}| > |a_1| \geq 0 \\ |a_n| & \text{if } |a_1| = |a_{n-1}| \leq 2|a_n| & (2) \\ |a_{n-1}| - |a_n| & \text{if } |a_1| = |a_{n-1}| > 2|a_n| & (3) \\ \frac{1}{2} \left( -|a_{n-1}| + \sqrt{(|a_{n-1}| + 2|a_n|)^2 + 4(|a_1| - |a_{n-1}|)|a_n|} \right) & (4) \\ \quad \text{if } |a_1| > |a_{n-1}| \geq 0. \end{cases}$$

If  $p_n$  is such that the inequalities

$$|a_1| > |a_{n-1}| > 2|a_n| \text{ and } (2|a_n| - |a_1|)^2 \geq |a_1|^2 - |a_{n-1}|^2 \quad (4')$$

both hold,  $p_n$  covers an annulus  $A(R_1, R_2) = \{z \mid R_1 \leq |z| \leq R_2\}$  in addition to the disk  $D_{R_0}$ , i.e.,  $p_n(D) \supseteq D_{R_0}$  and  $p_n(D) \supseteq A(R_1, R_2)$ . The radii  $R_1 > R_0$  and  $R_2 \geq R_1$  are given by the formula

$$R_{1,2} = \frac{1}{2} \left( |a_{n-1}| \mp \sqrt{(2|a_n| - |a_1|)^2 + |a_{n-1}|^2 - |a_1|^2} \right).$$

**Proof:** Using the notation of [3, §6.8, pp.491-496], it follows from the second Schur transform  $T^2 p$  of the polynomial  $p(z) = p_n(z) + a_0$  for some  $a_0 \neq 0$  that

$$\gamma_2 = (|a_0|^2 - |a_n|^2)^2 - |\overline{a_0} a_{n-1} - \overline{a_1} a_n|^2.$$

Hence,

$$\gamma_2 \leq \gamma_2^* = (|a_0|^2 - |a_n|^2)^2 - (|a_0||a_{n-1}| - |a_1||a_n|)^2.$$

$\gamma_2^*$  is a polynomial of degree four in  $|a_0|$  and satisfies  $\gamma_2^* \leq 0$  in the interval  $|a_n| \leq |a_0| \leq R_0$ . Hence, by Theorem 6.8b [3,p.493], the equation  $p_n(z) + a_0 = 0$  has a root in  $D$  for every  $a_0$  whose modulus falls into this interval, i.e.,  $p_n$  covers  $D_{R_0}$ . It is clear that there exist polynomials for which the inequalities (4') both hold. For such polynomials  $\gamma_2^* \leq 0$  in the disjoint intervals  $|a_n| \leq |a_0| \leq R_0$  and  $R_1 \leq |a_0| \leq R_2$ , where  $R_1 > R_0$ . Therefore, for any  $a_0$  whose modulus lies in one of these intervals, the equation  $p_n(z) + a_0 = 0$  has a root in  $D$  by Theorem 6.8b, i.e.,  $p_n$  covers the disk  $D_{R_0}$  and the annulus  $A(R_1, R_2)$ . This completes the proof of the lemma.

#### Remarks.

1. Similar propositions may be stated for "gap"-polynomials of the form  $p_n = a_l z^l + \dots + a_{n-2} z^{n-2} + a_n z^n$ , where  $a_n \neq 0$ ,  $a_l \neq 0$ ,  $l > 1$ , and  $|a_k| \geq 0$ ,  $k = l+1, \dots, n-2$ .
2. For quadratic polynomials  $p_2(z) = a_1 z + a_2 z^2$ ,  $a_1 \neq 0$ ,  $R_0$  is given by  $R_0 = |a_2|$  if  $|a_1| \leq 2|a_2|$  and  $R_0 = |a_1| - |a_2|$  if  $|a_1| > 2|a_2|$ . This result is sharp.
3.  $R_0 > |a_n|$  if cases (1), (3) or (4) occur.
4. The lower bound  $R_0$  is independent of the "intervening" coefficients  $a_2, a_3, \dots, a_{n-2}$  and in many cases sharp, e.g.,  $p_4(z) = z^4 + 3z^3 + 3z^2 + 3z$  covers  $D_{R_0}$  with  $R_0 = 2$ , but there is no  $R > 2$  such that  $p_4(D) \supseteq D_R$ .
5. Every polynomial  $p(z) = p_n(z) + a_0$ , where  $p_n$  satisfies the conditions of the lemma and  $a_0$  is a given constant, covers  $D$  for which  $R_0 \geq 1 + |a_0|$ . Since for the polynomial in Figure 3. this inequality holds, this remark provides an analytical proof of the "geometric evidence" that this polynomial covers  $D$ .
6. The lemma may also be invoked to show the existence of fixed points under certain circumstances. For example, let  $p(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0$  be a given polynomial and set  $\tilde{p}_n(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + (a_1 - 1)z$ . Then  $p(z) - z = \tilde{p}_n + a_0$ . If  $|a_{n-1}| + |a_1 - 1| > 0$ , the lemma applies to  $\tilde{p}_n$  and determines an  $\tilde{R}_0$ . If  $\tilde{R}_0 \geq |a_0|$ ,  $p$  has a fixed point in  $D$ .
7. For practical applications we have the following proposition.  
**Proposition.** Let  $p, p_n, \tilde{p}_n$  be defined as in Remarks 5. and 6. Then  $p$  covers  $D$  and has a fixed point in  $D$  if  $R_0 \geq 1 + |a_0|$  and  $\tilde{R}_0 \geq |a_0|$ .

Example:  $p(z) = z^n - 4iz^{n-1} + a_{n-2}z^{n-2} + \dots + a_2z^2 - 3z + (1 + \sqrt{2})e^{i\frac{\pi}{4}}$ .  
 For this example  $R_0 = 2 + \sqrt{2}$  and  $\bar{R}_0 = 3$ . Therefore,  $p$  covers  $D$  and has a fixed point in  $D$  for every  $n \geq 3$  and arbitrary  $a_2, \dots, a_{n-2}$ .  
 8.




Figure 4. The location of the zeros of  $G_n$ .

The following example shows that for each  $n \geq 3$  there are polynomials of degree  $n$  that cover  $D$  but do not have a fixed point in  $D$ . We define the polynomials

$$g_n(z) = z + 3(z - e^{i\alpha}(1 + n^{-4}))^n$$

$$\alpha = \begin{cases} 0 & \text{if } n = 2m, \quad m = 2, 3, \dots \\ \pi/2m & \text{if } n = 2m + 1, \quad m = 1, 2, \dots \end{cases}$$

and prove that they cover  $D$ . Clearly,  $g_n(z) \neq z$  for  $z \in D$ .

The polynomials  $g_3$  and  $g_4$  cover by Remark 5. Since the proof for the odd case is the same as for the even case, we consider only the latter and  $n$  will denote an even integer greater than or equal to 6. It follows first by Rouché's theorem that the zeros of the polynomial  $G_n(z) = g_n(z) - u$  for every  $u \in D$  are contained in the disk  $|z - (1 + n^{-4})| < 1 + n^{-4}$ . The circle  $C(n) = \{z \mid |z - (1 + n^{-4})| = 1 + n^{-4}\}$  with center  $c(n) = 1 + n^{-4}$  intersects the unit circle  $C$  in two points with  $x$ -coordinate  $x_0(n) = (2(1 + n^{-4}))^{-1}$  (see Figure 4.). The derivative  $G'_n$  vanishes on the real axis at the point



$$x_1(n) = 1 + n^{-4} - (3n)^{-1/n-1}.$$

This point  $x_1(n)$  lies to the left of  $x_0(n)$  for  $n \geq 6$ . Hence, by the theorem of Gauss-Lucas, there is at least one zero of  $G_n$  in the unit disk for every  $u \in D$ , i.e.,  $g_n$  covers  $D$ . This completes the proof.

This example shows that the principal result of BRL-TR-3063 cannot be generalized to polynomials of degree  $n \geq 3$  without additional assumptions. No such additional assumptions are known at present that are less stringent than those in the "encircling" theorem, its corollary, and the consequences of the covering lemma.

## REFERENCES

- [1] Lipka, S., Eine Verallgemeinerung des Rouchéschen Satzes, J. Reine Angew. Math. vol. 160(1929), pp. 143-150.
- [2] Marden, M., The Geometry of the Zeros of a Complex Polynomial, American Math. Soc., 1949.
- [3] Henrici, P., Applied and Computational Complex Analysis, vol. 1, John Wiley & Sons, 1974.
- [4] Hille, E., Analytic Function Theory, Ginn and Company, vol. 2, 1962.
- [5] Carathéodory, C., Conformal Representation, Cambridge Tracts No. 28, Cambridge University Press, 1963.
- [6] Pólya, G. and Szegő, G., Problems and Theorems in Analysis, vol. 1, Springer-Verlag, 1972.
- [7] Egerland, W. O. and Hansen, C. E., Fixed Points of Expansive Analytic Maps, Technical Report BRL-TR-3063, Nov. 1989.

## Appendix 1.

In this appendix we give a solution to Problem 6654 that appeared in the American Mathematical Monthly, Vol. 98, Number 3, March 1991, p. 273 as follows:

6654. Proposed by W. O. Egerland and C. E. Hansen, Aberdeen Proving Ground, Aberdeen, MD.

Suppose  $\omega$  is real,  $n$  is a positive integer greater than 1, and  $a_1, a_2, \dots, a_n$  are complex numbers with  $|a_k| < 1$  for  $k = 1, 2, \dots, n$ . Prove that the equation

$$e^{i\omega}(z - a_1)(z - a_2) \cdots (z - a_n) = z(1 - \bar{a}_1 z)(1 - \bar{a}_2 z) \cdots (1 - \bar{a}_n z)$$

has at least  $(n - 1)$  roots on the unit circle.

Solution by the Proposers. The roots of the given equation are the fixed points of the "Blaschke-Product"

$$B_n(z) = e^{i\omega} \prod_{k=1}^n \frac{z - a_k}{1 - \bar{a}_k z}.$$

We consider the cases 1.  $B_n(0) \neq 0$  and 2.  $B_n(0) = 0$  separately.

1. If  $B_n(0) \neq 0$ ,  $B_n$  has  $(n + 1)$  fixed points. Since  $B_n$  is analytic in  $U = \{z \mid |z| < 1\}$  and satisfies  $|B_n(z)| < 1$  for  $z \in U$ ,  $B_n$  belongs to the class of "analytic functions of bound 1 in the open unit disk". It is well-known that such functions can have at most one fixed point, if any, in  $U$ [1,2,3]. Therefore, in particular,  $B_n$  can have at most one fixed point  $z_0 \in U$ . Furthermore, it follows from the lemma of Schwarz-Pick[4] that the multiplicity of  $z_0$  is 1, by which is meant that the multiplicity of  $z_0$  as a zero of the function  $B_n(z) - z$  is 1. According to the lemma, the derivative  $B'_n(z)$  satisfies the inequality

$$(1 - |z|^2)|B'_n(z)| < 1 - |B_n(z)|^2 \quad (z \in U).$$

Setting  $z = z_0$ ,  $|B'_n(z_0)| < 1$ , and hence  $B'_n(z_0) - 1 \neq 0$  follows, i.e.,  $z_0$  is a zero of multiplicity 1 of  $B_n(z) - z$ . Finally, we observe that with the exception of  $z = 0, z = a_k, z = \bar{a}_k^{-1}, k = 1, 2, \dots, n$ ,  $B_n$  satisfies the identity

$$B_n(z)\overline{B_n(\bar{z}^{-1})} = 1,$$

so that if  $z_1$ ,  $|z_1| \neq 1$ , is a fixed point of  $B_n$ , the symmetric point  $\bar{z}_1^{-1}$  to  $z_1$  with respect to the unit circle is also a fixed point of  $B_n$ . A short calculation yields  $B'_n(z_1) = \overline{B'_n(\bar{z}_1^{-1})}$ , i.e., both fixed points have multiplicity 1. Since  $B_n$  can have at most one such pair of symmetric simple fixed points, the remaining  $(n - 1)$  fixed points of  $B_n$  lie necessarily on the unit circle.

2. If  $B_n(0) = 0$ ,  $B_n$  has  $n$  fixed points. These are  $z = 0$  and exactly  $(n - 1)$  points on the unit circle  $|z| = 1$ . This is immediate if  $a_1 = a_2 = \cdots = a_n = 0$ , and if  $a_1 = a_2 = \cdots = a_m = 0$ ,  $m < n$ , it follows from

$$\begin{aligned} B_n(z) - z &= z(e^{i\omega} z^{m-1} \prod_{k=m+1}^n \frac{z - a_k}{1 - \bar{a}_k z} - 1) \\ &= z(b_{n-1}(z) - 1) \end{aligned}$$

since  $|b_{n-1}(z)| \neq 1$  if  $|z| \neq 1$ . This completes the proof.

#### References.

1. Ash, R. B., Complex Variables, Academic Press, New York, 1971. Problem 3, p. 102.
2. Duncan, J., The Elements of Complex Analysis, John Wiley and Sons, New York, 1968. Problem 68, p. 239.
3. Biler, P., Witkowski, A., Problems in Mathematical Analysis, Marcel Dekker, New York, 1990. Problem 7.62, p. 85.
4. Heins, M., Complex Function Theory, Academic Press, New York, 1968, Exercise 7.3, pp. 226-227.

## Appendix 2.

**Lemma F.** Let

$$z^2 + az + b = 0$$

be a quadratic equation with roots  $z_1 = \rho_1 e^{i\alpha}$  and  $z_2 = \rho_2 e^{i\beta}$ .

Then

$$\begin{aligned} P &= (|b|^2 - 1)^2 - |\bar{a}b - a|^2 \\ &= (1 - \rho_1^2)(1 - \rho_2^2)((|b| - \cos(\alpha - \beta))^2 + \sin^2(\alpha - \beta)). \end{aligned}$$

**Proof.** Since  $a = -\rho_1 e^{i\alpha} - \rho_2 e^{i\beta}$  and  $b = \rho_1 \rho_2 e^{i(\alpha+\beta)}$ , we have

$$\bar{a}b - a = (1 - \rho_2^2)\rho_1 e^{i\alpha} + (1 - \rho_1^2)\rho_2 e^{i\beta}$$

and

$$|b|^2 - 1 = \rho_1^2 \rho_2^2 - 1 = (1 - \rho_1^2)(1 - \rho_2^2) - (1 - \rho_1^2) - (1 - \rho_2^2).$$

Hence

$$\begin{aligned} P &= (1 - \rho_1^2)^2(1 - \rho_2^2)^2 - 2(1 - \rho_2^2)(1 - \rho_1^2)^2 - 2(1 - \rho_1^2)(1 - \rho_2^2)^2 \\ &\quad + (1 - \rho_1^2)^2 + (1 - \rho_2^2)^2 + 2(1 - \rho_1^2)(1 - \rho_2^2) \\ &\quad - \rho_2^2(1 - \rho_1^2)^2 - \rho_1^2(1 - \rho_2^2)^2 - 2\rho_1\rho_2(1 - \rho_1^2)(1 - \rho_2^2)\cos(\alpha - \beta) \\ &= (1 - \rho_1^2)(1 - \rho_2^2)(\rho_1^2\rho_2^2 - 1 + 2 - 2\rho_1\rho_2\cos(\alpha - \beta)) \\ &= (1 - \rho_1^2)(1 - \rho_2^2)((|b| - \cos(\alpha - \beta))^2 + \sin^2(\alpha - \beta)). \end{aligned}$$

**Corollary 1.** The roots of the quadratic equation  $z^2 + az + b = 0$  have modulus greater than one if and only if the conditions  $|b| > 1$  and  $P > 0$  are satisfied. They have modulus less than one if and only if the conditions  $|b| < 1$  and  $P > 0$  are satisfied.

**Corollary 2.** A quadratic polynomial with both fixed points on the unit circle covers the unit disk if and only if it has the form

$$f(z) = z + i\lambda e^{-i\frac{\alpha+\beta}{2}}(z - e^{i\alpha})(z - e^{i\beta}), \quad \lambda \neq 0 \text{ and real.}$$

**INTENTIONALLY LEFT BLANK.**

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
2	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22304-6145	1	Commander U.S. Army Tank-Automotive Command ATTN: ASQNC-TAC-DIT (Technical Information Center) Warren, MI 48397-5000
1	Commander U.S. Army Materiel Command ATTN: AMCAM 5001 Eisenhower Ave. Alexandria, VA 22333-0001	1	Director U.S. Army TRADOC Analysis Command ATTN: ATRC-WSR White Sands Missile Range, NM 88002-5502
1	Commander U.S. Army Laboratory Command ATTN: AMSLC-DL 2800 Powder Mill Rd. Adelphi, MD 20783-1145	1	Commandant U.S. Army Field Artillery School ATTN: ATSF-CSI Ft. Sill, OK 73503-5000
2	Commander U.S. Army Armament Research, Development, and Engineering Center ATTN: SMCAR-IMI-I Picatinny Arsenal, NJ 07806-5000	(Class. only) 1	Commandant U.S. Army Infantry School ATTN: ATSH-CD (Security Mgr.) Fort Benning, GA 31905-5660
2	Commander U.S. Army Armament Research, Development, and Engineering Center ATTN: SMCAR-TDC Picatinny Arsenal, NJ 07806-5000	(Unclass. only) 1	Commandant U.S. Army Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905-5660
1	Director Benet Weapons Laboratory U.S. Army Armament Research, Development, and Engineering Center ATTN: SMCAR-CCB-TL Watervliet, NY 12189-4050	1	WL/MNOI Eglin AFB, FL 32542-5000
(Unclass. only) 1	Commander U.S. Army Rock Island Arsenal ATTN: SMCRI-TL/Technical Library Rock Island, IL 61299-5000		<u>Aberdeen Proving Ground</u>
1	Director U.S. Army Aviation Research and Technology Activity ATTN: SAVRT-R (Library) M/S 219-3 Ames Research Center Moffett Field, CA 94035-1000	2	Dir, USAMSAA ATTN: AMXSY-D AMXSY-MP, H. Cohen
1	Commander U.S. Army Missile Command ATTN: AMSMI-RD-CS-R (DOC) Redstone Arsenal, AL 35898-5010	1	Cdr, USATECOM ATTN: AMSTE-TC
		3	Cdr, CRDEC, AMCCOM ATTN: SMCCR-RSP-A SMCCR-MU SMCCR-MSI
		1	Dir, VLAMO ATTN: AMSLC-VL-D
		10	Dir, USABRL ATTN: SLCBR-DD-T

**No. of  
Copies    Organization**

1    **Commander  
U.S. Army Research Office  
ATTN: Dr. J. Andersen  
P.O. Box 12211  
Research Triangle Park, NC 27709-2211**



**USER EVALUATION SHEET/CHANGE OF ADDRESS**

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

- 1. BRL Report Number BRL-TR-3394 Date of Report September 1992
- 2. Date Report Received \_\_\_\_\_
- 3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- 4. Specifically, how is the report being used? (Information source, design data, procedure, source of ideas, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- 5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided, or efficiencies achieved, etc? If so, please elaborate. \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- 6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**CURRENT ADDRESS**

\_\_\_\_\_  
Name  
\_\_\_\_\_  
Organization  
\_\_\_\_\_  
Address  
\_\_\_\_\_  
City, State, Zip Code

7. If indicating a Change of Address or Address Correction, please provide the New or Correct Address in Block 6 above and the Old or Incorrect address below.

**OLD ADDRESS**

\_\_\_\_\_  
Name  
\_\_\_\_\_  
Organization  
\_\_\_\_\_  
Address  
\_\_\_\_\_  
City, State, Zip Code

(Remove this sheet, fold as indicated, staple or tape closed, and mail.)

**DEPARTMENT OF THE ARMY**  
Director  
U.S. Army Ballistic Research Laboratory  
ATTN: SLCBR-DD-T  
Aberdeen Proving Ground, MD 21005-5066

**OFFICIAL BUSINESS**

**BUSINESS REPLY MAIL**  
FIRST CLASS PERMIT No 0001, APG, MD

Postage will be paid by addressee.

Director  
U.S. Army Ballistic Research Laboratory  
ATTN: SLCBR-DD-T  
Aberdeen Proving Ground, MD 21005-5066



**NO POSTAGE  
NECESSARY  
IF MAILED  
IN THE  
UNITED STATES**

