

(2)



STARLAB
DEPARTMENT OF ELECTRICAL ENGINEERING / SEL
STANFORD UNIVERSITY • STANFORD, CA 94305

AD-A255 518



LONG-TERM AVERAGES OF GLOBALLY-
MEASURED ELF/VLF RADIO NOISE

by

M.M. Bowen
A.C. Fraser-Smith
P.R. McGill

SDTIC
ELECTE
SEP 02 1992
S A D

Technical Report E450-2

March 1992

This document has been approved
for public release and sale; its
distribution is unlimited.

Sponsored by
The Office of Naval Research
through
Contract No. N00014-81-K-0382 and Grant No. N00014-90-J-1080

92 9 01 003

84478

92-24198



208

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER E 450-2	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) LONG-TERM AVERAGES OF GLOBALLY-MEASURED ELF/VLF RADIO NOISE		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) M.M. BOWEN, A.C. FRASER-SMITH, P.R. MCGILL		8. CONTRACT OR GRANT NUMBER(s) CONTRACT No. N00014-81-K-0382 GRANT No. N00014-90-J-1080
9. PERFORMING ORGANIZATION NAME AND ADDRESS SPACE, TELECOMMUNICATION, AND RADIO SCIENCE LAB. DURAND BUILDING, STANFORD UNIVERSITY STANFORD, CA 94305-4090		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR089-171 and R&T No. 4149171
11. CONTROLLING OFFICE NAME AND ADDRESS OFFICE OF NAVAL RESEARCH, CODE 1114SP 800 QUINCY STREET ARLINGTON, VA 22217		12. REPORT DATE MARCH, 1992
		13. NUMBER OF PAGES 48
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) DISTRIBUTION UNLIMITED, APPROVED FOR PUBLIC RELEASE AND SALE.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) ELF NOISE VLF NOISE GLOBAL NOISE MEASUREMENTS LONG-TERM NOISE MEASUREMENTS SFERICS CHORUS HISS RADIO NOISE STATISTICS		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) IN THIS REPORT WE PRESENT THE RESULTS OF A STATISTICAL STUDY OF A YEAR OF DATA OBTAINED AS PART OF THE STAR LABORATORY'S GLOBAL SURVEY OF ELF/VLF RADIO NOISE (FREQUENCIES IN THE RANGE 10 HZ TO 32 KHZ). THE DATA INCLUDED IN THE STUDY WERE OBTAINED DURING THE ONE-YEAR PERIOD DECEMBER 1986 THROUGH NOVEMBER 1987 FROM SEVEN MEASUREMENT INSTRUMENTS ("RADIOMETERS") LOCATED AT BOTH HIGH AND LOW LATITUDES. AVERAGE AMPLITUDES AND VOLTAGE DEVIATION (V_d) STATISTICS FROM EACH OF THE SEVEN STATIONS WERE FIRST DIVIDED INTO FOUR THREE-MONTH		

BLOCK 20 CONT.

SEASONS AND THEN AVERAGED AT EVERY MEASUREMENT FREQUENCY IN TWO SEPARATE LONG-TERM AVERAGES.

THE FIRST LONG-TERM AVERAGE INVOLVED FURTHER DIVIDING THE MEASUREMENTS INTO THE SIX FOUR-HOUR TIME BLOCKS IN A DAY AND AVERAGING WITHIN THESE HOURLY BLOCKS ACROSS AN ENTIRE SEASON (E.G., ALL MEASUREMENTS MADE BETWEEN 0000 TO 0400 UT FOR A SEASON WERE AVERAGED TOGETHER). THIS PROCESS RESULTED IN A SET OF SIX SPECTRAL DISTRIBUTIONS FOR EACH SEASON. THUS, THE AVERAGES EMPHASIZE DIURNAL TRENDS IN THE NOISE AMPLITUDES AND V_d VALUES. THE SECOND LONG-TERM AVERAGE WAS OBTAINED BY AVERAGING ALL THE MEASUREMENTS TAKEN DURING A SEASON AND THUS OBTAINING ONE SPECTRAL DISTRIBUTION FOR EACH SEASON. THIS LONG-TERM AVERAGE SHOWS SEASONAL FEATURES IN THE NOISE AMPLITUDES AND V_d VALUES.

BOTH SETS OF LONG-TERM AVERAGES SHOW CHARACTERISTIC TRENDS IN THE MEASUREMENTS DUE TO THE MANNER IN WHICH NOISE PROPAGATES IN THE EARTH-IONOSPHERE WAVEGUIDE. THE AVERAGE AMPLITUDES OF THE MEASUREMENTS DECREASE WITH FREQUENCY, REACHING A MINIMUM AROUND 1 KHZ, AND RISE AGAIN TOWARDS 10 KHZ. THE V_d AMPLITUDES GENERALLY RISE WITH FREQUENCY EXCEPT FOR A DEPRESSION OF VALUES AROUND 1 KHZ.

**LONG-TERM AVERAGES OF GLOBALLY-MEASURED
ELF/VLF RADIO NOISE**

by

M. M. BOWEN, A. C. FRASER-SMITH, AND P. R. MCGILL

Technical Report E450-2

March 1992

Sponsored by

The Office of Naval Research

Contract No. N00014-81-K-0382 and Grant No. N00014-90-J-1080

ABSTRACT

In this report we present the results of a statistical study of a year of data obtained as part of the STAR Laboratory's global survey of ELF/VLF radio noise (frequencies in the range 10 Hz to 32 kHz). The data included in the study were obtained during the one-year period December 1986 through November 1987 from seven measurement instruments ("radiometers") located at both high and low latitudes. Average amplitude and voltage deviation (V_d) statistics from each of the six stations were first divided into four three-month seasons and then averaged at every measurement frequency to give two separate sets of long-term averages.

The first set of averages was obtained by further dividing the measurements into the six 4-hour time blocks in a day and averaging within these hourly blocks across an entire season (e.g., all measurements made between 0000 to 0400 UT for a season were averaged together). This process resulted in a set of six spectral distributions for each season. Thus, the averages emphasize diurnal trends in the noise amplitudes and V_d values. The second set of long-term average was obtained by averaging all measurements taken during a season and thus obtaining one spectral distribution for each season. This long-term average shows seasonal features in the noise amplitude and V_d values.

Both sets of long-term averages show characteristic trends in the measurements due to the manner in which noise propagates in the earth-ionosphere waveguide. The average amplitudes of the measurements decrease with frequency, reaching a minimum around 1 kHz, and rise again towards 10 kHz. The V_d amplitudes generally rise with frequency except for a depression of values around 1 kHz.

DTIC QUALITY INSPECTED 3

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By	
Distribution/	
Availability	
Dist	Avail Spec
A-1	

TABLE OF CONTENTS

	Page
ABSTRACT	iii
TABLE OF CONTENTS	v
1. INTRODUCTION	1
2. AMPLITUDE MEASUREMENTS	2
2.1 DATA ANALYSIS	2
2.2 DISCUSSION	3
3. V_d MEASUREMENTS	5
3.1 DATA ANALYSIS	5
3.2 DISCUSSION	5
4. CONCLUSION	6
ACKNOWLEDGEMENTS	6
REFERENCES	8
FIGURES	10

1. INTRODUCTION

As we have previously reported [1, 2, 3] our laboratory is presently conducting a global survey of extremely-low frequency (ELF; 5 Hz-3 kHz) and very-low frequency (VLF; 3 kHz-30 kHz) radio noise covering frequencies in the range 10 Hz-32 Hz. Noise statistics at all eight measurement locations are dominated by lightning-generated noises (predominantly sferics). Data from our three high latitude radiometers (Thule (TH; 76.5 N, 68.8 W) and Søndrestrømfjord (SS; 67.0° N, 50.1° W) in Greenland, and Arrival Heights (AH; 77.8° S, 193.3° W) in the Antarctic) include samples of polar hiss and greater amounts of other ELF/VLF radio noise of magnetospheric origin, such as chorus, than measurements made at our five lower latitude radiometers (Grafton, New Hampshire (43.6° N, 72.0° W); L'Aquila, Italy (43.4° N, 13.3° E); Stanford, California (37.4° N, 122.2° W); Kochi, Japan (33.3° N, 226.5° W); and Dunedin, New Zealand (45.8° S, 189.5° W)). A map showing the geographical locations of these stations is given in Figure 1.

The radio noise statistics computed continuously at each of the stations consist of the average, root-mean-square (rms), maximum, and minimum amplitudes in 16 narrow frequency bands (5% bandwidth) distributed over the ELF and VLF ranges (Table 1). They are computed at the end of every minute from 600 amplitude measurements made at the rate of 10 per second on the envelope of the noise signal emerging from each narrow-band filter (during calibration the number of measurements is reduced but never to less than 400 measurements/minute). In addition to these recordings, synoptic broadband ELF digital recordings are made for one minute every thirty minutes and synoptic broadband VLF analog recordings are made for one minute every hour. Processing of the narrow-band digital data can, with little additional computation, give the voltage deviation (V_d), effective antenna noise factor (F_a), and amplitude probability distributions (APD's). These various statistical quantities are widely used to characterize radio noise, and they are described in several reports issued by the International Radio Consultative Committee, or CCIR [4, 5]. Although we have made comparisons at 32 kHz between our noise data and the F_a statistics presented in the CCIR reports [6], we are unable to make comparisons at all frequencies because 14 of our 16 measurement frequencies are below the 10 kHz minimum frequency for the CCIR statistics.

TABLE 1. Center frequencies and bandwidths for the 16 channels of the ELF/VLF noise measurement systems.

Channel	Center Frequency	Bandwidth (5%)
1	10 Hz	0.5 Hz
2	30	1.5
3	80	4
4	135	6.75
5	275	13.75
6	380	19
7	500	25
8	750 Hz	37.5
9	1 kHz	50
10	1.5	75
11	2	100
12	3	150
13	4	200
14	8	400
15	10.2	510
16	32 kHz	1600 Hz

2. AMPLITUDE MEASUREMENTS

2.1. Data Analysis

The one-minute average data obtained from the stations can be processed in many different ways to give additional information about the morphology of ELF/VLF noise [7, 8]. One important form of processing we use is the computation of average amplitudes over long time intervals, such as three months. The upper panel of Figure 2 shows a three-month average

using data collected at Kochi, Japan (our station of lowest geographic latitude) during winter (December, January, February). Each day was divided into six periods of four hours and the one-minute average amplitudes were averaged together according to the time of day in which they were measured. (For example, all measurements taken between 0000 to 0400 Universal Time (UT) were averaged to obtain one value at each measurement frequency.) The result is the set of six spectral distributions that provide information about the diurnal variation of the ELF/VLF noise spectrum at Kochi during the winter of 1986. Figures 2-3 are similar plots showing the variations at Kochi in spring (March, April, May), summer (June, July, August) and autumn (September, October, November) of 1987. Graphs derived in an identical manner are presented in Figures 4-15 for the same 1986-1987 seasons at six other stations arranged in order of increasing geographic latitude: Stanford, L'Aquila, Dunedin, Søndrestrømfjord, Thule, and Arrival Heights. Although two of our stations are in the southern hemisphere we have used the northern hemisphere season names when referring to specific three-month periods in an effort to retain consistency in our nomenclature. Thus, we have used the season 'winter' to describe the months December, January, and February regardless of the hemisphere in which the station is located.

One-minute amplitude measurements were also averaged for each three-month season in the 1986-1987 year at each of the sixteen measurement frequencies in order to define any seasonal trends in the ELF/VLF noise. The results obtained for each station during the four seasons between December 1986 and November 1987 are shown in Figures 16-19.

2.2. Discussion

The ELF noise amplitudes measured at Thule and L'Aquila below 135 Hz are consistently an order of magnitude greater than those from other stations. Although Thule is considerably closer to a geomagnetic pole than any of our other stations [9], there is no reason to expect higher ELF noise amplitudes near a geomagnetic pole and there is no previous record of such measurements. At this time we tentatively ascribe the high amplitudes to broad band ELF interference from the nearby air base. The higher ELF noise amplitudes at L'Aquila are almost certainly due to interference from the Italian electric railroad system [10].

The amount of usable data collected by the radiometer at Thule is reduced to half by the

operation of the nearby Thule Ionosonde. The ionosonde operates in two-hour periods every other two-hour period (0000-0200 UT, 0400-0600 UT, ..., 2000-2200 UT). Although the primary frequencies produced by the ionosonde range from 10 to 30 kHz, we see contamination at all our measurement frequencies during its periods of operation and thus exclude from our statistical studies all measurements made during these periods.

The measurements made at 32 kHz by the radiometer at Stanford University during the year of data used in this study have been found to be contaminated by a narrowband signal or modulation product at ~ 31.5 kHz that appears to be man-made in origin. An examination of our broadband recordings has showed that this signal occurring sporadically with different strengths at different times throughout the 1986-1987 year.

The 10.2 kHz measurement channel of the radiometers coincides with one of the broadcast frequencies of the globally-located Omega navigation stations. As a result, the data from this channel are occasionally contaminated with the navigation signals. The extent of this contamination depends on the location of the radiometer with respect to the Omega stations and the propagation characteristics of the path between them. The signals are readily identified by examining one-minute synoptic VLF broadband recordings that are made automatically by each radiometer every hour.

The graphs of average noise amplitudes from each station show many of the same general trends. All exhibit a $\sim 1/f$ decrease of amplitude with frequency culminating in a depression of amplitudes around 1 kHz. At higher frequencies the amplitudes rise again, creating an elevation of amplitudes around 10 kHz. Much of this general morphology can be attributed to the characteristics of the earth-ionosphere waveguide. At ELF frequencies only the transverse electromagnetic (TEM) waves can propagate through the guide. As the frequency increases the attenuation of TEM waves in the earth-ionosphere waveguide also increases reducing the amplitude of noise measured at higher frequencies. Around 1-3 kHz the first-order modes of quasi-transverse electric (QTE) and quasi-transverse magnetic (QTM) waves can propagate. As the frequency increases, higher order QTE and QTM modes contribute to the propagation of noise [11]. Thus, the levels of noise begin to increase around 1-3 kHz when the QTE and QTM modes begin to contribute to noise propagation.

3. V_d MEASUREMENTS

3.1. Data Analysis

We processed seasonal averages of the one-minute V_d measurements for each of the six stations in a manner identical to the average amplitude measurement processing. One-minute averages of the V_d measurements were averaged over a three-month season for each of the six four-hour time divisions of a twenty-four-hour UT day at each of the sixteen measurement frequencies. Figures 20-33 show the results using all available data from December 1986 to November 1987 for the seven stations, Kochi, Stanford, L'Aquila, Dunedin, Søndrestrømsfjord, Thule, and Arrival Heights.

The seasonal variations may be examined by averaging the one-minute V_d measurements in each three-month season and comparing the seasons over an entire year. Figures 34-37 are the seasonal averages for the six stations from December 1986 to November 1987.

3.2. Discussion

V_d is a measure of the 'spikiness' of noise data and is computed by taking the ratio (in dB) of the rms amplitude to the average amplitude. For example, noise with Gaussian distributions of both amplitude and phase is characterized by a Rayleigh distribution and has a V_d of 1.05 dB [12]. Distributions containing a greater percentage of measurements at very high amplitudes have greater V_d amplitudes. Natural ELF/VLF noise contains sferics, or lightning-generated bursts of energy, that can have high amplitudes and thus, depending on the amount of lightning activity, can give values of V_d considerably higher than a Rayleigh distribution. Most of the V_d values measured at our stations lie in the range 1-10 dB. However, a sharp minimum occurs near 1 kHz at almost every station. The values of V_d in this minimum are often below those typical of a Rayleigh distribution, and they sometimes drop as low as 0.1 dB, as shown in the lower panel of Figure 35. Low values of V_d are known to occur when measurements are made of a sinusoidal signal within atmospheric noise. An increase in the signal-to-noise ratio results in a decrease in V_d [12]. We believe the reduction of natural noise around 1 kHz, due to the earth-ionosphere waveguide cutoff, allows the detection of very weak coherent signals, such as local man-made noise, that are normally

below the level of natural activity. Apart from the depression of V_d around 1-3 kHz, the values of V_d tend to increase with increasing frequency indicating that noise becomes more impulsive at higher frequencies.

4. CONCLUSION

We have presented long-term statistics of average amplitude and V_d statistics from our global array of ELF/VLF radio noise measurement systems. Data from six stations were individually averaged over a one year period from December 1986 to November 1987. The data were first averaged in periods of four hours over a three-month interval in order to show diurnal variations within a season. The data were also averaged across an entire three-month period in order to show seasonal variations during the year-long measurement period.

The long-term averages of average amplitude and V_d presented here show the effects of the earth-ionosphere waveguide on the propagation of radio noise. For example, the height and conductivity of the ionosphere determine the rate at which the TEM wave attenuates with frequency and the frequency at which higher modes begin to propagate. These effects are apparent in our data. The noise amplitudes tend to decrease with increasing frequency due to increased attenuation of the TEM wave. At 1-3 kHz and above, the QTE and QTM waves begin to propagate and the noise amplitudes increase. The V_d values tend to increase with increasing frequency indicating greater 'spikiness' in the noise. The prominent minimum of V_d values around 1 kHz is probably due to the reduction of natural noise levels allowing the measurement of local weak coherent signals.

ACKNOWLEDGEMENTS

This research was sponsored by the Office of Naval Research through Contract No. N00014-81-K-0382 and Grant No. N00014-90-J-1080. Support for the measurements at Thule, Greenland, was provided by Rome Air Development Center through Contract No. F10629-84-0043. Important logistics support for the measurements at Søndrestromfjord, Greenland, and Arrival Heights, Antarctica, was provided by the National Science Foundation (NSF) through NSF cooperative agreement ATM 88-22560 and NSF Grant DPP-8720167, respectively. We thank Tom Afferton and Paul R. McGill of our laboratory for their assistance with the data; Professors T. Ogawa and M. Kusunose of Kochi University for their cooperation in the mea-

surements in Japan; Mr. J. T. Turtle of Rome Air Development Center for his cooperation in the Thule measurements; Dr. Neil R. Thomson of the University of Otago for his cooperation in the New Zealand measurements; Drs A. Meloni and P. Palangio of the Istituto Nazionale di Geofisica for their cooperation in the Italian measurements, and Dr. J. D. Kelly of SRI International for facilitating the measurements at Søndrestrømfjord.

REFERENCES

1. Fraser-Smith, A. C., and R. A. Helliwell, "The Stanford University ELF/VLF radiometer project: measurement of the global distribution of ELF/VLF electromagnetic noise," *Proc. 1985 IEEE International Symp. on Electromagnetic Compatibility*, IEEE Catalog No. 85CH2116-2: 305-31, 1985.
2. Fraser-Smith, A. C., R. A. Helliwell, B. R. Fortnam, P. R. McGill, C. C. Teague, "A New Global Survey of ELF/VLF Radio Noise," Conf. on Effects of Electromagnetic Noise and Interference on Performance of Military Radio Communication Systems, Lisbon, Portugal, 26-30 October, 1987, Published in *AGARD Conference Proceedings No. 420: 4A-1-4A-7*, 1988.
3. Fraser-Smith, A. C., P. R. McGill, A. Bernadi, R. A. Helliwell, M. E. Ladd, "Global measurements of low-frequency radio noise," in *Environmental and Space Electromagnetics*, Toyko, Japan, pp 191-200, 1989.
4. C. C. I. R., "World distribution and characteristics of atmospheric radio noise," *Report 322*, International Radio Consultative Committee, International Telecommunication Union, Geneva, 1964.
5. C. C. I. R., "Characteristics and applications of atmospheric radio noise data," *Report 322-3*, International Radio Consultative Committee, International Telecommunication Union, Geneva, 1988.
6. Bower M. M., A. C. Fraser-Smith, "A comparison of measured 32 kHz radio noise amplitudes with CCIR model predictions," p. 247 in *Program and Abstracts*, National Radio Science Meeting (URSI), 7-10 January, 1992.
7. Fraser-Smith, A. C., P. R. McGill, R. A. Helliwell, S. Houery, "Radio noise measurements in the long-wave band at Thule, Greenland," *Report TADC-TR-89-94*, Rome Air Development Center, New York, 1989.
8. Fraser-Smith, A. C., T. Ogawa, P. R. McGill, A. Bernardi, M. E. Ladd, R. A. Helliwell, "Measurements of ELF/VLF radio noise in Japan", in *Environmental and Space Electromagnetics*, Toyko, Japan, pp 201-207, 1989.

9. Fraser-Smith, A. C., "Centered and eccentric geomagnetic dipoles and their poles, 1600-1985," *Rev. Geophysics*, 25, 1-16, 1987.
10. Fraser-Smith, A. C., and M. M. Bowen, "The natural background levels of 50/60 Hz radio noise," *IEEE Transactions on Electromagnetic Compatibility*, in press, 1992.
11. Poulsen, W. L., "Modeling of very low frequency wave propagation and scattering within the Earth-ionosphere waveguide in the presence of lower ionospheric disturbances," *Ph. D. Dissertation*, Department of Electrical Engineering, Stanford University, November 1991.
12. Huntoon, Z. McC., A. A. Giordano, "RMS-to-average deviation ratio for interference and atmospheric noise," *Proc. of the Fourth Symp. and Tech. Exhibition on Electromagnetic Compatibility*, Zurich, March 1981.

FIGURES




Figure 1. Locations of the eight ELF/VLF radiometer installations.





Figure 2. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Kochi, Japan, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Kochi, Japan, for Spring 1987.





Figure 3. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Kochi, Japan, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Kochi, Japan, for Autumn 1987.





Figure 4. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Stanford, California, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Stanford, California, for Spring 1987.





Figure 5. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Stanford, California, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Stanford, California, for Autumn 1987.





Figure 6. (Upper panel) Variation in average amplitudes of ELF/VLF noise at L'Aquila, Italy, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at L'Aquila, Italy, for Spring 1987.




Figure 7. (Upper panel) Variation in average amplitudes of ELF/VLF noise at L'Aquila, Italy, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Insufficient data were available at L'Aquila for Autumn 1987.





Figure 8. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Dunedin, New Zealand, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Dunedin, New Zealand, for Spring 1987.




Figure 9. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Dunedin, New Zealand, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Insufficient data were available at Dunedin for Autumn 1987.





Figure 10. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Søndrestrømfjord, Greenland, during Winter 1986. Measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Søndrestrømfjord, Greenland, for Spring 1987.




Figure 11. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Søndrestrømfjord, Greenland, during Autumn 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. No data were available at Søndrestrømfjord for Summer 1987.





Figure 12. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Thule, Greenland, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Measurements made during hours of ionosonde operation have been excluded. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Thule, Greenland, for Spring 1987.





Figure 18. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Thule, Greenland, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Measurements made during hours of ionosonde operation have been excluded. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Thule, Greenland, for Autumn 1987.





Figure 14. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Arrival Heights, Antarctica, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Arrival Heights, Antarctica, for Spring 1987.





Figure 15. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Arrival Heights, Antarctica, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Arrival Heights, Antarctica, for Autumn 1987.




Figure 16. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Kochi, Japan, during the 1986-1987 year. The measurements have been averaged at each frequency for the indicated season. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Stanford, California, for the 1986-1987 year.





Figure 17. (Upper panel) Variation in average amplitudes of ELF/VLF noise at L'Aquila, Italy, during the 1986-1987 year. The measurements have been averaged at each frequency for the indicated season. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Dunedin, New Zealand, for the 1986-1987 year.





Figure 18. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Søndrestrømfjord, Greenland, during the 1986–1987 year. The measurements have been averaged at each frequency for the indicated season. (Lower panel) Variation in average amplitudes of ELF/VLF noise at Thule, Greenland, for the 1986–1987 year. Measurements made during hours of ionosonde operation have been excluded.




Figure 19. (Upper panel) Variation in average amplitudes of ELF/VLF noise at Arrival Heights, Antarctica, during the 1986-1987 year. The measurements have been averaged at each frequency for the indicated season.





Figure 20. (Upper panel) Variations in V_d amplitudes of ELF/VLF noise at Kochi, Japan, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in V_d of ELF/VLF noise at Kochi, Japan, for Spring 1987.





Figure 21. (Upper panel) Variations in the V_g statistic for ELF/VLF noise measured at Kochi, Japan, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_g statistic for ELF/VLF noise measured at Kochi, Japan, for Autumn 1987.





Figure 22. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Stanford, California, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Stanford, California, for Spring 1987.





Figure 23. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Stanford, California, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Stanford, California, for Autumn 1987.





Figure 24. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at L'Aquila, Italy, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at L'Aquila, Italy, for Spring 1987.




Figure 25. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at L'Aquila, Italy, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Insufficient data were available at L'Aquila for Autumn 1987.





Figure 26. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Dunedin, New Zealand, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Dunedin, New Zealand, for Spring 1987.




Figure 27. (Upper panel) Variations in the V_g statistic for ELF/VLF noise measured at Dunedin, New Zealand, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Insufficient data were available at Dunedin for Autumn 1987.





Figure 28. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Søndrestørfjord, Greenland, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Søndrestørfjord, Greenland, for Spring 1987.




Figure 29. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Søndrestromsfjord, Greenland, during Autumn 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. No data were available at Søndrestromsfjord for Summer 1987.





Figure 30. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Thule, Greenland, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Measurements made during hours of ionosonde operation have been excluded. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Thule, Greenland, for Spring 1987.





Figure 31. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Thule, Greenland, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. Measurements made during hours of ionosonde operation have been excluded. (Lower panel) Variation in V_d of ELF/VLF noise at Thule, Greenland, for Autumn 1987.





Figure 32. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Arrival Heights, Antarctica, during Winter 1986. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Arrival Heights, Antarctica, for Spring 1987.





Figure 33. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Arrival Heights, Antarctica, during Summer 1987. The measurements have been averaged at each frequency for the indicated four-hour UT interval. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Arrival Heights, Antarctica, for Autumn 1987.





Figure 34. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Kochi, Japan, during the 1986–1987 year. The measurements have been averaged at each frequency for the indicated season. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Stanford, California, for the 1986–1987 year.





Figure 35. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at L'Aquila, Italy, during the 1986–1987 year. The measurements have been averaged at each frequency for the indicated season. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Dunedin, New Zealand, for the 1986–1987 year.




Figure 36. (Upper panel) Variations in the V_d statistic for ELF/VLF noise measured at Søndrestrømfjord, Greenland, during the 1986-1987 year. The measurements have been averaged at each frequency for the indicated season. (Lower panel) Variations in the V_d statistic for ELF/VLF noise measured at Thule, Greenland, for the 1986-1987 year. Measurements made during hours of ionosonde operation have been excluded.




Figure 37. (Upper panel) Variations in the V_g statistic for ELF/VLF noise measured at Arrival Heights, Antarctica, during the 1986-1987 year. The measurements have been averaged at each frequency for the indicated season.

DISTRIBUTION LIST

Organization	No. of Copies	Organization	No. of Copies
Administrative Grants Officer Office of Naval Research Resident Representative N63375 Administrative Contracting Officer Stanford University Rm 202 McCullough Bldg. Stanford, CA 94305-4055	1	Mr. Donald Butts JCS/J6E Rm ID 827A Washington, D.C. 20301	1
Director, Naval Research Laboratory Attn: Code 2627 Washington, D.C. 20375	1	Karl Cain (Code 832) Naval Ocean Systems Center San Diego, CA 92152-5000	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, VA 22314	4	Dr. Patrick Crowley Defense Nuclear Agency Headquarters Washington, D.C. 20305-1000	1
Mr. Martin R. Allerdig Pacific Sierra Research Corp. 1401 Wilson Blvd., Suite 1100 Arlington, VA 22209	1	Mr. Stanley F. Czajkowski Donna Gregg The Johns Hopkins University Applied Physics Laboratory/CSE Johns Hopkins Road Laurel, MD 20707	2
Dr. Peter Bannister (Code 3411) Naval Underwater Systems Center Fort Trumbull New London, CT 06320	1	Mr. Mark Deeble Space & Naval Warfare Systems Cmd PMW 152-61A Room 5N20 NC1 Washington, D.C. 20363-5100	1
Dr. John S. Belrose Communications Research Center 3701 Carling Avenue P.O. Box 11490 Station "H" Ottawa, Ontario K2H8S2 Canada	1	Dr. Jerry Ferguson (Code 542) Naval Ocean Systems Center San Diego, CA 92152-5000	1
Dr. J. Bond (Code 832) Naval Ocean Systems Center San Diego, CA 92152-5000	1	Dr. E. C. Field Pacific Sierra Research Corp. 12340 Santa Monica Blvd. Los Angeles, CA 90025	1
Dr. Richard Brandt (Code 1112) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000	1	Dr. Anthony J. Ferraro CSSL 316 Electrical Engineering East The Pennsylvania State University University Park, PA 16802	1
Charlie Brown, PMW-153 Space & Naval Warfare Systems Cmd Washington, D.C. 20363-5100	1	Dr. A. C. Fraser-Smith STARLAB Dept. of Electrical Engineering Stanford University Stanford, CA 94305-4055	1
Mr. Richard P. Buckner TCS, Inc. 1475 Richardson Drive, Suite 265 Richardson, TX 75080-4650	1	Greg Gharakhanian The Mitre Corporation 7525 Colshire Drive McLean, VA 22102	1

DISTRIBUTION LIST

Organization	No. of Copies	Organization	No. of Copies
Mr. Marvin Gantsweg RDA Corp. 6053 W. Century Blvd. Los Angeles, CA 90045	1	Mr. R. Gracen Joiner (Code 1114sp) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000	3
Ms. Kathy Grauer (Code 832) NRaD San Diego, CA 92152-5000	1	Mr. Martin E. Katz SAIC 8619 Westwood Center Dr. Vienna, VA 22182	1
Lt. Col. Robert Graney AFCSA/SASS Rm ID 431 - Pentagon Washington, D.C. 20330-5240	1	Dr. Paul Kossey AFGL/LID Hanscom AFB, MA 01731	1
Dr. Robert Helliwell STARLAB Dept. of Electrical Engineering Stanford University Stanford, CA 94305-4055	1	Per Kullstam PAIRCOM 8508 Shadeway Place Springfield, VA 22153	1
Mr. James Higbie The Johns Hopkins University Applied Physics Laboratory Johns Hopkins Road Laurel, MD 20707-60910	1	Dr. Wilford Lauber Communications Research Center 3701 Carling Avenue P.O. Box 11490 Station "H" Ottawa, Ontario K2H8S2 Canada	1
Mr. Vern Hildebrand Innovative Technologies 1133 Savoy Street San Diego, CA 92107	1	Dr. Lewis M. Linson SAIC-MS33 10260 Campus Point Drive San Diego, CA 92121	1
Dr. Umran Inan STARLAB Dept. of Electrical Engineering Stanford University Stanford, CA 94305-4055	1	Dr. Ray J. Lunnen, Jr. Applied Research Laboratory P.O. Box 30 State College, PA 16804	1
Dr. Ray Ingram (Code 3411) Naval Underwater Systems Center Fort Trumbull New London, CT 06320	1	Brad Lytle (MS: Z-435) The Mitre Corporation 7525 Colshire Drive McLean, VA 22102	1
Mr. Stan Jakubiak JCS/J6E Rm ID 834 - Pentagon Washington, D.C. 30301	1	Arnie Michelson GTE Government Systems 77 "A" Street, Bldg. 3-1551 Needham, MA 02919	1
Dr. David H. Johnson (Code 1223) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000	1	Mr. J. McCormick (PMA271A) Naval Air Systems Command Jefferson Plaza 1, Rm 582 Washington, D.C. 20361	1

DISTRIBUTION LIST

Organization	No. of Copies	Organization	No. of Copies
Mr. Edward G. McKay SAIC 8619 Westwood Center Dr. Vienna, VA 22182	1	Dan Sharp The Mitre Corporation Washington C3I Division 7525 Colshire Drive McLean, VA 22102	1
CDR L.E. Miller OCNO (NOP-941D1) Room 5A734 Pentagon Washington, D.C. 20350-5000	1	Mr. Wayne Sherman Attn: C4S/SSS Defense Communications Agency Washington, D.C. 20305-2000	1
Bill Moision NRaD San Diego, CA 92152-5000	1	Mr. Charles M. Smith The Mitre Corporation 7525 Colshire Drive Mail Stop 7-370 McLean, VA 22102	1
Mary Muir JHU/Applied Physics Lab. Rm 13N-536 Johns Hopkins Road Laurel, MD 20723	1	Dr. Gordon Soper OASD (C3, I) Room 3E163 Pentagon Washington, D.C. 20301-3040	1
Mr. David Niemoller SAIC 8619 Westwood Center Dr. Vienna, VA 22182	1	Dr. A. D. Spaulding NTIA/ITS 325 Broadway Boulder, CO 80303	1
Mr. Joe Orose (Code 3032) NADC Warminster, PA 18974	1	Paul Steichen Technology Services Corporation 4565 Hilton Parkway, Suite 204 Colorado Springs, CO 80907	1
Dr. Thomas P. Quinn DASD (C3, I) Room 3E160 Pentagon Washington, D.C. 20301-3040	1	Mr. Bradley L. Stubbs Defense Communications Agency Code: C4S/SSM Washington, D.C. 20305-2000	1
Dr. Juergen Richter (Code 532) NRaD San Diego, CA 92152-5000	1	Dr. David Sutkoff SAIC 8619 Westwood Center Dr. Vienna, VA 22182	1
Dr. David Sailors (Code 542) NRaD San Diego, CA 92152-5000	1	Dr. Albert Tomko (MS13N573) JHU/Applied Physics Laboratory Johns Hopkins Road Laurel, MD 20707	1
Hank Schmidt TSC 962 Wayne Avenue Suite 800 Silver Spring, MD 20910	1	Mr. John Turtle RADC/EEPS Hanscom AFB, MA 01731	1
Dr. Howard Shapiro OASD (C3, I) Room 3E163 Pentagon Washington, D.C. 20301-3040	1		

DISTRIBUTION LIST

Organization	No. of Copies	Organization	No. of Copies
Dr. Chris R. Warber Pacific Sierra Research Corp. 12340 Santa Monica Blvd. Los Angeles, CA 90025	1		
Mr. Don Wiltsy The Johns Hopkins University Applied Physics Laboratory/CSE Johns Hopkins Road Laurel, MD 20707	1		
Dr. Leon Wittwer Capt. Tom Bazzoli Defense Nuclear Agency RAAE Washington, D.C. 20305-1000	2		
LCDR. Rich Wooldridge Naval Air Systems Command PMA 271A Jefferson Plaza 1, Rm 582 Washington, D.C. 20361	1		
Lt. Col. Gary Zank OASD/PA & E Rm AE 286 - Pentagon Washington, D.C. 20301-1880	1		