

REPORT

AD-A256 505

Form Approved
OMB No 0704-0188

2

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed to collect the information, reviewing the collection of information, including suggesting changes to the collection, and reviewing the collection of information, including suggesting changes to the collection, and reviewing the collection of information, including suggesting changes to the collection.



Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed to collect the information, reviewing the collection of information, including suggesting changes to the collection, and reviewing the collection of information, including suggesting changes to the collection.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 10-14-92		3. REPORT TYPE AND DATES COVERED Technical Report	
4. TITLE AND SUBTITLE Synthesis of [(Me ₃ CCH ₂)Ga(PPh ₂) ₂] ₂ from [(Me ₃ CCH ₂)ClGaPPh ₂] ₃				5. FUNDING NUMBERS Grant: N00014-90-J-1530 R&T Code: 4135002	
6. AUTHOR(S) O. T. Beachley, Jr., John D. Malony, and Robin D. Rogers				8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No. 34	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry State University of New York at Buffalo Buffalo, NY 14214				10. SPONSORING / MONITORING AGENCY REPORT NUMBER n/a	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000				11. SUPPLEMENTARY NOTES Accepted for publication - <u>Organometallics</u>	
12a. DISTRIBUTION / AVAILABILITY STATEMENT This document has been approved for public release and sale; its distribution is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States government.				12b. DISTRIBUTION CODE n/a	
13. ABSTRACT (Maximum 200 words) The gallium phosphide [(Me ₃ CCH ₂)Ga(PPh ₂) ₂] ₂ has been prepared from [(Me ₃ CCH ₂)ClGaPPh ₂] ₃ and Li(CMe ₃) in an Et ₂ O/C ₆ H ₆ mixture at 0 °C and fully characterized by elemental analyses, physical properties, ¹ H NMR, ³¹ P NMR and IR spectroscopic data and an X-ray structural study. The identity of the gallium phosphide was confirmed by its independent synthesis from [(Me ₃ CCH ₂)BrGaPPh ₂] ₃ and KPPH ₂ in Et ₂ O/C ₆ H ₆ . The dimeric molecule crystallizes in the monoclinic spacegroup P $\bar{1}$ with cell dimensions of a = 10.106(2)Å, b = 11.146(6)Å, c = 12.266(2)Å, α = 93.91(2)°, β = 101.39(1)°, γ = 95.81(4)°, V = 1342.0Å ³ , and Z = 1. Diffraction data were collected on an Enraf-Nonius CAD-4 diffractometer, and the structure was refined to R = 3.8% and wR = 3.8% for those 3561 unique data with [F _o ≥ 5σ(F _o)]. (continued on next page)					
14. SUBJECT TERMS Gallium-phosphide, gallium chemistry, X-ray structural study.				15. NUMBER OF PAGES 17	
				16. PRICE CODE n/a	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL		

DTIC
ELECTE
OCT 20 1992
S C D

85175
92-27325
00352



92 1 138

The four-membered Ga₂P₂ ring has a planar conformation. The neopentyl groups are arranged trans to each other across the ring. The bridging Ga-P bond distances are 2.4568(9) and 2.4689(9)Å. The terminal Ga-P bond distance is 2.351(1)Å.

DISK CONTAINS INFORMATION FROM 1

SEARCHED
SERIALIZED
INDEXED
FILED
A-1

OFFICE OF NAVAL RESEARCH

Contract N-00014-90-J-1530

R&T Code 4135002

TECHNICAL REPORT NO. 34

Synthesis of $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ from $[(\text{Me}_3\text{CCH}_2)\text{ClGaPPh}_2]_3$

by

O. T. Beachley, Jr., John D. Maloney

and Robin D. Rogers

Prepared for Publication

in

Organometallics

State University of New York at Buffalo
Department of Chemistry
Buffalo, New York 14214

14 October 1992

Reproduction in whole or in part is permitted for
any purpose of the United States Government

*This document has been approved for public release
and sale; its distribution is unlimited

Synthesis of $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ from $[(\text{Me}_3\text{CCH}_2)\text{ClGaPPh}_2]_3$

by

O. T. Beachley, Jr.* and John D. Maloney

Department of Chemistry, State University of New York at Buffalo
Buffalo, NY 14214

and

Robin D. Rogers


Department of Chemistry, Northern Illinois University
DeKalb, IL 60115

Summary: The gallium phosphide $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ has been prepared from $[(\text{Me}_3\text{CCH}_2)\text{ClGaPPh}_2]_3$ and $\text{Li}(\text{CMe}_3)$ in an $\text{Et}_2\text{O}/\text{C}_6\text{H}_6$ mixture at 0°C and fully characterized by elemental analyses, physical properties, ^1H NMR, ^{31}P NMR and IR spectroscopic data and an X-ray structural study. The identity of the gallium phosphide was confirmed by its independent synthesis from $[(\text{Me}_3\text{CCH}_2)\text{BrGaPPh}_2]_3$ and KPPh_2 in $\text{Et}_2\text{O}/\text{C}_6\text{H}_6$. The dimeric molecule crystallizes in the monoclinic spacegroup $\text{P}\bar{1}$ with cell dimensions of $a = 10.106(2)\text{\AA}$, $b = 11.146(6)\text{\AA}$, $c = 12.266(2)\text{\AA}$, $\alpha = 93.91(2)^\circ$, $\beta = 101.39(1)^\circ$, $\gamma = 95.81(4)^\circ$, $V = 1342.0\text{\AA}^3$, and $Z = 1$. Diffraction data were collected on an Enraf-Nonius CAD-4 diffractometer, and the structure was refined to $R = 3.8\%$ and $wR = 3.8\%$ for those 3561 unique data with $[F_o \geq 5\sigma(F_o)]$. The four-membered Ga_2P_2 ring has a planar conformation. The neopentyl groups are arranged trans to each other across the ring. The bridging Ga-P bond distances are $2.4568(9)$ and $2.4689(9)\text{\AA}$. The terminal Ga-P bond distance is $2.351(1)\text{\AA}$.

The synthesis and characterization of heavier group 13 element compounds with three different substituents including two different organic substituents is of interest. The only examples of fully characterized compounds of this type are in indium chemistry and include $[\text{In}(\text{CH}_2\text{CMe}_3)(\text{CH}_2\text{SiMe}_3)\text{Cl}]_2$ ¹ and $\{\text{In}[\text{CH}(\text{SiMe}_3)_2](i\text{-Pr})\text{Cl}\}_2$.² Two other indium compounds of this general type, $[\text{In}(\text{CH}_2\text{CMe}_3)(\text{C}_6\text{H}_5)\text{Cl}]_2$ ¹ and $[(\text{Me}_3\text{CCH}_2)(\text{Me}_3\text{SiCH}_2)\text{InPEt}_2]_2$,¹ have been prepared but neither compound has been characterized by an X-ray structural study. When the syntheses of the closely related gallium compounds,¹ $\text{Ga}(\text{Me})(\text{CH}_2\text{CMe}_3)\text{Cl}$ and $\text{Ga}(\text{CH}_2\text{CMe}_3)(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}$, were attempted, impure products were isolated. The spectroscopic data suggested that ligand redistribution reactions to form symmetrized products had occurred. These observations suggested to us that the chlorine bridges between the gallium atoms in these compounds were apparently too weak to kinetically stabilize a dimer with two different organic substituents. Thus, our goal was to attempt the synthesis of an organogallium compound with two different organic substituents by using a bridging group that would be hopefully stronger than chlorine. The starting material³ $[(\text{Me}_3\text{CCH}_2)(\text{Cl})\text{GaPPh}_2]_3$ was selected for our synthetic studies. The bridging PPh_2 group was able to stabilize a trimer rather than a dimer, an observation consistent with stronger bridge bonding, but the chloride group would still be available for replacement by a new and different organic group.

The reaction of $[(\text{Me}_3\text{CCH}_2)\text{ClGaPPh}_2]_3$ ³ with LiCMe_3 in benzene/ Et_2O at 0 °C lead to the isolation of $(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$ in 55% yield based upon the monomeric gallium-phosphide. This reaction has been repeated and reproducible results have been obtained. The gallium-phosphorus product was characterized by its physical properties, an X-ray structural study and ¹H NMR, ³¹P NMR and IR spectroscopic data. The identity of

gallium-phosphorus product as $(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$ was also confirmed by its independent synthesis in 80% yield from the reaction of $[(\text{Me}_3\text{CCH}_2)\text{BrGaPPh}_2]_3$ ³ with KPPH_2 in benzene/ Et_2O at 0 °C (Equation 1). The product of this reaction was also fully



characterized including a partial elemental analysis (C and H). All data suggest that the products from the two reactions are one in the same. It is regrettable that the characterization data do not include cryoscopic molecular weight studies. The compound had insufficient solubility in benzene.

The structure of neopentylgallium-bis(diphenylphosphide) consists of dimeric units $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$. Similar non-bridging groups are arranged trans to each other across the ring. There are no abnormally close contacts in the unit cell. The labelling of the atoms in the molecule is depicted in Figure 1. The interatomic bond distances and angles are listed in Tables 1 and 2.

The four-membered Ga_2P_2 ring of $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ resides around a crystallographic center of inversion and is planar. The related compound⁴ $[(\text{Me}_3\text{CCH}_2)_2\text{GaPPh}_2]_2$ has a buckled or butterfly ring geometry whereas $[\text{Bu}_2^{\text{n}}\text{GaPBu}_2^{\text{t}}]_2$ ⁵ and $[\text{Bu}_2^{\text{t}}\text{GaP}(\text{H})(\text{C}_5\text{H}_9)]_2$ ⁶ have planar Ga_2P_2 rings. The bridging Ga-P bond distances in $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ are 2.4568(9)Å and 2.4689(9)Å whereas the terminal Ga-P distances are shorter at 2.351(1)Å. The bridging Ga-P bond distances are comparable to the distances in $[(\text{Me}_3\text{CCH}_2)_2\text{GaPPh}_2]_2$ ⁴ of 2.479(3) to 2.512(3)Å and the distances⁶ in $[\text{Bu}_2^{\text{t}}\text{GaP}(\text{H})(\text{C}_5\text{H}_9)]_2$ of 2.451(1)Å. The shorter terminal Ga-P bond

distances of 2.351(1)Å is comparable to the terminal Ga-P distances of 2.34(1)Å in the monomeric compound,⁷ Ga[P(H)(2,4,6-Bu₃^tC₆H₂)]₃ and of 2.326(4) and 2.323(5)Å in the other monomeric compound⁸ Bu^tGa[P(H)(2,4,6-Bu₃^tC₆H₂)]₂, the distance⁹ of 2.338(1)Å in [(Me₂NCH₂)₂C₆H₃}GaPSiPh₃]₂ and the distance of 2.360Å in gallium phcsphide.¹⁰ Thus, it appears unlikely that pi bonding is the reason for the terminal gallium-phosphorus distance being shorter than the bridging gallium-phosphorus distance.

Angles within the Ga₂P₂ ring are acute at gallium (P(1)-Ga-P(1a) = 84.87(3)°) and obtuse at phosphorus (Ga-P-Ga(a) = 95.13(3)°). The angles within the Ga₂P₂ ring in the closely related molecule⁴ [(Me₃CCH₂)₂GaPPh₂]₂ are related similarly. The tetrahedral geometry at the gallium(III) atom is irregular. The angle between the terminal and bridging phosphorus atoms at gallium (P(1)-Ga-P(2)) is 106.89(1)° whereas the terminal phosphorus-gallium-carbon angle (P(2)-Ga-C(1) = 131.6(1)°) is even larger. The angle between the α-carbon atoms of the terminal neopentyl groups at gallium⁴ in [(Me₃CCH₂)₂GaPPh₂]₂ are 124.35(50)° and 121.98(49)°.

The formation of [(Me₃CCH₂)Ga(PPh₂)₂]₂ from the reaction of [(Me₃CCH₂)(Cl)GaPPh₂]₃ with LiCMe₃ demonstrates that dissociation of the initial six-membered ring and rearrangement of the substituents bound to gallium must have occurred. The gallium-phosphorus ring must have come apart for a dimer to be formed from a trimer. Furthermore, ligand redistribution reactions must have occurred for a product with two gallium-phosphorus bonds in the simplest repeating unit to have been formed from a reactant with only one gallium-phosphorus bond in the simplest repeating unit. The diphenylphosphide group is apparently not a sufficiently strong bridging group in these

gallium compounds to prevent the break-up of the original six-membered ring. Similarly, the reaction of $[(\text{Me}_3\text{CCH}_2)(\text{Cl})\text{GaPPh}_2]_3$ with KPPh_2 to form $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ also confirms that the original six-membered is not maintained during this reaction either. These experimental observations clearly demonstrated that gallium phosphide compounds do not have to maintain their ring structure as they undergo substitution reactions.

The ^1H and ^{31}P NMR spectra of $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ are consistent with the presence of cis and trans isomers of dimeric molecules in d^6 -benzene solution. It is regrettable but it is not possible to definitively assign specific lines to a given isomer. However, one might expect on the basis of the solid state structural study and of steric arguments that the trans isomer might be the more abundant isomer in solution. The ^1H NMR spectrum revealed two singlets at 0.58 and 0.76 ppm for the methyl protons of the neopentyl groups and two multiplets at 1.13 and 1.63 for the methylene protons of the neopentyl groups. The lines at 0.58 and 1.13 ppm are the lines of higher intensity. The multiplets for the methylene protons arise from coupling with the terminal and bridging phosphorus atoms. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum consisted of two sets of two triplets each. The triplets of equal intensity at -22.7 and -45.9 ppm are assigned to the terminal and bridging phosphorus atoms, respectively, of the more abundant isomer. These assignments are based on the ^{31}P NMR spectra of $(\text{Me}_3\text{CCH}_2)_2\text{InPPh}_2$,⁴ $(\text{Me}_3\text{SiCH}_2)_2\text{InPPh}_2$ ¹¹ and $(\text{Me}_3\text{SiCH}_2)_2\text{GaPPh}_2$ ¹¹ which exist as monomer-dimer equilibrium mixtures. The ^{31}P NMR signal for the monomeric species was downfield of the signal for the dimeric species. Since the monomeric species would have a phosphorus atom analogous to a terminal phosphide in $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ and the phosphorus in the dimeric species is typical of a bridging phosphorus, the downfield signal is assigned to the terminal phosphorus atoms.

Triplets arise from phosphorus-31 coupling between the bridging and terminal phosphorus atoms. If the more abundant isomer in solution is the trans isomer, the trans isomer accounts for more than 85% of the species in solution.

EXPERIMENTAL

All compounds described in this investigation were extremely sensitive to oxygen and moisture and were manipulated in a standard vacuum line or under a purified argon atmosphere. Solvents were dried by conventional procedures. The compounds³ $(\text{Me}_3\text{CCH}_2)\text{ClGaPPh}_2$ and $(\text{Me}_3\text{CCH}_2)\text{BrGaPPh}_2$ were prepared and purified by literature methods. Tertbutyllithium was purchased as a 1.7 M solution in pentane from Aldrich. The pentane was removed by vacuum distillation and LiCMe_3 was purified by sublimation at 50 °C. Diphenylphosphine was purchased from Strem Chemicals, Inc., and was purified by distillation. Potassium diphenylphosphide¹² (KPPh_2) was prepared from excess diphenylphosphine and KH in diethyl ether. Elemental analyses were performed by Schwarzkopf Microanalytical Laboratory, Woodside, NY. Infrared spectra of Nujol mulls between CsI plates were recorded by means of a Perkin-Elmer 683 spectrometer. The ^1H NMR spectra were recorded at 300 MHz by using a Varian Gemini-300 spectrometer, respectively. Proton chemical shifts are reported in δ units (ppm) and are referenced to SiMe_4 at δ 0.00 ppm and C_6D_6 at δ 7.15 ppm. The ^{31}P NMR spectrum was recorded at 161.9 MHz on a Varian VXR-400 spectrometer. Proton-decoupled ^{31}P NMR spectra are referenced to 85% H_3PO_4 at δ 0.00 ppm. All samples for NMR spectra were contained in sealed NMR tubes. Melting points were observed in sealed capillaries and are uncorrected.

Reaction of $(\text{Me}_3\text{CCH}_2)\text{ClGaPPh}_2$ with LiCMe_3 . A side-arm dumper charged with 0.221 g (3.45 mmol) of LiCMe_3 was attached to a two-neck flask which contained 1.249 g (3.455 mmol) of $(\text{Me}_3\text{CCH}_2)\text{ClGaPPh}_2$ dissolved in a mixture of 10 mL of benzene and 20 mL of Et_2O . After the mixture was cooled to 0 °C, the LiCMe_3 was added slowly. The resulting mixture was stirred for 18 h at ambient temperature and then the solvents were removed by vacuum distillation. The two-neck flask was fitted with a medium frit equipped with a Schlenk receiving vessel. Three extractions with 30 mL of benzene provided a soluble brown goo. The brown goo was then washed twice with 15 mL of anhydrous pentane to leave a pentane insoluble, colorless solid (0.438 g, 0.946 mmol, 54.8% yield of $(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$). Recrystallization of $(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$ by using slow diffusion of anhydrous pentane into a saturated benzene solution afforded crystallographic quality crystals. **$(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$.** mp 206-210 °C (dec). ^1H NMR (C_6D_6 , δ): 0.58 (s, 8.8 H, $-\text{CMe}_3$), 0.76 (s, 1.1 H, $-\text{CMe}_3$), 1.13 (m, 2.0 H, $-\text{CH}_2-$), 1.63 (m, 0.3 H, $-\text{CH}_2-$). $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6 , δ): -22.7 (t, $^2J_{\text{PGaP}} = 27.5$ Hz, 9.0), -25.0 (t, $^2J_{\text{PGaP}} = 35.2$ Hz, 1.2), -45.9 (t, $^2J_{\text{PGaP}} = 29.1$ Hz, 8.2), -49.4 (t, $^2J_{\text{PGaP}} = 34.0$ Hz, 1.0). IR (Nujol mull cm^{-1}): 1582(m), 1569(w), 1479(m), 1432(m), 1377(m), 1365(m), 1359(m), 1236(vw), 1159(vw), 1134(w), 1104(vw), 1093(vw), 1070(vw), 1026(m), 1002(vw), 747(sh), 742(vs), 736(m), 721(w), 709(m), 700(s), 690(w), 595(w), 504(m), 479(m).

Synthesis of $(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$. To a solution of $(\text{Me}_3\text{CCH}_2)\text{BrGaPPh}_2$ (0.945 g, 2.33 mmol) in a mixture of 20 mL of benzene and 10 mL of Et_2O at 0 °C was added KPPH_2 (0.522 g, 2.33 mmol). The resulting mixture was stirred for 20 h at ambient temperature. Repetitive extractions with benzene provided a crude product which was washed with 15 mL of anhydrous pentane at ambient temperature to yield 0.867 g of

$(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$ (1.87 mmol, 80.4% based on $(\text{Me}_3\text{CCH}_2)\text{BrGaPPh}_2$). The product was recrystallized from a saturated benzene solution layered with pentane.

$(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2$. mp 218-219 °C (dec). ^1H NMR (C_6D_6 , δ): 0.60 (s, 9.1 H, $-\text{CMe}_3$), 0.78 (s, 0.9 H, $-\text{CMe}_3$), 1.15 (m, 2.0 H, $-\text{CH}_2-$), 1.63 (m, 0.2 H, $-\text{CH}_2-$). $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6 , δ): -22.4 (t, $^2J_{\text{PGaP}} = 28.7$ Hz, 10.4), -24.3 (t, $^2J_{\text{PGaP}} = 32.9$ Hz, 1.0), -45.6 (t, $^2J_{\text{PGaP}} = 28.3$ Hz, 7.7), -48.7 (t, $^2J_{\text{PGaP}} = 33.8$ Hz, 1.1). Anal. Calcd.: C, 68.13; H, 6.11. Found: C, 68.18; H, 6.09. IR (Nujol mull cm^{-1}): 1579(m), 1565(w), 1475(m), 1428(m), 1373(m), 1360(m), 1355(m), 1232(vw), 1155(w), 1130(w), 1099(vw), 1088(vw), 1067(vw), 1022(m), 998(vw), 745(sh), 738(vs), 732(s), 719(w), 705(m), 695(sh), 690(s), 592(w), 500(m), 475(m).

X-ray Data Collection, Structure Determination and Refinement for

$[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$. A transparent crystal of the title compound was mounted in a thin-walled glass capillary under Ar and transferred to the goniometer. The space group was determined to be either the centric $\text{P}\bar{1}$ or acentric P1. The subsequent solution and successful refinement of the structure was carried out in the centric space group $\text{P}\bar{1}$. A summary of data collection parameters is given in Table 3.

Least-squares refinement with isotropic thermal parameters led to $R = 0.075$. The geometrically constrained hydrogen atoms were placed in calculated positions 0.95\AA from the bonded carbon atom and allowed to ride on that atom with B fixed at 5.5\AA^2 . The methyl hydrogen atoms were included as a rigid group with rotational freedom at the bonded carbon atom ($\text{C-H} = 0.95\text{\AA}$, $B = 5.5\text{\AA}^2$). Refinement of nonhydrogen atoms with anisotropic temperature factors led to the final values of $R = 0.038$ and $R_w = 0.038$. The final values of the positional parameters are given in Table 4.

Acknowledgement. This work was supported in part by the Office of Naval Research (O.T.B.) and by a generous grant from the Eastman Kodak Co.

Supplementary Material. Listings of anisotropic thermal parameters and calculated positions of hydrogen atoms (pages); a listing of observed and calculated structure factors for $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ (pages). Ordering information is given on any current masthead page.

References

1. Beachley, O. T., Jr.; Maloney, J. D.; Churchill, M. R.; Lake, C. H. *Organometallics* **1991**, *10*, 3568.
2. Neumueller, B. *Naturforsch., B.: Chem. Sci.* **1991**, *46*, 1539.
3. Beachley, O. T., Jr.; Maloney, J. D.; Rogers, R. D. *J. Organomet. Chem.* (Submitted).
4. Banks, M. A.; Beachley, O. T., Jr.; Buttrey, L. A.; Churchill, M. R.; Fettinger, J. C. *Organometallics* **1991**, *10*, 1901.
5. Cowley, A. H.; Jones, R. A.; Benac, B. L.; Kidd, K. B.; Ekerdt, J. G.; Lee, J. Y.; Miller, J. E. *J. Am. Chem. Soc.* **1988**, *110*, 6248.
6. Heaton, D. E.; Jones, R. A.; Kidd, K. B.; Cowley, A. H.; Nunn, C. M. *Polyhedron* **1988**, *7*, 1901.
7. Cowley, A. H.; Jones, R. A.; Mardones, M. A.; Ruiz, J.; Atwood, J. L.; Bott, S. G. *Angew. Chem. Int. Ed. Engl.* **1990**, *29*, 1150.
8. Atwood, D. A.; Cowley, A. H.; Jones, R. A.; Mardones, M. A. *J. Am. Chem. Soc.* **1991**, *113*, 7050.
9. Cowley, A. H.; Jones, R. A.; Arif, A. M.; Benac, B. L.; Kidd, K. B.; Greets, R. L.; Power, J. M.; Schwab, S. T. *J. Am. Chem. Soc. Chem. Commun.* **1986**, 1543.
10. Wyckoff, R. W. G. "Crystal Structures", 2nd Edition; John Wiley, New York, 1963; Vol. 1, p. 108.
11. Beachley, O. T., Jr.; Kopasz, J. P. B.; Hunter, W. E.; Zhang, H.; Atwood, J. L. *J. Organomet. Chem.* **1987**, *325*, 69.
12. Beachley, O. T., Jr.; Tessier-Youngs, C. *Organometallics* **1983**, *2*, 796.

13. Sheldrick, G. M. SHELX76, a system of computer programs for X-ray structure determination as locally modified, University of Cambridge, England, 1976.
14. Sheldrick, G. M. SHELXS, *Acta Cryst.* **1990**, *A46*, 467.
15. "International Tables for X-ray Crystallography"; Vol. IV, Kynoch Press, Birmingham, England, 1974, pp. 72, 99, 149. (Present distributor: Kluwer Academic Publishers, Dordrecht and Boston.)

Table 1. Interatomic Distances (Å) for [(Me₃CCH₂)Ga(PPh₂)₂]₂.**(A) Gallium-Phosphorus Distances**

Ga-P(1)	2.4689(9)	Ga-P(2)	2.351(1)
Ga-P(1a) ^a	2.4568(9)		

(B) Gallium-Carbon Distances

Ga-C(1)	2.001(3)		
---------	----------	--	--

(C) Phosphorus-Carbon Distances

P(1)-C(6)	1.822(3)	P(2)-C(18)	1.835(4)
P(1)-C(12)	1.825(3)	P(2)-C(24)	1.835(4)

(D) Neopentyl Carbon-Carbon Distances

C(1)-C(2)	1.539(5)	C(2)-C(3)	1.507(5)
C(2)-C(4)	1.518(5)	C(2)-C(5)	1.508(5)

(E) Phenyl Carbon-Carbon Distances

C(6)-C(7)	1.392(4)	C(6)-C(11)	1.384(5)
C(7)-C(8)	1.370(5)	C(8)-C(9)	1.374(5)
C(9)-C(10)	1.366(5)	C(10)-C(11)	1.389(5)
C(12)-C(13)	1.398(5)	C(12)-C(17)	1.391(5)
C(13)-C(14)	1.390(5)	C(14)-C(15)	1.376(6)
C(15)-C(16)	1.366(6)	C(16)-C(17)	1.384(5)
C(18)-C(19)	1.385(5)	C(18)-C(23)	1.394(5)
C(19)-C(20)	1.384(5)	C(20)-C(21)	1.367(6)
C(21)-C(22)	1.363(6)	C(22)-C(23)	1.382(5)
C(24)-C(25)	1.382(5)	C(24)-C(29)	1.383(5)
C(25)-C(26)	1.388(5)	C(26)-C(27)	1.362(6)
C(27)-C(28)	1.356(6)	C(28)-C(29)	1.400(5)

a – Atoms designated by *a* are related by the crystallographic center of inversion.

Table 2. Angles (deg) for [(Me₃CCH₂)Ga(PPh₂)₂]₂.**(A) Angles around the Gallium Atom**

P(1)-Ga-P(1a)	84.87(3)	P(1)-Ga-C(1)	103.4(1)
P(1)-Ga-P(2)	106.89(3)	P(2)-Ga-C(1)	131.6(1)
P(2)-Ga-P(1a)	110.75(4)	C(1)-Ga-P(1a)	108.7(1)

(B) Angles around the Phosphorus Atom

Ga-P(1)-Ga(a)	95.13(3)	Ga-P(2)-C(18)	104.2(1)
Ga-P(1)-C(6)	120.8(1)	Ga-P(2)-C(24)	106.3(1)
Ga-P(1)-C(12)	108.1(1)	C(18)-P(2)-C(24)	104.0(2)
C(6)-P(1)-C(12)	104.2(2)		

(C) Gallium-Carbon-Carbon Angles

Ga-C(1)-C(2)	121.9(2)		
--------------	----------	--	--

(D) Phosphorus-Carbon-Carbon Angles

P(1)-C(6)-C(7)	120.4(3)	P(1)-C(6)-C(11)	121.5(3)
P(1)-C(12)-C(13)	119.4(3)	P(1)-C(12)-C(17)	121.8(3)
P(2)-C(18)-C(19)	124.9(3)	P(2)-C(18)-C(23)	117.3(3)
P(2)-C(24)-C(25)	118.9(3)	P(2)-C(24)-C(29)	122.9(3)

(E) Neopentyl Carbon-Carbon-Carbon Angles

C(1)-C(2)-C(3)	110.7(3)	C(1)-C(2)-C(4)	110.0(3)
C(1)-C(2)-C(5)	110.2(3)	C(3)-C(2)-C(4)	108.8(4)
C(3)-C(2)-C(5)	109.0(4)	C(4)-C(2)-C(5)	108.2(4)

Table 2. (cont.)

(E) Phenyl Carbon-Carbon-Carbon Angles

C(7)-C(6)-C(11)	117.9(3)	C(6)-C(7)-C(8)	121.1(4)
C(7)-C(8)-C(9)	120.2(4)	C(8)-C(9)-C(10)	120.0(4)
C(9)-C(10)-C(11)	120.1(4)	C(6)-C(11)-C(10)	120.7(4)
C(13)-C(12)-C(17)	118.7(3)	C(12)-C(13)-C(14)	119.8(4)
C(13)-C(14)-C(15)	120.4(4)	C(14)-C(15)-C(16)	120.2(4)
C(15)-C(16)-C(17)	120.3(4)	C(12)-C(17)-C(16)	120.6(4)
C(19)-C(18)-C(23)	117.8(4)	C(18)-C(19)-C(20)	120.9(4)
C(19)-C(20)-C(21)	120.5(4)	C(20)-C(21)-C(22)	119.4(4)
C(21)-C(22)-C(23)	121.1(4)	C(18)-C(23)-C(22)	120.3(4)
C(25)-C(24)-C(29)	117.9(4)	C(24)-C(25)-C(26)	121.2(4)
C(25)-C(26)-C(27)	120.2(5)	C(26)-C(27)-C(28)	119.8(4)
C(27)-C(28)-C(29)	120.7(5)	C(24)-C(29)-C(28)	120.2(4)

Caption to Figure

Labeling of atoms in $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$ (ORTEP diagram; 50% ellipsoids with hydrogen atoms eliminated for clarity).




Table 3. Crystal Data and Summary of Intensity Data Collection and Structure Refinement of $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$.

Molecular Formula	$\text{C}_{58}\text{H}_{62}\text{Ga}_2\text{P}_4$
Color/Shape	Colorless/parallelepiped
Mol. wt.	1022.5
Space group	$\bar{P}1$
Temp., °C (K)	20(293)
Cell Constants ^a	
a, Å	10.106(2)
b, Å	11.146(6)
c, Å	12.266(2)
α , deg	93.91(4)
β , deg	101.39(1)
γ , deg	95.81(4)
Volume, Å ³	1342.0
Z	1 (dimer)
$D_{\text{(calcd)}}$, g cm ⁻³	1.27
$\mu_{\text{(calcd)}}$, cm ⁻¹	12.19
Diffractometer/Scan	Enraf-Nonius CAD-4/ ω -2 θ
Range of relative transm. factors, %	91/100
Radiation, graphite monochromator	$\text{MoK}\alpha$ ($\lambda=0.71073$)
Max crystal dimensions, mm	0.30 × 0.33 × 0.40
Scan width	0.80 + 0.35tan θ
Standard reflections	300; 050; 007
Decay of standards	±2.5%
Reflections measured	4202
2 θ range, deg	2 ≤ 2 θ ≤ 50
Range of h,k,l	+12, ±13, ±14
Reflections observed [$F_o \geq 5\sigma(F_o)$] ^b	3561
Computer programs ^c	SHELX ¹³
Structure solution	SHELXS ¹⁴
No. of parameters varied	298
Weights	$[\sigma(F_o)^2 + 0.0001 F_o^2]^{-1}$
GOF	0.47
$R = \Sigma F_o - F_c / \Sigma F_o $	0.038
R_w	0.038
Largest feature final diff. map	0.4e ⁻ Å ⁻³

^aLeast-squares refinement of $((\sin\theta)/\lambda)^2$ values for 25 reflections $\theta > 19^\circ$.

^bCorrections: Lorentz-polarization and absorption (empirical, psi scan).

^cNeutral scattering factors and anomalous dispersion corrections from ref 15.

Table 4. Final Fractional Coordinates for $[(\text{Me}_3\text{CCH}_2)\text{Ga}(\text{PPh}_2)_2]_2$.

Atom	x/a	y/b	z/c	B(eqv) ^a
Ga	0.09386(4)	0.07098(3)	0.13463(3)	1.96
P(1)	0.02655(9)	-0.13588(7)	0.04173(7)	1.91
P(2)	0.33038(9)	0.11134(9)	0.15246(8)	2.29
C(1)	-0.0245(4)	0.0809(3)	0.2470(3)	2.54
C(2)	0.0015(4)	0.1843(3)	0.3412(3)	2.75
C(3)	0.1381(4)	0.1827(5)	0.4170(3)	4.13
C(4)	-0.1078(5)	0.1708(5)	0.4100(4)	4.92
C(5)	-0.0027(6)	0.3049(4)	0.2927(4)	5.27
C(6)	0.1540(3)	-0.2301(3)	0.0108(3)	2.02
C(7)	0.1226(4)	-0.3551(3)	-0.0097(3)	2.69
C(8)	0.2138(4)	-0.4263(3)	-0.0413(3)	3.46
C(9)	0.3387(4)	-0.3748(4)	-0.0536(4)	3.75
C(10)	0.3726(4)	-0.2524(4)	-0.0335(4)	3.67
C(11)	0.2803(4)	-0.1797(4)	-0.0022(3)	2.94
C(12)	-0.0635(4)	-0.2241(3)	0.1303(3)	2.19
C(13)	0.0104(4)	-0.2833(3)	0.2148(3)	2.96
C(14)	-0.0571(5)	-0.3483(4)	0.2844(3)	3.93
C(15)	-0.1958(5)	-0.3529(4)	0.2719(4)	4.27
C(16)	-0.2687(5)	-0.2939(4)	0.1907(4)	3.89
C(17)	-0.2036(4)	-0.2294(3)	0.1199(3)	2.87
C(18)	0.4006(4)	0.0093(3)	0.2544(3)	2.37
C(19)	0.3269(4)	-0.0559(4)	0.3193(3)	3.19
C(20)	0.3875(5)	-0.1340(4)	0.3914(4)	3.89
C(21)	0.5223(5)	-0.1470(4)	0.4014(4)	3.77
C(22)	0.5964(4)	-0.0835(4)	0.3383(4)	3.74
C(23)	0.5372(4)	-0.0070(4)	0.2639(4)	3.27
C(24)	0.3796(4)	0.2628(3)	0.2268(3)	2.48
C(25)	0.3238(4)	0.3601(4)	0.1798(4)	3.49
C(26)	0.3647(5)	0.4775(4)	0.2285(4)	4.40
C(27)	0.4631(6)	0.4989(4)	0.3232(4)	4.65
C(28)	0.5206(5)	0.4051(4)	0.3704(4)	4.39
C(29)	0.4804(4)	0.2861(4)	0.3224(3)	3.43

$$a_{\text{B(eqv)}} = 4/3[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33} + ab(\cos \gamma)\beta_{12} + ac(\cos \beta)\beta_{13} + bc(\cos \alpha)\beta_{23}]$$

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)*
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Elek Lindner (1)
Naval Command, Control and Ocean
Surveillance Center
RDT&E Division
San Diego, CA 92152-5000

Dr. Bernard E. Douda (1)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

Dr. Harold H. Singerman (1)
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Defense Technical Information
Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

* Number of copies to forward