

2

OFFICE OF NAVAL RESEARCH  
Grant No. N00014-90-J-1263  
RCT Project 4133002---05  
Technical Report #17

AD-A261 080



A SIMPLE MODEL OF ION SOLVATION WITH NON ADDITIVE CORES

by

W. R. Fawcett\* and L. Blum\*\*

Prepared for Publication in the  
Ukr. Acad. (M. Holovko festschrift)

DTIC  
SELECTED  
FEB 16 1993  
S B D

\*Department of Chemistry, University of California  
Davis, CA 95616

and

\*\*Department of Physics, University of Puerto Rico  
P.O. Box 23343  
Río Piedras, P.R. 00931-3343, USA

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

\*This document has been approved for public release  
and sale; its distribution is unlimited

\*This statement should also appear in Item 10 of Document Control  
Data - DD Form 1473. Copies of form available from cognizant  
contract administrator.

9

93-02826

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION		1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report #17		5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Physics Department University of Puerto Rico	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Río Piedras Puerto Rico 00931-3343		7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Chemistry Office of Naval Research	8b. OFFICE SYMBOL (If applicable) Code 472	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER RCT Project 4133002---05		
8c. ADDRESS (City, State, and ZIP Code) Arlington Virginia 22217-5000		10. SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO	PROJECT NO	TASK NO
11. TITLE (Include Security Classification) A Simple Model for Ion Solvation with Non Additive Cores				
12. PERSONAL AUTHOR(S) W.R. Fawcett and L. Blum				
13a. TYPE OF REPORT Paper	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1-15-93	15. PAGE COUNT 12	
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP			SUB-GROUP
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  The solvation of simple ions is studied with a model with non additive hard cores using the ion-dipole mean spherical approximation.				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Robert Nowak		22b. TELEPHONE (Include Area Code) (203)696-4410	22c. OFFICE SYMBOL ONR 472	

# A Simple Model for Ion Solvation with Non Additive Cores\*

**W. Ronald Fawcett**

Department of Chemistry

University of California

Davis, CA 95616

U.S.A.

and

**Lesser Blum**

Department of Physics

P.O. Box 23343

University of Puerto Rico

Rio Piedras, PR 00931

U.S.A.

\*In honor of Dr. Miroslav Holovko

DTIC QUALITY INSPECTED 3

<b>Accession For</b>	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## INTRODUCTION

Ionic solvation has been the subject of current interest for at least a century.<sup>1</sup> Modern experimental techniques<sup>2</sup> and the availability of powerful computers<sup>3</sup> have produced real progress in the understanding of this complex problem.

It is clear that in many problems the knowledge of the very complex details of the close range interactions are more than what one needs. In the interpretation of thermodynamic and kinetic parameters averages over large domains are predominant, and therefore, simple, yet controlled theories not only provide a way of fitting data, but also useful physical insight.

The success of the Marcus theory of electron transfer kinetics<sup>4</sup> in solution is a good example. In this theory the polarization of the solvent is evaluated from a continuum model for the solvent, that is, the structure of the solvent is completely ignored. The recent rather spectacular calculations of Chandler and his group<sup>5</sup> have conclusively shown that within certain limitations, this is not unreasonable provided that the radius of the activated complex is chosen in a suitable manner. There are also other examples where this is true. It was used by many investigators in the past, but in an entirely empirical way.<sup>6</sup>

In a series of recent papers<sup>7-10</sup> we have discussed the use of the Mean Spherical Approximation (MSA) in combination with a very simple asymmetric force model, which has a spherical repulsive core. The excellent correlations for the thermodynamic and kinetic parameters for a large number of solvents are more than mere coincidence. We believe that they are due to the fact that the MSA is not only a simple theory, but it also has some remarkable variational properties, that were discovered quite recently. Liquid state theories like the MSA and the HNC can be derived as variational problems of the free energy functional, which is written in terms of the Ornstein-Zernike direct correlation function. This view was introduced by Rosenfeld<sup>11-13</sup> for systems of hard objects in general, and hard spheres in particular. In that work the relation between the grand potential, the direct correlation function and the scaled particle theory was clearly shown. This analysis was extended to the case of charged hard spheres in general, and to the primitive model of electrolytes in particular.<sup>14,15</sup> It can be shown that the thermodynamics and

the pair correlation functions can be derived from a simple model in which the ions are replaced by charged shells and the dipoles by charged dipolar shells. We have shown that the direct correlation function in the MSA is just the electrostatic energy of the shells. Solving the MSA is thus equivalent to an electrostatics problem.

In the limit of high density and charge the Gibbs energy and the internal energy should diverge to the same order in the coupling parameter (charge/temperature ratio) while the entropy should diverge at a slower rate and therefore the Gibbs energy and the internal energy coincide in this limit. The mean spherical approximation (MSA) and the hypernetted chain approximation (HNC) satisfy these bounds, discovered by Onsager some fifty years ago. (This is not true for the Debye Hückel theory, where the entropy also diverges). Therefore, in this limit, the HNC and the MSA Gibbs energy, and its derivatives, coincide. This is a very gratifying feature, because the HNC, which is the more accurate theory, is in general difficult to solve numerically, while the MSA is analytical in most cases, and of a rather simple form.

In the asymptotic limit the excess electrostatic energy is identical to the exact Onsager lower bound, which is achieved by immersing the entire hard core system in an infinite neutral and perfectly conducting (liquid metal) fluid.<sup>14</sup> In the case of the primitive model of electrolytes, this is equivalent to smearing the charge on the surface of the ion. It was also shown<sup>14,16</sup> that the direct correlation function is equal to the interaction potential of two "smeared" ions of radius  $R_i$ ,  $R_j$  where  $R_i$  is the radius of ion  $i$  plus some screening length.

The Onsager process of introducing the infinite conductor, naturally decouples all the different components in the system which may differ in size, shape, charge distribution and relative orientation in space. As a result, the variational free energy functional in the high coupling limit diagonalizes, and the mathematical solution of the asymptotic problem is given in terms of the geometrical properties of the individual particles in the system.

The MSA Gibbs energy provides an appropriate functional from which the charge part of the direct correlation function is obtained by functional differentiation.

There is an actual numerical test case that proves the electrostatic analogy of the MSA. The very high charge and density limit for the one component plasma (point ions in a neutralizing background) was investigated using the HNC equation by Ng<sup>17</sup>, in a tour de force numerical calculation. The same results were obtained by Rosenfeld et al. using the analytical solution of the MSA.<sup>18</sup>

A second point is the use of spherical cores to represent non spherical objects. This also depends on the properties of interest. For the equation of state, it works even for very non spherical objects. There are numerous examples of sphericalization in the literature of molecular fluids. A notable example is given in the recent work of Williams et al.<sup>19</sup> The median diameter of a non spherical object yields a surprisingly good equation of state, not only for hard dumbbells but also for real gases at extremely high pressures.<sup>20,21</sup> Other examples are found in the extensive studies of Nezbeda and his group<sup>22,23</sup> on the structure of molecular liquids.

The analytical solution of the MSA for mixtures of ions and dipoles with a non spherical sticky potential was obtained some time ago<sup>24-26</sup> for the case of a sphere with a point dipole and a sticky potential of dipolar symmetry, which represents the specific non electrostatic interactions. In this work it was shown that the contribution of the sticky potential was merely a shift in the polarization parameter,  $\lambda$ . This work has been continued to include short ranged potentials of higher (tetrahedral) symmetry.<sup>27</sup>

In this spirit we have recently analyzed the solvation thermodynamics of monoatomic monovalent ions in water and in a number of polar solvents.<sup>7,10,28</sup> The basic idea comes from the observation that to a first approximation the corrections due to the molecular nature of the solvent result in a change in the screening length from  $1/r$  to  $1/r + \lambda\sigma_i/r \sigma_s$ , the shift being different for each ion, according to its size; because of geometrical packing considerations the number of solvent molecules that can be the first neighbors of a given ion,  $i$ , vary according to the ratio  $\sigma_i/\sigma_s$ . Consider now a realistic model of the solvent, with a soft repulsive core, and of non-spherical shape. Then, the distance of closest approach of a certain ion should be very different for the anion-solvent pair and the cation-solvent pair, even if the unsolvated anion and cation are of the

same size. The correct statical mechanical model for such a situation is the mixture with non-additive cores discussed in the literature.<sup>29</sup> A simple way of accounting for this is to assume that at low concentration the effective radius of the solvent is

$$r_{si} = r_s + \Delta_{si} \quad (1)$$

### The Model

In the MSA for a system of spherical ions and dipoles of arbitrary size<sup>30</sup>, the Gibbs energy of solvation in the limit of infinite dilution is<sup>31</sup>

$$G_{id} = - \frac{N_0(z_i e_0)^2}{8\pi \epsilon_0} \left(1 - \frac{1}{\epsilon_s}\right) \frac{1}{r_i + \delta^0} \quad (2)$$

Here,  $z_i$  is the valence of the ion,  $e_0$ , the fundamental charge,  $\epsilon_s$ , the static dielectric constant of the solvent,  $\epsilon_0$ , the permittivity of free space,  $r_i$ , the radius of the ion, and  $N_0$ , the Avogadro constant.  $\delta^0$  is equal to  $r_s/\lambda$ , where  $r_s$  is the radius of the solvent (the effective radius) and the MSA polarization parameter,  $\lambda$  is calculated from the dielectric constant of the pure solvent  $\epsilon_s$  using the Wertheim relationship<sup>32</sup>

$$\lambda^2 (\lambda + 1)^4 = 16 \epsilon_s \quad (3)$$

Although this value of the polarization parameter may be appropriate for estimating dipole-dipole interaction energy, it gives poor estimates of the Gibbs solvation energy when this is assumed to equal  $G_{id}$ .<sup>7,10,28</sup> Therefore, and in view of the above discussion we change eq. (2) to

$$G_{id}(C^+) = - \frac{N_0 (z_i e_0)^2}{8\pi \epsilon_0} \left(1 - \frac{1}{\epsilon_s}\right) \frac{1}{(r_i + \delta_c)} \quad (4)$$

where

$$\delta_c = \frac{r_s + \Delta_{sc}}{\lambda} \quad (5)$$

and for cations, and similarly, for monovalent monoatomic anions

$$G_{id}(A^-) = - \frac{N_0 (z_i e_0)^2}{8\pi \epsilon_0} \left(1 - \frac{1}{\epsilon_s}\right) \frac{1}{(r_i + \delta_A)} \quad (6)$$

where

$$\delta_A = \frac{r_s + \Delta_{sA}}{\lambda} \quad (7)$$

This separation and parameterization produces a much more sensible and physical model of  $G_{id}$  for ions in a number of solvents.

The second contribution to the Gibbs solution energy comes from repulsive dipole-dipole interactions<sup>25,33</sup> and is given by the equation

$$G_{dd} = \frac{(z_i e_0)^2 (\epsilon_s - 1)^2}{64\pi \epsilon_0 \epsilon_s (r_i + \delta_i)^2} \left[ \frac{4r_i + \delta_i \left( \frac{3\lambda + 2}{\lambda + 1} \right)}{\epsilon_s + \frac{\lambda(\lambda + 3)}{2(\lambda + 1)^2}} \right] \quad (8)$$

This expression may be simplified considerably when one considers the range of values typical for  $\lambda$ . For water whose dielectric constant is 78.3 at 25 °C,  $\lambda$  is equal to 2.65. The corresponding value of the ratio  $\lambda(\lambda+3)/2(\lambda+1)^2$  which appears in the denominator of the term in square brackets in eq. (8) is 0.56. This is negligible with respect to the value of  $\epsilon_s$  and may be neglected. The ratio appearing in the numerator of this term, namely,  $(3\lambda+2)/(\lambda+1)$  is equal to 2.7. Thus, the expression for  $G_{dd}$  may be rewritten as

$$G_{dd} = \frac{N_0(z_i e_0)^2}{64\pi \epsilon_0} \left( 1 - \frac{1}{\epsilon_s} \right)^2 \frac{(4r_i + 2.7 \delta_i)}{(r_i + \delta_i)^2} \quad (9)$$

The simplified expression has the advantage that the parameter  $\lambda$  does not appear on its own but only in  $\delta_i$ . This fact is important in applying the expression to experimental data. Finally, on the basis of previous work<sup>25</sup>, only a fraction  $\theta$  of the dipole-dipole term is used in estimating the Gibbs energy of solvation so that the final expression for  $G_s$  becomes

$$G_s = G_{id} + \theta G_{dd} \quad (10)$$

The expressions for the corresponding entropies are obtained from the temperature derivatives of the above Gibbs energies. For the ion-dipole term, the result for a cation is



$$S_{id}(C^+) = \frac{N_0(z_i e_0)^2}{8\pi \epsilon_0} \left[ \frac{1}{\epsilon_s^2} \frac{d\epsilon_s}{dT} \frac{1}{(r_i + \delta_c)} - \frac{1}{(r_i + \delta_c)^2} \left(1 - \frac{1}{\epsilon_s}\right) \frac{d\delta_c}{dT} \right] \quad (11)$$

where  $d\delta_c/dT$  is the temperature derivative of  $\delta_c$  which is obtained from the temperature derivative of  $\lambda_c$ . Comparing eqs. (4) and (11), it is easily shown that

$$S_{id}(C^+) \left(1 - \frac{1}{\epsilon_s}\right) = -\frac{G_{id}(C^+)}{\epsilon_s^2} \frac{d\epsilon_s}{dT} - \frac{8\pi \epsilon_0 G_{id}^2(C^+)}{N_0(z_i e_0)^2} \frac{d\delta_c}{dT} \quad (12)$$

The corresponding expression for an anion is

$$S_{id}(A^-) \left(1 - \frac{1}{\epsilon_s}\right) = -\frac{G_{id}(A^-)}{\epsilon_s^2} \frac{d\epsilon_s}{dT} - \frac{8\pi \epsilon_0 G_{id}^2(A^-)}{N_0(z_i e_0)^2} \frac{d\delta_A}{dT} \quad (13)$$

From these equations it follows that two additional parameters are required to estimate the entropy of solvation of a salt, namely, the temperature derivatives of the polarization parameters for the cation and anion.<sup>10,28</sup>

From the expression for  $G_{dd}$ , one obtains the following equation for the corresponding entropy:

$$S_{dd} = G_{dd} \left[ \frac{2}{\epsilon_s^2} \frac{d\epsilon_s}{dT} / \left(1 - \frac{1}{\epsilon_s}\right) + \frac{2.7 \delta_i}{\lambda} \frac{d\lambda}{dT} / (4r_i + 2.7 \delta_i) - \frac{2\delta_i}{\lambda} \frac{d\lambda}{dT} / (r_i + \delta_i) \right] \quad (14)$$

The derivative  $d\lambda/dT$  may be found from the Wertheim equation from which one obtains

$$\frac{d\lambda}{dT} = \left[ \frac{\lambda(1+\lambda)}{3\lambda+1} \right] \frac{1}{2\epsilon_s} \frac{d\epsilon_s}{dT} \quad (15)$$

Assuming that the fraction  $\theta$  is independent of temperature, the resulting expression for the entropy of solvation is

$$S_s = S_{id} + \theta S_{dd} \quad (16)$$

This model is now examined with respect to data for monoatomic monovalent ion solvation in water.

## Results and Discussion

In order to apply the above model to experimental data one must choose radii for the ions and an extrathermodynamic assumption for extracting single ion solvation energies from experimental values for the salts, namely, the alkali metal halides. Two sets of ionic radii give good fits between experiment and theory<sup>10</sup>, namely, the values given by Pauling<sup>34</sup> and those extracted from neutron and X-ray diffraction experiments.<sup>35</sup> The Pauling values are used here simply because they are more familiar. The extrathermodynamic assumptions used to separate experimental quantities into those for cations and anions were examined in detail for aqueous systems by Conway.<sup>36</sup> Values of the Gibbs energy and entropy of solvation for the alkali metal cations and halide anions estimated by Conway are summarized in Table 1 together with the Pauling radii for these ions.

In our previous analysis of the Gibbs solvation energy<sup>7,10</sup>, the contribution of dipole-dipole interactions was ignored ( $\theta = 0$  in eq. (10)). Analysis of the data for the alkali metal halides showed that the best values of  $\delta_c$  and  $\delta_A$  (eqs. (4) and (6)) assuming Pauling radii for the ions were 82.4 and 18.0 pm, respectively.<sup>10</sup> The quality of the fit between theory and experiment is really excellent, the standard deviation for the calculated values being 1.5 kJ mol<sup>-1</sup>, that is, significantly less than 1 percent. This result is illustrated in Fig. 1 in terms of a plot of  $-G_s$  against  $1/(r_i + \delta_i)$ . It is apparent that the values of  $G_s$  fit very well on a straight line in the region where data exist but that the slope of the best straight line through these data is slightly higher (75.2 kJ nm mol<sup>-1</sup>) than that predicted by theory (69.45 kJ nm mol<sup>-1</sup>). As a result, an ion with infinite radius is predicted to have a Gibbs solvation energy of 35.7 kJ mol<sup>-1</sup> when this simple model is applied. This can be attributed to the failure to account for dipole-dipole interactions which constitute a positive contribution to  $G_s$ .

The same data were re-examined on the basis of eq. (10) using the expression for  $G_{dd}$  given by eq. (9). The fit was carried out so that the resulting estimates of  $G_{id}$  and  $G_{dd}$  would go to zero for an ion of infinite radius. The resulting best values of  $\delta_c$  and  $\delta_A$  were much smaller than those obtained by the simpler analysis, and are recorded in Table 2. The corresponding estimates of  $G_{id}$  are plotted against  $1/(r_i + \delta_i)$  in Figure 2. As expected these estimates are larger in magnitude

than the experimental values of  $G_s$  also plotted on the same graph. Both plots are linear in  $1/(r_i + \delta_i)$  and have zero intercepts within the experimental standard deviation. Estimates of  $\theta G_{dd}$  for the same nine ions are also plotted in this figure. For the cations, this quantity varies from a high of 41.2 kJ mol<sup>-1</sup> for Li<sup>+</sup> to a low of 22.7 kJ mol<sup>-1</sup> for Cs<sup>+</sup>, and is approximately eight percent of the magnitude of  $G_{id}$ . In the case of the anions it represents a slightly higher fraction of the total Gibbs energy and varies from a high of 42.7 kJ mol<sup>-1</sup> for F<sup>-</sup> to a low of 30.8 kJ mol<sup>-1</sup> for I<sup>-</sup> ion. The values of  $\theta G_{dd}$  are also approximately linear in  $1/(r_i + \delta)$  with an intercept corresponding to zero for an infinitely large ion. The value of  $\delta$  used in estimating  $G_{dd}$  is that estimated on the basis of the Wertheim equation, namely, 51.6 pm. When the contributions,  $G_{id}$  and  $\theta G_{dd}$  are added one obtains estimates of  $G_s$  which agree with those obtained from experimental data to within a few kJ mol<sup>-1</sup>. Values of the parameters used in our calculations including the fraction  $\theta$  are summarized in Table 2.

A successful model for ionic solvation must also be able to estimate correctly the entropy of solvation. In order to make this calculation one must have estimates of the temperature derivatives of the parameters  $\delta_c$ ,  $\delta_A$  and  $\lambda$ . The latter was calculated using eq.(15) which gives a value of  $-2.48 \times 10^{-8} \text{ K}^{-1}$  for  $d\lambda/dT$  on the basis of the dielectric properties of water. Assuming  $r_s$ , the radius of a water molecule, is equal to 137 pm the corresponding value of  $d\delta/dT$  is 0.048 pm K<sup>-1</sup>. Values of  $\delta_c$  and  $\delta_A$  were determined from the single ion estimates of  $S_s$  given by Conway.<sup>13</sup> In the case of cations on the basis of eqs. (12) and (16), one may write

$$Y = \frac{d\delta_c}{dT} X \quad (17)$$

where

$$X = - \frac{8\pi \epsilon_0 G_{id}^2(C^+)}{N_0(z_i e_0)^2} \quad (18)$$

and

$$Y = S_s(C^+) \left(1 - \frac{1}{\epsilon_s}\right) + \frac{G_{id}(C^+)}{\epsilon_s^2} \frac{d\epsilon_s}{dT} + \theta S_{ss} \quad (19)$$

$S_c(C^+)$  being the solvation entropy of the cation. Similar equations can be written for anions using eqs. (13) and (16). A plot of  $Y$  against  $X$  for both ions is shown in Fig. 3. These data were fit to a straight line passing through zero (one parameter fit) as shown, the resulting slope which equals both  $d\delta_o/dT$  and  $d\delta_A/dT$  being  $0.032 \text{ pm K}^{-1}$ . It is important to note that the present analysis suggests that these temperature coefficients are equal in contrast to our previous conclusion based on an analysis which ignored dipole-dipole interactions. The estimated error in the slope is not large being about 4%. It should also be noted that the magnitude of  $d\delta_o/dT$  and  $d\delta_A/dT$  is about 65 percent of that found on the basis of the Wertheim equation (eq. (15)).

Using the above coefficients values of  $S_{id}$  were estimated for the alkali metal and halide ions and are plotted in Fig. 4 together with estimates of  $S_{dd}$  and experimental values of  $S_s$  against the experimental Gibbs solvation energy  $G_s$ . It is clear that the entropy of solvation is approximately a quadratic function of the Gibbs energy as suggested by the MSA model.

## References

1. Y. Marcus, *Ion Solvation*, J. Wiley, New York (1985).
2. J.E. Enderby and G.W. Neilson, *Rep. Progr. Phys.* **44**, 593 (1981).
3. M.P. Allen and D.J. Tildesley, *Computer Simulation of Liquids*, Oxford Science Publications, Oxford (1987).
4. R.A. Marcus, *J. Chem. Phys.* **24**, 966, 979 (1956).
5. D. Chandler, *Chimica Scripta*, **29A**, 61 (1989) and references therein.
6. J. O'M Bockris and A.K.N. Reddy, *Modern Electrochemistry*, Plenum Press, New York (1977), Vol. 1, Chapter 2, especially Sec. 2.2, and the references cited therein.
7. L. Blum and W.R. Fawcett, *J. Phys. Chem.* **96**, 408 (1992).
8. L. Blum, F. Vericat and W.R. Fawcett, *J. Chem. Phys.* **96**, 3039 (1992).
9. W.R. Fawcett and L. Blum, *Chem. Phys. Letters*, **187**, 173 (1991).
10. W.R. Fawcett and L. Blum, *J. Electroanal. Chem.* **328**, 333 (1992).
11. Y. Rosenfeld, *J. Chem. Phys.* **89**, 4272 (1988).
12. Y. Rosenfeld, *Phys. Revs. Lett.* **63**, 980 (1989).
13. Y. Rosenfeld, D. Levesque and J.J. Weis, *J. Chem. Phys.* **92**, 6818 (1990).
14. Y. Rosenfeld and L. Blum, *J. Phys. Chem.* **89**, 5149 (1985).
15. Y. Rosenfeld and L. Blum, *J. Chem. Phys.* **85**, 1556 (1986).
16. L. Blum and Y. Rosenfeld, *J. Stat. Phys.* **63**, 1177 (1991).
17. K.C. Ng, *J. Phys. Chem.* **61**, 2680 (1974).
18. Y. Rosenfeld, D. Levesque and J.J. Weis, *Phys. Rev.* **A39**, 3079 (1989).
19. G.O. Williams, J.L. Lebowitz and J.K. Percus, *J. Phys. Chem.* **88**, 6488 (1988).
20. E.A. Waisman, J.L. Lebowitz and D.A. MacGowan, *J. Chem. Phys.* **81**, 6096, 1984).
21. D.A. MacGowan, J.D. Johnson and M.S. Shaw, *J. Chem. Phys.* **82**, 3765 (1985).
22. I. Nezbeda, *Czech. J. Phys.* **30**, 601 (1980).
23. I. Nezbeda, W.R. Smith and S. Labik, *J. Chem. Phys.* **81**, 935 (1984).
24. L. Blum and D.Q. Wei, *J. Chem. Phys.* **87**, 555 (1987).

25. D.Q. Wei and L. Blum, *J. Chem. Phys.* **87**, 2999 (1987).
26. D.Q. Wei and L. Blum, *J. Chem. Phys.* **89**, 1091 (1988).
27. F. Vericat and L. Blum, unpublished results.
28. W.R. Fawcett and L. Blum, *J. Chem. Soc. Faraday Trans.*, in press.
29. J.L. Lebowitz and D. Zornick, *Phys. Revs.* xxxxxxxxx
30. L. Blum, *J. Stat. Phys.* **18**, 451 (1978).
- ~~32~~ ~~31~~. M.S. Wertheim, *J. Chem. Phys.* **55**, 4291 (1971).
- ~~31~~ ~~30~~. D.Y.C. Chan, D.J. Mitchell and B.W. Ninham, *J. Chem. Phys.* **70**, 2946 (1979).
- ~~33~~ ~~32~~. P. Garisto, P. Kusalik and G.N. Patey, *J. Chem. Phys.* **79**, 6294 (1983).
- ~~34~~ ~~33~~. L. Pauling, *J. Am. Chem. Soc.*, **49**, 765 (1927).
- ~~35~~ ~~34~~. Y. Marcus, *Chem. Rev.*, **88**, 1475 (1988)
- ~~36~~ ~~35~~. B.E. Conway, *J. Solution Chem.*, **7**, 721 (1978).