	ENTATION PAGE		Form Approved OMB No: 0704-0188
D-A263 363	timated to sverage 1 hour per response including the time for reviewing instructions, searching existing data sources no reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this his burden to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 jetterson the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503		
	PORT DATE 3. REPORT TYPE AND DATE 1 December 1992 Technical 6/1/9		192-5/31/93
What is Underneath? Movir	g Atoms and Molecules to F	ind Out	JNDING NUMBERS N00014-91-J-1630
UTHOR(S) P. S. Weiss and D. M. Eigler	DTIC		
ERFORMING ORGANIZATION NAME(S Department of Chemistry 152 Davey Laboratory The Pennsylvania State Univ University Park, PA 16802	AND ADDRE (S) APR 3 0 19 Persity	93 D 8. PI	RFORMING ORGANIZATION PORT NUMBER Report #5
PONSORING (MONITORING AGENCY)	AME(S) AND ADDRESS(ES)	10.5	PONSORING (MONITORING
Office of Naval Research Chemistry Program 800 N. Quincy Street Alexandria, VA 22217-500	0	Â	GENCY REPORT NUMBER
PREMENTARY NOTES Prepared for publication in M B. Vu Thien, N. Garcia and I	anipulations of Atoms in High K. Dransfeld, editors. (Kluwe	Fields and Tempe or Academic, 1993	ratures: Applications,
DISTRIBUTION / AVAILABILITY STATES Approved for public release. Distribution unlimited.	anipulations of Atoms in High K. Dransfeld, editors. (Kluwe	n Fields and Tempe or Academic. 1993	Pratures: Applications,
SUPPLEMENTARY NOTES Prepared for publication in M B. Vu Thien, N. Garcia and I DISTRIBUTION / AVAILABILITY STATES Approved for public release. Distribution unlimited.	anipulations of Atoms in High K. Dransfeld, editors. (Kluwe	a Fields and Tempe or Academic, 1993	pratures: Applications,
SUPPLEMENTARY NOTES Prepared for publication in M B. Vu Thien, N. Garcia and I DISTRIBUTION AVAILABILITY STATES Approved for public release. Distribution unlimited. Approved for public release. Distribution unlimited. We have used the scanning tu and Xe clusters in order to co initiation of the growth of a X This capability has also allowed the effect that these sites have 933	anipulations of Atoms in High C. Dransfeld, editors. (Kluwe MENT Inneling microscope (STM) to letermine their initial adsorp Xe surface overlayer, and to i ed us not only to examine the b e on the STM images.	p Fields and Tempe or Academic, 1993 12b. 12b. 12b. 12b. 12b. 12b. 12b. 12b.	such as benzene molecules such as benzene molecules is allowed us to image the twhich this growth begins. adsorbates, but also to learn
SUPPLEMENTARY NOTES Prepared for publication in M B. Vu Thien, N. Garcia and I DISTRIBUTION AVAILABILITY STATES Approved for public release. Distribution unlimited. ABSTRACT (Maximum 200 words) We have used the scanning tu and Xe clusters in order to continuation of the growth of a 2 This capability has also allowe the effect that these sites have UBJECT TERMS	anipulations of Atoms in High C. Dransfeld, editors. (Kluwe MENT Inneling microscope (STM) to letermine their initial adsorp Ke surface overlayer, and to i ed us not only to examine the b e on the STM images.	p Fields and Tempe or Academic, 1993 12b. 12b. 12b. 12b. 12b. 12b. 12b. 12b.	such as benzene molecules as allowed us to image the twhich this growth begins. adsorbates, but also to learn
SUPPLEMENTARY NOTES Prepared for publication in M B. Vu Thien, N. Garcia and I DISTRIBUTION AVAILABILITY STATES Approved for public release. Distribution unlimited. ABSTRACT (Maximum 200 words) We have used the scanning tu and Xe clusters in order to continuation of the growth of a X This capability has also allowed the effect that these sites have UBJECT TERMS	anipulations of Atoms in High C. Dransfeld, editors. (Kluwe MENT Inneling microscope (STM) to letermine their initial adsorp Ke surface overlayer, and to i ed us not only to examine the b e on the STM images.	p Fields and Tempe or Academic, 1993 12b. 0 move adsorbates tion site. This had dentify the sites at binding sites of the 93- 11111111111111111111111111111111111	such as benzene molecules as allowed us to image the twhich this growth begins. adsorbates, but also to learn -09120
SUPPLEMENTARY NOTES Prepared for publication in M B. Vu Thien, N. Garcia and I DISTRIBUTION AVAILABILITY STATES Approved for public release. Distribution unlimited. ABSTRACT (Maximum 200 words) We have used the scanning tu and Xe clusters in order to c initiation of the growth of a X This capability has also allows the effect that these sites have UBJECT TERMS IECURITY CLASSIFICATION 18. SEC DE REPORT	anipulations of Atoms in High C. Dransfeld, editors. (Kluwe MENT Inneling microscope (STM) to letermine their initial adsorp Ce surface overlayer, and to i ed us not only to examine the b e on the STM images. URITY CLASSIFICATION 19. SECU THIS PAGE	prove adsorbates tion site. This had dentify the sites at binding sites of the 93-	such as benzene molecule: such as benzene molecule: is allowed us to image the t which this growth begins adsorbates, but also to learr -09120

•

Prescribed by ANSI Std 239-18

OFFICE OF NAVAL RESEARCH

Grant #N00014-91-J-1630 R&T Code 413s002

Technical Report #5

What is Underneath? Moving Atoms and Molecules to Find Out

by

P. S. Weiss and D. M. Eigler

Department of Chemistry 152 Davey Laboratory The Pennsylvania State University University Park, PA 16802 and IBM Almaden Research Center 650 Harry Road San Jose, CA 95120

Prepared for publication in

Manipulations of Atoms in High Fields and Temperatures: Applications B. Vu Thien, N. Garcia, and K. Dransfeld, eds.

1 December 1992

Reproduction in whole, or in part, is permitted for any purpose of the United States Government.

This document has been approved for public release and sale: its distribution is unlimited.

WHAT IS UNDERNEATH? MOVING ATOMS AND MOLECULES TO FIND OUT

P. S. WEISS^{*†} and D. M. EIGLER* *IBM Research Division Almaden Research Center 650 Harry Road San Jose, CA 95120 USA

[†]Department of Chemistry The Pennsylvania State University University Park, PA 16802 USA

Accesi	on For	
NTIS DTIC Unarin Justific	CRA&I TAB ounced	2 0 0
By Distrib	ution /	
A	vailability (Codes
Dist A-1	Avail and Specia	l or

We have used the scanning tunneling microscope (STM) to move adsorbates such as benzene molecules and Xe clusters in order to determine their initial adsorption site. This has allowed us to image the initiation of the growth of a Xe surface overlayer, and to identify the sites at which this growth begins. This capability has also allowed us not only to examine the binding sites of the adsorbates, but also to learn the effect that these sites have on the STM images.

1. Introduction

Special surface sites such as steps and defects are thought to play an important and often dominant role in surface chemistry and in other surface processes such as diffusion [1]. We have previously shown how Xe atoms at dilute coverage first fill adsorption sites adjacent to steps, then form clusters on the terraces of the Pt{111} surface [2]. Using the STM tip to move the clusters allows subsequent imaging of the substrate sites at which the cluster islands nucleated and grew. In addition, we have shown how the appearance of an adsorbate in an STM image can vary according to binding site [3,4]. While this provides useful information on how electronic structure varies according to chemical environment, it also places constraints on the use of the STM in identifying molecules on surfaces. Moving the adsorbates once again allows examination of the initial binding sites.

2. Experimental

The experiments described here were conducted with an ultrahigh vacuum (UHV) scanning tunneling microscope at 4K. This microscope is equipped with a room temperature ultrahigh vacuum preparation and analysis chamber and a load lock. A schematic of the instrument is shown in Fig. 1. The STM is housed in a UHV chamber which itself is in an



Figure 1. Experimental set up showing the ultrahigh vacuum low temperature scanning tunneling microscope and associated vacuum chambers. The key is as follows: AES – Auger electron spectrometer, He(l) – liquid He dewar, He(g) – exchange gas chamber, I/V – electrometer, LEED – low energy electron diffractometer, STM – low temperature scanning tunneling microscope, Table – custom laser table supporting the instrument, and Translators – vertical and horizontal sample translators.

exchange gas chamber containing ca. 1 torr of He gas. The exchange gas chamber is immersed in a liquid He storage dewar with a five day capacity. We have shown using Johnson noise thermometry of the STM tunnel junction that this efficiently cools the STM to the temperature of the liquid He, *i.e.* to $T=4.5\pm0.5K[5]$. The STM itself is a modified louse design. The room temperature UHV preparation and analysis chamber is equipped with Auger electron spectroscopy, low energy electron diffraction, mass spectroscopy, ion sputtering, multiple sample transport, and rf induction. electron beam, and resistive heating. Samples are dosed by bleeding gas through sapphire leak valves into the upper room temperature chambers with the crystal in one of the upper room temperature chambers or in the microscope at 4K. Of particular importance here is the stability of the low temperature STM which allows us to turn off the feedback loop controlling the STM tip-sample separation for hours at a time and have the tip remain over the same atomic site during this time.





Figure 2. Topographic STM images of a 40Åx40Å area showing a cluster of Xe atoms adsorbed on Pt{111} in their initial adsorption site in the left frame, and after moving the cluster to reveal a defect in the right frame. Both images were recorded with a tip bias of 2 mV and a tunneling current of 5 pA. Two gray levels are highlighted to outline the positions of the five Xe atoms.

3. Results

3.1. MOVING XE CLUSTERS

We have found that Xe atoms preferentially bind at step edges on $Pt\{111\}$ [2]. After these sites are filled, the Xe atoms nucleate into small islands on the $Pt\{111\}$ terraces. In order to identify the nucleation sites of these islands, we have moved the Xe islands from their initial adsorption sites [2]. Our procedure for moving the islands was:

- 1) Positioning the STM tip over the outer perimeter of the island.
- 2) Turning off the feedback loop.
- 3) Moving the tip in towards the surface by 2.6Å.
- 4) Changing the tip bias voltage of the tip to $V=^{+}0.010V$ for 10 msec.
- 5) Returning the bias voltage to its original value ($^{-}0.002$ to $^{-}0.010$ V).
- 6) Returning the tip to its original height above the surface.

This procedure resulted in centering the island about the STM tip with approximately 50% efficiency [7]. By repeating this procedure, an island could be moved completely away from its initial adsorption site. An example of this is shown in Fig. 2. In all cases when this was done for the Xe islands on the Pt{111} surface, it was found that a depression of ≤ 0.1 Å in the STM images lay underneath the initial island positions. We have tentatively identified these point defects as surface carbon atoms [2]. The concentration of the point defects tracks the surface C contamination as measured by Auger spectroscopy. Theoretical calculations also indicate that a C atom on this surface would appear as a depression in an STM image [6].

3.2. MOVING BENZENE MOLECULES

Benzene adsorbed on $Pt\{111\}$ appears to exhibit three different types of images depending upon adsorption site [3]. This result is consistent with theoretical calculations of images of benzene on graphite and MoS_2 [9], and the known variety of binding sites for benzene on



Figure 3. Topographic STM images of a 50Å x 50Å region of Pt{111} which has been covered with 0.001 monolayers of benzene. Both images were recorded with a tip bias of $^{-10}$ mV and a tunneling current of 100 pA. Each bump corresponds to a single benzene molecule. The arrow in the left frame indicates the position of a benzene molecule before and after being moved, as described in the text. The right frame shows the surface after the molecule has been moved. Note that the two images are shifted somewhat laterally.

Pt{111} [10,11]. We have used the STM to move benzene from one of these sites to show that benzene which appears simply as a bump for low bias conditions does not sit at a defect. This is shown in Fig. 3. The left panel shows several benzene molecules as imaged initially. Then, one of the benzene molecules was lifted off the surface using the tip and redeposited back on the surface elsewhere (as shown by the arrow in the left panel). The right panel shows both the final position of the benzene molecule which has been moved and the substrate surface at this molecule's adsorption site. There does not appear to be a defect at this site. This is consistent with our interpretation of the different sites which we believe lead to the different images for benzene on this surface [3]. The procedure that we used to lift and replace the molecule was simply shutting off the feedback loop controlling the STM tip-sample separation, moving the STM tip in towards the surface 2.5Å for 50 msec, then pulling the tip back out and turning the feedback loop on.

While we were able to remove a number of molecules from the surface, this was the only case where we successfully dropped off the molecule afterwards. Still, by attempting to drop off the adsorbate far from the initial adsorption site so as to diminish the possibility of affecting this site, or by imaging with the molecule on the tip, the initial molecular adsorption site can be determined.

4. Conclusions

We have shown how it is both possible and useful to use the scanning tunneling microscope to move adsorbates so as to determine their surface adsorption sites. Knowledge of adsorption sites allows us to understand aspects of film growth and adsorption as well as the STM imaging process itself.

Acknowledgments

The authors would like to thank Norton Lang for helpful discussions, Joe Schlaegel and Les Harlow for help in the construction of the apparatus, and Stephan Stranick for help in preparing the figures. One of us (PSW) would like to acknowledge support from the National Science Foundation, the Office of Naval Research, and the Petroleum Research Fund (administered by the American Chemical Society).

References

- [1] e.g. G. A. Somorjai, <u>Chemistry in Two Dimensions</u> (Cornell University Press, Ithaca, New York, 1981).
- [2] P. S. Weiss and D. M. Eigler, 'Adsorption and Accommodation of Xe on Pt{111},' Phys. Rev. Lett. 69, 2240 (1992).
- [3] P. S. Weiss and D. M. Eigler, 'Site Dependence of the Apparent Shape of a Molecule in STM Images: Benzene on Pt{111},' to be published.
 [4] J. A. Stroscio and D. M. Eigler, 'Atomic and Molecular Manipulation with the
- [4] J. A. Stroscio and D. M. Eigler, 'Atomic and Molecular Manipulation with the Scanning Tunneling Microscope,' Science 254, 1319 (1991).
- [5] P. S. Weiss and D. M. Eigler, unpublished.
- [6] N. D. Lang, private communication.
- [7] N.B. A more efficient procedure was subsequently found for moving single Xe atoms on Pt{111} and Ni{110} [8].
- [8] D. M. Eigler and E. K. Schweizer, 'Positioning Single Atoms with a Scanning Tunneling Microscope,' *Nature* 344, 524 (1990); D. M. Eigler, E. K. Schweizer, C. Lutz, and P. Zeppenfeld, unpublished.
- [9] A. J. Fisher and P. E. Blöchl, 'Adsorption and STM Imaging of Benzene on Graphite and MoS₂,' to be published.
- [10] C. M. Mate, Ph. D. Thesis, 'A Molecular Surface Science Study of Adsorbates on Surfaces: Importance to Lubrication,' University of California, Berkeley (1986).
- [11] A. Wander, G. Held, R. Q. Hwang, G. S. Blackman, M. L. Xu, P. de Andres, M. A. Van Hove, G. A. Somorjai, 'A Diffuse LEED Study of the Adsorption Structure of Disordered Benzene on Pt{111},' Surface Sci. 249, 21 (1991).