

93-12439



A

JUN 03 1993

DTIC
ELECTE
D

Journal of Medical Virology 39:152-155 (1993)

2

Dobrava Virus as a New Hantavirus: Evidenced by Comparative Sequence Analysis

Shu-Yuan Xiao, Gordana Diglisic, Tatjana Avsic-Zupanc, and James W. LeDuc

Disease Assessment Division, U.S. Army Medical Research Institute of Infectious Diseases, Frederick, Maryland (S.-Y.X., G.D., J.W.L.); Institute of Microbiology, Medical Faculty of Ljubljana, Ljubljana, Slovenia (T.A.-Z.)

Dobrava virus, recently isolated from a yellow-neck mouse (*Apodemus flavicollis*), captured in a northern Slovenian village where severe cases of hemorrhagic fever with renal syndrome were recognized, was shown by serology and restriction enzyme digestion of PCR-amplified gene segments to be related to previously recognized hantaviruses. To investigate further the relationship of this new isolate to other hantaviruses, a portion of the medium (M) genome segment of Dobrava virus was amplified by PCR and the nucleotide sequence determined. Comparing the nucleotide sequence with the same gene region of other hantaviruses revealed an overall homology of 41.7%. A phylogenetic tree based on pairwise sequence similarity clearly showed that Dobrava virus is genetically distinct, and probably represents a new virus in the genus *Hantavirus* of the family Bunyaviridae. © 1993 Wiley-Liss, Inc.

KEY WORDS: Hantaan virus, hemorrhagic fever with renal syndrome, Slovenia, nucleotide sequence, RT-PCR

INTRODUCTION

The genus *Hantavirus* is comprised of at least 6 genetically and antigenically related but distinct viruses: Hantaan (HTN), Seoul (SEO), Puumala (PUU), Prospect Hill (PH), Thailand (THAI), and Thottapalayam (THP) [Lee et al., 1985; Schmaljohn et al., 1985; Xiao et al., 1992]. Some of these viruses are etiologic agents of the human disease, hemorrhagic fever with renal syndrome (HFRS). Recently, a new isolate, designated Dobrava virus, was obtained from a yellow-necked mouse, *Apodemus flavicollis*, captured in Slovenia [Avsic-Zupanc et al., 1992], where severe cases of HFRS were identified. Preliminary serological studies and polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) analysis demonstrated its relationship with the existing hantaviruses, suggesting that it might represent a new member of the genus *Hantavirus*, family Bunyaviridae [Avsic-Zupanc et al., 1992].

We now report the nucleotide sequence comparison of a PCR-amplified fragment from the M genomic segment of Dobrava virus with previously published sequences of other hantaviruses. Dobrava virus has found to be genetically distinct from other hantaviruses, adding further evidence that Dobrava represents a new hantavirus.

MATERIALS AND METHODS

Virus and Cell Culture

Dobrava virus was isolated from the lungs of an *Apodemus flavicollis* [Avsic-Zupanc et al., 1992]. The original specimens were collected from Dobrava village in Slovenia, where severe cases of HFRS were identified. Vero E6 cells were infected with the virus and processed for total cellular RNA extraction by using a guanidine isothiocyanate/cesium chloride method, as described previously [Xiao et al., 1991].

Reverse Transcriptase Polymerase Chain Reaction (RT-PCR) and Direct Sequencing of the Amplifier

Two primer sets used in the initial PCR amplification, HanG2-1:2 and HanG2-3:4, were derived from the glycoprotein G2 encoding region of the Hantaan virus M genome segment [Xiao et al., 1991]. The sizes of the expected amplifiers were 411 and 334 bp, respectively. RT-PCR reactions were done as described previously [Xiao et al., 1991]. Briefly, total cellular RNA was copied to cDNA with avian myeloblastosis virus reverse transcriptase, then amplified for 35 cycles with the *Taq* polymerase, in an automatic DNA Thermal Cycler (Perkin-Elmer-Cetus, Norwalk, CT).

The amplified product was subjected to preparative electrophoresis by using 1.2% low melting point (LMP) agarose gel (IBI, New Haven, CT) prepared with Tris-acetate-EDTA (TAE) buffer. The target band was cut out and DNA recovered by using the GENECLEAN Kit (BIO 101, La Jolla, CA). Ten microliters of double-stranded DNA (1.5 µg) was denatured with 2 µl of 2 N

Accepted for publication July 10, 1992.

Address reprint request to Dr. S.-Y. Xiao, Disease Assessment Division, U.S. Army Medical Research Institute of Infectious Diseases, Bldg. 1425, Fort Detrick, Frederick, MD 21702-5011

© 1993 WILEY-LISS, INC.

This document has been approved for public release and sale; its distribution is unlimited.

AD-A265 384



4

U

V

D

R




Fig. 1. Identification of cognate region for sequence comparison and dendrogram construction. Sequences were aligned against that of Hantaan 76-118 (Hash value = 6).

NaOH, ethanol precipitated, and dried briefly. Sequencing was done by the dideoxy chain termination method [Sanger et al., 1977], with the Sequenase Version 2.0 DNA Sequencing Kit from United States Biochemical (Cleveland, OH). Primers HanG2-1 and HanG2-2 were used for the sequencing reactions.

Sequence Analysis

Reference sequences were previously published and deposited in GenBank. Sequences examined included: Hantaan virus strains 76-118, Lee, and Hojo [Schmaljohn et al., 1987, 1988; Seoul virus strains HR8039 and SR-11 [Antic et al., 1991; Arikawa et al., 1990]; Puumala virus strains Sotkamo and Hällnäs-B1 [Giebel et al., 1989; Vapalahti et al., 1992]; Prospect Hill virus strain PHV-I [Parrington et al., 1991]; and Thailand virus strain Thai749 (Y-K. Chu et al., unpublished data). The Dobrava virus sequence was compared with all of these by the MacVector computer program from International Biotechnologies, Inc. (New Heaven, CT), to identify homologous regions in each sequence. The resulting sequence fragments were then compared with each other by using the CLUSTAL program of the PC/Gene software from IntelliGenetics, Inc. (Mountain View, CA). Pairwise similarity scores were calculated for all pairs of sequences and the degree of similarity used to form a tree (dendrogram) by clustering the sequences based on overall similarity.

RESULTS

Dobrava virus was successfully amplified with the HanG2-1:2, but not with the HanG2-3:4 primer pair, while both primer pairs amplified Hantaan virus strain 76-118 (data not shown). The amplicon generated with HanG2-1:2 was processed for sequencing. A sequence of 207 bp from the HanG2-1:2 amplicon was obtained and aligned against the whole M segment sequence of Han-

taan virus strain 76-118, along with those of 8 other hantaviruses. The cognate region was identified for each sequence as shown in Figure 1, corresponding to the region 2120 to 2330 in the Hantaan virus cDNA sequence. This region was extracted from all the sequences for further pairwise sequence comparison by the CLUSTAL program.

Alignment of the 10 nucleotide sequences is shown in Figure 2. A dendrogram was constructed based on the calculated scores of pairwise similarity (Fig. 3). The length of compared region was 211 bp, with an overall identity of 88 bp (41.7%). To ensure the reproducibility of this dendrogram without Dobrava virus, another 500 bp region from the glycoprotein G1 encoding gene of the completely sequenced strains was analyzed with the same strategy. An identical dendrogram, excluding Dobrava virus, resulted.

DISCUSSION

Seroepidemiological investigations have identified at least three distinct antibody reaction patterns among HFRS patients seen in Yugoslavia and Slovenia [Avsic-Zupanc et al., 1990; Gligic et al., 1992]. These include reactions specific for a virus similar or identical to prototype Hantaan virus, a second pattern specific for Puumala virus, and a third pattern that appears to be Dobrava virus-specific. In addition, examining Dobrava virus with monoclonal antibodies to various hantaviruses demonstrated a unique serological reactivity pattern for this virus. Finally, PCR-RFLP comparisons of Dobrava and other recognized hantaviruses likewise found that this virus is related to, yet distinct from, all previously recognized hantaviruses [Avsic-Zupanc et al., 1992].

The results show that at the genomic level, Dobrava virus shared homology with other hantaviruses. However, it was clearly distinct when the sequence of the

DOBRAVA	CCCATTAAATCAAGAACAATCAGTAGATCTTCACATAGAGATAAGGAGTCA	50
HTN	CCACTTGAGGAAGCACAAATCCATTGACCTACATATTGAAATAGAAGAACA	50
LEE	CCACTTGAAGAAGCACAAATCCATTGATCTACATATTGAAATAGAAGAACA	50
HOJO	CCACTTGAAG---CACAATCCATTGATCTACATATTGAAATAGAAGAACA	47
SR11	CCAGTTAATGACCAACAGAGTGTCTCATTGCATATAGAAATTGAAAGTCA	50
SEOUL	CCAGTTAATGACCAACAGAGTGTCTCATTGCACATAGAAATTGAAAGTCA	50
THAI749	CCAGTTAACACAGAACAAGGAGTTCATACATATAGAGATTGAAGAGCA	50
HALLNAS	CCAGCAAACGAACAAGAGAAAATCCCATTTCATCTGCAGTTAAGCAAACA	50
SOTKAMO	CCGGCAAATGAACAAGAAAAATACCATTTCATTTACAGATAAGTAAACA	50
PHV	---GCAAATGAGCAAAGAAATCCCTTTTCTTCCAGCTAGAGCGTCA	47
	. . . : : . . . : : : : : : : : : : . . . : . . . :	
DOBRAVA	AGGATTTT-CACAAGTGTTCATGCATTAGGTTCATTTGGTTTGGTGA-GAC	98
HTN	GACAAATGGGTGTTGATGTGCATGCTCTAGGACACTGGTTTGGTGGCGTC	100
LEE	AACAATGGGTGTAGATGTGCATGCTCTAGGACACTGGTTTGGTGGCGTC	100
HOJO	AACAATGGGTGTGGATGTACATGCTTATAGGACACTGGTTTGGTGGCGTC	97
SR11	AGGCATTGGTGTCTGCTGTTTCATCATCTTGGACATTGGTATGATGCAAGAT	100
SEOUL	AGGCATTGGTGTCTGATGTCCATCATCTTGGACATTGGTATGATGCAAGAT	100
THAI749	AGGAATGGTGTAGATGTCCATCACCTAGGCCATTGGTTTGGTGGCCGGC	100
HALLNAS	AGTGATTTCATGCAGAGATCCAGCAATTTAGGTTCATTGGATGGATGCTACAT	100
SOTKAMO	GGTGTTCATGCTGAAATTCACATCTAGGCCACTGGATGGATGCTACAT	100
PHV	GGTGTTCATGCAGAGATTCAAATTTAGGCCACTGGATGGATGGAACAT	97
	. . . : : : : . . . : : : : : : : : :	
DOBRAVA	TTAATTTAAAGACATCTTTTTCATTGTTATGGTGCATGCTACTAAGTATGAA	148
HTN	TTAACCTTAAAACATCCTTTTTCATGTTATGGGCTTGTACAAAAGTATGAA	150
LEE	TTAACCTTAAAACATCCTTTTTCATGTTATGGTGTGTTACAAAAGTATGAA	150
HOJO	TTAACCTTAAAACATCCTTTTTCATGTTATGGTGTGTTACAAAAGTATGAA	147
SR11	TGAATCTAAAACCTCATTTCATTGTTATGGTGCCTGCACAAAATATCAA	150
SEOUL	TGAATTTAAAACCTCATTTCATTGTTATGGTGCCTGCACAAAATATCAA	150
THAI749	TTAATTTAAAACATCATTCCATTGTTACGGGGCATGCTCAAAGTACCAG	150
HALLNAS	TTAATCTTAAAACCTGATTTTCATGCTATGGCTCATGTGAAAGTATGCT	150
SOTKAMO	TTAATCTAAAACCTGCTTTTTCATGTTATGGTTCATGTGAAAATATGCC	150
PHV	TTAATCTGAAAACATCATTCCATTGTTACGGTGTCTTGTGAAAAGTATGCT	147
	. . . : : : : : . . . : : : : : : : : : :	
DOBRAVA	TATCCTTGGCATAACGCAAAATGCCACTTTGAAAGGGATTTCGAGTATGA	198
HTN	TACCCTTGGCATACTGCAAAAGTGCCATTATGAAAGAGATTACCAATATGA	200
LEE	TACCCTTGGCATACTGCAAAAGTGCCACTATGAAAGGGATTACCAATATGA	200
HOJO	TACCCTTGGCATACTGCAAAAGTGCCACTATGAAAGGGATTACCAATATGA	197
SR11	TACCCATGGCACACTGCAAAATGCCATTTTGAGAAAAGATTATGAGTATGA	200
SEOUL	TATCCATGGCACACTGCAAAATGCCATTTTGAGAAAAGATTATGAGTATGA	200
THAI749	TACCCTTGGCATAACGCAAGTGTCACTTTGAAAAAGATTATGAATATGA	200
HALLNAS	TATCCTTGGCAGACAGCAGGTGTTTTCATAGAAAAGATTATGAATATGA	200
SOTKAMO	TACCCCTGGCAAAACAGCAGGATGTTTCGTGAAAAAGATTATGAATATGA	200
PHV	TATCCGTGGCAAACTGCAAAATGTTCTTAGAGAAAAGATTATGAATTTGA	197
	. . . : : : : : : : : : : : : :	
DOBRAVA	GAACA--TGGG	207
HTN	GACGAGCTGGG	211
LEE	GACAAGCTGGG	211
HOJO	GACAAGCTGGG	208
SR11	AAATAGCTGGG	211
SEOUL	AAATAGCTGGG	211
THAI749	AAATAGTGGG	211
HALLNAS	GACTGGTGGG	211
SOTKAMO	AACTGGTGGG	211
PHV	AAAGGGTGGG	208
	. . . : : : :	

Fig. 2. Multiple sequence alignment of the homologous region. The dash signs represent gaps.

PCR amplified segment is compared with other hantaviruses. This finding, combined with previously published observations of serological uniqueness, suggests that Dobrava is yet another distinct member of the genus.

Construction of phylogenetic trees based on nucleotide sequence similarity has been used to depict rela-

tionships of closely related viruses as an alternative to the traditional serotyping of viruses. Rico-Hesse et al. [1987], comparing nucleotide sequences of 150 bp, showed that this method of comparison was able to establish epidemiologically consistent relationship among widely distributed isolates of wild poliovirus type 1 genotypes. From the dendrogram constructed in




Fig. 3. Dendrogram based on the pairwise comparison of the hantavirus sequences. The horizontal distance to the node that connect 2 strains is proportional to the nucleotide sequence divergence between them.

this study, Dobrava virus was found to be most closely related to isolates of Hantaan, Seoul, and Thailand viruses; whereas it was more distantly related to isolates of Puumala and Prospect Hill viruses (Fig. 3). However, we showed it to be clearly separated from all of these viruses, in agreement with the serological results [Avcic-Zupanc et al., 1992]. Construction of a second dendrogram, based on the sequence analysis of a 500 bp segment from the G1 encoding region (excluding Dobrava virus), showed an identical relationship among these viruses, verifying the reliability of this partial sequencing analysis. In addition, similar relationships were recently reported by others who compared gene product sequences from the whole M segment of the hantaviruses with complete sequence information [Antic et al., 1992], further adding credence to our proposed placement of Dobrava virus.

The nucleotide sequence is the most reliable marker in identifying a virus; however, complete sequencing of all new isolates is neither practical nor necessary for this purpose. Data from this study showed that the overall interrelationship of hantaviruses documented from a region of approximately 211 bp is very similar (if not identical) to that from the whole M genome segment of hantavirus [Antic et al., 1992]. This provides a very useful and rapid tool for virus classification or typing, with the easy availability of purified DNA template generated by PCR.

ACKNOWLEDGMENTS

This work was carried out when Dr. S.-Y. Xiao held a senior research associateship from the National Research Council of the U.S.A.

REFERENCES

- Antic D, Lim B-U, Kang CY (1991): Molecular characterization of the M genomic segment of the Seoul 80-39 virus: Nucleotide and amino acid sequence comparisons with other hantaviruses reveal the evolutionary pathway. *Virus Research* 19:47-58.
- Antic D, Kang CY, Spik K, Schmaljohn C, Vapalahti O, Vaheri A (1992): Comparison of the deduced gene products of the L, M and S genome segments of hantaviruses. *Virus Research* 24:35-46.
- Arikawa J, Lapenotiere HF, Iacono-Connors L, Wang M, Schmaljohn CS (1990): Coding properties of the S and the M genome segments of Sapporo rat virus: Comparison to other causative agents of hemorrhagic fever with renal syndrome. *Virology* 176:114-125.
- Avcic-Zupanc T, Likar M, Novakovic S, Cizman B, Kraigher A, van der Groen G, Stojanovic R, Obradovic M, Gligic A, LeDuc JW (1990): Evidence of the presence of two hantaviruses in Slovenia, Yugoslavia. *Archives of Virology* 115:87-94.
- Avcic-Zupanc T, Xiao S-Y, Stojanovic R, Gligic A, van der Groen G, LeDuc JW (1992): Characterization of Dobrava virus: A hantavirus from Slovenia. *Journal of Medical Virology* 37.
- Giebel LB, Stohwasser R, Zöller L, Bautz EKJ, Darai G (1989): Determination of coding capacity of the M genome segment of nephropathia epidemica virus Hällnäs B1 by molecular cloning and nucleotide sequence analysis. *Virology* 172:498-505.
- Gligic A, Stojanovic R, Obradovic M, Hlaca D, Dimkovic N, Diglisic G, Lukac V, Ler Z, Bogdanovic R, Antonijevic B, Ropac D, Avcic-Zupanc T, LeDuc JW, Ksiazek T, Yanagihara R, Gajdusek DC (1992): Hemorrhagic fever with renal syndrome in Yugoslavia: Epidemiologic and epizootologic features of a nation-wide outbreak in 1989. *European Journal of Epidemiology* (in press).
- Lee PW, Gibbs CJ Jr., Gajdusek DC, Yanagihara R (1985): Serotypic classification of hantaviruses by indirect immunofluorescent antibody and plaque reduction neutralization tests. *Journal of Clinical Microbiology* 22:940-944.
- Parrington MA, Lee P-W, Kang CY (1991): Molecular characterization of the Prospect Hill virus M RNA segment: A comparison with the M RNA segments of other hantaviruses. *Journal of General Virology* 72:1845-1854.
- Rico-Hesse R, Pallansch MA, Nottay BK, Kew OM (1987): Geographic distribution of wild poliovirus type 1 genotypes. *Virology* 160:311-322.
- Sanger F, Nicklen S, Coulson AR (1977): DNA sequencing with chain-terminating inhibitors. *Proceedings of National Academy of Sciences of USA* 74:5463-5467.
- Schmaljohn CS, Hasty SE, Dalrymple JM, LeDuc JW, Lee HW, von Bonsdorff C-H, Brummer-Korvenkontio M, Vaheri A, Tasi TF, Regnery HL, Goldgaber D, Lee P-W (1985): Antigenic and genetic properties of viruses linked to hemorrhagic fever with renal syndrome. *Science* 227:1041-1044.
- Schmaljohn CS, Schmaljohn AL, Dalrymple JM (1987): Hantaan virus M RNA: Coding strategy, nucleotide sequence, and gene order. *Virology* 157:31-39.
- Schmaljohn CS, Arikawa J, Hasty SE, Rasmussen L, Lee HW, Lee PW, Dalrymple JM (1988): Conservation of antigenic properties and sequences encoding the envelope proteins of prototype Hantaan virus and two virus isolates from Korean hemorrhagic fever patients. *Journal of General Virology* 69:1949-1955.
- Vapalahti O, Kallio-Kokko H, Salonen E-M, Brummer-Korvenkontio M, Vaheri A (1992): Cloning and sequencing of Puumala virus Sotkamo strain S and M segments: Evidence for strain variation in hantaviruses and expression of the nucleocapsid protein. *Journal of General Virology* 73:829-838.
- Xiao S-Y, Chu Y-K, Knauer FK, Lofts RS, Dalrymple JM, LeDuc JW (1992): Comparison of hantavirus isolates using a genus-reactive primer pair polymerase chain reaction. *Journal of General Virology* 73:567-573.
- Xiao S-Y, Yanagihara R, Godec MS, Eldadah ZA, Johnson BK, Gajdusek DC, Asher DM (1991): Detection of hantavirus RNA in tissues of experimentally infected mice using reverse transcriptase-directed polymerase chain reaction. *Journal of Medical Virology* 33:277-282.