

NAVAL HEALTH RESEARCH CENTER

AD-A271 748



**THE BATTLE FOR HUE:
CASUALTY AND DISEASE RATES DURING URBAN WARFARE**

C. G. Blood

M. E. Anderson

2 DTIC
S ELECTE NOV 02 1993 D
E

Report No. 93-16

93-26415



Approved for public release: distribution unlimited.

93 11 1 02

NAVAL HEALTH RESEARCH CENTER
P. O. BOX 85122
SAN DIEGO, CALIFORNIA 92186 - 5122



NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

The Battle for Hue: Casualty and Disease Rates During Urban Warfare

**Christopher G. Blood
Marlisa E. Anderson***

DTIC QUALITY IMPACTED 8

**Medical Information Systems and
Operations Research Department**

**Naval Health Research Center
P.O. Box 85122
San Diego, CA 92186-5122**

***GEO-Centers, Inc.**

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Report No. 93-16 , supported by the Naval Medical Research and Development Command, Department of the Navy, under Work Unit No. M0095.005-6204. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of Defense, nor the U.S. Government. Approved for public release, distribution unlimited. To be presented at the Tenth International Symposium on Military Operational Research (10 ISMOR) to be held at the Royal Military College of Science, Shrivenham, Wiltshire, UK 7-10 September 1993. The theme of the symposium is OPERATIONAL RESEARCH AND FUTURE INTERNATIONAL SECURITY OPERATIONS

SUMMARY

Problem

Renewed nationalism with the ending of the Cold War has precipitated numerous conflicts between regions or countries which were formerly united. Hostilities between some ethnic and nationalistic factions have reached the point where regional security is threatened and United Nations military operations are required.

Objective

Because some U.N. operations may require the forcible removal of entrenched factions from urban settings, the present investigation seeks to determine the levels of medical casualties that might be sustained during urban warfare.

Approach

Casualty rates and illness incidence were examined for U.S. Marine forces participating in the re-taking of the city of Hue during the Tet offensive in 1968. The casualty rates were analyzed for different phases of the urban assault and contrasted with a different period of the Vietnam Conflict, and with the high intensity battle for Okinawa during World War II.

Results

Rates of casualties during the re-taking of Hue were highest during the two phases of the operation which required close-quarter fighting. The house-to-house fighting south of the river yielded a wounded rate of 37.9 per 1000 strength per day, while the fighting in the Inner City yielded a rate of 44.4. Rate of wounded during the "mopping-up" phase was 5.8. The rate of illness incidence was stable over the month-long operation and showed no concomitant increase with battle intensity.

Conclusion

Rates of casualties during a U.N.-led military operation may likely be high. It is essential that medical resource requirements be anticipated and pre-positioned prior to the first casualties being sustained.

The Battle for Hue: Casualty and Disease Rates During Urban Warfare

Renewed nationalism with the ending of the Cold War has yielded an explosion in internecine strife between countries or regions which were once united. Hostilities between Armenia and Azerbaijan, Turkey and Kurdistan, Pakistan and India, rival clan leaders within Somalia, and Yugoslavia and the surrounding republics have all erupted with deadly consequences. At the same time, the world community has put increasing pressure on the United Nations (U.N.) to take on the roles of disarming troops and "enforcing" peace in addition to its traditional peacekeeping function of monitoring cease-fire lines.

When opposing sides are intransigent, peacemaking or peace-enforcing may require U.N.-led military troops to forcibly end hostilities and impose a settlement between warring factions. Further, the use of such military force will at times likely need to be applied to cities and villages where one faction is firmly established. Therefore, it is quite possible that U.S. Marines, in addition to their amphibious power projection and crisis response functions, may be called upon to participate in an U.N. operation in which a goal is to force an entrenched enemy from an urban setting.

The present investigation examines rates of casualties and disease incidence incurred by U.S. Marines in the retaking of the city of Hue during the Vietnam Conflict. Hue was the ancient imperial capital and cultural seat of Vietnam. The city was divided by the

Huong River into two main sections (See Figure 1)¹. The area south of the river was residential and also included City Hall, Hue University, the train station, a Military Assistance Command, Vietnam (MACV) compound and other administrative agencies. The area north of the river contained the walled portion of Hue known as the Citadel. The Citadel was surrounded by canals and moats and was the site of the imperial palace. The barrier around the Citadel formed a 2,500 meter square and consisted of an outer stone wall five meters high and one to three meters thick, as well as an inner wall separated by dirt fill. In addition to the imperial palace, the Citadel contained an airport, houses, markets, and a large number of buildings. The North Vietnamese Army (NVA) infiltrated the city and took control as part of the Tet Offensive at the end of January, 1968. The urban assault that followed required the use of three Marine infantry battalions and 13 South Vietnamese battalions and lasted for one month.^{2,3,4} Given this duration and requisite force, the Hue operation has applicability to estimating the levels of casualties which might be sustained in a U.N.-led military operation within an urban environment.

METHOD

Medical incidence data were extracted from Unit Diaries of the three U.S. Marine infantry battalions (1/1/1, 1/5/1, 2/5/1) which




Figure 1. Schematic of the city of Hue, Vietnam

participated in the Battle for Hue. Unit Diaries, which are housed at Records Service Section, Headquarters U.S. Marine Corps, Quantico, VA, include the number of wounded in action (WIA), killed in action (KIA), Disease/Nonbattle Injuries (DNBI), dates of the casualties, and unit strength. Daily rates of wounded, killed, and DNBI incurred during the re-taking of Hue were computed and contrasted with pre- and post-Hue operations. Casualty and DNBI rates were calculated per 1000 strength per day.

Additionally, rates of casualties and DNBI were examined for three separate phases of the operation in which U.S. Marines were involved: 1) Fighting south of the Huong River, 2) Fighting in the Inner Citadel, and 3) Pursuit and "mopping up". Rates during the Hue battle were also contrasted with casualty incidence of the four-month period of peak U.S. Marine involvement in Vietnam (May through August 1968), as well as with rates sustained during the high intensity assault on Okinawa (April through June 1945). Muster Rolls, from which the Okinawa data were extracted, are maintained at the National Archives, Washington D.C.

RESULTS

CASUALTY RATES

Figure 2 is a display of the casualty rates of participating infantry units before, during the Battle of Hue (January 31 - March 2), and after the city was retaken. The daily wounded rate for these units during the 32 days of the battle was 17.5 per 1000 strength, and ranged from 1.6 to 45.5. The daily KIA rate was 2.2 and ranged from 0.0 to 9.6. For comparison purposes, casualty rates sus-

tained during the assault on Okinawa are shown in Figure 3. The daily wounded rate during the Okinawa assault was 6.57 per 1000 strength and ranged from 0.0 to 31.8. The daily KIA rate was 1.35 and varied between 0.0 and 6.4. Also shown for comparison purposes are the pulses and pauses in casualty incidence during the four-month period in which Marine involvement peaked in Vietnam (See Figure 4). The daily wounded rate during this period was 2.50, ranging from 0.0 to 13.9 while the KIA rate was 0.31.

Daily casualty incidence among troops involved in fighting in the urban area south of the river are graphed in Figure 5. The daily wounded rate during this first phase of the battle was 37.9 per 1000 strength, while the KIA rate was 3.1. The average strength during this eight-day period was 755 troops. Figure 6 is a display of the casualty rates for the five individual companies fighting south of the river; all but one company had a wounded rate which exceeded 35 per 1000 strength per day.

Figure 7 shows the daily casualty incidence among Marines fighting inside the Citadel during the second phase of the battle for Hue. The wounded rate during this ten-day period was 44.4 per 1000 strength per day, while the KIA rate was 7.8. The average strength during this phase inside the Citadel was 615 troops. Casualty rates among the individual companies fighting in the Citadel are shown in Figure 8; rates of WIA requiring evacuation ranged from 23 to 35 per 1000 strength per day.

Fluctuations in casualty incidence among troops during the mopping up phase of the battle for Hue are shown in Figure 9. The wounded rate during this seven-day period




Figure 2. Rates of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) among infantry battalions participating in the battle for Hue (January 31 - March 2)




Figure 3. Presentation rates of wounded in action (WIA) and killed in action (KIA) among infantry battalions during the Okinawa Operation (1945)




Figure 4. Presentation rates of wounded in action (WIA) and killed in action (KIA) among infantry battalions during a four month period of the Vietnam War (1968)




Figure 5. Rates of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) during the operations across the river at the battle for Hue




Figure 6. Rates of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) among five individual companies involved in Hue operations south of the river




Figure 7. Rates of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) during operations inside the citadel at the battle for Hue




Figure 8. Rates of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) among four individual companies involved in Hue operations inside the citadel




Figure 9. Rates of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) during mopping up operations at the battle for Hue

was 5.8 per 1000 strength per day, while the KIA rate was 0.93. The average strength during this phase was 2315 troops. Casualty rates among individual companies are shown in Table 1.

Figure 10 contrasts casualty rates of rifle/tank companies with Headquarters & Service (H&S) companies across the three phases of the battle; rates for H&S companies are from the mopping up phase exclusively. Daily rates of wounded among rifle companies exceeded 25 per 1000 strength during the three defined periods of the re-taking of Hue.

(H&S) companies across the three phases of the battle; rates for H&S companies are from the mopping up phase exclusively. Daily rates of wounded among rifle companies exceeded 25 per 1000 strength during the three defined periods of the re-taking of Hue.

Table 1. Comparisons of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) among individual companies involved in mopping up operations in the fighting for Hue.

Company	Rate per 1000 per day		
	WIA	WIANE	KIA
1	0.00	0.80	0.00
2	0.42	0.00	0.00
3	5.30	6.36	0.00
4	23.69	11.85	2.96
5	10.69	0.89	4.46
6	1.27	1.69	0.00
7	7.13	4.07	6.11
8	6.19	0.88	0.00
9	0.00	0.00	0.00
10	0.00	0.00	0.00
11	1.91	0.00	0.00
12	1.46	0.49	0.49
Overall	3.95	1.85	0.93




Figure 10. Comparisons of wounded in action (WIA), wounded not evacuated (WIANE), and killed in action (KIA) between Rifle/Tank Companies and Headquarters & Service (H&S) Companies Involved in the fighting for Hue.

DNBI RATES

Illness incidence during the battle for Hue was also examined and contrasted with DNBI incidence during the Okinawa assault and with another period of the Vietnam Conflict. Figure 11 is a display of the daily Disease/Nonbattle Injury (DNBI) rates and casualty (wounded and killed) rates among the battalions before, during, and after the Battle for Hue. While there were great fluctuations in the casualty rate during the battle for Hue, the DNBI rate was relatively stable over the three month period incorporating the Hue fighting. The DNBI rate was 0.92 per 1000 strength per day in the month prior to Hue, 0.98 during the Hue battle, and 0.92 in the month fol-

lowing the Hue battle.

The lack of major DNBI pulses during the re-taking of Hue is in contrast to large variations in DNBI incidence during the heavy intensity of the Okinawa assault. The rises in DNBI incidence paralleling increases in casualty rates are shown in Figure 12. The DNBI rate over the course of the 90 day Okinawa operation was 4.56 per 1000 strength per day. Figure 13 shows the DNBI and casualty incidence among infantry battalions during peak Marine involvement in Vietnam; as with the Hue data, DNBI incidence was relatively stable over the four-month period. The DNBI rate between May and August 1968 was 1.78.




Figure 11. Rates of disease and non-battle injuries (DNBI) and casualties among infantry battalions before, during, and after the battle for Hue.




Figure 12. Rates of disease and non-battle injury (DNBI) and casualties among infantry battalions during the Okinawa Operation (1945)




Figure 13. Rates of disease and non-battle injury (DNBI) and casualties among infantry battalions during a four month period of the Vietnam War (1968)

DISCUSSION

Tensions between nationalistic and/or ethnic factions in the former Republic of Yugoslavia, Cambodia, Central Asia, and the Middle East have increased dramatically with the end of the Cold War. The recent outbreaks of hostilities make it increasingly likely that United Nations members may vote to mount a military operation to make and enforce peace in one or more of these strife-ridden regions. Given the geographical areas where tensions are highest, it is also quite possible that a U.N.-led force may be called upon to drive a recalcitrant faction from an urban setting.

The present investigation examined rates of casualties and illness incidence sustained in the re-taking of the city of Hue during the Vietnam Conflict. Hue was the cultural capital and third largest city in Vietnam. During the Tet offensive of 1968, a division-sized force of the North Vietnamese Army infiltrated and took control of most of Hue. Three U.S. Marine battalions and 13 South Vietnamese battalions were involved in the fighting to regain the city. U.S. forces fought house-to-house in the battle south of the Huong River and would withdraw to the MACV compound at nightfall only to have

snipers come back in and re-claim positions. Fighting was at close quarters and slow, but after eight days the enemy was routed. Shortly thereafter, U.S. Marines moved into the inner city and fought to regain the southeast section. The WIA rates during the fighting south of the river and within the Citadel were, respectively, 37.9 and 44.4 per 1000 strength per day.

While a relationship between battle intensity and DNBI rates has been previously established⁵, there was no increase in DNBI incidence during the Hue offensive. Previous research suggested that the relationship between battle intensity and DNBI rates weakened with the tempo of the conflict. Apparently, increases in battle fatigue cases and illnesses tied to high levels of battlefield stress are more a function of sustained high intensity operation rather than a high intensity battle within a light-to-moderate intensity conflict.

Extensive medical resource planning for projected casualty and illness incidence during urban warfare is essential. While the demands of treating the DNBI cases among a U.N. force may be relatively minimal, the resources required to treat casualties sustained in close-quarter fighting are great.

REFERENCES

1. Pham VS, Le VD, Nguyen NH. The VC general offensive: General uprising Mau Than 1968. Material Printing Center, Vietnam. August 1968.
2. Christmas GR. A company commander reflects on operation Hue City. In: The Marines in Vietnam 1954-1973. U.S. Government Printing Office, Washington D.C. 1985.
3. Simmons EH. Marine Corps Operations in Vietnam, 1968. In: The Marines in Vietnam 1954-1973. U.S. Government Printing Office, Washington D.C. 1985.
4. Karnow S. Vietnam: A History. Viking Press, New York, N.Y. 1983.
5. Blood CG, Gauker, ED. The relationship between battle intensity and disease rates among Marine Corps infantry units. Military Medicine, 158 (5), 340-44, 1993.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0168
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0168), Washington, DC 20503.</p>			
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE August 1993	3. REPORT TYPE AND DATE COVERED Final Jan 93 to Aug 93	
4. TITLE AND SUBTITLE The Battle for Hue: Casualty and Disease Rates During Urban Warfare		5. FUNDING NUMBERS Program Element: 63706N Work Unit Number: M0095.005-6204	
6. AUTHOR(S) Christopher G. Blood, Marlisa E. Anderson		8. PERFORMING ORGANIZATION Report No. 93-16	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center P. O. Box 85122 San Diego, CA 92186-5122		9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Medical Research and Development Command National Naval Medical Center Building 1, Tower 2 Bethesda, MD 20889-5044	
10. SPONSORING/MONITORING AGENCY REPORT NUMBER			
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Daily rates of casualty and illness incidence sustained in the re-taking of the city of Hue during the North Vietnamese Tet Offensive of 1968 were examined. The daily wounded rate for the U.S. Marine battalions involved was 17.5 per 1,000 strength, and ranged from 1.6 to 45.5. The killed-in-action rate per 1,000 strength per day was 2.2, and ranged from 0.0 to 9.6. The wounded rate during the urban warfare of Hue was three times higher than during the high intensity battle for Okinawa and six-fold the wounded rate during normal Marine operations at the peak of the Vietnam Conflict. The disease/nonbattle injury (DNBI) rate remained steady over the course of the Hue operation at approximately 1.0 per 1,000 strength per day.			
14. SUBJECT TERMS Casualty rates, illness incidence, urban warfare, medical planning			15. NUMBER OF PAGES 14
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited