

AD-A272 664 NTATION PAGE

Form Approved
OMB No. 0704-0188

2



ated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, reviewing the collection of information, sending comments regarding this burden estimate or any other aspect of this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Washington, DC 20540, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1 AGENCY USE ONLY (Leave blank)	2 REPORT DATE	3 REPORT TYPE AND DATES COVERED WAR-143-72
4 TITLE AND SUBTITLE Lipid A and liposomes containing lipid A as adjuvants for vaccines - Chapter 18		5 FUNDING NUMBERS
6 AUTHOR(S) Carl R. Alving		
7 PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Division of Biochemistry Walter Reed Army Institute of Research Washington, DC 20307-5100		8. PERFORMING ORGANIZATION REPORT NUMBER
9 SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Medical Research & Development Command Fort Detrick, Frederick, MD 21701 Walter Reed Army Institute of Research Washington, DC 20307-5100		10 SPONSORING/MONITORING AGENCY REPORT NUMBER
11 SUPPLEMENTARY NOTES		
12a DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.		12b DISTRIBUTION CODE

13 ABSTRACT (Required)	14 ABSTRACT FOR
Literature review on lipid A as adjuvant	<input checked="" type="checkbox"/> CONFIDENTIAL <input type="checkbox"/> UNCLASSIFIED <input type="checkbox"/> OTHER
<p>DTIC QUALITY INSPECTED</p> <p>A-120</p>	

15 NUMBER OF PAGES Lipid A, Adjuvants, Vaccines, Liposomes	16 PRICE CODE
17 SECURITY CLASSIFICATION OF THIS PAGE Unclassified	18 SECURITY CLASSIFICATION OF ABSTRACT Unclassified
19 LIMITATION OF ABSTRACT	20 LIMITATION OF ABSTRACT

Bacterial Endotoxic Lipopolysaccharides

Volume II Immunopharmacology and Pathophysiology

Edited by

John L. Ryan, Ph.D., M.D.

Executive Director
Infectious Diseases Department
Merck Research Laboratories
West Point, Pennsylvania

David C. Morrison, Ph.D.

Associate Director, Cancer Center
Professor of Microbiology,
Molecular Genetics, and Immunology
University of Kansas Medical Center
Kansas City, Kansas



93-27236



CRC Press

Boca Raton Ann Arbor London Tokyo

93 11 0 0 97

Library of Congress Cataloging-in-Publication Data

Bacterial endotoxic lipopolysaccharides / editors, David C. Morrison,
John Ryan.

p. cm.

Includes bibliographical references and indexes.

Contents: v. 1. Molecular biochemistry and cellular biology -- v.

2. Immunopharmacology and pathophysiology.

ISBN 0-8493-6787-5 (v. 1). — ISBN 0-8493-6788-3 (v. 2)

1. Endotoxins. I. Morrison, David C., 1941- II. Ryan, John
Louis.

[DNLM: 1. Endotoxins. 2. Lipopolysaccharides—physiology. QW
630 B1285]

QP632.E4B23 1992

615.9 5299—dc20

DNLM/DLC

for Library of Congress

92-7865

CIP

This book represents information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. A wide variety of references are listed. Every reasonable effort has been made to give reliable data and information, but the author and the publisher cannot assume responsibility for the validity of all materials or for the consequences of their use.

Neither this book nor any part may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, microfilming, and recording, or by any information storage and retrieval system, without permission in writing from the publisher.

Direct all inquiries to CRC Press, Inc., 2000 Corporate Blvd., N.W., Boca Raton, Florida, 33431.

© 1992 by CRC Press, Inc.

International Standard Book Number 0-8493-6787-5 (vol. 1)

International Standard Book Number 0-8493-6788-3 (vol. 2)

Library of Congress Card Number 92-7865

Printed in the United States of America 1 2 3 4 5 6 7 8 9 0

Printed on acid-free paper

Chapter 18

**LIPID A AND LIPOSOMES CONTAINING LIPID A
AS ADJUVANTS FOR VACCINES**

Carl R. Alving

TABLE OF CONTENTS

I.	Adjuvant Activities of Lipopolysaccharide and Lipid A	430
II.	Mechanisms of Adjuvant Activity	430
A.	Insights from Synthetic Lipid A Analogs	430
B.	Role of Macrophages	430
C.	Effects of LPS on Antigen Presentation	431
D.	Effects on Suppressor T-Cells	431
III.	Vaccine Strategies	431
A.	Safety	433
1.	Lipid A Analogs	433
2.	Liposomes Containing Lipid A	434
B.	Enhancement of Immunogenicity	436
1.	Use of Liposomes	437
2.	Adjuvant Effects of Liposomes Containing Lipid A	437
3.	Liposomal Vaccine	438
	References	439

I. ADJUVANT ACTIVITIES OF LIPOPOLYSACCHARIDE AND LIPID A

The ability of Gram-negative bacterial lipopolysaccharide (LPS) to serve as an adjuvant to enhance immune responses to antigens has been recognized for more than 30 years.¹⁻³ LPS itself is also highly immunogenic. As evidence of this it was shown that LPS could sensitize mice to a secondary immune response with a few tens of molecules (picogram amounts)⁴ and could induce primary immune responses in rabbits after injection of only a few thousand molecules.⁵ Because of its potent immunogenicity LPS has been referred to as a "super antigen."⁴ Adjuvant activity of LPS was subsequently shown to reside in the lipid A moiety, and it was hypothesized that "LPS is a powerful immunogen because it is an antigen [polysaccharide] that carries its own adjuvant [lipid A]."⁶

The adjuvant activity of lipid A has been confirmed in many laboratories,⁷⁻¹¹ and activity has also been demonstrated with synthetic analogs of lipid A.¹²⁻¹⁴ Experiments with chemically modified LPS have led to the conclusion that only the lipid A portion is required for adjuvant activity, and adequate amounts of esterified and amidated lipid A fatty acids are also required.⁸

II. MECHANISMS OF ADJUVANT ACTIVITY

A. INSIGHTS FROM SYNTHETIC LIPID A ANALOGS

The availability of synthetic lipid A analogs has permitted analysis of minimal structural requirements for adjuvant activity.^{15,16} Synthetic analogs have also provided insights into the relatively small importance of the disaccharide structure compared with monosaccharide structures for adjuvant activity. In contrast, the fatty acid composition, including stereospecificity, is extremely critical for expression of adjuvant activity of lipid A.^{16,17} The number and composition of lipid A fatty acids would be expected to have an important effect on aggregation characteristics of lipid A and on the interactions of lipid A with cells, with liposomes, and with liposomal antigen. Lipid A fatty acids, therefore, might influence the adjuvant effects of liposomal lipid A in vaccine formulations that utilize liposomes as carriers. However, such influences of lipid A fatty acids cannot be easily predicted, and the relative efficacies of lipid A analogs can only be determined by empirical testing in candidate vaccine formulations.

B. ROLE OF MACROPHAGES

Early studies on the fate of radioactively labeled, parenterally injected LPS and lipid A demonstrated that both compounds rapidly left the blood and accumulated in macrophages in the liver and spleen.¹⁸ The concept that the adjuvant effect of LPS was related to its effect on macrophages was at least partly based on experiments involving transfer of murine macrophages

from donor animals to syngeneic recipient animals. Donor macrophages that had been allowed to ingest antigen *in vitro* enhanced the immune response to the antigen in the recipient animals, and enhancement was greatly stimulated by pretreating the macrophages with LPS.^{19, 21}

The adjuvant effects of LPS and lipid A on macrophages have now been well established. The interactions of LPS with macrophages cause myriad macrophage synthesis and secretory activities, and signal transduction mechanisms that are sometimes referred to as macrophage activation.²² In addition to the recruitment and activation of macrophages, it is also recognized that LPS and lipid A stimulate multiple cellular interactions. An important role of interactions of macrophages with T-lymphocytes has been proposed as a major basis for adjuvant effects of LPS.²³ The cellular interactions induced by LPS and lipid A are mediated by numerous secreted cytokines. Some of the complex interactions that occur between mediators and cells as a result of LPS and lipid A stimulation are illustrated in Figure 1.

C. EFFECTS OF LPS ON ANTIGEN PRESENTATION

In recent years the concept of the macrophage as an antigen presenting cell (APC) has emerged.²⁵ A key element in the process of antigen presentation is the participation of major histocompatibility gene complex (MHC) molecules. For induction of an immune response against foreign antigens, class II MHC molecules (also known as Ia molecules) are expressed by the APC, and recognition by helper T-cells of a complex of processed antigen with Ia on the surface of the APC is the initial event leading to an immune response.²⁵ It has now been demonstrated that LPS can stimulate induction of enhanced Ia expression by murine macrophages.²⁶⁻²⁸ Besides recruiting and stimulating macrophages to enhance the immune response against antigens, LPS also elicits macrophages that cause increased antigen catabolism. This latter process can lead to down-regulation of the immune response.²⁹

D. EFFECTS ON SUPPRESSOR T-CELLS

The immune response to pneumococcal polysaccharide is greatly amplified by the use of monophosphoryl lipid A (MPL).³⁰ The enhancing effect of MPL was attributed to the inactivation of suppressor T-cell activity, and this may represent a further adjuvant mechanism for lipid A.

III. VACCINE STRATEGIES

Among the most desirable attributes of a modern adjuvanted vaccine are that the vaccine should be safe and that the adjuvant should increase the specific efficacy of the vaccine to a sufficiently high level that the benefit of the adjuvant outweighs any risk and reactogenicity associated with the adjuvant.³¹ Although many of the most successful vaccines have had annoying but tolerable reactogenic characteristics, the evolution of vaccine technology

Stimulates or suppresses other T cells, B cells and macrophages; kills infected cells

Differentiates and produces antibodies




FIGURE 1. Secretion of mediators by an LPS-stimulated macrophage (center) sets in motion further secretory stimuli and biological effects by numerous cell types. Of particular interest for adjuvant development for vaccines are the effects of tumor necrosis factor (TNF), interleukin (IL) 1, IL 2, IL 4, IL 5, and IL 6, growth factor (TGF β), and interferon γ (IFN γ), on proliferation and differentiation of T- and B-lymphocytes that are crucial elements in the process of antigen presentation. (Redrawn and based on information from Reference 24.)

has now led to proliferation of myriad peptide and recombinant protein antigens that have inherently high levels of safety because of their purity. Unfortunately, simplified soluble protein antigens do not necessarily have high levels of inherent immunogenicity. In many cases this is because the original antigen is membrane-associated, and in its native form it and its associated neighboring environment are handled by the immune system as a particle. Particulate antigens undergo efficient processing and presentation by macrophages. Soluble antigens are much more likely to be processed by other types of APCs such as B-lymphocytes or dendritic cells. Macrophages, because of their complicated biological roles as scavenger and secretory cells, are more likely to induce potent immune responses to complex foreign antigens.

A. SAFETY

As noted earlier, macrophages are widely believed to serve as the target cell for LPS and lipid A, and also for many other types of adjuvants. Unfortunately, the same properties that stimulate secretory events leading to enhancement of the immune response can sometimes also cause undesirable local or systemic reactions that limit the potential usefulness of LPS and lipid A as adjuvants. One of the most common adverse reactions to vaccines is pyrogenicity, and it is perhaps *ironic in the present context* to note that in many instances the efficacy of the vaccine may be dependent on the presence of LPS, either as an unintended contaminant or as an integral part of the antigen, and the LPS may also be responsible for induction of pyrogenicity.

It is well known that different biological effects of LPS and lipid A can be related to different chemical moieties in the structure of LPS and lipid A. It has been argued that certain so-called "beneficial" effects may be identified with chemical structures that are different than the structures associated with "toxic" effects.¹⁷ The concept that nontoxic structures might be produced that retain the potent adjuvant effects of lipid A has generated optimism and interest in lipid A among investigators in the field of vaccinology.

I. Lipid A Analogs

Ribi and colleagues have demonstrated that an MPL formulation exhibits reduced lethality and pyrogenicity, but still retains antitumor and adjuvant activities.^{18,19,20} In the same context, it has also been a major goal in the lipid A biosynthesis field to produce lipid A analogs that would retain beneficial biological activities, including adjuvanticity, but lack endotoxic activity.^{17,21,22,23} Both of these approaches, *viz.*, chemically altered native lipid A and synthetic low toxicity analogs of lipid A, show considerable promise as adjuvants for vaccines. One possible limiting factor to these approaches would be if significant residual reactogenicity of the lipid A formulation were still observed at doses of lipid A that would be required for optimal immune responses. Intravenously administered MPL has been tested in a Phase I trial




FIGURE 2. Effect of infusion of lipid A ($10 \mu\text{g/kg}$) on leukocyte counts (mean \pm SEM) in rabbits. Lipid A that was in 0.5% triethylamine (TEA), or in liposomes, was diluted with 0.9% NaCl and infused during 1 h into four rabbits. Six controls received 0.5% TEA in 0.9% NaCl. (From Reference 34.)

in humans.³⁵ The maximal safe intravenous dose in humans was estimated to be $100 \mu\text{g/m}^2$, but even at $25 \mu\text{g/m}^2$ minor episodes of fever, gastrointestinal symptoms, and chills were often observed.³⁵ Most vaccine formulations are given intramuscularly or subcutaneously, and it is possible that these latter routes of administration might result in less reactogenicity than the intravenous route.

2. Liposomes Containing Lipid A

Another approach that has been employed in an effort to widen the gap between reactogenicity and adjuvanticity has been to incorporate lipid A into liposomes. Among the activities of lipid A that are reduced by incorporation into liposomes are included: Limulus lysate coagulation *in vitro*,^{36, 37} neutropenia in rabbits,³⁶ stimulation of both interleukin 1^{β} and tumor necrosis factor³⁸ by a macrophage cell line, pyrogenicity in rabbits,³⁹ and lethal toxicity to mice.⁴¹ Intravenously infused lipid A characteristically induces a profound neutropenia in rabbits, with a nadir reached several hours after initiation of infusion. As shown in Figure 2, neutropenia was not induced by liposomal lipid A.

One of the most common complaints associated with many vaccines is the occurrence of fever. In many cases pyrogenic reactions are due to the

TABLE I
Effect on Pyrogenicity of Incorporating Native Lipid A and Monophosphoryl Lipid A in Liposomes*

Lipid A type	Maximum nonpyrogenic dose ^b		Decrease in pyrogenicity of liposomal compared with free lipid A
	Free lipid A	Liposomal lipid A	
S. minnesota R595	0.008	1.7	214-fold
MP lipid A (MPL)	0.32	8.1	25-fold

* Data obtained from Reference 40.

^b Maximum nonpyrogenic dose was the maximum lipid A dose ($\mu\text{g}/\text{kg}$) that did not cause an increase in temperature of 0.6°C or greater in any of three rabbits over a 3-h period.

MP lipid A is a monophosphoryl lipid A fraction derived from *Staphylococcus aureus* R595 lipid A.

presence of LPS or lipid A, either as a contaminant or as an antigen in the vaccine. As mentioned earlier, MPL is less pyrogenic than native lipid A when tested in a rabbit pyrogenicity model. In experiments designed to explore the roles of physical environment and chemical structure on pyrogenicity, MPL was approximately 40-fold less pyrogenic than the native lipid A from which the MPL was derived (maximum nonpyrogenic dose of 0.32 compared with 0.008 $\mu\text{g}/\text{kg}$) (Table I). Liposomes further reduced the pyrogenicity by at least 25-fold (Table I), thus causing liposomal MPL to be 1000-fold less pyrogenic than free native lipid A. Vaccine grade liposomal MPL, prepared for human use according to Good Manufacturing Practices (GMP) as promulgated by the U.S. Food and Drug Administration, was 8-fold less pyrogenic yet (maximum nonpyrogenic dose of 66 μg of liposomal MPL per kilogram).¹¹ Inclusion of aluminum hydroxide (alum) as a nonliposomal adjuvant reduced the pyrogenicity of GMP liposomal MPL still further (approximately tenfold more).¹² It is, therefore, estimated that GMP liposomal MPL adsorbed to alum may be approximately 80,000-fold less pyrogenic than native lipid A alone.

Despite the virtual elimination of pyrogenicity, liposomal MPL still serves as an extremely potent adjuvant for inducing humoral immunity against liposomal protein antigens both in animals¹³ (Figure 3) and humans.¹⁴ The previously expressed concern that lipid A in liposomes would serve as an adjuvant only at pyrogenic concentrations, and that lipid A would, thus, be unacceptable for use in humans,¹⁵ has, therefore, been proved to be unfounded. The strong adjuvant effects of nonpyrogenic formulations of liposomal lipid A are also consistent with the observations mentioned earlier that adjuvant effects of certain synthetic lipid A analogs can be dissociated from pyrogenic effects.




FIGURE 3. Immune response of individual monkeys to liposomes containing a recombinant antigen containing epitopes from the circumsporozoite protein of *Plasmodium falciparum*. The liposomes contained either native lipid A (0.5 or 44 $\mu\text{g}/\text{kg}$, as indicated) or monophosphoryl lipid A (4 $\mu\text{g}/\text{kg}$) as an adjuvant. The liposomes were adsorbed with alum where indicated. The FSV-1 formulation (Falciparum Sporozoite Vaccine-1) consisted of the antigen alone adsorbed to alum, and this was the same vaccine that was previously tested in a human clinical trial. (From Reference 40.)

B. ENHANCEMENT OF IMMUNOGENICITY

In the context of vaccine usage an adjuvant is defined as a substance that enhances the specific immune response against an antigen. Research on the effector mechanisms of various adjuvants has revealed that different adjuvants may work at different stages or locations in the immune response. A rational strategy in formulating a vaccine would, therefore, be to enlist combinations of substances that could exert adjuvant effects at more than one point. This would allow additive effects to occur, and it might permit the usage of smaller amounts of each adjuvant component, thus promoting increased safety.

Because of the central role played by macrophages in adjuvant mechanisms of LPS and lipid A, a logical approach would be to promote increased delivery of LPS or lipid A to macrophages. One of the most effective methods currently being employed to deliver therapeutic substances to macrophages is to use liposomes as carriers.⁴⁴⁻⁴⁶

1. Use of Liposomes

Liposomes have been highly successful as carriers of antigens and adjuvants.⁴⁷ The theoretical basis for interactions of liposomes with the immune system and for enhancement of immune responses against liposomal protein antigens has been extensively reviewed elsewhere.⁴⁷ Liposomes provide a means to reconstitute hydrophobic antigens, or to encapsulate soluble antigens or peptides. They also provide a solid surface for the chemical attachment of antigens.

Intravenously injected liposomes are rapidly removed from the blood and are avidly ingested by macrophages in the liver and spleen where they undergo gradual degradation. The fate of intramuscularly or subcutaneously injected liposomes is dependent to a great extent on the scavenging characteristics of local tissue macrophages or macrophages recruited to the site in response to an inflammatory or adjuvant stimulus. For example, liposomal antigen (adenovirus hexon) that was injected intramuscularly was slowly released from the injection site with a half-life for clearance of 30 to 44 h.⁴⁸ It is, therefore, evident that a transient sustained release or depot effect contributes to the mechanism by which intramuscularly injected liposomal antigen interacts with the immune system. The depot effect of liposomes probably does not substantially enhance the adjuvant effect of lipid A compared with that observed with lipid A alone since it has been reported that subcutaneously injected lipid A alone has a half-life at the injection site of more than 15 d.⁴⁹

As noted earlier, liposomes also attenuate the toxic effects of lipid A, but the adjuvant effect of lipid A is retained even in the absence of pyrogenicity or other types of reactogenicity. It is, therefore, possible that the blocking of the toxicity of lipid A by liposomes is partly due to retardation of release of lipid A into the environment and also due to degradation of liposomal lipid A along with the liposomes in macrophages.

2. Adjuvant Effects of Liposomes Containing Lipid A

Early studies demonstrated that lipid A enhanced the murine immune responses against liposomal haptens consisting of lipid derivatives of either 2,4-dinitrophenyl or fluorescein groups.^{7,50} It was soon suggested that liposomes containing lipid A (or liposomes containing LPS) could stimulate immune responses against liposomal proteins,^{9,51} or even against lipid A as an antigen.⁹ Lipid A proved to be such a potent adjuvant that antibodies were even induced against both the liposomes themselves and liposomal phospholipid constituents.^{52,53}

Liposomes containing lipid A have been used as adjuvants for enhancing immune responses to cholera toxin;^{9,54,56} herpes simplex antigens, including glycoprotein-enriched antigens⁵⁷ and a peptide-palmitic acid conjugate;⁵⁷ Epstein-Barr virus membrane antigens;⁵⁸ *Plasmodium falciparum* sporozoite antigens, including a peptide-protein conjugate,⁵⁵ and a recombinant protein;^{40,59} an unconjugated 25-amino acid peptide from the active site of acetylcholinesterase;⁶⁰ and tumor-associated glycosphingolipids.⁶¹

One of the difficulties that is often encountered in adjuvant research lies in evaluating the relative efficacy of an adjuvant in comparison to other adjuvants. There are no standardized techniques available for this purpose because of the existence of multiple adjuvant mechanisms and immunological techniques. Different chemical and physical structures of adjuvants often preclude standardization of all the different possible variables. Comparative evaluation of adjuvants such as lipid A, analogs of lipid A, LPS, and muramyl dipeptide can be a heroic undertaking.⁶² However, some of these difficulties can be overcome with liposomes. Many variables can be simultaneously controlled by utilizing liposomes containing both the antigen and the adjuvant. In one study, direct comparison of the adjuvant effects of liposomal lipid A with liposomal lipophilic MDP derivatives demonstrated that, under the conditions employed, lipid A had stronger adjuvant effects than lipophilic MDP (Figure 4).⁵⁴

3. Liposomal Vaccine

The first injectable liposomal vaccine was administered to human volunteers in a Phase I trial in October 1989.⁴² The liposomes contained a recombinant protein antigen having repeat sequence epitopes from the circumsporozoite protein of *P. falciparum*. MPL was included in the liposomes as an adjuvant, and the liposomes containing antigen and MPL were also adsorbed to aluminum hydroxide to prolong the depot effect and provide further adjuvant activity. At the highest dose tested, 2.2 mg of liposomal MPL was administered with each intramuscular injection of vaccine (at 0, 12, and 20 weeks). Despite the extremely high doses of liposomal MPL (approximately 12-fold greater than the maximum safe dose of intravenous free MPL in humans),³⁵ the liposomal MPL exhibited virtually no acute toxic effects. The results from this phase I human trial have demonstrated that extremely high levels of specific IgG antibodies against the appropriate antigenic epitopes have been achieved.

The success of the human liposomal vaccine in providing exceptionally high levels of serum antibodies validates the concept that lipid A can be combined with other types of carriers and adjuvants to provide additive adjuvant activities. The high doses of MPL employed also demonstrate that potent adjuvant effects can be achieved in the absence of toxicity. Because of the success of this initial vaccine for including enhanced immunity, ad-




FIGURE 4. Immune response against cholera toxin (CT) alone, CT associated with complete Freund's adjuvant (CFA), liposomes, or liposomes containing lipid adjuvant. Each line is the mean of two rabbits immunized either with 5 μ g CT alone (i.v. injection), or with 5 μ g CT either emulsified with CFA (s.c. injection), attached to the surface of liposomes with no adjuvant, or attached to the surface of liposomes containing either lipid A or one of two forms of lipophilic MDP, $\text{CH}_3(\text{CH}_2)_{10}\text{CO-MDP}$ (stearoyl-MDP, or L18-MDP) or $[\text{CH}_3(\text{CH}_2)_{13}\text{CHCO-MDP}$ (B30-MDP) (i.v. injection). The maximum measurable level of radioimmunoassay (RIA) units in the assay was 10^7 . Therefore, for those data points for CFA and lipid A in which this latter value is illustrated, the actual values achieved were probably higher. (From Reference 55.).

ditional candidate human liposomal vaccines employing other antigens and MPL have been proposed, and several are now in advanced stages of development for initial clinical trials.

REFERENCES

1. Johnson, A. G., Gaines, S., and Landy, M., Studies on the O antigen of *Salmonella typhosa*. V. Enhancement of antibody response to protein antigens by the purified lipopolysaccharide, *J. Exp. Med.*, 103, 225, 1956.
2. Franzl, R. E. and McMaster, P. D., The primary immune response in mice. I. The enhancement and suppression of hemolysin production by a bacterial endotoxin. *J. Exp. Med.* 127, 1087, 1968.
3. Ada, G. L., Lang, P. G., and Plymin, G., Antigen in tissues. V. Effect of endotoxin on the fate of, and on the immune response to, serum albumin and to the albumin-antibody complexes, *Immunology*, 14, 825, 1968.

- 4 **Rudbach, J. A.**, Molecular immunogenicity of bacterial lipopolysaccharide antigens, establishing a quantitative system. *J. Immunol.*, 106, 993, 1971
- 5 **Landy, M. and Baker, P. J.**, Cytodynamics of the distinctive immune response produced in regional lymph nodes by Salmonella somatic polysaccharide. *J. Immunol.*, 97, 670, 1966
- 6 **Chiller, J. M., Skidmore, B. J., Morrison, D. C., and Weigle, W. O.**, Relationship of the structure of bacterial lipopolysaccharides to its function in mitogenesis and adjuvancy. *Proc. Natl. Acad. Sci. U.S.A.*, 70, 2129, 1973
- 7 **Dancey, G. E., Yasuda, T., and Kinsky, S. C.**, Enhancement of liposomal model membrane immunogenicity by incorporation of lipid A. *J. Immunol.*, 119, 1868, 1977
- 8 **Cho, Y., Tanamoto, K., Oh, Y., and Homma, J. Y.**, Differences of chemical structures of *Pseudomonas aeruginosa* lipopolysaccharide essential for adjuvanticity and antitumor and interferon inducing activities. *FEBS Lett.*, 105, 120, 1979
- 9 **Alving, C. R., Banerji, B., Clements, J. D., and Richards, R. L.**, Adjuvanticity of lipid A and lipid A fractions in liposomes, in *Liposomes and Immunobiology*, Tom. B. H. and Six. H. R., Elsevier North-Holland, Amsterdam, 1980, 67
- 10 **Johnson, A. G., Tomai, M., Solem, L., Beck, L., and Ribi, E.**, Characterization of a nontoxic monophosphoryl lipid A. *Rev. Infect. Dis.*, 9 (Suppl. 5), S512, 1987
- 11 **Baker, P. J., Hiernaux, J. R., Fauntleroy, M. B., Stashak, P. W., Prescott, B., Cantrell, J. L., and Rudbach, J. A.**, Ability of monophosphoryl lipid A to augment the antibody response of young mice. *Infect. Immun.*, 56, 3064, 1988
- 12 **Kumazawa, Y., Matsuura, M., Homma, J. Y., Nakatsuru, Y., Kiso, M., and Hasegawa, A.**, B cell activation and adjuvant activities of chemically synthesized analogues of the nonreducing sugar moiety of lipid A. *Eur. J. Immunol.*, 15, 199, 1985
- 13 **Kotani, S., Takada, H., Takahashi, I., Ogawa, T., Tsujimoto, M., Shimauchi, H., Ikeda, I., Okamura, H., Tamura, T., Harada, K., Tanaka, S., Shiba, T., Kusumoto, S., and Shimamoto, T.**, Immunobiological activities of synthetic lipid A analogs with low endotoxicity. *Infect. Immun.*, 54, 673, 1986
- 14 **Takada, H. and Kotani, S.**, Structural requirements of lipid A for endotoxicity and other biological activities. *CRC Crit. Rev. Microbiol.*, 16, 477, 1989
- 15 **Yasuda, T., Kanegasaki, S., Tsumita, T., Tadakuma, T., Homma, J. Y., Inage, M., Kusumoto, S., and Shiba, T.**, Biological activity of chemically synthesized analogues of lipid A. Demonstration of adjuvant effect in hapten sensitized liposomal system. *Eur. J. Biochem.*, 124, 405, 1982
- 16 **Takahashi, I., Kotani, S., Takada, H., Tsujimoto, M., Ogawa, T., Shiba, T., Kusumoto, S., Yamamoto, M., Hasegawa, A., Kiso, M., Nishijima, M., Amano, F., Akamatsu, Y., Harada, K., Tanaka, S., Okamura, H., and Tamura, T.**, Requirement of a properly acylated β -1,5-D-glucosamine disaccharide bisphosphate structure for efficient manifestation of full endotoxic and associated bioactivities of lipid A. *Infect. Immun.*, 65, 57, 1987
- 17 **Kumazawa, Y., Nakatsuka, M., Takimoto, H., Furuya, T., Nagumo, T., Yamamoto, A., Homma, J. Y., Inada, K., Yoshida, M., Kiso, M., and Hasegawa, A.**, Importance of fatty acid substituents of chemically synthesized lipid A subunit analogs in the expression of immunopharmacological activity. *Infect. Immun.*, 56, 149, 1988
- 18 **Howard, J. G., Rowley, D., and Wardlaw, A. C.**, Investigations on the mechanism of stimulation of non-specific immunity by bacterial lipopolysaccharides. *Immunology*, 1, 181, 1958
- 19 **Unanue, E. R., Askonas, B. A., and Allison, A. C.**, A role of macrophages in the stimulation of immune responses by adjuvants. *J. Immunol.*, 103, 71, 1969
- 20 **Spitznagel, J. K. and Allison, A. C.**, Mode of action of adjuvants: effects on antibody responses to macrophage-associated bovine serum albumin. *J. Immunol.*, 104, 128, 1970
- 21 **Allison, A. C.**, Mode of action of immunological adjuvants. *J. Reticuloendothelial Soc.*, 26, 619, 1979

22. **Karnovsky, M. L. and Lazdins, J. K.**, Biochemical criteria for activated macrophages. *J. Immunol.* 121, 809, 1978.
23. **McGhee, J. R., Farrar, J. J., Michalek, S. M., Mergenhagen, S. E., and Rosenstreich, D. L.**, Cellular requirements for lipopolysaccharide adjuvanticity. A role for both T lymphocytes and macrophage for *in vitro* responses to particulate antigens. *J. Exp. Med.* 149, 793, 1979.
24. **Old, L. J.**, Tumor necrosis factor. *Sci. Am.*, 258(5), 59, 1988.
25. **Unanue, E. R. and Allen, P. M.**, The basis for the immunoregulatory role of macrophages and other accessory cells. *Science*, 236, 551, 1987.
26. **Wentworth, P. A. and Ziegler, H. K.**, Induction of macrophage Ia expression by lipopolysaccharide and *Listeria monocytogenes* in congenitally athymic nude mice. *J. Immunol.* 138, 3167, 1987.
27. **Marshall, N. E. and Ziegler, H. K.**, Role of lipopolysaccharide in induction of Ia expression during infection with gram-negative bacteria. *Infect. Immun.* 57, 1556, 1989.
28. **Wentworth, P. A. and Ziegler, H. K.**, Modulation of macrophage Ia expression by lipopolysaccharide: stem cell requirements, accessory lymphocyte involvement, and IA-inducing factor production. *Infect. Immun.* 57, 2028, 1989.
29. **Cluff, C. W. and Ziegler, H. K.**, An early response to lipopolysaccharide is the elicitation of macrophages specialized for antigen degradation with negative regulatory effects on the induction of specific immune responses. *Infect. Immun.* 55, 1346, 1987.
30. **Baker, P. J., Hiernaux, J. R., Fauntleroy, M. B., Prescott, B., Cantrell, J. L., and Rudbach, J. A.**, Inactivation of suppressor T-cell activity by nontoxic monophosphoryl lipid A. *Infect. Immun.* 56, 1076, 1988.
31. **Edelman, R.**, Vaccine adjuvants. *Rev. Infect. Dis.* 2, 370, 1980.
32. **Nowotny, A.**, Review of the molecular requirements of endotoxic actions. *Rev. Infect. Dis.* 9 (Suppl. 5), S503, 1987.
33. **Takayama, K., Ribi, E., and Cantrell, J. L.**, Isolation of a nontoxic lipid A fraction containing tumor regression activity. *Cancer Res.* 41, 2654, 1981.
34. **Amano, K., Ribi, E., and Cantrell, J. L.**, Structural requirements of endotoxic glycolipid for antitumor and toxic activity. *J. Biochem.* 93, 1391, 1983.
35. **Vosika, G. J., Barr, C., and Gilbertson, D.**, Phase-I study of intravenous modified lipid A. *Cancer Immunol. Immunother.* 18, 107, 1984.
36. **Ramsey, R. B., Hamner, M. B., Alving, B. M., Finlayson, J. S., Alving, C. R., and Evatt, B. L.**, Effects of lipid A and liposomes containing lipid A on platelet and fibrinogen production in rabbits. *Blood*, 56, 307, 1980.
37. **Richardson, E. C., Banerji, B., Seid, R. C., Jr., Levin, J., and Alving, C. R.**, Interactions of lipid A and liposome-associated lipid A with *Limulus polyphemus* amoebocytes. *Infect. Immun.* 39, 1385, 1983.
38. **Dijkstra, J., Mellors, J. W., Ryan, J. L., and Szoka, F. C.**, Modulation of the biological activity of bacterial endotoxin by incorporation into liposomes. *J. Immunol.* 138, 2663, 1987.
39. **Dijkstra, J., Larrick, J. W., Ryan, J. L., and Szoka, F. C.**, Incorporation of LPS in liposomes diminishes its ability to induce tumoricidal activity and tumor necrosis factor secretion in murine macrophages. *J. Leukocyte Biol.* 43, 436, 1988.
40. **Richards, R. L., Swartz, G. M., Jr., Schultz, C., Hayre, M. D., Ward, G. S., Ballou, W. R., Chulay, J. D., Hockmeyer, W. T., Berman, S. L., and Alving, C. R.**, Immunogenicity of liposomal malaria sporozoite antigen in monkeys: adjuvant effects of aluminum hydroxide and non-pyrogenic liposomal lipid A. *Vaccine* 7, 506, 1989.
41. **Dijkstra, J., Mellors, J. W., and Ryan, J. L.**, Altered *in vivo* activity of liposome-incorporated lipopolysaccharide and lipid A. *Infect. Immun.* 57, 3357, 1989.
42. **Fries, L. F., Gordon, D. M., Richards, R. L., Egan, J. E., Hollingdale, M. R., Gross, M., Silverman, C., and Alving, C. R.**, Liposomal malaria vaccine in humans: a safe and potent adjuvant strategy. *Proc. Natl. Acad. Sci. U.S.A.* 89, 358, 1992.

43. Allison, A. C. and Byars, N. E., An adjuvant formulation that selectively elicits the formation of antibodies of protective isotypes and of cell-mediated immunity. *J. Immunol Methods*, 95, 157, 1986.
44. Alving, C. R., Delivery of liposome-encapsulated drugs to macrophages. *Pharmacol Ther.*, 22, 407, 1983.
45. Fidler, I. J., Macrophages and metastasis—a biological approach to cancer therapy: presidential address. *Cancer Res.*, 45, 4714, 1985.
46. Alving, C. R., Macrophages as targets for delivery of liposome-encapsulated antimicrobial agents. *Adv. Drug Deliv. Rev.*, 2, 107, 1988.
47. Alving, C. R., Liposomes as carriers of antigens and adjuvants. *J. Immunol Methods*, 140, 1, 1991.
48. Kramp, W. J., Six, H. R., and Kasel, J. A., Postimmunization clearance of liposome entrapped adenovirus type 5 hexon. *Proc. Soc. Exp. Biol. Med.*, 169, 135, 1982.
49. Yokochi, T., Inoue, Y., Yokoo, J., Kimura, Y., and Kato, N., Retention of bacterial lipopolysaccharide at the site of subcutaneous injection. *Infect. Immun.*, 57, 1786, 1989.
50. Tamauchi, H., Tadakuma, T., Yasuda, T., Tsumita, T., and Saito, K., Enhancement of immunogenicity by incorporation of lipid A into liposomal model membranes and its application to membrane-associated antigens. *Immunology*, 50, 605, 1983.
51. van Rooijen, N. and van Nieuwmege, R., Endotoxin enhanced adjuvant effect of liposomes, particularly when antigen and endotoxin are incorporated within the same liposome. *Immunol. Commun.*, 9, 747, 1980.
52. Schuster, B. G., Neidig, M., Alving, B. M., and Alving, C. R., Production of antibodies against phosphocholine, phosphatidylcholine, sphingomyelin, and lipid A by injection of liposomes containing lipid A. *J. Immunol.*, 122, 900, 1979.
53. Alving, C. R., Antibodies to liposomes, phospholipids and phosphate esters. *Chem. Phys. Lipids*, 40, 303, 1986.
54. Alving, C. R., Richards, R. L., Moss, J., Alving, L. I., Clements, J. D., Shiba, T., Kotani, S., Wirtz, R. A., and Hockmeyer, W. T., Effectiveness of liposomes as potential carriers of vaccines: applications to cholera toxin and human malaria sporozoite antigen. *Vaccine*, 4, 166, 1986.
55. Pierce, N. F., Sacci, J. B., Jr., Alving, C. R., and Richardson, E. C., Enhancement by lipid A of mucosal immunogenicity of liposome-associated cholera toxin. *Rev. Infect. Dis.*, 6, 563, 1984.
56. Naylor, P. T., Larsen, H. S., Huang, L., and Rouse, B. T., In vivo induction of anti-herpes simplex virus immune response by type I antigens and lipid A incorporated into liposomes. *Infect. Immun.*, 36, 1209, 1982.
57. Brynestad, K., Babbitt, B., Huang, L., and Rouse, B. T., Influence of peptide acylation, liposome incorporation, and synthetic immunomodulators on the immunogenicity of a 1-23 peptide of glycoprotein D of herpes simplex virus: implications for subunit vaccines. *J. Virol.*, 64, 680, 1990.
58. Morgan, A. J., Epstein, M. A., and North, J. R., Comparative immunogenicity studies on Epstein-Barr virus membrane antigen (MA) gp340 with novel adjuvants in mice, rabbits, and cotton-top tamarins. *J. Med. Virol.*, 13, 281, 1984.
59. Richards, R. L., Hayre, M. D., Hockmeyer, W. T., and Alving, C. R., Liposomes, lipid A, and aluminum hydroxide enhance the immune response to a synthetic malaria sporozoite antigen. *Infect. Immun.*, 56, 682, 1988.
60. Ogert, R. A., Gentry, M. K., Richardson, E. C., Deal, C. D., Abramson, S. N., Alving, C. R., Taylor, P., and Doctor, B. P., Studies on the topography of the catalytic site of acetylcholinesterase using polyclonal and monoclonal antibodies. *J. Neurochem.*, 55, 756, 1990.
61. Brodin, T., Thurin, J., Stromberg, N., Karlsson, K., and Sjogren, H. O., Production of oligosaccharide-binding monoclonal antibodies of diverse specificities by immunization with purified tumor-associated glycolipids inserted into liposomes with lipid A. *Eur. J. Immunol.*, 16, 951, 1986.

- 62 **Kotani, S., Takada, H., Tsujimoto, M., Ogawa, T., Mori, Y., Sakuta, M., Kawasaki, A., Inage, M., Kusumoto, S., Shiba, T., and Kasai, N.** Immunobiological activities of synthetic lipid A analogs and related compounds as compared with those of bacterial lipopolysaccharide. Re: glycolipid, lipid A, and muramyl dipeptide. *Int. J. Immun.* 42: 258, 1983.
- 63 **Alving, C. R. et al.**, unpublished data.