NPS-OR-93-015

# NAVAL POSTGRADUATE SCHOOL

AD-A273 215

# Monterey, California



## TBMs AND THE FLAMING DATUM PROBLEM

Alan Washburn

September 1993

Approved for public release; distribution is unlimited.

Prepared for: Ballistic Missile Defense Office Washington, D.C. 20301-7100







#### NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943-5000

Rear Admiral T. A. Mercer Superintendent Harrison Shull Provost

This report was prepared for and funded by Ballistic Missile Defense Office, Washington, D.C.

Reproduction of all or part of this report is authorized.

This report was prepared by:

ALAN WASHBURN Professor of Operations Research

Reviewed by:

Released by:

PETER PURDUE Professor and Chairman Department of Operations Research

rarto

PAUL J. MARTO Dean of Research

REPORT D	Form Approved OMB No. 0704-0188				
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of I collection of information, including suggestions for reducing this burden, to Washington Headquariers Services, Directionate for Information Operations and Reports, 1215 Jeffers Davis Highway, Suite 1204, Artington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188). Washington, DC 20503.					
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE September 1993	3. REPORT TYPE AND DA Technical	ATES COVERED		
4. TITLE AND SUBTITLE TBMs and the Flaming	Datum Problem	<b></b>	5. FUNDING NUMBERS		
6. AUTHOR(S)			PE1731804.1180		
Alan Washburn					
7. PERFORMING ORGANIZATION NAM	8. PERFORMING ORGANIZATION REPORT NUMBER				
Naval Postgraduate Sc Monterey, CA 93943	NPS-OR-93-015				
9. SPONSORING / MONITORING AGEN	10. SPONSORING / MONITORING AGENCY REPORT NUMBER				
Ballistic Missile Defens Washington, D.C. 203					
11. SUPPLEMENTARY NOTES	<u> </u>		L		
Approved for public re					
13. ABSTRACT (Maximum 200 word	(s)	<u></u>	L		
Theater Ballistic Miss because the missile's The situation is simila datum" near which A describes how some s of locating the TBM la	ile launching systems a track can be extrapolate ar to one where a subma SW forces may concent simple analytic methods auncher.	re vulnerable just a ed backwards to the arine torpedoes a s rate a search for th s adapted from AS	after a missile is launched e location of the launcher. hip, thus creating a "flaming e submarine. This report W can be applied to the task		
14. SUBJECT TERMS TBM, SCUD, SEARCH			15. NUMBER OF PAGES 13 16. PRICE CODE		
17. SECURITY CLASSIFICATION	18. SECURITY CLASSIFICATION	19. SECURITY CLASSIFICA	TION 20. LIMITATION OF ABSTRACT		
OF REPORT	OF THIS PAGE I Inclassified	OF ABSTRACT	тп		
NSN 7540-01-280-5500		Unicid Sined	Stendard Form 298 (Rev. 2-89)		

### TBMs AND THE FLAMING DATUM PROBLEM

Alan Washburn Naval Postgraduate School Monterey, CA 93943-5000

#### ABSTRACT

Theater Ballistic Missile launching systems are vulnerable just after a missile is launched because the missile's track can be extrapolated backwards to the location of the launcher. The situation is similar to one where a submarine torpedoes a ship, thus creating a "flaming datum" near which ASW forces may concentrate a search for the submarine. This report describes how some simple analytic methods adapted from ASW can be applied to the task of locating the TBM launcher.

· · ·	 •	

Accesion For			
NTIS	CRA&I	M	
DTIC	TAB		- 1
Unannourised			
Justification			
By Distribution (			
Availability Codes			
Dist	Avail and/or Special		
<b>A</b> 1			
A-			

#### **1. THE TBM PROBLEM**

SCUD launches during Desert Storm were a significant nuisance to the Allies, and could have been worse had the missiles been armed with other than conventional warheads. SCUD missiles are launched from Transporter Erector Launcher vehicles (TELs), the kind of system that can be expected with any Theater Ballistic Missile (TBM). TELs normally attempt to avoid detection, since they are subject to air attack if caught in the open. Launching the missile requires the TEL to be in the open, so the Allies spent considerable time and effort in "SCUD hunting" – attempting to find and destroy the TELs. This effort was not notably successful, (Cohen, 1993) so in the aftermath there has been some effort to explain why and perhaps fix whatever the problem was, since similar situations involving TBMs could occur in the future. This report is devoted to a simple model of one method of dealing with TBMs, the "flaming datum" attack that may succeed in destroying the TEL right after the launch of a missile.

One would prefer, of course, to find the TEL *before* the missile is launched, rather than after. If the TELs have sufficient area to roam in, however, or if they are difficult to distinguish from innocent traffic, it may not be possible to search the available area rapidly enough to have an appreciable effect. There is an AntiSubmarineWarfare (ASW) precedent for this situation. Submarines are difficult to detect without some kind of initial cue as to location, so one strategy for protecting ships from torpedo attacks is to wait until an attack occurs, which establishes a "flaming datum" near which there must be a submarine, and then react quickly to detect and attack the submarine. In doing so one is protecting all the ships that the submarine would have attacked in the future, if not the current victim. Such tactics are effective against submarines because each submarine would attack many ships if left alone; sinking a submarine right after a torpedo attack is

almost as good as sinking it beforehand, statistically speaking. These tactics are also potentially effective against TELs, since a TBM launch is easily detected and extrapolated back to where the TEL must have been when it occurred. The only problem is to get to the launch site quickly and effectively. The chances of doing this are the subject of sections 2 and 3 of this report.

Nothing has been said above about the possibility that TELs may have shelters of some kind that protect them either from observation or attack. Such shelters could do much to tilt the game in favor of the TEL force. In the worst case the shelters would protect the TELs from air attack, and TBMs could be launched from the shelter or so near it as to require very little time in the open. In that case very little can be done from the air. In intermediate cases shelters might themselves be subject to air attack, or might merely protect TELs from observation, or significant travel away from the shelter might be required to launch a TBM. Analysis is possible in these intermediate cases, but it should be emphasized that there are no shelters or protective emplacements of any kind in the model described below, where the TEL's only hope after launch of a TBM is to become lost in an area so large that further search for it is pointless. If the environment is such that a TEL, after launching a missile, says to himself, "Now I have to get away from here in a hurry before searchers arrive", then the methods described below are applicable. If the TEL instead says, "Now I have to hurry back to the shelter", then they are not applicable.

#### 2. FLAMING DATUM ATTACKS

Assume that a launch takes place at time 0, and that the launching vehicle (hereafter the "target") immediately proceeds away from the launch site with speed U. The target necessarily remains inside a circle with radius Ut at time t after the launch, the Farthest-On-Circle or FOC. At time  $t_1$ , a searcher arrives and begins searching in the FOC at speed  $V_S$  and with sweep width W. The sweep width W is

assumed to hold everywhere in the region, so there is no specific place for the target to hide. The target may nonetheless escape detection because the area of the FOC expands quadratically with time whereas the area covered by the searcher expands only linearly. A probabilistic model of detection can be based on this observation. In this model detections are assumed to happen in a nonhomogeneous Poisson Process where the rate of detection  $\lambda(t)$  is the ratio of the rate of covering area ( $V_SW$ ) to the area of the FOC at time t:

$$\lambda(t) = V_S W / (\pi U^2 t^2). \tag{1}$$

If the searcher searches between  $t_1$  and  $t_2$ , the average number of detections  $n(t_1, t_2)$  is the integral of  $\lambda(t)$  between those limits:

$$n(t_1, t_2) = V_S W / (\pi U^2) (1/t_1 - 1/t_2).$$
<sup>(2)</sup>

Since the actual number of detections is a Poisson random variable, the probability of at least one detection is  $1-\exp(-n(t_1, t_2))$ , the desired detection probability. Limited testing of this model in an abstract situation where military officers played the roles of target and searcher in 295 replications is in agreement with it. Figure 1 is taken from Washburn (1989).

Formula (2) is much more sensitive to  $t_1$  than to  $t_2$ . Assume for simplicity that  $t_2$  is so large compared to  $t_1$  that it can safely be assumed to be infinite (note that  $n(t_1, t_2)$  does not approach infinity with  $t_2$  – even searching "forever" will not necessarily detect the target). In that case the detection probability is

$$PD(t_1) = 1 - \exp(-V_S W / (\pi U^2 t_1)).$$
(3)

Formula (3) will be assumed to govern detections in the next section.

#### 3. AVERAGING THE DETECTION PROBABILITY

Formula (3) applies only when the time late  $t_1$  is given. In practice the time late will vary depending on how near the nearest searcher is to the launch (only the nearest searcher is assumed to respond even if several happen to detect or are

informed of the launch). We will use the upper case letter T to refer to time late in this section to emphasize the fact that it is random (but some nonrandom quantities such as W are also capitalized).



Let d be the average density of searchers per unit area. Typically d would be calculated by some formula like

$$d = Nf/A \tag{4}$$

where N is the number of searchers assigned to patrol the region in which targets can be expected, A is the area of that region, and f is the fraction of the time during which patrolling actually occurs. If a given searcher can patrol for 8 hours out of every 24 and if launches can happen anytime, then f is 1/3. If launches happen entirely at night (12 hours out of 24), then f would be 2/3 because the searchers would patrol only at night. If launches could happen anytime, but if the sweep width W were different between day and night, then determination of f would require solution of a two-person-zero-sum game.

A rough analysis can be made by assuming that *d* is the density of a twodimensional Poisson field of searchers whenever the launch occurs, and that launches are so widely separated in time as to be considered independent events. Probably the latter assumption is worse than the former, since TELs are in reality motivated to make multiple simultaneous launches that will exhaust the patrolling searchers (as well as any defense that might be mounted against the missiles). In any case, the Poisson assumption determines the distribution of *S*, the distance from the launch to the nearest patroller. The event (*S*>*s*) is the event that the number of patrollers in a circle of radius *s* about the launch is 0. Since the average number of patrollers in that circle is  $\pi ds^2$ ,  $Prob(S>s)=exp(-\pi ds^2)$ . Since *T* is related to *S* by  $T=S/V_T$ , where  $V_T$  is the transit speed of the closest searcher, the density function of *T* can be determined by differentiation of the cumulative distribution function to be  $f_T(t)=2\pi t dV_T^2 exp(-\pi dt^2V_T^2)$ . (5)

T turns out to be a Rayleigh random variable. The average detection probability in a flaming datum attack is then

$$E(PD(T)) = \int_{0}^{\infty} \left[ 1 - \exp\left(-V_{S}W / \left(\pi U^{2} t\right) \right) \right] f_{T}(t) dt.$$
(6)

By letting

$$x = V_S V_T W \sqrt{d/\pi} / U^2 , \qquad (7)$$

and substituting  $u = \pi dt^2 V_T^2$  in (6), equation (6) can be expressed in dimensionless form as

$$E(PD(T)) = 1 - \int_{0}^{\infty} \exp\left(-\left(u + x/\sqrt{u}\right)\right) du.$$
(8)



Let (8) be called simply P(x), the detection probability as a function of x. Figure 2 shows the function P(x), calculated numerically, using MATLAB (1990).

Figure 2

If the TBM launch is detected by some system other than the responding aircraft, then some allowance must be made for a communications delay c. The average response time without the delay is  $E(T)=1/(2V_T\sqrt{d})$ . Define an "equivalent transit speed"  $V_T'$  to be such that  $1/(2V_T'\sqrt{d})=c+1/(2V_T\sqrt{d})$ , so that

$$V_T' = V_T / (1 + 2cV_T \sqrt{d}). \tag{9}$$

Then replacing  $V_T$  by  $V'_T$  in calculating x makes a rough correction for the communications delay. A more accurate analysis would acknowledge that the total response time including the communications delay is no longer a Rayleigh random variable when c is included.

In summary, although there are a wide variety of input parameters that influence the detection probability, it is only the dimensionless coverage parameter x in (7) that is ultimately of any importance. Given x, the detection probability is a simple graphical lookup. Conner, et al. (1993) give a more general model of TEL detection that would accept this probability as an input.

#### 4. AN EXAMPLE INVOLVING SCUD DETECTION

Except for W, which depends on the sensors involved and has therefore been chosen arbitrarily, the following assumptions are roughly characteristic of detection of SCUD launches within a country the size of Iraq by a fleet of 4 aircraft:

$$N = 4 \text{ aircraft}$$
  

$$f = .1 = \text{fraction of the time actually on patrol}$$
  

$$A = 100,000 \text{ square miles}$$
  

$$V_S = 100 \text{ mph}$$
  

$$V_T = 300 \text{ mph}$$
  

$$W = 10 \text{ miles}$$
  

$$U = 20 \text{ mph}$$

These parameters result in a coverage parameter x=.846, from which one would conclude that 1-P(.846)=.65, so 65% of the launches should result in ultimate detection by a pursuing aircraft.

If there were a communications delay c=15 minutes=.25 hour, then the equivalent transit speed  $V'_T$  according to (9) would be 231 mph. This would reduce x to .651, and 1-P(.651)=.58. Without the communications delay the average response time is already 50 minutes, so adding 15 minutes to it does not make a lot of difference.

#### 5. REFERENCES

£

1

Conner, G. W., M. Ehlers, and K. Marshall, Countering Short Range Ballistic Missiles, Naval Postgraduate School NPSOR-93-009, 1993.

Cohen, E. A. et al. Gulf Warfare Air Power Survey, Secretary of the Air Force, Washington, D. C., 1993.

Mathworks, Inc., MATLAB User's Guide, 1990.

Washburn, A. R., Search and Detection (2nd edition), Operations Research Society of America (1989).

#### INITIAL DISTRIBUTION LIST

.

. .

.

•

1.	Defense Technical Information Center2 Cameron Station Alexandria, VA 22304-6145
2.	Dudley Knox Library (Code 52)2 Naval Postgraduate School Monterey, CA 93943-5002
3.	Office of Research Administration (Code 08)1 Naval Postgraduate School Monterey, CA 93943-5000
4.	Department of Operations Research (Code OR)1 Naval Postgraduate School Monterey, CA 93943-5000
5.	Prof. Alan Washburn (Code OR/Ws)5 Department of Operations Research Naval Postgraduate School Monterey, CA 93943-5000
6.	Prof. Kneale T. Marshall (Code OR/Mt)
7.	CAPT George W. Conner, USN (Code OR/Co)
8.	Dr. David Israel
9.	Office of the Chief of Naval Operations1 N-813 (Capt. Jack Kerwin) Washington, DC 20350
10.	Center for Naval Analyses