

AD-A274 875

EFFECTS OF POTASSIUM CHANNEL BLOCKERS ON THE NEGATIVE  
INOTROPIC RESPONSES INDUCED BY CROMAKALIM AND PINACIDIL  
IN GUINEA PIG ATRIUM(U) DEFENCE SCIENCE AND TECHNOLOGY  
ORGANIZATION CANBERRA (AUSTRALIA) W LAU

1/1

UNCLASSIFIED

NL







AIM


Association for Information and Image Management

1100 Wayne Avenue, Suite 1100  
Silver Spring, Maryland 20910




301/587-8202



Centimeter



Inches



MANUFACTURED TO AIM STANDARDS  
BY APPLIED IMAGE, INC.

# AD-A274 875



## REPORT DOCUMENTATION PAGE

1

1 AGENCY USE ONLY		2 REPORT DATE 1992		3 TYPE/DATES COVERED	
4 TITLE AND SUBTITLE EFFECTS OF POTASSIUM CHANNEL BLOCKERS ON THE NEGATIVE INOTROPIC RESPONSES INDUCED BY CROMAKALIM AND PINACIDIL IN GUINEA PIG ATRIUM				5 FUNDING NUMBERS	
6 AUTHOR WAI-MAN LAU					
7 FORMING ORG NAMES/ADDRESSES DEFENCE SCIENCE AND TECHNOLOGY ORGANIZATION, MATERIALS RESEARCH LABORATORY, PO BOX 50, ASCOT VALE VICTORIA 3032 AUSTRALIA				8 PERFORMING ORG. REPORT NO	
9 SPONSORING/MONITORING AGENCY NAMES AND ADDRESSES					
11 SUPPLEMENTARY NOTES					
12 DISTRIBUTION/AVAILABILITY STATEMENT <b>DISTRIBUTION STATEMENT A</b>				12B DISTRIBUTION CODE	
13. ABSTRACT (MAX 200 WORDS): THE K <sup>+</sup> CHANNEL OPENERS CROMAKALIM AND PINACIDIL INDUCED A CONCENTRATION-DEPENDENT REDUCTION IN ATRIAL CONTRACTION FORCE WITH EC <sub>50</sub> VALUES OF 25 ± 2 AND 37 ± 2 UMOL/L, RESPECTIVELY. THIS DEPRESSANT EFFECT WAS ANTAGONISED BY 50 UMOL/L TACRINE WHICH DISPLACED THE CONCENTRATION-RESPONSE CURVES OF CROMAKALIM AND PINACIDIL TO THE RIGHT. THE RESPECTIVE DR <sub>50</sub> VALUES WERE 3.8 AND 2.3. INCREASING THE TACRINE CONCENTRATION (100 AND 500 UMOL/L) PRODUCED NO ADDITIONAL EFFECT ON THE CONCENTRATION-RESPONSE RELATIONSHIPS. ADDITION OF 1 UMOL/L ATROPINE ENHANCED THE ANTAGONISM DUE TO TACRINE BY INCREASING THE DR VALUE FROM 3.8 TO 6.5 FOR CROMAKALIM AND FROM 2.3 TO 5.2 FOR PINACIDIL. GLIBENCLAMIDE, AN ATP-SENSITIVE K <sup>+</sup> CHANNEL BLOCKER, COMPETITIVELY INHIBITED THE NEGATIVE INOTROPIC EFFECTS OF CROMAKALIM AND PINACIDIL. THE RESPECTIVE DISSOCIATION CONSTANTS FOR GLIBENCLAMIDE AGAINST CROMAKALIM AND PINACIDIL WERE 0.57 AND 0.35 UMOL/L. NEITHER APAMIN NOR VARIATION IN EXTERNAL CA <sub>2+</sub> CONCENTRATION AFFECTED THE NEGATIVE INOTROPIC EFFECTS OF THE K <sup>+</sup> CHANNEL OPENERS. IT WAS SUGGESTED THAT THE MECHANICAL EFFECTS OF CROMAKALIM AND PINACIDIL ARE MEDIATED THROUGH THE ATP-SENSITIVE K <sup>+</sup> CHANNELS IN THE HEART.					
14 SUBJECT TERMS				15 NUMBER OF PAGES 7	
				16 PRICE CODE	
17 SECURITY CLASS. REPORT UNCLASSIFIED		18 SEC CLASS PAGE UNCLASSIFIED	19 SEC CLASS ABST.	20 LIMITATION OF ABSTRACT	

### 94-01490



998

**DISTRIBUTION STATEMENT A**  
 Approved for public release  
 Distribution Unlimited

**DTIC**  
**ELECTE**  
 JAN 13 1994

**S B D**

# 94 1 12 065

**Pharmacology**

Editors: K.-F. Sewing, Hannover;  
E.S. Vesell, Hershey, Pa.

**Reprint**

Publisher: S. Karger AG, Basel  
Printed in Switzerland

**Original Paper**

Pharmacology 1992;45:9-16

*Wai-Man Lau*

Materials Research Laboratory,  
Defence Science and Technology  
Organisation, Ascot Vale,  
Victoria, Australia

**Effects of Potassium  
Channel Blockers on the  
Negative Inotropic Responses  
Induced by Cromakalim and  
Pinacidil in Guinea Pig Atrium**

**Key Words**

Cromakalim  
Pinacidil  
Tacrine  
Glibenclamide  
ATP-sensitive K<sup>+</sup> channel,  
cardiac  
Atrial contraction  
K<sup>+</sup> channel openers

**Abstract**

The K<sup>+</sup> channel openers cromakalim and pinacidil induced a concentration-dependent reduction in atrial contraction force with EC<sub>50</sub> values of 25 ± 2 and 37 ± 2 μmol/l, respectively. This depressant effect was antagonised by 50 μmol/l tacrine which displaced the concentration-response curves of cromakalim and pinacidil to the right. The respective DR<sub>50</sub> values were 3.8 and 2.3. Increasing the tacrine concentration (100 and 500 μmol/l) produced no additional effect on the concentration-response relationships. Addition of 1 μmol/l atropine enhanced the antagonism due to tacrine by increasing the DR<sub>50</sub> value from 3.8 to 6.5 for cromakalim and from 2.3 to 5.2 for pinacidil. Glibenclamide, an ATP-sensitive K<sup>+</sup> channel blocker, competitively inhibited the negative inotropic effects of cromakalim and pinacidil. The respective dissociation constants for glibenclamide against cromakalim and pinacidil were 0.57 and 0.35 μmol/l. Neither apamin nor variation in external Ca<sup>2+</sup> concentration affected the negative inotropic effects of the K<sup>+</sup> channel openers. It was suggested that the mechanical effects of cromakalim and pinacidil are mediated through the ATP-sensitive K<sup>+</sup> channels in the heart.

**Introduction**

Regulation of potassium channels in cardiac tissues is important in regulating normal heart functions, such as the control of cardi-

ac contractility and the determination of cell membrane potential. Cromakalim (BRL 34915), a benzopyrane, and pinacidil, a cyanoguanidine, belong to a new class of anti-hypertensive agents [1, 2] which relax vascu-

Received  
May 8, 1991  
Accepted  
October 29, 1991

Dr. Tony Wai-Man Lau  
Materials Research Laboratory  
Defence Science and Technology Organisation  
PO Box 50, Ascot Vale 3032, Vic. (Australia)

© 1992  
S. Karger AG, Basel  
0031-7012/92/  
0451-0009\$2.75/0

lar and cardiac smooth muscles by activation of K<sup>+</sup> channels [3, 4]. Increasing the efflux of K<sup>+</sup> results in the hyperpolarisation of the cell membranes, shortening of action potentials and reduction in the force of muscle contractions. These properties of cromakalim and pinacidil form the basis of their anti-hypertensive actions [5].

As there are many types of K<sup>+</sup> channels in the heart, all involved in the regulation of electrophysiological and contractile responses [6]; it will be of interest to find out the type of K<sup>+</sup> channels activated by cromakalim and pinacidil. We have previously shown that the centrally acting anti-cholinesterase, tacrine (THA), blocks the K<sup>+</sup> channels in the heart [7]. Glibenclamide, a sulphonylurea, was reported to antagonize cromakalim in guinea pig isolated trachealis muscle [8] and to block the ATP-dependent K<sup>+</sup> channel in insulin-secreting cells [9]. Studying the antagonism of the K<sup>+</sup> channel openers with tacrine and glibenclamide might provide information to characterise the K<sup>+</sup> channels activated by cromakalim and pinacidil in guinea pig atrium. In this study, the effects of potassium channel blockers on the atrial muscle relaxation induced by cromakalim and pinacidil were investigated. The results obtained are interpreted in relation to their activities at the K<sup>+</sup> channels.

## Materials and Methods

### *Guinea Pig Atrial Preparations*

Left atrial preparations were surgically removed from female guinea pigs weighing 250–400 g as described by Freeman and Turner [10]. Preparations were then placed in an organ bath containing heart Ringer's solution (composition mmol/l: NaCl 115, KCl 4.6, CaCl<sub>2</sub> 1.8, MgSO<sub>4</sub> 1.2, NaH<sub>2</sub>PO<sub>4</sub> 1.2, NaHCO<sub>3</sub> 22, glucose 22, pH = 7.4) at 37 °C. The buffer was bubbled with carbogen (95% O<sub>2</sub> and 5% CO<sub>2</sub>) continuously. The preparations were allowed to equilibrate for 30 min before the experiment commenced.

The atria were stimulated at 2 Hz and 10 V, and the tension of isometric contraction was recorded by a Shinkoh U-gauge (UL-2-120). The signal outputs were modulated by a Coulbourn transducer (S72-75) and recorded by a Graphtech linear recorder (WR-3071). Tension development was constant within the period of the experiments which lasted for approximately 3 h.

### *Data Analysis*

The concentration-response curves were constructed by adding drugs cumulatively to the organ bath. Data are presented as means ± SEM and analysed according to the methods (ANOVA and Bonferroni) suggested by Wallenstein et al. [11].


### *Drugs*

Cromakalim was donated by Beecham, UK, and pinacidil was a gift from Leo, Denmark. Tacrine was obtained from the Institute of Drug Technology, Melbourne, Australia. Apamin, atropine and glibenclamide were purchased from Sigma, St. Louis, Mo., USA.

## Results

### *Effects of Cromakalim and Pinacidil on Atrial Contractility*

Both cromakalim and pinacidil depressed the force of atrial contraction in a concentration-dependent manner. The effect of cromakalim was just perceptible at 10 µmol/l and rapidly reached a maximum suppression of approximately 80% of the control at 40 µmol/l. The effective concentration to reduce the force of contraction by 50% (EC<sub>50</sub>) for cromakalim was 25 ± 2 µmol/l (fig. 1). Pinacidil showed a similar concentration-response relationship to cromakalim except that its potency was slightly weaker. This is indicated by the higher EC<sub>50</sub> of 37 ± 2 µmol/l for pinacidil (fig. 1). There is however one noticeable difference between the negative inotropic effects of cromakalim and pinacidil. It took more than 30 min for the force of contraction to return to the control level after the



**Fig. 1.** The negative inotropic effects of cromakalim (●) and pinacidil (▲). Data are obtained from six experiments. Means  $\pm$  SEM.

**Table 1.** Effects of varying concentrations of tacrine, and tacrine (100  $\mu$ mol/l) plus atropine (1  $\mu$ mol/l) on the negative inotropic responses induced by cromakalim and pinacidil

	Control	Tacrine			Tacrine (100 $\mu$ mol/l) + atropine (1 $\mu$ mol/l)
		50 $\mu$ mol/l	100 $\mu$ mol/l	500 $\mu$ mol/l	
Pinacidil					
EC <sub>50</sub> , $\mu$ mol/l	36.8 $\pm$ 1.3	85.4 $\pm$ 3.1*	86.5 $\pm$ 4.2*	84.2 $\pm$ 0.5*	192.1 $\pm$ 2.2*, **
Cromakalim					
EC <sub>50</sub> , $\mu$ mol/l	27.5 $\pm$ 1.1	105.6 $\pm$ 2.6*	106.1 $\pm$ 2.7*	106.9 $\pm$ 1.0*	178.9 $\pm$ 1.7*, **

Each data point is calculated from five experiments and is expressed as mean  $\pm$  SEM. Data are analysed by ANOVA and Bonferroni methods according to Wallenstein et al. [11]. \*  $p < 0.01$  vs. control, \*\*  $p < 0.01$  vs. tacrine alone.


wash-out of pinacidil, but the depression was totally and rapidly reversed when cromakalim was removed.

#### *Interaction with Tacrine*


Tacrine at 50  $\mu$ mol/l caused a parallel shift of the concentration-response curves of pinacidil and cromakalim to the right. The dose ratio at 50% reduction in the force of atrial

contraction (DR<sub>50</sub>) for cromakalim was 3.8, while that for pinacidil was 2.3. Increasing the concentration of tacrine to 100 and 500  $\mu$ mol/l did not produce an additional effect on the concentration-response curves of cromakalim and pinacidil (table 1), suggesting the antagonism by tacrine is independent of its concentration.

**Fig. 2.** The effects of 100  $\mu\text{mol/l}$  tacrine ( $\square$ ), and 100  $\mu\text{mol/l}$  tacrine + 1  $\mu\text{mol/l}$  atropine ( $\blacksquare$ ) on the negative effect of pinacidil.  $\bullet$  = Controls. Data are obtained from five experiments. Means  $\pm$  SEM.




**Fig. 3.** The effects of 100  $\mu\text{mol/l}$  tacrine ( $\square$ ); and 100  $\mu\text{mol/l}$  tacrine + 1  $\mu\text{mol/l}$  atropine ( $\blacksquare$ ) on the negative inotropic effect of cromakalim.  $\bullet$  = Controls. Data are obtained from five experiments. Means  $\pm$  SEM.



*Interaction with Tacrine and Atropine*  
To investigate if the cholinergic effects of tacrine are involved in antagonising the depressant effects induced by cromakalim and pinacidil, their concentration-response relationships were studied in the presence of 100  $\mu\text{mol/l}$  tacrine and 1  $\mu\text{mol/l}$  atropine, an anti-muscarinic agent. The concentration-response curves for cromakalim and pinacidil

were both shifted further to the right (fig. 2, 3). The  $\text{DR}_{50}$  value was increased from 3.8 to 6.5 for cromakalim and from 2.3 to 5.2 for pinacidil.

*Interactions with Glibenclamide*  
Glibenclamide itself produced a concentration-dependent reduction in atrial force of contraction. The effect was just perceptible at



**Fig. 4.** Schild plot of the antagonism of the negative inotropic effect of pinacidil (a) and cromakalim (b) by glibenclamide. Data are obtained from thirteen (a) and twelve experiments (b). Means  $\pm$  SEM. DR = Dose ratio.

1  $\mu\text{mol/l}$  which reduced the force of the control by 8% and reached the maximum at 20  $\mu\text{mol/l}$  with a reduction of 55%. The negative inotropic effects of cromakalim and pinacidil were both antagonised by glibenclamide by displacing their concentration-response curves to the right in a concentration-dependent manner (1–50  $\mu\text{mol/l}$ ). Analysis of the data with Schild plot [12] showed a linear relationship between the (dose ratio-1) and glibenclamide concentrations against the two potassium channel openers (fig. 4). The slope of the plot for cromakalim was 0.90 and for pinacidil was 0.88 which were not significantly different from unity ( $p < 0.05$ ). The apparent dissociation constant ( $K_a$ ) for glibenclamide was 0.57  $\mu\text{mol/l}$  against cromakalim and 0.35  $\mu\text{mol/l}$  against pinacidil. These comparable dissociation constants suggested that they may be acting at a common site.

#### *Interactions with Apamin and Exogenous $\text{Ca}^{2+}$*

Apamin (10–100 nmol/l) had no effect on the depressant effects of cromakalim and pinacidil. Similarly, reducing the  $\text{Ca}^{2+}$  content to one third of the original strength of the heart Ringer's solution or increasing it by 50% produced no effect on the negative inotropism induced by the two  $\text{K}^+$  channel openers.

#### **Discussion**

Activation of the  $\text{K}^+$  channels in cardiac sarcolemma by cromakalim and pinacidil has been suggested to cause a shortening of action potential, reducing the time for  $\text{Ca}^{2+}$  influx and thus resulting in a reduction in the contraction force [13]. Our results for cromakalim and pinacidil agreed with this proposal. Both of them, even at higher concentrations, could not completely abolish atrial contraction. This is different to the suppression caused by diltiazem and nifedipine, both of which are  $\text{Ca}^{2+}$  channel blockers and are capa-



ble of totally relaxing atrial muscle [14]. The residual contraction of the atrium in the presence of high concentrations of cromakalim and pinacidil is thus an indication that these  $K^+$  channel openers do not directly affect the inward  $Ca^{2+}$  current. The potency of the relaxing properties of cromakalim and pinacidil in guinea pig atrium is much weaker than that reported in canine atrial muscle [15], in guinea pig trachea [16, 17], in rabbit arteries [18] and in human skeletal muscle fibres [19]. It is not known whether the difference in potency is linked to differences in channel properties between different tissues or species.

Tacrine antagonized the cromakalim- and pinacidil-induced negative inotropic effects in a concentration-independent manner (table 1). This effect is apparently not arisen from a simple competitive antagonism at a site common to tacrine, cromakalim and pinacidil. Tacrine was found to block the inwardly rectifying  $K^+$  current in guinea pig isolated ventricular myocytes [20] and inhibit the slow outward  $K^+$  current in the peptidergic neurons of snail [21]. If these are the primary action sites for tacrine, our results suggest that cromakalim and pinacidil do not activate these  $K^+$  channels. The uncompetitive antagonism observed is probably a functional rather than a specific one. Tacrine has been reported to inhibit acetylcholinesterase and to block muscarinic acetylcholine (M-ACh) receptors [7, 22]. Inhibition of the cholinesterase by sub-micromolar concentration of tacrine ( $K_i = 1.8 \mu\text{mol/l}$ ) would result in a transient accumulation of acetylcholine and increase the frequency of M-ACh receptor activation. This would lead to an increase in  $K^+$  efflux, thereby opposing the effects of  $K^+$  channel blockade due to tacrine. However, at  $100 \mu\text{mol/l}$  tacrine, all M-ACh receptors would have been blocked. Thus the enhancement of tacrine by atropine to antagonise the

negative inotropic effects of the two  $K^+$  channel openers is therefore inexplicable in the context of M-ACh receptor blockade.

Atropine is a potent muscarinic antagonist, with an affinity of  $2 \times 10^9 \text{ l/mol}$  for M-ACh receptors [23]. It may be possible that excessive atropine ( $1 \mu\text{mol/l}$ ) used in this study displaced tacrine at the M-ACh sites, releasing free tacrine to block the  $K^+$  channels. Additional work is required to clarify this point.

Glibenclamide had been found to block ATP-sensitive  $K^+$  channels in insulin-secreting cells [9], pancreatic  $\beta$ -cells [24] and cardiac muscles [25]. It was also reported that glibenclamide antagonised competitively with cromakalim and pinacidil in vascular tissues and airway smooth muscles [26, 27]. Our results demonstrated that glibenclamide antagonised the negative inotropic effects of cromakalim and pinacidil on guinea pig atrium. The Schild plots give a slope of 0.90 for cromakalim and 0.88 for pinacidil suggesting the antagonism is of the simple competitive type. The apparent dissociation constants for glibenclamide to act against cromakalim ( $0.57 \mu\text{mol/l}$ ) and pinacidil ( $0.35 \mu\text{mol/l}$ ) agree well with the values reported in other studies [27, 28]. It is also noted that the values of the two constants are comparable to each other, hence suggesting that cromakalim, pinacidil and glibenclamide could interact at a common site. This site is most likely the atrial ATP-sensitive  $K^+$  channels which are also the primary site for glibenclamide [25]. Further experiments in the presence of apamin, which blocks small conductance  $Ca^{2+}$ -activated  $K^+$  channels [29], or by varying the concentration of exogenous  $Ca^{2+}$  produced no changes in the muscle relaxation by cromakalim and pinacidil. It appears that cromakalim and pinacidil do not open  $Ca^{2+}$ -activated  $K^+$  channels.

The current study demonstrated that the mechanical effects of cromakalim and pinacidil were inhibited competitively by glibenclamide; uncompetitively by tacrine and were unaffected by apamin or by varying the exogenous  $Ca^{2+}$  contents. The excitation-contraction coupling in atrial muscles has not been fully understood. If the mechanical effects of cromakalim and pinacidil are primarily associated with their potassium channel opening

properties, it can be established that they mediate the depression through opening of the ATP-sensitive  $K^+$  channels.

### Acknowledgement

The author wishes to thank Dr. Shirley Freeman for reviewing the manuscript, Beecham, UK, for a gift of cromakalim and Leo, Denmark, for a gift of pinacidil.

### References

- 1 Cohen ML, Colbert WE: Comparison of the effects of pinacidil and its metabolite pinacidil-N-oxide in isolated smooth and cardiac muscle. *Drug Dev Res* 1986;7:111-124.
- 2 Cook NS: The pharmacology of potassium channels and their therapeutic potential. *Trends in Pharmacol Sci* 1988;9:21-28.
- 3 Scholtysek G: Evidence for inhibition by BRL 34915 of  $K^+$  conductance in cardiac muscle. *Naunyn Schmiedebergs Arch Pharmacol* 1987;335:692-696.
- 4 Osterrieder W: Modification of  $K^+$  conductance of heart cell membrane by BRL 34915. *Naunyn Schmiedebergs Arch Pharmacol* 1988;337:93-97.
- 5 Weston AH, Bray KM, Duty S, McHarg AD, Newgreen DT, Johnson JS: In vitro studies on the mode of action of pinacidil. *Drugs* 1988;36(suppl 7):10-28.
- 6 Pelzer D, Trautwein N: Currents through ionic channels in multicellular cardiac tissue and single heart cells. *Experientia* 1987;43:1153-1162.
- 7 Freeman SE, Lau WM, Szilagyi M: Blockade of a cardiac  $K^+$  channel by tacrine: Interactions with muscarinic and adenosine receptors. *Eur J Pharmacol* 1988;154:59-66.
- 8 Murray MA, Boyle JP, Small RC: Cromakalim-induced relaxation of guinea-pig isolated trachealis: Antagonism by glibenclamide and by phentolamine. *Br J Pharmacol* 1989;98:865-874.
- 9 Schmid-Antomarchi H, Weille J, Fosset M, Lazdumski M: The receptor for antidiabetic sulfonylureas controls the activity of the ATP-modulated  $K^+$  channel in insulin-secreting cells. *J Biol Chem* 1987;262:15840-15844.
- 10 Freeman SE, Turner RJ: Phase-plane trajectories of atrial cell action potentials: Effects of temperature reduction. *Cardiovasc Res* 1974;8:451-459.
- 11 Wallenstein S, Zucker CL, Fleiss JL: Some statistical methods useful in circulation research. *Circ Res* 1980;47:1-9.
- 12 Arunlakshana O, Schild HO: Some quantitative uses of drug antagonists. *Br J Pharmacol* 1959;14:48-58.
- 13 Yanagisawa T, Hashimoto H, Taira N: The negative inotropic effect of nicorandil is independent of cyclic GMP changes: A comparison with pinacidil and cromakalim in canine atrial muscle. *Br J Pharmacol* 1988;95:393-398.
- 14 Fleckenstein A: Calcium antagonists and calcium agonists: Fundamental criteria and classification; in Fleckenstein A, Van Breemen C, Grop R, Hoffmeister F (eds): *Cardiovascular Effects of Dihydropyridine-Type Calcium Antagonists and Agonists*. Berlin, Springer, 1985, pp 3-31.
- 15 Yanagisawa T, Hashimoto H, Taira N: Interaction of potassium channel openers and blockers in canine atrial muscle. *Br J Pharmacol* 1989;97:753-762.
- 16 Allen SL, Boyle JP, Cortiji J, Foster RW, Morgan GP, Small RC: Electrical and mechanical effects of BRL 34915 in guinea-pig isolated trachealis. *Br J Pharmacol* 1986;89:395-405.
- 17 Mellekjaer S, Nielsen-Kudsk JE, Nielsen CB, Sigsgaard C: A comparison of the relaxant effects of pinacidil in guinea-pig trachea, aorta and pulmonary artery. *Eur J Pharmacol* 1989;167:275-280.
- 18 Post JM, Smith JM, Jones AW: BRL 34915 (cromakalim) stimulation of  $^{42}K$  efflux from rabbit arteries is modulated by calcium. *J Pharmacol Exp Ther* 1989;250:591-597.
- 19 Spuler A, Lehmann-Horn F, Gräfe P: Cromakalim (BRL 34915) restores in vitro the membrane potential of depolarized human skeletal muscle fibres. *Naunyn Schmiedebergs Arch Pharmacol* 1989;339:327-331.

- 20 Osterrieder W: 9-Amino-1,2,3,4-tetrahydroacridine (THA) is a potent blocker of cardiac potassium channels. *Br J Pharmacol* 1987;92:521-525.
- 21 Drukarch B, Kits KS, Vander Meer EG, Lodder JC, Stoof JC: 9-Amino-1,2,3,4-tetrahydroacridine (THA) an alleged drug for the treatment of Alzheimer's disease inhibits acetylcholinesterase activity and slow outward K<sup>+</sup> current. *Eur J Pharmacol* 1987;141:153-157.
- 22 Ho AKS, Freeman SE: Anticholinesterase activity of tetrahydroaminacrine (THA) in regard to succinylcholine hydrolysis. *Nature* 1965; 205:1118-1119.
- 23 Birdsall NJM, Hulme EC: Biochemical studies on muscarinic acetylcholine receptors. *J Neurochem* 1976;27:7-16.
- 24 Zunkler BJ, Lenzen S, Manner K, Panten U, Trube G: Concentration-dependent effects of tolbutamide, meglitinide, glipizide, glibenclamide and diazoxide on ATP-regulated K<sup>+</sup> currents in pancreatic B-cells. *Naunyn Schmiedeberg's Arch Pharmacol* 1988;337:225-230.
- 25 Sanguinetti MC, Scott AL, Zingaro GJ, Siegl PKS: BRI 34915 (cromakalim) activates ATP-sensitive K<sup>+</sup> current in cardiac muscle. *Proc Natl Acad Sci USA* 1988;85:8360-8364.
- 26 Nielsen-Kudsk JE, Bang L, Bronsgaard AM: Glibenclamide blocks the relaxant action of pinacidil and cromakalim in airway smooth muscle. *Eur J Pharmacol* 1990;180:291-296.
- 27 Masuzawa K, Matsuda I, Asano M: Evidence that pinacidil may promote the opening of ATP-sensitive K<sup>+</sup> channels yet inhibit the opening of Ca<sup>2+</sup>-activated K<sup>+</sup> channels in K<sup>+</sup>-contracted canine mesenteric artery. *Br J Pharmacol* 1990;100:180-184.
- 28 Quast U, Cook NS: In vitro and in vivo comparison of two K<sup>+</sup> channel openers, diazoxide and cromakalim, and their inhibition by glibenclamide. *J Pharmacol Exp Ther* 1989; 250:261-270.
- 29 Blatz AL, Magleby KL: Single-apamin-blocked Ca-activated K<sup>+</sup> channel of small conductance in cultured rat skeletal muscle. *Nature* 1986;323:718-720.

DTIC QUALITY INSPECTED 8

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	21

**END  
FILMED**

**DATE:**

*2-94*

**DTIC**