ARMY PROJECT ORDER NO: 90PP0807

TITLE:



AD-A276 135

PRINCIPAL INVESTIGATOR: Ronald I. Herning, Ph.D.

CONTRACTING ORGANIZATION: National Institutes of Health NIDA/Addiction Research Center Francis Scott Key Medical Center P.O. Box 5180 Baltimore, Maryland 21224

REPORT DATE: November 4, 1993

TYPE OF REPORT: Final Report

- PREPARED FOR: U.S. Army Medical Research and Development Command, Fort Detrick Frederick, Maryland 21702-5012
- DISTRIBUTION STATEMENT: Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

94-05763

DTIC QUALITY INGRECTED L

2

23

Best Available Copy

r			_	
REPORT DOCUMENTATION PAGE			Fi O	orm Approved MB No. 070 4-0188
Public reporting burden for this collection of gathering and maintaining the data needed, collection of information, including suggestic Davis Highway, Suite 1204, Arlington, VA 22	information is estimated to average 1 hour pe and completing and reviewing the collection of his for reducing this burden to Washington He 2024302, and to the Office of Management and	r response, including the time for re information – Send comments rega adduarters Services, Directorate fo a Budget, Paperwork Reduction Proj	rviewing instruct rging this burder r information Op ect (0704-0188), 1	ions, searching existing data sources, i estimate or any other aspect of this erations and Reports, 1215 Jefferson Washington, DC 20503
1. AGENCY USE ONLY (Leave b)	ank) 2. REPORT DATE	3. REPORT TYPE AN	D DATES CO	VERED
A TITLE AND SUBTITLE	4 November 199	3 Final Repor	t (11/1)	5/89-9/30/92) G NUMBERS
Performance Norm Use in Males (NP	s on NPPAB for Lim PABNRM)	ited Drug	Army Project Order 90PP0807	
6. AUTHOR(S)			060300	22
Ronald I. Herning, Ph.D.			3M6300 WUDA33	2D995.BG.200 5390
7. PERFORMING ORGANIZATION	NAME(S) AND ADDRESS(ES)		8. PERFOR	MING ORGANIZATION
National Institu NIDA/Addiction R Francis Scott Ke P.O. Box 5180	tes of Health esearch Center y Medical Center		REPORT	NUMBER
Baltimore, Maryland 21224 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research & Development Command Fort Detrick Frederick, Maryland 21702-5012			10. SPONSC AGENCI	RING/MONITORING / REPORT NUMBER
11. SUPPLEMENTARY NOTES	STATEMENT		12b. DISTRI	BUTION CODE
Approved for pub distribution unl	lic release; imited			
13. ABSTRACT (Maximum 200 wo	rds)			
A Neurophysiological Performat tasks was designed by the Off for cognitive ERPs for tasks P2 and P3 components of cognitive two Continuous Performance ta used to test within day (2 te components recorded in this bill exceeded the reliability of t letter set size. The measur appears stable enough to built data will be useful in the ap in using this normative ERP of (N=10) who denied illicit drug	Ince Assessment Test (NPAT) bat ice of Miliary Performance Ass in this battery was obtained fi tive event-related potential w sks and the Sternberg Memory t sts on the same day) and betwee attery of tasks was high (.7 to he percent correct values in a rement of the ERP components, d norms for the ERPs components, d norms for the ERPs components plication of NPAT battery. We ata we were able to identify co g use.	tery of sensory and cog essment Technology (OPM rom 37 normal subjects. ere determined for the ask with two set sizes. en day reliability. The 0.9). The reliability ll tasks except the Ster in particular the P3, in s for each of the tasks but only was the test-rei ognitive processing alto	nitive even MT). Test- The test- auditory ran The Cronbu- e test-retes of the P3 a rnberg memory a this batton in the batt liability of erations in	t-related potential (ERP) retest reliability data retest reliability of N1, re event monitoring task, ach alpha coefficient was st reliability of the ERP amplitude measurements ry task with a three ery of cognitive tasks tery. Such normative ERP f the ERPs data high, but a group of drug abusers
14 SURIECT TERMS			I 16	
Performance, Dru	gs, Neurophysiolog	y, Sensory Evo	ked	
Potential, Cogni	tive Evoked Potent	ials Neurophys	iolo-	. PRICE CODE
GIC NOTES, COGNI 17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	KA V 19. SECURITY CLASSIFIC OF ABSTRACT	ATION 20	LIMITATION OF ABSTRACT
Unclassified NSN 7540-01-280-5500	Unclassified	Unclassified	Stand	Unlimited and Form 298 (Rev 2-89)

,

²⁹⁸⁻¹⁰²

FOREWORD

Opinions, interpretations, conclusions, and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

<u>Citations of commercial organizations and trade names in this</u> report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) have adhered to policies of applicable Federal Law 45CFR46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institute of Health.

Investigator's Signature

Accesi	on For	
NTIS DTIC Unann Justific	CRA&I TAB ounced cation	6 0 0
By Distrib	ution (
A	vailability	Codes
Dist A-1	Avail and Specia	/ 01

FINAL REPORT

 Army Project Order: 90PP0807
 Report Date: 4 Nov 93
 Reporting Period: 15 November 1989 - 30 September 1992
 PI.: Ronald I. Herning, Ph.D. 5. Telephone No. (410) 550-1477
 Agency: National Institute on Drug Abuse, Addiction Research Center, P.O. Box 5180, Baltimore MD. 21224

7. Project Title: Drug Use in Males (NPPABNRM)

8. Current Staff with percent of effort on each project.

Herning, R.I. - 50% Guo, X - 30%

Koeppl, B -30% Graber, E. (unpaided volunteer)-10%

9. Approximately PI expenditures 1 Sept 1991 to date:

Personal Services: 0 Travel: 0 Other Services: 0

10. Comments on administrative and logistical matters:

Both Dr. Brigham and Ms. Glover, who were responsible for subject testing, left the project in 1991. The loss of key personnel made it impossible to continue testing on this project at that time. Testing was further delayed when the acting Etiology Branch chief was replaced by a permanent chief. In October 1992, Dr. Herning was transferred to the Medical Affairs Branch where the project was completed.

iii

Table of Contents

Page	No.

I.	Test-Retest Event-Related Potential Data1
II.	Introduction1
III.	Methods3
IV.	Results
v.	Discussion and Conclusions21
VI.	References25

ing in Northern

iv

i

Introduction

The use of Event Related Potential (ERP) methodology in the assessment of cognitive processing and performance has excellent Sensory and cognitive stages of human information potential. processing can be directly measured by brain potentials during cognitive tasks. These ERP measures complement more traditional measures of reaction time and correctness. Brain potentials are related to cognitive processes as such as selective attention (Hanson & Hillyard, 1980; Naatanen et al., 1993; Teder et al., 1993), working memory (Gevin & Cutillo, 1993), stimulus evaluation (Donchin & Coles, 1988; Polich, 1989) and other aspects of information processing (Johnson, 1993). The brain potentials are sensitive to the effects of drugs (Herning et al., 1979; 1985, 1987; Coons et al., 1981; Peloquin and Klorman, 1986; Brunaghin et al., 1987; Fitzpatrick et al., 1988), fatigue or habituation (Knott, 1985; Ditraglia & Polich, 1991), changes in dual task performance (Wickens et al., 1983) and other conditions of interest to the military.

The ERP methodology is not new. Its application to military performance questions is, likewise, not new. Its application to applied problems has suffered from a lack of standardization of task parameters, ERP recording procedures and thorough characterization of the subject populations.

Some guidelines for standardization were outlined under the Office of Miliary Performance Assessment Technology (OPMAT). A test battery of sensory and cognitive ERP tasks was designed with

specific stimulus parameters and recording procedures (Reeves et al., 1991). The sensory ERP paradigms have norms since they are also used in clinical diagnosis of medical disease. Test-retest reliability was been reported for the auditory rare event monitoring task or oddball task as it is often called (Sklare & Lynn, 1984; Polich, 1986; Fabianai et al., 1987; Segalowitz & Barnes, 1993), but not the other cognitive tasks. Whether or not norms for cognitive ERPs can be constructed remains unclear. This project was designed to collect normative ERP data on some of neurophysiologic cognitive tasks specified in the OMPAT, Level 1, battery as well as provide test-retest reliability data on normal non-drug using males. The project was extended to females.

The research plan called for testing 40 to 50 non-drug using subjects on four tasks from the Neurophysiologic Workload Test The tasks included the Auditory Rare Event Battery (NWTB). Monitoring Task, the Continuous Performance Task (two versions) and the Sternberg Memory Task with two memory loads. These cognitive ERP tasks were administered and ERP data was collected using OPMAT quidelines. In addition, DSM-IIIR interviews and other psychometric testing were obtained to screen for current psychiatric illness and drug use. An urine toxicology for illict drugs was also obtained. Half of the subjects were tested on two day study to evaluate the test-retest reliability of these brain potentials over two test days. The other half of the subjects were tested to evaluate test-retest reliability on the same day. The ERP data from the first day of the two day study will be combined

with the one study to increase the sample size.

The collection of normative ERP data on non-drug using persons is complicated by cigarette smoking. While nicotine is certainly is a drug, normative ERP data on smokers who do not use other drugs is of importance. Nicotine is particular problematic since the effects of nicotine deprivation on the EEG and performance and their reversal by smoking are well documented (Knott, 1985). We proposed to extend our collection of normative cognitive ERP data to include non-deprived smokers. We feel the development of norms on these cognitive tasks for both smokers and non smokers may increase the success of this project by reducing the variability in the otherwise non-drug using group.

Methods

Subjects

Fifty-six subjects passed a screening interview designed and administered by Nova Research, a contractor responsible for obtaining human subjects for research at the Addiction Research Center. Nine subjects of these subjects failed show up for testing or were screened out for current psychiatric problems using the Computerized Diagnostic Interview, Revised (CDISR: Blouin, 1985) with a followup interview or NIMH Diagnostic Interview Schedule, Version III Revised (DSM-IIIR: Robins et al., 1989). Since the urine toxicologies lagged behind ERP testing, ten additional

subjects, who reported no illicit drug use, but had positive urine toxicologies were tested on first day of ERP test battery. The ERP and performance data from this group were compared to the non-drug using subjects, but the data was not used in the test-retest reliability calculations.

The second second

Thus, we collected normative ERP data on 37 subjects who (1) had no current Axis I (except nicotine dependence) or Axis II diagnosis as determined by an interview using DSM-IIIR criteria, (2) were healthy (health history interview given by a nurse or counsellor) and (3) had a negative urine toxicology screen for amphetamine, barbiturates, benzodiazepines, cocaine, opiates, phencyclidine and cannabinoids. The mean age of the sample was 27.5 years (5.6 years standard deviation) and the mean years of formal education was 12.5 years (2.1 years standard deviation). Eighteen males and nineteen females were tested. Twenty of the 37 subjects were tested on the cognitive ERP tasks (Auditory Rare Event Monitoring Task, two Continuous Performance Tasks, Sternberg Memory Task (2 set sizes) on two different days with two test sessions each day. The remainding subjects were tested on the one day test-retest procedure using the same tasks administered twice Sixteen of the 37 subjects were nicotine on the same day. dependent and were allowed to smoke if they chose to any time during the study except during of the 25 minute test battery. They were asked if they wanted to smoke before and after the ERP test battery.

Experimental Procedures

On each day of ERP testing, the subjects completed the St. Mary's Sleep Questionnaire, the Profile of Mood State questionnaire and the Beck Depression Inventory. We evaluated test-retest reliability of the ERP measures on the same day and between test days. On the first day test day, the subjects were instructed on the tasks and given a practice test before the actual test session. One hour after the start of the first test session they were retested on each task in the battery. The second test day was similar except that there were no practice sessions.

Cognitive Information Processing Tasks. In the Auditory Rare Event Monitoring (AREM) Task, tones were delivered binaurally through TDH-39 earphones. The subject was required to count rare tones from a series of 1000 Hz and 2000 Hz tones (80 dB SPL, 50 ms in duration, 2.0 s between tones). The 1000 Hz or rare tones occurred about 20% of the time. After the subject heard the series of tones for four minutes, he reported his count to the researcher at the end of the session. A practice session preceded the first formal test session. The N100, P200 and P300 of AREM were measured from the frequent and rare tone ERP. The ERP peaks were measured in the following latency ranges: N100 (50-180 ms), P200 (100-250 ms) and P300 (250-700 ms).

The Sternberg Memory Task was originally developed by Sternberg (1975). In this task, evoked responses were elicited visually by presenting the letter on a TV monitor. The subject was required to memorize three or six letter sets which appeared on the monitor for 30 seconds. A series of probe letters was then displayed on the screen, one at a time. The probe letters subtended 10° of visual angle, remained on the screen for 600 ms and were presented at a rate of one every two seconds. The mean luminance of the screen was 40 cd/m². If the probe letter was one that was part of the memory set, then the subjects were instructed to press a button with their preferred hand. When a presented letter was not in the memory set, then the subjects were required to press a button with their non preferred hand. A practice session proceeded the first formal test session. P200 and P300 components of the Sternberg Memory Task were measured using the same latency ranges as the AREM Task. Task performance was evaluated by reaction time and percent correct.

Two Continuous Performance Tasks (CPT) were also performed. They were not in the OPMAT, Level 1, battery. The first CRT task or CPT-X was similar to the Sternberg Memory Task with a one letter memory set. During this task the subjects were required to press the push button with their preferred hand when they saw an X on the TV monitor. The stimulus characteristics were identical to the Sternberg task. No practice session was given. The second CPT task was the Paired Letter CPT. During this task the subjects were required to press the push button with their preferred hand when they saw an letter repeat (two of the same letter in a row) on the The stimulus characteristics were identical to the TV monitor. Sternberg task and no practice session was given. The P200 and P300 components of the visual ERP were measured an in the Sternberg

Memory Task. Task performance was evaluated by reaction time and percent correct.

BRP Recording and Measurement. The ERPs were recorded from F_z, C_z, and P_z (the International 10-20 System: Jasper, 1958). The reference electrode was on right ear tip. Electrooculogram (EOG) was recorded from side of the left eye and from above the left eye referred to the left ear tip in order to monitor eye movement Isoground electrode was placed on forehead. artifacts. Silver/silver chloride electrodes were used at all locations. Each electrode impedance was kept below 10 K ohm. Trials with EOG artifact were rejected. Testing was performed in a sound attenuated, electrically shielded chamber. The EEG and EOG were recorded with Grass (Model 75P11) amplifiers with a .1 to 100 Hz half amplitude band pass and a 60 Hz notch filter. The EEG and EOG channels were sampled at 200 Hz/channel with 8 microsecond interchannel delay. Single trial EEG and EOG data was saved on The EEG and EOG were processed for artifact and averaged line. after the test session. The target and non-target stimuli were averaged separately for each task.

Statistical Analysis. Separate Analyses of Variance (ANOVA) were performed on each ERP and performance measure from the four tasks (AREM, CRT-X, CPT-Paired Letter, Sternberg Memory Task). These ANOVAs were used to test for significant differences between test sessions within a day (N=37) and between test days (N=20). Each ANOVA included the between subject (nonsmoker versus smoker) factor and within subject factors: stimulus type (target versus

non-target), set size (3 versus 6 letter, Sternberg Task only) and electrode (F_z, C_z, P_z) . The ANOVAs also guided in the calcolation of the reliability coefficients. If the smokers and non-smokers significantly differed, separate coefficients would be calculated for each group.

Tests of the assumptions for within subject ANOVAs were made as recommended by Vasey and Thayer (1987) and corrected probability values (Greenhouse & Geisser, 1959) are reported in all cases.

The test-retest reliability of the ERPs collected in each of the tasks was determined with the intraclass reliability coefficient (Winer, 1972) and Cronbach's alpha coefficient (Nunnally, 1978). Both analyses were similar and we report only the alpha coefficients.

Finally, the ERPs and performance of the drug abusers who denied using illicit drugs were compared to that of non-drug using subjects. This analysis was not originally planned, this group is of interest to both the military and the ARC. These subjects denied using drugs on three occasions (telephone interview, on site screening interview and the psychiatric interview), but they had illicit drugs in their urine the day that they were tested on the ERP test battery. Thus, they provided a comparison group who might have altered ERPs and performance.

Results

Comparison of Non-Smokers with Smokers. Of the many

8

statistical tests made in the one day and two day comparisons, few significant differences were noted between non-smokers and smokers (the group factor) in the ERP components. The smokers differed from non-smokers on the amplitude of the P2 component of the ERP in the visual tasks. For P2 amplitude in the CPT-X and CPT-Paired Letter tasks, the group main effect was significant for the one day (p<.01)and a number of interactions involving group were statistically significant for the two day comparisons. P2 amplitude for the smokers was larger than the non smokers. **P2** latency was also longer for the smokers than the non-smokers in the CPT-X task (significant main effect for group), in the CPT-Pair the letter task (significant main effect for group) and in Sternberg task (significant group by electrode interaction). For N1 latency, the group by test time by electrode interaction was significant for two day (p<.01), but not the one day comparisons in the AREM task. For P3 amplitude in the CPT-Paired Letter task, the group by day by time interaction was a significant in the two day comparison, but there was not indication of group differences in P3 amplitude in the one test where the sample size was larger.

Comparisons between Test Times. Within day statistically significant differences between test sessions were observed only for P2 amplitude or latency in all tests. However, no differences were observed between test session for N1 amplitude, N1 latency, P3 amplitude, P3 latency, reaction time and percent correct. P3 amplitude and latency at the P_z for the target stimulus in all the tasks is plotted in Figures 1 and 2 respectively. Reaction time to







P3 Latency



Figure 2. P3 latency at Pz is plotted for each task. Bars are standard errors. The left column of each pair is time 1. The right is time 2.

- - -

REACTION TIME



Figure 3. Reaction time is plotted for each task. Bars are standard errors. The left column of each pair is time 1. The right is time 2.

PERCENT CORRECT



Figure 4. Percent correct is plotted for each task. Bars are standard errors. The left column of each pair is time 1. The right is time 2.

....



Figure 5. P3 amplitude at Pz is plotted for each task. Bars are standard errors. The left column of each pair is day 1. The right is day 2.

P3 LATENCY



Figure 6. P3 latency at Pz is plotted for each task. Bars are standard errors. The left column of each pair is day 1. The right is day 2. The star indicates a significant difference between test days (p < .05).

15

REACTION TIME

- Te

33



Figure 7. Reaction time is plotted for each task. Bars are standard errors. The left column of each pair is day 1. The right is day 2.

PERCENT CORRECT



Figure 8. Percent correct is plotted for each task. Bars are standard errors. The left column of each pair is day 1. The right is day 2. Stars indicate a significant difference between days (p < .05).

the target stimulus and percent correct is plotted in Figures 3 and 4.

a she in the second

Comparisons between Test Days. Statistically significant differences between test days were observed only for P2 amplitude or latency on all tests. However, no differences were observed between test days for N1 amplitude, N1 latency, P3 amplitude, P3 latency, reaction time and percent correct. P3 amplitude and latency at the P_z for the target stimulus in all the tasks is plotted in Figures 5 and 6 respectively. Reaction time to the target stimulus and percent correct is plotted in Figures 7 and 8.

Reliability Estimates for ERP and Performance Data. The alpha reliability coefficients for the two test times on the same day are listed in Table 1. Included in the table are ERP measures for the target stimulus. The reliability was also list for the reaction time and percent correct measures. Most often the reliability of the amplitude of the ERP component was higher than that of the latency. The reliability coefficients for ERP amplitude measures ranged to .4465 to .8251. The Sternberg Memory task with set size 3 had the lowest values. P2 amplitude, likewise, had the lower values than other ERP amplitude measures. The alpha reliability coefficients for the between day comparisons for the ERP and performance measures are listed in Table 2. Lower reliability values were observed for N1 and P2 amplitude as compared to P3 amplitude. Except for the Sternberg Memory task with set size 3, the reliability coefficients for P3 amplitude ranged from .7714 to .9015 across the cognitive tasks.

Table 1 Alpha Reliability Coefficients for Measures On the Same Day (N=37)

.

. •

.

AREM	Task			
	N1 at F_z Amplitude P2 at C_z Amplitude P3 at P_z Amplitude	.7837 .6516 .7284	Latency Latency Latency	.6102 .6009 .6089
CPT-3	(Task			
	P2 at C, Amplitude P3 at P, Amplitude Percent Correct Reaction Time	.7539 .7629 .2714 .8375	Latency Latency	.5650 .6108
CPT-I	aired Letter Task			
	P2 at C_z Amplitude P3 at P_z Amplitude Percent Correct Reaction Time	.8046 .7836 .7780 .8618	Latency Latency	4088 .8755
Sternberg Memory Task				
Set	Size 3 P2 at C _z Amplitude P3 at P _z Amplitude Percent Correct Reaction Time	.4465 .5606 .7216 .8233	Latency Latency	.4978 .5053
Set	Size 6 P2 at C _z Amplitude P3 at P _z Amplitude Percent Correct Reaction Time	.6755 .8251 .8119 .9017	Latency Latency	.5909 .5405

19

Table 2 Alpha Reliability Coefficients for Measures between Days (N=20)

۰

C.

AREM	Task				
	N1 at F_z Amplitude P2 at C_z Amplitude P3 at P_z Amplitude	.5667 .5031 .8364	Latency Latency Latency	.8928 .6995 .5723	
CPT-2	(Task				
	P2 at C _z Amplitude P3 at P _z Amplitude Percent Correct Reaction Time	.6631 .9015 .6860 .8713	Latency Latency	.1627 .5914	
CPT-Paired Letter Task					
	P2 at C, Amplitude P3 at P, Amplitude Percent Correct Reaction Time	.6298 .7845 .6173 .8607	Latency Latency	.7425 .4505	
Sternberg Memory Task					
Set	Size 3 P2 at C, Amplitude P3 at P, Amplitude Percent Correct Reaction Time	.6589 .6868 .6770 .7503	Latency Latency	.3983 .8157	
Set	Size 6 P2 at C _z Amplitude P3 at P _z Amplitude Percent Correct Reaction Time	.1710 .7714 .7234 .7533	Latency Latency	0711 .4868	

20

Ì

-

and the second se

Comparison of ERPs from Normal Subjects with Subjects Who Used Drugs. ERP differences in the ERPs and performance measures were observed between the normal subjects (non-smokers and smokers) and drug abusers who denied illicit drug use. On all tasks reaction time was significantly longer for the drug abusers (CPT-X: group main effect [p<.05]; CPT-Paired Letter: group main effect [p<.001];</pre> Sternberg: group by set by condition interaction [p<.05]). **P3** amplitude differed among the three groups on the AREM (group by time by electrode interaction, p< .05) and CRT-X (group main effect, p<.01) tasks. The P3 amplitude data for the P_z electrode is plotted in Figure 9. While the P3 amplitude for the non-smokers and smokers drops slightly from the first to the second test, P3 amplitude for the drug abusers is low at the first test time and is significantly increased at the second test time. The drug abusers have the smallest P3 of the three groups in the CPT-X task.

Discussion and Conclusions

The test-retest reliability of N1, P2 and P3 components of cognitive event-related potential were determined for the auditory rare event monitoring task (Oddball task), two Continuous Performance Tasks (CPT-X: respond for an X; CPT-Paired Letter: respond to any letter which repeats) and the Sternberg Memory task with three and six letter memory sets. The Cronbach alpha coefficient was used to test within day (2 tests on the same day) and between day reliability. The alpha was .70 or higher for P3 amplitude for all tasks in both the within and



Figure 9. P3 amplitude is plotted for the 3 groups of subjects. Bars are standard errors. The star indicates a significant difference between time 1 and time 2 for the drug abusers.

between day comparisons except for the Sternberg memory task with a three letter set. The reliability of P3 amplitude in these four tasks compare favorably with previously reported findings on the Oddball task (Polich, 1986; Fabiani et al, 1987; Segalowitz and Barnes, 1993). We are the first to report reliability measures for ERP components for other three cognitive tasks. The reliability of P3 latency (.54-.87 within day, .45-.81 between days) over our four tasks, likewise, is similar to data published by the same researchers for the Oddball task.

Only Segalowitiz and Barnes (1993) tested the reliability of the N1 and P2. Our reliability values for both amplitude and latency of these components were higher than those Segalowitz and Barnes reported, but lower than we found for P3 amplitude. P2 amplitude was found to significantly differ between the nonsmokers and non-deprived smokers in our sample. P3 amplitude did not. Thus, the lower reliability values for P2 amplitude in our sample and those in Segalowitz study may be due to the increased variability resulting from smoking status.

When the low reliability values for the Sternberg memory task with a three letter set size are not considered, comparisons can made among the within and between day alpha measures. N1 and P2 amplitude measurements were more reliable when made on the same day. P3 amplitude measures are more reliable when measured across days than within a day for the Oddball and both CPT tasks. No clear patterns were observed for N1, P2 and P3 latency.

The reliability of the P3 amplitude measurements exceeded the

reliability of the percent correct values in all tasks except the Sternberg memory task with a three letter set size. The reliability of reaction time measurements most often were higher than similar values for P3 amplitude and latency. However, Donchin and Coles (1988) have suggested that P3 latency and reaction time may be independent. Thus, the comparison of P3 and reaction time may not be appropriate.

Sec. Sec.

The normative ERP data collected in this study proved useful when it was compared to ERPs collected from the group of drug abusers who denied using illicit drug during the screening process for this study. The test-retest pattern for the drug abusers significantly differed from non-smokers and cigarette smokers who did not use illicit drugs. Not only was the test-reliablity of the ERPs data high, but in using this normative ERP data we were able to indentify cognitive processing alterations in a group of drug abuser who denied illict drug use.

In conclusion, the test-retest reliability of the ERP components recorded in this battery of four different cognitive tasks is high and comparable with data collected by other researchers on only a single task. The measurement of the ERP components, in particular the P3, in this battery of cognitive tasks appears stable enough to build norms for the ERPs components for each of the tasks in the battery.

References

- Blouin, A (1985) Computerized Psychiatric Diagnosis (CDISR) Ottawa Civic Hospital: Ottawa.
- Brumaghim JT, Klorman R, Strauss, J, Levine JD, Goldstein MG (1987). What aspects of information processing are affected by methylphenidate? Findings on performance and P3b from two studies. **Psychophysiology 24**:361-373.
- Coons HW, Peloquin LJ, Klorman R, Ryan RM, Bauer LO, Perlmutter RA, Salzman, L.F. (1981). Effects of methylphenidate on young adults' event related potentials. Electroenceph clin Neurophysiol 51:373-387
- Ditraglia GM, Polich J (1991) P300 and introverted/extraverted personality types. **Psychophysiology 28**: 177-184.
- Donchin E, Coles MGH (1988) Is the P300 component a manifestation of context updating. **Behav Brain Sci** 38:387-401.
- Fabiani M, Gratton G, Karis D, Donchin E (1987) Definition, identification, and reliabilty of the measurement of the P300 component of the event-related brain potential. In PK Ackles, JR Jennings & MGH Cloes (Eds.) Advances in psychophysiology (vol2, pp1-78) London:JAI Press.
- Fitzpatrick P, Klorman R, Brumaghim JT, Keefover RW (1988). Effects of methylphenidate on stimulus evaluation and response processes: Evidence from performance and event-related potentials. Psychophysiology 25:292-307.
- Gevins A, Cutillo B (1993) Spatiotemporal dynamic of component processes in working memory. Electrocencep clin Neurophysiol 87: 128-143.
- Greenhouse SW, Geisser S 1959 On methods in the analysis of profile data. **Psychometrika 24**:95-152.
- Hanson JC, Hillyard SA (1980) Endogenous brain potentials associated with attention. EEG and Clin Neurophysiol 49:277-290.
- Herning RI, Hooker WD, Jones RT (1987). Cocaine effects on electroencephalographic cognitive event-related potentials and performance. EEG and Clin Neurophysiol 66:34-42
- Herning RI, Jones RT, Hooker WD, Tulunay FC (1987) Information processing components of the auditory event related potentials are reduced by cocaine. Psychopharmacology 87:178-185.

- Herning RI, Jones RT, Peltzman DJ (1979) Changes in human event related potentials with prolonged delta-9-tetrahydrocannabinol (THC) use. Electroencephalogr clin Neurophysiol 27:556-570.
- Jasper HH (1958) The 10-20 electrode system of the International Federation. Electroencephalogr clin Neurophysiol 10:371-375.
- Johnson R (1993) On neural generators of the P300 component of the event-related potential. **Psychophysiology 30**:90-97.
- Klorman R, Coons HW, Borgstedt AD (1989) Effects of methylphenidate on adolescents with childhood history of attention-deficit disorder: II. Clinical findings. J Am Acad Child Psychiatry 34: 231-246.
- Knott VJ (1985) Tobacco effects on cortical evoked potentials to distracting stimuli. **Neuropsychobiology 13**:74-80.
- Nataanen R, Paavilainen P, Tiitnen H, Jiang D, Alho K. (1993) Attention and mismatch negativity. **Psychophysiology 30**:436-450.
- Nunnally J (1978) **Psychometric Theory.** 2nd Edition New York: McGraw-Hill.
- Polich J (1986) Normal variation of P300 from auditory stimuli. Electroencephalogr clin Neurophysiol 65:236-240.
- Polich J (1989) Frequency, intensity, and duration as determinants of P300 from auditory stimuli. J Clin Neurophysiol 6:277-286.
- Reeves DL, Stanny RR, Wilson GF, Herning RI, Pickworth W, VanOrden KF, Caldwell J.A. (1991) The OMPAT level I Neurophysiological Performance Assessment Battery: NPPAB. Naval Aerospace Medical Research Laboratory: Monograph 43. Pensacola, Florida.
- Robin L, Helzer J, Cottler L, Golding E (1989) NIMH Diagnostic Interview Schedule, Version III Revised.
- Segalowitz SJ, Barnes KL (1993) The reliability of ERP components
 in the auditory oddball paradigm. Psychophysiology
 30:451-459.
- Sklare DA, Lynn GE (1984) Latency of the P3 event-related potential: Normative aspect and within subject variability. Electroencephalogr clin Neurophysiol 59:420-424.

Sternberg S (1975) Memory scanning: New findings and current controversies. Q J Exp Psychol 27:1-32.

- Teder W, Alho K, Reinikainen K, Naatanen R (1993) Interstimulus interval and selective-attention effect on auditory ERPs: "N1 enhancement" versus processing negativity. **Psychophysiology 30**: 71-81.
- Vasey MW and Thayer JF (1987) The continuing problem of false positive in repeated measures ANOVA in psychophysiology: A multivariate solution. Psychophysiology 24:479-486.
- Wickens C, Kramer A, Vanesse L, Donchin E (1982) The performance of concurrent tasks: a psychophysiological analysis of the reciprocity of information processing resources. Science 221:1080-1082.
- Winer BJ (1972) Statistical Principles in Experiment Design Mc Graw-Hill: New York, 1972.