AD-A277 372	TATION P	AGE distribution	- brarr	Form Acore	é J
$\Lambda \square = \Delta / / / - \Delta / /$			n uni i	OMB VC 21	4-0188
	, all standard fragmants and an all standard fragmants an all standard fragmants and an all standard fragmants an all standard	respirate including their meirorial normarich i Send comments regard dduarteis vervices Li remorate for Budger Paperwork Redum or Proje		um onri lear Hings Beniest mateloris Operations and se Bill Weshingt in	an an an ghailte an stainn an stainn an ghailte an stainn an stainn an stainn an stainn an stainn an stainn an stainn an stainn an stainn an stainn an stainn an stainn an stainn a an stainn an
	T DATE	3. REPORT TYPE AND	DATES		Final
TITLE AND SUBTITLE	January 1994	1 December 19	5. FUND	ING NUMBERS	r 1993
		and and			
lime-resolved ik spectrosco	opy, energy tra	insier, and	AFO	SR90-0055	
state-to-state collision dy	ynamics of atmo	spheric species			50
AUTHOR(S,			Tas	K # 2303/	65
David J. Nesbitt					
PERFORMING ORGANIZATION NAME(S) AN	D ADDRESS(ES)		8. PERFC	RMING ORGA	NIZATION
University of Colorado			REPO	(T NUMBER	
Boulder, CO 80309-0019			40000		00 E 7
			ACURE	9 4	0051
SPONSORING MONITORING AGENCY NAM	E(S) AND ADDRESS(ES	,	10. SPON	SORING MON	ITORING
AFOSR/NC	14-14 B	· · · · ·	• • -	~ ^ ^	0110
Building 410				94- 0	AIIN
BOLLING AFB DC	FIECTE				
20332-0440	E				
	U B				
DISTRIBUTION AVAILABILITY STATEMEN	T		12b. DIS1	RIBUTION CO	DE
Approved for public release	e:				
•• •• • • • • • •					
distribution is unlimited					
distribution is unlimited	,				
distribution is unlimited					
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFC)SR grant have	been in the follo	owing	areas 1)	We have
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFC veloped a new method for st)SR grant have tate-to-state c	been in the follo collisional energy	owing y tran	areas l) sfer in c	We have rossed
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFC veloped a new method for st lecular beams, based on high)SR grant have tate-to-state c 3h sensitivity	been in the follo collisional energy direct IR laser a	owing y tran absorp	areas l) sfer in c tion, and	We have rossed allowing
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFC veloped a new method for st lecular beams, based on hig solute cross sections to be	OSR grant have tate-to-state c gh sensitivity e obtained by m	been in the follo collisional energy direct IR laser a measurement of <u>abs</u>	owing y tran absorp solute	areas 1) sfer in c tion, and number d	We have rossed allowing ensities
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFC veloped a new method for st lecular beams, based on hig <u>solute</u> cross sections to be the scattering species (CH ve been developed to study	OSR grant have tate-to-state of sensitivity obtained by m H_2 , H_2 0, HF wit	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> th rare gases). 2)	owing y tran absorp solute) Pola	areas 1) sfer in c tion, and number d rization d modulat	We have rossed allowing ensities methods
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AF(veloped a new method for st lecular beams, based on hig <u>solute</u> cross sections to be the scattering species (CH ve been developed to study 5 kHz) of a single mode dic	DSR grant have tate-to-state of gh sensitivity e obtained by m t ₄ , H ₂ O, HF wit alignment of J ode laser, and	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> th rare gases). 2) states, based or used to study col	owing y tran absorp solute) Pola n rapi llisio	areas l) sfer in c tion, and number d rization d modulat nal align	We have rossed allowing ensities methods ion ment of
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFC veloped a new method for st lecular beams, based on high solute cross sections to be the scattering species (CH ve been developed to study 5 kHz) of a single mode dic 2 in supersonic jets. 3) High	OSR grant have tate-to-state of sensitivity e obtained by m H ₄ , H ₂ O, HF wit alignment of J ode laser, and igh resolution	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> th rare gases). 2) states, based or used to study col flash kinetic spe	owing y tran absorp solute) Pola n rapi llisio ectros	areas 1) sfer in c tion, and number d rization d modulat nal align copy has	We have rossed allowing ensities methods ion ment of been used
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on hig solute cross sections to be the scattering species (CF ve been developed to study 5 kHz) of a single mode dic 2 in supersonic jets. 3) Hz investigate reaction kinet	OSR grant have tate-to-state of sensitivity obtained by m i ₄ , H ₂ O, HF wit alignment of J ode laser, and igh resolution tics of OH radi	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> h rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocar	owing y tran absorp solute) Pola n rapi llisio ectros rbons,	areas l) sfer in c tion, and number d rization d modulat nal align copy has as well	We have rossed allowing ensities methods ion ment of been used as to
ABSTRACT (Maximum 200 words) Achievements under the AFC veloped a new method for st lecular beams, based on hig solute cross sections to be the scattering species (Ch ve been developed to study '5 kHz) of a single mode dic '2 in supersonic jets. 3) Hz investigate reaction kinet	OSR grant have tate-to-state of sensitivity obtained by m t ₄ , H ₂ O, HF wit alignment of J ode laser, and igh resolution tics of OH radi	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> h rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 3 and	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at	We have rossed allowing ensities methods ion ment of been used as to 193 nm and
ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on his solute cross sections to be the scattering species (CF ve been developed to study 5 kHz) of a single mode did in supersonic jets. 3) Hz investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening	OSR grant have tate-to-state of the sensitivity to obtained by m talignment of J ode laser, and igh resolution tics of OH radi the for photofrance of of the sense of OH	been in the follo collisional energy direct IR laser a beasurement of <u>abs</u> th rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 and to IR	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "airglow	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detectio
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on high solute cross sections to be the scattering species (CF ve been developed to study 5 kHz) of a single mode dic 2 in supersonic jets. 3) High investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have bee asuring state-resolved energy	OSR grant have tate-to-state of the sensitivity to obtained by m the obtained by m the solution of J ode laser, and ligh resolution tics of OH radi tics of OH radi tics for photofra ing rates of OH ten measured. 5)	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> h rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant A new capability "bot atom" coll	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 3 and to IR y has	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "airglow been deve	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detection loped for shell
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on hig solute cross sections to be the scattering species (CF ve been developed to study 5 kHz) of a single mode did 2 in supersonic jets. 3) Hz investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have bee asuring state-resolved ener dicals. By absolute absort	OSR grant have tate-to-state of sensitivity e obtained by m H ₄ , H ₂ O, HF wit alignment of J ode laser, and igh resolution tics of OH radi ds for photofra- ng rates of OH en measured. 5) rgy transfer in bance measureme	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> th rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant A new capability "hot atom" coll:	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 and to IR y has isions of hi	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "aifglow been deve of open gh resolu	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detectio loped for shell tion
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on his solute cross sections to be the scattering species (CF ve been developed to study 5 kHz) of a single mode did 2 in supersonic jets. 3) Hz investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have been asuring state-resolved energicals. By absolute absort ppler profiles, both <u>integr</u>	OSR grant have tate-to-state of sensitivity obtained by m 4, H ₂ O, HF wit alignment of J ode laser, and igh resolution tics of OH radi is for photofran is for photofran g rates of OH en measured. 5) rgy transfer in bance measurement cal and differe	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> h rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant A new capability whot atom" coll: ents and analysis ential cross sect	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 3 and to IR y has isions of hi ions t	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "airglow been deve of open gh resolu o a speci	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detectio loped for shell tion fic final
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on high solute cross sections to be the scattering species (CF ve been developed to study 5 kHz) of a single mode dic 2 in supersonic jets. 3) High investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have been asuring state-resolved energy dicals. By absolute absort ppler profiles, both <u>integri</u> state are obtained for Cl 4	OSR grant have tate-to-state of the sensitivity obtained by m d, H ₂ O, HF wit alignment of J ode laser, and igh resolution tics of OH radi ds for photofra- ng rates of OH en measured. 5) rgy transfer in bance measurement ral and differe	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> h rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan agmentation of HNG radical relevant A new capability "hot atom" coll: ents and analysis ential cross sect	owing y tran absorp solute) Pola n rapi llisio ectros rbons, to IR y has isions of hi ions t	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "airglow been deve of open gh resolu o a speci	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detectio loped for shell tion fic final
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for si lecular beams, based on hig solute cross sections to be the scattering species (CH ve been developed to study 5 kHz) of a single mode did 2 in supersonic jets. 3) Hy investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have been asuring state-resolved energy dicals. By absolute absort ppler profiles, both integristate are obtained for CL 4 SUBJECT TERMS	OSR grant have tate-to-state of sensitivity obtained by m i ₄ , H ₂ O, HF wit alignment of J ode laser, and igh resolution tics of OH radi is for photofra- ing rates of OH en measured. 5) rgy transfer in bance measurement ral and different	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> h rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant A new capability "hot atom" coll: ents and analysis ential cross sections.	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 and to IR y has isions of hi ions t	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "aifglow been deve of open gh resolu o a speci	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detection loped for shell tion fic final OF PAGES
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for si lecular beams, based on high solute cross sections to be the scattering species (Chi ve been developed to study '5 kHz) of a single mode dic '1 supersonic jets. 3) Hy investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have been asuring state-resolved energy dicals. By absolute absort oppler profiles, both integristate are obtained for Cl 4 SUBJECT TERMS infrared airglow; energy tr Dopplerimetry	OSR grant have tate-to-state of sensitivity e obtained by m I_4 , H_2O , HF wit alignment of J ode laser, and igh resolution tics of OH radii is for photofra- ing rates of OH en measured. 5) rgy transfer in bance measurement ral and different HCl scattering transfer; potent	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> th rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant A new capability "hot atom" coll: ents and analysis ential cross sections.	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 and to IR y has isions of hi ions t	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "äifglow been deve of open gh resolu o a speci 15. NUMBER 17 16. PRICE CO	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detectio loped for shell tion fic final OF PAGES
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on high solute cross sections to be the scattering species (CH ve been developed to study 5 kHz) of a single mode dic 1 n supersonic jets. 3) Hy investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have been asuring state-resolved energy dicals. By absolute absort ppler profiles, both integristate are obtained for CL 4 SUBJECT TERMS infrared airglow; energy tr Dopplerimetry SECURITY CLASSIF:CATION 18. SECUR	OSR grant have tate-to-state of sensitivity e obtained by m I ₄ , H ₂ O, HF wit alignment of J ode laser, and igh resolution tics of OH radi is for photofra- ing rates of OH en measured. 5) rgy transfer in bance measurement ral and different HCl scattering cansfer; potent	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> is tates, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant A new capability "hot atom" coll: ents and analysis ential cross sections of the section of	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 and to IR y has isions of hi ions t	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "aifglow been deve of open gh resolu o a speci 15. NUMBER 17 16. PRICE CO	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detectio loped for shell tion fic final OF PAGES DE
distribution is unlimited ABSTRACT (Maximum 200 words) Achievements under the AFG veloped a new method for st lecular beams, based on high solute cross sections to be the scattering species (CF ve been developed to study '5 kHz) of a single mode did '1 supersonic jets. 3) Hy investigate reaction kinet tain absolute quantum yield 8 nm. 4) Pressure broadening d characterization have bee asuring state-resolved energy dicals. By absolute absort oppler profiles, both integr state are obtained for C1 4 SUBJECT TERMS infrared airglow; energy tr Dopplerimetry SECURITY CLASSIFICATION 18 SECUR OF THM	OSR grant have tate-to-state of sensitivity e obtained by m 4, H ₂ O, HF witt alignment of J ode laser, and igh resolution tics of OH radi is for photofran is for photofran grates of OH en measured. 5) rgy transfer in bance measureme tal and differed HCl scatterin cansfer; potent	been in the follo collisional energy direct IR laser a measurement of <u>abs</u> th rare gases). 2) states, based or used to study col flash kinetic spe cal with hydrocan gmentation of HNG radical relevant A new capability "hot atom" coll: ents and analysis ential cross sections. (g. dial energy surface 19. SECURITY CLASSIFIC OF ABSTRACT	owing y tran absorp solute) Pola n rapi llisio ectros rbons, 0 and to IR y has isions of hi ions t	areas 1) sfer in c tion, and number d rization d modulat nal align copy has as well H ₂ O ₂ at "airglow been deve of open gh resolu o a speci 15. NUMBER 17 16. PRICE CO	We have rossed allowing ensities methods ion ment of been used as to 193 nm and " detectio loped for shell tion fic final OF PAGES DE

5)

· . ,

Abstract:

Achievements under the AFOSR grant have been in the following areas 1) We have developed a new method for state-to-state collisional energy transfer in crossed molecular beams, based on high sensitivity direct IR laser absorption, and allowing <u>absolute</u> cross sections to be obtained by measurement of <u>absolute</u> number densities of the scattering species (CH_4 , H_2O , HF with rare gases). 2) Polarization methods have been developed to study alignment of J states, based on rapid modulation (75 kHz) of a single mode diode laser, and used to study collisional alignment of CO_2 in supersonic jets. 3) High resolution flash kinetic spectroscopy has been used to investigate reaction kinetics of OH radical with hydrocarbons, as well as to obtain absolute quantum yields for photofragmentation of HNO₃ and H₂O₂ at 193 nm and 248 nm. 4) Pressure broadening rates of OH radical relevant to IR "airglow" detection and characterization have been measured. 5) A new capability has been developed for measuring state-resolved energy transfer in "hot atom" collisions of open shell radicals. By absolute absorbance measurements and analysis of high resolution Doppler profiles, both integral and differential cross sections to a specific final J state are obtained for Cl + HCl scattering.

Acces	sion For		
NTIS	CRA&I		
DTIC	Tab		
Unann	cunced		
Jacuification			
By	ibut long		
Aves.	lability	Qoğes	
	Aveil and	1/ur	
Dist	Special	L	
A-1			

Approved for public release; distribution unlimited.

SUMMARY OF RESEARCH ACCOMPLISHMENTS UNDER CURRENT AFOSR FUNDING (1990-1993)

The efforts over the present granting period have been in the following areas: i) to exploit the fully operational, high resolution flash kinetic spectrometer to probe transient kinetics, photofragmentation dynamics, and collisional broadening for OH radical, ii) to develop the capability for state-to-state collisional energy transfer in crossed molecular beams via high sensitivity, direct absorption of a single mode IR probe laser, iv) implement fast polarization modulation methods for detecting M, alignment in rotational states via direct IR absorption, and v) develop high resolution IR laser Dopplerimetry methods for measuring velocity resolved Cl + HCl scattering in open shell collision systems. Some highlights of the past three years are briefly summarized below.

A. State-to-state Collisional Energy Transfer in Crossed Supersonic Expansions

As a major thrust of this current granting period, we have developed a high resolution IR method of monitoring state-to-state collisional energy transfer in low density, crossed supersonic jets. The method involves i) supersonic cooling of the IR absorber into a well defined initial rotational state, ii) scattering from this state into various final states by collisions with a second supersonic jet, and iii) probing the final state distribution by fractional absorption of the F-center or diode laser light in the intersection region of the two expansions. By virtue of dual beam subtraction and near shot noise limited absorption methods, the IR detection sensitivity ($\leq 10^6 MHz$) is sufficiently high to observe collisionally excited species present at the very low densities needed to ensure single inelastic collision conditions. As result of the small IR Doppler widths and intrinsically high resolution of the IR probe laser, the method offers better than 0.01 cm⁻¹ resolution of the final scattering states. This reflects more than three orders of magnitude improvement over current time-of-flight methods that probe kinetic energy loss distributions in the scattered flux. One of the clear advantages of such an IR based technique is its general applicability to a nearly unlimited choice of collision systems. This capability is of particular importance to IR "airglow" phenomena, and opens up for detailed study many atmospherically relevant IR emitters such as CH₄, H₂O, CO₂, O₃, N₂O, etc. which are inaccessible to more sensitive LIF and MPI optical methods.

One configuration of the crossed jet apparatus is schematically shown in Fig. 1. Two pulsed valves are positioned at right angles to the tunable F-center probe laser, which is multipassed up to 36 times in a polarization preserving Herriot cell. Typical distances from the jet to intersection

٢



Fig. 1. Schematic of the current crossed jet apparatus for state-resolved energy transfer

region are 10-15 cm, with target densities on the order of 10^{11} /cm³, and collider densities between 0.5×10^{12} /cm³, respectively. High resolution scans over individual Doppler broadened transitions are performed, and the scattered state density integrated the over all final velocities. In order to ensure single inelastic collision conditions, the final state signals are monitored as a function of the stagnation pressure behind the second jet, and which demonstrate a linear dependence on collision density over the range of stagnation pressures included in the analysis.

We have performed a series of state-resolved scattering studies for CH₄ colliding with Ar, Ne and He, and have initiated parallel studies on the corresponding HF and H₂O + rare gas systems as well. The CH₄ scattering target offers a particularly novel advantage, since due to nuclear spin $(I_H = 1/2)$ statistics the molecules cool down into the lowest energy level of the three distinct symmetry species (see Fig. 2), which are not interconverted by collisions in the intersection region. Consequently, one is able to perform three scattering experiments simultaneously, out of the A $(J^*=0)$, F $(J^*=1)$ and E $(J^*=2)$ rotational symmetry manifolds of CH₄. A similar advantage is obviously offered in any system with symmetrical, non-zero spin nuclei, e.g. H₂O, NH₃, CF₄, etc.

Sample data for CH₄ scattered from $J''=0(A_1)$ into $J'=3(A_2)$, $4(A_1)$, and $6(A_2$ and $A_1)$ by collisions with He (E_{con} = 460 cm⁻¹) are shown in Fig. 3(a,b), and verify the anticipated linear





Fig. 2. Relevant nuclear spin symmetries (A,E, F) and collisional pathways for the CH₄ rotational manifold.

dependence on backing pressure. <u>Relative</u> state-to-state collision cross sections are readily obtained from the slope ratios for the various curves. However, one major advantage of a high resolution IR probe technique is that <u>absolute</u> number densities in the scattering region can be easily determined from the fractional absorbances and known IR line strengths. Specifically, by "doping" the scattering beam with trace amounts of an IR absorber, the column integrated scattering densities (integrated over the multipass region) can be reliably determined. With a dual, skimmed beam geometry to better define initial collision velocities in the intersection region, the apparatus will also be able to measure differential state-to-state cross sections as well via high resolution Dopplerimetry on the scattered species. It is worth noting that at this level of quantum state detail in CH₄, many of the simple quantum and energy gap "law" predictions are strongly violated. For example, the j=3-0 (A symmetry) cross section is almost three times as large as the j=3-1 (F symmetry) cross section, despite both a smaller energy and quantum "gap."



Fig. 3. Sample data for He + CH₄ scattering at $E_{max} = 460 \text{ cm}^{-1}$

5

STATE-TO-STATE SCATTERING:

As one additional example, inelastic scattering data for Ar + H₂O out of the lowest "para" $(J_{K_aK_c} = 0_{00})$ and "ortho" $(J_{K_aK_c} = 1_{01})$ levels are shown in Fig 4. The plot indicates the total loss of the initial quantum state (column integrated densities) as a function of stagnation pressure of the scattering rare gas. Note the factor of 3 difference in the initial populations due to the 3:1 nuclear spin statistics in H₂O.

We have purposefully chosen relatively "simple" collision systems for initial investigation, in order to facilitate the most rigorous connection between the experimental data and a theoretical description of the dynamics. In collaboration with Prof. Jeremy Hutson from the University of Durham, we therefore have extended full QM scattering codes to include both i) atom + spherical top and ii) atom + diatomic collision partners. From the T_d symmetry of CH₄, the lowest order form of the potential can be written as an radially dependent expansion in linear combinations of spherical harmonics, i.e.

$$V(R,\chi,\theta) = V_0(R) + V_3(R) \cdot T_3(\theta,\chi) + V_4(R) \cdot T_4(\theta,\chi) + ...$$
(3)



Fig. 4. Ar + H₂O scattering at $E_{com} = 500 \text{ cm}^{-1}$. The \Box and \odot represent total loss out of the l_{01} (ortho) and 0_{00} (para) levels of water, respectively.

The predicted state-to-state cross sections can then be obtained from a trial form of the potential by both approximate coupled states and full close coupling calculations. The previous analysis by Secrest of the time-of-flight scattering data was able to infer the lowest order V₃ anisotropy in the He + CH₄ potential. Full close coupling calculations on this potential are in rough qualitative agreement with the present scattering results, but indicate considerable discrepancies, particularly in the J dependent trends for a given nucelar spin symmetry and for the higher ΔJ collisions. These higher ΔJ cross sections contain detailed information on V₄ and higher order anisotropies, which we are currently using to refine the CH₄-rare gas intermolecular potentials. In particular, we find that the ratio of V₃ to V₄ terms controls dramatically influences the relative cross sections into the various fine structure levels, quantities that are directly measured in the experimental data.

B. IR Polarization Studies of Rotational M, Alignment

The high spectral resolution in the above apparatus permits cross sections for <u>J-state</u> changing collisions to be studied in considerable detail. An even finer probe of the anisotropy in the intermolecular potential can be obtained from studies of the simultaneous <u>M₁-state</u> changing collisions. In an anisotropic velocity field such as exists in a crossed beam, these processes can result in a macroscopic <u>alignment</u> of the overall final state rotation by a single inelastic collision, and can be studied via polarization dependent absorption methods. In the course of the next granting period, we will implement such polarization modulation methods in the crossed beam geometry, and thereby probe the final $|J, M_j\rangle$ state distributions via the infrared "birefringence" of the scattered molecules. In addition to providing a more state-to-state description of collision dynamics, these anisotropic scattering processes may provide a novel source of highly aligned states for further studies of <u>stereospecific</u> collision and chemical reaction phenomena.

As a necessary initial experiment, we have used high resolution IR diode lasers to investigate the collisional formation of aligned J states induced simply by relative velocity slip in a seeded supersonic expansion. The present experimental method (see Fig. 5) involves i) fast, amplitude preserving modulation of the probe IR laser polarization with an 80 kHz photoelastic ZnSe crystal, ii) polarization dependent attenuation of the probe light by molecules in the jet, followed by iii) phase sensitive detection of this modulation on the transmitted light. Our test system has been CO₂ expanded in H₂, He and Ar diluent, taking advantage of the strong ν_3 absorption band at 4.3 μ m, a single mode laser diode which conveniently accesses this band, and the relative simplicity of quantum calculations for "ball + stick" collisional scattering. Furthermore, natural abundance CO₂ has zero spin nuclei,



Fig. 5. Apparatus schematic for measuring rotational alignments

and hence dipolar couplings between \underline{J} and \underline{I} , which can cause \underline{J} to precess in the lab frame, vanish identically. Experimentally we directly measure the polarizance, i.e.

$$P = \frac{A_{\perp} - A_{\parallel}}{A_{\perp} + A_{\parallel}}$$
(4)

where A_{\perp} and A_{\parallel} are the measured absorbances for light polarized perpendicular and parallel to the expansion axis, respectively. This polarizance can be directly related to the alignment parameter via

$$A_0 = \frac{4P}{3-P} \cdot \frac{1}{h^{(2)}(J',J'')}$$
(5)

where $h^{(2)}$ is a function only of the upper/lower states of the probe transition. This alignment parameter is conventionally used to determine the second moment of an M_J distribution in P_t(cos θ). The first moment of such a distribution corresponds to rotational <u>orientation</u>, and can be obtained from analogous measurements with modulated, <u>circularly</u> polarized light. With the present modulated absorption method, we have demonstrated a sensitivity to fractional polarizance in the jet samples of $\leq 0.1\%$.

A sample set of data (J=0,2,4) for 6.4% CO₂ in He at 700 Torr is shown in Fig. 6. As expected, there is a null alignment signal for J=0, which serves as an essential diagnostic at these



Fig. 6. Typical alignment scans over CO₂ transitions (v₃ asymmetric stretch)

high sensitivity levels for stray birefringence, optical feedback to the diode laser, etc. For any J > 0, however, there are quite significant alignment signals that increase monotonically with J to a asymptotic maximum value ($A_0 \approx -0.10$) by J=8, corresponding to a roughly 3:2 enhancement of the low $|M_1|$ levels over that of an isotropic distribution. A similar asymptotic trend in J was also observed by Bergmann and coworkers in Na, expansions. This alignment is negative, which is consistent with an excess of CO₂ molecules rotating with J perpendicular to the expansion axis. By virtue of the relative velocity slip, the alignment signals are also sensitive to the mass of the expansion gas, dropping by 5 fold between He and Ar. However, despite the $\sqrt{2}$ greater relative velocity slip, we observe a 2-fold reduction in CO_2 alignment signals in H_2 vs. He expansions. This suggests a higher efficiency for M, scrambling collisions with structured vs. unstructured diluent gases, in contrast to the results of Friedrich et al. for L/He and L/H₂ expansions. Interestingly, the maximum alignment values are in good agreement with the predictions from a steady state classical model developed by Zare and coworkers, taking eccentricity parameters from the recent He + CO_2 intermolecular potential by Beneventi et al. Considering that CO, has a relatively small eccentricity, it seems quite probable that even more highly aligned molecular samples would be experimentally accessible in the IR for more "rod-like" species such as cumulated acetylenes. Furthermore, this magnitude of alignment for CO₂ develops for even modest backing pressures of He (i.e. 200-1000 Torr), and thus it is also quite probable that many supersonic jet pinhole experiments are being performed, quite unintentionally, on non-isotropic molecular samples. Interestingly, however, we have observed no measurable alignments on species to date in a slit expansion geometry.

C. Absolute OH Photolysis Quantum Yields

The production of CM via photolysis of HNO₃ and H_2O_2 has received considerable attention in recent years due to its in portance as a precursor for laboratory studies of open shell van der Waals complexes, vibrationally mediated photodissociation, inelastic energy transfer and reactive kinetics, with additional interest in HNO₃ as a stratospheric source of OH from unfiltered solar photolysis in the 190-220 nm region. Nevertheless, absolute quantum yields for these photolytic processes have been quite difficult to determine, due primarily to the difficulty in establishing a reliable probe of absolute OH radical concentrations. Consequently, most studies have been based on relative measurements, calibrated against standard reference systems such as H_2O_2 photolysis at 248 nm.

With a reliable dipole moment function for OH experimentally measured (0.70 Å $\leq R \leq$ 1.76 Å) from the prior work of Nelson et al. in our laboratories, the integrated absorption cross sections for OH (v' \leftarrow v") vibrational transitions have been determined as a function of J", ΔJ , and F=1 or 2 spin orbit states. From a high resolution scan over the spectral line profile, therefore, absolute OH concentrations in a given quantum state can be measured actinometrically in the flash kinetic spectrometer from the absorbance of the narrow band IR laser in a known path length. By repeated measurements over the range of thermalized OH quantum states, both the equilibrium nature of the distribution and the total density of $[OH]_{tot}$ can be determined. From additional measurements on the spatial profile and homogeneity of the excimer beam, one can also simply obtain the concentration of absorbed UV photons, i.e. [UV photons]_tot. The ratio of these two quantities provides an absolute measure of the quantum yield, i.e.

$$\Phi = [OH]_{uv} / [UV \text{ photons}]_{uv}$$
(6)

With this direct method, we have performed absolute quantum yield measurements for HNO_3 and H_2O_2 at 193 nm and 248 nm, and find each quantum yield to be less than 100% photon efficient (see Table I). This includes photolysis of H_2O_2 at 248 nm, for which a quantum yield of 2.0 (i.e. unit

Wavelength	Precursor	Φ	
193 nm	HNO,	0.47±0.06	
248 nm	HNO,	0.75 ± 0.10	
193 nm	H ₂ O ₂	1.22 ± 0.13	
248 nm	H_2O_2	1.58 ± 0.23	

Table I. Photolysis Quantum Yields

efficiency for breaking the O-O bond) has often been used as the reference system in many relative measurements. The HNO₃ system at 193 nm is particularly dramatic, with the production of OH actually constituting less than half $(42\pm6\%)$ of all contributing photolysis channels. A sample data plot for 193 nm is shown in Fig. 7, with the dashed line corresponding to a 100% photodissociation quantum yield shown for comparison. The presence of significant non-OH producing channels in HNO₃ photolysis has since been confirmed by Curl and coworkers, (via HONO production), and Lee and coworkers (via O atom production). The presence of O(³P), O(¹D) and H(²S) channels in HNO₃ photolysis has been recently quantified from resonance fluorescence studies by Ravishankara and coworkers.

E. Collisional Line Broadening in OH (v=1-0) Transitions

There has been a considerable interest in developing accurate intermolecular potentials for open shell systems. A particular focus has been the Ar-OH complex, for which a high level ab initio potential presently exists and the electronically excited potential surface has been extensively investigated. The corresponding ground state potential has proven more difficult to probe, although there have been some very challenging SEP and IR-UV double resonance studies by Lester and



Fig. 7. Sample data plot for determining the quantum yield for OH production from HNO₃ photolysis at 193 nm.

coworkers. Alternative, non-spectroscopic tests of this ab initio Ar + OH potential for the ground state would be quite useful, as well as for the more complex systems such as $N_2 + OH$ and $O_2 + OH$ which are of more direct relevance to airglow phenomena.

We have therefore investigated collisional broadening on the v=1-0 transition for X(II) OH radicals in the presence of He, Ar, N₂, and O₂. This flows naturally from the above photolysis quantum yield studies, since high resolution scans over individual transitions of the transient OH radical are necessary in order to obtain the integrated absorbances. The high resolution spectral data are analyzed both as i) a Voigt convolution over Gaussian (thermal) and Lorentzian (collisional broadening) contributions and ii) a Dicke narrowed line shape, as a function of buffer gas pressure, end-over-end rotational angular momentum (N), spin orbit (F=1,2), and lambda doublet (\pm) state for each of the four buffer gases.

A sample presentation of the pressure broadening results is shown in Fig. 8 for the P(2.5)1⁻ transition, which demonstrates an ordering (i.e. $k^{N_2} > k^{O_2} > k^{A_r} > k^{H_e}$) that is consistent with well depths for the corresponding N₂, O₂, Ar and He complexes with HF. The complete set of results as a function of N reveals several interesting trends. i) The pressure broadening is strongly rotational state dependent, decreasing with increasing N. This is consistent with a similar J dependence in pressure



Fig. 8. Pressure broadening results for the P(2.5)1' transition in OH(v=1-0) in the presence of N₂, O₂, Ar and Hc buffer gas

broadening rates observed in the HF + Ar and Ne systems, and would support a metastability of high rotational levels of OH to collisional relaxation. ii) With the exception of He + OH, the pressure broadening rates depend only to the end-over-end angular momentum, N, and within experimental error are insensitive to either spin orbit or lambda doublet state. It is interesting to note that this issue of lambda doublet propensities has relevance in interstellar dynamics, since some models for population inversions in the OH maser require unequal cross sections for collisional excitation into the two components. iii) In addition to the pressure broadening, we have observed and analyzed collisional IR Dicke narrowing for the first time in an open shell hydride (Ar + OH). We are presently collaborating with Dr. Sheldon Greene to predict the J-dependent line broadening and narrowing effects from open shell quantum scattering calculations, and thereby independently test the accuracy or the ab initio Ar + OH ground state potential energy surface.

E. Flash Kinetic Spectroscopy of Cl Radical Reactions

The steady state concentrations of ozone destroying Cl radicals in the stratosphere is strongly linked to the removal rates by reaction with hydrocarbons, i.e.

$$CI + RH \rightarrow HCI(v) + R$$
 (7)

where RH has been generated at the earth's surface and survives propagation upward through the atmosphere. The predominant hydrocarbon species at these altitudes is methane (1 ppm) with larger, non-methane hydrocarbons at 3-4 orders of magnitude lower concentrations (0.1-1 ppb). The rates for Cl + non-methane hydrocarbons, however, are more than 3 orders of magnitude <u>faster</u> than for Cl + CH₄, and hence these larger hydrocarbon species can also contribute to the removal of Cl radical.

We have used the IR flash kinetic spectrometer to investigate the absolute rates of these reactions via production of HCl. The Cl atoms are formed by excimer laser photolysis of CCl₄ or CFCl₃, thermalized by excess buffer gas, and the rate of hydrogen abstraction probed by monitoring the time resolved, single exponential appearance of the HCl(v=0) product under fully rotationally and vibrationally relaxed conditions. The results for the series of C₁-C₄ hydrocarbons are shown in Table II. The observed rates for Cl + CH₄ agree quantitatively with a wide range of previous studies, and

RH	kohorved	k medeling	
methane	0.0105(4)	0.0104	
ethane	4.42(26)	5.7	
propane	9.08(52)	16	
isobutane	9.66(31)	14	
n-butane	13.3(10)	21	

Table II. Cl + RH rates (10^{11} cm³/molecule/sec)

^{a)} from a recent compilation by W. B. Demore et al.

in particular supports the value used in current atmospheric models. However, the present work indicate significantly <u>slower</u> H abstraction rates for the C_2 - C_4 species, with as much as 30%-70% discrepancies from previous measurements. These measurements have been tested extensively i) over 2 orders of magnitude in buffer gas pressures, ii) over 1-2 orders of magnitude in [Cl]/[RH] ratios, iii) for a range of buffer gases to quench any vibrationally excited states, and iv) for several thermally populated J states of HCl. Due to the relevance of these reactions in current atmospheric models, these results are rather surprising and are providing a stimulus for reinvestigation of these systems via alternative techniques. The discrepancy for Cl + ethane (see Fig. 9) is of particular concern due to its use as reference system in relative reaction rate measurements.



Fig. 9. Stem-Volmer plot for H atom abstraction of ethane by Cl radicals

F. Velocity resolved scattering in open shell (Cl + HCl) collision via high resolution IR laser Dopplerimetry

As a recent thrust of the flash kinetic spectrometer, we have been investigating energy transfer in "hot atom" collisions, using the high resolution IR laser to probe both i) the final quantum states and ii) the final velocity distributions. The initial system we have studied is Cl + HCl, monitoring the HCl (v=1-0) transition on a series of R branch transitions up to J = 12. The Cl atoms are prepared by excimer laser photolysis of Cl₂ molecules at 308 nm, which generates a Cl atom velocity of ≈ 2000 m/s. If one includes the relative motion of the initial Cl₂ and HCl species, this translates into a center of mass collision energy for Cl + HCl of E_{com} ~ 3500 cm⁻¹, sufficient to surmount the 3000 cm⁻¹ activation barrier to Cl-H-Cl, H atom exchange (see Fig. 10). Although the reactive component is still relatively small at these energies, the collisions significantly sample "frustrated" reactive attempts, and thus state resolved energy transfer can probe the entrance and exit channels to reaction.

By measuring absolute absorbances, and integrating over the velocity profile, we have measured the net integral cross sections for collisional excitation into a given final state J from a room temperature distribution ($< J_{initial} > _{XOK} = 3$) in the flow cell. The results are summarized in Fig. 11, and demonstrate a monotonic decrease in cross section with increasing J. In particular, the maximum J excited is 12, far from J=18 accessible at these center of mass collision energies. The



Fig. 10. Schematic of the Cl + HCl collision system for state-resolved energy transfer on open shell potential surfaces



Fig. 11. Absolute integral cross sections (in $Å^2$) for scattering of thermal HCl into a final J level

inefficiency of this hot atom energy transfer process is not due to angular momentum constraints, but instead is a reflection of non-colinear "heavy-light-heavy" collision dynamics. In essence, optimum collision geometries for rotational energy transfer require the incident Cl atom to interact most strongly with the H end of HCl, which does not effectively slow the Cl atom in the Cl-HCl center of mass frame.

Such a heavy-light-heavy collision model would predict strongly forward scattering of the Cl atom, which we have investigated via time resolved high resolution Dopplerimetry on the recoiling HCl quantum states. In order to implement these ideas, we have developed Green's function methods for analyzing the Doppler profiles in the single collision regime, and thereby extracting absolute stateto-state differential cross sections into a given final HCl J state. The results are shown in Fig 12, and clearly demonstrate the anticipated trend toward strongly backward scattered (i.e. $\cos(\Theta) \approx -1$) HCl product. We are currently investigating trajectory calculations on a state-of-the-art Cl-H-Cl surface to compare with both the integral and differential cross section results, as well as to verify the key elements of the collision dynamics. 117+ tears that a to



Fig. 12. Absolute differential cross sections (in Å²/steradian) for scattering of thermal HCl into a final J^{*} level. Note the strong propensity for $\cos(\Theta) \approx -1$, i.e. predominantly forward scattering of the Cl atoms in the center-ofmass frame