

1

NAVAL HEALTH RESEARCH CENTER

AD-A279 547



MUSCULOSKELETAL AND MEDICAL MORBIDITY
ASSOCIATED WITH RIGOROUS PHYSICAL TRAINING

DTIC
S ELECTE D
MAY 24 1994
F

J. M. Linenger

S. Flinn

B. Thomas

C. W. Johnson

94-15502



APJ

Report No. 91-35

DTIC QU...

Approved for public release: distribution unlimited.



NAVAL HEALTH RESEARCH CENTER
P. O. BOX 85122
SAN DIEGO, CALIFORNIA 92186 - 5122



NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

Musculoskeletal and Medical Morbidity Associated with Rigorous Physical Training

Jerry M. Linenger, M.D., M.P.H., Ph.D., *Scott Flinn, M.D., *Bruce Thomas, M.D., and Chrisanna Weech Johnson, M.P.H.

*Division of Epidemiology, Department of Health Sciences and Epidemiology, Naval Health Research Center, San Diego, California 92186, and *Naval Special Warfare Center, Naval Amphibious Base, Coronado, San Diego, California, U.S.A.*

Abstract: In a prospective, epidemiological study, we describe the distribution and rates of medical conditions and musculoskeletal injuries suffered during training by U.S. Navy SEAL (Sea-Air-Land) Special Warfare trainees. The subjects were consecutive classes of candidates, healthy males, aged 18 to 31, contributing 482 trainee-months at risk. They were undergoing a physically demanding 25-week Naval Special Warfare Center indoctrination training during 1991 in Coronado, California. The main outcome measures were training-related medical conditions and musculoskeletal injuries resulting in a visit to the outpatient medical clinic. Combined medical conditions and musculoskeletal injuries occurred at a rate of 61.4 cases per 100 trainee-months at risk. Visits were nearly evenly split between medical conditions and musculoskeletal injuries. Respiratory complaints accounted for almost half of the medical conditions; overuse injuries (primarily, iliotibial band syndrome, patellofemoral syndrome, and stress fracture) accounted for 90% of the musculoskeletal injuries. We conclude that strenuous, sustained physical training results in a high incidence of medical conditions and musculoskeletal injury in trainees. Given this high morbidity, recommending limits on the amount and intensity of exercise seems prudent. Further research should examine the etiology, predisposing factors, therapy, and prevention of exercise-induced medical conditions and injuries. **Key Words:** Physical fitness—Epidemiology—Overuse injury—Morbidity—Physical training—Musculoskeletal. *Clin J Sport Med* 1993;3(4):229-34.

Documented multiple health benefits from regular physical activity are the prevention and management of coronary heart disease, diabetes mellitus, osteoporosis, obesity, and mental health problems (26). Regular physical activity has been promoted as a necessary and important component of disease prevention, health promotion, and improved quality of life (2,7,17,26). Yet, significant gaps in knowl-

edge remain concerning the short-term morbidity associated with physical activity.

Exercise-related soft-tissue injury rates and medical conditions connected to exercise have been reported in select populations, but the studies often suffer from methodological difficulties. Rate comparisons are difficult between studies, usually because only the cumulative frequency of injuries are reported over variable training periods (4,9,13,19). The diagnoses are often grouped into nonspecific categories such as knee injuries, sprains, and tendinitis, which further complicate direct comparisons. In retrospective studies using medical record review to obtain data, the population at risk is difficult to ascertain. In some such cases, rates cannot be calculated.

We investigated outpatient medical conditions and musculoskeletal injuries associated with an ex-

Received June 29, 1992. Accepted August 10, 1992.

Address correspondence and reprint requests to Chrisanna Weech Johnson, Division of Epidemiology, Department of Health Sciences and Epidemiology, Naval Health Research Center, P.O. Box 85122, San Diego, California 92186, U.S.A.

Work of U.S. Government. Not subject to copyright in the United States.

The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the U.S. Government.

tremely rigorous, sustained physical training program. The U.S. Navy SEALs (Sea-Air-Land) comprise an elite group of Naval Special Warfare personnel whose origins date back to 1943. At that time, the first group of volunteers from the Naval Construction Battalions fought during the invasion of Sicily. Since then, SEALs have played vital roles in unconventional warfare, counter-guerrilla warfare, and clandestine operations in maritime and riverine environments. Mission completion depends on the highest level of physical and mental stamina. These requirements dictate that initial training of prospective SEALs be among the most difficult and physically demanding military training programs in the world (1).

Favorable research characteristics of the SEAL trainee population include standardized training regimens, complete participation in all events, and a well-delineated population. This is in sharp contrast to individual or team sport populations where player exposure to physical stress varies depending on the degree of player participation and the specific position played (12,20,28). Conversely, the closed environment of military training allows accurate determination of population-at-risk.

We determined the rate of medical conditions and musculoskeletal injuries among U.S. Navy SEAL trainees at the Naval Special Warfare Center, Coronado, California, between April and June 1991. We prospectively examined the distribution and rates of cases presenting to the clinic by specific International Classification of Disease (ICD-9) diagnosis (25).

METHODS

Study group

All subjects were SEAL candidates training between April and June, 1991. Eligibility for selection to training included the following criteria: male, between the ages of 18 and 31, passing a screening physical fitness test, and completing a combat swimmer physical examination.

Prior to official training, all candidates engage in structured and mandatory fitness preparation for a period of 2 to 7 weeks, varying according to reporting date and degree of fitness on arrival. Formal training continues for 25 weeks and consists of three instructional phases that emphasize different strengths. Phase 1 involves 9 weeks of physical conditioning that requires pool and ocean swims of up to 2 miles, 4-mile timed runs, obstacle course, and inflatable raft seamanship. The sixth week of this phase, known as "Hell Week," consists of five and a half virtually sleepless days of nonstop training. Because of the nature of training during this week, injuries are treated by medical personnel in the field who follow the trainees for that purpose. Consequently, data from this week were inaccessible for our study.

Conditioning continues during the second and

third phases but has a different focus. Phase 2, which is 7 weeks in duration, emphasizes basic combat diving and swimming. The third and final phase of formal training involves 9 weeks of land warfare, including activities like rappelling. The final 5 weeks of this phase were spent off-site. Any diagnoses made during this period were not recorded nor were population days at risk counted for our study. During all three phases, physical training intensified progressively. In summary, individuals were at risk and under observation (for study purposes) for 22 to 27 weeks.

Data collection

We used a specially designed computer-based data collection system to record all patient encounters at the medical clinic during the study period. The software program featured user-friendly pull-down menus, preset defaults, and memory search features. A personal computer located at the clinic was linked via modem to a central VAX computer where the actual-time data entry was monitored closely by research personnel to ensure completeness.

We collected the following information on all patients who presented to the clinic for care: name, social security number, sex, date of birth, race, rate/rank, date reported to training, date of encounter, current phase of training, new or follow-up visit, and diagnosis. One of two physicians confirmed all diagnoses, which were recorded by specific ICD-9 code.

Injury codes were organized in the computer software by anatomical location—such as knee, shoulder, foot—on pull-down menus. If the specific diagnosis could not be found on the menu, a special code was entered and a comment provided in free text. Research staff would then classify the diagnosis by referring to the ICD-9 text and, where appropriate, add the new ICD-9 code and descriptive text to the menu. We displayed daily patient logs locally and at the research facility to help ensure accuracy and completeness.

We obtained counts of the population at risk used to calculate incidence rates from daily census figures provided by the Naval Special Warfare Center personnel department. The population at risk included only those trainees who were physically located at the base and who would present to the clinic if injured.

Injury was defined as any complaint or disability that developed during the study period and required a clinic visit, involved the musculoskeletal system, resulted from training, and was assigned an injury-related ICD-9 code.

Overuse injury, a subset of musculoskeletal injury, was defined as nonacute musculoskeletal injuries sustained over a period of time. Collaboration with clinical medical personnel who understand the SEAL training experience and etiology of injuries

during training consulted in classification of specific ICD-9 codes as overuse injuries.

Medical conditions were defined, similar to musculoskeletal injury, as any complaint that required a clinic visit and was assigned a medically related diagnostic code. The *International Classification of Disease, Clinical Modification*, 9th edition code book describes the criteria used to classify each specific injury or medical condition (25).

Analysis of data

We summarized demographic characteristics of the population and grouped diagnoses into one of two categories: medical conditions (noninjury) and musculoskeletal injuries. Incidence rates were expressed as the number of cases per 100 trainee-months at risk.

RESULTS

SEAL trainees were exclusively male, predominantly Caucasian and enlisted, and between the ages of 18 and 31.

During the study period, 482 trainee-months at risk and 296 cases were accumulated (Table 1). One hundred trainee-months can be interpreted as the equivalent of 100 troops training for 30 days. Combined medical conditions and musculoskeletal injuries occurred at a rate of 61.4 cases per 100 trainee-months at risk. Visits were nearly evenly split between medical conditions (31.7 cases per 100 trainee-months) and musculoskeletal injuries (29.7 cases per 100 trainee-months).

Respiratory conditions accounted for nearly half (14.7 cases per 100 trainee-months) of the medical disorder cases (Table 2). Acute bronchitis, common cold, strep throat, and sinusitis were the leading diagnoses in this category.

In the grouping of the 143 musculoskeletal injuries by anatomical location, over three-quarters involved the lower limb (Table 3). Iliotibial band syndrome, patellofemoral syndrome, and stress fracture occurred most frequently among specific injury diagnoses. Overuse injury accounted for >90% of musculoskeletal injury, with most cases occurring at the knee (Table 4).

TABLE 1. Incidence^a of musculoskeletal and medical disorders among SEAL trainees at NAVSPECWAR, Coronado, April-June 1991

| Disorder | Frequency | Incidence ^a (new cases/100 trainee-months) |
|--------------------------|-----------|---|
| Total | 296 | 61.4 |
| Medical (noninjuries) | 153 | 31.7 |
| Musculoskeletal injuries | 143 | 29.7 |

SEAL, Sea-Air-Land; NAVSPECWAR, Naval Special Warfare Center.

^a Incidence of new cases/100 trainee-months accumulated over 482 trainee-months.

TABLE 2. Incidence^a of medical disorders (exclusive of musculoskeletal injuries) among SEAL trainees at NAVSPECWAR, Coronado, April-June 1991

| Disorder | ICD-9 Code | Frequency | Incidence ^a |
|-----------------------------|---------------|-----------|------------------------|
| Medical conditions | | 153 | 31.7 |
| Respiratory | | 71 | 14.7 |
| Bronchitis | 49000 | 27 | 5.6 |
| Common cold | 46000 | 21 | 4.4 |
| Strep throat | 03400 | 8 | 1.6 |
| Sinusitis | 46100 | 6 | 1.2 |
| Pharyngitis/sore throat | 46200 | 4 | 0.8 |
| Influenza | 48710 | 2 | 0.4 |
| Tonsillitis | 46300 | 1 | 0.2 |
| Upper respiratory infection | 46590 | 1 | 0.2 |
| Pneumonia | 48600 | 1 | 0.2 |
| Gastrointestinal | | 28 | 5.8 |
| Gastroenteritis | 55890 | 21 | 4.4 |
| Diarrhea, functional | 56450 | 2 | 0.4 |
| Hemorrhoids | 45530 | 1 | 0.2 |
| Ulcer, gastric | 53190 | 1 | 0.2 |
| Gastritis | 53540 | 1 | 0.2 |
| Cholecystitis | 57500 | 1 | 0.2 |
| Abdominal pain | 78900 | 1 | 0.2 |
| Head, ear, nose, and throat | | 23 | 4.8 |
| Otitis externa | 38010 | 5 | 1.0 |
| Otitis serous | 38100 | 4 | 0.8 |
| Eustachian tube dysfunction | 38181 | 4 | 0.8 |
| Otitis media | 38200 | 3 | 0.6 |
| Eye infection | 36000 | 1 | 0.2 |
| Ear impaction | 38040 | 1 | 0.2 |
| Tympanic membrane rupture | 38420 | 1 | 0.2 |
| Conduction disorder | 42690 | 1 | 0.2 |
| Syncopal episode | 78020 | 1 | 0.2 |
| Headache | 78400 | 1 | 0.2 |
| Nosebleed | 78470 | 1 | 0.2 |
| Dermatological | | 4 | 0.8 |
| Wart | 07810 | 1 | 0.2 |
| Impetigo | 68400 | 1 | 0.2 |
| Ingrown toenail | 70300 | 1 | 0.2 |
| Folliculitis | 70480 | 1 | 0.2 |
| Urogenital | | 3 | 0.6 |
| Genital herpes | 05410 | 1 | 0.2 |
| Gonorrhea | 09800 | 1 | 0.2 |
| Pediculosis pubis | 13290 | 1 | 0.2 |
| Other | | 22 | 4.6 |
| Patient not seen | V6430 | 11 | 2.3 |
| Administrative encounters | V6890 | 4 | 0.8 |
| Consultation | V6590 | 3 | 0.6 |
| Hypothermia | 99160 | 1 | 0.2 |
| Typhoid shot | V0310 | 1 | 0.2 |
| Chest x-ray reading | V7250 | 1 | 0.2 |
| Skin test for tuberculosis | V7410 | 1 | 0.2 |

SEAL, Sea-Air-Land; NAVSPECWAR, Naval Special Warfare Center; ICD-9, International Classification of Disease.

^a Incidence expressed as number of cases per 100 trainee-months at risk.

DISCUSSION

Although increasing physical activity has many beneficial long-term health effects, it can also result in significant short-term morbidity. Musculoskeletal injury (especially overuse injury) and medical conditions (especially involving the respiratory system) occurred at very high rates. The lack of diagnostic specificity in previous studies made direct comparison of rates difficult; but it is apparent that the morbidity rates are higher in this population than in most others reported (4,5,9,13).

Although this study was not designed to assess the etiology of these clinical conditions, attributing

TABLE 3. Incidence^a of musculoskeletal disorders among SEAL trainees at NAVSPECWAR, Coronado, April-June 1991

| Disorder | ICD-9 Code | Frequency | Incidence ^a |
|-------------------------------|------------|-----------|------------------------|
| Musculoskeletal | | 143 | 29.7 |
| Lower limb | | 112 | 23.2 |
| Ankle/foot | | 34 | 6.0 |
| Sprain/strain, Achilles | 84509 | 5 | 1.0 |
| Tendinitis, ant/post tibialis | 72672 | 5 | 1.0 |
| Tendinitis/bursitis, Achilles | 72671 | 4 | 0.8 |
| Blister, noninfected | 91720 | 3 | 0.6 |
| Blister, infected | 91730 | 3 | 0.6 |
| Tendinitis/bursitis, peroneal | 72679 | 3 | 0.6 |
| Sprain/strain, foot, cuboid | 84519 | 2 | 0.4 |
| Cellulitis, foot | 68270 | 1 | 0.2 |
| Tenosynovitis | 72706 | 1 | 0.2 |
| Bunion/bunionette | 72710 | 1 | 0.2 |
| Plantar fasciitis | 72871 | 1 | 0.2 |
| Stress fracture | 73310 | 1 | 0.2 |
| Pes planus | 73400 | 1 | 0.2 |
| Sprain, ankle, unspecified | 84500 | 1 | 0.2 |
| Sprain, ankle, deltoid | 84501 | 1 | 0.2 |
| Strain/sprain, foot, other | 84510 | 1 | 0.2 |
| Lower leg | | 16 | 3.3 |
| Stress fracture | 73310 | 11 | 2.3 |
| Periostitis | 73350 | 3 | 0.6 |
| Sprain/strain, lower leg | 84490 | 1 | 0.2 |
| Abrasion/friction burn | 91900 | 1 | 0.2 |
| Knee | | 49 | 10.2 |
| Iliotibial band syndrome | 72662 | 21 | 4.4 |
| Patellofemoral syndrome | 71770 | 16 | 3.3 |
| Contusion, knee | 92411 | 5 | 1.0 |
| Sprain, lateral collateral | 84400 | 2 | 0.4 |
| Tendinitis, patella | 72664 | 1 | 0.2 |
| Tendinitis, knee | 72709 | 1 | 0.2 |
| Periostitis | 73350 | 1 | 0.2 |
| Sprain, medial collateral | 84410 | 1 | 0.2 |
| Sprain, cruciate | 84420 | 1 | 0.2 |
| Hip/thigh | | 13 | 2.7 |
| Sprain/strain, hip | 84380 | 7 | 1.4 |
| Stress fracture | 73310 | 3 | 0.6 |
| Cellulitis, hip and thigh | 68260 | 1 | 0.2 |
| Hamstring pull | 84300 | 1 | 0.2 |
| Contusion, thigh | 92400 | 1 | 0.2 |
| Upper limb | | 16 | 3.3 |
| Wrist/hand | | 1 | 0.2 |
| Strain/sprain, IP joint | 84213 | 1 | 0.2 |
| Elbow/forearm | | 7 | 1.4 |
| Tendinitis, elbow | 72639 | 5 | 1.0 |
| Epicondylitis | 72632 | 2 | 0.4 |
| Shoulder/arm | | 8 | 1.6 |
| Sprain/strain, biceps | 84080 | 3 | 0.6 |
| Shoulder instability | 71881 | 1 | 0.2 |
| Tendinitis, rotator cuff (RC) | 72610 | 1 | 0.2 |
| Subacromial bursitis | 72619 | 1 | 0.2 |
| Sprain/strain/tear, RC | 84040 | 1 | 0.2 |
| Contusion, shoulder | 92300 | 1 | 0.2 |
| Trunk | | 15 | 3.1 |
| Back/pelvis | | 9 | 1.9 |
| Low back sprain | 84690 | 7 | 1.4 |
| Strain/sprain, pelvis | 84880 | 1 | 0.2 |
| Contusion, back | 92230 | 1 | 0.2 |
| Chest | | 6 | 1.2 |
| Costochondritis | 73390 | 3 | 0.6 |
| Contusion | 92210 | 3 | 0.6 |

SEAL, Sea-Air-Land; NAVSPECWAR, Naval Special Warfare Center.

^a Incidence expressed as number of cases per 100 trainee-months at risk.

the higher rates to the extreme physical activity seems appropriate. Other studies confirm this relationship: For example, among triathletes, a group known for their intense and rigorous training habits,

overuse injury rates up to 86% of participants have been reported (16).

Overuse injury accounted for >90% of all musculoskeletal injury. The distribution of overuse injuries observed in this study was similar to that seen among civilian runners. The rank order of specific musculoskeletal injuries observed is generally consistent with patterns described by other athletic and military investigators, although iliotibial band syndrome (ITBS) occurred more frequently in our subjects (4,13,22,23). Conceivably, the familiarity of SEAL clinic providers with ITBS and other musculoskeletal injuries may have influenced diagnosis and, consequently, rates of injury. However, the rates of injury are not felt to be artificially high, and diagnostic procedures agree with accepted standards in sports medicine.

Overuse injuries are caused by repetitive microscopic trauma or overloading of musculoskeletal structures that leads to inflammation and secondary pain (20). These forces individually are insufficient to cause injury, but the cumulative effect exceeds the adaptive capacity of the body (23). For example, while running, there are ~1000 to 1200 foot strikes per mile each at 1.5 to 3 times bodyweight, putting tremendous cumulative stress on the musculoskeletal system (23). Several factors believed to contribute to overuse injury include persistent high intensity training and sudden increases in intensity (23). When an injury or re-injury cycle has been established, symptoms often do not resolve until the cycle is broken by resting the injured extremity.

Known predisposing factors for overuse injury prevalent among SEAL trainees include progressively increased running mileage, changes in running surface, running on uneven surfaces or in soft sand, and running in boots (8). Participation in training events for SEAL trainees is mandatory. Conversely, distance runners have the luxury of longer recovery periods, rest breaks, and individually varying training intensity. SEAL trainees are

TABLE 4. Incidence^a of overuse injuries by anatomical location among SEAL trainees at NAVSPECWAR, Coronado, April-June 1991

| Disorder | Frequency | Incidence ^a |
|------------------------|-----------|------------------------|
| Total overuse injuries | 129 | 26.8 |
| Knee | 44 | 9.1 |
| Ankle/foot | 33 | 6.8 |
| Lower leg | 15 | 3.1 |
| Hip/thigh | 11 | 2.3 |
| Back | 8 | 1.6 |
| Shoulder/arm | 7 | 1.4 |
| Elbow/forearm | 7 | 1.4 |
| Chest | 3 | 0.6 |
| Wrist/hand | 1 | 0.2 |

SEAL, Sea-Air-Land; NAVSPECWAR, Naval Special Warfare Center.

^a Incidence expressed as number of cases per 100 trainee-months at risk.

known to continue training in spite of injuries ("play hurt"), possibly exacerbating minor subclinical injuries (14,17). Experts believe risk factors for injury might also include: prior physical conditioning, proper warm-up and stretching, race, body composition, history of previous injury, nutrition, temporal aspects of injury in relation to training cycle, type of activity or training situation precipitating a particular injury, and anatomical factors (e.g., Q angle, pes cavus/planus, leg length discrepancies) (9,10,15,21,22,24,27).

Rates of medical conditions compared to musculoskeletal injury were particularly interesting. Respiratory conditions accounted for half of the medical conditions—and occurred at a rate more than double the 1987 U.S. incidence rate for all acute respiratory conditions (14.7 versus 6.7 cases per 100 person months) (18).

High rates of respiratory illness are known to occur with overcrowding, cigarette smoking (including passive smoking), colder monthly temperatures, progressively younger age, and among females (18). The SEAL trainees lived and trained in close quarters, smoked infrequently (not permitted during training), trained in San Diego during late spring/early summer (warm climate), averaged 20 years of age, and were male. Although this study was not designed to weigh the contribution of these combined factors, provide a nontraining comparison group, nor examine medical conditions, the rate observed seems so remarkable that it deserves further attention.

Possible explanations of the higher rate of respiratory illness include the stressful environment of training and the intense level of sustained exercise. As measured by the Daily Hassle Scale, a tool used by other researchers to examine the effects of stress on individuals, the high-stress groups have been shown to have significantly more episodes and more symptom-days than low stress groups in some studies (6). While low-intensity exercise appears to be beneficial for the immune system, high intensity and long duration exercise can result in immunosuppression and adverse effects on immune function (3). Overtraining may lead to temporary susceptibility to infection (11).

Throughout this study, the computer-based data collection system proved to be highly flexible and dependable, increasing our confidence in the reliability of the findings. We used actual-time entry of specific diagnoses, eliminating many of the classification biases inherent in studies using chart review methods to obtain data.

The intense physical training and progressive physical demands involved in SEAL training probably exceed those of almost any other group (1). We conclude that exercise-related morbidity in this group was substantial, probably representing the quantitative extreme in comparison to other exercising groups. Given the high morbidity shown to be

associated with extreme physical-conditioning activity, recommending limits on the amount and intensity of exercise seems prudent.

Application of epidemiological methods, as in this study, provide the logic and methods to quantitatively identify factors contributing to the risk of illness and injury. Presently, risk factors of musculoskeletal injury and medical conditions during intense physical activity are not well defined. These factors need to be systematically delineated for primary prevention to be possible. To better understand and minimize the negative impact of exercise-related morbidity, further research should examine the etiology, predisposing factors, therapy, and prevention of exercise-induced medical conditions and injuries.

Acknowledgment: We thank Capt C. Pollard, MC, USN, HM2 W. Higgs, HM2 K. Pundt, and Ensigns M. Olcott, T. Bednarczyk, N. Delaney, L. Savoia, L. Lamb, P. Singson, and J. Hamilton, MC, USNR, for their contributions to the execution of this study.

This work was supported by the Naval Medical Research and Development Command, Department of the Navy, under Work Unit MM33P30.002-6005. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the U.S. Government. This report is approved for public release, distribution unlimited.

REFERENCES

1. Barnes L, Strauss RH. The U.S. Navy Seal team: total commitment to fitness. *Phys Sports Med* 1986;14:176-83.
2. Bouchard C, Shephard RJ, Stephens T, et al., eds. *Exercise, fitness, and health: a consensus of current knowledge*. Champaign, Illinois: Human Kinetics Books, 1990.
3. Brouns F, ed. *Advances in nutrition and top sport*. *Med Sport Sci* Basel: Karger, 1991:85.
4. Clement DB, Tauton JE, Smart GW, McNicol KL. A survey of overuse running injuries. *Phys Sports Med* 1981;9:47-58.
5. Gordon NF, Hugo EP, Cilliers JF. The South African Defence Force Physical Training Programme. Part III. Exercise-related injuries sustained at an SADF Basic Training Centre. *S Afr Med J* 1986;69:491-4.
6. Graham NMH, Douglas RM, Ryan P. Stress and acute respiratory infection. *Am J Epidemiol* 1986;124:389-401.
7. Harris SS, Caspersen CJ, DeFries GH, Estes EH Jr. Physical activity counseling for healthy adults as a primary preventive intervention in the clinical setting: report for the U.S. Preventive Services Task Force. *JAMA* 1980;261:3590-8.
8. Hooper M. Iliotibial band friction syndrome—a common injury. *Practitioner* 1989;333:948-51.
9. Jones DC, James SL. Overuse injuries of the lower extremity: shin splints, iliotibial band friction syndrome, and exertional compartment syndromes. *Clin Sports Med* 1987;6:273-90.
10. Jones BH, Harris JM, Vinh TN, Rubin C. Exercise-induced stress fractures and stress reactions of bone: epidemiology, etiology, and classification. *Exerc Sport Sci Rev* 1989;17:379-422.
11. Keast D, Cameron K, Morton AR. Exercise and the immune response. *Sports Med* 1988;5:248-67.
12. Keller CS, Noyes FR, Buncher CR. The medical aspects of soccer injury epidemiology. *Am J Sports Med* 1987;15:230-7.

13. Kowal DM. Nature and causes of injuries in women resulting from an endurance training program. *Am J Sports Med* 1980;8:265-9.
14. Macera CA, Pate RR, Woods J, Davis BA, Jackson KL. Postrace morbidity among runners. *Am J Prev Med* 1991;7:194-8.
15. Marti B, Vader JP, Minder CE, Abelin T. On the epidemiology of running injuries. The 1984 Bern grand-prix study. *Am J Sports Med* 1988;16:285-93.
16. Massimino FA, Armstrong MA, Hiller WDB, Laird R, O'Toole ML. Training and injury patterns in triathletes. *Med Sci Sports Exerc* 1986;18:S86.
17. Matarazzo JD, Weiss SM, Herd JA, Miller NE, Weiss SM, eds. *Behavioral health: a handbook of health enhancement and disease prevention*. New York: John Wiley & Sons, 1984.
18. Powell KE, Kohl HW, Caspersen CJ, Blair SN. An epidemiological perspective on the cases of running injuries. *Phys Sports Med* 1986;14:100-14.
19. Schoenborn CA, Marano M. Current estimates from the National Health Interview Survey, United States, 1987. In: *Vital and health statistics*. Series 10, No. 166. (PHS) Washington: National Center for Health Statistics, 1988.
20. O'Toole ML, Hiller DB, Smith RA, Sisk TD. Overuse injuries in ultraendurance triathletes. *Am J Sports Med* 1989;17:514-8.
21. Pollock ML, Gettman LR, Milesis CA, Bah MD, Durstine L, Johnson RB. Effects of frequency and duration of training on attrition and incidence of injury. *Med Sports Sci* 1977;9:31-6.
22. Stacey RJ, Hungerford RL. A method to reduce work-related injuries during basic recruit training in the New Zealand Army. *Mil Med* 1984;149:318-20.
23. Tauton JE, McKenzie DC, Clement DB. The role of biomechanics in the epidemiology of injuries. *Sports Med* 1988;6:107-20.
24. Tomlinson JP, Lednar WM, Jackson JD. Risk of injury in soldiers. *Mil Med* 1987;152:60-4.
25. U.S. Department of Health and Human Services, Public Health Service, Health Care Financing Administration. *International classification of disease, clinical modification*. 9th ed. Washington: DHHS, 1989.
26. U.S. Department of Health and Human Services, Public Health Service. *Healthy people 2000: national health promotion and disease prevention objectives*. Washington: DHHS, 1990.
27. Volpin G, Petronius G, Hoerer D, Stein H. Lower limb pain and disability following strenuous activity. *Mil Med* 1989;154:294-7.
28. Watson MD, DiMartino PP. Incidence of injuries in high school track and field athletics and its relation to performance ability. *Am J Sports Med* 1987;50:251-4.

| | |
|----------------------|-------------------------------------|
| Accession For | |
| NTIS CRA&I | <input checked="" type="checkbox"/> |
| ERIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification _____ | |
| By _____ | |
| Distribution / _____ | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A-1 | 20 |

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

| | | | | |
|--|--|---|---|--|
| 1. AGENCY USE ONLY (Leave blank) | | 2. REPORT DATE December 1991 | 3. REPORT TYPE AND DATE COVERED Final dates: APR - JUN 91 | |
| 4. TITLE AND SUBTITLE MUSCULOSKELETAL AND MEDICAL MORBIDITY ASSOCIATED WITH RIGOROUS PHYSICAL TRAINING | | | 5. FUNDING NUMBERS Program Element: 62233N Work Unit Number: MM33P30.002-6005 | |
| 6. AUTHOR(S) Jerry Linenger, Scott Flinn, Bruce Thomas, and Chrisanna Weech Johnson | | | 8. PERFORMING ORGANIZATION Report No. 91-35 | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center P. O. Box 85122 San Diego, CA 92186-5122 | | | 10. SPONSORING/MONITORING AGENCY REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Medical Research and Development Command National Naval Medical Center Building 1, Tower 2 Bethesda, MD 20889-5044 | | | 11. SUPPLEMENTARY NOTES Published in: <u>Clinical Journal of Sports Medicine</u> , 1993, 3(4), 229-234 | |
| 12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited. | | 12b. DISTRIBUTION CODE | | |
| 13. ABSTRACT (Maximum 200 words) The distribution and rates of medical conditions and musculoskeletal injuries suffered during training by U.S. Navy SEAL Special Forces trainees were described in a prospective, descriptive epidemiological study. We studied consecutive classes of candidates undergoing the physically-demanding 25-week Navy Special Warfare indoctrination training during Spring 1991 in Coronado, CA. Healthy males, ages 19 to 31, contributed 482 trainee-months at risk. Combined medical conditions and musculoskeletal injuries occurred at a rate of 61.4 cases per 100 trainee-months at risk. Visits were nearly evenly split between medical conditions (31.7 cases/100 trainee-months) and musculoskeletal injuries (29.7 cases/100 trainee-months). Respiratory complaints accounted for almost half (14.7 cases/100 trainee-months) of the medical conditions, while overuse injuries (primarily iliotibial band syndrome, patellofemoral syndrome, and stress fracture) accounted for 90% of the musculoskeletal injuries. | | | | |
| 14. SUBJECT TERMS physical fitness, injury, epidemiology, military, U.S. Navy, SEALS, overuse injury, physical training, respiratory | | | 15. NUMBER OF PAGES 6 | |
| | | | 16. PRICE CODE | |
| 17. SECURITY CLASSIFICATION OF REPORT Unclassified | 18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified | 19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified | 20. LIMITATION OF ABSTRACT Unlimited | |