

AD-A280 115



TECHNICAL REPORT
NATICK/TR-94/019

AD _____

EFFECT OF HYPERBARIC CARBON DIOXIDE
ON SPORES AND VEGETATIVE CELLS
OF *BACILLUS STEAROTHERMOPHILUS*

DTIC
ELECTE
JUN 13 1994
S F D

By
Chester T. Roskey*
Anthony Sikes

*Framingham State College
Framingham, MA 01701

94-18004



May 1994

FINAL REPORT
March - September 1992

Approved for Public Release; Distribution Unlimited

UNITED STATES ARMY NATICK
RESEARCH, DEVELOPMENT AND ENGINEERING CENTER
NATICK, MASSACHUSETTS 01760-5000

SUSTAINABILITY DIRECTORATE

94 6 10 112

DTIC QUALITY INSPECTED 1

DISCLAIMERS

The findings contained in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

DESTRUCTION NOTICE

For Classified Documents:

Follow the procedures in DoD 5200.22-M, Industrial Security Manual, Section II-19 or DoD 5200.1-R, Information Security Program Regulation, Chapter IX.

For Unclassified/Limited Distribution Documents:

Destroy by any method that prevents disclosure of contents or reconstruction of the document.

REPORT DOCUMENTATION PAGE			Form Approved OMB No 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE May 1994		3. REPORT TYPE AND DATES COVERED Final 30 Mar - 30 Sep 92
4. TITLE AND SUBTITLE Effect of Hyperbaric Carbon Dioxide on Spores and Vegetative Cells of <u>Bacillus Stearothermophilus</u>			5. FUNDING NUMBERS PR: TB040 PE: 040 WU: AH52 TA: AG: FTB1298	
6. AUTHOR(S) Dr. Chester T. Roskey* & Dr. Anthony Sikes				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Natick RD&E Center ATTN: SATNC-WRA Kansas Street Natick, MA 01760-5018			8. PERFORMING ORGANIZATION REPORT NUMBER NATICK/TR-94/019	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Army Research Office 200 Park Drive Research Triangle Park, NC 27709			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES *Framingham State College 100 State Street Framingham, MA 01701				
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>Spore suspensions in water, 0.05M pH 7 phosphate buffer, 0.05M pH 4.5 and 0.05M pH 4.0 acetate buffer, and antibiotic assay media supplement broth (AAMS) were subjected to 50 psi to 1100 psi carbon dioxide at various temperatures in Parr bombs for up to 96 hours. Spores were resistant to high pressure carbon dioxide treatment at each pH and at both high and low temperatures. Spore suspensions that were first heat activated and then incubated at 55°C for a short time increased sensitivity to destruction by CO₂. Although spores were resistant to destruction by hyperbaric CO₂ low pressure (50 psi) CO₂ was seen to completely inhibit spore germination and outgrowth. Inhibition to germination was found to be reversible. Spore germination was found to be impaired in carbonated AAMS broth. However, the addition of 50 mM NaHCO₃ to AAMS enhanced germination and outgrowth.</p> <p>Vegetative cells were shown to be sensitive to CO₂ treatment. Complete destruction of 10⁶ cells in AAMS broth was obtained by exposure to 400 psi CO₂ for 1.5 hours.</p>				
14. SUBJECT TERMS BACILLUS STEAROTHERMOPHILUS THERMOPHILIC BACTERIA THERMOPHILIC SPOILAGE		VEGETATIVE CELLS CARBON DIOXIDE SPORES SHELF LIFE		15. NUMBER OF PAGES 39
		PERISHABLE COMMODITIES		16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT SAR	

TABLE OF CONTENTS

LIST OF FIGURES.....	v
LIST OF TABLES.....	vii
PREFACE.....	ix
INTRODUCTION.....	1
METHODS AND MATERIALS.....	3
RESULTS AND DISCUSSION.....	6
CONCLUSIONS.....	26
REFERENCES.....	29

Accession For	
NTIS CRASH	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

LIST OF FIGURES

Figure	Page
1. Effect of exposure to 200, 300, and 400 psi CO ₂ for one hour at 25°C on survival of vegetative cells of <u>Bacillus stearothermophilus</u>	7
2. Effect of exposure to 200 psi CO ₂ for 3.5 hours at 25°C on survival of vegetative cells of <u>Bacillus stearothermophilus</u>	8
3. Effect of exposure to 400 psi CO ₂ or N ₂ at 25°C on survival of vegetative cells of <u>Bacillus stearothermophilus</u>	9
4. Effect of exposure to 875 psi CO ₂ and N ₂ at 25°C for 1.25 hours on survival of vegetative cells of <u>Bacillus stearothermophilus</u>	10
5. Effect of exposure to 850-875 psi CO ₂ at 90°C on viability of <u>Bacillus stearothermophilus</u> spores suspended in 0.05M acetate buffer pH 4.5.	15
6. Effect of carbonation on germination and outgrowth of <u>Bacillus stearothermophilus</u> spores in AAMS broth during incubation at 55°C.	21
7. Effect of 54 mM NaHCO ₃ on germination of <u>Bacillus stearothermophilus</u> spores in AAMS during incubation at 55°C.	23
8. Ultraviolet absorption spectra of culture supernates obtained from:	
(A) BS vegetative cells in phosphate buffer, pH 7.0 for 17 hours at 25°C under ambient air.	
(B) BS vegetative cells in phosphate buffer, pH 7.0 for 17 hours at 25°C under 850 psi CO ₂	25

LIST OF TABLES

Table		Page
1.	The effect of hyperbaric carbon dioxide on viability of <u>Bacillus stearothermophilus</u> spores.	13
2.	The effect of heat activation on survival of <u>Bacillus stearothermophilus</u> spores exposed to 800-875 psi CO ₂ for 21 hours at 25°C.	16
3.	The effect of DMSO activation on survival of <u>Bacillus stearothermophilus</u> spores exposed to 900 psi CO ₂ for 20 hours at 25°C.	18
4.	The effect of heat activation and incubation for three hours at 55°C on survival of <u>Bacillus stearothermophilus</u> spores exposed to 900 psi CO ₂ for 67 hours at 25°C.	19
5.	The effect of 1000 psi CO ₂ on activated and non activated spores of <u>Bacillus stearothermophilus</u> suspended in AAMS broth and stored at 55°C for 18 hours.	20

PREFACE

This study was conducted during the months of May through August 1992, by Dr. Chester Roskey , Framingham State College, Framingham, MA, under the general supervision of Dr. Anthony Sikes (project officer), U. S. Army Natick Research, Development and Engineering Center, Natick, MA. The research was sponsored by the U.S. Army Research Office of Scientific Services Program administered by Battelle (Delivery Order 259, Contract No. DAAL03-91-C-0034). The work was funded under the project titled "Hyperbaric Preservation," DA project AH5240D00.

Dr. Roskey's research was designed to ascertain the following: (1) determine the effects of pressurized CO₂ on the microbial activity of the thermophilic spoilage bacterium, Bacillus stearothermophilus, (2) evaluate the effects of gas mixtures, e.g., CO₂:O₂:N₂, at different partial pressures on both vegetative and sporeforming bacteria in model food systems and ration items, (3) select optimum atmosphere/pressures as a result of time-temperature studies and (4) apply optimum condition to test potential of shelf life extension of perishable foods.

Effect of Hyperbaric Carbon Dioxide on Spores and Vegetative Cells of Bacillus stearothermophilus.

Introduction

Carbon dioxide inhibits the growth of many microorganisms. This fact has been exploited in preventing bacterial food spoilage through modified atmosphere packaging (MAP). The ability of high concentrations of CO₂ to retard the growth of spoilage flora in meat, poultry, and fish is well documented (1, 3, 17, 18, 20, 21).

The inhibitory effect of CO₂ increases when it is applied under pressure (3, 8, 10, 11, 15,). Enfors and Molin reported that germination of C. sporogenes was inhibited slightly at 4 atmospheres and almost completely at 10 atmospheres, whereas germination of C. perfringens was slightly stimulated at 4 atmospheres, unaffected at 10 atmospheres and stopped at 25 atmospheres of pure CO₂ (10). Doyle revealed that at atmospheric pressure 100% CO₂ delayed toxin production by C. botulinum. Increasing the pressure to 8.8 atmospheres caused a further delay in toxin production. Pressurized CO₂ was also lethal to C. botulinum with the viability decreasing two log cycles during eight weeks of storage under 8.8 atmospheres. Loss of viability was directly related to pressure of CO₂ and length of exposure (8). Haas et al, investigated the effect of hyperbaric CO₂ on food microbiota and found that the total count of microorganisms in foods can be reduced by treatment with CO₂ under pressure. These workers determined that 900 psi CO₂ had the greatest effect and that inhibition decreased with decreasing water activity. They also reported that low pH acted synergistically with CO₂ and that microbicidal activity was enhanced at higher temperature (15). However Blickstad et al, had reported earlier that the inhibitory effect of CO₂ was greater at lower temperatures (3).

The capacity of the almost inert spores to spoil food depends on their ability to germinate, outgrow and achieve extensive vegetative multiplication in the food. Interruption of this chain

of events at any point will prevent spoilage. Bacillus stearothermophilus (BS) is a ubiquitous sporeformer. As a thermophilic sporeformer it is of particular concern as a potential spoilage microbe in thermally-processed foods. The bacterium can grow under both aerobic and anaerobic conditions, and ferments sugars (2). Spores of BS may be present in "commercially sterile" foods and may be responsible for "flat-sour" spoilage of low acid foods (14).

The objectives of the present study were to investigate the effect of hyperbaric CO₂ on survival and growth of vegetative cells of BS as well as its effect on viability, germination, and outgrowth of endospores of BS.

METHODS AND MATERIALS

Test bacteria

Bacillus stearothermophilus ATCC 12980 (Type strain).

Vegetative cells were maintained on Antibiotic Assay Media Supplement (AAMS) agar slants at 4°C. Vegetative cells were cultivated in AAMS broth at 55°C. Spore suspensions of BS were provided by A. Sikes of the U.S. Army Natick Research, development and Engineering Center. Spore suspensions were prepared as described by Feeherry et al (13).

Hyperbaric treatment

Cells or spores were suspended in either sterile distilled deionized water, 0.05M phosphate buffer pH 7.0, 0.05M acetate buffer pH 4.5, 0.05M acetate buffer pH 4.0, or AAMS broth, in 16 mm by 150 mm screw-capped culture tubes. The caps were loosely secured to permit free gas exchange. The tubes were placed in the cylinder of 1650 ml Parr cell disruption bombs (Parr Instrument Company, 211 Fifty-Third Street, Moline, Illinois 61265 USA). After loading, the cylinders were sparged with either nitrogen or carbon dioxide at 200 psi for 1 minute to remove air. The cylinders were then pressurized (50 psi-1000 psi) and then stored at various temperatures (3°, 25°, 55°, 65°, 75°, 90°, and 100°C) for 1 to 96 hours.

Activation of spores

Spores were activated by subjecting suspensions in water or AAMS broth to flowing steam for 15 minutes. Some spore suspensions were activated by suspension in 9% or 90% v/v aqueous dimethylsulphoxide (DMSO).

Preparation of germinated spores

Heat-activated spores suspended in AAMS broth were incubated for 3 hours at 55°C.

Carbonation

Sixty ml of ice-chilled AAMS broth were aseptically transferred to an iced sterile 250 ml cylinder. A sterile 10 ml pipette was fitted to the regulator hose of a CO₂ tank. The pipette was placed into the broth and CO₂ bubbled through the media at 50 psi for 4 minutes. The pH of the carbonated AAMS broth was 5.44.

Effect of sodium bicarbonate

A 5% NaHCO₃ solution was prepared and filter sterilized. One ml of the filter-sterilized solution was added to each of three tubes containing 10 ml of AAMS broth.

One ml of sterile water was added to each of three tubes containing 10 ml of AAMS as a control. Each of the six tubes was inoculated with 0.1 ml of a BS spore suspension. All tubes were then incubated at 55°C for 24 hours. The pH of the NaHCO₃ supplemented broth was 7.6.

UV absorption spectra

One hundred ml of AAMS broth was inoculated with 1.0 ml of an overnight culture of BS in AAMS and then incubated for 7 hours at 55°C. Cells were harvested by centrifugation at 1000 X g for 20 minutes. Harvested cells were washed three times in phosphate buffer and then resuspended in buffer.

Cell suspensions were then subjected to 850 psi CO₂ for 17 hours at 25°C. Cell suspensions similarly prepared were stored under ambient conditions for the same time as controls. After incubation the CO₂ treated and control cultures were centrifuged at 1000 X g for 30 minutes and the UV absorption spectrum of the supernates was determined. UV absorption analysis was provided by K. Whitburn (U.S. Army Natick Research, Development and Engineering Center).

Enumeration

Cells or spores were enumerated on AAMS agar spread plates. Dilutions were made in either water or 0.05 M phosphate buffer pH 7.0. All dilutions were made with a Gilson pipetman P200 or P1000 fitted with appropriate tips. All counts reported represent the mean of duplicate or triplicate determinations.

RESULTS AND DISCUSSION

Effect on vegetative cells

Hyperbaric CO₂ was shown to have a biocidal effect on vegetative cells of BS. Destruction of vegetative cells was shown to be dependent on both time of exposure, as well as CO₂ pressure. The effect of a 1 hour exposure to CO₂ at 0, 300, 400 and 500 psi at 25°C is shown in Figure 1. Carbon dioxide treatment for 1 hour at 400 and 500 psi resulted in more than a two log cycle reduction in viability of B. stearothermophilus cells (from log 6.7 to log 4.2). Control cells (log 6.7) at ambient conditions did not decrease in viability in this interval. There was no apparent difference in lethality due to increasing CO₂ pressure during the 1 hour treatment period, when the CO₂ pressure was 300 psi.

Extending the exposure period, at a lower pressure (200 psi), beyond 1 hour resulted in a further decline in viability of vegetative cells. The effect of exposure time on survival of vegetative cells exposed to 200 psi over a 3.5 hour period was greater than 400 psi for only 1 hour as depicted in Figure 2. The data indicates that increasing exposure time can result in increasing lethality at lower pressures. The population of vegetative cells decreased over four log cycles (log 6.3 to log 2.0) due to 200 psi CO₂ treatment for 3.5 hours. The data presented in Figure 3 indicated that complete destruction of vegetative cells was achieved through hyperbaric CO₂ treatment at 400 psi by increasing the time from 1 h (Fig. 1) to 1.5 h. Nitrogen gas at the same pressure had no adverse effect on viability of BS vegetative cells.




Figure 1. Effect of exposure to 300, 400, and 500 psi CO₂ for one hour at 25°C on survival of vegetative cells of *Bacillus stearothermophilus*.




Figure 2. Effect of exposure to 200 psi CO₂ for 3.5 hours at 25°C on survival of vegetative cells of Bacillus stearothermophilus.




Figure 3. Effect of exposure to 400 psi CO₂ or N₂ at 25°C on survival of vegetative cells of Bacillus stearothermophilus.




Figure 4. Effect of exposure to 875 psi CO₂ and N₂ at 25°C for 1.25 hours on survival of vegetative cells of *Bacillus stearothermophilus*.

Since complete destruction of B. stearothermophilus vegetative cells culture was attained by exposure to 400 psi CO₂ for 1.5 hours, one would expect that higher pressures could achieve the same effect in less time. However, when vegetative cells were subjected to 875 psi CO₂ for 1.25 hours at room temperature (Figure 4) there was only a four log cycle reduction in vegetative cells (log 6.45 to log 2.63). Viable cells were still recovered. Cells subjected to N₂ at 875 psi and cells kept under ambient conditions for the same period were unaffected.

These results are in agreement with those of other workers (1, 3, 7, 15,17) who reported that carbon dioxide under pressure was lethal to vegetative cells of food spoilage microorganisms, and rate of lethality was dependent on the amount of pressurized CO₂ and the length of exposure. Haas et al (15) found that all microorganisms on chives, thyme, parsley, mint, and spoiled apple juice were killed by a 30-minute exposure to 800 psi CO₂ at elevated temperature (45°C). In the present study, B. stearothermophilus vegetative cells were not completely destroyed by a 75 minute exposure to 875 psi at a lower temperature (25°C). Survival of the B. stearothermophilus treated cells in this study may be attributed to the lower temperature treatment. Haas et al. (15) reported that higher temperatures enhanced the bactericidal effect of CO₂. Enfors and Molin (12), on the other hand, showed that there was greater CO₂ inhibition of the growth rate of Pseudomonas fragi and Bacillus cereus at lower temperatures.

Effect on endospores

Spores of B. stearothermophilus proved to be remarkably resistant to hyperbaric CO₂ treatment. The effect of CO₂ and N₂ on the survival of BS spores under varying conditions of temperature and pH is shown in Table 1.

When B. stearothermophilus spores at a population density of ca. log 4.7 were suspended in distilled water, 0.05M acetate buffer pH 4.0, 0.05M acetate buffer pH 4.5, or 0.05M phosphate buffer pH 7.0, and then subjected to hyperbaric CO₂ for 1 to 96 hours, there was no apparent spore destruction. Aliquots of the same spore suspensions subjected to ambient gas conditions and those subjected to hyperbaric N₂ were similarly unaffected.

There is agreement among workers regarding the effect of temperature on the bactericidal activity of CO₂. Blickstad et al, (3) found that CO₂ inhibited the microbial flora of pork to a greater extent at 4°C than at 14°C. Daniels et al (6) attributed increased inhibition by CO₂ at lower temperature to increased solubility of the gas in the water phase. However, higher temperatures enhanced the microbicidal action of CO₂ (15). The results obtained during the present investigation revealed that neither high (95°-100°C) nor low temperature (3°C) had any effect on the sporicidal activity of hyperbaric CO₂.

Blocher and Busta (4) stated that the heat resistance (D-value) of spores decreased as the pH of the medium decreased. Haas et al (15) reported that lower pH acted synergistically with CO₂ pressure. These workers found that 800 psi CO₂ at 70°C killed spores of Clostridium sporogenes suspended in thioglycolate broth when the pH of the broth was between 2.5 and 3.0, but did not kill spores when the pH of the broth was 4.0.

Table 1. The effect of hyperbaric carbon dioxide on viability of Bacillus stearothermophilus spores.

pH	Temp. °C	Pressure psi	Time hrs	Survival Fraction		
				$\log N_{CO_2}^b/N_0^a$	$\log N_{Ac}/N_0^b$	$\log N_{N_2}^d/N_0$
4	3	800-550	68	0.98	0.98	_____
4	25	840-890	68	0.94	0.93	_____
4	55	840-1050	68	0.99	0.87	_____
4	65	800-900	45	0.98	0.86	0.82
4.5	3	820-540	96	1.04	1.06	1.06
4.5	25	830-850	96	1.01	0.98	_____
4.5	65	840-900	23	0.99	0.82	0.89
4.5	90	850-875	22	0.55	0.78	_____
7	25	800-930	23	1.01	0.96	1.01
7	55	940-1050	46	1.01	0.80	0.81
7	65	940-975	20	1.00	0.81	_____
7	75	875-1000	26	1.06	0.91	_____
7	100	900-1100	1	1.09	1.08	_____

^aN₀ = initial population (mean initial population was log 4.7)

^bN_{CO₂} = number present after CO₂ treatment.

^cN_A = number present after exposure to ambient gas atmosphere for the treatment period.

^dN_{N₂} = number present after exposure to N₂ gas at the same pressure as CO₂ for the same time.

Examination of the data presented in Table 1 reveals that spores were able to tolerate exposure to hyperbaric CO₂ for 1-96 h when suspended in buffer at pH 4.0, pH 4.5, and pH 7.0. However, high temperature (90°C) and low pH (4.5) were found to act synergistically with hyperbaric CO₂ to reduce viable spores. Figure 5 shows that 850-875 psi CO₂ treatment over a 22 hour period resulted in nearly a two log cycle reduction in viability of spores (log 4.34 to log 2.40). The viability of spores held at ambient gas conditions at pH 4.5 and 90°C was also diminished by the treatment (log 4.34 to log 3.40) but to a lesser extent. Additional studies should be done to more clearly characterize the synergistic effects of high temperature, low pH, and hyperbaric CO₂.

Since unactivated spore suspensions proved to be resistant to destruction by hyperbaric CO₂ treatment, attempts were made to evaluate the effect of hyperbaric CO₂ on spores that had undergone "activation." Accordingly, spore suspensions in 0.05M phosphate buffer, pH 7.0 were activated by subjecting the suspensions to flowing steam for 5, 10, 15, 20, and 25 minutes. Viable counts were made immediately before and after steam activation and after exposure to 800-875 psi CO₂ at 25°C for 21 hours. The effect of heat activation on survival of spores exposed to hyperbaric CO₂ is shown in Table 2. Examination of the data in Table 2 reveals that hyperbaric CO₂ treatment was not effective in reducing the viability of heat activated B. stearothermophilus spores.




Figure 5. Effect of exposure to 850-875 psi CO₂ at 90°C on viability of Bacillus stearothermophilus spores suspended in 0.05M acetate buffer pH 4.5.

Table 2. The effect of heat activation on survival of Bacillus stearothermophilus spores exposed to 800-875 psi CO₂ for 21 hours at 25°C.

Time in steam (minutes)	CFU/ml ^a after steam treatment	CFU/ml after CO ₂ treatment	% Reduction
0	46.8 x 10 ³	32 x 10 ³	0
5	7.1 x 10 ³	30 x 10 ³	0
10	13.6 x 10 ³	20 x 10 ³	0
15	24.3 x 10 ³	25.5 x 10 ³	0
20	23.8 x 10 ³	22.5 x 10 ³	5.5
25	26.4 x 10 ³	20.5 x 10 ³	22.0

^a CFU = colony forming units

The maximum reduction was only 22.0 % of spores activated for 25 min. However, activation treatment of B. stearothermophilus for 60 minutes at pH 2.0 and 60°C showed a marked decrease in their heat resistance (9). Additional studies to evaluate the effect of hyperbaric CO₂ on heat activated BS spores at low pH may prove fruitful.

Widdowson (22) reported that reduction of the S-S linkages in spore protein may lead to rapid activation. It was found that pretreatment of a spore suspension with 90% v/v aqueous solution of dimethylformamide produced rapid activation of B. pantothenicus spores even at 4°C.

Widdowson (22) also reported that dimethylsulphoxide (DMSO) was effective in producing rapid activation at room temperature. She suggested that the decrease in the amount of H-bonding in the spore brought about by these compounds might be expected to result in an increase of sensitivity to damaging agents. With this in mind, an experiment was designed to assess the effect of hyperbaric CO₂ on DMSO activated spores. Spore suspensions were made in water, 9% v/v aqueous DMSO, and in 90% aqueous DMSO. Viable spore populations were determined initially and after 20 hours at 25°C under both ambient and hyperbaric (900 psi) conditions. The effect of CO₂ on DMSO activated spores is shown in Table 3. The data presented reveals that spores activated by suspension in DMSO were not more susceptible to destruction by exposure to hyperbaric CO₂ than were spores suspended in water.

There was no apparent difference between the survival of B. stearothermophilus spores treated by exposure to 900 psi for 20 hours at 25°C and controls that were exposed to ambient air under the same conditions.

Since spore suspensions as well as "activated" spore suspensions proved to be resistant to hyperbaric CO₂ treatment, activated spore suspensions were incubated (in AAMS and water) at 55°C for 3 hours in order to allow time for germination prior to being subjected to hyperbaric CO₂.

The effect of heat activation and incubation for three hours at 55°C on the survival of heat activated and germinated B. stearothermophilus spores to 900 psi CO₂ for 67 hours at 25°C is shown in Table 4.

Table 3. The effect of DMSO activation on survival of Bacillus stearothermophilus spores exposed to 900 psi CO₂ for 20 hours at 25°C.

Spores suspended in	Initial count CFU/ml	% Survival after 20 hours at 25°C CFU/ml	
		Ambient	CO ₂
H ₂ O	100 x 10 ³	28	19
9% DMSO	75 x 10 ³	44	33
90% DMSO	54 x 10 ³	76	91

Germination and growth of spores resulted in a 100 fold increase in viable counts on AAMS. Carbon dioxide treatment of the activated and germinated spore suspension resulted in a two log cycle reduction in the population. Heat activated and germinated spores that were subjected to ambient gas conditions for the same period did not exhibit a similar decrease in viability. Spores suspended in water, heat activated, and incubated, responded as did spores suspended in AAMS broth. Although there was no evidence of germination and outgrowth of the spores in the water suspension, the spores nevertheless were susceptible to destruction by hyperbaric CO₂.

Table 4. The effect of heat activation and incubation for three hours at 55°C on survival of *Bacillus stearothermophilus* spores exposed to 900 psi CO₂ for 67 hours at 25°C.

Spore Suspension in	Initial count CFU/ml	Count after 3 hours at 55°C CFU/ml	Count after 67 hours at 25°C CFU/ml	
			Ambient	CO ₂
AAMS ^a	40x10 ³	61x10 ⁵	28x10 ⁵	73x10 ³
H ₂ O	48x10 ³	18x10 ³	11x10 ³	71x10 ¹

^a AAMS = antibiotic assay medium supplement

Effect of hyperbaric CO₂ on germination and outgrowth

The experimental work reported has shown that spores are virtually indestructible to exposure to CO₂ under the conditions tested. An experiment was undertaken to determine the effect of CO₂ on germination and outgrowth of BS spores in AAMS broth. The results of this study are shown in Table 5. Three replicates of each sample were prepared. Initially, each of the 12 tubes contained ca. 29 x 10³ CFU/ml of BS spores. After 18 hours at 55°C, all tubes stored under ambient gas conditions were turbid. All tubes that were incubated under 1000 psi CO₂ were clear. Carbon dioxide at 1000 psi had prevented the germination and outgrowth of BS spores. When the spore suspensions that had received CO₂ treatment underwent further incubation under ambient conditions at 55°C, all these tubes became turbid.

Table 5. The effect of 1000 psi CO₂ on activated and non activated spores of Bacillus stearothermophilus suspended in AAMS broth and stored at 55°C for 18 hours.

Spore suspension in AAMS broth	Appearance of broth after 18 hours at 55°C	
	Control Ambient	Treated CO ₂ 1000 psi
Heat Activated	Turbid	Clear
Not Activated	Turbid	Clear

Although CO₂ had prevented germination and outgrowth of spores, it did not kill them or impair the spores ability to undergo further development under more favorable ambient conditions.

Additional tests were performed to ascertain the lowest CO₂ pressure that would prevent germination and outgrowth of BS spores. Triplicate tubes of AAMS broth were inoculated with ca. 10³ spores and then stored at 50 psi, 75 psi, 100 psi, 200 psi, 400 psi, and 600 psi CO₂. Triplicate tubes were also inoculated and stored under 50 psi and 600 psi nitrogen. All samples were incubated for from 16 to 87 hours at 55°C. All samples that were treated with CO₂ remained clear while all controls (33 tubes) that were incubated concurrently under ambient conditions became turbid. Samples that were incubated at 55°C under a nitrogen atmosphere (50 psi or 600 psi) behaved as did ambient controls.




Figure 6. Effect of carbonation on germination and outgrowth of Bacillus stearothermophilus spores in AAMS broth during incubation at 55°C.

When BS spores were suspended in carbonated AAMS broth and incubated at 55°C, growth and germination were adversely affected. The effect of carbonation on the growth of BS spores at 55°C is depicted in Figure 6. The data obtained indicate that carbonation of the media completely prevented germination and outgrowth for 6 hours. After 6 hours the spores overcame the inhibitory effect. Carbonation of the media caused the pH to drop from 6.7 to 5.4. Inhibition of germination may be due to the lower pH of the media. However, vegetative growth of BS is reported to be rapid in many foods of pH above 5.0 if held at an elevated temperature (2). Blocher and Busta (4) reported that the minimum pH for germination of BS spores at 55°C is less than 5.4.

Cross et al (5) reported that germination of four strains of *Bacillus* in a yeast-dextrose broth medium was inhibited by bicarbonate at 5, 25, and 50 mM concentration. Similar findings were reported by Hachisuka et al, who reported that 50mM NaHCO₃ inhibit germination of *Bacillus subtilis* spores (16). The effect of 54 mM NaHCO₃ on germination and outgrowth of BS spores in AAMS broth was evaluated and the results are shown in Figure 7. The BS spores were not inhibited by bicarbonate, but rather growth and germination were enhanced. Enhancement of germination may be attributed to the buffering capacity of the NaHCO₃. The pH of the bicarbonate supplemented media was 7.0 after incubation, while the pH of the unsupplemented media was 4.7. The data suggests that NaHCO₃ be added to AAMS routinely to optimize cultivation of BS.

UV absorption spectra of supernates of CO₂ treated and control BS vegetative cells.

Shigehisa et al (19) reported that high hydrostatic pressure (3000-6000 atm) killed all microorganisms tested except *Bacillus cereus* spores.




Figure 7. Effect of 54 mM NaHCO₃ on germination of *Bacillus stearothermophilus* spores in AAMS during incubation at 55°C.

Examination of the supernatant fluid of E. coli cells subjected to such hydrostatic pressure revealed they leaked cytoplasmic RNA that was detected by analysis of the UV absorption spectrum of the culture supernate. Supernates obtained from vegetative cells that had been subjected to 850 psi CO₂ for 17 hours, as well as supernates obtained from control cells, were analyzed by UV spectroscopy. Results of this analysis are shown in Figure 8. The UV spectra revealed no significant difference in the amount of DNA/RNA present in the supernatant fluid of ambient and CO₂ treated cells. However, there was a greater absorbance in the 210-240 nm range in the CO₂ treated than in the ambient sample supernate (19). Further analysis by gas chromatography of the nature of the molecules absorbing in the 210-240 nm range may reveal information regarding the nature of the lesion induced by hyperbaric CO₂ on vegetative cells.




Figure 8. Ultraviolet absorption spectra of culture supernates obtained from:
(A) BS vegetative cells in phosphate buffer, pH 7.0 for 17 hours at 25°C under ambient air.
(B) BS vegetative cells in phosphate buffer, pH 7.0 for 17 hours at 25°C under 850 psi CO₂.

CONCLUSIONS

The objectives of the present study were to determine the effect of hyperbaric carbon dioxide on survival and growth of vegetative cells of Bacillus stearothermophilus as well as its effect on viability, germination and outgrowth of spores.

Vegetative cells were found to be sensitive to damage by relatively low pressure CO₂ treatment for a short time. Complete destruction of 10⁶ cells in AAMS broth was obtained by exposure to 400 psi CO₂ for 1.5 hours.

Spores, however, were found to be unaffected by high pressure carbon dioxide treatment. Spores remained viable after exposure to 800 - 1100 psi CO₂ for as long as 96 hours. Spores survived CO₂ treatment at low (3°C) as well as high (100°C) temperatures. Spores survived CO₂ treatment at pH 4, 4.5, and 7. However, data obtained suggests that high pressure (850-875 psi) CO₂ exposure at high temperature (>90°C) act synergistically to compromise spore viability.

Spores that had undergone a heat activation followed by a 3 hour incubation at 55°C were susceptible to destruction by CO₂. Germination was not necessary for susceptibility to CO₂, for heat activated spores suspended in water were as readily destroyed by CO₂ exposure as were those suspended in a rich nutrient media.

Carbon dioxide at low pressure (50 psi) inhibited the germination and outgrowth of spores. The inhibition was found to be reversible. Spores that had been inhibited germinated when exposed to ambient conditions. These findings suggest that flat sour spoilage of thermal-processed foods could be avoided by incorporating CO₂ at low pressure (50 psi) in the package.

Spore germination was impaired in carbonated AAMS broth. However, the addition of 50 mM NaHCO₃ to AAMS broth enhanced rather than impaired germination and outgrowth.

An attempt was made to characterize the nature of the lesion inflicted by CO₂ on vegetative cells. There was no evidence of a difference in DNA/RNA leakage indicated by analysis of the ultraviolet absorption spectra of culture supernates of CO₂ treated and control cells.

REFERENCES

1. Baker, R. C., R. A. Quereshi and J. H. Hotchkiss. 1986. Effect of an elevated level of carbon dioxide containing atmosphere on the growth of spoilage and pathogenic bacteria at 2°, 5°, and 13°C. *Poult. Sci.* 65: 729-737.
2. Bergey's Manual of Determinative Bacteriology. 8th edition, 1984. The Williams and Wilkins Company, Baltimore, MD., pp. 529-540.
3. Blickstad, E., S. O. Enfors, and G. Molin. 1981. Effect of hyperbaric carbon dioxide pressure on the microbial flora of pork stored at 4° or 14°C. *J. Appl. Bacteriol.* 50: 493-504.
4. Blocher, J. C., and F. F. Busta. 1983. Bacterial spore resistance to acid. *Food Technol.* 37(11): 87-99.
5. Cross, G., J. Wolf and A. N. Barker. 1973. The effect of sodium bicarbonate on the germination of spores of some Bacillus species, pp 199-205. In A. N. Barker G. W. Gould and J. Wolf (ed.), Spore Research, Academic Press, New York.
6. Daniels, J. A., R. Krishnamurthi and S. Rizvi. 1985. A review of effect of carbon dioxide on microbial growth and food quality. *J. Food Protection* 48: (6) 532-537.
7. Dixon, N. M., and D. B. Kell. 1989. A Review. The inhibition by CO₂ of the growth and metabolism of micro-organisms. *J. Appl. Bacteriol.* 67: 109-136.
8. Doyle, M. P. 1983. Effect of carbon dioxide on toxin production by Clostridium botulinum. *Eur. J. Appl. Microbiol. Biotechnol.* 17: 53-56.
9. El-Mabsout, Y. E., and K. E. Stevenson. 1979. Activation of Bacillus stearothermophilus spores at low pH. *J. Food Sc.* 44: 705-709.
10. Enfors, S. O., and G. Molin. 1978. The influence of high concentration of carbon dioxide on the germination of bacterial spores. *J. Appl. Bacteriol.* 45: 279-285.

11. Enfors, S. O., and G. Molin. 1978. Mechanism of the inhibition of spore germination by inert gases and carbon dioxide. In Spores VII, ed. Chambliss, G. & Vary, J.C. pp. 80-84. Washington: Society of Microbiology.
12. Enfors, S. O., and G. Molin. 1980. Effect of high concentrations of carbon dioxide on the growth rate of Pseudomonas fragi, Bacillus cereus, and Streptococcus cremoris. J. Appl. Bacteriol. 48: 409-416.
13. Feeherry, F. E., Munsey, D. T., and D. B. Rowley. 1987. Thermal inactivation and injury of Bacillus stearothermophilus spores. Appl. and Environ. Microbiol. 53: 365-367.
14. Jay, J. M. 1986. Characteristics and growth of thermophilic microorganisms. Ch. 24. In "Modern Food Microbiology," J. M. Jay (Ed.), p. 593. Van Nostrand Reinhold Co., New York, NY.
15. Haas, G. J., H. E. Prescott, E. Dudley, R. Dik, C. Hintlian, and L. Keane. 1989. Inactivation of microorganisms by carbon dioxide under pressure. J. Food Safety. 9: 253-265.
16. Hachisuka, Y., N. Kato, and N. Asano. 1956. Inhibitory effect of sodium carbonate on spore germination of Bacillus subtilis. J. Bacteriology. 71: 250-251.
17. Madden, R. H. and B. Moss, 1987. Extension of shelf life of minced beef by storage in vacuum packages with carbon dioxide. J. Food Protect. 50: 229-233.
18. Molin, G. 1983. The resistance to carbon dioxide of some food related bacteria. Eur. J. Appl. Microbiol. Biotechnol. 18: 214-217.
19. Shigehisa, T., T. Ohmori, A. Saito, S. Taji and R. Hayashi. 1991. Effects of high hydrostatic pressure on characteristics of pork slurries and inactivation of microorganisms associated with meat and meat products. Int. J. Food Microbiol. 12: 207-216.
20. Silliker, J. H. The influence of atmospheres containing elevated levels of CO₂ on growth of psychrotrophic organisms in meat and poultry. Ch. 37. 370-375.

21. Thomas, Y. O., Kraft, A. A., Rust, R. E., and Hotchkiss, D. K. 1984. Effect of carbon dioxide flushing and packaging methods on the microbiology of packaged chickens. J. Food Sci. 49: 1367-1371.
22. Widdowson, J. P. 1967. Bacillus pantothenicus spores activated by dimethylformamide and dimethylsulphoxide. Nature. 214: 812-813.