

1961

NBS MONOGRAPH 19

AD-A280 150



Best Available Copy

COPY 1

Atomic Energy Levels In Crystals

DTIC
ELECTE
Date 1 1994
S B D

15088 N

15088-1

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited



LIBRARY COPY

JUN 5 1961

94-16041



9096

U.S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS

DTIC QUALITY INSURED 1

94 5 27 007

0
9

THE NATIONAL BUREAU OF STANDARDS

Functions and Activities

The functions of the National Bureau of Standards are set forth in the Act of Congress, March 3, 1901, as amended by Congress in Public Law 619, 1950. These include the development and maintenance of the national standards of measurement and the provision of means and methods for making measurements consistent with these standards; the determination of physical constants and properties of materials; the development of methods and instruments for testing materials, devices, and structures; advisory services to government agencies on scientific and technical problems; invention and development of devices to serve special needs of the Government; and the development of standard practices, codes, and specifications. The work includes basic and applied research, development, engineering, instrumentation, testing, evaluation, calibration services, and various consultation and information services. Research projects are also performed for other government agencies when the work relates to and supplements the basic program of the Bureau or when the Bureau's unique competence is required. The scope of activities is suggested by the listing of divisions and sections on the inside of the back cover.

Publications

The results of the Bureau's work take the form of either actual equipment and devices or published papers. These papers appear either in the Bureau's own series of publications or in the journals of professional and scientific societies. The Bureau itself publishes three periodicals available from the Government Printing Office: The Journal of Research, published in four separate sections, presents complete scientific and technical papers; the Technical News Bulletin presents summary and preliminary reports on work in progress; and Basic Radio Propagation Predictions provides data for determining the best frequencies to use for radio communications throughout the world. There are also five series of nonperiodical publications: Monographs, Applied Mathematics Series, Handbooks, Miscellaneous Publications, and Technical Notes.

Information on the Bureau's publications can be found in NBS Circular 460, Publications of the National Bureau of Standards (\$1.25) and its Supplement (\$1.50), available from the Superintendent of Documents, Government Printing Office, Washington 25, D.C.

UNITED STATES DEPARTMENT OF COMMERCE • Luther H. Hodges, *Secretary*
NATIONAL BUREAU OF STANDARDS • A. V. Astin, *Director*

Atomic Energy Levels in Crystals

John L. Prather



National Bureau of Standards Monograph 19

Issued February 24, 1961

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington 25, D.C. - Price 60 cents

Preface

This present work had its inception in 1954, shortly after Professor G. H. Dieke of The Johns Hopkins University, with the support of the U.S. Atomic Energy Commission, initiated a program of experimental studies of the sharp line absorption spectra of crystals. The effort was directed principally to rare earth salt crystals, but some work was also done on the ruby and uranyl salts. Dr. Dieke suggested to the author that a theoretical study to accompany the experimental program would be desirable and encouraged the undertaking of such a study.

It rapidly became apparent on surveying the problem that the techniques of group theory would be of central importance in such a study. Early in 1955, Dr. Charles M. Herzfeld, then of the Naval Research Laboratory in Washington, addressed the Spectroscopy Seminar at Hopkins on the techniques of group theory with particular application to the paramagnetic resonance of salts of the iron group. This was the beginning of a continuing association with Dr. Herzfeld in the study of group theory, an association which has been a source of great personal and professional satisfaction to the author to the present time.

In August of 1955, through the courtesy of Dr. F. G. Brickwedde, then Chief of the Heat and Power Division of the National Bureau of Standards, and Dr. R. P. Hudson, Chief of the Cryogenic Physics Section of the Bureau, the author was appointed a Guest Worker at the Bureau, and office space was provided and the extensive library facilities of the Bureau made available. This appointment continued when Dr. Herzfeld was appointed Chief of the Heat Division.

The present work is largely an exhaustive collection of the results of group theory which apply to the theory of atomic energy levels in crystals, together with a careful, and detailed, but not especially abstract development of the general theory to permit an intelligent application of the results to experimental problems. Dr. Herzfeld, in collaboration with Professor Paul H. Meijer of The Catholic University of America, has written a review on the general subject "Group Theory and Crystal Field Theory", soon to be published, treating the subject from a more abstract viewpoint of modern algebra and concentrating on the fundamental aspects of the theory. This latter work and the present one form a complementary pair of works on the general topic. It is hoped that, together, they will meet the needs of most workers in the field.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Contents

	Page
1. Introduction.....	1
2. The electrostatic field.....	3
2.1. Expansion in tesseral harmonics.....	3
2.2. Symmetry.....	6
2.3. Preferred coordinate systems.....	9
2.4. Cubic point groups.....	12
3. Groups, characters, and representations.....	14
3.1. Representations and quantum mechanics.....	14
3.2. Rotations in three dimensions.....	14
3.3. Group theory and the free atom or ion.....	18
3.4. Group theory and the perturbed ion.....	20
3.5. Group characters and the rep D_7	21
3.6. Noncubic crystallographic groups.....	24
3.7. Noncubic double groups.....	25
3.8. Reps and characters.....	27
3.9. Infinite and cubic groups.....	29
3.10. Table of group characters.....	30
4. Degeneracy, selection rules, and polarization.....	37
4.1. Splitting of free ion levels.....	37
4.2. Kramers degeneracy.....	50
4.3. Orthogonal states and vanishing matrix elements.....	52
4.4. Selection rules and polarization.....	53
5. Matrix elements.....	65
5.1. The reduced matrix element.....	65
5.2. Wigner coefficients.....	66
5.3. $L-S$ coupling and the one-electron approximation.....	72
5.4. Rare earth ground terms.....	73
6. Applications.....	74
6.1. General considerations.....	74
6.2. A crystal field.....	77
6.3. The fluorescence spectrum of the terbium ion.....	78
7. Bibliography.....	82

List of Tables

1. Normalized Legendre functions.....	4
2. Normalized tesseral harmonics.....	5
3. Structure of noncubic point groups.....	8
4. Symmetry elements in tesseral harmonics.....	9
5. Potential coefficients present for various symmetries.....	10
6. Transformation of tesseral harmonics.....	11
7. A character table for the reps D_7	24
8. A table of group characters.....	31
9. Type and number of perturbed levels.....	38
10. Correlation tables.....	46
11. Reduction of the cubic groups.....	48
12. Multipole moment transformations.....	56
13. Selection rules.....	57
14. Angular dependence of multipole components.....	64
15. Wigner coefficients.....	67
16. Reduced matrix elements.....	75
17. A representative crystal field.....	77
18. Fluorescence spectrum of $TbCl_3 \cdot 6H_2O$	79

Atomic Energy Levels in Crystals^{1,2}

John L. Prather³

Discrete energy levels observed within certain crystals are treated as due to perturbations of the energy levels of the free ion by an electrostatic field arising from the crystal lattice. The analytic procedures for determining the field from the charge configuration are given, and the resulting fields are classified according to their symmetry. After a general survey of group-theoretical ideas, the applicable groups are analyzed in detail, and characters appropriate for both integral and half-integral angular momenta of the free ion are tabulated. These are applied to the determination of the number and type of levels arising from a free ion level with $J \leq 8$. The results of this analysis are tabulated, as are the selection rules for electric dipole, magnetic dipole, and electric quadrupole transitions. Calculation of the perturbation matrix elements by the use of Wigner and Racah coefficients is discussed. Examples of the application of these several techniques to specific problems are given.

1. Introduction

In contrast to the sharp lines or resolvable bands characteristic of the optical absorption spectra of free atomic and molecular systems, such spectra for solids consist for the most part of broad absorption regions or bands without any resolvable structure. These spectra are therefore of little used for analysis of the details of the solids for which they are observed, revealing only general information such as the presence or absence of certain molecular groupings. Nevertheless, it was early recognized in the development of optical spectroscopy that certain naturally occurring minerals did in fact show a characteristic line absorption spectrum. The earliest observations were on the crystals of salts of the iron group elements and of the rare earth elements, though later observations have shown similar spectra in crystals of all elements characterized by a partially filled inner shell of electrons. In this present work we shall be concerned primarily with the rare-earth spectra, though we shall sketch the application of the development to other possible cases.

The earliest data on these spectra, covering about 50 years before 1905 are reviewed by Kayser (see bibliography). Subsequent to the discovery of the Zeeman effect in 1895, J. Becquerel initiated an extensive investigation on the effects of magnetic fields on the absorption lines of natural rare earth crystals, their polarization, and particularly their rotation of the plane of polarization of light (Faraday effect). These measurements, carried out both at room temperature and at liquid air temperature, are summarized in his paper of 1908. In this same year, Becquerel and Onnes reported on the spectra and magnetic effects at liquid hydrogen temperatures, and measurements at liquid helium temperatures were finally reported in 1926 by Becquerel, Onnes, and de Haas. du Bois and Elias (1908, 1911) also

reported work on chromium, cobalt, and uranium crystals as well as on the rare earths.

The year 1929 marked the beginning of an extensive program of investigation of these spectra at various temperatures by Spedding, Freed, and collaborators. The principal experimental advance over earlier work lay in the use of synthetic crystals of more definite composition than the natural crystals, which usually consisted of a mixture of rare earths of varying composition. Hence there had been little if any reliability in intercomparison of the results obtained from various crystals, even of the same nominal type. This was a considerable advance over the previous attempts at a systematic experimental approach to these spectra, though by today's standards even these crystals were not of satisfactory purity. These measurements were also much aided by continuing developments in the theory of these spectra, which we shall trace below.

All of these spectra show the same general pattern and behavior upon reducing the temperature. The absorption lines for various elements fall into groups in several regions of the spectrum—the general wavelength regions where the lines fall depend on the rare earth involved, while details concerning the number of lines, the exact position, intervals, Zeeman effect, etc., depend on the negative ions involved and on the details of the crystal structure. In comparison with room temperature measurements, those at reduced temperatures show generally sharper lines, together with a slight overall shift of the groups to the red and increased separation of the lines within a group (Spedding and Bear, 1932). The number of lines also changes; certain "high temperature lines" disappear, while others appear or become more intense at lower temperatures. Such lines have also been observed in solutions of the ions, though these lines are generally broader than in the crystals.

Although there was some theoretical speculation on these spectra, no theoretically satisfactory contributions save the derivation of empirical energy level diagrams could be made prior to the advent of quantum mechanics. The first major

¹ Based on a dissertation submitted to the Faculty of Philosophy of The Johns Hopkins University in partial fulfillment of the requirements for the degree Doctor of Philosophy.

² Supported in part by the U.S. Air Force, through the Air Force Office of Scientific Research of the Air Research and Development Command, under Contract No. CSO and A 690-56-21.

³ Guest worker, formerly from The George Washington University, now with Beloit College, Beloit, Wis.

contribution directly related to the rare earth spectra was the calculation by Hund (1925) of the room temperature paramagnetic susceptibility of the rare earth crystals on the assumption that the crystals consisted of an independent statistical assemblage of ions, characterized by an unfilled shell of $4f$ electrons which became increasingly filled as one proceeded up the periodic table. The ground states of these ions were calculated on the assumption that Hund's rules for free atoms were valid, i.e., that of all the terms arising from the given $4f^n$ configuration, that with the greatest multiplicity had the lowest energy, and of those of this greatest multiplicity, the largest L value was lowest. For a given $L-S$ term, the resulting multiplet was taken as normal in the first half of the series (the $4f$ shell being less than half filled) and as inverted in the last half of the series. The agreement with experiment was quite good, save in the cases of europium and samarium, and these two discrepancies were corrected by the second order calculations of Van Vleck and Frank (1929), Hund having considered only the lowest level of the ground multiplet as contributing to the susceptibility. This agreement with experiment not only gave evidence for the electronic structure of the rare earth ions in the crystals, but also suggested that the effects on the magnetic properties of the ion due to inclusion in the crystal were small in comparison with kT at 300° K and also small in comparison with the multiplet splitting of the ground $L-S$ term.

It was perhaps natural to assume that the line spectra observed were similarly due to the spectra of the free ions, modified to some slight degree by their inclusion in the crystal, but the first suggestion to this effect was made by Brunetti and Ollano (1929), who considered the crystal as a perturbing electrostatic field on the free ion spectrum. This idea was also developed by Freed and Spedding (1929) in initiating their program of experimental observations noted above. The classic paper in these early years of the theory is that by Bethe (1929). Bethe pointed out that the details of the theoretical treatment depended on the strength of this external crystal field compared to the other interactions present (principally the Coulomb interaction of the electrons and the spin-orbit interaction), and that the symmetry alone of the external field, independent of its magnitude, would determine the number and type of levels arising from a given level of the free ion. These were specifically computed by Bethe for certain symmetries, emphasizing in the process the importance of group-theoretical ideas for such calculations, and the properties of one-electron wave functions under these same crystal fields were also calculated. These ideas were later extended to treat the Zeeman effect in crystals (Bethe, 1930). At about the same time, Kramers (1929, 1930) considering principally the Faraday effect, formulated the fundamental requirement

that these electrostatic fields arising from the crystal must leave every level of an odd-electron system at least twofold degenerate.

Although it seemed clear that the idea of an electrostatic perturbation acting upon the levels of the free ion was the key to the problem, it was not at all certain what transitions were responsible for the lines actually observed. The situation was complicated by utter ignorance of the actual free ion spectrum, and even as recently as 1958 only that of CeIV is known (Lang, 1936). The earliest suggestions were that the transition involved a $4f$ electron excited to either $5d$ or $6s$ (Spedding, 1931; Spedding, 1933). These were based principally on analogy with the selection rules applicable to free ions. However, the sharpness of the lines indicates a freedom from interaction with other ions which would not be expected from such levels (involving $5d$ or $6s$) having a fair probability of being at appreciable distances from the nucleus. Furthermore, the general intensity of the lines relative to the high concentration of ions in a solid yields transition probabilities corresponding to transitions forbidden in the usual type of atomic spectra (Van Vleck, 1937). For this reason, it is today generally considered that the lines come from transitions between terms arising from the ground $4f^n$ configuration. This view is strengthened by the simplicity of such sharp line spectra in the salts of cerium and ytterbium (Dieke and Crosswhite, 1956) both of which have only one excited level from the ground configurations $4f^1$ and $4f^{13}$, respectively. Van Vleck also made some estimates of the relative probability of transitions due to electric quadrupole radiation, magnetic dipole radiation, electric dipole radiation arising from absence of a static center of symmetry in the crystal field (which destroys the characterization of levels as odd or even), and electric dipole radiation arising from destruction of a static center of symmetry by lattice vibrations. It was concluded that there was no clear quantitative reason to reject completely any of these as being not responsible for the lines observed.

During this period (the 1930's) detailed calculations were made by Penney and Schlapp (1932, Schlapp and Penney) based on the assumption of a cubic crystal field, adjusting the parameters to fit observed magnetic susceptibility data. Although good agreement with experiment in this regard was obtained, it is in marked disagreement with the spectral data. This may be ascribed to the insensitivity of susceptibility data (a statistical average over all ions present) to details of the crystal field. The same objection also applies to attempts to derive the crystal field from specific heat data (Ahlberg et al., 1937). It is now known that the symmetry of the cubic system is too high to account for the spectroscopic observations.

Apart from theoretical considerations of the rare earth spectra alone, several general theoretical contributions have been made which are applicable in major part to the determination of the energy

levels of rare earth ions in crystals. Since these will be referred to in detail later in this work, we shall merely mention them for general background at this point. The general applicability of group theory to problems of atomic structure has been emphasized by Weyl (1928), clarified by Eckart (1930), and treated at length by Wigner (1931). The calculations of Bethe (1929) for certain symmetry groups were extended by Wigner (1930) to other symmetries, though this was in connection with problems of molecular vibrations. The monumental work of Condon and Shortley (1935) is invaluable for the study of the free ion spectrum.

While experimental data on many different crystals were collected by many observers, using many different techniques (absorption spectra, Zeeman effect, fluorescence spectra, magnetic measurements, specific heat data) during the following years, the next major advance was the application of paramagnetic resonance techniques to the rare earth crystals. A general review of this work may be found in the paper by Bleaney and Stevens (1953), while additional data are given in the supplementary article by Bowers and Owen (1955). This method has the advantage of giving data of high accuracy and resolution—its major limitation is in the fact that with rare exceptions only data on the ground level in the crystal are obtained.

The next major advance in the general theory was the work of Racah (1942, 1942a,b) on complex spectra, which provided very powerful general techniques for the solution of problems in many-electron systems. His results have been made applicable to specific numerical calculations by the recent appearance of numerical tables by Biedenharn (1952), Simon (1954), and Simon et al. (1954). While the procedures of Racah have been used in several fields, and their applicability to the problem of rare earth spectra in crystals has been mentioned by Elliott and Stevens (1953), no widespread use of them has been made in crystal spectra problems. The general work of Bethe (1929) on the applications of group theory to the problem has subsequently been applied by various authors to special cases, but no general overall survey of the possibilities has been given. The results have been obtained by Hellwege (1948, 1948a-d) without the explicit use of group theory, but much elegance and conciseness is lost thereby, and the results are presented in a rather cumbersome form. A general survey of these and many other aspects of the problem by Fick and Joos (1957) has recently appeared.

2. The Electrostatic Field

2.1. Expansion in Tesseral Harmonics

Let us consider, in a given frame of reference, an ion with a charge q located at the point described in spherical coordinates as (R, α, β) . The potential at the point (r, θ, ϕ) may be expressed as a series of Legendre polynomials in the variable $\cos \gamma$,

Recent renewed interest in these spectra and the associated properties of the ions has generated a requirement for a comprehensive survey and development of the theoretical techniques required for an analysis of such spectra. While the theory in its most general terms is known, its applications in the past have either been concerned with a particular salt or ion, or have used only a portion of the theoretical equipment available, are widely scattered through the literature, and their usefulness is limited by differences in conventions of sign, phase, normalization, notation, etc. It is the purpose of this paper to provide a comprehensive theoretical approach to the analysis of these spectra in a form particularly adapted to the requirements of the experimental workers. In the interest of meeting this aim, mathematical rigor and long purely algebraic manipulations have for the most part been omitted, except where necessary for an understanding of the concepts and procedures involved. It is hoped that it will be of value to experimentalists in the field who have had only a general background in the ordinary theory of the atomic spectra of free atomic systems.

Since the general problem is that of computing the effect on the free ion of the electrostatic perturbations arising from the crystal lattice, we begin by considering in section 2 the development of this potential in a power series of tesseral harmonics, in explicit terms of the charge and position of the ions giving rise to the field. An important property of such fields is their symmetry, and this will be discussed in detail. In section 3, we shall consider the connection between the theory of groups and quantum mechanics. We begin by sketching the group-theoretical ideas associated with the problem of the free atom. This leads naturally into those modifications caused by the inclusion of the ion in the crystal lattice. The properties of the groups involved in the present problem are then considered in detail. In section 4, we present explicitly the results of the considerations of section 3 to the degeneracy of levels, selection rules and polarization of lines in the spectrum of the crystal. In section 5, we consider the calculation of the matrix elements of the perturbing electrostatic potential. Finally, we conclude in section 6 by a review of the various factors influencing the experimentally observed spectra, correlated with the spectra of the free ion, and some examples of the application of the techniques to specific problems.

where γ is the angle between the directions given by α, β , and θ, ϕ . The unit vectors in the two directions are $\sin \alpha \cos \beta \mathbf{i} + \sin \alpha \sin \beta \mathbf{j} + \cos \alpha \mathbf{k}$, and $\sin \theta \cos \phi \mathbf{i} + \sin \theta \sin \phi \mathbf{j} + \cos \theta \mathbf{k}$, respectively. Thus, $\cos \gamma$ is the scalar product of the two unit vectors and $\cos \gamma = \sin \theta \sin \alpha \cos \beta \cos \phi + \sin \theta \sin \alpha \sin \beta \sin \phi + \cos \theta \cos \alpha = \sin \alpha \sin \theta \cos(\phi - \beta) +$

$\cos \theta \cos \alpha$. The potential is given, for $r \leq R$, by

$$V = q \sum_{N=0}^{\infty} \frac{r^N}{R^{N+1}} P_N(\cos \gamma). \quad (2.1)$$

The total potential is the sum of similar contributions from each ion of the crystal lattice. In the above expression, the contribution of each ion is described with reference to a different axis, and it is convenient to refer each contribution to the same axis, which we choose to be the z -axis of the given frame of reference.

This may be accomplished through the use of the addition theorem of spherical harmonics, which expresses $P_N(\cos \gamma)$ in terms of α , β , θ , and ϕ (Stratton, 1941).

$$P_N(\cos \gamma) = P_N(\cos \alpha) P_N(\cos \theta) + 2 \sum_{M=1}^N \frac{(N-M)!}{(N+M)!} P_N^M(\cos \alpha) P_N^M(\cos \theta) \cos M(\phi - \beta). \quad (2.2)$$

The potential due to the i -th charge can then be put into the form

$$V_i = q_i \sum_{N=0}^{\infty} \frac{r^N}{R^{N+1}} \left[P_N(\cos \alpha_i) P_N(\cos \theta) + 2 \sum_{M=1}^N \frac{(N-M)!}{(N+M)!} P_N^M(\cos \alpha_i) P_N^M(\cos \theta) (\cos M\phi \cos M\beta_i + \sin M\phi \sin M\beta_i) \right]. \quad (2.3)$$

The total potential may then be obtained by summing over all charges in the lattice.

As used above, the $P_N^M(\cos \theta)$ are not normalized in the quantum-mechanical sense, since

$$\int_{-1}^1 P_N^M(\cos \theta) P_L^K(\cos \theta) d(\cos \theta) = \delta_{N,K} \delta_{M,L} \frac{2}{2N+1} \frac{(N+M)!}{(N-M)!}$$

and it is consequently convenient to define

$$\Theta_N^M = \sqrt{\frac{2N+1}{2} \frac{(N-M)!}{(N+M)!}} P_N^M(\cos \theta). \quad (2.4)$$

Tabulated functions Θ_N^M for N up to and including 6 are given in table 1. Similarly, the functions $\frac{1}{\sqrt{\pi}} \sin M\phi$, $\frac{1}{\sqrt{\pi}} \cos M\phi$, and $\frac{1}{\sqrt{2\pi}}$ constitute an orthonormal set of functions over the interval $[0, 2\pi]$ in the variable ϕ . We shall define the following tesseral harmonics, which are functions of position on the surface of the unit sphere, and may be evaluated either from θ , ϕ , or from x , y ,

TABLE 1. Normalized Legendre functions

$\Theta_0^0 = \frac{\sqrt{2}}{2}$	$\Theta_6^0 = \frac{3\sqrt{2}}{16} (35 \cos^4 \theta - 30 \cos^2 \theta + 3)$
$\Theta_1^0 = \frac{\sqrt{6}}{2} \cos \theta$	$\Theta_6^1 = \frac{3\sqrt{10}}{8} (7 \cos^3 \theta - 3 \cos \theta) \sin \theta$
$\Theta_1^1 = \frac{\sqrt{3}}{2} \sin \theta$	$\Theta_6^2 = \frac{3\sqrt{5}}{8} (7 \cos^2 \theta - 1) \sin^2 \theta$
$\Theta_2^0 = \frac{\sqrt{10}}{4} (3 \cos^2 \theta - 1)$	$\Theta_6^3 = \frac{3\sqrt{70}}{8} \cos \theta \sin^3 \theta$
$\Theta_2^1 = \frac{\sqrt{15}}{2} \sin \theta \cos \theta$	$\Theta_6^4 = \frac{3\sqrt{35}}{16} \sin^4 \theta$
$\Theta_2^2 = \frac{\sqrt{15}}{4} \sin^2 \theta$	$\Theta_6^5 = \frac{\sqrt{22}}{16} (63 \cos^2 \theta - 70 \cos^2 \theta + 15 \cos \theta)$
$\Theta_3^0 = \frac{\sqrt{14}}{4} (5 \cos^3 \theta - 3 \cos \theta)$	$\Theta_6^6 = \frac{\sqrt{165}}{16} (21 \cos^4 \theta - 14 \cos^2 \theta + 1) \sin \theta$
$\Theta_3^1 = \frac{\sqrt{42}}{8} (5 \cos^2 \theta - 1) \sin \theta$	$\Theta_7^0 = \frac{\sqrt{1,155}}{8} (3 \cos^3 \theta - \cos \theta) \sin^2 \theta$
$\Theta_3^2 = \frac{\sqrt{105}}{4} \sin^2 \theta \cos \theta$	$\Theta_7^1 = \frac{\sqrt{770}}{32} (9 \cos^2 \theta - 1) \sin^3 \theta$
$\Theta_3^3 = \frac{\sqrt{70}}{8} \sin^3 \theta$	$\Theta_7^2 = \frac{3\sqrt{385}}{16} \cos \theta \sin^4 \theta$
	$\Theta_7^3 = \frac{3\sqrt{154}}{32} \sin^5 \theta$
	$\Theta_8^0 = \frac{\sqrt{26}}{32} (231 \cos^4 \theta - 315 \cos^2 \theta + 105 \cos^2 \theta - 5)$
	$\Theta_8^1 = \frac{\sqrt{273}}{16} (33 \cos^5 \theta - 30 \cos^3 \theta + 5 \cos \theta) \sin \theta$
	$\Theta_8^2 = \frac{\sqrt{2,730}}{64} (33 \cos^4 \theta - 18 \cos^2 \theta + 1) \sin^2 \theta$
	$\Theta_8^3 = \frac{\sqrt{2,730}}{32} (11 \cos^3 \theta - 3 \cos \theta) \sin^3 \theta$
	$\Theta_8^4 = \frac{3\sqrt{91}}{32} (11 \cos^2 \theta - 1) \sin^4 \theta$
	$\Theta_8^5 = \frac{3\sqrt{2,002}}{32} \cos \theta \sin^5 \theta$
	$\Theta_8^6 = \frac{\sqrt{6,006}}{64} \sin^6 \theta$

and z . They are given as functions of x , y , and z in table 2:

$$\begin{aligned} S_N^M &= \Theta_N^M \frac{\sin M\phi}{\sqrt{\pi}}, S_N^0 = 0, \\ C_N^M &= \Theta_N^M \frac{\cos M\phi}{\sqrt{\pi}}, C_N^0 = \frac{1}{\sqrt{2\pi}} \Theta_N^0. \end{aligned} \quad (2.5)$$

TABLE 2. Normalized tesseral harmonics

$C_0^2 = \frac{\sqrt{6}}{2} \frac{1}{\sqrt{2\pi}} \frac{z}{r}$	$C_0^3 = \frac{\sqrt{14}}{4} \frac{1}{\sqrt{2\pi}} \frac{2z^3 - 3xz^2 - 3y^2z}{r^3}$	$C_0^4 = \frac{\sqrt{1,155}}{8} \frac{1}{\sqrt{\pi}} \frac{(2z^2 - x^2 - y^2)(x^2 - y^2)z}{r^5}$
$C_1^2 = \frac{\sqrt{3}}{2} \frac{1}{\sqrt{\pi}} \frac{x}{r}$	$C_1^3 = \frac{\sqrt{42}}{8} \frac{1}{\sqrt{\pi}} \frac{4z^2x - x^3 - xy^2}{r^3}$	$S_0^4 = \frac{\sqrt{1,155}}{8} \frac{1}{\sqrt{\pi}} \frac{(2z^2 - x^2 - y^2)(2xyz)}{r^5}$
$S_1^2 = \frac{\sqrt{3}}{2} \frac{1}{\sqrt{\pi}} \frac{y}{r}$	$S_1^3 = \frac{\sqrt{42}}{8} \frac{1}{\sqrt{\pi}} \frac{4z^2y - x^2y - y^3}{r^3}$	$C_1^4 = \frac{\sqrt{770}}{32} \frac{1}{\sqrt{\pi}} \frac{(8z^2 - x^2 - y^2)(x^2 - 3xy^2)}{r^5}$
$C_2^2 = \frac{\sqrt{10}}{4} \frac{1}{\sqrt{2\pi}} \frac{2z^2 - x^2 - y^2}{r^2}$	$C_2^3 = \frac{\sqrt{105}}{4} \frac{1}{\sqrt{\pi}} \frac{zx^2 - zy^2}{r^3}$	$S_1^4 = \frac{\sqrt{770}}{32} \frac{1}{\sqrt{\pi}} \frac{(8z^2 - x^2 - y^2)(3x^2y - y^3)}{r^5}$
$C_1^3 = \frac{\sqrt{15}}{2} \frac{1}{\sqrt{\pi}} \frac{xz}{r^2}$	$S_2^3 = \frac{\sqrt{105}}{4} \frac{1}{\sqrt{\pi}} \frac{2xyz}{r^3}$	$C_0^5 = \frac{3\sqrt{385}}{16} \frac{1}{\sqrt{\pi}} \frac{(x^4 - 6x^2y^2 + y^4)z}{r^5}$
$S_2^3 = \frac{\sqrt{15}}{2} \frac{1}{\sqrt{\pi}} \frac{yz}{r^2}$	$C_3^3 = \frac{\sqrt{70}}{8} \frac{1}{\sqrt{\pi}} \frac{x^3 - 3xy^2}{r^3}$	$S_2^5 = \frac{3\sqrt{385}}{16} \frac{1}{\sqrt{\pi}} \frac{(x^2 - y^2)(4xyz)}{r^5}$
$C_0^4 = \frac{\sqrt{15}}{4} \frac{1}{\sqrt{\pi}} \frac{x^2 - y^2}{r^2}$	$S_3^3 = \frac{\sqrt{70}}{8} \frac{1}{\sqrt{\pi}} \frac{3x^2y - y^3}{r^3}$	$C_1^5 = \frac{3\sqrt{154}}{32} \frac{1}{\sqrt{\pi}} \frac{x^5 - 10x^3y^2 + 5xy^4}{r^5}$
$S_3^4 = \frac{\sqrt{15}}{4} \frac{1}{\sqrt{\pi}} \frac{2xy}{r^2}$		$S_3^5 = \frac{3\sqrt{154}}{32} \frac{1}{\sqrt{\pi}} \frac{y^5 - 10y^3x^2 + 5yx^4}{r^5}$
		$C_2^5 = \frac{\sqrt{26}}{32} \frac{1}{\sqrt{2\pi}} \frac{16x^5 - 120(x^2 + y^2)x^3 + 90(x^2 + y^2)^2z^2 - 5(x^2 + y^2)^3}{r^5}$
$C_0^5 = \frac{3\sqrt{2}}{16} \frac{1}{\sqrt{2\pi}} \frac{8z^5 + 3x^4 + 3y^4 - 24xz^3 - 24y^2z^3 + 6x^2y^2}{r^5}$	$C_1^5 = \frac{\sqrt{273}}{16} \frac{1}{\sqrt{\pi}} \frac{[8z^5 - 20z^3(x^2 + y^2) + 5z(x^2 + y^2)^2]x}{r^5}$	
$C_1^5 = \frac{3\sqrt{10}}{8} \frac{1}{\sqrt{\pi}} \frac{(4z^2 - 3x^2 - 3y^2)(xz)}{r^4}$	$S_1^5 = \frac{\sqrt{273}}{16} \frac{1}{\sqrt{\pi}} \frac{[8z^5 - 20z^3(x^2 + y^2) + 5z(x^2 + y^2)^2]y}{r^5}$	
$S_1^5 = \frac{3\sqrt{10}}{8} \frac{1}{\sqrt{\pi}} \frac{(4z^2 - 3x^2 - 3y^2)(yz)}{r^4}$	$C_2^5 = \frac{\sqrt{2,730}}{64} \frac{1}{\sqrt{\pi}} \frac{[16z^4 - 16(x^2 + y^2)z^2 + (x^2 + y^2)^2](x^2 - y^2)}{r^5}$	
$C_2^5 = \frac{3\sqrt{5}}{8} \frac{1}{\sqrt{\pi}} \frac{(6z^2 - x^2 - y^2)(x^2 - y^2)}{r^4}$	$S_2^5 = \frac{\sqrt{2,730}}{64} \frac{1}{\sqrt{\pi}} \frac{[16z^4 - 16(x^2 + y^2)z^2 + (x^2 + y^2)^2](2xy)}{r^5}$	
$S_2^5 = \frac{3\sqrt{5}}{8} \frac{1}{\sqrt{\pi}} \frac{(6z^2 - x^2 - y^2)(2xy)}{r^4}$	$C_3^5 = \frac{\sqrt{2,730}}{32} \frac{1}{\sqrt{\pi}} \frac{(8z^2 - 3x^2 - 3y^2)(x^2 - 3y^2)(xz)}{r^5}$	
$C_3^5 = \frac{3\sqrt{70}}{8} \frac{1}{\sqrt{\pi}} \frac{(x^2 - 3y^2)(xz)}{r^4}$	$S_3^5 = \frac{\sqrt{2,730}}{32} \frac{1}{\sqrt{\pi}} \frac{(8z^2 - 3x^2 - 3y^2)(3x^2 - y^2)(yz)}{r^5}$	
$S_3^5 = \frac{3\sqrt{70}}{8} \frac{1}{\sqrt{\pi}} \frac{(3x^2 - y^2)(yz)}{r^4}$	$C_4^5 = \frac{3\sqrt{91}}{32} \frac{1}{\sqrt{\pi}} \frac{(10z^2 - x^2 - y^2)(x^4 - 6x^2y^2 + y^4)}{r^5}$	
$C_4^5 = \frac{3\sqrt{35}}{16} \frac{1}{\sqrt{\pi}} \frac{x^4 - 6x^2y^2 + y^4}{r^4}$	$S_4^5 = \frac{3\sqrt{91}}{32} \frac{1}{\sqrt{\pi}} \frac{(10z^2 - x^2 - y^2)(4x^2y - 4xy^2)}{r^5}$	
$S_4^5 = \frac{3\sqrt{35}}{16} \frac{1}{\sqrt{\pi}} \frac{4xy(x^2 - y^2)}{r^4}$	$C_5^5 = \frac{3\sqrt{2,002}}{32} \frac{1}{\sqrt{\pi}} \frac{(x^5 - 10x^3y^2 + 5xy^4)z}{r^5}$	
$C_0^6 = \frac{\sqrt{22}}{16} \frac{1}{\sqrt{2\pi}} \frac{8z^6 - 40z^4(x^2 + y^2) + 15(x^2 + y^2)^2z}{r^6}$	$S_5^5 = \frac{3\sqrt{2,002}}{32} \frac{1}{\sqrt{\pi}} \frac{(y^5 - 10y^3x^2 + 5yx^4)z}{r^5}$	
$C_1^6 = \frac{\sqrt{165}}{16} \frac{1}{\sqrt{\pi}} \frac{(8z^4 + x^4 + y^4 + 2x^2y^2 - 12x^2z^2 - 12y^2z^2)x}{r^5}$	$C_0^6 = \frac{\sqrt{6,006}}{64} \frac{1}{\sqrt{\pi}} \frac{x^6 - 15x^4y^2 + 15x^2y^4 - y^6}{r^6}$	
$S_1^6 = \frac{\sqrt{165}}{16} \frac{1}{\sqrt{\pi}} \frac{(8z^4 + x^4 + y^4 + 2x^2y^2 - 12x^2z^2 - 12y^2z^2)y}{r^5}$	$S_1^6 = \frac{\sqrt{6,006}}{64} \frac{1}{\sqrt{\pi}} \frac{(3x^2 - y^2)(x^2 - 3y^2)(2xy)}{r^6}$	

With these substitutions, the potential may now be written in the form

$$V = \sum_{N=0}^{\infty} r^N \sum_{M=0}^N (C_N^M C_N^M + S_N^M S_N^M) \quad (2.6)$$

where the coefficients C_N^M and S_N^M depend on the size and location of the neighboring ions and are given explicitly by

$$C_N^M = \frac{4\pi}{2N+1} \sum_i \frac{q_i}{R_{i,N+1}^M} C_N^M(\alpha_i, \beta_i) \quad (2.7a)$$

$$S_N^M = \frac{4\pi}{2N+1} \sum_i \frac{q_i}{R_{i,N+1}^M} S_N^M(\alpha_i, \beta_i) \quad (2.7b)$$

We recall here that the i -th ion, charge q_i , is located in spherical coordinates at (R_i, α_i, β_i) . The terms $C_N^M(\alpha_i, \beta_i)$ and $S_N^M(\alpha_i, \beta_i)$ represent the quantities (2.5) evaluated for the several pairs of angles involved in the summation.

We shall also have occasion to utilize the complex form of (2.5), defining

$$Y_N^M = \frac{(-1)^M}{\sqrt{2}} (C_N^M + iS_N^M), \quad Y_N^{-M} = \frac{1}{\sqrt{2}} (C_N^M - iS_N^M). \quad (2.8)$$

and

$$Y_N^0 = C_N^0$$

This will similarly require, in lieu of (2.7a, b) the coefficients

$$Y_N^M = \frac{(-1)^M}{\sqrt{2}} (C_N^M - iS_N^M), \quad Y_N^{-M} = \frac{1}{\sqrt{2}} (C_N^M + iS_N^M). \quad (2.9)$$

This expression (2.6) for the potential diverges for $r > R$. As we shall see later, however, we shall require only terms for N up to and including 6, and we shall assume that the wave function of the free ion vanishes sufficiently rapidly for large r that negligible error is made by integrating to infinite distances.

2.2. Symmetry

The effects of the electrostatic field on the free ion spectrum may be separated into two categories: first, those depending on the qualitative nature of the field (its shape or symmetry), which is expressed through the mere presence or absence of certain of the C_N^M or S_N^M ; and second, the quantitative details which depend on the sign and magnitude of the C_N^M and S_N^M . It is convenient to consider separately these two characteristics of the field.

There are several standard and equivalent techniques for describing the symmetry of the electrostatic field. For example, given an arbitrary point (x, y, z) of the field, one may indicate the coordinates of all points having the same value of the potential, such as $(x, -y, z)$, $(-x, y, z)$, $(-x, -y, z)$. Another widely used scheme is to

indicate by a symbolic notation the operations which convert a given initial point into one of equal potential. Thus, in the example just given, the operations are respectively a reflection in the x - z plane, a reflection in the y - z plane, and a rotation of π about the z axis, and may be designated as σ_{xz} , σ_{yz} , and C_2 . Finally, for a given point (x, y, z) one may give those 3×3 matrices which transform the original point into one of equal potential. If one is thoroughly familiar with the notation and properties of the various rotations, reflections, and the inversion, there is no reason to prefer one of these to the other. If one does not have this familiarity, the use of the matrices has the advantage that the effect of two successive operations can be computed by matrix multiplication without recourse to geometric arguments. Thus, if $\mathbf{r}' = \mathbf{A}\mathbf{r}$ and $\mathbf{r}'' = \mathbf{B}\mathbf{r}'$ for all \mathbf{r} , then $\mathbf{r}'' = \mathbf{A}\mathbf{B}\mathbf{r}$ is also a point of equal potential as is $\mathbf{r}'' = \mathbf{B}\mathbf{A}\mathbf{r}$. For the problem to be considered here, there are only a few basic matrices which must be considered. These, together with their symbolic notation (Schönflies, 1923) are:

$$C_\infty = \begin{pmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad C_6 = \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$C_4 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad C_3 = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$C_2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad C'_2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

$$E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad I = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

Thus, the point (or vector) $\mathbf{r} = (x, y, z)$ under the operation C_∞ becomes

$$\mathbf{r}' = (x \cos \phi - y \sin \phi, x \sin \phi + y \cos \phi, z).$$

We see that this is equivalent to a rotation of the vector about the z -axis by an angle ϕ , counterclockwise (i.e., by $+\phi$ in terms of the right-hand

screw convention) when looking toward the origin along the positive z -axis. This may also be interpreted as a new description of the old point (x, y, z) in terms of a new coordinate system where the coordinate system has been rotated by the angle ϕ in the opposite sense. In figure 1, we have initially (in two dimensions only) the vector r in the




FIGURE 1

x, y frame. The operation C_n will then give r' in the x, y frame. However, this is clearly the same as r described in the rotated frame x', y' . We shall use both interpretations in the present work.

$$IC_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} = \sigma_h$$

$$IC_2^2 = \begin{pmatrix} 1 & -3/2 & 0 \\ 3/2 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} = \sigma_h C_6 = S_6$$

$$IC_2' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \sigma_v$$

$$IC_2'^2 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} = \sigma_v C_4 = S_4$$

There are only a finite number of possible combinations of these basic operations which are of interest in the present work. These are diagrammed in table 3 and are known as crystallographic point groups, since each of the combinations constitutes a mathematical group, the elements of which leave the potential of a point unchanged. In interpreting table 3, the starting point is the column headed C_n . The groups C_n for the values of n for which we are interested, consist of the operation C_n and its n distinct powers, so a total of n symmetry operations are represented in the group C_n . This set of operations may be enlarged by adding the element I (moving along the diagonal to the left), as is indicated under the heading $C_n \times I$. This nota-

The designation C_n represents a n -fold axis of symmetry ($n=2,3,4,6$), since the matrix is obtained from C_∞ by restricting ϕ to $2\pi/n$. Each rotation by $2\pi/n$ yields an equivalent point in the potential field and, including the original one, there are n such altogether. The element C_2 is a twofold axis of symmetry about the y -axis—it is characteristic of the collections of these symmetry operations to be emphasized here (the noncubic point groups) that there is at most one axis having more than twofold symmetry, and hence rotations about the y -axis can be restricted to those of order two. The element I , the inversion, is noteworthy in the above as being the only one with a determinant -1 , all others being $+1$. Viewed as an operation on a coordinate system, the inversion represents the transformation to left-handed rather than the usual right-handed system.

In addition to the symmetry operations given above, there are operations which may be represented as the product of the element I with those rotations given above. If we are considering a collection of these symmetry operations containing both I and the rotation, no fundamentally new information is obtained by considering the products, but there are certain collections of symmetry operations which include some of these product operations but not the factor operations separately. These product operations, together with their symbolic notation, are as follows:

tion represents the fact that I commutes with all powers of C_n . The group C_n may also be enlarged by adding the element C_2 (moving along the diagonal to the right). Here, the heading $\{C_n C_2\}$ represents that the various operations do not now necessarily commute, but that all possible combinations of them and their powers are included. Both of these enlarged groups have $2n$ elements. To the latter set, I may now be added (moving farther to the right), obtaining $4n$ operations. From these groups containing I , in some cases groups of lower order may be obtained by suppressing half the elements, including the element I itself but not all products containing I . For example, let us consider the group C_2 , which contains the elements C_2 and E . If we add I , we

TABLE 3. Structure of noncubic point groups

Order	Sub-group	$C_n \times I$	C_n	$\{C_n, C_2\}$	$\{C_n, C_2\} \times I$	Sub-group
1			E			
2	$C_2 = \sigma_h$		I	C_2		
3				C_3		
4	S_4			C_2	D_2	C_{2v}
6	C_{3h}			C_3	D_3	C_{3v}
8				C_4	D_4	C_{4v}
12				C_6	D_6	C_{6v}
16					D_{4h}	C_{2v}
24					D_{6h}	C_{3v}
∞		$C_{\infty h}$	C_{∞}	D_{∞}	$D_{\infty h}$	$C_{\infty v}$

obtain C_{2h} , containing C_2 , E , C_2I , and I . From this we obtain C_2 by considering only E and $C_2I = \sigma_h$. Similarly, by adding C_2' to C_2 , we obtain D_2 , which involves E , C_2 , C_2' , and C_2C_2' . By the further inclusion of I , we obtain D_{2h} with E , C_2 , C_2' , C_2C_2' , I , IC_2 , IC_2' , and IC_2C_2' . By suppressing I , IC_2 , C_2' , and C_2C_2' , the remaining four elements E , C_2 , IC_2' ($=\sigma_v$) and $C_2\sigma_v$ constitute the group C_{2v} .

There are a few points which should be made clear in this connection. First, in the abstract mathematical sense, these groups are not all distinct. For example, the group C_2 contains the elements C_2 and E , while the group I contains the elements I and E , and they both have the same multiplication table. On the other hand, there is some degree of distinction which should be made between the $E = C_2^2$ (which we might call the identity in rotation space) and the $E = I^2$ (which we might call the identity in inversion space). Strictly speaking, the identity element of the groups such as $C_n \times I$ is the product of these two separate identity elements.

We shall now investigate, term by term, the symmetry possessed by the tesseral harmonics tabulated above. The overall symmetry of a given potential will then be at least that of those symmetry elements common to each term of the potential expansion. Conversely, if the symmetry is known from other considerations, we shall know what terms must be included to describe such a condition.

C_{∞} : For this element to be present, the potential must be independent of ϕ , a condition which is

met only by terms with $M=0$ for all N .

C_n : For invariance here, $V(\phi) = V(\phi + 2\pi/n)$. Since the terms involve $\sin M\phi$ or $\cos M\phi$, it is clear that $n=M$, $M/2$, $M/3$, for integral values only, etc. We shall specify $n=M$, since the other cases are covered by the powers of C_n which are also present in any group.

C_2' : A point with the coordinates (r, θ, ϕ) under the transformation C_2' becomes $(r, \pi - \theta, \pi - \phi)$. Since Θ_N^M is a polynomial of degree $N-M$ in $\cos \theta$ and degree M in $\sin \theta$, and $\cos(\pi - \theta) = -\cos \theta$, $\sin(\pi - \theta) = \sin \theta$, then Θ_N^M changes as $(-1)^{N-M}$. However, $\sin M(\pi - \phi) = \sin M\pi \cos M\phi - \cos M\pi \sin M\phi = (-1)^{M+1} \sin M\phi$ and $\cos M(\pi - \phi) = \cos M\pi \cos M\phi + \sin M\pi \sin M\phi = (-1)^M \cos M\phi$. Hence C terms have this symmetry for N even and S terms have it for N odd.

I : Under this operation, the point (x, y, z) becomes $(-x, -y, -z)$, and it is clear from inspection of table 2 that terms with even N have this symmetry, while terms with odd N do not.

σ_h : This operation results, in rectangular coordinates, in the substitution of $-z$ for z , and, in spherical coordinates, in the substitution of $\pi - \theta$ for θ . This is the same as the θ substitution in C_2' and supplies a factor $(-1)^{N-M}$. Hence, N and M must both be even or both odd for this element.

σ_v : A vertical plane of symmetry exists at $\phi = \beta$ if $V(r, \theta, \beta + \alpha) = V(r, \theta, \beta - \alpha)$ for all values of α . For S terms, we have $\sin M(\beta + \alpha) = \sin M\beta \cos M\alpha + \cos M\beta \sin M\alpha$ and $\sin M(\beta - \alpha) = \sin M\beta \cos M\alpha - \cos M\beta \sin M\alpha$. This yields the requirement that $\cos M\beta \sin M\alpha = -\cos M\beta \sin M\alpha = 0$. For this to be independent of α , $\cos M\beta = 0$, $M\beta = \pi/2, 3\pi/2, 5\pi/2, \dots$ and planes of symmetry exist at $\beta = \pi/2M, 3\pi/2M, \dots, (2M-1)\pi/2M$. Similarly for C terms, we obtain the requirement that $\sin M\beta \sin M\alpha = 0$, $M\beta = 0, \pi, 2\pi, 3\pi, \dots$ and planes of symmetry exist at $\beta = 0, \pi/M, 2\pi/M, (M-1)\pi/M$. We shall find it convenient to distinguish these sets of planes, describing those arising from C terms as σ_c and those from S terms as σ_s . Note that the set σ_c for a given even M includes both σ_c and σ_s for $M/2$, and that any vertical plane is a plane of symmetry if $M=0$. Let us emphasize that the $x-z$ plane ($\beta=0$) is a plane of symmetry for all C terms, while the $y-z$ plane ($\beta = \pi/2$) is a plane of symmetry for C terms if M is even, for S terms if M is odd.

S_4 : This operation transforms (r, θ, ϕ) into $(r, \pi - \theta, \phi + (\pi/2))$ and Θ_N^M transforms as $(-1)^{N-M}$. S terms involve $\sin M(\phi + (\pi/2)) = \sin M\phi \cos M\pi/2 + \cos M\phi \sin M\pi/2$. M odd yields a $\cos M\phi$ term which we reject, and M even yields $(-1)^{M/2} \sin M\phi$. Thus S terms transform as $(-1)^{N-1/2M}$ for M even. For C terms, the θ contribution is the same as for S terms and $\cos M(\phi + (\pi/2)) = \cos M\phi \cos M\pi/2 - \sin M\phi \sin M\pi/2$. As for S terms, we reject M odd and in ϕ obtain $(-1)^{M/2} \cos M\phi$ for M even. C terms then transform as $(-1)^{N-1/2M}$,

just as S terms. The overall requirement for S_4 is then that both M and $N - \frac{1}{2}M$ must be even.

S_4 : Since this element is defined as C_2I , it is necessary and sufficient that both C_2 and I be present for this element to be present, and it need not be considered separately.

Based on the above analysis and discussion, the symmetry elements present in each of the tesseral harmonics of table 2 have been given in table 4. An analysis of table 4 will indicate the potential coefficients (2.7) which must be included to represent a potential appropriate to the groups of table 3, and the results of this analysis are given in table 5. Two points are worthy of note in the use of table 5. In each case where, for a given M and N both C and S are shown, the first S appearing is shown in parentheses. This represents the fact that, though both are consistent with the required symmetry, the S term may be suppressed by a suitable rotation about the z -axis. This may be done, of course, for any one C, S pair, but only for one. All other indicated combinations of C and S must then be allowed, and this one S omitted. Second, with the potential terms considered, it will be noted that the groups D_6 and C_{6h} cannot be distinguished from the group D_{6h} of higher symmetry, nor can C_6 be distinguished from C_{6v} . Further analysis of the potential terms shows that the coefficients S_7^6 , allowed in D_6 , and S_8^6 in C_{6h} , neither of which is allowed in D_{6h} , are the lowest order terms separating these three groups. Similarly, S_7^6 in C_6 will separate it from C_{6v} , where this coefficient is forbidden. Finally, it will be noted that the groups $D_\infty, C_{\infty h},$ and C_∞ have been omitted from the table. These symmetries do not arise from the electrostatic fields considered here.

2.3. Preferred Coordinate Systems

In considering the expansion of the potential (2.6) and the calculation of the coefficients (2.7), it is clear that the coefficients (2.7) will depend upon the choice of the coordinate system used.

On the other hand, it is also clear that all expansions of the field (2.6) must be physically equivalent, and that a choice between two differing descriptions of the same field is merely a matter of choosing the simplest description. Following universal convention, the principal axis of symmetry in the preceding considerations has been taken to be the z -axis, and in the calculation of the coefficients (2.7) this will have to be determined by an inspection of the given configuration of charges, as will also the location of any y -axis (C_2). Any other choice of axis will, in general, yield a more complicated expansion, and the symmetry will be *apparently* (but not *actually*) lower.

It would be well to discuss further the point mentioned in the preceding subsection concerning the planes of symmetry. For a given N and M , both C terms and S terms describe planes of symmetry, differing merely in their orientation with respect to the vertical coordinate planes. Clearly, any linear combination of these two terms also represents a set of planes of symmetry, at some intermediate angle. If only one N and M (other than $M=0$) is present, the linear combination will represent no more information than will either one alone. The general preference arises from the fact that the actual calculation of matrix elements in section 5 will be done through the Y terms (2.8) rather than the C and S terms. It will be seen from (2.9) that the suppression of S terms (as far as possible) will yield real coefficients for the matrix elements. A conventional preference for C terms then implies the $x-z$ plane as the preferred vertical plane of symmetry, and the y -axis for the operation C_2' , since IC_2' is then a reflection in this preferred plane. Sometimes (e.g., D_{3h}), we shall admit S terms and keep C_2' as the y -axis rather than adopt some other axis in the plane as C_2' .

Once an expansion has been obtained in a given coordinate system, that in any other coordinate system may in principle be obtained from the fact that each of the terms (2.5) or (2.8) in the new coordinate system will be expressible as a sum of

TABLE 4. Symmetry elements in tesseral harmonics

$M=$	0	1	2	3	4	5	6	
$N=1$ C	$C_\infty\sigma_z$	$\sigma_\lambda\sigma_z$						
	$C_\infty IC_2\sigma_z S_1\sigma_\lambda$	$\sigma_z IC_2$	$C_2 IC_2\sigma_\lambda\sigma_z$					
	$C_\infty\sigma_z$	$\sigma_\lambda\sigma_z$	$C_2\sigma_z S_1$	$C_2\sigma_\lambda\sigma_z$				
					$C_2 IC_2\sigma_z S_1$	$C_2 IC_2\sigma_\lambda\sigma_z S_1$		
					$C_2\sigma_z S_1$	$C_2\sigma_z$	$C_2\sigma_\lambda\sigma_z$	
		$C_\infty IC_2\sigma_z S_1\sigma_\lambda$	$\sigma_z IC_2$	$C_2 IC_2\sigma_\lambda\sigma_z$	$C_2 IC_2\sigma_z S_1$	$C_2 IC_2\sigma_\lambda\sigma_z S_1$	$C_2 IC_2\sigma_z$	$C_2 IC_2\sigma_\lambda\sigma_z S_1$
S		$\sigma_\lambda\sigma_z C_2$						
		$\sigma_z I$	$C_2 I\sigma_\lambda\sigma_z$					
		$\sigma_\lambda\sigma_z C_2$	$C_2 C_2\sigma_z S_1$	$C_2 C_2\sigma_\lambda\sigma_z$				
					$C_2 I\sigma_z S_1$	$C_2 I\sigma_\lambda\sigma_z S_1$		
		$\sigma_z I$	$C_2 I\sigma_\lambda\sigma_z$	$C_2 C_2\sigma_z S_1$	$C_2 C_2\sigma_\lambda\sigma_z$	$C_2 C_2\sigma_z S_1$	$C_2 C_2\sigma_\lambda\sigma_z$	
		$\sigma_\lambda\sigma_z C_2$	$C_2 I\sigma_\lambda\sigma_z$	$C_2 I\sigma_z S_1$	$C_2 I\sigma_\lambda\sigma_z S_1$	$C_2 I\sigma_z$		$C_2 I\sigma_\lambda\sigma_z S_1$

TABLE 6a. Transformation of tesseral harmonics

O'	O	C_1	C_1	S_1
C_1		0	1	0
C_1		0	0	1
S_1		1	0	0

TABLE 6b. Transformation of tesseral harmonics

O'	O	C_2	C_2	C_2	S_2	S_2
C_2		$-\frac{1}{2}$	0	$\frac{\sqrt{3}}{2}$	0	0
C_2		0	0	0	0	1
C_2		$-\frac{\sqrt{3}}{2}$	0	$-\frac{1}{2}$	0	0
S_2		0	1	0	0	0
S_2		0	0	0	1	0

TABLE 6d. Transformation of tesseral harmonics

O'	O	C_3	C_3	C_3	C_3	C_3	S_3	S_3	S_3	S_3
C_3		$\frac{3}{8}$	0	$-\frac{\sqrt{5}}{4}$	0	$\frac{\sqrt{35}}{8}$	0	0	0	0
C_3		0	0	0	0	0	$-\frac{\sqrt{2}}{4}$	0	$\frac{\sqrt{14}}{4}$	0
C_3		$\frac{\sqrt{5}}{4}$	0	$-\frac{1}{2}$	0	$-\frac{\sqrt{7}}{4}$	0	0	0	0
C_3		0	0	0	0	0	$-\frac{\sqrt{14}}{4}$	0	$-\frac{\sqrt{2}}{4}$	0
C_3		$\frac{\sqrt{35}}{8}$	0	$\frac{\sqrt{7}}{4}$	0	$\frac{1}{8}$	0	0	0	0
S_3		0	$-\frac{3}{4}$	0	$\frac{\sqrt{7}}{4}$	0	0	0	0	0
S_3		0	0	0	0	0	$-\frac{\sqrt{2}}{4}$	0	$\frac{\sqrt{14}}{4}$	0
S_3		0	$-\frac{\sqrt{7}}{4}$	0	$-\frac{3}{4}$	0	0	0	0	0
S_3		0	0	0	0	0	$-\frac{\sqrt{14}}{4}$	0	$-\frac{\sqrt{2}}{4}$	0

the $2N+1$ terms of the same value of N but of different M in the old coordinate system. While general formulas have been derived for such relations (Wigner, 1931), a detailed study has been carried out for a second coordinate system O' obtained by a rotation of O by -120° about the $(i+j+k)$ axis. This is equivalent to a cyclic permutation $x \rightarrow y \rightarrow z \rightarrow x$ of the variables; i.e., the functions $C_N^M(y, z, x)$ and $S_N^M(y, z, x)$ were evaluated in terms of $C_N^M(x, y, z)$ and $S_N^M(x, y, z)$. The results of this evaluation are presented as the matrices of table 6. It should be noted that these matrices are orthogonal.

TABLE 6c. Transformation of tesseral harmonics

O'	O	C_3	C_3	C_3	C_3	S_3	S_3	S_3
C_3		0	$-\frac{\sqrt{6}}{4}$	0	$\frac{\sqrt{10}}{4}$	0	0	0
C_3		0	0	0	0	$-\frac{1}{4}$	0	$\frac{\sqrt{15}}{4}$
C_3		0	$-\frac{\sqrt{10}}{4}$	0	$-\frac{\sqrt{6}}{4}$	0	0	0
C_3		0	0	0	0	$-\frac{\sqrt{15}}{4}$	0	$-\frac{1}{4}$
S_3		$-\frac{\sqrt{6}}{4}$	0	$\frac{\sqrt{10}}{4}$	0	0	0	0
S_3		0	0	0	0	0	1	0
S_3		$-\frac{\sqrt{10}}{4}$	0	$-\frac{\sqrt{6}}{4}$	0	0	0	0

TABLE 6c. Transformation of lesser harmonics

O' \ O	C_1	C_2	C_3	C_4	C_5	C_6	S_1	S_2	S_3	S_4	S_5
C_1	0	$\frac{\sqrt{15}}{8}$	0	$-\frac{\sqrt{70}}{16}$	0	$\frac{3\sqrt{14}}{16}$	0	0	0	0	0
C_2	0	0	0	0	0	0	$\frac{1}{8}$	0	$-\frac{\sqrt{42}}{16}$	0	$\frac{\sqrt{210}}{16}$
C_3	0	$\frac{\sqrt{7}}{4}$	0	$-\frac{\sqrt{6}}{8}$	0	$-\frac{\sqrt{30}}{8}$	0	0	0	0	0
C_4	0	0	0	0	0	0	$\frac{\sqrt{42}}{16}$	0	$-\frac{13}{16}$	0	$-\frac{3\sqrt{5}}{16}$
C_5	0	$\frac{\sqrt{21}}{8}$	0	$\frac{9\sqrt{2}}{16}$	0	$\frac{\sqrt{10}}{16}$	0	0	0	0	0
C_6	0	0	0	0	0	0	$\frac{\sqrt{210}}{16}$	0	$\frac{3\sqrt{5}}{16}$	0	$\frac{1}{16}$
S_1	$\frac{\sqrt{15}}{8}$	0	$-\frac{\sqrt{7}}{4}$	0	$\frac{\sqrt{21}}{8}$	0	0	0	0	0	0
S_2	0	0	0	0	0	0	0	$-\frac{1}{2}$	0	$\frac{\sqrt{3}}{2}$	0
S_3	$\frac{\sqrt{70}}{16}$	0	$-\frac{\sqrt{6}}{8}$	0	$-\frac{9\sqrt{2}}{16}$	0	0	0	0	0	0
S_4	0	0	0	0	0	0	0	$-\frac{\sqrt{3}}{2}$	0	$-\frac{1}{2}$	0
S_5	$\frac{3\sqrt{14}}{16}$	0	$\frac{\sqrt{30}}{8}$	0	$\frac{\sqrt{10}}{16}$	0	0	0	0	0	0

2.4. Cubic Point Groups

In addition to the 27 noncubic point groups of table 3, there are 5 point groups known as the cubic groups. While these are not known to arise in natural rare earth crystals, they are important in the consideration of crystals of other types; e.g., the iron group, and some data on rare earth ions have been obtained by the inclusion of the ion in a foreign lattice of cubic symmetry (Low, 1958). We shall therefore complete the discussion by considering these groups.

The five groups are assigned the symbols O , $O_h=(O \times I)$, T , $T_h=(T \times I)$, and T_d , but only four distinct abstract groups are involved since T_d and O are isomorphic. The group O has 24 elements and represents the purely rotational symmetry of the cube (or octahedron), while O_h , 48 elements, represents the full symmetry of the cube. T is of order 12 and represents the pure rotational symmetry of the tetrahedron, while T_d , a subgroup of O_h with 24 elements, represents the full symmetry of the regular tetrahedron. T_h is also of order 24.

These groups represent a higher degree of symmetry than our previous considerations have suggested. In particular, O includes fourfold axes along x , y , and z , threefold axes along the principal diagonal of each octant, and twofold axes along

the diagonals of the coordinate planes. T has 4 threefold axes, of which at most one can be along a coordinate axis, but has also three mutually perpendicular twofold axes. Rotations of these groups will be generally designated C_n for rotations $\frac{2\pi}{n}$, and they may be further distinguished by primes. However, the distinctive conventions of section 2.2 which are appropriate for D_{nh} and its subgroups, cannot be adhered to for these high symmetries. These symmetries cannot arise from considering the symmetry elements common to a sum of terms as in the previous cases, but depend upon certain fixed ratios in the coefficients. In other words, these may be viewed as higher symmetries arising from "accidental" values of the coefficients in a case of lower symmetry.

As a particular example, let us consider the group D_4 , representing the rotational symmetry of a rectangular prism, with potential coefficients (table 5) C_2^z , C_4^z , C_2^x , C_2^y , S_4^z , and C_2^x . The x and y axes are equivalent, but the long dimension, the z axis, will generally be different. If the prism actually becomes a cube, the x , y , and z axes will all be equivalent. Hence we expect that any combination of these terms which is invariant under a substitution $x \rightarrow y \rightarrow z \rightarrow x$ will represent a higher symmetry, in this case O . This is just the substitution considered in table 6. We see, for

TABLE 6f. Transformation of tesseral harmonics

O'	O	C_0^0	C_2^0	C_4^0	C_6^0	C_8^0	C_{10}^0	C_{12}^0	C_{14}^0	C_{16}^0	S_2^0	S_4^0	S_6^0	S_8^0	S_{10}^0	S_{12}^0	S_{14}^0	S_{16}^0
C_0^0		$-\frac{5}{16}$	0	$\frac{\sqrt{210}}{32}$	0	$-\frac{3\sqrt{7}}{16}$	0	$\frac{\sqrt{462}}{32}$	0	0	0	0	0	0	0	0	0	0
C_2^0		0	0	0	0	0	0	0	0	$\frac{\sqrt{10}}{16}$	0	$-\frac{\sqrt{3}}{4}$	0	$\frac{3\sqrt{22}}{16}$	0	0	0	0
C_4^0		$-\frac{\sqrt{210}}{32}$	0	$\frac{17}{32}$	0	$-\frac{\sqrt{30}}{32}$	0	$-\frac{3\sqrt{55}}{32}$	0	0	0	0	0	0	0	0	0	0
C_6^0		0	0	0	0	0	0	0	0	$\frac{9}{16}$	0	$-\frac{\sqrt{30}}{8}$	0	$-\frac{\sqrt{55}}{16}$	0	0	0	0
C_8^0		$-\frac{3\sqrt{7}}{16}$	0	$\frac{\sqrt{30}}{32}$	0	$\frac{13}{16}$	0	$\frac{\sqrt{66}}{32}$	0	0	0	0	0	0	0	0	0	0
C_{10}^0		0	0	0	0	0	0	0	0	$\frac{\sqrt{165}}{16}$	0	$\frac{\sqrt{22}}{8}$	0	$\frac{\sqrt{3}}{16}$	0	0	0	0
C_{12}^0		$-\frac{\sqrt{462}}{32}$	0	$-\frac{3\sqrt{55}}{32}$	0	$-\frac{\sqrt{66}}{32}$	0	$-\frac{1}{32}$	0	0	0	0	0	0	0	0	0	0
S_2^0		0	$\frac{5}{8}$	0	$-\frac{3\sqrt{10}}{16}$	0	$\frac{\sqrt{66}}{16}$	0	0	0	0	0	0	0	0	0	0	0
S_4^0		0	0	0	0	0	0	0	$\frac{\sqrt{10}}{16}$	0	$-\frac{9}{16}$	0	$\frac{\sqrt{165}}{16}$	0	0	0	0	0
S_6^0		0	$\frac{3\sqrt{10}}{16}$	0	$-\frac{1}{16}$	0	$-\frac{\sqrt{165}}{16}$	0	0	0	0	0	0	0	0	0	0	0
S_8^0		0	0	0	0	0	0	0	$\frac{\sqrt{3}}{4}$	0	$-\frac{\sqrt{30}}{8}$	0	$-\frac{\sqrt{22}}{8}$	0	0	0	0	0
S_{10}^0		0	$\frac{\sqrt{66}}{16}$	0	$\frac{\sqrt{165}}{16}$	0	$\frac{5}{16}$	0	0	0	0	0	0	0	0	0	0	0
S_{12}^0		0	0	0	0	0	0	0	$\frac{3\sqrt{22}}{16}$	0	$\frac{\sqrt{55}}{16}$	0	$\frac{\sqrt{3}}{16}$	0	0	0	0	0

example, that C_4^0 becomes a combination of C_2^0 and C_0^0 . Since C_4^0 is not allowed in D_4 , we suppress the second degree terms and proceed to those of the fourth degree. Although both C_4^0 and C_2^0 yield C_4^0 , if the coefficients are in the ratio $C_4^0:C_2^0 = \sqrt{7}:\sqrt{5}$, the C_4^0 will cancel and the combination $C_4^0 + \sqrt{\frac{5}{7}} C_2^0$ will be invariant under the operation of table 6. Similarly, if $C_6^0:C_4^0 = -1:\sqrt{7}$, C_6^0 and C_4^0 arising from the transformation will vanish and $C_6^0 - \sqrt{7} C_4^0$ will also be invariant. S_4^0 will admit S_2^0 , forbidden in D_4 , so S_4^0 must also vanish. We conclude then that under these conditions, we have but two coefficients, C_2^0 and C_0^0 left to fix the strength of the field, and the symmetry is O rather than D_4 . In fact, all of the terms allowed also contain I as a symmetry element, and the symmetry is even higher, i.e., O_h . The distinction between O and O_h will arise from potential terms of higher order.

On the other hand, the cube may be considered as standing on a corner with a threefold axis along

the z axis, arising as a special case of D_3 symmetry. The allowed coefficients will be generally C_0^0 ; S_2^0 ; C_4^0 ; C_6^0 ; S_4^0 ; C_8^0 ; C_{10}^0 , and C_{12}^0 . Since we do not have data corresponding to table 6 to apply our previous method, we shall set up a charge configuration having the desired symmetry and orientation, evaluate the coefficients (2.7) and determine the necessary relations among them. This has been done in detail as an example in section 6—we report merely the result at this point. The coefficients C_2^0 , S_4^0 , and S_6^0 must vanish, while the ratios among the nonvanishing coefficients are $C_4^0:C_0^0 = -7.2\sqrt{35}$ and $C_6^0:C_2^0:C_0^0 = 24.2\sqrt{105}:\sqrt{462}$. In this case as well, C_4^0 and C_0^0 are the only adjustable parameters to fix the strength of the field.

The group T may likewise be considered either as a special case of D_2 , where the x , y , and z axes all become equivalent, or as a special case of C_3 with the principal threefold axis of the figure vertical. In the former case, the allowed coefficients are: C_0^0 ; C_2^0 ; S_2^0 ; C_4^0 ; C_6^0 ; C_8^0 ; S_4^0 ; S_6^0 ; and C_{10}^0 , C_{12}^0 , and C_{14}^0 . A consideration of these terms and table 6 shows that the following conditions

must be met:

$$\begin{aligned} C_3^0 = C_3^2 = 0 \quad S_3^2 \text{ is arbitrary} \quad C_4^2 = 0 \quad S_4^2 = S_4^6 = 0 \\ C_4^0 : C_4^4 = \sqrt{7} \cdot \sqrt{5} \quad C_4^2 : C_4^6 = -1 : \sqrt{7} \\ \text{and } C_3^2 : C_3^4 = -11 : \sqrt{55}. \end{aligned}$$

The constants S_3^2 , C_4^0 , C_4^2 and C_3^2 can be considered the variable coefficients describing the strength of the field. The symmetry becomes T_h if the only one of these terms lacking the element I , S_3^2 , is suppressed. If we start from the case of D_{2d} symmetry instead of merely D_2 , we obtain T_d symmetry but must suppress the coefficient C_3^2 , leaving S_3^2 , C_4^0 , and C_4^2 .

In evaluating the coefficients for the threefold axis vertical, we have again had recourse to the establishment of a specific configuration, namely, that of the regular tetrahedron itself. This will yield terms for T_d rather than T or T_h . Configurations for these latter symmetries are considerably more complicated, and have not been evaluated for this orientation. The nonvanishing coefficients must meet the following conditions:

$$\begin{aligned} C_3^0 : C_3^2 = \sqrt{5} : 2 \quad C_4^0 : C_4^2 = -\sqrt{35} : 5 \quad \text{and} \\ C_6^0 : C_6^2 : C_6^4 = 4\sqrt{2310} : 77\sqrt{5} : 35\sqrt{22}. \end{aligned}$$

3. Groups, Characters, and Representations

3.1. Representations and Quantum Mechanics

Let us consider a Hamiltonian H , and ψ_i ($i=1, 2, \dots, n$) an n -fold degenerate solution of the Schrodinger equation $H\psi_i = E\psi_i$. Let us further consider a group of transformations (generally rotations or the inversion) which may be applied to this equation. If an element of the group is P , with inverse P^{-1} , then $PH\psi_i = PE\psi_i$, which may be rewritten as

$$(PHP^{-1})(P\psi_i) = E(P\psi_i). \quad (3.1)$$

This we interpret as yielding a transformed Hamiltonian (PHP^{-1}) and a transformed solution $P\psi_i$ of the Schrodinger equation. Let us now assume that the Hamiltonian is invariant under the group of operations; i.e., that $PHP^{-1} = H$ or $PH = HP$, and that Q is another operation of the group. Then (3.1) becomes $H(P\psi_i) = E(P\psi_i)$, and $P\psi_i$ must be a solution of the original wave equation with the same energy; i.e., it must be a linear combination of the original wave functions

$$P\psi_i = \sum_j P_{ij} \psi_j. \quad (3.2)$$

Similarly $Q\psi_i = \sum_j Q_{ij} \psi_j$, and $(QP)\psi_i = \sum_j (QP)_{ij} \psi_j$.

Thus the effect of each operation of the group may be expressed as a $n \times n$ unitary matrix. The importance of the concept becomes most evident when we consider the product QP as the operation P followed by the operation Q :

$$\begin{aligned} Q(P\psi_i) &= Q\left(\sum_j P_{ij} \psi_j\right) = \sum_k Q_{ik} \sum_j P_{kj} \psi_j \\ &= \sum_j \sum_k Q_{ik} P_{kj} \psi_j. \end{aligned} \quad (3.3)$$

Thus, the matrix for (QP) is the product of the matrix for P by that for Q . These unitary matrices are called a "representation of the group" because any relation between elements of the group is also a relation between the matrices corresponding to the group elements.

Clearly, the precise form of the matrices depends on the particular choice of original wave functions, and any other set, related to the original set by a unitary transformation may be selected and will yield a transformed representation of the group. If it is possible to find a transformation such that the group operations cause only m of the n wave functions to transform among themselves, and the remaining $n-m$ functions among themselves, the representation is said to be (fully) reducible, otherwise irreducible. Following Melvin (1956), we shall for brevity speak of an irreducible representation as a "rep". If the representation is a rep, the degeneracy is said to be essential, since the n wave functions may be transformed into each other purely by symmetry operations. If the representation can be reduced into two or more reps, the degeneracy is essential between wave functions belong to the same rep, but "accidental" when wave functions belonging to different reps are concerned, since wave functions belonging to different reps are not related to each other purely by symmetry operations. Accidental degeneracy is either due to a purely fortuitous consequence of the numerical parameters of the system under consideration, or to the presence of additional symmetry not previously considered (sometimes called "excess degeneracy"). An example of this latter will be considered in connection with Kramers' theorem (section 4.2), which states that all levels of an atomic system with an odd number of electrons will be at least twofold degenerate under the influence of external fields of purely electrostatic character. In this case, it will be seen that wave functions belonging to different reps of the rotation-reflection group may actually be related by symmetry with respect to time reversal.

3.2. Rotations in Three Dimensions

One of the transformations in which we shall be particularly interested is that of an arbitrary rotation of the coordinate axes in three dimensions, so we shall now examine this in some detail. Let us

consider a fixed (x, y, z) reference frame, and a movable frame (ξ, η, ζ) , the origin of the latter being fixed at the origin of the (x, y, z) frame. We assume initially that the $\xi, \eta,$ and ζ axes coincide with the $x, y,$ and z axes, respectively. Any rotation will move the (ξ, η, ζ) frame to some new orientation which is uniquely determined by the rotation performed. We are interested in the various ways of specifying the orientation of the (ξ, η, ζ) frame and its relation to the rotation involved.

It is clear that there will be three independent quantities to be specified, two to fix the direction of the axis of rotation and one to fix the angle of rotation about this axis. Consider a vector \mathbf{r} fixed in the (ξ, η, ζ) frame rotated through an angle ω about an axis fixed in (x, y, z) containing

$$D_1(\mathbf{u}, \omega) = \begin{pmatrix} \cos \omega + (1 - \cos \omega)\alpha^2 & (1 - \cos \omega)\alpha\beta - \gamma \sin \omega & (1 - \cos \omega)\alpha\gamma + \beta \sin \omega \\ (1 - \cos \omega)\alpha\beta + \gamma \sin \omega & \cos \omega + (1 - \cos \omega)\beta^2 & (1 - \cos \omega)\beta\gamma - \alpha \sin \omega \\ (1 - \cos \omega)\alpha\gamma - \beta \sin \omega & (1 - \cos \omega)\gamma\beta + \alpha \sin \omega & \cos \omega + (1 - \cos \omega)\gamma^2 \end{pmatrix} \quad (3.5)$$

and \mathbf{r}, \mathbf{r}' are the column vectors (ξ, η, ζ) or (x, y, z) , and (x', y', z') , respectively.

Since \mathbf{r} is fixed in the (ξ, η, ζ) frame, its components in that frame are the same before and after the rotation; they are the same as its coordinates (x, y, z) before the rotation in the fixed frame.

The direction of the (ξ, η, ζ) frame may be determined then by applying this relation to the vectors $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$ in turn.

The set of all three dimensional rotations constitutes a group which we shall designate as \mathfrak{R}_3 . The set of real matrices $D_1(\mathbf{u}, \omega)$ is orthogonal, hence unitary, and meets the requirements of section 3.1 for a representation of the group. (Note that D_1 is a particular element of D_1 .) It is important that the group of matrices D_1 be not confused with the group of abstract operations \mathfrak{R}_3 , but this requires a clear understanding of the difference between a group and any particular representation of the group. In elementary work, a three-dimensional rotation is invariably thought of as an operation on a vector. This is adequate in these elementary cases, since this leads, as we have seen, to a particularly simple faithful representation (a unique matrix for each operation) of the group. However, we have already mentioned in section 2.3 that an element of \mathfrak{R}_3 will induce not only the linear substitution D_1 among the three components of a vector, but also a linear substitution D_N among the $2N+1$ components of the tesseral harmonics of degree N (table 2). All these substitutions are equally well representations of \mathfrak{R}_3 , and they are faithful for $N > 0$. Geometrically, a vector may be resolved into its components, which are the tesseral harmonics of degree 1, which lie along the three coordinate axes, so that a linear substitution among the components of the tesseral harmonics of degree 1

the unit vector \mathbf{u} with direction cosines (α, β, γ) . Observe that \mathbf{r} may be resolved into a component parallel to \mathbf{u} , $(\mathbf{r} \cdot \mathbf{u})\mathbf{u}$, and a component perpendicular to \mathbf{u} , $(\mathbf{r} - (\mathbf{r} \cdot \mathbf{u})\mathbf{u})$, and that the vector $(\mathbf{u} \times \mathbf{r})$ is perpendicular to both and has the same magnitude $r \sin(\mathbf{u}, \mathbf{r})$ as the latter.

The component of \mathbf{r} along \mathbf{u} , $(\mathbf{r} \cdot \mathbf{u})\mathbf{u}$ will be unchanged, while the perpendicular component $\mathbf{r} - (\mathbf{r} \cdot \mathbf{u})\mathbf{u}$ will be rotated through the angle ω into $(\mathbf{r} - (\mathbf{r} \cdot \mathbf{u})\mathbf{u}) \cos \omega + (\mathbf{u} \times \mathbf{r}) \sin \omega$. Thus,

$$\mathbf{r}' = \mathbf{r} \cos \omega + (1 - \cos \omega)(\mathbf{r} \cdot \mathbf{u})\mathbf{u} + (\mathbf{u} \times \mathbf{r}) \sin \omega. \quad (3.4)$$

This may be written as the matrix equation $\mathbf{r}' = D_1(\mathbf{u}, \omega)\mathbf{r}$, where

necessarily implies a corresponding linear substitution among the components of a vector, which we interpret as a rotation of the vector. Likewise, any linear combination of the $2N+1$ tesseral harmonics of degree N (or any quantity having similar transformation properties) may be viewed as a vector in a space of $2N+1$ dimensions, with the $2N+1$ tesseral harmonics serving as the unit vectors. An element of \mathfrak{R}_3 may also be represented by a rotation matrix in this $2N+1$ dimensional space. Our choice of properly normalized tesseral harmonics ensures that the resulting matrices are orthogonal. In terms of this concept, the matrices of table 6 are the matrices D_N for $N=1, 2, 3, 4, 5, 6$ and

$$\mathbf{u} = \frac{1}{\sqrt{3}}(\mathbf{i} + \mathbf{j} + \mathbf{k}), \quad \omega = \frac{2\pi}{3}.$$

A general rotation may also be described by a point along the direction of \mathbf{u} and at a distance ω from the origin. All rotations are thus represented as points within or on the surface of a sphere of radius π . More precisely, the points represent the results of the rotation, while the details of the rotation are fixed by prescribing a path from the origin (representing the original orientation) to this particular end point. It is important to notice that all points of the closed sphere represent distinct orientations, save those on the surface where diametrically opposed pairs of points represent the same orientation, being attained by a rotation π about oppositely directed axes. Consequently, there are two essentially different types of paths by which one can go from the origin to another point within the sphere. Thus, if it is desired to reach the point $(0, \frac{\pi}{2}, 0)$ in the (\mathbf{u}, ω) space, this could be accomplished by a single rotation $\frac{\pi}{2}$ about

the y -axis, and the resulting path is a straight line from $(0, 0, 0)$ to $(0, \frac{\pi}{2}, 0)$. On the other hand, a rotation by π about the $-y$ axis will be represented by a straight line from $(0, 0, 0)$ to $(0, -\pi, 0)$. Since the latter point is equivalent to $(0, \pi, 0)$, a further rotation $\frac{\pi}{2}$ will result in a path from $(0, \pi, 0)$ to $(0, \frac{\pi}{2}, 0)$. We have thus reached $(0, \frac{\pi}{2}, 0)$ by two different paths. These are essentially different since there is no continuous deformation of the paths, keeping the end points fixed, which will cause them to coincide. Paths which can thus be made to coincide are said to be of the same homotopy class, otherwise they are of a different homotopy class. There are only two homotopy classes for rotation in three dimensions. We shall call them P_0 (for paths involving no points on the surface of the sphere, or an even number of pairs of such points) and P_1 (paths involving an odd number of pairs of points on the surface of the sphere).

It is frequently more convenient to describe directly the orientation of the (ξ, η, ζ) frame than to describe the rotation as above; i.e., we give directly the angles between the two frames. The general technique is due to Euler, and the three angles which must be specified are known as Euler's angles. There are a great number of different choices which can be made, and indeed have been made by various authors, a freedom which complicates considerably comparisons between different authors. We shall follow here what seems to be the most frequent choice by recent American writers on the subject (Edmonds, 1957; Rose, 1957). The position of the ζ axis will be specified by θ , the angle between ζ and z , and by ϕ , the angle between x and the projection of ζ in the $x-y$ plane. These are just the usual spherical coordinates of the unit vector along the ζ axis in the (x, y, z) system. For the third angle, we shall

use ψ , the angle between η and the intersection of the $x-y$ and the $\xi-\eta$ planes. Experimental workers (Dieke and Crosswhite, 1956) have found convenient an alternate choice χ , the angle between ξ and the intersection of the $x-z$ and $\xi-\eta$ planes. These angles are related by the equation $\chi = \chi_0 + \psi$, where $\tan \chi_0 = \cos \theta \tan \phi$. If $\theta = 90^\circ$, then $\chi = \psi$.

A major advantage of Euler's angles is that any given orientation of the (ξ, η, ζ) frame can be obtained by three rotations in succession about the coordinate axes. For these rotations, the matrices (3.5) take on a particularly simple form, and their product, expressing the general form of (3.5) in terms of Euler's angles, may be computed directly. The three rotations are:

1. A rotation by ϕ about the z, ζ axes (which initially coincide). The $\xi-\eta$ plane remains in the $x-y$ plane, but the $\xi-\zeta$ and the $\eta-\zeta$ planes are rotated by the angle ϕ .

2. A rotation by θ about the η axis. This will bring the ζ axis into its final position, as specified by ϕ and θ . The $\xi-\eta$ plane will no longer be in the $x-y$ plane, but their intersection is the η axis.

3. A rotation by ψ about the ζ axis. This will bring the ξ and η axes to their final orientation.

In performing these rotations, it will be noted that each one is carried out in the (ξ, η, ζ) frame, which is generally in a different position each time as a result of the preceding rotation in the sequence. The same final configuration can be obtained by rotating about the (x, y, z) frame axes, provided, that the sequence of the angles is reversed. In other words, a rotation of the (ξ, η, ζ) frame first about the z -axis by the angle ψ , then about the y axis by θ , and finally about the x -axis again by the angle ϕ , will yield exactly the same final orientation of the (ξ, η, ζ) frame. This may be established either by purely geometric considerations, or by specific multiplication of the matrices involved. Using this latter sequence of angles, we obtain from (3.5)

if

$$\mathbf{u} = (0, 0, 1), \quad \omega = \phi \text{ or } \psi \quad \text{or} \quad \mathbf{u} = (0, 1, 0), \quad \omega = \theta$$

$$\mathbf{R}_\phi = \begin{pmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \mathbf{R}_\theta = \begin{pmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{pmatrix} \quad \mathbf{R}_\psi = \begin{pmatrix} \cos \psi & -\sin \psi & 0 \\ \sin \psi & \cos \psi & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

and their product $\mathbf{R}_\psi \mathbf{R}_\theta \mathbf{R}_\phi$ yields the matrices of the representation D_1 of \mathfrak{R}_3 , now in terms of the three Euler angles,

$$\mathbf{D}_1(\phi, \theta, \psi) = \begin{pmatrix} \cos \phi \cos \theta \cos \psi - \sin \phi \sin \psi & -\cos \phi \cos \theta \sin \psi - \sin \phi \cos \psi & \cos \phi \sin \theta \\ \sin \phi \cos \theta \cos \psi + \cos \phi \sin \psi & -\sin \phi \cos \theta \sin \psi + \cos \phi \cos \psi & \sin \phi \sin \theta \\ -\sin \theta \cos \psi & \sin \theta \sin \psi & \cos \theta \end{pmatrix} \quad (3.5a)$$

Since these matrices are orthogonal, the inverse of a given matrix is its transpose. In terms of u, ω , this amounts to reversing either the vector u or the sign of the angle of rotation. In terms of the Euler angles, the inverse amounts to reversing the sign of the angles and also their sequence. In general, we shall not indicate the variables for D_1 (or for D_N), since either set may be used.

A third extremely important description of a three-dimensional rotation arises from the Cayley-Klein parameters. Let us consider the general 2×2 matrix with complex elements, which we shall write in the form

$$\begin{pmatrix} ae^{i\alpha} & be^{i\beta} \\ ce^{i(\beta+\gamma)} & de^{i(\alpha+\delta)} \end{pmatrix}$$

where a, b, c , and d are real nonnegative numbers, and α, β, γ , and δ are real numbers. This matrix contains eight arbitrary parameters. If we require that it be a unitary matrix U , four conditions of restraint on these parameters are imposed, and there remain but four independent quantities. These four conditions are: $a=d$; $b=c$; $a^2+b^2=1$; and $2\alpha+\delta=2\beta+\gamma+\pi$. The further condition that $\det U=+1$ requires that $2\alpha+\delta=0$, leaving three independent quantities. We can then write

$$U = \begin{pmatrix} ae^{i\alpha} & be^{i\beta} \\ -be^{-i\beta} & ae^{-i\alpha} \end{pmatrix},$$

remembering that $a^2+b^2=1$. The independent parameters are then, essentially, a, α , and β .

Further, let

$$R = \begin{pmatrix} \xi & \xi - i\eta \\ \xi + i\eta & -\xi \end{pmatrix}$$

be a Hermitian matrix with a trace of zero, ξ, η , and ζ being arbitrary real numbers. A unitary transformation

$$URU^{-1} = R' \quad (3.6)$$

will leave invariant the Hermitian property, the trace, and the determinant, so R' can be written

$$R' = \begin{pmatrix} z & x - iy \\ x + iy & -z \end{pmatrix}.$$

Furthermore, $\det R' = -(x^2 + y^2 + z^2) = \det R = -(\xi^2 + \eta^2 + \zeta^2)$. If we interpret the sets of numbers (ξ, η, ζ) and (x, y, z) as the coordinates of a point in the movable (ξ, η, ζ) system and the fixed (x, y, z) system, respectively, we see that the transformation leaves unchanged the distance of the point from the origin, so that the transformation can be interpreted as a rotation. The matrix U can be explicitly calculated in terms of the Euler angles by finding the matrices U_* , U_θ , and U_ψ , so that

$U = U_* U_\theta U_\psi$. Thus,

$$U_* = \begin{pmatrix} e^{-i\psi/2} & 0 \\ 0 & e^{+i\psi/2} \end{pmatrix}$$

$$U_\theta = \begin{pmatrix} \cos \frac{\theta}{2} & -\sin \frac{\theta}{2} \\ \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix} \quad U_\psi = \begin{pmatrix} e^{-i\psi/2} & 0 \\ 0 & e^{+i\psi/2} \end{pmatrix}$$

$$U = \begin{pmatrix} e^{-\frac{i(\phi+\psi)}{2}} \cos \frac{\theta}{2} & -e^{-\frac{i(\phi-\psi)}{2}} \sin \frac{\theta}{2} \\ e^{\frac{i(\phi-\psi)}{2}} \sin \frac{\theta}{2} & e^{\frac{i(\phi+\psi)}{2}} \cos \frac{\theta}{2} \end{pmatrix} \quad (3.7)$$

Just as in the previous discussion distinguishing between \mathfrak{R}_3 and D_1 , we have here an abstract group of unitary unimodular transformations in a complex space of two dimensions, and a particular representation of the group through the matrices (3.7). The abstract group we shall designate as U_2 , the particular representation by the collection of U matrices (3.7) as $D_{1/2}$. The elements of the representation $D_{1/2}$ may be given either in terms of the Euler angles as in (3.7), or in terms of the unit vector u and the angle ω , though we shall not give the latter form here explicitly. Since it will not generally be necessary to indicate the variables, we shall generally write the matrices merely as $D_{1/2}$ (instead of U) to parallel our earlier distinction between D_1 and D_3 .

Let us consider the implication of this transformation in more detail. We first observe that the matrix R can be written in the form

$$R = \xi \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \eta \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} + \zeta \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

where the 2×2 matrices are the Pauli matrices corresponding to the x, y , and z -components, respectively, of the spin angular momentum operator. If we let (ξ, η, ζ) be a unit vector, the matrix R is the operator corresponding to the projection of the spin angular momentum in the direction given by the unit vector (ξ, η, ζ) . Let us consider for simplicity U for $\theta = \psi = 0$. $R' = URU^{-1}$ may similarly be written

$$R' = (\xi \cos \phi - \eta \sin \phi) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$+ (\eta \cos \phi + \xi \sin \phi) \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} + \zeta \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

This corresponds to our interpretation that the vector with components (ξ, η, ζ) , fixed in the (ξ, η, ζ) frame, is given a rotation about the z - ζ axis, positive in terms of the right hand screw

convention. R' is now the operator corresponding to the projection of spin in the new direction, still described in the movable system by (ξ, η, ζ) , but described in the fixed (x, y, z) system by $x = \xi \cos \phi - \eta \sin \phi$, $y = \eta \cos \phi + \xi \sin \phi$, $z = \zeta$. Note that the Pauli matrices retain their form, i.e., they are expressed in the fixed (x, y, z) frame. However, we can also write

$$R' = \xi \begin{pmatrix} 0 & e^{-i\phi} \\ e^{i\phi} & 0 \end{pmatrix} + \eta \begin{pmatrix} 0 & -ie^{-i\phi} \\ ie^{i\phi} & 0 \end{pmatrix} + \zeta \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

where ξ, η , and ζ retain their original values, and the Pauli spin matrices have been transformed to the form appropriate to the new ξ, η, ζ axes. Recalling that this is a rotation by $+\phi$, this should be equivalent to a rotation of the physical system, the fixed x, y, z axes, and the spin functions described therein, by $-\phi$. In other words, generally, a transformation of the spin operators σ , by U is equivalent to a transformation of the spin functions themselves by U^{-1} . More formally, consider $\phi = R\psi$, where ϕ and ψ are spin wave functions. This expression, under the transformation U , yields

$$U\phi = URU^{-1}U\psi \quad \text{or} \quad \phi' = R'\psi'$$

which represents merely the transformation of each side of the expression to a new frame of reference; in other words, essentially the same angular momentum measurement. From the standpoint of a simple rotation of R to measure angular momentum in the new direction fixed by R' as considered in the original frame of reference, we are interested in the quantity $R'\psi' = URU^{-1}\psi = X$ which is generally different from ϕ . This new measurement may be transformed by U^{-1} to give

$$U^{-1}X = U^{-1}[URU^{-1}]UU^{-1}\psi = R[U^{-1}\psi].$$

In other words, the rotation of R to R' , URU^{-1} gives the same result as the inverse transformation of the spin wave functions by U^{-1} . Since we are usually interested in transforming wave functions, we shall write $D_{\frac{1}{2}} = U^{-1}$ rather than $D_{\frac{1}{2}} = U$.

The presence of the half-angle functions in this transformation reflects the very special properties of the spin transformations. In particular, the rotation $\phi' = \phi + 2\pi$, $\theta' = \theta + 2\pi$, $\psi' = \psi + 2\pi$ clearly yields an orientation of axes identical to that designated by ϕ, θ, ψ . On the other hand, it may be seen that $D_{\frac{1}{2}}(\phi', \theta', \psi') = -D_{\frac{1}{2}}(\phi, \theta, \psi)$. Thus, for each change of orientation in ordinary three-dimensional space, there are two matrices corresponding to this in the two-dimensional spin space. For this reason, the set of matrices $D_{\frac{1}{2}}$ does not constitute a representation of the three-dimensional

rotation group in the stricted mathematical sense of the term. It has become customary to refer to $D_{\frac{1}{2}}$ as a "two-valued" or "ambiguous" representation, and the abstract group U_2 as the double rotation group. If a rotation is specified by giving the initial and final configuration of axes, the matrices $D_{\frac{1}{2}}$ are fixed apart from the sign. The sign may also be fixed if the detailed course of the rotation is also prescribed. Clearly, the matrix $D_{\frac{1}{2}}$ is a continuous function of the variables ϕ, θ, ψ and becomes the unit matrix for no rotation at all. In other words, if the initial and final axes are the same, we know only that the appropriate $D_{\frac{1}{2}}$ matrix is either the unit 2×2 matrix or its negative, while the detailed knowledge that no rotation at all has occurred (or one involving a path in the $(\mathbf{u}\omega)$ space of homotopy class P_0) enables us to fix upon the unit matrix as the appropriate one. Generally, if two axes are related by the Euler angles ϕ, θ, ψ we will select $D_{\frac{1}{2}}(\phi, \theta, \psi)$ if the transformation is of homotopy class P_0 , and $-D_{\frac{1}{2}}(\phi, \theta, \psi)$ if of class P_1 .

3.3. Group Theory and the Free Atom or Ion

The Hamiltonian for a free atom contains the following types of terms:

1. Kinetic energy terms $\frac{1}{2m} \mathbf{p}_i^2$
2. Central field potential terms $-\frac{Ze^2}{r_i} + V'(r_i)$
3. Electrostatic repulsion terms $\frac{e^2}{r_{ij}} - V'(r_{ij})$
4. Spin-orbit interaction terms $\zeta(r_i) \mathbf{l}_i \cdot \mathbf{s}_i$

In the above, the $V'(r_i)$ is selected so that the overall effect of the third term is minimized. In addition to the terms of the free ion Hamiltonian, we shall also be interested in the effects of the crystal lattice potential V discussed in the preceding section and of an external magnetic field \mathbf{B} . Consequently, we shall be interested in the operations which leave invariant these various terms in the Hamiltonian.

Let us first consider terms (1) and (2) only. In this case, each electron is considered to move independently of the detailed motion of the other electrons present, their mutual interaction appearing in an averaged form in the $V'(r_i)$ term. Term (1) gives the Laplacian operator, which may be viewed as a scalar product of two vector operators and is therefore invariant under an arbitrary rotation of coordinate axes about the force center. Since second derivatives are involved, it is also invariant under the inversion. Similarly, the second term involves only the scalar magnitude of \mathbf{r} which is invariant under the same two operations. The solutions of the Schrodinger equation for one electron for these two terms only are

characterized by the quantum numbers n , l , m_l , and m_s , and have a degeneracy equal to $2(2l+1)$, there being $2l+1$ values of m_l and 2 values of m_s , all belonging to the same energy and all related to each other by the particular choice of coordinate axis. These $2l+1$ functions are just the functions (2.8). Thus, a rotation of the coordinate axes will cause each of the $2l+1$ functions in the new set of axes to be expressed in terms of all $2l+1$ functions referred to the old axes, yielding a $2l+1$ dimensional representation of the rotation as in section 3.2. It is shown in the general theory of groups that for all nonnegative integral values of l these representations are in fact reps. General formulas for the matrix elements in terms of Euler's angles have been given by Wigner (1931). Specific examples of these matrices using a real basis (2.5) rather than the complex basis (2.8) have already been given in table 6. It is further to be noted that, for $l > 0$, the representation is faithful, i.e., there is a unique matrix corresponding to each rotation. We shall designate the rep by D_l , its elements by D_l .

For a single electron, as in hydrogen or in the alkali-type atoms or ions, term (3) in the Hamiltonian vanishes. So long as term (4) is neglected, the transformation of the $2l+1$ values of m_l and the two values of m_s are completely independent of each other, and may even be described with respect to different coordinate axes. The general transformation may be represented by a matrix of $2(2l+1)$ dimensions, the direct product of D_{2l} and D_1 . Upon the introduction of term (4), the independent transformation of spin and orbit no longer leaves the Hamiltonian invariant. The invariance of the scalar product of two operators (e.g., $\mathbf{l} \cdot \mathbf{s}$) is founded upon the assumption that each is referred to the same set of axes. The group of allowed transformations is now less general, since both spin and orbit must simultaneously undergo the same rotation. The $2(2l+1)$ dimensional matrix is still an element of a representation of the group of allowed transformations, but it is no longer irreducible, and it may be transformed by a new choice of wave functions into a diagonal matrix of sub-matrices of dimensionality $2(l+\frac{1}{2})+1$ and $2(l-\frac{1}{2})+1$. The correct choice of wave functions is precisely that dictated by the usual transformation from a m_l, m_s to j, m_j representation in one-electron spectra. The wave functions are determined by the coefficients known variously as Wigner, Clebsch-Gordon, or vector addition coefficients, and yield allowed values of the total angular momentum $j=l+\frac{1}{2}$ or $j=l-\frac{1}{2}$.

A similar argument is applicable to the case of two electrons. Without terms (3) and (4) in the Hamiltonian, all transformations are independent and most generally yield a matrix of $2^2(2l_1+1)(2l_2+1)$ dimensions. The introduction of term (3) now excludes those transformations affecting differently the space parts of the two one-electron wave functions, and the space portion of the

matrix can be reduced into sub-matrices D_L of dimensionality $2L+1$, where L ranges from l_1+l_2 to $|l_1-l_2|$. The restrictions of the Pauli principle require that the four-dimensional spin matrix be simultaneously reduced, yielding a three-dimensional submatrix D_1 and the unit 1 by 1 matrix D_0 . This reduction of the spin transformation matrix corresponds to the well-known existence of triplet and singlet states in two-electron spectra. For our immediate purposes, perhaps the most noteworthy point is the disappearance of the ambiguity in sign of the spin transformations. This is a general characteristic of atoms or ions with an even number of electrons. The reps of U_2 are infinite in number and are of both even and odd dimensionality. Those of even dimensionality include $D_{\frac{1}{2}}$ and others designated as $D_{3/2}, D_{5/2}$, etc., the dimensionalities being 2, 4, 6, . . . , respectively. These representations are all faithful, and hence all are double-valued representations of the three-dimensional rotation group. The odd dimensional reps are the $D_{0,1,2,3}$. . . previously discussed as reps of the three-dimensional rotation group, but these latter are not faithful. Aside from the identity rep D_0 , any element D_J , J integral, corresponds to $\pm D_J$, J half-integral. This is a point of greatest importance when considering the representations of the point groups, as we shall see shortly.

The introduction of term (4) into the Hamiltonian now requires the reduction of the $(2S+1) \cdot (2L+1)$ dimensional direct product into a diagonal array of sub-matrices D_J , where $J=L+S, L+S-1, \dots, |L-S|$. Here, the integral S yields integral J and the spin ambiguity remains suppressed. In certain atoms, the magnitude of the various terms in the Hamiltonian requires that term (4) be introduced before (3). The reduction of the various matrices must then be accomplished in a different order, corresponding to the $j-j$ coupling scheme rather than the $L-S$ scheme discussed above. For 3 or more electrons, the details can become increasingly complicated, but the general features remain the same. In particular, integral J 's without spin ambiguity appear for systems with an even number of electrons, while half-integral J 's corresponding to double-valued representations of pure rotations appear for systems with an odd number of electrons.

There is another symmetry which must be considered at this point, that of the inversion. Terms (1), (2), (3) in the Hamiltonian depend only on the length of a vector or vector operator and hence are invariant under the transformation from a right-handed to a left-handed coordinate system. The angular momentum operator, orbital or spin, has the transformation properties of a cross product of two vectors, i.e., an axial vector or pseudovector. Although the two vectors will change sign on inversion, their product will not and the vector operators (pseudovectors) \mathbf{l} and \mathbf{s} as well as the Hamiltonian term $\mathbf{l} \cdot \mathbf{s}$ are also invariant. It is

important to note at this point the distinction between a vector and its description. Let us consider the vector $\mathbf{r}=(x, y, z)$ and $\mathbf{t}=(u, v, w)$, with $\mathbf{r}\times\mathbf{t}=(yw-zv, zu-xw, xv-yu)$. If we now switch to a left-handed system, the vector \mathbf{r} remains the same, but it is now described by $(-x, -y, -z)$. Likewise, \mathbf{t} remains the same but its description becomes the negative of the original. On the other hand, the product $\mathbf{r}\times\mathbf{t}$ retains the same description, but is consequently a different vector since it is now in a left-handed system. The distinction between a quantity and its description is a fundamental one which must on occasion be made very carefully. If we consider a one-electron wave function $\psi(x, y, z, s)$, the description of this state in the inverted frame is $\psi(-x, -y, -z, s)$, the description of an angular momentum being invariant. ψ is a homogeneous polynomial of degree l in the variables x, y, z and the transformation under the inversion is that of $(-1)^l$ and that of a n -electron system $(-1)^{2l}$. Strictly speaking, of course, Σl_i will not have a definite numerical value when terms (3) and (4) in the Hamiltonian are introduced, but only down through term (2). Nevertheless, the introduction of (3) and (4), invariant under the inversion, will admit only wave functions of the same value of $(-1)^{2l}$ which may be used to describe states of the free atom or ion. In addition to J , the states may be designated even or odd, (g or u , respectively) depending on $(-1)^{2l}$.

3.4. Group Theory and the Perturbed Ion

It is the primary purpose of this paper to consider in detail the effects of terms in the Hamiltonian arising from the crystal lattice perturbation V and from the external magnetic field B . There are two techniques which may be used in the solution of the problem, both of which must be used in a complementary fashion for a complete discussion. One technique is to apply further the theory of groups, reducing the representation of the state of the free ion according to the reps of the appropriate electrostatic-magnetic field symmetry group. This technique will yield the number of levels, their description in terms of their behavior under the allowed symmetry operations, and selection and polarization rules for various types of transitions. The method will not yield purely numerical details, such as the relative positions of levels, their variation with electric and magnetic field intensities, or transition probabilities (save for the vanishing forbidden transitions). The other technique is to calculate to some degree of approximation the matrix elements of the perturbing potentials in terms of the free ion wave functions, interpreting the results in the framework of the usual perturbation theory. In the final analysis, this latter procedure will yield not only the numerical details not obtained from the former technique, but can be so interpreted as to yield all of the results that the former technique will give. Nevertheless, the group-theoretical ap-

proach is one of great generality and provides a very concise systematic framework for stating the results obtained. Furthermore, the principles of group theory (or essentially similar arguments) are used to determine what matrix elements of the perturbing potentials vanish if the second approach is used, and to calculate the magnitude of those which do not vanish. Hence, the two approaches are not really so divergent as might at first sight appear to be the case. In this work, the contributions to the problem which can be made with the use of group theory will be studied in detail before proceeding to the alternate considerations of calculations involving specific matrix elements. While it is not the intent of this paper to furnish a general treatise on group theory, and a general acquaintance with at least the basic definitions will be presumed, certain concepts will be examined in detail and applied as the necessity arises.

Let us consider the very simple group C_3 , consisting of the elements C_3, C_3^2 , and E . We shall describe the group in terms of the effect of its operations on the basis triangle of figure 2a, taking C_3 as a counterclockwise rotation of the triangle by 120° . In the present configuration, the operation C_3 will cause the vertex (1) to appear at the point previously occupied by (2), (2) at that occupied by (3), and also (3) \rightarrow (1). We may shorten this to $C_3: (1)\rightarrow(2)\rightarrow(3)$. We may also describe the operations in terms of their effect on the triangle in figure 2b, where we see that we also obtain $C_3: (1)\rightarrow(2)\rightarrow(3)$. Notice that figure 2b differs from figure 2a only in that C_3 has been applied to the basis triangle before we use it to describe the group. We could also have applied C_3^2 to figure 2a before describing the operation, and again would have obtained $C_3: (1)\rightarrow(2)\rightarrow(3)$. A study of the operation C_3^2 will show that, using any of these basis triangles, we obtain $C_3^2: (1)\rightarrow(3)\rightarrow(2)$ as its description. Thus, if we restrict ourselves to transformations within the group under discussion, there is no transformation which will cause C_3^2 and C_3 to have the same description in terms of their effect on the basis triangle. Let us now consider the group D_3 , where we have added to our original set of three elements three rotations of 180° about axes in the x - y plane, described in terms again of figure 2a as $C_2: (2)\rightarrow(3)$; $C_2': (1)\rightarrow(2)$; and $C_2'': (1)\rightarrow(3)$. Still restricting ourselves to transformations within the group, let us consider the effect of C_3 in terms of the basis of figure 2c, derived from figure 2a by the application of C_2 . Here we see that we obtain $C_3: (1)\rightarrow(3)\rightarrow(2)$, but this is the same as C_3^2




FIGURE 2

described in terms of figure 2a. We thus see that in our first case C_2 and C_3 were essentially different, in that none of our allowed transformations permit them to have the same description, while in the second case at least part of this essential difference has been lost. This difference is expressed by the notion of class. In the first case C_2 and C_3 are said to be in a different class, while in the second case they are said to be in the same class.

More generally, if we consider the general transformation, $P' = TPT^{-1}$ and limit ourselves to transformations T belonging to the group of P , we see that P' is also in the group of P , and is defined as being in the class of P . For each T of the group a P' is obtained, not necessarily distinct, and the complete set of P' constitutes the class of P . Clearly, if P and T commute, $P' = P$. In an Abelian group, where all elements commute, every element is in a class by itself. Conversely, if P' is distinct from P , then P and T must not commute. The separation of a group into classes may be accomplished either by geometric considerations such as outlined above, or by formal computation. The formal computation is aided by the following considerations. Let N be the set of group elements which commute with P . It may be shown that this set of elements is in fact a subgroup (possibly the entire group) called the normalizer of P . Any element of N as the transforming element yields a $P' = P$. Let K be a group element not in N (K does not exist if the normalizer is the whole group). Then K as the transforming element T yields a P' different from P , but any element in the complex KN yields the same P' as does K . Similarly, if L is a group element not in N or KN , it will yield still another P'' , but all elements in LN will yield the same P'' . Generally, if the group order is g and the order of the normalizer is n , the ratio g/n , an integer, is called the index of N , and the group may be split into g/n complexes, one of which is the subgroup N . All of the elements of the class of P may then be obtained by transforming P with an element from each of these g/n complexes. Hence the class of P has g/n distinct elements.

These considerations may be applied to the group D_3 discussed above as an example. The group order is 6, and the elements which commute with C_2 are E, C_2, C_3^2 . Hence the index of the normalizer is 2, and there are two elements in the class of C_2 , one of which is C_2 itself, the other of which may be obtained by transforming C_2 with any one of C_3, C_3^2 , or C_3^3 . To accomplish this latter we need the multiplication table for the group elements, usually summarized as $C_3^3 = C_3^2 = E, C_3 C_2 = C_2 C_3^2$. The other element in the class of C_2 is $C_2 C_3 C_2^{-1} = C_2 C_3 C_2 = C_2 C_3 C_3^2 = C_3^2$, as was geometrically shown above. C_2 commutes only with itself and E , so the order of its normalizer is 2, its index is 3, and there are three elements in the class of C_2 . These are C_2 and the products of C_2 with C_3 and C_3^2 .

If we have a set of g unitary matrices forming a representation of a group of order g , the set of

traces of these g matrices is said to be the character of the representation. The importance of the concept lies in the invariance of the trace under a unitary transformation, $\text{Tr } P' = \text{Tr } UPU^{-1} = \text{Tr } P$. We see immediately that if U is a matrix of the representation, then P' is an element of the class of P and all elements of the same class have the same trace. If U is a matrix reducing the representation into two or more reps, the character of the representation is the sum of the characters of the component reps. This latter fact will allow us to determine what reps may be present in a given representation without actually finding the particular U necessary to accomplish the reduction.

This latter procedure is precisely what we have outlined above in sketching the application of group-theoretical ideas to the description of states of free atoms or ions, and what we wish to examine in detail when considering the effect of the introduction of the crystal field V and the external magnetic field B . Before the introduction of the external fields, the Hamiltonian is invariant under arbitrary rotation (of the simple or double group) and inversion, and the wave functions are transformed by D_j and $(-1)^{2j}$, respectively. After the introduction of the external fields, the Hamiltonian is now invariant only with respect to subgroups of the original group of transformations. The original rep of the complete group will not in general be a rep of this smaller group but merely a representation, reducible into reps of the smaller group. This is the expression of the fact that the original $2J+1$ fold degenerate level will under the influence of these external fields split into several levels of lower degeneracy. The character of the original representation is the sum of the characters of the reps into which the state splits. We shall now outline the method whereby this determination of the component reps may be accomplished.

3.5. Group Characters and the Rep- D_j

Let us consider a finite group of g elements, with the elements separated into classes. There will, in general, be several reps for this group. Let us write the trace in the i -th rep and j -th class as χ_{ij} . For a given i, j will assume values from 1 to n , where n is the number of classes, and χ_{ij} will be a set of numbers, generally complex and not necessarily all distinct. Strictly speaking, the set of numbers χ_{ij} for all j and a given i constitutes the character, while the number χ_{ij} for a particular i and j is merely the trace of a matrix, and one component of the character. This may be emphasized by considering the character as a vector χ_i in a space of n dimensions, with the n components, χ_{ij} . This is a distinction which is usually either overlooked or ignored. We shall now introduce a weighted trace, ξ_{ij} defined as

$\sqrt{\frac{N_j}{g}} \chi_{ij}$, where N_j is the number of elements in the class j , and, correspondingly, a weighted character ξ_i . Fundamental to our further considera-

tions will be the orthogonality theorem for these weighted characters, which we shall state without proof, referring to Wigner (1931) for details. This theorem states the following:

1. The weighted character ξ_i for a given i is a normalized vector in the Hermitian sense in a space of n dimensions. Symbolically

$$\xi_i \cdot \xi_i = \sum_{j=1}^n \xi_i^* \xi_{ij} = 1. \quad (3.8)$$

2. The weighted characters belonging to different reps are orthogonal in the Hermitian sense.

$$\xi_i \cdot \xi_k = \sum_{j=1}^n \xi_i^* \xi_{kj} = \delta_{ik}. \quad (3.9)$$

3. The set of weighted characters ξ_i for all i forms a complete set in the space of n dimensions, in terms of which the weighted character of any representation may be expanded. If Ξ is such a

weighted character, with components $\Xi_j = \sqrt{\frac{N_j}{g}} X_j$,

$$\Xi = \sum_{i=1}^n a_i \xi_i \quad \text{and} \quad a_i = \xi_i \cdot \Xi. \quad (3.10)$$

Two important conclusions may be immediately drawn from this theorem. We observe that, since the weighted characters form a complete set in a space of n dimensions, there must be n of them. In other words, the index i as well as j must assume values from 1 to n , and the number of reps is equal to the number of classes of the group. The orthogonality theorem suggests that we consider these weighted characters as the elements of a $n \times n$ matrix. The orthonormality (3.8) and (3.9) in the rows (i.e., in the index i) implies a similar relation in the columns (i.e., in the index j). In

particular, we have $\sum_{i=1}^n \xi_{iE}^* \xi_{iE} = 1$

where $j=E$ refers to the class of the unit element. The unit element is always in a class by itself, so $N_E=1$. The matrix corresponding to the unit element in any rep is the unit matrix of the dimensionality d_i of the rep, so $\chi_{iE}=d_i$ and $\xi_{iE} = \frac{d_i}{\sqrt{g}}$.

Consequently,

$$\sum_{i=1}^n \xi_{iE}^* \xi_{iE} = 1 \quad \text{implies} \quad \sum_{i=1}^n d_i^2 = g. \quad (3.11)$$

For the finite groups under consideration, there will be a unique solution to this relation in terms of integral values of d_i . The dimensionality of the reps (and hence the maximum allowable essential degeneracy) will be determined by the

order and class structure of the group. Thus, for the group D_3 discussed above, with six elements and three classes, we have $2^2+1+1=6$, and the maximum essential degeneracy will be 2

We shall begin our discussion of the character of specific reps by considering the reps D_J , initially limiting J to 1. The ideas of class, character, and orthogonality just discussed were with specific reference to finite groups, but are applicable to infinite groups with but minor modification, the most far-reaching being in the orthogonality relations. In this case the finite sums cited are replaced by integrals over parameters defining the elements of the group. We shall not require the theorem for infinite groups to be discussed here.

A matrix D_1 expresses a transformation of a vector \mathbf{r} (ξ, η, ζ) into a vector \mathbf{r}' (x, y, z). The transformation is orthogonal, leaving the length of the vector unchanged, and may be viewed as a rotation about an axis along $\mathbf{r} \times \mathbf{r}'$ with an angle of

rotation ω such that $\cos \omega = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}| |\mathbf{r}'|}$. To resolve the

group elements into classes note that we can first rotate the vector \mathbf{r} into the x - y plane and, if necessary, perform a rotation about \mathbf{r} so that \mathbf{r}' will also lie in the x - y plane. We thus transform D_1 into a rotation about the z axis where it will, from (3.5) for $\mathbf{U}=(0, 0, 1)$ assume the form

$$R_1(\omega) = \begin{pmatrix} \cos \omega & -\sin \omega & 0 \\ \sin \omega & \cos \omega & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We see that all rotations with the same angle of rotation are in the same class, and that the trace is $1+2 \cos \omega$. It is important to observe that the fundamental interval of ω and the complete range of the character, here a vector with an infinite number of components, may be obtained with ω between the limits 0 and π . This reflects the fact that a rotation, $-\omega$ about a given axis is equivalent to a rotation ω about an oppositely directed axis. Thus every rotation is in the same class as its inverse.

Preparatory to considering other integral values of J , let us transform $R_1(\omega)$ to the purely diagonal form $R'_1(\omega)$ given by

$$R'_1(\omega) = \begin{pmatrix} e^{+i\omega} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-i\omega} \end{pmatrix}.$$

This is the matrix describing the effect of a rotation about the z axis on the three states (+1, 0, -1) of a p electron. For a d electron, we know

that such a rotation may be described by the matrix

$$\mathbf{R}'_2(\omega) = \begin{pmatrix} e^{+2i\omega} & 0 & 0 & 0 & 0 \\ 0 & 0^{+i\omega} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & e^{-i\omega} & 0 \\ 0 & 0 & 0 & 0 & e^{-2i\omega} \end{pmatrix}.$$

Since there is a unique 1 to 1 correspondence between the matrices \mathbf{D}_1 and \mathbf{D}_2 , it follows that any matrix \mathbf{D}_2 may be brought into the form $\mathbf{R}'_2(\omega)$ by the transformation corresponding to that bringing \mathbf{D}_1 into $\mathbf{R}'_1(\omega)$. Here the trace is $1+2 \cos \omega+2 \cos 2\omega$. A similar argument holds for other integral J . A general expression $\chi_J(\omega)$ is then given

by $\sum_{j=0}^J e^{i j \omega}$. This is a geometrical progression in

which the first term is $e^{-i J \omega}$, the common ratio is $e^{i \omega}$, and which contains $2J+1$ terms. The general formula for the sum of such a progression gives

$$\frac{e^{-i J \omega}}{1-e^{i \omega}} (1-e^{i(2J+1)\omega}) = \chi_J(\omega)$$

which may be simplified to

$$\chi_J(\omega) = \frac{\sin(2J+1)\frac{\omega}{2}}{\sin\frac{\omega}{2}}. \quad (3.12)$$

We see from an inspection of this formula that the relations considered for D_1 hold generally for integral J , namely, $\chi(\omega) = \chi(-\omega)$, and that ω on the interval $(0, \pi)$ covers the allowed range.

Let us now consider the group for J half-integral, initially limiting ourselves to $J = \frac{1}{2}$.

Any unitary matrix may be diagonalized, and

\mathbf{D}_1 will assume the particular form $\begin{pmatrix} a & 0 \\ 0 & a^* \end{pmatrix}$

where $|a|$ is $+1$. This diagonalized form of \mathbf{D}_1 can be written as

$$\begin{pmatrix} e^{+i\frac{\omega}{2}} & 0 \\ 0 & e^{-i\frac{\omega}{2}} \end{pmatrix}.$$

By equating the trace of this matrix to that of the nondiagonalized matrix (3.7), we obtain

$$2 \cos \frac{\omega}{2} = 2 \cos \left(\frac{\phi + \psi}{2} \right) \cos \frac{\theta}{2},$$

a relation between the angle of rotation ω and the Euler angles ϕ, θ, ψ . We see that in this group, as well as in that with integral J , every element and its inverse are in the same class, since the character does not depend on the sign of ω and that the classes are determined by the parameter ω . The character of this rep is $2 \cos \omega/2$. The

range of the parameter $\omega/2$ in defining the group element is $(0, \pi)$ corresponding to ω in three dimensions on the interval $(0, 2\pi)$, again confirming the double nature of this group in describing real rotations. Let ω_0 be the angle, less than π , relating two orientations of three-dimensional axes, the corresponding diagonalized two-dimensional matrix being

$$\mathbf{R}_1(\omega_0) = \begin{pmatrix} e^{i\frac{\omega_0}{2}} & 0 \\ 0 & e^{-i\frac{\omega_0}{2}} \end{pmatrix}$$

with character $2 \cos \omega_0/2$. The other two-dimensional matrix corresponding to the same orientation of three-dimensional axes is then given by

$$\begin{aligned} \mathbf{R}_1(\omega_0 + 2\pi) &= \begin{pmatrix} e^{i(\frac{\omega_0}{2} + 2\pi)} & 0 \\ 0 & e^{-i(\frac{\omega_0}{2} + 2\pi)} \end{pmatrix} \\ &= \begin{pmatrix} -e^{+i\frac{\omega_0}{2}} & 0 \\ 0 & -e^{-i\frac{\omega_0}{2}} \end{pmatrix} = -\mathbf{R}_1(\omega_0). \end{aligned}$$

The character of $\mathbf{R}_1(\omega_0 + 2\pi) = 2 \cos(\omega_0/2 + \pi) = -2 \cos \omega_0/2$. Hence these two elements of D_1 corresponding to the same three-dimensional rotation have different traces and belong to different classes, unless the trace is zero. This latter occurs for $\omega_0 = \pi$, at which point the two matrices become

$$\mathbf{R}_1(\pi) = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad \text{and} \quad \mathbf{R}_1(3\pi) = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}.$$

For real rotations of π , the two elements are in the same class, because they are then reciprocal elements.

For higher half-integral values of J , with a unique 1 to 1 correspondence with the elements of the group D_1 , a similar reduction to diagonal form must be obtainable, yielding

$$\begin{pmatrix} e^{+iJ\omega} & 0 & \dots & 0 & 0 \\ 0 & e^{+i(J-1)\omega} & \dots & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \dots & e^{-i(J-1)\omega} & 0 \\ 0 & 0 & \dots & 0 & e^{-iJ\omega} \end{pmatrix}$$

The trace of this matrix is a geometrical progression of the same form as the earlier one for integral J , so the general formula (3.12) will still be valid.

Values of $\chi_J(\omega)$ have been calculated for ω and J of interest and are presented in table 7. In further confirmation of previous discussion, for integral J , $\omega/2$ need only range from 0 to $\pi/2$, since the numerator is an odd polynomial of the denominator, and the numerator and denominator will change sign at the same time. For half

TABLE 7. A character table for the reps D_J

$J \backslash \omega$	0°	60° $2\pi/6$	90° $2\pi/4$	120° $2\pi/3$	180° π	240° $4\pi/3$	270° $6\pi/4$	300° $10\pi/6$	360° 2π
0	1	1	1	1	1	1	1	1	1
1/2 1	2 3	$\sqrt{3}$ 2	$\sqrt{2}$ 1	1 0	0 -1	-1 0	$-\sqrt{2}$ 1	$-\sqrt{3}$ 2	-2 3
3/2 2	4 5	$\sqrt{3}$ 1	0 -1	-1 -1	0 1	1 -1	0 -1	$-\sqrt{3}$ 1	-4 5
5/2 3	6 7	0 -1	$-\sqrt{2}$ -1	0 1	0 -1	0 1	$\sqrt{2}$ -1	0 -1	-6 7
7/2 4	8 9	$-\sqrt{3}$ -2	0 1	1 0	0 1	-1 0	0 1	$\sqrt{3}$ -2	-8 9
9/2 5	10 11	$-\sqrt{3}$ -1	$\sqrt{2}$ 1	-1 -1	0 -1	1 -1	$-\sqrt{2}$ 1	$\sqrt{3}$ -1	-10 11
11/2 6	12 13	0 1	0 -1	0 1	0 1	0 1	0 -1	0 1	-12 13
13/2 7	14 15	$\sqrt{3}$ 2	$-\sqrt{2}$ -1	1 0	0 -1	-1 0	$\sqrt{2}$ -1	$-\sqrt{3}$ 2	-14 15
15/2 8	16 17	$\sqrt{3}$ 1	0 1	-1 -1	0 1	1 -1	0 1	$-\sqrt{3}$ 1	-16 17

integral J , the numerator will be a product of an odd function of $\sin \omega/2$ and an odd function of $\cos \omega/2$. No new values are obtained by going to negative values of ω , since again the numerator and denominator will both change sign together at $\omega=0$. However, the odd terms in $\cos \omega/2$ change sign at $\omega/2=\pi/2$, so that the range $(0, 2\pi)$ in ω will give distinct traces in this latter case. This is due, of course, to the fact that the division between the two homotopy classes of D_1 occurs at $\omega=\pi$ rather than at $\omega=0$.

3.6. Noncubic Crystallographic Groups

We shall now consider the details of the subgroups of D_J in which we shall be interested. Upon the introduction of the crystal field and/or the external magnetic field the general rotation-inversion invariance of the free ion Hamiltonian is destroyed and only those transformations which leave invariant terms representing these external fields are admissible. These were defined in section 2 in terms of the elements $D_1 \times I$. Because of the unique 1 to 1 correspondence between $D_1 \times I$ and $D_J \times I$ for integral $J > 0$, the structure as a group of the allowed transformations for all integral $J > 0$ is the same as those of the perturbing fields. This correspondence does not hold for J half-integral. In this case we again select out of $D_J \times I$ those transformations which correspond in terms of $D_1 \times I$ to invariance of the perturbing fields. However, there are two elements of $D_J \times I$, J half-integral, corresponding to the same transformation in terms of $D_1 \times I$, so there will be

twice as many elements in the corresponding subgroups. For this reason the structure of the subgroups for half-integral J will be different from those for integral J and must be considered separately. We shall initially limit ourselves to those subgroups appropriate for integral J .

We shall examine more closely the detailed structure of those groups listed in table 3, excluding temporarily the infinite groups listed at the bottom of the table and the cubic point groups. These we divide into four general types:

1. The cyclic groups C_n
 2. The dihedral groups D_n
 3. The holohedric groups $C_n \times I$, $D_n \times I$
 4. The hemihedric groups, which are the subgroups of the holohedric groups which do not contain I alone but only in products with rotations.
- The cyclic groups C_n are Abelian, of order n . The elements of the group are the various powers of C_n with $C_n^n = E$. Every element is in a class by itself; consequently, there are n reps for the group.
- The dihedral groups D_n , $n > 1$, are of order $2n$, and include the n powers of C_n and products of these n elements with the elements C_2' . Products of C_n and C_2' are related by $C_2' C_n = C_n^{-1} C_2'$. Hence the group is non-Abelian for $n > 2$. If $n=2$, every element is in a class by itself, otherwise the class structure is more complicated. For $n > 2$, the element C_n^n and its inverse C_n^{n-1} are in the same class. If n is odd, there are $n-1$ powers of C_n (eliminating the unit element in a class by itself) which are grouped by pairs into $\frac{n-1}{2}$ classes.

The n products of C_2^k with the powers of C_n are all in the same class, so we have a total of

$$2 + \frac{n-1}{2} = \frac{n+3}{2}$$

classes. For $n > 2$ and even, the element $C_n^{n/2}$ is its own inverse and like E is in a class by itself. The other $n-2$ powers of C_n fall into $\frac{n-2}{2}$

classes, each with two elements. The n products of the form $C_n^k C_2^l$ will in this case fall into two classes, one with k even, the other with k odd. Here the total number of classes is given by

$$2 + \frac{n-2}{2} + 2 = 3 + \frac{n}{2}$$

The case for $n=3$ has been discussed above (section 3.4) in detail in connection with the definition of class. In the present work, this is the only case we shall be concerned with, though a little geometric consideration on the symmetry of the pentagon, heptagon, etc, will confirm the above discussion of odd n . Since n is odd, an axis of the form C_2^k will pass through a vertex and the opposite side of a n -gon. For n even, such axes will join either opposite pairs of vertices or sides, and no operation of the group either represents a rotation about a line joining a vertex and a side, or will transform a rotation about one type of axis into one about the other.

The holohedric groups $D_n \times I$, $C_n \times I$ contain twice as many elements as D_n , C_n and hence are of order $4n$ or $2n$, respectively. Since I commutes with all elements, the group $C_n \times I$ is Abelian. It is interesting to note that for odd n it is also cyclic since the group can be written in terms of the powers of $C_n I$, with $(C_n I)^n = I$. In the case of $D_n \times I$, if A, B , and X are elements of D_n such that $A = X B X^{-1}$, the element $I X$ will similarly transform B into A . Thus the addition of I to the group elements does not affect the class structure of those elements not containing I . On the other hand $I A = X (I B) X^{-1} = (I X) (I B) (I X)^{-1}$ so the class structure of those elements containing I exactly reproduces that of those elements not containing I .

Let us consider the relations between D_3 , $D_3 \times I$ ($= D_{3d}$) and the hemihedric subgroup C_{3v} . D_3 contains six elements in three classes as follows: $E; C_3, C_3^2; C_2, C_2, C_2$. D_{3d} contains these three classes and three more: $I; IC_3, IC_3^2; IC_2, IC_2, IC_2$. We will now consider which elements must be suppressed to eliminate I as an element of the group without reverting to D_3 . Clearly, I must itself be eliminated, as well as the class of $C_2 I$, since $(C_2 I)^2 = I$. If the class of IC_2 is eliminated we are back to D_3 , so we retain these three elements, noting that we cannot obtain I from powers of these elements as was done with IC_3 since the square of any of them is the unit element. In looking at the classes not containing I , we must retain E to have a group, and the class of C_3 must be retained, since $C_3^2 = (IC_3) (IC_3)$ and both of the latter are retained. On the other hand, $C_2 (IC_2) = I$, so we conclude that the class of C_2 must

also be eliminated, so we obtain for the class structure of C_{3v} : $E; C_3, C_3^2; IC_3, IC_3, IC_3$ which is the same as that of D_3 (i.e., the unit element, two elements of order three, and three elements of order two). Furthermore, the multiplicative relations between the elements of C_{3v} are the same as those between corresponding elements of D_3 . The product of two elements of C_{3v} , neither containing I or both containing I clearly follows the parallel relations in D_3 . If only one contains I , say $IC_2 C_3$, its products with the elements E, C_3, C_3^2 yield I times the products of E, C_3, C_3^2 with the element $C_2 C_3$ of D_3 . We shall see later the importance of the fact that not only is the class structure the same, but that the corresponding multiplication table is the same (though the latter implies the former) for the two groups.

We may generalize these considerations, and assert that the structure of any hemihedric group is the same as that of the pure rotation group from which it is derived through the intermediary holohedric group of double order. Such groups are said to be isomorphic. The only difference, mathematically superficial, is in the geometric designation of the operations involved. The primary significance of this for the present is that the number of actually independent groups to be considered is considerably less than the 27 finite groups enumerated in table 3, and that only the structures of the pure rotation groups and the holohedric groups have to be considered.

3.7. Noncubic Double Groups

We now turn our attention to the double groups, which we shall designate as ΠC_{2n} , ΠD_{2n} , etc., depending on the corresponding subgroup of $D_1 \times I$. This notation will be used when it is desired to particularly emphasize the double group. Usually we shall merely imply these groups through the use of the symbols C_{2n} , D_{2n} , etc., in connection with half-integral J . Some of these groups were first considered by Bethe (1929). The problem was again considered by Opechowski (1940), who clarified many of the ideas involved. We shall begin with the groups ΠC_n . The element of $D_{1/2}$ for $\phi = 2\pi/n$, $\theta = \psi = 0$ is, from (3.7), $[D_{1/2} = U^{-1}]$ the matrix

$$\begin{pmatrix} e^{+i\pi/n} & 0 \\ 0 & e^{-i\pi/n} \end{pmatrix}$$

This is obviously the element of a cyclic group of order $2n$. We shall call this ΠC_n , observing that

ΠC_n^2 is given by the matrix $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ which

we usually designate by R . The unit element is then R^2 . As in the case of C_n , each element of ΠC_n is in a class by itself. The groups C_{2n} and ΠC_n are isomorphic. The inverse of ΠC_n^2 is $\Pi C_n^{2n-2} = R_{\Pi C_n}^{-2}$.

The double dihedral groups consist of the powers

of πC_n , and the powers and products of the matrix $D_n(0, \pi, 0) = \pi C_n' = \begin{pmatrix} 0 & +1 \\ -1 & 0 \end{pmatrix}$ with πC_n and

its powers. The element $\pi C_n'$ is of order 4; its square is R . We note from direct multiplication of the corresponding matrices that $(\pi C_n')(\pi C_n) = (\pi C_n^{-1})(\pi C_n')$. Groups such as this are called dicyclic groups, and have the defining relations as abstract groups $A^n = B^2 = R$, $R^2 = E$, $A^{-1}B = BA$, a group of order $4n$. The class structure may be determined from the matrices themselves, or from the defining relations of the abstract group. A pictorial argument is here inapplicable ab initio, but we shall investigate the relations between the structure of D_n and πD_n . We shall use the defining relations.

The elements which commute with A clearly include the $2n$ powers of A , but no element involving B will commute with A (unless $A^{-1} = A$, which implies that $A = R$ and that $n = 1$, which is excluded, since $n \geq 2$). Thus the order of the normalizer of A is $2n$, its index is two, and there are two elements in the class of A , which we see immediately from the defining relations are A and A^{-1} . Generally, A^k and A^{-k} will be in the same class, except when $k = n$, since $A^n = R$ commutes with all elements and is in a class by itself.

The normalizer of B consists of B , R , B^3 , and E , of order 4 and index n , so there are n elements in the class of B , obtained by transforming B with the first n powers of A . We have $ABA^{-1} = A^2B$; $A^2BA^{-2} = A^4B$; etc., or generally the even powers of A , times B . We have left over the class of AB , with normalizer AB , R , $(AB)^3$, E . The n elements in the class of AB may likewise be obtained by transforming by the first n powers of A , yielding generally the odd powers of A times B . Summarizing, there is the class of E with one element, the class of R with one element, the $(n-1)$ classes of A^k with two elements, the class of B with n elements, and the class of AB with n elements. Altogether there are $n+3$ classes. We notice that the class structure is the same for both odd and even n .

It will now be interesting to compare the class structure of πD_n with that of the group D_n from which it is derived. Let us write the defining relations of D_n in terms of the abstract elements A and B as $A^n = B^2 = E$; $AB = BA^{-1}$, so that we distinguish corresponding elements by italics. Let us further, in πD_n , write powers of A greater than n in terms of products of powers of A less than n and the element R . Thus, $A^{n+k} = RA^k$, and $A^{-k} = A^{2n-k} = RA^{n-k}$. Let us consider the class of A , which is (A, A^{-1}) . There are four elements in πD_n corresponding to these two, A and RA to the first, $A^{-1} = RA^{n-1}$ and A^{n-1} to the second. These four elements are in separate classes by pairs (A, RA^{n-1}) and (RA, A^{n-1}) . If n is even, these are both classes of order $2n$. If n is odd, A is of order $2n$, but RA is of order n . Similarly, the class of A^k , A^{-k} will generally

split into (A^k, RA^{n-k}) and (A^{-k}, RA^k) . If n is even, these are of the same order; if n is odd, these are of the same order for even k , and of different order for odd k . The sole exception to this will occur if $k = n/2$ (which will require n to be even), where there is but one element in the class $(A^{n/2})$ and the two elements $A^{n/2}$, $RA^{n/2}$ are in the same class.

Let us now consider the classes of B and B . If n is odd, the class of B contains A^2B , A^4B , . . . $A^{n-1}B$, RA^2B , RA^4B . . . $RA^{n-2}B$, but it does not contain RB , nor does it contain AB . The class of AB also contains A^3B , . . . $A^{n-2}B$, RB , RA^3B . . . $RA^{n-1}B$. On the other hand, the class of B contains B , AB , A^2B , . . . $A^{n-1}B$, so the class of B will split into two classes, one containing B and RA^2B , the other containing AB and RB (and other elements as indicated above). If n is even the class of B will still include only the even powers of A times B , but $A^n = R$ will now be included, so B and RB are now in the same class. The class of B now involves only B and the even powers of A times B , so for n even the class of B will not split, but will merely contain twice as many elements.

We may summarize this behavior by the following diagram, where we give in one column the typical classes of D_n , in the next column the corresponding classes of πD_n , for both odd and even n .

n odd	
D_n	πD_n
E	(E) (R)
(A^k, A^{-k})	(A ^k , RA ^{n-k}) (A ^{n-k} , RA ^k)
$(B, AB, A^2B \dots)$	(B, RAB, A^2B \dots) (RB, AB, RA^2B \dots)
n even	
(E)	(E) (R)
(A^k, A^{-k})	(A ^k , RA ^{n-k}) (A ^{n-k} , RA ^k)
$(A^{n/2})$	(A ^{n/2} , RA ^{n/2})
$(B, A^2B, A^4B \dots)$	(B, RB, A^2B, RA^2B \dots)
$(AB, A^3B, A^5B \dots)$	(AB, RAB, A^3B, RA^3B, \dots)

We see that the class structure of πD_n follows the same pattern for both odd and even n ; the pattern varies in the case of D_n , in particular, for classes of elements of order 2. Results of the comparison may be summarized into the following rules, first given by Opechowski:

1. For each class of D_n of order other than 2, there are two classes of πD_n each having the same number of elements as the class of D_n .
2. If there is but one class in D_n of order 2 (n is odd), this will in πD_n follow rule (1).
3. If there are two or more classes in D_n of order 2, (n is even) i.e., there are two or more mutually perpendicular two-fold axes, these classes

are real, i.e., +1 or -1, they are designated as "A" if the element for rotation about the z -axis is +1, "B" if the element is -1. They may be further distinguished by subscripts 1, 2 according as the element for rotation about the y -axis is +1 or -1; g or u for the inversion; or (') or (") for +1, -1, respectively, for reflection in the x - y plane. While this is the general policy to which the molecular notation conforms, it is usually abandoned in the case of the group D_2 or D_{2h} . This is because the x , y , and z axes are distinguishable only by convention, and the reps are called B_1, B_2, B_3 . This does not seem to be warranted; even the assignment of subscripts in the B 's must be done by convention, so one may as well conform to the general rule. If the elements are not all real, we use the symbol C_k , according to the quantity $(\sqrt[n]{1})^k$. The general procedures of molecular spectroscopy merge a complex rep with its conjugate, which must also appear, as a two-dimensional rep designated as "E". We are interested in distinguishing these reps because of their separation when a magnetic field is applied.

The double groups do not appear in problems of molecular vibrations. Their reps are of dimensionality four, two, and one. That one of dimensionality four is the appropriate collection of the elements of the rep $D_{3/2}$ of U_3 , so we designate it as $D_{3/2}$ here also. Some of those of dimensionality two which appear are similarly collections of elements of $D_{3/2}$, which we continue to designate as $D_{3/2}$; those two dimensional reps not so designated are assigned the symbol ${}_2S_k$, where "S" represents the idea of "spin" or "specific", the anterior subscript for a two-dimensional rep, and the posterior one (if necessary) by convention. Similarly, the one-dimensional reps are designated "S", with a distinguishing subscript according to the root of 1 involved in the rep.

The reps of the cyclic groups C_n are all one-dimensional. It is clear that just as the group consists of the n powers of C_n , so also will the n powers of $\sqrt[n]{1}$ serve as a rep. It is similarly clear that the n powers of $(\sqrt[n]{1})^k$ will also serve as a rep, where k may be 2, 3, 4, . . . n . If $k=n$, we designate it by A , a general notation for a rep where every rotation about the z -axis is the unit one-dimensional matrix. If n is even, for $k=n/2$ we use B , similarly a general notation for a rep where the element of the rep corresponding to the basic group element C_n is the 1×1 matrix (-1) . In other words, A is a symmetric rep, B an anti-symmetric rep. Note that C_{n-2} is the complex conjugate rep to C_n .

The double cyclic groups πC_n are isomorphic with the group C_{2n} , and hence the reps are the various powers of $(\sqrt[n]{1})^k$. For k even, we designate the rep by $C_{k/n}$ or A or B since it will then be also the corresponding rep of C_n . For k odd, we shall call the rep S_k (S for spin or specific) to emphasize that it is peculiar to the double group. S_k and S_{2n-k} are complex conjugate reps. We note that, if $\sqrt[n]{1} = \omega, \omega^2, 1$ then $\sqrt[n]{1} = -\omega^2, \sqrt[n]{1} = -i\omega$.

The dihedral groups are less easily analyzed. The reps (and characters) for D_2 were obtained above in connection with the illustration of the application of the factor group concept. In terms of numbering there used, we shall describe (4) as A_1 , (1) as A_2 , (2) as B_1 , and (3) as B_2 . Here again, A is used for reps which are totally symmetric for rotations about the z -axis, being distinguished by subscripts according to the behavior for rotations about the secondary y -axis. B is used where the basic rotation about the z -axis is antisymmetric, and again distinguishing various behavior for rotations about the y -axis.

The group D_2 contains the invariant subgroup (E, C_2, C_2') , so two reps are obtained as the rep (1, 1) and (1, -1) of the factor group. Since (3.11) admits of the solution $2^2 + 1^2 + 1^2 = 6$, the other rep is two-dimensional and its character may be deduced from the orthogonality relation for group characters. The one dimensional reps are designated as A_1 and A_2 , while two-dimensional reps generally are designated as E .

For D_4 , 8 elements in 5 classes, eq (3.11) admits the solution $2^2 + 1^2 + 1^2 + 1^2 + 1^2 = 8$, so there are four 1-dimensional reps and one 2-dimensional rep. Since (E, C_4, C_2, C_2') is an invariant subgroup, two reps are again the reps of the factor group (1, 1) and (1, -1), designated A_1 and A_2 . We may also verify that (E, C_2') is an invariant subgroup, whose factor group is isomorphic with D_2 , having the structure $(E, C_2') (C_4, C_2) (C_2, C_2' C_2') (C_4 C_2, C_2' C_2')$. A_1 and A_2 of D_2 give no new reps, but B_1 and B_2 of D_2 will give us new reps, which we shall designate as B_1 and B_2 in this case also. The two-dimensional rep E may now be found from the orthogonality condition.

In the case of D_6 , 12 elements in 6 classes, $2^2 + 2^2 + 1^2 + 1^2 + 1^2 + 1^2 = 12$. The group C_6 is an invariant subgroup, and the (1, 1) and (1, -1) of its factor group C_2 are A_1 and A_2 , respectively of D_6 . The group C_3 is also an invariant subgroup, with factor group D_2 , whose reps B_1 and B_2 give B_1 and B_2 of D_6 . Finally, the group (E, C_2) is an invariant subgroup with factor group D_3 , whose rep E will yield a new rep E_2 of D_6 . We find from the orthogonality condition the rep E_1 . The reps are numbered according to the reps of D_{2n} , with which they may be correlated (table 10).

For the double groups generally, cyclic or dicyclic, (E, R) is an invariant subgroup whose factor group is the corresponding single group, so the reps of the single group are also reps of the double group, as we have already seen in the case of the cyclic groups. For any half-integral J , the traces of $D_J(E)$ and $D_J(R)$ are $(2J+1)$ and $-(2J+1)$, respectively. Hence we see that we cannot resolve a faithful representation for half-integral J into a sum of reps, any one of which has the same trace for E and R . This latter is clearly the case for those reps derived from the invariant subgroup (E, R) , namely those which are also reps of the single group, so we conclude that they will not be useful for describing

states arising from an odd number of electrons. Those reps which are peculiar to the double groups we shall call specific reps (the basis for the "S" notation in the cyclic groups used earlier); we include the non-specific reps for the completeness required by the orthogonality theorem and for the treatment of selection rules.

ΠD_2 contains 8 elements in 5 classes, so $2^2+1^2+1^2+1^2+1^2=8$. The four 1-dimensional reps are the reps of D_2 , so the only specific rep is two-dimensional. It is obvious that this must be equivalent to the appropriate elements of D_{2h} , or in other words, the representation is its own rep. We shall emphasize this by designating the rep as D_{2h} .

ΠD_3 has 12 elements in 6 classes, whence $2^2+2^2+1^2+1^2+1^2+1^2=12$. Half of these are the reps of D_3 , so there are two specific one-dimensional reps and one specific two-dimensional rep. We observe that $(E, \Pi C_3, R_{\Pi} C_3)$ is an invariant subgroup. Its associated complexes (cosets) are $(\Pi C_2, \Pi C_{2h}, R_{\Pi} C_{2h}, R_{\Pi} C_2)$, $(\Pi C_3, R, R_{\Pi} C_3)$ and $(\Pi C_{2h}, R_{\Pi} C_2, R_{\Pi} C_3, R_{\Pi} C_{2h})$ yielding a factor group isomorphous with C_4 . The reps A and B of C_4 yield no new reps, but C_1 and C_3 will yield specific reps which we shall distinguish as S_1 and S_3 . Because of these two 1-dimensional reps into which the representation from D_{3h} could perhaps split, we cannot conclude that the representation from D_{3h} is also a rep here, but this proves to be actually the case if the two-dimensional character is calculated from the orthogonality relations. As before, we call this rep D_{3h} .

ΠD_4 has 16 elements in 7 classes. Since $2^2+2^2+2^2+1^2+1^2+1^2+1^2=16$, we find on subtracting the reps of D_4 that there are but two specific reps, both two-dimensional. One of these must be D_{4h} ; the other, derived from the orthogonality relations, we shall call ${}_2S$.

ΠD_6 has 24 elements in 9 classes. Equation (3.11) gives us $2^2+2^2+2^2+2^2+2^2+1^2+1^2+1^2+1^2=24$. Eliminating the reps of D_6 , we have three specific reps, all two-dimensional. One of these must be D_{6h} . Unfortunately, we cannot obtain either of the other two characters by the methods used so far, and an alternate procedure must be used. We obtain from table 7 the character of the representation arising from $D_{3/2}$, which is four-dimensional and hence reducible. By application of the expansion theorem (3.10) we see that $D_{3/2}$ is included once in the representation. The other rep involved we shall call ${}_2S_1$. The final rep may now be obtained from the orthogonality theorem as ${}_2S_2$.

The holohedric groups, single or double, are defined as the direct product of the group $(E, I) = I$ with a pure rotation group. The reps may be obtained as products of the reps $(1, 1)$ and $(1, -1)$ of I with those of the pure rotation group. The reps are designated like those of the pure rotation group, with the additional subscript "g" if $(1, 1)$ is used and "u" if $(1, -1)$ is used.

It has already been pointed out that the hemi-

hedric groups are isomorphous with the corresponding (in the sense of table 3) pure rotation group and hence have the same reps. Most of these hemihedric reps are described just like the corresponding pure rotation reps. The exceptions are the groups D_{2h} , C_{2h} , C_4 , and S_4 . For the first three of these, the reps are classified according to behavior under the operation $IC_2 = \sigma_h$, symmetric reps carrying a (') while antisymmetric reps have ("). D_{3h} is further subclassified according to C_2' . This difference in notation is peculiar to the single groups—the specific reps of all hemihedric double groups will be designated like those of the corresponding pure rotation group. In the case of S_4 , isomorphous with C_4 , the generating element is taken as the operation $IC_2 = C_4 IC_2 = C_4 \sigma_h$, which is usually considered as the fundamental rotation-reflection operation. This is why the rep C_1 of S_4 corresponds to C_3 of C_4 rather than to C_1 of C_4 .

3.9. Infinite and Cubic Groups

We have finally to discuss the infinite groups involving infinitesimal rotations about the z -axis, namely C_{∞} , $C_{\infty h}$, D_{∞} , $D_{\infty h}$, $C_{\infty v}$, and the cubic groups. In C_{∞} , the group operation consists of arbitrary rotations $C(\phi)$ such that $C(\phi_1) C(\phi_2) = C(\phi_2) C(\phi_1) = C(\phi_1 + \phi_2)$, and that $C(2\pi) = C(0)$. If k is any number, $e^{ik\phi}$ will meet the first requirement, but the second limits k to a real integer, positive or negative, including zero. If $k=0$, we shall call the rep A , otherwise C_k . For the group ΠC_{∞} , the first of the above conditions holds, but the second is replaced by $C(2\pi) = -C(0)$. Here k must be half-integral, positive or negative, and we designate the reps by S_{2k} ($2k$ is an odd integer). $C_{\infty h}$ is $C_{\infty} \times I$, so the reps for this group will carry the subscript "g" or "u".

In D_{∞} , the group elements consist of the rotations $C(\phi)$ and the rotation C_2' , such that $C_2' C(\phi) = C(-\phi) C_2'$. Considering C_{∞} as an invariant subgroup, the reps $(1, 1)$ and $(1, -1)$ of the factor group C_2 yield reps A_1 and A_2 of D_{∞} . As discussed under C_{∞} , $e^{ik\phi}$ will serve as a rep insofar as $C(\phi)$ is concerned, but the operation C_2' makes $e^{ik\phi}$ and $e^{ik(-\phi)}$ equivalent. Except when $k=0$ (which gives A_1 and A_2) our rep for $C(\phi)$ must have the two-dimensional form

$$\begin{pmatrix} e^{ik\phi} & 0 \\ 0 & e^{-ik\phi} \end{pmatrix}$$

and C_2' will assume the form

$$\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

which will transform $C(\phi)$ into $C(-\phi)$. It may be shown that there are no other reps. These reps we shall call E_k , where k is a positive integer. In the case of ΠD_{∞} , the reps become

$$\Pi C(\phi) = \begin{pmatrix} e^{ik\phi/2} & 0 \\ 0 & e^{-ik\phi/2} \end{pmatrix}, \quad \Pi C_2' = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

and we shall designate them as ${}_2S_k$ (k odd). Reps of D_{2k} will have the additional subscripts "g" or "u", while reps of the hemihedric group C_{2k} will be designated like those of D_{2k} .

In the group O , there are 24 elements in 5 classes. These are the class of E , 1 element; the class of C_4 , 6 elements; the class of C_2 , 3 elements; the class of C_2' , the body diagonal axes, 8 elements; and the class of C_2'' (here, C_2'' is not along the y -axis), 6 elements. (3.11) gives $3^2+3^2+2^2+1^2+1^2=24$. The set of elements (E , $3C_2$, $8C_2'$) is an invariant subgroup with factor group of order 2. Its reps (1,1) and (1, -1) are the reps A_1 and A_2 of O . If we consider D_J for $J=1$, we find that neither A_1 nor A_2 is in the representation. Since it cannot be reduced to A_1+E or A_2+E (E being the two-dimensional rep), it must itself be one of the three-dimensional reps. These are generally designated as T —this particular rep we shall call T_1 . D_J , $J=2$, contains neither A_1 , A_2 , or T_1 , so it must reduce to $E+T_2$, but we cannot separate them. On going to $J=3$, we find from this seven-dimensional representation, A_1 and T_1 accounting for four. The remaining three must then be T_2 which, subtracted from the $J=2$ case leaves E . The groups O_h and T_d follow our previous considerations on pure rotation, holohedric, and hemihedric groups. The reps of T_d parallel those of O .

The double group ${}_2O$ has 48 elements. The splitting into classes, as compared with the group O , follows the same rules as for the dihedral groups given in section 3.7, so the class structure is: E (1); R (1); ${}_2C_4$ (6); R_2C_4 (6); ${}_2C_2$ (6); ${}_2C_2'$ (8); R_2C_2' (8); ${}_2C_{2d}$ (12). (3.11), after eliminating the reps of O , indicates three reps for ${}_2O$, two 2-dimensional and one 4-dimensional. As usual, D_{2k} is one of these reps. $D_{3/2}$ in this case is also a rep, the four-dimensional one. The remaining rep obtained from orthogonality we call ${}_2S$.

The group T , 12 elements, may be separated into the class E (1); C_2 (3); C_3 (4); and C_3' (4). It is interesting to note that, in spite of the relatively high symmetry, a rotation C_3 and its inverse $C_3^{-1}=C_3'$ are in different classes. (3.11) yields $3^2+1^2+1^2+1^2=12$. The elements (E , $3C_2$) constitute an invariant subgroup, with a factor group of order 3, C_3 . The reps A , C_1 and C_2 of T are thus the one-dimensional reps A , C_1 , and C_2 of T . The rep T may then be obtained from orthogonality and completeness considerations. The group ${}_2T$, 24 elements divides (section 3.7) into E (1); R (1); ${}_2C_2$ (6); ${}_2C_3$ (4); R_2C_3 (4); ${}_2C_2'$ (4); and R_2C_2' (4). Subtracting off the reps of T , we have $2^2+2^2+2^2=12$, indicating reps we shall call $D_{3/2}$, ${}_2S_1$, and ${}_2S_2$. As the notation suggests, $D_{3/2}$ is its own rep. Unfortunately, our previous techniques for finding reps fail in this case, since the reps ${}_2S_1$ and ${}_2S_2$ always appear together and cannot be separated. We must therefore proceed to a more general method.

The product of any two classes of a group $[C_i]$

and $[C_j]$ with N_i and N_j elements, respectively, will be a total of $N_i N_j$ elements, not necessarily all distinct, which may themselves be written as a sum of classes $[C_k]$, or $[C_i][C_j]=\sum N_{i,j,k}[C_k]$,

with $N_{i,j,k}$ a small nonnegative integer. A particular k may not appear at all, or may appear more than once. It may be shown (Murnaghan, 1938) that this expansion implies the following relation between the traces of the i -th rep:

$$N_i N_j \chi_{iB} \chi_{jB} = \chi_{iB} \sum_k N_{i,j,k} N_k \chi_{kB}$$

where χ_{iB} is the trace of the unit element, i.e., the dimension of the rep. By developing in detail the multiplication table of a group, sufficient relations may be found to calculate the complete character table of any group. In the present case, we already have all but two of the seven possibilities for ${}_2T$. A solution of these equations for $\chi_{iB}=2$ will then give the desired characters.

3.10. Table of Group Characters

The foregoing considerations have been summarized in table 8, a table of group characters for both the single and double groups in which we shall be interested. Each portion of the table is constructed around one of the single holohedric groups. Typical elements of the various classes of the group, together with an indication of the total number of elements in the class, are shown to the right of the group designation, while the designations of the several reps are shown under the group designation. For convenience, we also give the notation of Bethe (1929). The characters for the various classes and reps are the entries in the table. The classes, reps, and characters for the corresponding pure rotation group may be obtained by dropping all classes containing I and the subscripts "g" and "u" on the reps.

Above and to the left of each basic character table are shown the classes and reps of the corresponding hemihedric groups. The appropriate group designations are given upward and to the left of that of the basic holohedric group, while the class structure is given along the same row and the several reps in the same column, so that both classes and reps correspond to those of the holohedric group. A careful study of these will indicate very clearly the relations between a holohedric group and its hemihedric derivatives.

The characters and reps for the specific reps of the double groups are given in table 8 immediately below the reps of the single groups. Because of the frequent splitting of the classes of the single group into two classes discussed above, there will in such cases be two characters for each class of the single group. If A is a class element of the single group, the first double group character will be that for the class of ${}_2A$, the second for that of R_2A . Upon inspection, it will be seen that such pairs of characters have opposite signs, while those arising from classes of the single group which do not split are all zero.

TABLE 8. A table of group characters

$C_{\infty v}$	$D_{\infty h}$	E	∞IC_i		I	$2IC(\phi)$	∞IC_i	E	C(φ)	I	IC(φ)
			$2C(\phi)$	∞C_i							
A_1	A_{1g}	1	1	1	1	1	1	1	1	1	1
	A_{1u}	1	1	-1	-1	-1	-1	1	1	-1	-1
A_2	A_{2g}	1	-1	1	1	1	1	1	$e^{i\phi}$	1	$e^{i\phi}$
	A_{2u}	1	-1	-1	-1	-1	-1	1	$e^{i\phi}$	-1	$-e^{i\phi}$
E_1	E_{1g}	2	0	2	$2 \cos \phi$	0	$2 \cos \phi$	1	$e^{-i\phi}$	1	$e^{-i\phi}$
	E_{1u}	2	0	-2	$-2 \cos \phi$	0	$-2 \cos \phi$	1	$e^{-i\phi}$	-1	$-e^{-i\phi}$
E_2	E_{2g}	2	0	2	$2 \cos 2\phi$	0	$2 \cos 2\phi$	1	$e^{i2\phi}$	1	$e^{i2\phi}$
	E_{2u}	2	0	-2	$-2 \cos 2\phi$	0	$-2 \cos 2\phi$	1	$e^{i2\phi}$	-1	$-e^{i2\phi}$
E_n	E_{ng}	2	0	2	$2 \cos n\phi$	0	$2 \cos n\phi$	1	$e^{in\phi}$	1	$e^{in\phi}$
	E_{nu}	2	0	-2	$-2 \cos n\phi$	0	$-2 \cos n\phi$	1	$e^{in\phi}$	-1	$-e^{in\phi}$
${}_2S_1$	${}_2S_{1g}$	2	0	2	$2 \cos \phi/2$	0	$2 \cos \phi/2$	1	$e^{i\phi/2}$	1	$e^{i\phi/2}$
	${}_2S_{1u}$	2	0	-2	$-2 \cos \phi/2$	0	$-2 \cos \phi/2$	1	$e^{i\phi/2}$	-1	$-e^{i\phi/2}$
${}_3S_2$	${}_3S_{2g}$	2	0	2	$2 \cos 3\phi/2$	0	$2 \cos 3\phi/2$	1	$e^{i3\phi/2}$	1	$e^{i3\phi/2}$
	${}_3S_{2u}$	2	0	-2	$-2 \cos 3\phi/2$	0	$-2 \cos 3\phi/2$	1	$e^{i3\phi/2}$	-1	$-e^{i3\phi/2}$
${}_2S_n$	${}_2S_{ng}$	2	0	2	$2 \cos n\phi/2$	0	$2 \cos n\phi/2$	1	$e^{in\phi/2}$	1	$e^{in\phi/2}$
	${}_2S_{nu}$	2	0	-2	$-2 \cos n\phi/2$	0	$-2 \cos n\phi/2$	1	$e^{in\phi/2}$	-1	$-e^{in\phi/2}$
	(n odd)							1	$e^{-in\phi/2}$	1	$e^{-in\phi/2}$
								1	$e^{-in\phi/2}$	-1	$-e^{-in\phi/2}$
											(n odd)

TABLE 8. A table of group characters—Continued

D _{3h}	E						3C ₂							
	E	2C ₆		2C ₃		3C ₂		E	2C ₆		2C ₃		3C ₂	
		1	1	1	1	1	1		1	1	1	1	1	1
A ₁	1	1	1	1	1	1	1	1	1	1	1	1	1	1
A ₂	1	1	1	1	1	1	1	1	1	1	1	1	1	1
A ₁ '	1	1	1	1	1	1	1	1	1	1	1	1	1	1
A ₂ '	1	1	1	1	1	1	1	1	1	1	1	1	1	1
E'	2	1	-1	1	-1	1	0	0	0	0	0	0	0	0
E''	2	1	-1	1	-1	1	0	0	0	0	0	0	0	0
D _{3h}	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₃)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₄)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₅)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₆)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₇)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₈)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₉)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₀)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₁)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₂)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₃)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₄)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₅)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₆)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₇)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₈)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₁₉)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₀)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₁)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₂)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₃)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₄)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₅)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₆)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₇)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₈)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₂₉)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2
D _{3h} (Γ ₃₀)	2	1	-1	1	-1	1	2	-2	2	-2	2	-2	2	-2

TABLE 8. A table of group characters—Continued

D_{3d}		E	$2C_3$	C_2	$2C_2'$	$2C_2''$	$2C_2'''$					
		E	$2C_3$	C_2	$2C_2'$	$2C_2''$	$2C_2'''$	I	$2C_3$	$2C_2'$	$2C_2''$	
A_1	$A_{1g}(\Gamma_1)$	1	1	1	1	1	1	1	1	1	1	1
A_2	$A_{1u}(\Gamma_2)$	1	1	1	-1	-1	-1	-1	-1	-1	-1	-1
B_1	$A_{2g}(\Gamma_3)$	1	1	1	1	1	1	1	1	1	1	1
B_2	$A_{2u}(\Gamma_4)$	1	1	1	-1	-1	-1	-1	-1	-1	-1	-1
E	$B_{1g}(\Gamma_5)$	2	0	-2	0	0	0	2	0	0	0	0
D_{3d}	$B_{1u}(\Gamma_6)$	2	$\sqrt{2}$	0	0	0	0	2	$\sqrt{2}$	0	0	0
2S	$B_{2g}(\Gamma_7)$	2	$\sqrt{2}$	0	0	0	0	2	$\sqrt{2}$	0	0	0
	$B_{2u}(\Gamma_8)$	2	$-\sqrt{2}$	0	0	0	0	2	$-\sqrt{2}$	0	0	0
	$E_g(\Gamma_9)$	2	0	-2	0	0	0	2	0	0	0	0
	$E_u(\Gamma_{10})$	2	0	-2	0	0	0	2	0	0	0	0
	$D_{3d}^{(u)}$	2	$\sqrt{2}$	0	0	0	0	2	$\sqrt{2}$	0	0	0
	2S_g	2	$-\sqrt{2}$	0	0	0	0	2	$-\sqrt{2}$	0	0	0
	2S_u	2	$\sqrt{2}$	0	0	0	0	2	$\sqrt{2}$	0	0	0

C_{3v}		E	$2C_3$	$3C_2$	$3C_2'$			
		E	$2C_3$	$3C_2$	$3C_2'$	I	$2C_3$	$3C_2'$
A_1	A_{1g}	1	1	1	1	1	1	1
A_2	A_{1u}	1	1	1	-1	-1	-1	-1
E	A_{2g}	2	-1	0	0	2	-1	0
	E_u	2	-1	0	0	2	-1	0
S_1	S_{1g}	1	-1	i	-i	1	-1	i
	S_{1u}	1	-1	-i	i	1	-1	-i
S_2	S_{2g}	1	-1	i	-i	1	-1	i
	S_{2u}	1	-1	-i	i	1	-1	-i
D_{3d}	$D_{3d}^{(u)}$	2	-2	0	0	2	-2	0
	$D_{3d}^{(g)}$	2	-2	0	0	2	-2	0

TABLE 8. A table of group characters—Continued

C _n	E		C ₂		IC ₂ C ₂		IC ₂ C ₂		I	IC ₂	IC ₂	IC ₂	IC ₂
	D _{2h}	E	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂					
A ₁	A _{1g}	1	1	1	1	1	1	1	1	1	1	1	1
A ₂	A _{1u}	1	1	1	1	1	1	1	-1	-1	-1	-1	-1
	A _{2g}	1	1	-1	-1	-1	-1	-1	1	1	1	1	1
	A _{2u}	1	1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1
B ₁	B _{1g}	1	-1	1	1	1	1	1	1	-1	-1	-1	-1
B ₂	B _{1u}	1	-1	1	1	1	1	1	-1	-1	-1	-1	-1
	B _{2g}	1	-1	-1	-1	-1	-1	-1	1	1	1	1	1
	B _{2u}	1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1
D _{2h}	D _{2hg}	2	-2	0	0	0	0	0	2	-2	-2	0	0
	D _{2hu}	2	-2	0	0	0	0	0	-2	2	2	0	0

C _n	E		C ₂		C ₂		C ₂		C ₂		C ₂		C ₂		C ₂	
	E	IC ₂	IC ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂	C ₂
C ₁	C _{1g}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	C _{1u}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
C ₂	C _{2g}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	C _{2u}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
A'	A' _g	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	A' _u	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
S ₁	S _{1g}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	S _{1u}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
S ₂	S _{2g}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	S _{2u}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
S ₃	S _{3g}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	S _{3u}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
S ₄	S _{4g}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	S _{4u}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
S ₅	S _{5g}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	S _{5u}	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

S ₁	I		IC ₁		C ₂		IC ₂		I	IC ₁	IC ₂	IC ₃
	C _{1A}	E	C ₁	IC ₁	C ₂	IC ₂	C ₂	IC ₂				
C ₂	1	1	i	i	-1	-1	-i	-i	1	i	-1	-i
C _{1A}	1	1	-1	-1	-1	-1	-i	-i	-1	-1	-1	-1
B ₂	1	1	-1	-1	-1	-1	-i	-i	-1	-1	-1	-1
B _{1A}	1	1	-1	-1	-1	-1	-i	-i	-1	-1	-1	-1
C ₁	1	1	-i	-i	-1	-1	i	i	1	-1	-1	-1
C _{2A}	1	1	-i	-i	-1	-1	i	i	1	-1	-1	-1
A ₂	1	1	-1	-1	-1	-1	-i	-i	-1	-1	-1	-1
A _{1A}	1	1	-1	-1	-1	-1	-i	-i	-1	-1	-1	-1
S ₂	1	-1	$\sqrt{2}$	$-\sqrt{2}$	i	-i	$i\sqrt{2}$	$-i\sqrt{2}$	1	$\sqrt{2}$	$-\sqrt{2}$	$i\sqrt{2}$
S _{1A}	1	-1	$\sqrt{2}$	$-\sqrt{2}$	i	-i	$i\sqrt{2}$	$-i\sqrt{2}$	1	$\sqrt{2}$	$-\sqrt{2}$	$i\sqrt{2}$
S _{2A}	1	-1	$i\sqrt{2}$	$-i\sqrt{2}$	i	-i	$\sqrt{2}$	$-\sqrt{2}$	1	$i\sqrt{2}$	$-i\sqrt{2}$	$\sqrt{2}$
S _{1A}	1	-1	$i\sqrt{2}$	$-i\sqrt{2}$	i	-i	$\sqrt{2}$	$-\sqrt{2}$	1	$i\sqrt{2}$	$-i\sqrt{2}$	$\sqrt{2}$
S ₂	1	-1	$-\sqrt{2}$	$\sqrt{2}$	i	-i	$-i\sqrt{2}$	$i\sqrt{2}$	1	$-\sqrt{2}$	$\sqrt{2}$	$-i\sqrt{2}$
S _{1A}	1	-1	$-\sqrt{2}$	$\sqrt{2}$	i	-i	$-i\sqrt{2}$	$i\sqrt{2}$	1	$-\sqrt{2}$	$\sqrt{2}$	$-i\sqrt{2}$
S _{2A}	1	-1	$i\sqrt{2}$	$-i\sqrt{2}$	i	-i	$\sqrt{2}$	$-\sqrt{2}$	1	$i\sqrt{2}$	$-i\sqrt{2}$	$\sqrt{2}$
S _{1A}	1	-1	$i\sqrt{2}$	$-i\sqrt{2}$	i	-i	$\sqrt{2}$	$-\sqrt{2}$	1	$i\sqrt{2}$	$-i\sqrt{2}$	$\sqrt{2}$

S ₂	E	C ₂	C ₂	IC ₂	IC ₂	IC ₂	IC ₂	IC ₂	IC ₂	IC ₂	IC ₂	IC ₂
C _{1A}	1	ω	ω	ω^2	ω^2	ω	ω	ω^2	ω^2	ω	ω	ω^2
C _{1A}	1	ω	ω	ω^2	ω^2	ω	ω	ω^2	ω^2	ω	ω	ω^2
C _{2A}	1	ω	ω	ω^2	ω^2	ω	ω	ω^2	ω^2	ω	ω	ω^2
C _{2A}	1	ω	ω	ω^2	ω^2	ω	ω	ω^2	ω^2	ω	ω	ω^2
A ₂	1	1	1	1	1	1	1	1	1	1	1	1
A _{1A}	1	1	1	1	1	1	1	1	1	1	1	1
S _{1A}	1	-1	$-\omega^2$	ω^2	ω	ω	$-\omega$	$-\omega$	$-\omega^2$	ω^2	ω	$-\omega$
S _{1A}	1	-1	$-\omega^2$	ω^2	ω	ω	$-\omega$	$-\omega$	$-\omega^2$	ω^2	ω	$-\omega$
S _{2A}	1	-1	-1	1	1	1	-1	-1	-1	1	1	-1
S _{2A}	1	-1	-1	1	1	1	-1	-1	-1	1	1	-1
S _{2A}	1	-1	$-\omega$	ω	ω^2	ω^2	$-\omega$	$-\omega$	$-\omega^2$	ω^2	ω	$-\omega^2$
S _{2A}	1	-1	$-\omega$	ω	ω^2	ω^2	$-\omega$	$-\omega$	$-\omega^2$	ω^2	ω	$-\omega^2$

TABLE 8. A table of group characters—Continued

C_4	C_{2A}	E		IC_2		I		IC_2	
		E	C_2	I	IC_2	I	IC_2		
A''	B_g	1	-1	1	-1	1	-1	1	-1
	B_u	1	-1	-1	1	-1	1	-1	1
A'	A_g	1	1	1	1	1	1	1	1
	A_u	1	1	-1	-1	-1	-1	-1	-1
S_1	S_{1g}	1	-1	i	$-i$	1	-1	i	$-i$
	S_{1u}	1	-1	i	$-i$	-1	1	$-i$	i
S_3	S_{3g}	1	-1	$-i$	i	1	-1	$-i$	i
	S_{3u}	1	-1	$-i$	i	-1	1	i	$-i$

S_2		E	I
A_u		1	-1
A_g		1	1

T_d	O_h	E		$6IC_4$		$3C_2$		$6IC_2'$		$8C_3'$		I		$6IC_4$		$3IC_2$		$6IC_2'$		$8IC_3'$	
		E	$6C_4$	$3C_2$	$6C_2'$	$8C_3'$	I	$6IC_4$	$3IC_2$	$6IC_2'$	$8IC_3'$	I	$6IC_4$	$3IC_2$	$6IC_2'$	$8IC_3'$					
A_1	$A_{1g}(\Gamma_1)$	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	A_{1u}	1	1	1	1	1	1	1	1	1	1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1
A_2	$A_{2g}(\Gamma_2)$	1	-1	1	-1	1	-1	1	-1	1	-1	1	-1	1	-1	1	-1	1	-1	1	-1
	A_{2u}	1	-1	1	-1	1	-1	1	-1	1	-1	-1	1	-1	1	-1	1	-1	1	-1	1
E	$E_g(\Gamma_3)$	2	0	2	0	-1	2	0	-2	0	2	0	0	2	0	0	-1	2	0	0	-1
	E_u	2	0	2	0	-1	-2	0	2	0	-2	0	0	-2	0	0	1	-2	0	0	1
T_1	$T_{1g}(\Gamma_4)$	3	1	-1	-1	0	-3	1	-1	-1	-3	1	-1	-1	-1	0	0	-3	1	-1	0
	T_{1u}	3	1	-1	-1	0	-3	1	-1	-1	-3	1	-1	-1	0	0	0	-3	1	-1	0
T_2	$T_{2g}(\Gamma_5)$	3	-1	-1	1	0	3	-1	1	1	3	-1	1	-1	1	0	0	3	-1	1	0
	T_{2u}	3	-1	-1	1	0	-3	1	1	1	-3	-1	1	-1	1	0	0	-3	-1	1	0
D_{3d}	$D_{3d}(\Gamma_6)$	2	-2	$\sqrt{2}$	$-\sqrt{2}$	0	0	1	-1	2	-2	$\sqrt{2}$	$-\sqrt{2}$	0	0	1	-1	2	-2	$\sqrt{2}$	$-\sqrt{2}$
	$D_{3d}u$	2	-2	$\sqrt{2}$	$-\sqrt{2}$	0	0	1	-1	-2	2	$-\sqrt{2}$	$\sqrt{2}$	0	0	-1	1	-2	2	$-\sqrt{2}$	$\sqrt{2}$
$2S$	$2S_g(\Gamma_7)$	2	-2	$-\sqrt{2}$	$\sqrt{2}$	0	0	1	-1	2	-2	$-\sqrt{2}$	$\sqrt{2}$	0	0	1	-1	2	-2	$-\sqrt{2}$	$\sqrt{2}$
	$2S_u$	2	-2	$-\sqrt{2}$	$\sqrt{2}$	0	0	1	-1	-2	2	$\sqrt{2}$	$-\sqrt{2}$	0	0	-1	1	-2	2	$\sqrt{2}$	$-\sqrt{2}$
D_{3d2}	$D_{3d2}(\Gamma_8)$	4	-4	0	0	0	0	-1	1	4	-4	0	0	0	0	-1	1	4	-4	0	0
	$D_{3d2}u$	4	-4	0	0	0	0	-1	1	-4	4	0	0	0	0	-1	1	-4	4	0	0

T_h	E	$3C_2$	$4C_3$	$4C_3'$	I	$3IC_2$	$4IC_3$	$4IC_3'$
A_g	1	1	1	1	1	1	1	1
A_u	1	1	1	1	-1	-1	-1	-1
C_{1g}	1	1	ω	ω^2	1	1	$-\omega$	$-\omega^2$
C_{1u}	1	1	ω	ω^2	-1	-1	$-\omega$	$-\omega^2$
C_{2g}	1	1	ω^2	ω	1	1	$-\omega^2$	$-\omega$
C_{2u}	1	1	ω^2	ω	-1	-1	$-\omega^2$	$-\omega$
T_g	3	-1	0	0	3	-1	0	0
T_u	3	-1	0	0	-3	1	0	0
D_{3d}	2	-2	0	1	-1	0	1	-1
$D_{3d}u$	2	-2	0	1	-1	0	-1	1
$2S_{1g}$	2	-2	0	ω	$-\omega$	0	$-\omega$	$-\omega$
$2S_{1u}$	2	-2	0	ω	$-\omega$	0	$-\omega$	$-\omega$
$2S_{2g}$	2	-2	0	ω^2	$-\omega^2$	0	$-\omega^2$	$-\omega^2$
$2S_{2u}$	2	-2	0	ω^2	$-\omega^2$	0	$-\omega^2$	$-\omega^2$

4. Degeneracy, Selection Rules, and Polarization

4.1. Splitting of Free Ion Levels

We are now in a position to apply the results of our previous considerations to the problem of degeneracy in the presence of a perturbing external field. The general procedure has been covered in the previous section; we consider here the specific example of a free ion level described as $4_u(J=4, (-1)^{2l} = -1)$ in a field of C_{6v} symmetry. From table 8 we obtain the class structure of C_{6v} as E, $2C_6$, $2C_3$, C_2 , $3IC_2'$, $3IC_2''$. The traces of the matrices of the rotation group $J=4$ corresponding to the first four of these classes may be obtained directly from table 7. Those for the last two classes will be -1 times the traces of the corresponding pure rotation. We obtain the following for the character table of a *reducible* representation of C_{6v} . We also include for immediate convenience the reps A_1 and A_2 .

	E	$2C_6$	$2C_3$	C_2	$3IC_2'$	$3IC_2''$
4_u	9	-2	0	1	-1	-1
A_1	1	1	1	1	1	1
A_2	1	1	1	1	-1	-1

The expansion theorem for group characters (3.10) was given in terms of the weighted traces

$$\xi_{ij} = \sqrt{\frac{N_j}{g}} \chi_{ij}$$

and

$$\Xi_j = \sqrt{\frac{N_j}{g}} X_j$$

but it is usually more convenient to use the traces directly. Thus $a_i = \xi_i \cdot \Xi$ (3.10) becomes $ga_i = \sum_j N_j \chi_{ij} X_j$. Using this latter form, we obtain for A_1 : $12a_{A_1} = 1(1)(9) + 2(1)(-2) + 2(1)(0) + 1(1)(1) + 3(1)(-1) + 3(1)(-1) = 0$, so A_1 will not appear in the reduction. Likewise, $12a_{A_2} = 1(1)(9) + 2(1)(-2) + 2(1)(0) + 1(1)(1) + 3(-1)(-1) + 3(-1)(-1) = 12$, so A_2 will appear in the reduction once.

The results of this and similar calculation for all the subgroups of D_7 of present interest are given in table 9. As in table 8, the results for the pure rotation groups are not given separately, and may be obtained from the associated holohedric group by omitting the "g"- "u" distinction. Only the group I has been omitted; this has no symmetry apart from the inversion, and states are either A_g or A_u . The integral range of J is from 0 to 8, while half-integral J ranges from $\frac{1}{2}$ to $1\frac{1}{2}$. If only electric fields constitute the perturbation on the free ion, the appropriate group may be obtained from table 5 or from crystallographic considera-

tions. For a pure magnetic field, we shall consider only a uniform field with symmetry $C_{\infty v}$. In case both fields are present, the appropriate group is the group of those symmetry operations common to the electric and magnetic fields. Unless the magnetic field is parallel to one of the symmetry axes of the crystal field, there will in general be no symmetry common to both perturbations.

In accordance with our previous discussions, the physical implication of this reduction is that all one dimensional reps appearing in the reduction represent states arising from the free ion level J , which are not related by symmetry to any of the other states so arising. These states, then, are either nondegenerate or "accidentally" degenerate, the latter being due to numerical values of the parameters hence very improbable in any actual case. Any two-dimensional rep represents a pair of states from the original $2J+1$ which are essentially degenerate, being related by symmetry operations, but not degenerate with the other $2J-1$ states. Similarly for any reps of higher dimension. If a particular rep appears only once in the reduction the state or states it represents may be uniquely associated with a particular combination of the original $2J+1$ states. If the rep appears more than once there will be two or more states of the original $2J+1$ having the same symmetry properties, and the appropriate states in the crystal may be a linear combination of these. This can only be determined from the numerical parameters involved. These considerations are modified by Kramers degeneracy to be considered in detail in section 4.2. Certain pairs of reps are noted in table 9 as being Kramers conjugate states. These states, even though belonging to different reps, are degenerate if the perturbing field is purely electrostatic.

It is often of considerable interest to follow the behavior of a given level during the introduction of perturbing fields of increasingly lower symmetry. For example, one may wish to see the effect on the levels under D_{3d} symmetry of a reduction of symmetry to C_{3v} . This may be accomplished by the correlation table, table 10. Since most of the groups with which we are presently concerned are subgroups of $D_{\infty h}$, we start with the reps of $D_{\infty h}$. For each group of lower symmetry, the designation of the level or levels is given in the appropriate column. Again, the pure rotation groups are not given separately, requiring only the suppression of "g"- "u" in the associated holohedric group. The procedure involved here is the application to the characters of the reps of $D_{\infty h}$ precisely the same procedures as were applied to the rotation-inversion group characters in constructing Table 9. Notice that here, in many cases, the rep of $D_{\infty h}$ is still a rep of the subgroup, since the levels frequently remain twofold degenerate.

For an example, let us consider the state 2_u under the influence of a crystal field of symmetry

TABLE 9. Type and number of perturbed levels

		0		1		2		3		4		5		6		7		8	
		g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u
D _{∞h}	A _{1g}	1				1				1				1				1	
	A _{1u}		1				1				1				1				1
	A _{2g}			1				1				1				1			
	A _{2u}				1				1				1				1		
	E _{1g}			1		1		1		1		1		1		1		1	
	E _{1u}				1		1		1		1		1		1		1		1
	E _{2g}					1		1		1		1		1		1		1	
	E _{2u}						1		1		1		1		1		1		1
	E _{3g}							1		1		1		1		1		1	
	E _{3u}								1		1		1		1		1		1
	E _{4g}									1		1		1		1		1	
	E _{4u}										1		1		1		1		1
	E _{5g}											1		1		1		1	
	E _{5u}												1		1		1		1
	E _{6g}													1		1		1	
	E _{6u}														1		1		1
E _{7g}															1		1		
E _{7u}																1		1	
E _{8g}																	1		
E _{8u}																		1	
C _{∞v}	A ₁	1		1		1		1		1		1		1		1		1	
	A ₂		1		1		1		1		1		1		1		1		1
	E ₁			1		1		1		1		1		1		1		1	
	E ₂				1		1		1		1		1		1		1		1
	E ₃						1		1		1		1		1		1		1
	E ₄								1		1		1		1		1		1
	E ₅										1		1		1		1		1
	E ₆												1		1		1		1
E ₇															1		1		
E ₈																1		1	
C _{∞h}	A _g	1		1		1		1		1		1		1		1		1	
	A _u		1		1		1		1		1		1		1		1		1
	C _{±1g}			1		1		1		1		1		1		1		1	
	C _{±1u}				1		1		1		1		1		1		1		1
	C _{±2g}				1		1		1		1		1		1		1		1
	C _{±2u}					1		1		1		1		1		1		1	1
	C _{±3g}						1		1		1		1		1		1		1
	C _{±3u}							1		1		1		1		1		1	1
	C _{±4g}								1		1		1		1		1		1
	C _{±4u}									1		1		1		1		1	1
	C _{±5g}										1		1		1		1		1
	C _{±5u}											1		1		1		1	1
	C _{±6g}												1		1		1		1
	C _{±6u}													1		1		1	1
	C _{±7g}														1		1		1
	C _{±7u}															1		1	1
C _{±8g}																	1		
C _{±8u}																		1	

Note: The states C_{ng}, C_{-ng}, and C_{nu}, C_{-nu}, are Kramers conjugate states.

TABLE 9. Type and number of perturbed levels—Continued

	0		1		2		3		4		5		6		7		8	
	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u
	D_{4h}	1	1	1	1	1	1	1	1	1	1	1	1	2	2	1	1	2
A_{1g}																		
A_{1u}																		
A_{2g}			1				1					1	1	1	2	1	1	2
A_{2u}				1				1					1	1		2		1
B_{1g}							1		1		1	1	1	1	1	1	1	1
B_{1u}							1		1	1	1	1	1	1	1	1	1	1
B_{2g}								1		1	1	1	1	1	1	1	1	1
B_{2u}								1		1	1	1	1	1	1	1	1	1
E_{1g}			1		1		1		1		2	2	2	3	3	3	3	3
E_{1u}				1		1		1		2	2	2	2	2	3	3	3	3
E_{2g}					1		1		2		2	2	2	2	2	2	3	3
E_{2u}					1		1		2		2	2	2	2	2	2	3	3
C_{4v}	1	1	1	1	1	1	1	1	1	1	1	2	2	1	2	2	2	1
A_1																		
A_2																		
B_1																		
B_2																		
E_1			1	1	1	1	1	1	2	2	2	2	2	3	3	3	3	3
E_2					1	1	1	1	2	2	2	2	2	2	2	3	3	3
D_{3h}	1	1	1	1	1	1	1	1	1	1	1	2	2	1	2	2	2	1
A'_1																		
A'_2																		
A''_1			1		1		1		1		1	1	1	2	1	1	1	2
A''_2				1		1		1	1	1	1	1	1	1	2	1	1	1
E'			1	1	1	1	1	2	2	2	2	2	2	3	3	3	3	3
E''					1	1	1	1	2	2	2	2	2	3	2	3	3	3
D_{4h}	1	1	1	1	1	1	1	2	2	1	2	2	1	1	3	3	3	2
A_{1g}																		
A_{1u}																		
A_{2g}			1				1	1	2	1	2	1	2	2	2	2	2	2
A_{2u}				1				1	1		2	1	1	2	2	2	2	2
B_{1g}					1		1	1	1	1	2	2	2	2	2	2	2	2
B_{1u}						1	1	1	1	1	1	2	2	2	2	2	2	2
B_{2g}					1		1	1	1	1	2	2	2	2	2	2	2	2
B_{2u}						1	1	1	1	1	1	2	2	2	2	2	2	2
E_g			1	1	1	2	2	2	2	3	3	3	4	4	4	4	4	4
E_u					1	1	2	2	2	2	3	3	3	4	4	4	4	4
C_{4v}	1	1	1	1	1	1	1	2	2	1	2	2	1	2	3	3	3	2
A_1																		
A_2																		
B_1					1	1	1	1	1	1	2	2	2	2	2	2	2	2
B_2					1	1	1	1	1	1	2	2	2	2	2	2	2	2
E			1	1	1	2	2	2	2	3	3	3	4	4	4	4	4	4
D_{3d}	1	1	1	1	1	1	1	2	2	1	2	2	1	2	3	3	3	2
A_1																		
A_2																		
B_1			1		1		1	1	2	1	2	2	2	2	2	2	2	3
B_2				1		1		1	1	1	2	2	1	2	2	2	2	2
E			1	1	1	2	2	2	2	3	3	3	4	4	4	4	4	4

TABLE 9. Type and number of perturbed levels—Continued

		0		1		2		3		4		5		6		7		8	
		g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u
D _{3d}	A _{1g}	1				1		1		2		1		3		2		3	
	A _{1u}		1				1		1		2		1		3		2		3
	A _{2g}			1				2		1		2		2		3		2	
	A _{2u}				1				2		1		2		2		3		2
E _g	E _g			1		2		2		3		4		4		5		6	
	E _u				1		2		2		3		4		4		5		6
C _{3g}	A ₁	1		1		1		1		2		1		2		2		3	
	A ₂		1		1		2		2		1		2		3		3		3
	E ₁			1		2		2		3		4		4		5		6	
D _{2h}	A _{1g}	1				2		1		3		2		4		3		5	
	A _{1u}		1				2		1		3		2		4		3		5
	A _{2g}			1		1		2		2		3		3		4		4	
	A _{2u}				1		1		2		2		3		3		4		4
	B _{1g}			1		1		2		2		3		3		4		4	
	B _{1u}				1		1		2		2		3		3		4		4
B _{2g}	B _{2g}			1		1		2		2		3		3		4		4	
	B _{2u}				1		1		2		2		3		3		4		4
C _{2g}	A ₁	1		1		2		1		2		2		3		3		4	
	A ₂		1		1		2		2		3		3		4		4		5
	B ₁			1		1		2		2		3		3		4		4	
	B ₂			1		1		2		2		3		3		4		4	
C _{4h}	A _g	1		1		1		1		1		1		3		3		3	
	A _u		1		1		1		1		1		1		3		3		3
	B _g						2		2		2		2		2		2		2
	B _u							2		2		2		2		2		2	
	C _{1g}			1		1		1		1		2		2		3		3	
	C _{1u}				1		1		1		1		2		2		3		3
	C _{2g}					1		1		2		2		2		2		3	
	C _{2u}						1		1		2		2		2		2		3
	C _{4g}					1		1		2		2		2		2		3	
C _{4u}						1		1		2		2		2		2		3	
C _{5g}	C _{5g}			1		1		1		1		2		2		3		3	
	C _{5u}				1		1		1		1		2		2		3		3
Note: The states (C _{1g} , C _{3g}), (C _{1u} , C _{3u}), (C _{2g} , C _{4g}), (C _{2u} , C _{4u}) are Kramers conjugate states.																			
C _{4h}	A'	1		1		1		1		2		1		2		3		3	
	A''		1		1		1		2		1		2		3		3		3
	C' ₁			1		1		1		1		2		2		2		3	
	C'' ₁				1		1		1		1		2		2		3		3
C' ₂	C' ₂			1		1		1		1		2		2		2		3	
	C'' ₂				1		1		1		1		2		2		3		3

Note: The states (C'₁, C''₁) and (C'₂, C''₂) are Kramers conjugate states.

TABLE 9. Type and number of perturbed levels—Continued

	0		1		2		3		4		5		6		7		8	
	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u
	S_0	1	1	1	1	1	1	3	3	3	3	3	3	5	5	5	5	5
A_g A_u																		
C_{1g} C_{1u}			1	1	2	2	2	2	3	3	4	4	4	4	5	5	6	6
C_{2g} C_{2u}			1	1	2	2	2	2	3	3	4	4	4	4	5	5	6	6

Note: The states (C_{1g} , C_{2g}) and (C_{1u} , C_{2u}) are Kramers conjugate states.

C_{13}	A_g	1	1	1	1	1	1	3	3	3	3	3	3	3	5	5
	A_u															
	B_g				2	2	2	2	2	2	2	2	4	4	4	4
	B_u					2	2	2	2	2	2	2	4	4	4	4
	C_{1g}		1	1	1	1	2	2	2	3	3	3	3	4	4	4
	C_{1u}			1	1	1	2	2	2	2	3	3	3	4	4	4
	C_{2g}		1	1	1	1	2	2	2	3	3	3	4	4	4	4
	C_{2u}				1	1	2	2	2	2	3	3	3	4	4	4

Note: The states (C_{1g} , C_{2g}) and (C_{1u} , C_{2u}) are Kramers conjugate states.

S_1	A	1	1	1	2	1	2	3	2	3	2	3	4	3	4	5	4
	B		1	1	1	2	1	2	3	2	3	4	3	4	3	4	5
	C_1			1	1	1	1	2	2	2	3	3	3	4	4	4	4
	C_2			1	1	1	1	2	2	2	3	3	3	4	4	4	4

Note: The states C_1 and C_2 are Kramers conjugate states.

C_{2A}	A_g	1	1	3	3	5	5	5	7	7	7	9	9
	A_u												
	B_g		2	2	3	4	4	6	6	6	8	8	8
	B_u			2	2	4	4	6	6	6	8	8	8

C_3	A'	1	1	2	3	3	4	5	4	5	6	7	6	7	8	9	8
	A''			2	3	4	3	4	5	6	5	6	7	8	7	8	9

O_A	A_{1g}	1	1					1	1			1	1			1	1
	A_{1u}																
	A_{2g}						1					1	1	1	1		
	A_{2u}							1					1	1	1		
	E_g				1			1	1	1	1	1	1	1	2	2	2
	E_u					1			1	1	1	1	1	1	1	2	2
	T_{1g}		1				1	1	1	2	2	1	1	2	2	2	2
	T_{1u}			1				1	1	2	2	1	1	2	2	2	2
	T_{2g}				1	1		1	1	1	1	2	2	2	2	2	2
	T_{2u}					1		1	1	1	1	2	2	2	2	2	2

TABLE 9. Type and number of perturbed levels—Continued

		0		1		2		3		4		5		6		7		8	
		g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u
T _g	A ₁	1						1	1	1				1	1	1	1	1	
	A ₂		1								1			1	1				1
	E					1	1			1	1	1	1	1	1	1	1	2	2
	T ₁ T ₂			1	1	1	1	1	1	1	1	2	1	1	2	2	2	2	2
T _h	A _g A _u	1	1					1	1	1	1			2	2	1	1	1	1
	C _{1g} C _{1u}					1	1			1	1	1	1	1	1	1	1	2	2
	C _{2g} C _{2u}					1	1			1	1	1	1	1	1	1	1	2	2
	T _g T _u			1	1	1	1	2	2	2	2	3	3	3	3	4	4	4	4

Note: The states (C_{1g}, C_{1u}) and (C_{2g}, C_{2u}) are Kramers conjugate states.

		$\frac{1}{2}$		$\frac{3}{2}$		$\frac{5}{2}$		$\frac{7}{2}$		$\frac{9}{2}$		$\frac{11}{2}$		$\frac{13}{2}$		$\frac{15}{2}$	
		g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u
D _{∞h}	² S _{1g}	1		1		1		1		1		1		1		1	
	² S _{1u}		1		1		1		1		1		1		1		1
	² S _{3g}			1		1		1		1		1		1		1	
	² S _{3u}				1		1		1		1		1		1		1
	² S _{5g}					1		1		1		1		1		1	
	² S _{5u}						1		1		1		1		1		1
	² S _{7g}							1		1		1		1		1	
	² S _{7u}								1		1		1		1		1
	² S _{9g}									1		1		1		1	
	² S _{9u}										1		1		1		1
	² S _{11g}											1		1		1	
	² S _{11u}												1		1		1
	² S _{13g}													1		1	
	² S _{13u}														1		1
C _{∞v}	² S ₁	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	² S ₃			1	1	1	1	1	1	1	1	1	1	1	1	1	1
	² S ₅					1	1	1	1	1	1	1	1	1	1	1	1
	² S ₇							1	1	1	1	1	1	1	1	1	1
	² S ₉									1	1	1	1	1	1	1	1
	² S ₁₁											1	1	1	1	1	1
	² S ₁₃													1	1	1	1
	² S ₁₅															1	1

TABLE 9. Type and number of perturbed levels—Continued

	$\frac{1}{2}$		$\frac{3}{2}$		$\frac{5}{2}$		$\frac{7}{2}$		$\frac{9}{2}$		$\frac{11}{2}$		$\frac{13}{2}$		$\frac{15}{2}$		
	g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u	
$C_{\infty h}$	$S_{\pm 1g}$	1		1		1		1		1		1		1		1	
	$S_{\pm 1u}$		1		1		1		1		1		1		1		1
	$S_{\pm 3g}$			1		1		1		1		1		1		1	
	$S_{\pm 3u}$				1		1		1		1		1		1		1
	$S_{\pm 5g}$					1		1		1		1		1		1	
	$S_{\pm 5u}$						1		1		1		1		1		1
	$S_{\pm 7g}$						1		1		1		1		1		1
	$S_{\pm 7u}$							1		1		1		1		1	
	$S_{\pm 9g}$								1		1		1		1		1
	$S_{\pm 9u}$									1		1		1		1	
	$S_{\pm 11g}$										1		1		1		1
	$S_{\pm 11u}$											1		1		1	
	$S_{\pm 13g}$												1		1		1
	$S_{\pm 13u}$													1		1	
	$S_{\pm 15g}$														1		1
$S_{\pm 15u}$															1		

Note: The states (S_{ng} , S_{-ng}) and (S_{nu} , S_{-nu}) are Kramers conjugate states.

D_{6h}	$D_{3/2g}$	1		1		1		1		1		2		2		3		3		3		3					
	$D_{3/2u}$		1		1		1		1		1		1		1		3		3		3		3				
	${}^2S_{1g}$			1		1		1		1		2		2		2		2		2		2		3			
	${}^2S_{1u}$				1		1		1		1		2		2		2		2		2		2		3		
D_{6g}	${}^2S_{2g}$					1		2		2		2		2		2		2		2		2		2		2	
	${}^2S_{2u}$						1		2		2		2		2		2		2		2		2		2		
D_{3h}	$D_{3/2g}$	1		1		1		1		1		2		2		2		2		2		2		2		2	
	2S_1			1		1		1		1		2		2		2		2		2		2		2		2	
	2S_2		1		1		1		2		1		2		1		2		2		2		3		2		3
D_{4h}	$D_{3/2g}$	1		1		1		2		3		3		3		3		3		3		4		4		4	
	$D_{3/2u}$		1		1		1		2		2		3		3		3		3		3		3		3		4
	2S_g			1		2		2		2		2		3		3		4		3		4		4		4	
D_{4g}	2S_u				1		2		2		2		2		2		3		3		4		4		4		4
	$D_{3/2g}$	1		1		1		2		2		3		3		3		3		3		4		4		4	
D_{2d}	$D_{3/2g}$	1		1		1		2		2		3		3		3		3		3		4		4		4	
	2S		1		1		2		1		2		2		2		3		3		3		4		3		4
D_{3d}	$D_{3/2g}$	1		1		2		3		3		4		5		5		5		5		5		5		5	
	$D_{3/2u}$		1		1		2		3		3		4		5		5		5		5		5		5		5
	S_{1g}			1		1		1		1		2		2		2		2		2		2		2		3	
	S_{1u}				1		1		1		1		2		2		2		2		2		2		2		3
	S_{3g}			1		1		1		1		2		2		2		2		2		2		2		2	

Note: The states (S_{1g} , S_{3g}) and (S_{1u} , S_{3u}) are Kramers conjugate states.

TABLE 9. Type and number of perturbed levels—Continued

		$\frac{1}{2}$		$\frac{3}{2}$		$\frac{5}{2}$		$\frac{7}{2}$		$\frac{9}{2}$		$\frac{11}{2}$		$\frac{13}{2}$		$\frac{15}{2}$	
		g	u	g	u	g	u	g	u	g	u	g	u	g	u	g	u
C _{3v}	D _{3/2}	1	1	1	1	2	2	3	3	3	3	4	4	5	5	5	5
	S ₁			1	1	1	1	1	1	2	2	2	2	2	2	3	3
	S ₃			1	1	1	1	1	1	2	2	2	2	2	2	3	3

Note: The states (S₁, S₃) are Kramers conjugate states.

C _{4v}	S _{1g}	1		1		1		2		3		3		3		4	
	S _{1u}		1		1		1		2		3		3		3		4
	S _{3g}			1		2		2		2		3		3		4	
	S _{3u}			1		2		2		2		3		3		4	
	S _{5g}			1		1		2		2		2		3		4	
	S _{5u}	1	1	1	1	1	1	2	2	3	3	3	3	3	4	4	4

Note: The states (S_{1g}, S_{1u}), (S_{3g}, S_{3u}), (S_{5g}, S_{5u}) are Kramers conjugate states.

S ₄	S ₁		1	1	1	2	1	2	2	2	3	3	3	4	3	4	4
	S ₃	1		1	1	1	2	2	2	3	2	3	3	3	4	4	4
	S ₅	1		1	1	1	2	2	2	3	2	3	3	3	4	4	4
	S ₇		1	1	1	2	1	2	2	2	3	3	3	4	3	4	4

Note: The states (S₁, S₇) and (S₂, S₃) are Kramers conjugate states.

D _{2d}	D _{3/2g}	1		2		3		4		5		6		7		8	
	D _{3/2u}		1		2		3		4		5		6		7		8
C _{6v}	S _{1g}	1		1		1		1		1		2		3		3	
	S _{1u}		1		1		1		1		1		2		2		3
	S _{3g}			1		1		1		1		2		2		2	
	S _{3u}			1		1		1		1		2		2		2	
	S _{5g}					1		2		2		2		2		2	
	S _{5u}					1		1		2		2		2		2	
	S _{7g}					1		1		2		2		2		2	
	S _{7u}					1		1		2		2		2		2	
S _{9g}			1		1		1		1		2		2		2		
S _{9u}			1		1		1		1		2		2		2		
S _{11g}	1		1		1		1		1		1		2		3		
S _{11u}		1		1		1		1		1		1		2		3	

Note: The states (S_{1g}, S_{11g}), (S_{1u}, S_{11u}), (S_{3g}, S_{9g}), (S_{3u}, S_{9u}), (S_{5g}, S_{7g}), (S_{5u}, S_{7u}) are Kramers conjugate states.

C _{3h}	S ₁	1		1		1		1		2		2		2		3	
	S ₃			1		1		1		1		2		2		2	
	S ₅		1		1	1	1	2	1	2	1	2	2	2	3	2	3
	S ₇		1		1	1	1	2	1	2	1	2	2	2	3	2	3
	S ₉			1	1	1	1	1	1	2	2	2	2	2	2	3	3
S ₁₁	1		1		1		1		1		2		2		3		

Note: The states (S₁, S₁₁), (S₃, S₉), (S₅, S₇) are Kramers conjugate states.

TABLE 9. Type and number of perturbed levels—Continued

	$\frac{1}{2}$		$\frac{3}{2}$		$\frac{5}{2}$		$\frac{7}{2}$		$\frac{9}{2}$		$\frac{11}{2}$		$\frac{13}{2}$		$\frac{15}{2}$		
	<i>g</i>	<i>u</i>	<i>g</i>	<i>u</i>	<i>g</i>	<i>u</i>	<i>g</i>	<i>u</i>	<i>g</i>	<i>u</i>	<i>g</i>	<i>u</i>	<i>g</i>	<i>u</i>	<i>g</i>	<i>u</i>	
	S_4																
S_{1g}	1		1		2		3		3		4		5		5		5
S_{1u}		1		1		2		3		3		4		5		5	5
S_{3g}			2		2		2		4		4		4		6		6
S_{3u}				2		2		2		4		4		4		6	6
S_{5g}	1		1		2		3		3		4		5		5		5
S_{5u}		1		1		2		3		3		4		5		5	5

Note: The states (S_{1g} , S_{3g}), (S_{1u} , S_{3u}) are Kramers conjugate states. The states (S_{3g} , S_{3u}) appear an even number of times as Kramers conjugate pairs.

C_{2v}	D_{3g}	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8
C_{2h}	S_{1g}	1		2		3		4		5		6		7		8	
	S_{1u}		1		2		3		4		5		6		7		8
	S_{3g}	1		2		3		4		5		6		7		8	
	S_{3u}		1		2		3		4		5		6		7		8

Note: The states (S_{1g} , S_{3g}) and (S_{1u} , S_{3u}) are Kramers conjugate states.

O_h	D_{3g}	1						1		1		1		1		1	
	D_{3u}		1						1		1		1		1		1
	D_{5g}			1		1		1		2		2		2		3	
	D_{5u}				1		1		1		2		2		2		3
	$2S_g$					1		1				1		2		2	
	$2S_u$						1		1			1		2		2	
T_d	D_{3g}	1		1		1		1		1		1		1		1	
	D_{3u}		1		1		1		1		2		2		2		3
	$2S_g$					1		1		1		1		2		2	
	$2S_u$						1		1		1		2		2		3
T_h	D_{3g}	1				1		2		1		2		3		2	
	D_{3u}		1				1		2		1		2		3		2
	$2S_{1g}$			1		1		1		2		2		2		3	
	$2S_{1u}$				1		1		1		2		2		2		3
	$2S_{2g}$			1		1		1		2		2		2		3	
	$2S_{2u}$				1		1		1		2		2		2		3

Note: The states ($2S_{1g}$, $2S_{2g}$) and ($2S_{1u}$, $2S_{2u}$) are Kramers conjugate pairs of states.

TABLE 10—Correlation tables

D _{00A}	C ₀₀	C _{00A}	D _{0A}	D _{0A}	D _{0A}	C _{0A}	D _{0A}	D _{0A}	C _{0A}	D _{0A}	D _{0A}	D _{0A}	C _{0A}	C _{0A}	S ₁	C _{0A}	C _{0A}	C _{0A}	
A _{1E} A _{1u}	A ₁ A ₂	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}	A B	A ₁ A ₂	A _{1E} A _{1u}	A _{1E} A _{1u}	A _{1E} A _{1u}
A _{2E} A _{2u}	A ₂ A ₁	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}	A B	A ₂ A ₁	A _{2E} A _{2u}	A _{2E} A _{2u}	A _{2E} A _{2u}
E _{1E} E _{1u}	E ₁ E ₁	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	E _{1E} E _{1u}	C ₁ C ₁ C ₁ C ₁	E E	C _{1E} C _{1u} C _{1u}	C _{1E} C _{1u} C _{1u}	C _{1E} C _{1u} C _{1u}
E _{2E} E _{2u}	E ₂ E ₂	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}	B B B B B	E E	E _{2E} E _{2u}	E _{2E} E _{2u}	E _{2E} E _{2u}
E _{3E} E _{3u}	E ₃ E ₃	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}	C ₂ C ₂ C ₂ C ₂	E E	E _{3E} E _{3u}	E _{3E} E _{3u}	E _{3E} E _{3u}
E _{4E} E _{4u}	E ₄ E ₄	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}	C ₃ C ₃ C ₃ C ₃	E E	E _{4E} E _{4u}	E _{4E} E _{4u}	E _{4E} E _{4u}
E _{5E} E _{5u}	E ₅ E ₅	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}	C ₄ C ₄ C ₄ C ₄	E E	E _{5E} E _{5u}	E _{5E} E _{5u}	E _{5E} E _{5u}
E _{6E} E _{6u}	E ₆ E ₆	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}	C ₅ C ₅ C ₅ C ₅	E E	E _{6E} E _{6u}	E _{6E} E _{6u}	E _{6E} E _{6u}
E _{7E} E _{7u}	E ₇ E ₇	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}	C ₆ C ₆ C ₆ C ₆	E E	E _{7E} E _{7u}	E _{7E} E _{7u}	E _{7E} E _{7u}
E _{8E} E _{8u}	E ₈ E ₈	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}	C ₇ C ₇ C ₇ C ₇	E E	E _{8E} E _{8u}	E _{8E} E _{8u}	E _{8E} E _{8u}
E _{9E} E _{9u}	E ₉ E ₉	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}	C ₈ C ₈ C ₈ C ₈	E E	E _{9E} E _{9u}	E _{9E} E _{9u}	E _{9E} E _{9u}

D_{2h} , where table 5 shows that potential coefficients $C_2^0 C_4^0 C_6^0$; $C_2^0 C_4^0 C_6^0$; $C_4^0 C_6^0$ and C_6^0 are present, and table 9 shows that there are two states A_{1u} and one each of A_{2u} , B_{1u} , and B_{2u} , indicating that the original fivefold degeneracy of the state 2_u has been destroyed. If the terms C_2^0 , C_4^0 , and C_6^0 are more significant than the rest, an initial approximation using these terms only leads to D_{2h} symmetry, with states A_{1u} , E_{1u} , and E_{2u} . The inclusion of C_2^0 will reduce the symmetry to D_{2h} , but the states retain their designation. On the other hand, the inclusion of C_4^0 will reduce the symmetry to D_{2h} , leave the A_{1u} level unaffected, leave the E_{1u} level unaffected save for a redesignation to E_u , and cause the E_{2u} level to split to B_{1u} and B_{2u} . If we now include all terms, we see that E_u of D_{2h} splits to B_{1u} and B_{2u} of D_{2h} , and that the states B_{1u} and B_{2u} of D_{2h} become A_{1u} and A_{2u} , respectively, in D_{2h} . This A_{1u} state can interact with the A_{1u} state from the original D_{2h} approximation, so that a rigid correlation with the states of D_{2h} has been lost.

Table 10 has been so constructed that the same process may be followed for still lower order groups. For every column in the table, its subgroups are found further to the right in the table (or in the same column without the "g"-"u"), and the designation of any level under increasingly lower symmetry may be followed across the table. Entries of the same line have significance only with respect to the allowed subgroups of any given group. Thus, there is no correlation between parallel entries in D_{2h} and D_{2h} , though either may be correlated with D_{2h} . All splittings of double-degenerate reps have been correlated to ensure consistency. Since the basic entries of the table are the reps of D_{2h} , any desired rep of some other group must be found from the table by inspection

but its behavior under lower symmetry does not depend upon the rep of D_{2h} from which it arose. In a manner of speaking, the columns of table 10 are first differences of the entries of table 9.

Table 11 shows the reduction of the reps of the cubic group into the reps of their own cubic subgroups, and into reps of the noncubic groups. Here, we have considered the two possibilities which may arise in destroying cubic symmetry. Thus, the group O may be reduced to D_4 by destroying the threefold axes, or to D_3 by destroying the fourfold axes. In either case, the principal axis remaining is considered to be the principal axis of symmetry.

While our discussion has been that appropriate to the weak-field case (sections 3.3, 3.4), the results are immediately applicable to both the intermediate field case and the strong field case. In the former, the external perturbation is stronger than the spin-orbit interaction. Consequently, instead of reducing the representation $D_L \times D_S$ into those reps of the group leaving invariant the spin-orbit term in the Hamiltonian, i.e., D_J , we reduce the representation into those reps of the group leaving invariant the crystal field potential V . Since the spin is not included in the Hamiltonian at this stage of the approximation, we reduce D_L only into reps X_i . This may be done by using table 9 for L instead of J (here, the double groups will not appear, since L must be integral). The rep for the wave function, including spin, will then be $X_i \times D_S$. We shall parallel spectroscopic notation for such a state by using the appropriate symbol from table 8 for the rep X_i , with an anterior superscript $2S+1$ for D_S , the whole being placed in parentheses. If the spin-orbit interaction is now intro-

TABLE 11. Reduction of the cubic groups

O_h	T_h	T_d	D_{4h}	D_{2d}	T_h	D_{2h}	S_6	T_d	T	D_{2d}	C_{3v}
A_{1g}	A_g	A_1	A_{1g}	A_{1g}	A_g	A_{1g}	A_g	A_1	A	A_1	A_1
A_{1u}	A_u	A_2	A_{1u}	A_{1u}	A_u	A_{1u}	A_u	A_2	A	B_1	A_2
A_{2g}	A_g	A_2	B_{1g}	A_{2g}	C_{1g}	A_{2g}	C_{1g}	E	$C_1 C_2$	$A_1 B_1$	E
A_{2u}	A_u	A_1	B_{1u}	A_{2u}	C_{1u}	A_{1u}	C_{1u}	T_1	T	$A_2 E$	$A_2 E$
E_g	$C_{1g} C_{2g}$	E	$A_{1g} B_{1g}$	E_g	C_{2g}	A_{1g}	C_{2g}	T_2	T	$B_2 E$	$A_1 E$
E_u	$C_{1u} C_{2u}$	E	$A_{1u} B_{1u}$	E_u	C_{2u}	A_{1u}	C_{2u}	$D_{3/2}$	$D_{3/2}$	$D_{3/2}$	$D_{3/2}$
T_{1g}	T_g	T_1	$E_g A_{2g}$	$E_g A_{2g}$	T_g	$A_{2g} B_{1g} B_{2g}$	$A_g C_{1g} C_{2g}$	$2S$	$D_{3/2}$	$D_{3/2}$	$D_{3/2}$
T_{1u}	T_u	T_2	$E_u A_{2u}$	$E_u A_{2u}$	T_u	$A_{2u} B_{1u} B_{2u}$	$A_u C_{1u} C_{2u}$	$2S$	$2S_1 2S_2$	$D_{3/2} 2S$	$D_{3/2} S_1 S_3$
T_{2g}	T_g	T_2	$E_g B_{2g}$	$E_g A_{1g}$	$D_{3/2g}$	$D_{3/2g}$	$S_{1g} S_{5g}$				
T_{2u}	T_u	T_1	$E_u B_{2u}$	$E_u A_{1u}$	$D_{3/2u}$	$D_{3/2u}$	$S_{1u} S_{5u}$				
$D_{3/2g}$	$D_{3/2g}$	$D_{3/2}$	$D_{3/2g}$	$D_{3/2g}$	$2S_{1g}$	$D_{3/2g}$	$S_{1g} S_{3g}$				
$D_{3/2u}$	$D_{3/2u}$	$2S$	$D_{3/2u}$	$D_{3/2u}$	$2S_{1u}$	$D_{3/2u}$	$S_{1u} S_{3u}$				
$2S_g$	$D_{3/2g}$	$2S$	$2S_g$	$D_{3/2g}$	$2S_{2g}$	$D_{3/2g}$	$S_{3g} S_{5g}$				
$2S_u$	$D_{3/2u}$	$D_{3/2}$	$2S_u$	$D_{3/2u}$	$2S_{2u}$	$D_{3/2u}$	$S_{3u} S_{5u}$				
$D_{3/2g}$	$2S_{1g} 2S_{2g}$	$D_{3/2}$	$D_{3/2g} 2S_g$	$D_{3/2g} S_{1g} S_{3g}$							
$D_{3/2u}$	$2S_{1u} 2S_{2u}$	$D_{3/2}$	$D_{3/2u} 2S_u$	$D_{3/2u} S_{1u} S_{3u}$							

duced, the rep $X_i \times D_3$ is now merely a representation, so that it must be reduced according to the overall symmetry of the problem, which is still that of the crystal field V . If X_i is one-dimensional, the splitting which arises is that of the spin states only through D_3 , and the angular momentum is said to be quenched; if X_i is of dimensionality greater than 1 but less than $2L+1$, the orbital angular momentum is partially quenched. This verbally depicts the effect on the orbital angular momentum of the strong crystal potential field V , which destroys (X_i one-dimensional) or limits ($1 < X_i < 2L+1$) any further influence on the angular momentum either by spin-orbit interaction or by a moderate external magnetic field. Still another intuitive view is that the crystal potential partially or completely fixes the orientation of the angular momentum in the lattice, so that these other influences have no further effect.

As an example, let us consider the 7F_2 level of the europium trivalent ion, the symbol indicating a reduction of $D_3 \times D_3$ to D_2 by spin-orbit interaction. A weak perturbing field D_{4h} will (table 9) split this 7F_2 level into states described as $({}^7F)A_{1g}$; $({}^7F)B_{1g}$; $({}^7F)B_{2g}$; and $({}^7F)E_g$. An intermediate case field D_{4h} will split the 7F term (table 9 for 7F_2) into $({}^7A_2)$; $({}^7B_{1g})$; $({}^7B_{2g})$; and $2({}^7E_g)$, a total of $(2S+1)(2L+1)=49$ states. The introduction of the spin-orbit coupling will now cause each of these to be split further. Consider the level $({}^7A_2)$. The rep A_{2g} in D_{4h} has the character (1, 1, 1, -1, -1, 1, 1, -1, -1), while the representation from D_3 has the character (parity even, since it is a spin representation) (7, -1, -1, -1, -1, 7, -1, -1, -1), so the representation $A_{2g} \times D_3$ has the character (7, -1, -1, 1, 1, 7, -1, -1, 1) and is reducible into A_{1g} ; B_{1g} ; B_{2g} ; and $2E_g$. The final states may be designated $({}^7A_{2g})A_{1g}$; $({}^7A_{2g})B_{1g}$; $({}^7A_{2g})B_{2g}$; and $2({}^7A_{2g})E_g$. This notation parallels that of the usual $L-S$ coupling scheme. The labels inside the parentheses indicate the approximate transformation properties of the state for transformations of spin and orbit separately which leave invariant the Hamiltonian, while the final designation is the accurate specification of the behavior of the state under the simultaneous transformation of spin and orbit.

In the strong field case, we introduce the crystal field in our sequence of approximations (section 3.3) before considering the details of the electrostatic interaction of the electrons, retaining only the initial central field average $V'(r_i)$. We now specify our initial energy by giving only the configuration, and the reduction of the representations begins at this point. As a matter of notation, following Mulliken (1955), we shall designate the reps and characters for a single electron by lower case letters (e.g., a_{2g}). If there is but one electron, there is no distinction between the intermediate field and strong field cases, and we proceed as outlined for that case.

We shall illustrate the procedure for the case of

2 nonequivalent electrons, and then consider the restrictions imposed on equivalent electrons by the operation of the Pauli principle. We shall take the configuration pp in a field of C_{2v} symmetry. The threefold degeneracy of a p state (apart from spin) is completely lifted, and we obtain (table 9 for 1_u) a_1 ; b_1 ; and b_2 for each electron, so the configuration breaks up into $(a_1)(a_1)$; $(a_1)(b_2)$; $(b_2)(a_1)$; etc., a total of 9 possibilities corresponding to the original 9 orbital possibilities in pp . The spin transformation will be $D_{3/2} \times D_{3/2}$. The introduction of the details of electrostatic interaction cannot cause any further splitting of the orbital wave function, since all the states are one dimensional, though a relabeling may occur. The requirement for totally antisymmetric wave functions will require that the spin transformation $D_{3/2} \times D_{3/2}$ be reduced to D_0 and D_1 , giving the usual singlet and triplet states, respectively. Thus, $(a_1)(a_1)$ becomes $(a_1)(a_1)({}^1A_1)$ and $(a_1)(a_1)({}^3A_1)$; $(b_1)(b_2)$ becomes $(b_1)(b_2)({}^1A_2)$ and $(b_1)(b_2)({}^3A_2)$; etc. Note that this latter state is distinct from $(b_2)(b_1)({}^3A_2)$, which is also an allowed state in the reduction. The further introduction of spin-orbit interaction now requires the reduction of $X_i \times D_0$ for the singlet states, $X_i \times D_1$ for the triplet states, in accordance with the overall symmetry of the problem, which is still that of the external crystal field potential V . Consider $(b_1)(b_2)({}^3A_2)$. The character of A_2 is (1, 1, -1, -1), while the character of D_1 in C_{2v} (even parity for a spin transformation) is (from table 7) (3, -1, -1, -1), so the character to be reduced is (3, -1, 1, 1), giving A_1 , B_1 , and B_2 . We may write the final states, then, as $pp(b_1)(b_2)({}^3A_2)A_1$, or $pp(b_1)(b_2)({}^3A_2)B_1$, for example.

If the electrons are equivalent, i.e., p^2 , we write $(a_1)^2$ instead of $(a_1)(a_1)$, and we cannot distinguish $(a_1)(b_2)$ from $(b_2)(a_1)$. Furthermore, the Pauli principle requires the spin state of opposing spins for the orbital states $(a_1)^2$; $(b_1)^2$; and $(b_2)^2$, so that these appear as singlets only. The remainder of the orbital states may appear either as singlets or as triplets. Apart from these considerations, the development follows that of the nonequivalent electrons. A possible final state would be written $p^2(b_1)^2({}^1A_1)A_1$.

We have selected here a particularly simple case in that there is no orbital degeneracy present in the reduction of D_1 for a p electron. The requirements of the Pauli principle are then particularly easy to satisfy. If the reduction yields a two-dimensional rep, or remains three dimensional, the development must be followed more carefully. We introduce the procedure by considering, in the concepts of the present work, the familiar operation of the Pauli principle in labeling the levels of the configuration p^2 itself. We imagine a uniform magnetic field such that the resulting perturbation in the Hamiltonian, $-\mathbf{M} \cdot \mathbf{B}$, has the symmetry $C_{\infty v}$. The degeneracy of each p electron is lifted, and we have (table 9) the states a_u , c_{1u} and c_{-1u} for each electron. Because of the Pauli

principle, the state $(c_{1u})^2 C_{2g}$ occurs with anti-parallel spins only; i.e., a singlet state. We see that no state C_{2g} for n greater than 2 arises. We then infer from table 9 that there exists a singlet level $L=2$ in the configuration when the magnetic field is removed. Similarly, we deduce a triplet $L=1$ and a singlet $L=0$.

A similar process can be used if the crystal field leaves degeneracy in the 1-electron wave functions. The differences are two: first, it is generally more convenient to remove the degeneracy by the imaginary application of an electrostatic potential of a symmetry which is a subgroup of that actually present; second, the correlation of states in this additionally perturbed problem with those of the primary perturbation is not unique, and several alternatives must be considered.

Let us further consider p^2 , this time under the symmetry C_{4v} , where I_u reduces to a_1 and e . Our problem really lies in the study of $(e)^2$, since $(a_1)(e)$ can yield either triplet or singlet, and $(a_1)^2$ is necessarily a singlet. Without the considerations of the Pauli principle, $(e)^2$ yields A_1 , A_2 , B_1 , and B_2 , both triplet and singlet, and the Pauli principle will suppress certain of these terms. Since (e) , including spin, represents 4 states, we expect $\frac{4 \cdot 3}{2} = 6$ states altogether. Let us reduce the symmetry to C_{2v} ; (e) becomes $b_1 + b_2$, and the allowed orbital states are $(b_1)^2$; $(b_2)^2$; and $(b_1)(b_2)$. The first two are both A_1 and necessarily singlet, while the third is A_2 , either singlet or triplet, so we have our total of six states. If we examine table 10 for the relations between these states of C_{2v} and states of C_{4v} , we see that A_1 in the former implies either A_1 or B_1 in the latter. Since we have A_1 twice in C_{2v} and cannot have either A_1 or B_1 more than once from $(e)^2$ in C_{4v} , we conclude that both are present in C_{4v} as singlet terms. However, the singlet, or the triplet, A_2 implies either A_2 or B_2 , and we cannot decide which is the correct choice. If we reduce the symmetry to C_4 instead of C_{2v} , (e) reduces to (c_1) and (c_3) , yielding singlets $(c_1)^2$ and $(c_3)^2$, and singlet and triplet $(c_1)(c_3)$. The first two give B states, while the third gives A states. Examination of table 10 shows that B in C_4 implies either B_1 or B_2 in C_{4v} . Again, since we have two of the former, and are allowed at most one of each of the latter, we conclude the presence of singlet B_1 and singlet B_2 , together with the singlet A_1 previously deduced. The triplet state in C_{2v} implies either A_2 or B_2 in C_{4v} , while the triplet state in C_4 implies either A_1 or A_2 . Obviously, the only choice consistent with both is the triplet A_2 . We have, thus, 1A_1 ; 3A_2 ; 1B_1 ; and 1B_2 . The balance of the problem then follows previous discussion. In general, each problem must be considered as an individual case.

4.2. Kramers Degeneracy

There is one very important symmetry operation, in addition to spatial rotations and the

inversion, which should be considered at this time. This is the symmetry of "time-reversal". This was initially considered by Kramers (1929, 1930) in connection with the rotation of the plane of polarization of polarized light. The matter has been thoroughly considered by Wigner (1932) in the nonrelativistic case in which we are here interested, and more recently by Klein (1952). We shall outline here the principal concepts involved, referring the reader to the original papers for additional details.

Classically, the reversal of time in a mechanical system is accomplished by the reversal of the direction of all velocities (or momenta) by the substitution $\mathbf{r}' = \mathbf{r}$, $\mathbf{p}' = -\mathbf{p}$. Quantum mechanically, we assume that this is also time reversal, if a similar substitution $\mathbf{s}' = -\mathbf{s}$ is made for any spin angular momentum involved. In the case of a free atom, the Hamiltonian $H(\mathbf{r}, \mathbf{p}, \mathbf{s})$ involves \mathbf{p} and \mathbf{s} in kinetic energy terms with \mathbf{p}^2 and spin-orbit interaction terms with $\mathbf{p} \cdot \mathbf{s}$, both of which are invariant under the time-reversal operation. This invariance of the Hamiltonian remains under the influence of external perturbations of a purely electrostatic character, in which case the vector potential \mathbf{A} may be set equal to zero and the scalar potential $V(\mathbf{r})$ introduced. This invariance is destroyed by the application of magnetic fields—a term $\mathbf{p} \cdot \mathbf{A}$ arises in the Hamiltonian, which is linear in \mathbf{p} and changes sign on time reversal.

In general, then, we shall seek an operator \mathbf{K} such that $\mathbf{K}H(\mathbf{r}, \mathbf{p}, \mathbf{s})\mathbf{K}^{-1} = H(\mathbf{r}, -\mathbf{p}, -\mathbf{s})$, and $\mathbf{K}\Psi(\mathbf{r}, \mathbf{s}, t) = \Psi(\mathbf{r}, -\mathbf{s}, -t)$. Since the operator \mathbf{K} represents a transformation of our wave functions, we shall require it to be a unitary operator. If we

consider the spin-free theory, with $\Psi(\mathbf{r}, t) = \psi(\mathbf{r})e^{-\frac{iEt}{\hbar}}$, we see that the operation $t \rightarrow -t$ is accomplished merely by taking the complex conjugate, an operation we shall call \mathbf{K}_0 . Insofar as the spin-free theory is concerned, this is sufficient; the \mathbf{p} operator $-i\hbar\nabla$ becomes $-\mathbf{p} = +i\hbar\nabla$ upon taking the complex conjugate. Let us now consider the Pauli spin theory for one electron. The Pauli spin matrices

$$\mathbf{s}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \mathbf{s}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \mathbf{s}_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

are both unitary and Hermitian. The transformation $\mathbf{K}_0 \mathbf{s}_y \mathbf{K}_0^{-1} = -\mathbf{s}_y$, since \mathbf{s}_y is purely imaginary, but \mathbf{s}_x and \mathbf{s}_z , both real, are invariant under \mathbf{K}_0 . Hence, in order to effect the reversal of \mathbf{s}_x and \mathbf{s}_z , we seek yet another operator, which must also be a unitary operator \mathbf{U} , such that it commutes with \mathbf{r} , \mathbf{p} , \mathbf{s}_y , and \mathbf{K}_0 , but anticommutes with \mathbf{s}_x and \mathbf{s}_z so that $\mathbf{U} \mathbf{s}_x \mathbf{U}^{-1} = -\mathbf{s}_x$ and similarly for \mathbf{s}_z . This latter requirement is met by \mathbf{s}_y itself, or by some product of \mathbf{s}_y with a complex number a of modulus 1. To ensure commutation with \mathbf{K}_0 , the product $a\mathbf{s}_y$ must be real, which follows if $a=i$, so that $\mathbf{U} = i\mathbf{s}_y$, and $\mathbf{K} = \mathbf{K}_0 i\mathbf{s}_y$. Note that the operator \mathbf{U} is still unitary but no

longer Hermitian. For n electrons,

$$U = (i)^n (s_y)_1 (s_y)_2 (s_y)_3 \dots$$

Physically, it is clear that time reversal and any rotation-inversion operation commute. Mathematically, the inclusion of the operator \mathbf{K} in an extended group of the form $D_J \times I \times K$ cannot be accomplished since \mathbf{K} is not a linear operator. The basic rules for the operator \mathbf{K} were given by Wigner as:

1. $\mathbf{K}^2\psi = \mathbf{U}\mathbf{K}_0\mathbf{U}\mathbf{K}_0\psi = \mathbf{U}^2\psi = (-1)^n\psi$, since \mathbf{U} is real, $\mathbf{K}_0^2=1$, and $s_y^2=1$.
2. $\mathbf{K}(a\psi + b\phi) = a^*\mathbf{K}\psi + b^*\mathbf{K}\phi$. This shows the nonlinearity of \mathbf{K} .
3. $(\phi, \psi) = (\mathbf{K}_0\psi, \mathbf{K}_0\phi) = (\mathbf{U}\mathbf{K}_0\psi, \mathbf{U}\mathbf{K}_0\phi) = (\mathbf{K}\psi, \mathbf{K}\phi)$.

Applying (3) and (1), we have $(\mathbf{K}\psi, \psi) = (\mathbf{K}\psi, \mathbf{K}\mathbf{K}\psi) = (\mathbf{K}\psi, \mathbf{K}^2\psi) = (-1)^n(\mathbf{K}\psi, \psi)$. We see that the Hermitian scalar product must be zero if n is odd, or that ψ and $\mathbf{K}\psi$ are then orthogonal and hence linearly independent, yet have the same energy if $\mathbf{K}\mathbf{H} = \mathbf{H}\mathbf{K}$. This is the content of Kramers Theorem as usually stated, namely, that every level for an odd-electron system is at least twofold degenerate under the influence of external perturbations of a purely electrostatic character. We must examine further the inclusion of the operator \mathbf{K} into our previous considerations on group representations.

We shall consider the following three cases (Frobenius and Schur, 1906; Wigner 1932):

1. The rep R is real, or may be put into some equivalent form which is wholly real. This refers to the elements \mathbf{R} of the rep not merely the traces.
2. R cannot be made real, and R^* is not equivalent to it.
3. R cannot be made real, but R^* is equivalent to R .

For the finite groups a calculation of $\frac{1}{g} \sum_{\mathbf{R}} \chi(\mathbf{R}^2)$ will yield 1, 0, or -1 in cases (1), (2), or (3), respectively. Case (1) occurs for integral J in D_J and in all its subgroups save those one-dimensional reps where complex numbers appear explicitly in the character table (table 8). The only such specific reps of the double groups are S_{3g} , S_{2n} of ΠS_6 and S_3 of ΠC_3 . Case (2) holds in those one-dimensional reps where there are obviously complex characters. The only two-dimensional reps in this category are the pairs $({}_2S_{1,2}, {}_2S_{2,1})$, $({}_2S_{1,2}, {}_2S_{2,1})$, and $({}_2S_{1,2}, {}_2S_{2,1})$ in the groups ΠT_h and ΠT . Case (3) does not hold for any reps of the single groups, but holds for all those of the double groups not in (1) or (2).

In case (1) let us consider a group operation \mathbf{Q} such that $\mathbf{Q}\psi_i = \sum Q_{ij}\psi_j$ and $\mathbf{K}\mathbf{Q}\psi_i = \mathbf{Q}\mathbf{K}\psi_i = \sum Q_{ij}\mathbf{K}\psi_j$, so the function $\mathbf{K}\psi$ will transform like ψ under the operations of the group. Let us further consider $\phi_i = \psi_i + \mathbf{K}\psi_i$ and $\Omega_i = i(\psi_i - \mathbf{K}\psi_i)$. Clearly ϕ_i and Ω_i transform only among themselves under the group operations. Furthermore, if there is an even number of electrons, $\mathbf{K}\phi_i = \mathbf{K}\psi_i + \mathbf{K}^2\psi_i = \mathbf{K}\psi_i$

$+\psi_i = \phi_i$, and $\mathbf{K}\Omega_i = \mathbf{K}i\psi_i - \mathbf{K}i\mathbf{K}\psi_i = -i\mathbf{K}\psi_i + i\psi_i = \Omega_i$, so the two sets of functions ϕ_i and Ω_i are not related by time reversal or otherwise, and any degeneracy present is presumably only an accidental degeneracy. Practically speaking, this means that our set of wave functions may be chosen to be real, and that in such a case time reversal yields no connections among the set of real functions that are not already present as a result of the rotation-inversion symmetry. If there is an odd number of electrons, $\mathbf{K}\phi_i = \mathbf{K}\psi_i - \psi_i \neq \phi_i$, and we cannot so separate the ϕ and Ω sets of wave functions as we did above. Hence the rep must appear twice, once for ψ and once for $\mathbf{K}\psi$ orthogonal to it. The only actual such cases are the reps S_{3g} , S_{2n} of S_6 and S_3 of C_3 and it may be verified from table 9 for S_6 that they do in fact appear only an even number of times in any reduction of a group of half-integral J .

In case (2), $\mathbf{Q}\psi_i = \sum Q_{ij}\psi_j$ yields $\mathbf{K}\mathbf{Q}\psi_i = \mathbf{Q}\mathbf{K}\psi_i = \sum Q_{ij}^*\mathbf{K}\psi_j$. Since the rep R^* is not equivalent to R , ψ and $\mathbf{K}\psi$ belong to different reps of the group and hence are orthogonal (section 4.3), but have the same energy. Hence a separation of a rep such as E of C_{3v} into C_1 and C_2 of C_3 by an electrostatic field does not imply a removal of the degeneracy. It is complex conjugate reps such as these which are usually grouped together into a single two-dimensional rep by workers in the field of molecular spectroscopy.

The equivalence of R and R^* in case (3) requires the existence of a unitary matrix \mathbf{S} such that $\mathbf{S}\mathbf{R}\mathbf{S}^{-1} = \mathbf{R}^*$ or $\mathbf{S}^*\mathbf{R}^*(\mathbf{S}^{-1})^* = \mathbf{R}$, so $\mathbf{S}^*\mathbf{S}\mathbf{R}\mathbf{S}^{-1}(\mathbf{S}^{-1})^* = \mathbf{R}$ or $\mathbf{S}^*\mathbf{S}\mathbf{R} = \mathbf{R}\mathbf{S}^*\mathbf{S}$. We see that $\mathbf{S}^*\mathbf{S}$ commutes with every \mathbf{R} and hence must be a multiple of the unit matrix, say $c\mathbf{E}$, with c of absolute value 1. With $\mathbf{S}^*\mathbf{S} = c\mathbf{E}$, we have $\mathbf{S}^* = c\mathbf{S}^{-1} = c(\tilde{\mathbf{S}})^*$, where $\tilde{\mathbf{S}}$ is the transposed matrix of \mathbf{S} . This is a consequence of the unitary property of \mathbf{S} . The conjugate of this relation is $\mathbf{S} = c^*\tilde{\mathbf{S}}$, and the transpose of this gives $\tilde{\mathbf{S}} = c^*\mathbf{S}$, finally giving $\mathbf{S} = c^*(c^*)\mathbf{S}$, so $(c^*)^2 = 1$ or $c = \pm 1$. If $c = +1$, we have \mathbf{S} with $+1$ along the minor diagonal and zero elsewhere—in this case $\mathbf{R} = \mathbf{R}^*$ and case (1) applies, so we take $c = -1$. Then \mathbf{S} will consist of alternate $+1$ and -1 along the minor (nonprincipal) diagonal. Incidentally, this requires that the dimensionality of \mathbf{S} be even.

Let us put \mathbf{S} in the form such that $+1$ is in the upper right corner, and let us label the rows and columns of \mathbf{S} and the \mathbf{R} matrices by the numbers $j, j-1, j-2, \dots, -j$, where j is half-integral.

Then

$$S_{pe} = (-1)^{j-p}\delta_{p(-e)} \text{ and } (S^{-1})_{ab} = (-1)^{j-a+1}\delta_{a(-b)}.$$

The transformation $\mathbf{S}\mathbf{R}\mathbf{S}^{-1}$ is then as follows:

$$\begin{aligned} (\mathbf{R}\mathbf{S}^{-1})_{ab} &= \sum_k R_{ak}(S^{-1})_{kb} = \sum_k R_{ak}(-1)^{j-k+1}\delta_{k(-b)} \\ &= (-1)^{j+1}R_{a(-b)} \end{aligned}$$

and

$$(SRS^{-1})_{cd} = \sum_k (-1)^{j-c} \delta_{c(-k)} (-1)^{j+d+1} R_{k(-d)} \\ = (-1)^{2j+1+d-e} R_{(-e)(-d)} = (-1)^{d-e} R_{(-e)(-d)}.$$

This latter term must then also be $(R^*)_{cd}$, since \mathbf{R} and \mathbf{R}^* are equivalent by the transformation \mathbf{S} .

We now consider

$$\mathbf{Q}\psi_a = \sum_b Q_{ab} \psi_b$$

and

$$\mathbf{K}\mathbf{Q}\psi_a = \mathbf{K}\mathbf{Q}\psi_a \\ = \sum_b Q_{ab}^* \mathbf{K}\psi_b = \sum_b (-1)^{b-a} Q_{(-a)(-b)} \mathbf{K}\psi_b$$

or

$$\mathbf{Q}(-1)^a \mathbf{K}\psi_a = \sum_b Q_{(-a)(-b)} (-1)^b \mathbf{K}\psi_b.$$

If we compare this with the transformation on ψ_{-a} obtaining $\mathbf{Q}\psi_{-a} = \sum_b Q_{(-a)(-b)} \psi_{-b}$ we see that $(-1)^a \mathbf{K}\psi_a$ will transform like ψ_{-a} under the group operations, and that $(-1)^{-a} \mathbf{K}\psi_{-a}$ will transform like ψ_a . The linear combination $\phi_a = \psi_a + (-1)^{-a} \mathbf{K}\psi_{-a}$ will transform in a similar fashion under the group operations. Further, we observe that

$$\mathbf{K}\phi_a = \mathbf{K}\psi_a + (-1)^a \mathbf{K}^2 \psi_{-a} = \mathbf{K}\psi_a - (-1)^a \psi_{-a} \\ = -(-1)^a [\psi_{-a} + (-1)^a \mathbf{K}\psi_a] = -(-1)^a \phi_{-a},$$

since our convention has "a" half-integral, $(-1)^a$ is purely imaginary and $\mathbf{K}_0(-1)^a = (-1)^{-a} = -(-1)^a$, and $\mathbf{K}^2 \psi_a = -\psi_a$. The important point is that the set of functions $\phi_a (a=j, j-1, \dots, -j)$ transform among themselves under the group operations, and that the operator \mathbf{K} merely yields a wave function which, apart from a phase factor, is already in the set. Similarly, if $\Omega_a = i(\psi_a - (-1)^a \mathbf{K}\psi_{-a})$ the Ω_a will also transform among themselves under the group operations and time reversal, but will not involve the ϕ_a . Hence we conclude that the introduction of time reversal does not require any additional essential degeneracy in case (3).

For emphasis, we shall summarize the cases where we find that the symmetry of time reversal (with external electrostatic fields only) gives us degeneracy beyond that expected from previous considerations.

1. States belonging to complex conjugate reps are degenerate. This is the case for both an odd or even number of electrons.

2. The reps S_{3g}, S_{3u} of S_6 and S_3 of C_3 , arising from odd-electron systems will always occur twice, once for ψ and once for its orthogonal but degenerate conjugate $\mathbf{K}\psi$.

Finally, let us consider

$$\mathbf{K} = (i)^n (S_{\nu})_1 (S_{\nu})_2 \dots (S_{\nu})_n \mathbf{K}_0$$

operating on a wave function consisting of 2^n

terms of the form

$$\psi = f(\mathbf{r}) \zeta_1(S_1) \zeta_2(S_2) \dots \zeta_n(S_n).$$

Since

$$S_{\nu} \zeta(S) = (-1)^{\frac{1}{2}n-S} i^S \zeta(-S), \quad i S_{\nu} \zeta(S) = (-1)^{\frac{1}{2}n+S} \zeta(-S)$$

and

$$\mathbf{K}\psi = f^*(\mathbf{r}) (-1)^{\frac{n}{2}+2S} \zeta_1(-S_1) \zeta_2(-S_2) \dots \zeta_n(-S_n).$$

It will be noted that \mathbf{K} will commute with \mathbf{L}^2 , \mathbf{S}^2 and \mathbf{J}^2 , but not with \mathbf{L}_z , \mathbf{S}_z or \mathbf{J}_z . If $f(\mathbf{r})$ belongs to the eigenvalue L and M_L , $f^*(\mathbf{r})$ will belong to L , $-M_L$ if the usual phase factor $(-1)^{M_L}$ is supplied. Similarly, the original spin terms, belonging to S , M_S will belong to S , $-M_S$ with the phase factor $(-1)^{\frac{n}{2}+M_S}$. Thus

$$\mathbf{K}\psi(L, M_L, S, M_S) \\ = (-1)^{M_L} (-1)^{M_S} (-1)^{\frac{n}{2}} \psi(L, -M_L, S, -M_S).$$

The transformation from L, M_L, S, M_S to L, S, J, M_J is accomplished by the Clebsch-Gordan coefficients (section 5.2), which have phase factors such that

$$C_{M_L M_S}^{L S J} = (-1)^{L+S-J} C_{-M_L -M_S}^{L S J}$$

In a state described by J, M_J , we must also introduce this factor, and we obtain finally

$$\mathbf{K}\psi(L, S, J, M_J) \\ = (-1)^{M_L} (-1)^{M_S} (-1)^{\frac{n}{2}} (-1)^{L+S-J} \psi(L, S, J, -M_J) \\ = (-1)^{J-M_J} (-1)^{L+S+\frac{n}{2}} \psi(L, S, J, -M_J).$$

The factor $(-1)^{L+S+\frac{n}{2}}$ we shall neglect, but the

term $(-1)^{J-M_J}$ is of some importance, since we shall be considering states which are a linear combination of states with the same J and different M_J . There will be a physically significant shift of relative phases among the components of such states when passing to the time-reversed state.

4.3. Orthogonal States and Vanishing Matrix Elements

Before proceeding to the consideration of selection rules and polarization, we shall digress here briefly to consider certain aspects of the calculation of matrix elements of the form

$$\int \psi_a^* X \psi_b d\tau$$

which will be required at this time, deferring to the next section additional details. Let us assume initially that $X=1$. Let ψ_a and ψ_b belong to reps (1) and (2) respectively of the appropriate symmetry group, so that any group

operation Q will yield $Q\psi_a = \sum_i Q_{ia}^{(1)}\psi_i^{(1)}$ and $Q\psi_b = \sum_j Q_{jb}^{(2)}\psi_j^{(2)}$, where $Q_{ia}^{(1)}$ and $Q_{jb}^{(2)}$ are the same if the two reps (1) and (2) are the same, otherwise they are distinct. The operation Q on $\psi_i^*\psi_a$ then gives $\sum_{j,k} Q_{ij}^{(2)*} Q_{ka}^{(1)} \psi_j^* \psi_k = \sum_{j,k} Q_{ba, jk} \psi_j^* \psi_k^{(1)}$. The matrix with elements $Q_{ba, jk}$ is called the direct product of $Q^{(2)*}$ and $Q^{(1)}$, written $Q^{(2)* \times (1)}$. If these latter are of dimensionalities n_1 and n_2 , then the direct product is a $n_1 n_2 \times n_1 n_2$ matrix. Obviously, $Q^{(2)* \times (1)}$ is an element of a representation of the group—its trace is $\sum_{ab} Q_{ba, ba} = \sum_b Q_{bb}^{(2)*} \sum_a Q_{aa}^{(1)}$, the product of the traces of the two "factor" matrices.

In general, $Q^{(2)* \times (1)}$ will be either a rep or a sum of reps, in which latter case it may be reduced by the methods of the preceding section. In either case, we obtain $\sum_{j,k} Q_{ba, jk} \psi_j^* \psi_k^{(1)} = \sum_d Q_{cd}^{(3)*} \psi_d^{(3)} + \sum_f Q_{ef}^{(4)*} \psi_f^{(4)} \dots$ where $\psi_d^{(3)}, \psi_f^{(4)}$, etc., are the combinations of $\psi_j^* \psi_k^{(1)}$ appropriate to the reduced representation. Our original integral then becomes $\sum_d Q_{cd}^{(3)*} \int \psi_d^{(3)} d\tau + \sum_f Q_{ef}^{(4)*} \int \psi_f^{(4)} d\tau \dots$

The basic argument we shall use here is that our integral, i.e., the matrix element, represents a physically significant quantity and hence cannot depend on the choice of any symmetrically equivalent reference frame. Consequently, the terms of the form $\sum Q_{cd} \int \psi_d^{(3)} d\tau$ must either actually be independent of Q or must vanish. If the rep (3) is actually independent of Q , it is then in fact the completely symmetric or identity rep of the group, and we must ascertain the conditions under which the rep is present in the representation $Q^{(2)* \times (1)}$.

The identity rep has the character +1 for every group operation, so the orthogonality theorem for group characters tells us that $\frac{1}{g} \sum_Q \sum_{ab} Q_{ba, ba}$ is the number of times the symmetric rep occurs in the reduction. But we recall that $\sum_Q \sum_{ab} Q_{ba, ba} = \sum_Q \left(\sum_b Q_{bb}^{(2)*} \right) \left(\sum_a Q_{aa}^{(1)} \right)$, and this latter expression is merely the orthogonality theorem applied to the characters of the reps (1) and (2), which is zero unless (1)=(2). Hence we conclude the very important result, that states belonging to different reps of the appropriate symmetry group are orthogonal.

The example just considered took $X=1$. If X itself, either a function or operator, transforms as a rep or sum of reps of the symmetry group, the product $X\psi_a^{(1)}$ must first be reduced. If this reduction does not contain the rep (2), then the integral must vanish.

4.4. Selection Rules and Polarization

Selection rules and polarization rules are intimately connected, and we shall consider them together. We shall consider in detail the effect of an incident perturbing electromagnetic radiation superimposed upon the static electric and/or magnetic field already present. In a region of space not including the sources of the field, the perturbing radiation is usually described by its associated vector potential A , with the auxiliary requirement $\nabla \cdot A = 0$, which will allow us to set the scalar potential $V=0$. The fields E and B are then given by $E = -\frac{1}{c} \frac{\partial A}{\partial t}$ and $B = \nabla \times A$. This vector potential modifies the classical Hamiltonian term for one electron, $\frac{1}{2m} p^2$, to

$$\frac{1}{2m} \left(p - \frac{e}{c} A \right)^2 = \frac{1}{2m} \left(p^2 - \frac{e}{c} p \cdot A - \frac{e}{c} A \cdot p + \frac{e^2}{c^2} A^2 \right).$$

Neglecting the term in A^2 , we obtain as the perturbation term

$$\begin{aligned} H' &= -\frac{e}{2mc} (p \cdot A + A \cdot p) \\ &= -\frac{e}{2mc} [A \cdot p + A \cdot p + (p \cdot A)]. \end{aligned}$$

Since p is, apart from a constant factor, the vector operator ∇ , the term $(p \cdot A)$ vanishes and our final $H' = -\frac{e}{mc} A \cdot p$. This acts on each electron individually, and the total perturbation requires that this be added for all electrons in the atom. If a

system is initially in a state $\Psi_a = \psi_a e^{-\frac{iE_a t}{\hbar}}$ the probability $C_b C_b^*$ that the system under the influence of the time-dependent perturbation H' will at a later time be in the state Ψ_b is determined by

$$\frac{dC_b}{dt} = -\frac{i}{\hbar} \int \Psi_b^* H' \Psi_a d\tau$$

so we shall investigate this integral.

The components of A are functions of (x, y, z, t) . We shall assume that the variation of A with t is harmonic. We shall not require explicitly the harmonic factor $e^{i\omega t}$ in the following development and shall generally omit further reference to it; i.e., $A(x, y, z, t) = A(x, y, z) e^{i\omega t}$. For the wavelengths of light in which we are interested, λ is of the order of 10^{-4} cm. In comparison with the order of size of the atom, 10^{-8} cm, to a first approximation the variation of the components of A with (x, y, z) may frequently be neglected. We shall see that this approximation may be inadequate for our purposes, and we will improve this approximation by developing the components of the vector po-

tential in a Taylor's series about the origin. This will yield

$$A_z(x, y, z) = A_z^0 + x \left(\frac{\partial A_z}{\partial x} \right)_0 + y \left(\frac{\partial A_z}{\partial y} \right)_0 + z \left(\frac{\partial A_z}{\partial z} \right)_0,$$

where the derivatives are evaluated at the origin. Similar relations hold for A_x and A_y . To this approximation, H' becomes

$$\begin{aligned} H' = & -\frac{e}{mc} \sum \left[(A_z^0 p_x + A_y^0 p_y + A_x^0 p_x) + \left(\frac{\partial A_z}{\partial x} \right) x p_x \right. \\ & + \left(\frac{\partial A_z}{\partial y} \right) y p_x + \left(\frac{\partial A_z}{\partial z} \right) z p_x + \left(\frac{\partial A_y}{\partial x} \right) x p_y \\ & + \left(\frac{\partial A_y}{\partial y} \right) y p_y + \left(\frac{\partial A_y}{\partial z} \right) z p_y + \left(\frac{\partial A_x}{\partial x} \right) x p_x \\ & \left. + \left(\frac{\partial A_x}{\partial y} \right) y p_x + \left(\frac{\partial A_x}{\partial z} \right) z p_x \right]. \end{aligned}$$

The first three of these terms will involve matrix elements of the components of \mathbf{p} only, since \mathbf{A}^0 is a constant. The general quantum-mechanical equations of motion allow $\int \Psi^* \mathbf{p} \Psi d\tau$ to be written

as $\frac{im}{\hbar} (E_0 - E_1) \int \Psi^* \mathbf{r} \Psi d\tau$. Note that our integral

now involves the time-independent wave functions. Thus, these three terms can be written in terms of the components of the vector \mathbf{r} rather than of \mathbf{p} . Since \mathbf{er} is the classical electric dipole, transitions induced by these three terms are called electric dipole transitions. Let us assume temporarily that there is no external perturbation apart from the radiation, so that J and parity remain good quantum numbers. Let us consider an initial state transforming as J_0 , $J_0 \neq 0$. Each of the vector components x , y , and z will transform as 1_u (i.e., as the components of an odd state $J=1$). Applying our previous considerations,

the character of the rep J_1 is $\sum_{J'} e^{iM\phi}$ for both proper and improper rotations, while that for 1_u is $(e^{-i\phi} + 1 + e^{i\phi})$ for proper rotations and (-1) times this for improper rotations. The product of the characters for pure rotation yields

$$\begin{aligned} (e^{-i\phi} + 1 + e^{i\phi}) \left(\sum_{J'} e^{iM\phi} \right) = & \sum_{-(J+1)}^{J+1} e^{iM\phi} \\ & + \sum_{-J}^J e^{iM\phi} + \sum_{-(J-1)}^{J-1} e^{iM\phi}, \end{aligned}$$

and (-1) times this for the improper rotations. It is immediately clear that the direct product $1_u \times J_0$ yields $(J-1)_u + J_u + (J+1)_u$, and (from section 4.3) our final state arising from the transition must be one of these. This is equivalent to the well-known selection rule $\Delta J = 0, \pm 1$, and a particular example of Laporte's rules, where

even must combine with odd terms. If $J=0$, $1_u \times 0_u = 1_u$, and we obtain the restriction that $J=0$ does not combine with $J=0$.

The other set of 9 terms in H' does not itself transform under a single rep of the rotation group, and must be separated into terms, each one of which does so transform. The 9 terms consist of the general products of the 3 components of the vector \mathbf{r} with those of the vector \mathbf{p} . It is well known that three linearly independent terms arise from the vector product $\mathbf{r} \times \mathbf{p}$, and a fourth arises from the scalar product $\mathbf{r} \cdot \mathbf{p}$, leaving five terms, constituting the components of an irreducible tensor of rank 2 (in contrast to the original set of 9 components, which is also a tensor but not irreducible). In the terminology of the preceding paragraph, we have $1_u \times 1_u = 2_g + 1_g + 0_g$. In order to exhibit these more explicitly, let us

consider $\left(\frac{\partial A_z}{\partial y} \right) y p_x + \left(\frac{\partial A_y}{\partial x} \right) x p_y$. By adding

and subtracting $\left(\frac{\partial A_z}{\partial y} \right) x p_y + \left(\frac{\partial A_y}{\partial x} \right) y p_x$ the expression may be brought in to the form

$$\begin{aligned} \frac{1}{2} \left(\frac{\partial A_z}{\partial y} + \frac{\partial A_y}{\partial x} \right) (y p_x + x p_y) \\ + \frac{1}{2} \left(\frac{\partial A_y}{\partial x} - \frac{\partial A_z}{\partial y} \right) (x p_y - y p_x), \end{aligned}$$

and a similar relation may be derived for the other 2 pairs of similar form. This process amounts to finding the antisymmetric portion of the original 9 components, viewed as a 3×3 matrix. We observe that the first factor of the second term is the z component of $\nabla \times \mathbf{A} = \mathbf{B}$ and that the second factor is the z component of $\mathbf{r} \times \mathbf{p} = \mathbf{L}$, so, including the other 2 pairs of terms, we have a contribution

$-\frac{e}{2mc} \mathbf{B} \cdot \mathbf{L}$ to H' . Since the quantity $-\frac{e}{2mc} \mathbf{L}$

is the magnetic moment operator, transitions arising from these terms are called magnetic dipole transitions. The components of \mathbf{L} (a pseudovector) transform as 1_g , in contrast to the electric dipole's 1_u , so here even states combine with even, or odd with odd. We may note at this point that we have not yet included any terms in our radiative perturbation representing the interaction of the radiation with the spin of the electron. This may be accomplished at this point (in a more or less ad hoc manner)

by using $-\frac{e}{2mc} (\mathbf{L} + 2\mathbf{S})$ as the interaction with

the magnetic part of the incident radiation, though this will not change the transformation properties of the term, since spin is also an angular momentum operator with the transformation 1_g . It will be recalled from section 3.2 that the transformation (3.6) is equivalent to a rotation among the components of the spin angular momentum

operator (then called R) and hence equivalent to a transformation like the three components of a state $J=1$.

In considering the other six linearly independent terms, we again have recourse to the general quantum mechanical equations of motion. In particular, we obtain

$$\int \Psi_i^*(x p_x) \Psi_a d\tau = \frac{im}{\hbar} (E_b - E_a) \int \psi_b(\frac{1}{2}x^2) \psi_a d\tau$$

and

$$\int \Psi_i^*(x p_y + y p_x) \Psi_a d\tau = \frac{im}{\hbar} (E_b - E_a) \int \psi_b(xy) \psi_a d\tau.$$

If we further simplify the notation by writing A_{xx} for $(\frac{\partial A_x}{\partial x})$ and A_{xy} for $(\frac{\partial A_x}{\partial y} + \frac{\partial A_y}{\partial x})$ our remaining perturbation can be written in terms of the matrix elements of

$$(A_{xx}x^2 + A_{yy}y^2 + A_{zz}z^2 + A_{xy}xy + A_{xz}xz + A_{yz}yz).$$

In lieu of x^2 , y^2 , and z^2 , let us consider the linearly independent combinations

$$\begin{aligned} r^2 &= x^2 + y^2 + z^2 \\ Z^2 &= 2z^2 - x^2 - y^2 \\ \rho^2 &= x^2 - y^2 \end{aligned}$$

with the inverse relations

$$\begin{aligned} x^2 &= \frac{1}{2} \rho^2 + \frac{1}{2} r^2 - \frac{1}{2} Z^2 \\ y^2 &= \frac{1}{2} r^2 - \frac{1}{2} Z^2 - \frac{1}{2} \rho^2 \\ z^2 &= \frac{1}{2} Z^2 + \frac{1}{2} r^2. \end{aligned}$$

Substituting, we have as our perturbation the matrix elements of

$$\begin{aligned} r^2 (\frac{1}{2} A_{xx} + \frac{1}{2} A_{yy} + \frac{1}{2} A_{zz}) + Z^2 (\frac{1}{2} A_{zz} - \frac{1}{2} A_{xx} - \frac{1}{2} A_{yy}) \\ + \rho^2 (\frac{1}{2} A_{xx} - \frac{1}{2} A_{yy}) + A_{xy}xy + A_{xz}xz + A_{yz}yz. \end{aligned}$$

Of these six terms, that in r^2 is the O_0 term, invariant under rotation and inversion, and here actually zero, since $A_{xx} + A_{yy} + A_{zz} = \text{div } \mathbf{A} = 0$. Notice that the terms Z^2 , xy , etc., are, apart from normalization factors, the tesseral harmonics of degree 2 (table 2).

The remaining five terms are the components of the quadrupole moment, and transitions arising from them are called electric quadrupole transitions. The general selection rule is here given by a consideration of the \mathcal{Q}_2 transformation properties of the quadrupole moment. If $J \geq 2$,

$\mathcal{Q}_2 \times J_0 = (J+2)_0 + (J+1)_0 + J_0 + (J-1)_0 + (J-2)_0$, so the general selection rule is $\Delta J = \pm 2, \pm 1, 0$. If $J=1$, $\mathcal{Q}_2 \times 1_0 = \mathcal{Q}_2 + \mathcal{Q}_2 + 1_0$, so $J=1$ does not combine with $J=0$. Likewise, $\mathcal{Q}_2 \times 0_0 = \mathcal{Q}_2$, so $J=0$ does not combine with $J=0$ or $J=1$.

We have up to this point considered that the symmetry present was that of the full three-dimensional rotation-inversion group. If this is not the case, the 3 electric dipole components, the 3 magnetic dipole components, and the 5 electric quadrupole components will belong to different reps of the symmetry group. These may be determined from table 9 for 1_u , 1_g , and \mathcal{Q}_2 , respectively, and the detailed correlation may in most cases be obtained from table 10. The use of table 10 after an initial application of table 9 for the group $D_{\infty h}$ (for noncubic fields) is perhaps most convenient. The reduction to $D_{\infty h}$ will separate both 1_u and 1_g into one 1-dimensional rep and one 2-dimensional rep, while \mathcal{Q}_2 yields one 1-dimensional rep and two 2-dimensional reps. These may be followed to lower symmetries with table 10. Details may be obtained from table 8. For example, L , belonging to 1_g , yields, in $D_{\infty h}$, L_z belonging to A_{2g} , while L_x and L_y both belong to E_{1g} . Upon reducing the symmetry to C_{4v} , L_z now becomes A_2 while L_x and L_y now both belong to E . If we consider the reduction from $D_{\infty h}$ to $C_{\infty v}$, we see that E_{1g} splits into C_{1g} and C_{-1g} . However, these do not represent the transformation of either L_x or L_y , but of $L_x + iL_y$ and $L_x - iL_y$, respectively. In this case, and also for the subgroups of $C_{\infty v}$, we shall consider these latter, along with L_z , as the three linearly independent components, each of which involves only one rep of the symmetry group. Similarly, we shall resolve the electric dipole term \mathbf{r} into x , y , and z for most groups, using $x + iy$, $x - iy$, and z for the subgroups of $C_{\infty v}$, and the electric quadrupole components into Z^2 , xz , yz , ρ^2 , xy , or, alternatively, Z^2 , $z(x + iy)$, $z(x - iy)$, $(x + iy)^2$, and $(x - iy)^2$. These resolutions are presented in table 12.

As usual, there are two possibilities for the cubic groups, one in which the z -axis is an even-fold axis and one in which it is an odd-fold axis. This makes a difference only for the quadrupole components, since the electric and magnetic dipole components remain indistinguishable under these symmetries. If we consider the z -axis to be an even-fold axis, i.e., C_4 for O and O_h , S_4 for T_d , C_2 for T_h and T , the five real quadrupole components remain a satisfactory basis for the resolution in all cases except for two components in T_h and T , where we require the normalized components

$$\frac{1}{\sqrt{2}} [C_2 + iC_2] \quad \text{and} \quad \frac{1}{\sqrt{2}} [C_2 - iC_2]$$

transforming as C_{1g} and C_{2g} respectively.

If we take the z -axis to be the C_3 axis in these cases, we find that these quadrupole components, the tesseral harmonics of degree 2, do not serve as a basis for the reduction of the \mathcal{Q}_2 representation

TABLE 12. Multipole moment transformations

	x	y	z	L_x	L_y	L_z	Z^2	xz	yz	ρ^2	xy
$D_{\infty h}$ $C_{\infty v}$	E_{1u} E_1	E_{1u} E_1	A_{2u} A_1	E_{1g} E_1	E_{1g} E_1	A_{2g} A_2	A_{1g} A_1	E_{1g} E_1	E_{1g} E_1	E_{2g} E_2	E_{2g} E_2
O_h T_d	T_{1u} T_2	T_{1u} T_2	T_{1u} T_2	T_{1g} T_1	T_{1g} T_1	T_{1g} T_1	E_g E	T_{2g} T_2	T_{2g} T_2	E_g E	T_{2g} T_2
T_h D_{2h}	T_u E_{1u}	T_u E_{1u}	T_u A_{2u}	T_g E_{1g}	T_g E_{1g}	T_g A_{2g}	^(a) A_{1g}	T_g E_{1g}	T_g E_{1g}	^(a) E_{2g}	T_g E_{2g}
D_{4h} C_{4v}	E_u E_1	E_u E_1	A_{2u} A_1	E_g E_1	E_g E_1	A_{2g} A_2	A_{1g} A_1	E_g E_1	E_g E_1	B_{1g} E_2	B_{2g} E_2
D_{3h} D_{3d}	E'_u E_u	E'_u E_u	A''_u A_{2u}	E'_g E_g	E'_g E_g	A''_g A_{2g}	A'_g A_{1g}	E'_g E_g	E'_g E_g	E'_g E_g	E'_g E_g
D_{2d} C_{2v}	E E	E E	B_2 A_1	E E	E E	A_2 A_2	A_1 A_1	E E	E E	B_1 B_1	B_2 B_2
D_{2h} C_{2h}	B_{2u} E	B_{1u} E	A_{2u} A_1	B_{2g} E	B_{1g} E	A_{2g} A_2	A_{1g} A_1	B_{1g} E	B_{2g} E	A_{1g} E	A_{2g} E
C_{2v} C_{2h}	B_1 B_u	B_2 B_u	A_1 A_u	B_2 B_g	B_1 B_g	A_2 A_g	A_1 A_g	B_1 B_g	B_2 B_g	A_1 A_g	A_2 A_g
C_3	A'	A'	A''	A''	A''	A'	A'	A''	A''	A'	A'
	$x+iy$	$x-iy$	z	L_x+iL_y	L_x-iL_y	L_z	Z^2	$(x+iy)z$	$(x-iy)z$	$(x+iy)^2$	$(x-iy)^2$
$C_{\infty h}$	C_{1u}	C_{-1u}	A_u	C_{1g}	C_{-1g}	A_g	A_g	C_{1g}	C_{-1g}	C_{2g}	C_{-2g}
C_{3h} C_{4h}	C_{1u} C_{1u}	C_{2u} C_{3u}	A_u A_u	C_{1g} C_{1g}	C_{2g} C_{3g}	A_g A_g	A_g A_g	C_{1g} C_{1g}	C_{2g} C_{3g}	B_{2g} B_{2g}	C_{3g} B_{2g}
C_{2h} S_6	C'_1 C_{1u}	C'_2 C_{2u}	A'' A_u	C''_1 C_{1g}	C'_2 C_{2g}	A' A_g	A' A_g	C''_1 C_{1g}	C'_2 C_{2g}	C'_2 C_{2g}	C'_1 C_{1g}
S_4	C_1	C_3	B	C_3	C_1	A	A	C_3	C_1	B	B

^a See text.

of the quadrupole interaction. For example, the reduction of \mathcal{Q}_z under O_h yields the following as a basis:

$$E: \left[-\frac{\sqrt{6}}{3} C_2 + \frac{\sqrt{3}}{3} C_3 \right]; \left[-\frac{\sqrt{6}}{3} S_2 + \frac{\sqrt{3}}{3} S_3 \right]$$

$$T_2: \left[\frac{\sqrt{3}}{3} C_2 + \frac{\sqrt{6}}{3} C_3 \right]; \left[\frac{\sqrt{3}}{3} S_2 + \frac{\sqrt{6}}{3} S_3 \right]; \left[C_2 \right]$$

While this, of course, yields the same polarization results in the physical sense, the trigonal aspect will not be considered further due to its formal mathematical complexities.

Because of the different transformation properties of these components, they will most generally permit transitions between different states of the perturbed ion. The nonvanishing matrix elements may be deduced by the procedures outlined above (section 4.3). These are the selection rules which are applicable in a case of definite symmetry. Furthermore, the relative amount of any particular component of the multipole interaction present

will depend upon the vector potential, and in particular, upon the direction of propagation and state of polarization of the incident radiation. A detailed examination of this latter feature leads to the polarization rules for absorption of radiation.

The selection rules for most of the groups in which we shall be interested (omitting only the very simplest) are given in table 13. For each rep in the left column, the states to which a particular component of the multipole interaction will permit transitions is shown in the column under that multipole component. Generally, the table parallels table 8, in that closely related groups are shown with but minor modifications on the same table. Results for pure rotation may be obtained by dropping the "g"—"u" distinction in the tables. In a few cases, the changes in notation and convention are sufficiently far-reaching to require a separate table. It is particularly interesting to note the table for $C_{\infty h}$, appropriate to a uniform magnetic field. If we denote the general rep by C_{M_p} ($C_0=A$), we see that electric dipole transitions have $\Delta M=0$,

TABLE 13. Selection rules

D_{00A}	z	x, y	L_z	L_x, L_y, xz, yz	Z^2	ρ^2, xy
A_{1g} A_{1u}	A_{2u} A_{2g}	E_{1u} E_{1g}	A_{2g} A_{2u}	E_{1g} E_{1u}	A_{1g} A_{1u}	E_{2g} E_{2u}
A_{2g} A_{2u}	A_{1u} A_{1g}	E_{1u} E_{1g}	A_{1g} A_{1u}	E_{1g} E_{1u}	A_{2g} A_{2u}	E_{2g} E_{2u}
E_{1g} E_{1u}	E_{1u} E_{1g}	$E_{2u}A_{1u}A_{2u}$ $E_{2g}A_{1g}A_{2g}$	E_{1g} E_{1u}	$E_{2g}A_{1g}A_{2g}$ $E_{2u}A_{1u}A_{2u}$	E_{1g} E_{1u}	$E_{1g}E_{2g}$ $E_{1u}E_{2u}$
E_{2g} E_{2u}	E_{2u} E_{2g}	$E_{3u}E_{1u}$ $E_{3g}E_{1g}$	E_{2g} E_{2u}	$E_{3g}E_{1g}$ $E_{3u}E_{1u}$	E_{2g} E_{2u}	$E_{4g}A_{1g}A_{2g}$ $E_{4u}A_{1u}A_{2u}$
E_{3g} E_{3u}	E_{3u} E_{3g}	$E_{4u}E_{2u}$ $E_{4g}E_{2g}$	E_{3g} E_{3u}	$E_{4g}E_{2g}$ $E_{4u}E_{2u}$	E_{3g} E_{3u}	$E_{5g}E_{1g}$ $E_{5u}E_{1u}$

(Pattern repeats for higher states)

${}^2S_{1g}$ ${}^2S_{1u}$	${}^2S_{1u}$ ${}^2S_{1g}$	${}^2S_{1u} {}^2S_{3u}$ ${}^2S_{1g} {}^2S_{3g}$	${}^2S_{1g}$ ${}^2S_{1u}$	${}^2S_{1g} {}^2S_{3g}$ ${}^2S_{1u} {}^2S_{3u}$	${}^2S_{1g}$ ${}^2S_{1u}$	${}^2S_{3g} {}^2S_{5g}$ ${}^2S_{3u} {}^2S_{5u}$
${}^2S_{3g}$ ${}^2S_{3u}$	${}^2S_{3u}$ ${}^2S_{3g}$	${}^2S_{1u} {}^2S_{5u}$ ${}^2S_{1g} {}^2S_{5g}$	${}^2S_{3g}$ ${}^2S_{3u}$	${}^2S_{1g} {}^2S_{5g}$ ${}^2S_{1u} {}^2S_{5u}$	${}^2S_{3g}$ ${}^2S_{3u}$	${}^2S_{1g} {}^2S_{7g}$ ${}^2S_{1u} {}^2S_{7u}$
${}^2S_{5g}$ ${}^2S_{5u}$	${}^2S_{5u}$ ${}^2S_{5g}$	${}^2S_{3u} {}^2S_{7u}$ ${}^2S_{3g} {}^2S_{7g}$	${}^2S_{5g}$ ${}^2S_{5u}$	${}^2S_{3g} {}^2S_{7g}$ ${}^2S_{3u} {}^2S_{7u}$	${}^2S_{5g}$ ${}^2S_{5u}$	${}^2S_{1g} {}^2S_{9g}$ ${}^2S_{1u} {}^2S_{9u}$

(Pattern repeats for higher states)

For C_{00} , delete "g" and "u" and read the upper left corner as:

C_{00}	z
A_1	A_1
A_2	A_2

D_{00A}	z	x, y	L_z	L_x, L_y, xz, yz	Z^2	ρ^2, xy
A_{1g} A_{1u}	A_{2u} A_{2g}	E_{1u} E_{1g}	A_{2g} A_{2u}	E_{1g} E_{1u}	A_{1g} A_{1u}	E_{2g} E_{2u}
A_{2g} A_{2u}	A_{1u} A_{1g}	E_{1u} E_{1g}	A_{1g} A_{1u}	E_{1g} E_{1u}	A_{2g} A_{2u}	E_{2g} E_{2u}
B_{1g} B_{1u}	B_{2u} B_{2g}	E_{2u} E_{2g}	B_{2g} B_{2u}	E_{2g} E_{2u}	B_{1g} B_{1u}	E_{1g} E_{1u}
B_{2g} B_{2u}	B_{1u} B_{1g}	E_{2u} E_{2g}	B_{1g} B_{1u}	E_{2g} E_{2u}	B_{2g} B_{2u}	E_{1g} E_{1u}
E_{1g} E_{1u}	E_{1u} E_{1g}	$A_{1u}A_{2u}E_{2u}$ $A_{1g}A_{2g}E_{2g}$	E_{1g} E_{1u}	$A_{1g}A_{2g}E_{2g}$ $A_{1u}A_{2u}E_{2u}$	E_{1g} E_{1u}	$B_{1g}B_{2g}E_{1g}$ $B_{1u}B_{2u}E_{1u}$
E_{2g} E_{2u}	E_{2u} E_{2g}	$B_{1u}B_{2u}E_{1u}$ $B_{1g}B_{2g}E_{1g}$	E_{2g} E_{2u}	$B_{1g}B_{2g}E_{1g}$ $B_{1u}B_{2u}E_{1u}$	E_{2g} E_{2u}	$A_{1g}A_{2g}E_{2g}$ $A_{1u}A_{2u}E_{2u}$
$D_{\frac{1}{2}g}$ $D_{\frac{1}{2}u}$	$D_{\frac{1}{2}u}$ $D_{\frac{1}{2}g}$	${}^2S_{1u}D_{\frac{1}{2}u}$ ${}^2S_{1g}D_{\frac{1}{2}g}$	$D_{\frac{1}{2}g}$ $D_{\frac{1}{2}u}$	${}^2S_{1g}D_{\frac{1}{2}g}$ ${}^2S_{1u}D_{\frac{1}{2}u}$	$D_{\frac{1}{2}g}$ $D_{\frac{1}{2}u}$	${}^2S_{1g} {}^2S_{2g}$ ${}^2S_{1u} {}^2S_{2u}$
${}^2S_{1g}$ ${}^2S_{1u}$	${}^2S_{1u}$ ${}^2S_{1g}$	${}^2S_{2u}D_{\frac{1}{2}u}$ ${}^2S_{2g}D_{\frac{1}{2}g}$	${}^2S_{1g}$ ${}^2S_{1u}$	${}^2S_{2g}D_{\frac{1}{2}g}$ ${}^2S_{2u}D_{\frac{1}{2}u}$	${}^2S_{1g}$ ${}^2S_{1u}$	${}^2S_{2g}D_{\frac{1}{2}g}$ ${}^2S_{2u}D_{\frac{1}{2}u}$
${}^2S_{2g}$ ${}^2S_{2u}$	${}^2S_{2u}$ ${}^2S_{2g}$	${}^2S_{1u} {}^2S_{2u}$ ${}^2S_{1g} {}^2S_{2g}$	${}^2S_{2g}$ ${}^2S_{2u}$	${}^2S_{1g} {}^2S_{2g}$ ${}^2S_{1u} {}^2S_{2u}$	${}^2S_{2g}$ ${}^2S_{2u}$	${}^2S_{1g}D_{\frac{1}{2}g}$ ${}^2S_{1u}D_{\frac{1}{2}u}$

For C_{00} , drop "g"—"u" and amend upper left corner to read:

C_{00}	z
A_1	A_1
A_2	A_2
B_1	B_1
B_2	B_2

TABLE 13. Selection rules—Continued

D_{3A}	z	x, y	L_z	L_x, L_y, xz, yz	Z^2	ρ^2	xy
A_{1g} A_{1u}	A_{2g} A_{2u}	E_u E_g	A_{2g} A_{2u}	E_g E_u	A_{1g} A_{1u}	B_{1g} B_{1u}	B_{2g} B_{2u}
A_{2g} A_{2u}	A_{1g} A_{1u}	E_u E_g	A_{1g} A_{1u}	E_g E_u	A_{2g} A_{2u}	B_{2g} B_{2u}	B_{1g} B_{1u}
B_{1g} B_{1u}	B_{2g} B_{2u}	E_u E_g	B_{2g} B_{2u}	E_g E_u	B_{1g} B_{1u}	A_{1g} A_{1u}	A_{2g} A_{2u}
B_{2g} B_{2u}	B_{1g} B_{1u}	E_u E_g	B_{1g} B_{1u}	E_g E_u	B_{2g} B_{2u}	A_{2g} A_{2u}	A_{1g} A_{1u}
E_g E_u	E_u E_g	$A_{1u}, A_{2u}, B_{1u}, B_{2u}$ $A_{1g}, A_{2g}, B_{1g}, B_{2g}$	E_g E_u	$A_{1g}, A_{2g}, B_{1g}, B_{2g}$ $A_{1u}, A_{2u}, B_{1u}, B_{2u}$	E_g E_u	E_g E_u	E_g E_u
D_{3g} D_{3u}	D_{3g} D_{3u}	${}^2S_u, D_{3u}$ ${}^2S_g, D_{3g}$	D_{3g} D_{3u}	${}^2S_g, D_{3g}$ ${}^2S_u, D_{3u}$	D_{3g} D_{3u}	2S_g 2S_u	2S_g 2S_u
2S_g 2S_u	2S_u 2S_g	${}^2S_u, D_{3u}$ ${}^2S_g, D_{3g}$	2S_g 2S_u	${}^2S_g, D_{3g}$ ${}^2S_u, D_{3u}$	2S_g 2S_u	D_{3g} D_{3u}	D_{3g} D_{3u}

For C_{3v} : Drop "g" and "u" and read z in Z^2 column.
 For D_{2d} : Drop "g" and "u" and read z in xy column.

D_{3A}	z	x, y, ρ^2, xy	L_z	L_x, L_y, xz, yz	Z^2
A_1' A_2'	A_1'' A_2''	E' E''	A_1' A_2'	E'' E'	A_1' A_2'
A_1'' A_2''	A_2' A_1'	E'' E'	A_2'' A_1''	E' E''	A_1'' A_2''
E' E''	E'' E'	A_1', A_2', E' A_1'', A_2'', E''	E' E''	A_1'', A_2'', E'' A_1', A_2', E'	E' E''
D_{3g} 2S_1 2S_2	2S_2 2S_1 D_{3g}	${}^2S_1, {}^2S_2$ $D_{3g}, {}^2S_2$ $D_{3g}, {}^2S_1$	D_{3g} 2S_1 2S_2	$D_{3g}, {}^2S_1$ ${}^2S_2, D_{3g}$ ${}^2S_1, {}^2S_2$	D_{3g} 2S_1 2S_2

± 1 , with change in parity, magnetic dipole transitions have $\Delta M=0, \pm 1$, with no change in parity, while electric quadrupole transitions have $\Delta M=0, \pm 1, \pm 2$, with no change in parity.

In order to examine in detail the polarization effects, let us consider a set of orthogonal unit vectors in spherical coordinates.

$$\mathbf{r}_0 = \sin \theta \cos \phi \mathbf{i} + \sin \theta \sin \phi \mathbf{j} + \cos \theta \mathbf{k}$$

$$\theta_0 = \frac{\partial \mathbf{r}_0}{\partial \theta} = \cos \theta \cos \phi \mathbf{i} + \cos \theta \sin \phi \mathbf{j} - \sin \theta \mathbf{k}$$

$$\phi_0 = \frac{1}{\sin \theta} \frac{\partial \mathbf{r}_0}{\partial \phi} = -\sin \phi \mathbf{i} + \cos \phi \mathbf{j}$$

TABLE 13. Selection rules—Continued

D_{2d}	z	x, y	L_z	ρ^2, xy L_x, L_y xz, yz	Z^2
A_{1g} A_{1u}	A_{2u} A_{2g}	E_u E_g	A_{2g} A_{2u}	E_g E_u	A_{1g} A_{1u}
A_{2g} A_{2u}	A_{1u} A_{1g}	E_u E_g	A_{1g} A_{1u}	E_g E_u	A_{2g} A_{2u}
E_g E_u	E_u E_g	$A_{1u} A_{2u} E_u$ $A_{1g} A_{2g} E_g$	E_g E_u	$A_{1g} A_{2g} E_g$ $A_{1u} A_{2u} E_u$	E_g E_u
S_{1g} S_{1u}	S_{2u} S_{2g}	$D_{3/2u}$ $D_{3/2g}$	S_{2g} S_{2u}	$D_{3/2g}$ $D_{3/2u}$	S_{1g} S_{1u}
S_{2g} S_{2u}	S_{1u} S_{1g}	$D_{3/2u}$ $D_{3/2g}$	S_{1g} S_{1u}	$D_{3/2g}$ $D_{3/2u}$	S_{2g} S_{2u}
$D_{3/2g}$ $D_{3/2u}$	$D_{3/2u}$ $D_{3/2g}$	$S_{1u} S_{2u} D_{3/2u}$ $S_{1g} S_{2g} D_{3/2g}$	$D_{3/2g}$ $D_{3/2u}$	$S_{1g} S_{2g} D_{3/2g}$ $S_{1u} S_{2u} D_{3/2u}$	$D_{3/2g}$ $D_{3/2u}$

For C_{3v} : Drop "g" and "u" and read z in Z^2 column.

D_{2d}	z	x	y	L_x, xy	L_y, yz	L_z, xz	Z^2, ρ^2
A_{1g} A_{1u}	A_{2u} A_{2g}	B_{2u} B_{2g}	B_{1u} B_{1g}	A_{2g} A_{2u}	B_{2g} B_{2u}	B_{1g} B_{1u}	A_{1g} A_{1u}
A_{2g} A_{2u}	A_{1u} A_{1g}	B_{1u} B_{1g}	B_{2u} B_{2g}	A_{1g} A_{1u}	B_{1g} B_{1u}	B_{2g} B_{2u}	A_{2g} A_{2u}
B_{1g} B_{1u}	B_{2u} B_{2g}	A_{2u} A_{2g}	A_{1u} A_{1g}	B_{2g} B_{2u}	A_{2g} A_{2u}	A_{1g} A_{1u}	B_{1g} B_{1u}
B_{2g} B_{2u}	B_{1u} B_{1g}	A_{1u} A_{1g}	A_{2u} A_{2g}	B_{1g} B_{1u}	A_{1g} A_{1u}	A_{2g} A_{2u}	B_{2g} B_{2u}
$D_{3/2g}$ $D_{3/2u}$	$D_{3/2u}$ $D_{3/2g}$	$D_{3/2u}$ $D_{3/2g}$	$D_{3/2u}$ $D_{3/2g}$	$D_{3/2g}$ $D_{3/2u}$	$D_{3/2g}$ $D_{3/2u}$	$D_{3/2g}$ $D_{3/2u}$	$D_{3/2g}$ $D_{3/2u}$
-----	-----	y, L_x yz	x, L_y xz	L_x, xy	-----	-----	z, Z^2, ρ^2

C_{2v} : Omit "g"—"u" and read bottom of column.

S_6	$x+iy$	$x-iy$	z	$(x+iy)z$ L_x+iL_y $(x-iy)^2$	$(x-iy)z$ L_x-iL_y $(x+iy)^2$	L_z, Z^2
A_g A_u	C_{1u} C_{1g}	C_{2g} C_{2u}	A_u A_g	C_{1g} C_{1u}	C_{2u} C_{2g}	A_g A_u
C_{1g} C_{1u}	C_{2u} C_{2g}	A_u A_g	C_{1u} C_{1g}	C_{2g} C_{2u}	A_g A_u	C_{1g} C_{1u}
C_{2g} C_{2u}	A_u A_g	C_{1u} C_{1g}	C_{2u} C_{2g}	A_g A_u	C_{1g} C_{1u}	C_{2g} C_{2u}
S_{1g} S_{1u}	S_{3u} S_{3g}	S_{5u} S_{5g}	S_{1u} S_{1g}	S_{3g} S_{3u}	S_{5g} S_{5u}	S_{1g} S_{1u}
S_{3g} S_{3u}	S_{5u} S_{5g}	S_{1u} S_{1g}	S_{3u} S_{3g}	S_{5g} S_{5u}	S_{1g} S_{1u}	S_{3g} S_{3u}
S_{5g} S_{5u}	S_{1u} S_{1g}	S_{3u} S_{3g}	S_{5u} S_{5g}	S_{1g} S_{1u}	S_{3g} S_{3u}	S_{5g} S_{5u}

TABLE 13. Selection rules—Continued

$C_{\infty A}$	$x+iy$	$x-iy$	z	L_x+iL_y $(x+iy)z$	L_x-iL_y $(x-iy)z$	$L_x Z^2$	$(x+iy)^2$	$(x-iy)^2$
A_x	C_{1u}	C_{-1u}	A_u	C_{1x}	C_{-1x}	A_x	C_{2x}	C_{-2x}
A_u	C_{1x}	C_{-1x}	A_x	C_{1u}	C_{-1u}	A_u	C_{2u}	C_{-2u}
C_{1x}	C_{2u}	A_u	C_{1u}	C_{2x}	A_x	C_{1x}	C_{3x}	C_{-1x}
C_{1u}	C_{2x}	A_x	C_{1x}	C_{2u}	A_u	C_{1u}	C_{3u}	C_{-1u}
C_{-1x}	A_u	C_{-2u}	C_{-1u}	A_x	C_{-2x}	C_{-1x}	C_{1x}	C_{-3x}
C_{-1u}	A_x	C_{-2x}	C_{-1x}	A_u	C_{-2u}	C_{-1u}	C_{1u}	C_{-3u}
C_{2x}	C_{3u}	C_{1u}	C_{2u}	C_{3x}	C_{1x}	C_{2x}	C_{4x}	A_x
C_{2u}	C_{3x}	C_{1x}	C_{2x}	C_{3u}	C_{1u}	C_{2u}	C_{4u}	A_u
C_{-2x}	C_{-1u}	C_{-3u}	C_{-2u}	C_{-1x}	C_{-3x}	A_x	C_{-4x}	C_{-4x}
C_{-2u}	C_{-1x}	C_{-3x}	C_{-2x}	C_{-1u}	C_{-3u}	A_u	C_{-4u}	C_{-4u}
C_{3x}	C_{4u}	C_{2u}	C_{3u}	C_{4x}	C_{2x}	C_{3x}	C_{5x}	C_{1x}
C_{3u}	C_{4x}	C_{2x}	C_{3x}	C_{4u}	C_{2u}	C_{3u}	C_{5u}	C_{1u}
C_{-3x}	C_{-2u}	C_{-4u}	C_{-3u}	C_{-2x}	C_{-4x}	C_{-3x}	C_{-1x}	C_{-5x}
C_{-3u}	C_{-2x}	C_{-4x}	C_{-3x}	C_{-2u}	C_{-4u}	C_{-3u}	C_{-1u}	C_{-5u}

(Pattern for higher $C_{n,x,u}$ repeats)

S_{1x}	S_{3u}	S_{-1u}	S_{1u}	S_{3x}	S_{-1x}	S_{1x}	S_{5x}	S_{-3x}
S_{1u}	S_{3x}	S_{-1x}	S_{1x}	S_{3u}	S_{-1u}	S_{1u}	S_{5u}	S_{-3u}
S_{-1x}	S_{1u}	S_{-3u}	S_{-1u}	S_{1x}	S_{-3x}	S_{-1x}	S_{3x}	S_{-5x}
S_{-1u}	S_{1x}	S_{-3x}	S_{-1x}	S_{1u}	S_{-3u}	S_{-1u}	S_{3u}	S_{-5u}
S_{3x}	S_{5u}	S_{1u}	S_{3u}	S_{5x}	S_{1x}	S_{3x}	S_{7x}	S_{-1x}
S_{3u}	S_{5x}	S_{1x}	S_{3x}	S_{5u}	S_{1u}	S_{3u}	S_{7u}	S_{-1u}
S_{-3x}	S_{-1u}	S_{-5u}	S_{-3u}	S_{-1x}	S_{-5x}	S_{-3x}	S_{1x}	S_{-7x}
S_{-3u}	S_{-1x}	S_{-5x}	S_{-3x}	S_{-1u}	S_{-5u}	S_{-3u}	S_{1u}	S_{-7u}
S_{5x}	S_{7u}	S_{3u}	S_{5u}	S_{7x}	S_{3x}	S_{5x}	S_{9x}	S_{1x}
S_{5u}	S_{7x}	S_{3x}	S_{5x}	S_{7u}	S_{3u}	S_{5u}	S_{9u}	S_{1u}
S_{-5x}	S_{-3u}	S_{-7u}	S_{-5u}	S_{-3x}	S_{-7x}	S_{-5x}	S_{-1x}	S_{-9x}
S_{-5u}	S_{-3x}	S_{-7x}	S_{-5x}	S_{-3u}	S_{-7u}	S_{-5u}	S_{-1u}	S_{-9u}

(Pattern for higher $S_{n,x,u}$ repeats)

$C_{\infty A}$	z	$x+iy$	$x-iy$	$L_x Z^2$	L_x+iL_y $(x+iy)z$	L_x-iL_y $(x-iy)z$	$(x+iy)^2$	$(x-iy)^2$
A_x	A_u	C_{1u}	C_{5u}	A_x	C_{1x}	C_{5x}	C_{2x}	C_{4x}
A_u	A_x	C_{1x}	C_{5x}	A_u	C_{1u}	C_{5u}	C_{2u}	C_{4u}
C_{1x}	C_{1u}	C_{2u}	A_u	C_{1x}	C_{2x}	A_x	B_x	C_{5x}
C_{1u}	C_{1x}	C_{2x}	A_x	C_{1u}	C_{2u}	A_u	B_u	C_{5u}
C_{2x}	C_{2u}	B_u	C_{1u}	C_{2x}	B_x	C_{1x}	C_{4x}	A_x
C_{2u}	C_{2x}	B_x	C_{1x}	C_{2u}	B_u	C_{1u}	C_{4u}	A_u
B_x	B_u	C_{4u}	C_{2u}	B_x	C_{4x}	C_{2x}	C_{5x}	C_{1x}
B_u	B_x	C_{4x}	C_{2x}	B_u	C_{4u}	C_{2u}	C_{5u}	C_{1u}
C_{4x}	C_{4u}	C_{5u}	B_u	C_{4x}	C_{5x}	B_x	A_x	C_{2x}
C_{4u}	C_{4x}	C_{5x}	B_x	C_{4u}	C_{5u}	B_u	A_u	C_{2u}
C_{5x}	C_{5u}	A_u	C_{4u}	C_{5x}	A_x	C_{4x}	C_{1x}	B_x
C_{5u}	C_{5x}	A_x	C_{4x}	C_{5u}	A_u	C_{4u}	C_{1u}	B_u
S_{1x}	S_{1u}	S_{3u}	S_{11u}	S_{1x}	S_{3x}	S_{11x}	S_{5x}	S_{9x}
S_{1u}	S_{1x}	S_{3x}	S_{11x}	S_{1u}	S_{3u}	S_{11u}	S_{5u}	S_{9u}
S_{3x}	S_{3u}	S_{5u}	S_{1u}	S_{3x}	S_{5x}	S_{1x}	S_{7x}	S_{11x}
S_{3u}	S_{3x}	S_{5x}	S_{1x}	S_{3u}	S_{5u}	S_{1u}	S_{7u}	S_{11u}
S_{5x}	S_{5u}	S_{7u}	S_{3u}	S_{5x}	S_{7x}	S_{3x}	S_{9x}	S_{1x}
S_{5u}	S_{5x}	S_{7x}	S_{3x}	S_{5u}	S_{7u}	S_{3u}	S_{9u}	S_{1u}
S_{7x}	S_{7u}	S_{9u}	S_{5u}	S_{7x}	S_{9x}	S_{5x}	S_{11x}	S_{3x}
S_{7u}	S_{7x}	S_{9x}	S_{5x}	S_{7u}	S_{9u}	S_{5u}	S_{11u}	S_{3u}
S_{9x}	S_{9u}	S_{11u}	S_{7u}	S_{9x}	S_{11x}	S_{7x}	S_{1x}	S_{5x}
S_{9u}	S_{9x}	S_{11x}	S_{7x}	S_{9u}	S_{11u}	S_{7u}	S_{1u}	S_{5u}
S_{11x}	S_{11u}	S_{1u}	S_{9u}	S_{11x}	S_{1u}	S_{9x}	S_{3x}	S_{7x}
S_{11u}	S_{11x}	S_{1x}	S_{9x}	S_{11u}	S_{1u}	S_{9u}	S_{3u}	S_{7u}

TABLE 13. Selection rules—Continued

C_{3A}	z	$x+iy$	$x-iy$	$L_x Z^2$	L_x+iL_y $(x+iy)z$	L_x-iL_y $(x-iy)z$	$(x-iy)^2$ $(x+iy)^2$
A_x	A_u	C_{1u}	C_{3u}	A_x	C_{1x}	C_{3x}	B_x
A_u	A_x	C_{1x}	C_{3x}	A_u	C_{1u}	C_{3u}	B_u
C_{1x}	C_{1u}	B_u	A_u	C_{1x}	B_x	A_x	C_{3x}
C_{1u}	C_{1x}	B_x	A_x	C_{1u}	B_u	A_u	C_{3u}
B_x	B_u	C_{3u}	C_{1u}	B_x	C_{3x}	C_{1x}	A_x
B_u	B_x	C_{3x}	C_{1x}	B_u	C_{3u}	C_{1u}	A_u
C_{3x}	C_{3u}	A_u	B_u	C_{3x}	A_x	B_x	C_{1x}
C_{3u}	C_{3x}	A_x	B_x	C_{3u}	A_u	B_u	C_{1u}
S_{1x}	S_{1u}	S_{3u}	S_{7u}	S_{1x}	S_{3x}	S_{7x}	S_{5x}
S_{1u}	S_{1x}	S_{3x}	S_{7x}	S_{1u}	S_{3u}	S_{7u}	S_{5u}
S_{3x}	S_{3u}	S_{5u}	S_{1u}	S_{3x}	S_{5x}	S_{1x}	S_{7x}
S_{3u}	S_{3x}	S_{5x}	S_{1x}	S_{3u}	S_{5u}	S_{1u}	S_{7u}
S_{5x}	S_{5u}	S_{7u}	S_{3u}	S_{5x}	S_{7x}	S_{3x}	S_{1x}
S_{5u}	S_{5x}	S_{7x}	S_{3x}	S_{5u}	S_{7u}	S_{3u}	S_{1u}
S_{7x}	S_{7u}	S_{1u}	S_{5u}	S_{7x}	S_{1x}	S_{5x}	S_{3x}
S_{7u}	S_{7x}	S_{1x}	S_{5x}	S_{7u}	S_{1u}	S_{5u}	S_{3u}

S_4	$\frac{(x+iy)^2}{z(x-iy)^2}$	$\frac{L_x-iL_y}{(x+iy)(x-iy)z}$	$\frac{L_x+iL_y}{(x-iy)(x+iy)z}$	$L_x Z^2$
A	B	C_1	C_3	A
C_1	C_3	B	A	C_3
B	A	C_3	C_1	B
C_3	C_1	A	B	C_1
S_1	S_5	S_3	S_7	S_1
S_3	S_7	S_5	S_1	S_3
S_5	S_1	S_7	S_3	S_5
S_7	S_3	S_1	S_5	S_7

C_{3A}	z	$\frac{x+iy}{(x-iy)^2}$	$\frac{x-iy}{(x+iy)^2}$	$L_x Z^2$	$\frac{L_x+iL_y}{(x+iy)z}$	$\frac{L_x-iL_y}{(x-iy)z}$
A'	A''	C_1'	C_2''	A'	C_1''	C_2''
A''	A'	C_1''	C_2'	A''	C_1'	C_2'
C_1'	C_1''	C_2''	A'	C_1'	C_2''	A''
C_1''	C_1'	C_2'	A''	C_1''	C_2'	A'
C_2'	C_2''	A'	C_1'	C_2'	A''	C_1''
C_2''	C_2'	A''	C_1''	C_2''	A'	C_1'
S_1	S_7	S_9	S_5	S_1	S_3	S_{11}
S_3	S_9	S_{11}	S_7	S_3	S_5	S_1
S_5	S_{11}	S_1	S_9	S_5	S_7	S_3
S_7	S_1	S_3	S_{11}	S_7	S_9	S_5
S_9	S_3	S_5	S_1	S_9	S_{11}	S_7
S_{11}	S_5	S_7	S_3	S_{11}	S_1	S_9

TABLE 13. Selection rules—Continued

C_A	x, y	z	L_x, L_y xz, yz	L_z, Z^2 xy, ρ^2
A_z A_u	B_u B_z	A_u A_z	B_z B_u	A_z A_u
B_z B_u	A_u A_z	B_u B_z	A_z A_u	B_z B_u
S_{1z} S_{1u}	S_{3u} S_{3z}	S_{1u} S_{1z}	S_{3z} S_{3u}	S_{1z} S_{1u}
S_{3z} S_{3u}	S_{1u} S_{1z}	S_{3u} S_{3z}	S_{1z} S_{1u}	S_{3z} S_{3u}

O_A	r	L	Z^2, ρ^2	xz, yz, xy
A_{1z} A_{1u}	T_{1u} T_{1z}	T_{1z} T_{1u}	E_z E_u	T_{2z} T_{2u}
A_{2z} A_{2u}	T_{2u} T_{2z}	T_{2z} T_{2u}	E_z E_u	T_{1z} T_{1u}
E_z E_u	$T_{1u} T_{2u}$ $T_{1z} T_{2z}$	$T_{1z} T_{2z}$ $T_{1u} T_{2u}$	$A_{1z} A_{2z} E_z$ $A_{1u} A_{2u} E_u$	$T_{1z} T_{2z}$ $T_{1u} T_{2u}$
T_{1z} T_{1u}	$A_{1u} E_u T_{1u} T_{2u}$ $A_{1z} E_z T_{1z} T_{2z}$	$A_{1z} E_z T_{1z} T_{2z}$ $A_{1u} E_u T_{1u} T_{2u}$	$T_{1z} T_{2z}$ $T_{1u} T_{2u}$	$A_{2z} E_z T_{1z} T_{2z}$ $A_{2u} E_u T_{1u} T_{2u}$
T_{2z} T_{2u}	$A_{2u} E_u T_{1u} T_{2u}$ $A_{2z} E_z T_{1z} T_{2z}$	$A_{2z} E_z T_{1z} T_{2z}$ $A_{2u} E_u T_{1u} T_{2u}$	$T_{1z} T_{2z}$ $T_{1u} T_{2u}$	$A_{1z} E_z T_{1z} T_{2z}$ $A_{1u} E_u T_{1u} T_{2u}$
$D_{3/2z}$ $D_{3/2u}$	$D_{3/2u} D_{3/2z}$ $D_{3/2z} D_{3/2u}$	$D_{3/2z} D_{3/2u}$ $D_{3/2u} D_{3/2z}$	$D_{3/2z}$ $D_{3/2u}$	$2S_z D_{3/2z}$ $2S_u D_{3/2u}$
$2S_z$ $2S_u$	$2S_u D_{3/2u}$ $2S_z D_{3/2z}$	$2S_z D_{3/2z}$ $2S_u D_{3/2u}$	$D_{3/2z}$ $D_{3/2u}$	$D_{3/2z} D_{3/2z}$ $D_{3/2u} D_{3/2u}$
$D_{3/2z}$ $D_{3/2u}$	$2S_u D_{3/2u} D_{3/2z}$ $2S_z D_{3/2z} D_{3/2u}$	$2S_z D_{3/2z} D_{3/2u}$ $2S_u D_{3/2u} D_{3/2z}$	$2S_z D_{3/2z} D_{3/2z}$ $2S_u D_{3/2u} D_{3/2u}$	$2S_z D_{3/2z} D_{3/2z}$ $2S_u D_{3/2u} D_{3/2u}$

and the inverse relations

$$\mathbf{i} = \sin \theta \cos \phi \mathbf{r}_0 - \sin \theta \sin \phi \phi_0$$

$$\mathbf{j} = \sin \theta \sin \phi \mathbf{r}_0 + \cos \theta \sin \phi \theta_0 + \cos \theta \phi_0$$

$$\mathbf{k} = \cos \theta \mathbf{r}_0 - \sin \theta \theta_0$$

Let us further consider \mathbf{A} as a plane polarized wave, incident toward the origin along the vector \mathbf{r}_0 [\mathbf{r}_0 points to the source of \mathbf{A}], and having components along θ_0 and ϕ_0 perpendicular to the direction of propagation.

$$\mathbf{A} = (A \cos \psi \theta_0 + A \sin \psi \phi_0) \exp i(\sigma[\mathbf{r} \cdot \mathbf{r}_0] + \omega t).$$

In the terminology of the Euler angles of section 3.2, the plane polarized light is incident along the ζ axis with \mathbf{A} parallel to the ξ axis. At the origin,

\mathbf{A}^0 may be resolved into components as follows:

$$A_x^0 = A e^{i\omega t} (\cos \psi \cos \theta \cos \phi - \sin \psi \sin \phi)$$

$$A_y^0 = A e^{i\omega t} (\cos \psi \cos \theta \sin \phi + \sin \psi \cos \phi)$$

$$A_z^0 = A e^{i\omega t} (-\cos \psi \sin \theta).$$

The vector \mathbf{A} changes in space only along the direction \mathbf{r}_0 , and hence a variation along the x -axis, for example, will change \mathbf{A} according to the component of the variation along \mathbf{r}_0 . Formally,

$$\frac{\partial \mathbf{A}}{\partial x} = \frac{\partial \mathbf{A}}{\partial [\mathbf{r} \cdot \mathbf{r}_0]} \frac{\partial [\mathbf{r} \cdot \mathbf{r}_0]}{\partial x}.$$

Letting

$$\frac{\partial \mathbf{A}}{\partial [\mathbf{r} \cdot \mathbf{r}_0]} = \mathbf{A}',$$

TABLE 13. Selection rules—Continued

T_A	r	L xx, yz, xy	$Z^2 + i\rho^2$	$Z^2 - i\rho^2$
A_x A_y	T_x T_y	T_x T_y	C_{1x} C_{1y}	C_{2x} C_{2y}
C_{1x} C_{1y}	T_x T_y	T_x T_y	C_{2x} C_{2y}	A_x A_y
C_{2x} C_{2y}	T_x T_y	T_x T_y	A_x A_y	C_{1x} C_{1y}
T_x T_y	$A_x C_{1x} C_{2x} T_x$ $A_y C_{1y} C_{2y} T_y$	$A_x C_{1x} C_{2x} T_x$ $A_y C_{1y} C_{2y} T_y$	T_x T_y	T_x T_y
$D_{\frac{1}{2}x}$ $D_{\frac{1}{2}y}$	$D_{\frac{1}{2}x} {}_2S_{1x} {}_2S_{2x}$ $D_{\frac{1}{2}y} {}_2S_{1y} {}_2S_{2y}$	$D_{\frac{1}{2}x} {}_2S_{1x} {}_2S_{2x}$ $D_{\frac{1}{2}y} {}_2S_{1y} {}_2S_{2y}$	${}_2S_{1x}$ ${}_2S_{1y}$	${}_2S_{2x}$ ${}_2S_{2y}$
${}_2S_{1x}$ ${}_2S_{1y}$	$D_{\frac{1}{2}x} {}_2S_{1x} {}_2S_{2x}$ $D_{\frac{1}{2}y} {}_2S_{1y} {}_2S_{2y}$	$D_{\frac{1}{2}x} {}_2S_{1x} {}_2S_{2x}$ $D_{\frac{1}{2}y} {}_2S_{1y} {}_2S_{2y}$	${}_2S_{2x}$ ${}_2S_{2y}$	$D_{\frac{1}{2}x}$ $D_{\frac{1}{2}y}$
${}_2S_{2x}$ ${}_2S_{2y}$	$D_{\frac{1}{2}x} {}_2S_{1x} {}_2S_{2x}$ $D_{\frac{1}{2}y} {}_2S_{1y} {}_2S_{2y}$	$D_{\frac{1}{2}x} {}_2S_{1x} {}_2S_{2x}$ $D_{\frac{1}{2}y} {}_2S_{1y} {}_2S_{2y}$	$D_{\frac{1}{2}x}$ $D_{\frac{1}{2}y}$	${}_2S_{1x}$ ${}_2S_{1y}$

T_B	r xx, yz, xy	L	Z^2, ρ^2
A_1 A_2	T_2 T_1	T_1 T_2	E E
E T_1	$T_1 T_2$ $A_2 E T_1 T_2$	$T_1 T_2$ $A_1 E T_1 T_2$	$A_1 A_2 E$ $T_1 T_2$
T_2 $D_{\frac{1}{2}}$	$A_1 E T_1 T_2$ ${}_2S D_{3/2}$	$A_2 E T_1 T_2$ $D_{\frac{1}{2}} D_{3/2}$	$T_1 T_2$ $D_{3/2}$
${}_2S$ $D_{3/2}$	$D_{\frac{1}{2}} D_{3/2}$ $D_{\frac{1}{2}} {}_2S D_{3/2}$	${}_2S D_{3/2}$ $D_{\frac{1}{2}} {}_2S D_{3/2}$	$D_{3/2}$ $D_{\frac{1}{2}} {}_2S D_{3/2}$

we have

$$\frac{\partial A}{\partial x} = A'(\mathbf{i} \cdot \mathbf{r}_0), \quad \frac{\partial A}{\partial y} = A'(\mathbf{j} \cdot \mathbf{r}_0), \quad \frac{\partial A}{\partial z} = A'(\mathbf{k} \cdot \mathbf{r}_0).$$

Obviously, the resolution of A' into components is parallel to that of A , so we obtain the following 9 terms involving the derivatives of A :

$$\frac{\partial A_x}{\partial x} = A'(\cos \psi \cos \theta \cos \phi - \sin \psi \sin \phi)(\sin \theta \cos \phi)$$

$$\frac{\partial A_x}{\partial y} = A'(\cos \psi \cos \theta \cos \phi - \sin \psi \sin \phi)(\sin \theta \sin \phi)$$

$$\frac{\partial A_x}{\partial z} = A'(\cos \psi \cos \theta \cos \phi - \sin \psi \sin \phi)(\cos \theta)$$

$$\frac{\partial A_y}{\partial x} = A'(\cos \psi \cos \theta \sin \phi + \sin \psi \cos \phi)(\sin \theta \cos \phi)$$

$$\frac{\partial A_y}{\partial y} = A'(\cos \psi \cos \theta \sin \phi + \sin \psi \cos \phi)(\sin \theta \sin \phi)$$

$$\frac{\partial A_y}{\partial z} = A'(\cos \psi \cos \theta \sin \phi + \sin \psi \cos \phi)(\cos \theta)$$

$$\frac{\partial A_z}{\partial x} = A'(-\cos \psi \sin \theta)(\sin \theta \cos \phi)$$

$$\frac{\partial A_z}{\partial y} = A'(-\cos \psi \sin \theta)(\sin \theta \sin \phi)$$

$$\frac{\partial A_z}{\partial z} = A'(-\cos \psi \sin \theta)(\cos \theta).$$

These may now be combined as previously defined (page 55) to yield

$$A_{xx} = \frac{1}{2} A'(\cos \psi \sin 2\theta \cos^2 \phi - \sin \psi \sin \theta \sin 2\phi)$$

$$A_{yy} = \frac{1}{2} A'(\cos \psi \sin 2\theta \sin^2 \phi + \sin \psi \sin \theta \sin 2\phi)$$

$$A_{zz} = \frac{1}{2} A'(-\cos \psi \sin 2\theta)$$

$$A_{xy} = \frac{1}{2} A'(\cos \psi \sin 2\theta \sin 2\phi + 2 \sin \psi \sin \theta \cos 2\phi)$$

$$A_{xz} = A'(\cos \psi \cos 2\theta \cos \phi - \sin \psi \cos \theta \sin \phi)$$

$$A_{yz} = A'(\cos \psi \cos 2\theta \sin \phi + \sin \psi \cos \theta \cos \phi)$$

$$B_x = -A'(\cos \psi \sin \phi + \sin \psi \cos \theta \cos \phi)$$

$$B_y = A'(\cos \psi \cos \phi - \sin \psi \cos \theta \sin \phi)$$

$$B_z = A'(\sin \psi \sin \theta).$$

We may note that $A_{xx} + A_{yy} + A_{zz}$ is in fact zero, and that the angular factors for the components of B are the same as those for A with a rotation of 90° in ψ , showing the mutual perpendicularity of B and A . In order to obtain our final five quadrupole components, there is a question of normalization which must be considered. Strictly speaking, the appropriate "base vectors" for the quadrupole components are the normalized tesseral harmonics of table 2 for $N=2$. Since we shall be interested in relative intensities, we may omit a common factor, which we shall here choose to be $\frac{\sqrt{15}}{2\sqrt{\pi}}$. With the omission of this factor, xy , yz , and z^2 will still be the proper basis, but for the other 2, where we have heretofore used $x^2 - y^2$ and $2z^2$

$-x^2-y^2$, we must now include factors of $\frac{1}{2}$ and $\frac{1}{2\sqrt{3}}$ respectively, so the balance of the term with these two bases will become multiplied by the reciprocals of these factors. We shall continue to denote these two renormalized bases as ρ^2 and Z^2 .

Finally, we must consider the resolution of the multipole interactions into the (complex) base vectors appropriate to the group $C_{\infty h}$ and its subgroups with complex reps. The terms in z , L_x , and Z^2 will be unaffected by this change. In lieu of x and y , we shall use $\frac{1}{\sqrt{2}}(x+iy)$ and $\frac{1}{\sqrt{2}}(x-iy)$, for L_x and L_y a similar combination, and for ρ^2 and xy , $\frac{1}{\sqrt{2}}(\rho^2+ixy)$ and $\frac{1}{\sqrt{2}}(\rho^2-ixy)$. This requires the use of the following linear combinations of the previous terms:

$$\begin{aligned} A_+^0 &= \frac{\sqrt{2}}{2} (A_x^0 - iA_y^0) & A_-^0 &= \frac{\sqrt{2}}{2} (A_x^0 + iA_y^0) \\ B_+^0 &= \frac{\sqrt{2}}{2} (B_x^0 - iB_y^0) & B_-^0 &= \frac{\sqrt{2}}{2} (B_x^0 + iB_y^0) \\ A_{x+} &= \frac{\sqrt{2}}{2} (A_{xz} - iA_{yz}) & A_{x-} &= \frac{\sqrt{2}}{2} (A_{xz} + iA_{yz}) \\ A_{2+} &= \frac{\sqrt{2}}{2} (A_{\rho^2} - iA_{xy}) & A_{2-} &= \frac{\sqrt{2}}{2} (A_{\rho^2} + iA_{xy}) \end{aligned}$$

These are all summarized in table 14.

Let us now consider some of the conclusions of this analysis. We shall discuss in detail six cases.

1. Only one component of the eleven considered is effective in producing the transition. In this case, the rate of absorption will be proportional to the Hermitian square of the matrix element, and hence proportional to the Hermitian square of the appropriate angular dependence factor of table 14. Such a case is exemplified by the $B_{1g}-B_{2g}$ transition in D_{4h} , which are joined only by L_x . The angular dependence is thus $(\sin \psi \sin \theta)^2$.

2. Two components only of the eleven join the states, these two belonging to a doubly degenerate rep and consequently arising from the same type of multipole interaction (e.g., in D_{4h} , A_{1g} is joined to E_{2g} by both xy and ρ^2). Then the total matrix element will involve the sum of the two angular factors for the two multipole components, and the angular dependence will be proportional to the Hermitian square of the sum of these angular factors.

3. Two components only join the states involved, arising from different multipole interactions (e.g., A_1 of $C_{\infty h}$ will join itself by either z or Z^2). The matrix element will then include the sum of the angular factors for the two components, but one will be multiplied by a factor "q" determined by the numerical ratio of the strength of

TABLE 14. Angular dependence of multipole components

1. Electric Dipole	
$A_+^0 = A(-\cos \psi \sin \theta)$	
$A_x^0 = A(\cos \psi \cos \theta \cos \phi - \sin \psi \sin \phi)$	
$A_y^0 = A(\cos \psi \cos \theta \sin \phi + \sin \psi \cos \phi)$	
$A_+^1 = A\left(\frac{\sqrt{2}}{2}\right)(\cos \psi \cos \theta - i \sin \psi)e^{-i\phi}$	
$A_-^1 = A\left(\frac{\sqrt{2}}{2}\right)(\cos \psi \cos \theta + i \sin \psi)e^{i\phi}$	
2. Magnetic Dipole	
$B_x = A'(\sin \psi \sin \theta)$	
$B_y = A'(-\cos \psi \sin \phi - \sin \psi \cos \theta \cos \phi)$	
$B_z = A'(\cos \psi \cos \phi - \sin \psi \cos \theta \sin \phi)$	
$B_+ = A'\left(\frac{\sqrt{2}}{2}\right)(-\sin \psi \cos \theta - i \cos \psi)e^{-i\phi}$	
$B_- = A'\left(\frac{\sqrt{2}}{2}\right)(-\sin \psi \cos \theta + i \cos \psi)e^{i\phi}$	
3. Electric Quadrupole	
$A_{zz} = A'\left(\frac{\sqrt{3}}{2}\right)(-\cos \psi \sin 2\theta)$	
$A_{xx} = A'(\cos \psi \cos 2\theta \cos \phi - \sin \psi \cos \theta \sin \phi)$	
$A_{yy} = A'(\cos \psi \cos 2\theta \sin \phi + \sin \psi \cos \theta \cos \phi)$	
$A_{\rho^2} = \frac{1}{2}A'(\cos \psi \sin 2\theta \cos 2\phi - 2 \sin \psi \sin \theta \sin 2\phi)$	
$A_{xy} = \frac{1}{2}A'(\cos \psi \sin 2\theta \sin 2\phi + 2 \sin \psi \sin \theta \cos 2\phi)$	
$A_{z+} = A'\left(\frac{\sqrt{2}}{2}\right)(\cos \psi \cos 2\theta - i \sin \psi \cos \theta)e^{-i\phi}$	
$A_{z-} = A'\left(\frac{\sqrt{2}}{2}\right)(\cos \psi \cos 2\theta + i \sin \psi \cos \theta)e^{i\phi}$	
$A_{x+} = A'\left(\frac{\sqrt{2}}{4}\right)(\cos \psi \sin 2\theta - 2i \sin \psi \sin \theta)e^{-2i\phi}$	
$A_{x-} = A'\left(\frac{\sqrt{2}}{4}\right)(\cos \psi \sin 2\theta + 2i \sin \psi \sin \theta)e^{2i\phi}$	

the two types of interaction, and is fixed by explicit calculations and not by general symmetry considerations. The angular variation will then be proportional to the Hermitian square of this sum, and will contain q and q^2 . If one type of interaction is very much stronger than the other, then q will be small and it may be neglected. In the present problem of crystal spectra, this is not necessarily the case, and each case must be examined in detail.

4. Two components join the states in question, arising from the same multipole interaction, but the components belong to one-dimensional reps (as distinguished from (2)). For example, in D_{2h} , A_{2g} is joined to itself by both Z^2 and ρ^2 . The details here are very similar to case (3). The matrix element will include the sum of the angular factors. However, in contrast to (1), the term Z^2 is not related by symmetry to ρ^2 , so one angular factor will again be multiplied by a factor q , expressing the relative strength of the interaction, and the Hermitian square will again contain both q and q^2 . Generally, one may only expect that q will in this case be nearer 1 than in case (3).

5. Only one component joins the states, but another component joins another pair of states, both of which are respectively degenerate with the first pair. This is a frequent situation in

$C_{\infty h}$ and its subgroups. Here the transitions proceed independently, and the total absorption will be proportional to the sum of the Hermitian squares of the angular factors, rather than to the Hermitian square of the sum as in previous cases.

6. If the states are joined by three components, all belonging to the same triply degenerate rep, as is the case for both electric and magnetic dipole transitions in the cubic groups, there are

no polarization effects.

We shall not discuss in detail more complicated cases. In general, the angular dependence will include factors expressing the relative strengths of the transitions arising from different dipole interactions, as well as from nondegenerate interactions of the same order. In specific cases, these relative strengths may be computed from the methods of the following section.

5. Matrix Elements

5.1. The Reduced Matrix Element

We now turn our attention to the procedure alternate to that which has occupied our attention in the preceding two sections, namely, the calculation of the matrix elements of the perturbing potentials of the crystal field and their interpretation. These potentials were discussed in detail in section 2. We have already touched upon certain general features of the problem in section 4.3—we must now examine this situation in more detail, and also determine the simplifying relationships which exist among the nonvanishing elements.

So far in our discussion of the transformation properties of the several wave functions belonging to a degenerate rep of a group, we have emphasized the equivalence of these several wave functions, as well as the properties of the rep (notably the character and class structure) which were invariant under a unitary transformation. We shall now depart from this procedure. We are now interested in picking out a particular set of linearly independent wave functions, in terms of which specific matrices representing the operations of the group may be written down. This is equivalent to the introduction of a specific system of base vectors in ordinary vector analysis.

Since we start with the wave functions of the free ion, described by the quantum number J , with $2J+1$ linearly independent wave functions, we select for our basis, functions distinguished by the quantum number M (or M_J will sometimes be used for clarity) where $M=J, J-1, J-2, \dots, -J$, and such that the matrix representing rotations about the z -axis is diagonal with elements $Me^{iM\phi}$. We have seen from the preceding section that wave functions belonging to different reps (here, with different J values) are orthogonal—we see that these $2J+1$ functions themselves belong to different reps of the group $C_{\infty h}$, and hence are mutually orthogonal within a given J . These wave functions we can write as $|\alpha JM\rangle$ (in the notation of Dirac), where α represents all quantum numbers other than J and M . To emphasize that such a state is one component of a $2J+1$ dimension rep of the rotation group, we may also write in the form $(\alpha J)|JM\rangle$. A different choice of axes will leave the (scalar) portion of the function unaltered, but will convert the normalized com-

ponent of the rep $|JM\rangle$ into a combination of the same J and different M .

For these calculations, it will be convenient to use the complex potentials Y_{ν}^{μ} (2.9) rather than the real C_{ν}^{μ} and S_{ν}^{μ} (2.5) as our perturbation. We note here that these perturbing potentials actually act upon the space coordinates of the individual electrons, so the total perturbation is a sum of such over all electrons. Clearly, if these potentials obey a certain transformation law in the coordinates of one electron, their sum will similarly transform when all the coordinates of all electrons undergo the simultaneous transformations implied by the quantum number J , or when only the space coordinates of all electrons undergo the simultaneous transformation implied by the quantum number L . Our problem has therefore become that of computing the sum of terms of the form

$$\langle \alpha' J' M' | r^{\nu} Y_{\nu}^{\mu} | \alpha J M \rangle \quad (5.1)$$

We shall for the time being disregard the factor r^{ν} .

According to section 4.3, we know that the matrix element will vanish unless J' is one of the values J'' arising from the reduction of the product $D_J \times D_{\nu}$, namely, $J+\nu, J+\nu-1, \dots, |J-\nu|$. Let us expand the product $Y_{\nu}^{\mu} | \alpha J M \rangle$ into a sum of terms

$$\sum_{M''} C_{M''}^{J''} Y_{\nu}^{\mu} | \alpha J \rangle | J'' M'' \rangle \quad (5.2)$$

where the C -coefficients are the Wigner coefficients to be discussed further in section 5.2, the term $(Y_{\nu}^{\mu} | \alpha J)$ represents the portion of the product invariant under a rotation of the axes, and $| J'' M'' \rangle$ represents the M'' -th component of the normalized rep $D_{J''}$ of the rotation group. We now know that the remaining term of (5.1), $\langle \alpha' J' M' | = (\alpha' J') \langle J' M' |$ is orthogonal to all states except $| J' M' \rangle$, i.e.,

$$\langle J' M' | J'' M'' \rangle = \delta_{J', J''} \delta_{M', M''} \quad (5.3)$$

so we obtain, from (5.2) and (5.3),

$$\langle \alpha' J' M' | Y_{\nu}^{\mu} | \alpha J M \rangle = C_{M''}^{J''} (\alpha' J') (Y_{\nu}^{\mu} | \alpha J). \quad (5.4)$$

In this expression, parity is included in the α, α' . If ν is even, α and α' must be of the same parity; if odd, of opposite parity.

The importance of this step is that we have placed into the term $C_{M'M}^{\alpha J}$, the dependence of the matrix element on M , M' , and μ , and the "reduced matrix element" $\langle \alpha' J' || Y_\mu || \alpha J \rangle$ contains those portions of the interaction not dependent on these quantum numbers. If the matrix element is evaluated for one particular M , M' , and μ , then the reduced matrix element can be obtained, and from it the matrix element for any other M , M' , and μ . This relation was initially given by Eckart (1930) and again by Wigner (1931) and Racah (1942a). There is some divergence in the literature on normalization and phase factors for these reduced matrix elements. We have here followed Eckart, Wigner, Rose, and others, while Racah, Biedenharn (1952), and Simon et al. (1954) have used as reduced matrix element a quantity greater by a factor $\sqrt{2J'+1}$.

Eckart and Wigner have based their derivation of this result on the ideas of the representations of the rotation group, while Racah has developed the algebraic application of the angular momentum operators J_x , J_y , and J_z , which may be interpreted as differential rotations about the x , y , or z -axes, respectively. Since the vectorial character of a vector operator T with components T^n , T^0 , T^{-1} is fixed by the general commutation rules $[J_z, T^n] = nT^n$ and $[J_x \pm iJ_y, T^n] = \sqrt{(1 \mp n)(1 \pm n + 1)} T^{n \pm 1}$, a generalized irreducible tensor operator of degree k has been defined as one having $2k+1$ components and satisfying the commutation rules $[J_z, T_k^n] = nT_k^n$ and $[J_x \pm iJ_y, T_k^n] = \sqrt{(k \mp n)(k \pm n + 1)} T_k^{n \pm 1}$. Our quantity Y_μ^J is a specific example of such an operator.

Stevens (1952) has emphasized a geometrical interpretation of this result, and has given what amounts to a technique for calculating certain of the Wigner coefficients in the special case $J'=J$. The behavior under rotation of any quantity which may be expressed in terms of variously oriented coordinate systems, may be described by one or more components along a set of mutually orthogonal base vectors in an infinite number of dimensions. Subspaces of dimension 1, 2, 3, . . . $2J+1$ determine a particular D_J , while the individual vectors within the subspaces correspond to the various rows of the D_J . A product of two components, each of which lies along one of these base vectors (such as the $Y_\mu^J | \alpha J M \rangle$ being considered) may itself be resolved into components by means of the Wigner coefficients; all save the one parallel to the third quantity $\langle \alpha' J' M' |$ will vanish in the matrix element. Stevens also pointed out that it is possible to form irreducible tensors J_ν^J from the noncommuting components of J by taking the $2\nu+1$ totally symmetric products of degree ν . This is analogous to the process we have already used several times in connection with the reduction of the general product of two vectors. We then have

$$\langle \alpha' J' M' | J_\nu^J | \alpha J M \rangle = C_{M'M}^{\alpha J \nu} (J || J_\nu || J),$$

since this reduced matrix element is diagonal in both α and J , and independent of α . These matrix elements may themselves be calculated, giving a set of numbers proportional to the Wigner coefficients. Certain tables based on this procedure have been given by Stevens (1952), by Elliott and Stevens (1953a), and by Judd (1955).

5.2. Wigner Coefficients

The Wigner coefficients $C_{M_1 M_2 M}^{\alpha J}$, frequently written $(J_1 J_2 J M | J_1 J_2 M_1 M_2)$ are the elements of the unitary matrix C which will reduce the $(2J_1+1) \times (2J_2+1)$ dimensional direct product of the components of D_{J_1} and D_{J_2} and which will provide the proper linear combinations of the products of these components $|J_1 M_1 \rangle$ and $|J_2 M_2 \rangle$ appropriate to the reduced component $|J M \rangle$. The columns of the C matrix are distinguished by the double indices (M_1, M_2) , while the rows bear the indices (J, M) . The phase conventions implied in (2.8) ensure that the matrix elements may be chosen to be real. The inverse C^{-1} is merely the transposed matrix. The orthogonality relations for a real unitary (=orthogonal) matrix then give us

$$\sum_{M_1, M_2} (J_1 J_2 M_1 M_2 | J_1 J_2 J M) (J_1 J_2 M_1 M_2 | J_1 J_2 J' M') = \delta_{J J'} \delta_{M M'} \quad (5.5a)$$

$$\sum_{J, M} (J_1 J_2 M_1 M_2 | J_1 J_2 J M) (J_1 J_2 M_1' M_2' | J_1 J_2 J M) = \delta_{M_1 M_1'} \delta_{M_2 M_2'} \quad (5.5b)$$

The product of $|J_1 M_1 \rangle$ and $|J_2 M_2 \rangle$ will, under a rotation about the z -axis through ϕ , be multiplied by $e^{iM_1 \phi} e^{iM_2 \phi} = e^{i(M_1 + M_2) \phi}$ and hence $M = M_1 + M_2$, and all coefficients not meeting this requirement are zero. The index M is frequently suppressed when written in the form $C_{M_1 M_2}^{\alpha J}$. Because of this relation between the M_1 and M_2 , the sum in (5.5a) is in effect a sum over M_1 only (or M_2 only).

A general formula for these coefficients has been given by Wigner, who carried out in detail the process outlined above of reducing the direct product matrix. Condon and Shortley have approached the problem from the standpoint of transforming eigenstates of two commuting angular momenta J_1 and J_2 with quantum numbers $J_1 M_1$ and $J_2 M_2$ into eigenstates of $J = J_1 + J_2$ with quantum numbers J and M . They pointed out that the vector addition coefficients may in principle be obtained from the initial condition $(J_1 J_2 J_1 J_2 | J_1 J_2 [J_1 + J_2] [J_1 + J_2]) = 1$ and from a successive application of the $J_x - iJ_y$ and $J_x + iJ_y$ operators, but that a general formula is difficult to obtain. Racah (1942a) utilizing as well the operator $J_x + iJ_y$ has provided a purely algebraic derivation of a general formula for the coefficients which is relatively convenient and symmetric in the various parameters. Based upon this for-

mula, Simon (1954) has provided a numerical table of these coefficients, but no J greater than $\frac{7}{2}$ is included. Condon and Shortley have given formulas for the cases where $J_2 = \frac{1}{2}, 1, \frac{3}{2},$ and $2,$ while Falkoff et al. (1952) give $J_2 = 3.$

For actual computations, a numerical table of Wigner coefficients is required. The only such table known to the writer is that of Simon (1954). Unfortunately, this covers an inadequate range of values for rare-earth spectra, and relies on symmetry properties of the Wigner coefficients, involving factors of the form $\left(\frac{2J+1}{2J_1+1}\right)^{1/2},$ which are usually somewhat awkward. To meet partially this need for the present application, table 15, containing the coefficients $C_{M\mu}^{J\nu J}$ to 6 decimals for $J \leq 8, \nu = 2, 4, 6$ and $\mu = 0, 2, 3, 4, 6$ consistent with the ν involved has been computed. These will permit calculation of matrix elements diagonal in J and between states of the same parity.

In the case $\mu = 0, 3,$ and $6,$ the values for $M = J - \mu$ were calculated from the formula of Racah and those for other values of M were calculated by the use of the tables of Stevens and others noted above (section 5.1). For $\mu = 4$ and $\nu = 4, \mu = 2,$ factors corresponding to those of Stevens were computed and used in a similar process. For $\mu = 2, \nu = 6$ this became too unwieldy and the formula of Racah was used directly for all entries. Coefficients were all checked for obedience to the requirement

$$\sum_M (C_{M\mu}^{J\nu J})^2 = \frac{2J+1}{2\nu+1}$$

and selected entries were checked for

$$\sum_M (C_{M\mu}^{J\nu J})(C_{M\mu}^{J\nu J}) = 0.$$

It is believed that errors do not exceed 2 in the sixth decimal. In a very few entries a seventh decimal was carried.

It is to be noted that only positive values of μ are given, and only entries for $M \geq -\frac{\mu}{2}.$ To obtain other values required, the following special cases of the general symmetry relations are required (diagonal in J and ν even):

$$C_{-M-\mu}^{J\nu J} = C_{M\mu}^{J\nu J} \quad (5.6a)$$

$$C_{(M+\mu)-\nu}^{J\nu J} = (-1)^\mu C_{M\mu}^{J\nu J} \quad (5.6b)$$

$$C_{-(M+\mu)\mu}^{J\nu J} = (-1)^\mu C_{M\mu}^{J\nu J}. \quad (5.6c)$$

The relation (5.6c) is especially noteworthy: we observe that (for half-integral J)

$$C_{-\frac{1}{2}\frac{1}{2}}^{J\nu J} = (-1)^3 C_{-\frac{3}{2}\frac{3}{2}}^{J\nu J}$$

and hence must be zero. This is intimately associated with the essential degeneracy of Kramers discussed earlier.

TABLE 15. Wigner coefficients

		$C_{M\mu}^{J\nu J}$									
$J \backslash M$	0	1	2	3	4	5	6	7	8		
1	-0.632456	0.316228									
2	-.534522	-.267261	0.534522								
3	-.516398	-.387298	0	0.645497							
4	-.509647	-.433200	-.203859	.178376	0.713506						
5	-.506369	-.455732	-.303822	-.050637	.303822	0.759554					
6	-.504525	-.468488	-.360375	-.180188	.072075	.396412	0.792825				
7	-.503382	-.476415	-.395515	-.260680	-.071912	.170790	.467426	0.817996			
8	-.502625	-.481682	-.418854	-.314140	-.167542	.020943	.251312	.523568	0.837708		
$J \backslash M$	1/2	3/2	5/2	7/2	9/2	11/2	13/2	15/2			
3/2	-0.447214	0.447214									
5/2	-.478091	-.119523	0.597614								
7/2	-.487950	-.292770	.097590	0.683130							
9/2	-.492366	-.369274	-.123092	.246183	0.738549						
11/2	-.494728	-.409917	-.240296	.014135	.353377	0.777429					
13/2	-.496139	-.434122	-.310087	-.124035	.124035	.434122	0.806228				
15/2	-.497050	-.449712	-.355036	-.213022	-.023669	.213022	.497050	0.828417			

TABLE 15. Wigner coefficients—Continued

$C_{M_0}^{JM}$

$J \backslash M$	0	1	2	3	4	5	6	7	8
2	0. 534523	-0. 356348	0. 089087,						
3	. 426402	. 071067	-. 497469	0. 213201					
4	. 402291	. 201146	-. 245844	-. 469340	0. 312893				
5	. 392232	. 261488	-. 065372	-. 392232	-. 392232	0. 392232			
6	. 386953	. 294821	. 050672	-. 248756	-. 442232	-. 304035	0. 456052		
7	. 383807	. 315270	. 127428	-. 126413	-. 357408	-. 441175	-. 217795	0. 508189	
8	. 381771	. 328747	. 180281	-. 031814	-. 254514	-. 413585	-. 413585	-. 137862	0. 551447
$J \backslash M$	1/2	3/2	5/2	7/2	9/2	11/2	13/2	15/2	
5/2	0. 308606	-0. 462909	0. 154303						
7/2	. 341882	-. 113961	-. 493829	0. 265908					
9/2	. 354787	. 059131	-. 335077	-. 433629	0. 354787				
11/2	. 361298	. 154842	-. 167745	-. 425815	-. 348394	0. 425815			
13/2	. 365073	. 212959	-. 043944	-. 310988	-. 446201	-. 260284	0. 483384		
15/2	. 367467	. 250811	. 044718	-. 196371	-. 390798	-. 429684	-. 176929	0. 530786	

$C_{M_0}^{J_0 J}$

$J \backslash M$	0	1	2	3	4	5	6	7	8
3	-0. 482804	0. 362103	-0. 144841	0. 024140 ₂					
4	-. 373979	. 018699	. 411377	-. 317882	0. 074795 ₈				
5	-. 347368	-. 104210	. 312631	. 251842	-. 416842	0. 130263			
6	-. 335531	-. 167765	. 184542	. 360696	. 067106	-. 461355	0. 184542		
7	-. 329015	-. 205635	. 082254	. 324080	. 289534	-. 090479	-. 470492	0. 235246	
8	-. 324985	-. 230198	. 005416	. 251864	. 346651	. 176034	-. 211240	-. 457688	0. 281654
$J \backslash M$	1/2	3/2	5/2	7/2	9/2	11/2	13/2	15/2	
7/2	-0. 241402	0. 434524	-0. 241402	0. 048280 ₆					
9/2	-. 273115	. 204836	. 341393	-. 375533	0. 102418				
11/2	-. 286829	. 057366	. 358536	. 157756	-. 444585	0. 157756			
13/2	-. 294280	. 036785	. 272209	. 334007	-. 016185	-. 469376	0. 210410		
15/2	-. 298843	-. 099614	. 179306	. 346658	. 235090	-. 155398	-. 466195	0. 258997	

TABLE 15. Wigner coefficients—Continued

$C_{M^2}^{JM}$

$J \backslash M$	-1	0	1	2	3	4	5	6
1	0. 774597							
2	. 654653	0. 534522						
3	. 632455	. 577350	0. 408248					
4	. 624188	. 592156	. 495434	0. 330289				
5	. 620173	. 599144	. 535891	. 429669	0. 277350			
6	. 617915	. 603023	. 558291	. 483494	. 377965	0. 239046		
7	. 616515	. 605405	. 572054	. 516377	. 438160	. 336817	0. 210042	
8	. 615587	. 606977	. 581136	. 538028	. 477567	. 399561	. 303489	0. 187317

$J \backslash M$	-1/2	1/2	3/2	5/2	7/2	9/2	11/2	
3/2	0. 632456							
5/2	. 621059	0. 462910						
7/2	. 617213	. 534522	0. 365148					
9/2	. 615457	. 564075	. 460566	0. 301511				
11/2	. 614509	. 579365	. 508862	. 402291	0. 256776			
13/2	. 613941	. 588349	. 537087	. 459933	. 356263	0. 223607		
15/2	. 613572	. 594088	. 555088	. 496486	. 418079	. 319313	0. 198030	

$C_{M^2}^{JM}$

$J \backslash M$	-1	0	1	2	3	4	5	6
2	-0. 563436	0. 345033						
3	-. 449467	-. 123091	0. 302233					
4	-. 424052	-. 245844	. 136989	0. 560968				
5	-. 413449	-. 299572	0	. 358057	0. 554700			
6	-. 407884	-. 328395	-. 110558	. 183513	. 449089	0. 532554		
7	-. 404568	-. 345779	-. 180745	. 056476	. 308827	. 495260	0. 505389	
8	-. 402422	-. 357114	-. 227940	-. 035172	. 187317	. 391802	. 515831	0. 477567

$J \backslash M$	-1/2	1/2	3/2	5/2	7/2	9/2	11/2	
5/2	-0. 345033	0. 462910						
7/2	-. 372194	. 053722	0. 550482					
9/2	-. 381690	-. 116608	. 285631	0. 560968				
11/2	-. 386244	-. 204837	. 099950	. 410891	0. 544705			
13/2	-. 388809	-. 256681	-. 022676	. 252439	. 476312	0. 519238		
15/2	-. 390406	-. 289807	-. 105958	. 126363	. 354689	. 507935	. 491412	

TABLE 15. Wigner coefficients—Continued

C_{JM}^{JJ}

$J \backslash M$	-1	0	1	2	3	4	5	6
3	0.494727	-0.361298	0.127738					
4	.383214	0	-.434524	0.289683				
5	.355945	.122813	-.302079	-.308257	0.397959			
6	.343817	.184542	-.163087	-.376632	-.147214	0.465530		
7	.337140	.220710	-.058394	-.316265	-.335450	0	0.505389	
8	.333010	.243919	.017964	-.232843	-.361685	-.247027	.121960	0.526926

$J \backslash M$	-1/2	1/2	3/2	5/2	7/2	9/2	11/2
7/2	.255476	-.442498	.215917				
9/2	.285631	-.186989	-.381691	.349825			
11/2	.298081	-.035129	-.359967	-.227663	.435942		
13/2	.304672	.058394	-.255871	-.365192	-.070719	.488252	
15/2	.308643	.119537	-.156366	-.349645	-.294427	.064249	.517994

C_{JM}^{JJ}

$J \backslash M$	-1	0	1	2	3	4	5
2	-0.527046						
3	-.265908	-0.564076					
4	-.186989	-.469339	-0.494727				
5	-.146176	-.392232	-.506369	-0.423659			
6	-.120654	-.335111	-.472034	-.490098	-0.363803		
7	-.102998	-.291972	-.431342	-.493065	-.457141	-0.315165	
8	-.089984	-.258460	-.393234	-.474495	-.486664	-.420595	-0.275723

$J \backslash M$	-1/2	1/2	3/2	5/2	7/2	9/2
5/2	-0.577350					
7/2	-.402015	-0.531816				
9/2	-.311649	-.498639	-0.458029			
11/2	-.255476	-.442497	-.501745	-0.392232		
13/2	-.216867	-.391883	-.487398	-.474594	-0.338200	
15/2	-.188584	-.349644	-.457792	-.492112	-.438906	-0.294427

C_{JM}^{JJ}

$J \backslash M$	-1	0	1	2	3	4	5
3	0.469340	-0.221249					
4	.270973	.323875	-0.409673				
5	.201802	.376036	.058255	-0.487398			
6	.163087	.360696	.248129	-.147214	-0.509963		
7	.137637	.332940	.322787	.079066	-.258078	-0.505389	
8	.119407	.304863	.347875	.209882	-.071755	-.372080	-0.487838

TABLE 15. Wigner coefficients—Continued

$J \backslash M$	-1/2	1/2	3/2	5/2	7/2	9/2	
7/2	0. 442498	-0. 334497					
9/2	. 373979	. 186990	-0. 458029				
11/2	. 316163	. 324509	-. 053661	-0. 503382			
13/2	. 272507	. 357008	. 163319	-. 223634	-0. 509963		
15/2	. 239074	. 354605	. 270834	0	-. 333849	-0. 497673	

$C_{M'}^{JM}$

$J \backslash M$	-2	-1	0	1	2	3	4
2	0. 745356						
3	. 594588	0. 460566					
4	. 560968	. 494727	0. 312893				
5	. 546941	. 506369	. 392232	0. 226455			
6	. 539583	. 511894	. 432629	. 313114	0. 171499		
7	. 535193	. 514990	. 456490	. 366006	. 253968	0. 134387	
8	. 532354	. 516920	. 471881	. 401022	. 310630	. 209427	0. 108148
$J \backslash M$	-3/2	-1/2	1/2	3/2	5/2	7/2	
5/2	0. 577350						
7/2	. 550482	0. 376051					
9/2	. 539792	. 440738	0. 264443				
11/2	. 534366	. 469339	. 349825	0. 196116			
13/2	. 531214	. 484930	. 397958	. 281399	0. 151247		
15/2	. 529213	. 494472	. 428225	. 336926	. 230164	0. 120199	

$C_{M'}^{JM}$

$J \backslash M$	-2	-1	0	1	2	3	4
3	-0. 541947	0. 349825					
4	-. 419790	-. 079333	0. 501745				
5	-. 389919	-. 195539	. 233021	0. 504505			
6	-. 376632	-. 248128	. 067106	. 382472	0. 465530		
7	-. 369318	-. 277194	-. 037801	. 247517	. 441640	0. 417311	
8	-. 364795	-. 295182	-. 107785	. 137400	. 354765	. 454447	0. 370540
$J \backslash M$	-3/2	-1/2	1/2	3/2	5/2	7/2	
7/2	-0. 312893	0. 458029					
9/2	-. 334497	. 102418	0. 512092				
11/2	-. 341495	-. 055544	. 322921	0. 487398			
13/2	-. 344698	-. 141599	. 167849	. 419971	0. 441641		
15/2	-. 346452	-. 194225	. 056068	. 308798	. 452034	0. 393445	

TABLE 15. Wigner coefficients—Continued

		C_{JM}^{JM}					
$J \backslash M$		-3	-2	-1	0	1	2
3		0. 733799					
4		. 568399	0. 429669				
5		. 527952	. 457220	0. 273241			
6		. 509963	. 465530	. 345246	0. 184542		
7		. 500060	. 469098	. 383017	. 260982	0. 130491	
8		. 493934	. 470948	. 405906	. 310016	. 200114	0. 095673 ₆
$J \backslash M$		-5/2	-3/2	-1/2	1/2	3/2	
7/2		0. 554700					
9/2		. 518874	0. 339683				
11/2		. 503381	. 397958	0. 223100			
13/2		. 495034	. 423922	. 299758	0. 154399		
15/2		. 489956	. 438230	. 344798	. 228062	0. 111283	

5.3. L-S Coupling and the One-Electron Approximation

It is well to emphasize at this point that the relation (5.4) involves no approximations—these we shall introduce in evaluating the reduced matrix element. The first of these is the approximation of L-S coupling, i.e., the state $|\alpha JM\rangle$ may be written $|\alpha JM\rangle = \sum_{M_L} |\beta LSM_L M_S\rangle (LSM_L M_S | LSJM)$, and consequently,

$$\langle \alpha J' M' | Y^{\dagger} | \alpha JM \rangle = \sum_{M_L M_L'} (L' S' J' M' | L' S' M_L' M_S') \langle \beta' L' S' M_L' M_S' | Y^{\dagger} | \beta L S M_L M_S \rangle (LSM_L M_S | LSJM) \quad (5.7)$$

We now recall that Y^{\dagger} does not act upon the spin coordinates of the system, that it is thus a scalar quantity with respect to S, and therefore

$$\langle \beta' L' S' M_L' M_S' | Y^{\dagger} | \beta L S M_L M_S \rangle = \delta_{SS'} \delta_{M_L M_S'} \langle \beta' L' S' M_L' M_S' | Y^{\dagger} | \beta L S M_L M_S \rangle \quad (5.8)$$

This will reduce (5.7) to the equation

$$\langle \alpha' J' M' | Y^{\dagger} | \alpha JM \rangle = \delta_{SS'} \sum_{M_L M_L'} (L' S' J' M' | L' S' M_L' M_S') \langle \beta' L' S' M_L' M_S' | Y^{\dagger} | \beta L S M_L M_S \rangle (LSM_L M_S | LSJM) \quad (5.9)$$

All other matrix elements vanish—in particular we see that there are no matrix elements between terms of different multiplicities. Also we see that

(5.4) is again applicable, and obtain

$$\langle \beta' L' S' M_L' M_S' | Y^{\dagger} | \beta L S M_L M_S \rangle = (L' \nu L' M_L' | L' \nu M_L \mu) (\beta' L' S' | | Y^{\dagger} | | \beta L S) \quad (5.10)$$

Hence, we can write (5.9) in the form

$$\langle \beta' L' S' J' M' | Y^{\dagger} | \beta L S J M \rangle = \sum_{M_L M_L'} (L' S' J' M' | L' S' M_L' M_S') (L' \nu L' M_L' | L' \nu M_L \mu) (LSM_L M_S | LSJM) (\beta' L' S' | | Y^{\dagger} | | \beta L S) \quad (5.11)$$

and this expression is also (5.4). If we multiply both expressions by $(J' \nu J' M' | J' \nu M' \mu)$, introduce the requirement that $M_L' = M_L + \mu$, and sum over M , the expression on the right of (5.4) will reduce to $(\alpha' J' | | Y^{\dagger} | | \alpha J)$ by (5.5a) while the right side of (5.11) becomes

$$\sum_{M, M_L} (J' \nu J' M' | J' \nu M' \mu) (L' S' J' M' | L' S' (M_L + \mu) M_S) (L' \nu L' [M_L + \mu] | L' \nu M_L \mu) (LSM_L M_S | LSJM) (\beta' L' S' | | Y^{\dagger} | | \beta L S) \quad (5.12)$$

Such a sum of the product of 4 Wigner coefficients appears recurrently in the theory of complex spectra. The properties of such expressions were first studied extensively by Racah (1942a), and they have consequently become known as Racah coefficients. In the present application the essential significance of (5.11) is that the two states JM and $J'M'$ are “uncoupled” into $LSM_L M_S$ and $L'S'M_L' M_S'$ so that the effect of the Y^{\dagger} , acting only on the L part (i.e., the space coordinates) may be determined. The Racah coeffi-

cients appear generally in connection with such uncoupling or recoupling of angular momenta. The Racah coefficient in (5.12) depends on six parameters J, J', L, L', S , and ν ; in spite of the explicit appearance of the projection quantum numbers M, M', M_L, M_S , and μ , they are eliminated in the summation.

Although a general algebraic formula for these coefficients is quite unwieldy, numerical tables have been computed by Biedenharn (1952) and by Simon et al. (1954). The first of these carries the entries in a closed fractional form, the other has a ten-place decimal form. In order to clarify difference in notation, let us rewrite (5.12) as

$$(\alpha' J' || Y_\nu || \alpha J) = \sum_{M, M_L} C_{M, M_L}^{J, J'} C_{(M_L + \nu) M_S}^{L', S} C_{M_L \mu}^{J, L'} C_{M_L M_S}^{L, S} (\beta' L' S || Y_\nu || \beta L S). \quad (5.13)$$

Omitting algebraic details, the use of the symmetry relations for the Wigner coefficients will permit this to be transformed into

$$(-1)^{L+S-J'-\nu} \sqrt{\frac{(2J+1)(2L'+1)}{(2S+1)(2\nu+1)}} \sum_{M, M_L} C_{M, M_L}^{J, L, S} C_{(M+M_L)(M'-M-M_L)}^{S, L'} C_{M(M'-M)}^{J, L'} (\beta' L' S || Y_\nu || \beta L S)$$

and a direct comparison with the notation of this with eq (1), page x of Simon et al. will give

$$(\alpha' J' || Y_\nu || \alpha J) = (-1)^{L+S-J'-\nu} \sqrt{\frac{(2J+1)(2L'+1)}{(2S+1)(2\nu+1)}} W(JLJ'L'; S\nu) (\beta' L' S || Y_\nu || \beta L S). \quad (5.14)$$

The six-parameter expression W is tabulated in Simon et al. for $J, J' \leq 15/2$; $L, L' \leq 9/2$ (though of course a half-integral L has no significance in the present application) $S \leq 3$ and $\nu \leq 8$, though the useful range of values may be extended through the use of the symmetry relations given in the introduction to the tables.

We are now in a position to make comparisons between quantities which depend only on the ratios of the matrix elements for a given $\beta' L' S$, $\beta L S$ combination, but for additional details we must evaluate the reduced matrix element $(\beta' L' S || Y_\nu || \beta L S)$. In principle, this may be accomplished in a way similar to the derivation of (5.13) i.e., we now assume that the state $|LSM_L M_S\rangle$ may be written as a sum of antisymmetrized products of one-electron wave functions $|nlm_l m_s\rangle$ and the matrix element (5.10) can be expressed in terms of one-electron matrix elements such as

$$\langle n'l'm'_l m'_s | Y_\nu | nlm_l m_s \rangle.$$

Unfortunately, this procedure has several limi-

tations. First, there is the additional assumption that the state involves only one configuration of the free ion. In the present case, it is expected that configurations other than $4f^n$ will interact with the surroundings and other rare earth ions to such an extent that they will no longer be characterized by sharp energy levels, but by broad levels or bands. The sharpness of the observed lines suggests that the contribution of the other configurations to the wave functions of the free ion is, at least as far as the energy is concerned, negligible. It is quite possible that contributions from other configurations are responsible for features such as intensities and polarization; this will be discussed further in the following section.

An additional disadvantage is that, unlike the transformation from a JM representation to an LS representation, there is no general formula for passage from the $LSM_L M_S$ representation of a state to its one-electron representation. The general method of Gray and Wills (Condon and Shortley 5^s) may be used, but each configuration must be considered by itself, and no general formula can be derived.

5.4. The Rare Earth Ground Terms

In the evaluation of the matrix element (5.10) we are free to select any consistent set of the projection quantum numbers; we shall therefore consider the specific element $[M_L=L'; M_S=S]$

$$\langle \beta' L' S L' S | Y_\nu | \beta L S L S \rangle = (L_\nu L' L' | L_\nu L \mu) (\beta' L' S || Y_\nu || \beta L S). \quad (5.15)$$

By selecting the maximum values of the projection quantum numbers, we shall find that the transformation to the one-electron representations is thereby simplified considerably.

For any particular rare earth, the analysis is best carried forward in a systematic manner, beginning with the ground state. In accordance with Hund's rules, for $4f^n$ we select the highest allowed multiplicity, and then the highest L consistent with this multiplicity. The corresponding antisymmetrized products of one-electron wave functions are usually written $\left\{ \begin{smallmatrix} + & - & + \\ m_1 & m_1 & m_2 \end{smallmatrix} \dots \right\}$; for example, in this notation we start with $\left\{ \begin{smallmatrix} + & + & + \\ 3 & 2 & 1 \end{smallmatrix} \right\}$ for Nd, $\left\{ \begin{smallmatrix} + & + & + & + \\ 3 & 2 & 1 & -2 \end{smallmatrix} \right\}$ for Eu, $\left\{ \begin{smallmatrix} + & - & + & + \\ 3 & 3 & 2 & 1 \end{smallmatrix} \right\}$ for Tb, etc. While the analysis may be carried out for any configuration, we shall consider the specific example of Nd.

Let us begin with $\left\{ \begin{smallmatrix} + & + & + \\ 3 & 2 & 1 \end{smallmatrix} \right\} = |^4 I 6 \frac{3}{2}\rangle$. The application of the symmetric operator $\mathcal{L} = L_x - iL_y$ (Condon and Shortley 5^s3) yields $\left\{ \begin{smallmatrix} + & + & + \\ 3 & 2 & 0 \end{smallmatrix} \right\} = |^4 I 5 \frac{3}{2}\rangle$.

A second application of \mathcal{L} yields $2\sqrt{3} \left\{ \begin{smallmatrix} + & + \\ 3 & 2 \end{smallmatrix} \right\} + \sqrt{10} \left\{ \begin{smallmatrix} + & + & + \\ 3 & 1 & 0 \end{smallmatrix} \right\} = \sqrt{22} |^4 I 4 \frac{3}{2}\rangle$. The state $\frac{1}{\sqrt{22}} \left[\sqrt{10} \left\{ \begin{smallmatrix} + & + & + \\ 3 & 1 & 0 \end{smallmatrix} \right\} + \sqrt{3} \left\{ \begin{smallmatrix} + & + \\ 3 & 2 \end{smallmatrix} \right\} \right]$

$\left\{ \begin{smallmatrix} ++ \\ 32-1 \end{smallmatrix} \right\} - 2\sqrt{3} \left\{ \begin{smallmatrix} +++ \\ 310 \end{smallmatrix} \right\}$ is orthogonal to $|^4I 4 \frac{3}{2}\rangle$ and is also a quartet with $M_L=4$ and hence is the state $|^4G 4 \frac{3}{2}\rangle$. Repeated application of \mathcal{L} will yield the states $|^4F 3 \frac{3}{2}\rangle$, $|^4D 2 \frac{3}{2}\rangle$, and $|^4S 0 \frac{3}{2}\rangle$, all of which arise from $4f^3$ (Condon and Shortley able 1'). Similarly, beginning with $\left\{ \begin{smallmatrix} +++ \\ 332 \end{smallmatrix} \right\} = |^2L 8 \frac{1}{2}\rangle$ we can obtain the state $|^2K 7 \frac{1}{2}\rangle$. There are a total of four states with $M_L=6$, $M_S=\frac{1}{2}$, namely $|^2L 6 \frac{1}{2}\rangle$, $|^2K 6 \frac{1}{2}\rangle$, $|^2I 6 \frac{1}{2}\rangle$ and $|^4I 6 \frac{1}{2}\rangle$. The first two are obtained by a further application of \mathcal{L} while the latter is obtained from $|^4I 6 \frac{3}{2}\rangle$ by the operator $\mathcal{S}_- = S_x - iS_y$. The fourth state is then $|^2I 6 \frac{1}{2}\rangle$.

In endeavoring to continue this process to states where $M_L=5$, we find that in addition to those arising from higher values of L or S , that there are two additional states with $M_L=5$, $M_S=\frac{1}{2}$. They must be therefore both described as $|^2H 5 \frac{1}{2}\rangle$, and additional labels are necessary to distinguish them. The usual procedure is to explicitly diagonalize the submatrix of electrostatic interaction between the two states (Condon and Shortley 7⁸), but further work of Racah (1949) has shown that a more abstract classification of states based on group theory is possible. The former technique has been applied by Judd (1955) to the specific case of Eu.

In any perturbation calculation, the matrix elements first to be considered are those between different states of a degenerate level. In the rare earth ions, and in crystal spectra generally, the one most important level is the ground level. While the great significance of optical spectroscopy, in contrast to magnetic resonance measure-

ments, lies in data provided concerning excited levels, a general ion-by-ion analysis of the excited levels is a major undertaking, and only the explicit details for the ground multiplet will be developed here. As a specific example, we shall further consider the case of Nd, where (5.15) becomes

$$\begin{aligned} & \langle 4f^3 \ ^4I 6 \frac{3}{2} | Y_0^0 | 4f^3 \ ^4I 6 \frac{3}{2} \rangle \\ & = \langle \left\{ \begin{smallmatrix} +++ \\ 321 \end{smallmatrix} \right\} | Y_0^0 | \left\{ \begin{smallmatrix} +++ \\ 321 \end{smallmatrix} \right\} \rangle. \end{aligned} \quad (5.16)$$

We shall here insert again the factor r^r in the potential, omitted since (5.1) since it is invariant under the transformations we have been considering up to this point. Again recalling that $r^r Y_0^0$ is an electrostatic perturbation upon the individual electrons, we see that it is a spin-free quantity of Condon and Shortley type F (6⁶⁹), and (5.16)

becomes $\sum_{m=1}^3 \langle 4f^+ m | r^r \Theta_0^0 | 4f^+ m \rangle$, where $\Theta_0^0 = \frac{1}{\sqrt{2\pi}}$

and Θ_0^0 is given in table 1. The spin terms of the matrix element give +1, the ϕ terms give merely $\frac{1}{\sqrt{2\pi}}$ and the θ integral is

$$\int_0^\pi \Theta_0^0 \Theta_0^0 \Theta_0^0 \sin \theta \, d\theta = \sqrt{\frac{2\nu+1}{2}} \, c^r(3m, 3m)$$

where c^r is given in table 1⁶ of Condon and Shortley. For the time being, we shall write the r^r integral as $\langle r^r \rangle$.

By evaluating (5.14) and (5.15) for all the rare earths, the reduced matrix elements for the ground levels apart from the term $\langle r^r \rangle$ have been computed and presented in table 16. The reduced L - S matrix elements are included principally for information, though they are directly useful in cases where the electrostatic perturbation is greater than the spin-orbit interaction, or where the electrostatic and spin-orbit perturbations must be considered simultaneously.

6. Applications

6.1. General Considerations

In this concluding section, we shall discuss further the general significance of the preceding sections in the interpretation of crystal spectra, the connection mentioned at the beginning of section 3.4 between the group-theory technique of sections 3 and 4, and the matrix element calculations of section 5, and shall illustrate the application of these general techniques to specific problems.

Generally, a spectrum may be considered as understood if the spacing of the energy levels is known, the effect upon them of a magnetic field,

selection and polarization rules and relative intensities of lines, and all these are correlated with the structure of the atom responsible for the spectrum. These may all in principle be obtained from a knowledge of the initial and final states of the transition involved in the spectral line. Unfortunately, our knowledge of these initial and final states is largely a matter of approximation which is not always entirely satisfactory. This is true both for the free ions and for the perturbed ions in the crystal. Let us review briefly the situation for the free ion.

The good quantum numbers for the wave functions of the free ion (section 3.3) are the parity

TABLE 16. *Reduced matrix elements*

Element	<i>L</i>	<i>S</i>	<i>J</i>	$\nu=2$	$\nu=4$	$\nu=6$	
Ce ⁺⁺⁺	3	1/2		-0.325732	0.360856	-0.491065	
Yb ⁺⁺⁺							
			5/2	-.301569	.261168	0	
			7/2	-.307788	.289329	-.245532	
Pr ⁺⁺⁺	5	1		-.276822	-.261529	.455018	
Tm ⁺⁺⁺							
				6	-.265205	-.224931	.321183
				5	-.249840	-.174353	.136505
			4	-.260040	-.208628	.261268	
Nd ⁺⁺⁺	6	3/2		-.106082	-.168698	-.642369	
Er ⁺⁺⁺							
				9/2	-.098820	-.131239	-.356506
				11/2	-.091864	-.096145	-.099840
				13/2	-.093887	-.106106	-.169019
			15/2	-.101524	-.144946	-.457704	
Pm ⁺⁺⁺	6	2		.106082	.168698	.642369	
Ho ⁺⁺⁺							
				4	.095469	.115742	.260356
				5	.084208	.054866	-.090686
				6	.083901	.061345	-.087596
				7	.089965	.088311	.062990
				8	.100398	.139515	.420886
Sm ⁺⁺⁺			5	5/2		.276822	.261529
Dy ⁺⁺⁺							
		5/2			.217802	.102885	0
		7/2			.160051	-.033771	.252972
		9/2			.155288	-.038079	.220158
		11/2			.174559	-.007942	.210017
		13/2			.208638	.070737	.112678
		15/2			.253811	.193261	-.228851
Eu ⁺⁺⁺	3	3				.325732	-.360856
Tb ⁺⁺⁺							
				0	0	0	0
				1	-.199469	0	0
				2	-.012363	-.150786	0
				3	-.054289	.060143	-.081844
				4	.032147	.171347	.259348
				5	.138409	.130764	-.227510
				6	.265202	-.168697	.064237
Gd ⁺⁺⁺			0	7/2		0	0

Note 1: Where no *J* value is given, the tabulated quantity is $(\beta LS||Y_{\nu}||\beta LS)$, otherwise $(\beta LSJ||Y_{\nu}||\beta LSJ)$.
 Note 2: For the second ion of the above pairs, all signs change.

"*g*" or "*u*", the total angular momentum *J* and its component *M*. The strict selection rules $\Delta J=0, \pm 1$ ($0 \rightarrow 0$) for electric and magnetic dipole radiation, and $\Delta J=0, \pm 1, \pm 2$ ($0 \rightarrow 0, 0 \rightarrow 1$) for electric quadrupole radiation, with change in parity in the first case, no change in the last two, were calculated in section 4.4. There is no physical significance to a selection rule on *M* in the absence of a nonspherical perturbation. If the spin-orbit interaction is small, *L* and *S* approximately describe a state and we obtain selection rules in *L* similar to those in *J*, while in *S* we obtain $\Delta S=0$ for electric dipole or quadrupole transitions, $\Delta S=0, \pm 1$ ($0 \rightarrow 0$) for magnetic dipole

transitions. Thus, consider a transition probability for electric dipole radiation involving the matrix element $\langle \beta' L' S J' M' | e r | \beta L S J M \rangle$. If $J'=J+1$ so the transition is allowed in *J*, but $L'=L+2$, it will be forbidden in *L*, and will be nonzero only if the spin-orbit interaction has mixed terms with spatial angular momentum *L*+1 in either *L* or *L'* or both. The application of a magnetic field splits a state into its 2*J*+1 components with a splitting determined to a first approximation by the matrix element of $-\frac{e}{2mc} (\mathbf{L}+2\mathbf{S}) \cdot \mathbf{B}$, yielding the usual expression for the Landé *g*-factor. Selection rules for *M* now become sig-

nificant, and we have $\Delta M=0, \pm 1$ for either type of dipole radiation, $\Delta M=0, \pm 1, \pm 2$ for quadrupole transitions, and their associated polarization effects.

The perturbing potential V (2.6) will influence a state $|\alpha JM\rangle$ in general by four types of matrix elements, the effects of which we shall examine in detail. These are:

1. Matrix elements diagonal in α, J, M .
2. Matrix elements diagonal in α, J .
3. Matrix elements diagonal in α .
4. Matrix elements nondiagonal in α , though perhaps diagonal in J and M . These matrices (or rather, approximations to them) may be calculated by the methods of section 5, though the scope of table 15 itself will suffice only for elements of types (1), (2), and those of (4) diagonal in parity and J . We will now examine the effect on the original level of these several matrix elements.

Matrix elements of type (1) arise from potential terms with ν even, $\mu=0$. Their effect is to split the level into pairs of states $\pm M$ (and the single level $M=0$ for integral J) in the ratios indicated by table 15. The inclusion of these matrix elements only results in an initial axial field approximation which is sometimes useful. The selection rules on J and parity, as well as any approximate rules on L and S , remain valid, as do those on M corresponding to the free ion in a magnetic field. This case corresponds closely to a reduction in symmetry from D_2 to D_{2h} , which differs through the inclusion of matrix elements of types (3) and (4) which are diagonal in M .

Elements of type (2) arise from potential terms with ν even but $\mu \neq 0$, still with the ratios determined by table 15 and of the same order of magnitude as those of type (1). The solution of the secular determinant for elements of (1) and (2) together will yield the first approximation to the energies of the perturbed states as well as the zero-order linear combination of states with various M required for higher order perturbation calculations. States with M values differing by μ will interact and M will no longer be a good quantum number for these states. However, if $2J$ is less than the lowest μ involved, there will be no matrix elements of this type and the rules for (1) only will still be valid. This situation will occur for levels of low J in fields of high symmetry. Generally, the selection rules for J and α will still hold. Selection rules on M will apply to those states which do not involve a combination of M values, and can be applied to the components of those states which do involve a mixing of two or more M states.

Type (3) matrix elements still involve only potential terms with ν even, since α includes the parity quantum number. The addition of these matrix elements to our scheme destroys the validity of J as a quantum number which has held for (1) and (2). These elements are those between various J values of a given $L-S$ term,

since they are still diagonal in α . In order of magnitude, these elements are as large as those of types (1) and (2), since all three come from the same reduced $L-S$ matrix element (table 16) and differ only in the associated Racah and Wigner coefficients (5.14, 5.4). Their effect is numerically somewhat smaller, since terms they contribute in perturbation theory are divided by the energy differences between the unperturbed levels. A state of given $J=J_0$ will be mixed directly with other J values such that $|J-J_0| \leq \nu$. The selection rules for the free ion J are now either poor or completely worthless, depending on the numerical parameters. Since both the upper and lower states of a transition can involve a range of J , the effective selection rule is merely $\Delta J \leq 2\nu + 1$.

Matrix elements of type (4) will generally be numerically the smallest of the four types. They arise from all terms in the potential (2.6), this being the only one of the four where the terms with odd ν are included. Such elements, at least as far as the ground state is concerned, involve relatively distant levels, so the influence on the splitting of the ground state will be small and lie perhaps more in the overall position of the level with respect to the unperturbed level. Perhaps the most important effect of such elements is on the selection rules and the associated polarization effects, and in the intensity calculations. Just as elements of type (3) destroyed J as a quantum number, so will these destroy α as a quantum number. α includes the specification of the configuration and consequently the parity of a state. In the most general case, there will be matrix elements for even ν between $4f^n$ and higher even configurations, such as $4f^{n-2}5d^2$, and for odd ν between $4f^n$ and $4f^{n-1}5d$. These latter elements destroy the parity of the state and lift the restrictions of Laporte's rule for electric dipole transitions. Intensities corresponding to electric dipole transitions between states principally $4f^n$ in character must, therefore, be computed from these extremely small inter-configurational matrix elements and are therefore extremely weak; nevertheless, all crystal spectra are in any absolute sense quite weak and such transitions must be considered in the case of any nonholohedric perturbing potential.

The further effect of a uniform magnetic field will depend to a large extent on the degeneracy allowed by the electrostatic perturbation. The magnetic field does not allow any remaining degeneracy; hence, levels degenerate in the electrostatic field, including the Kramers conjugate states, will be split by a term linear in the magnitude of the field B which can be computed from a solution of the associated secular determinant. The average magnetic moment of any nondegenerate state is zero (Van Vleck, 1932; Klein, 1952) so these states will show no first order effect for weak magnetic fields. If the crystal field leaves two or more levels relatively close together, a case of intermediate or strong magnetic field approximation can be observed and the level will show a

finite displacement from its $B=0$ position. Unless the magnetic field and crystal field symmetry axes are parallel, there will generally be no symmetry and no selection rules. Marked variation of intensities of lines has been observed in the crystal Zeeman effect. A more specific example of some of these considerations will be given in section 6.2.

6.2. A Crystal Field

The direct approach to a crystal problem is to obtain the crystal structure by diffraction measurements, geometric considerations, or other crystallographic techniques. These techniques do not locate the lighter ions with sufficient accuracy, particularly water of crystallization, which contribute greatly to the crystal field. As an illustration of the procedure, let us consider a possible short range structure for rare-earth chlorides of the form $XCl_3 \cdot 6H_2O$ (H. S. Robertson, informal communication) where 6 Cl^- (three from adjoining X) are in two horizontal planes equidistant from X , forming an equilateral triangle in each plane, the two triangles being rotated 60° to each other about a vertical axis. The water molecules [considered as dipoles] are then located along the $X-Cl$ lines.

While detailed calculations were not carried out due to uncertainties in the various distance parameters, certain features of considerable interest may be obtained through partial application of (2.7). The above charge array may be described by

$$\begin{aligned} \alpha_{1-9} &= \theta & \alpha_{10-18} &= \pi - \theta & \beta_{1-3} &= 0 \\ \beta_{4-6} &= \frac{2\pi}{3} & \beta_{7-9} &= \frac{4\pi}{3} & \beta_{10-12} &= \frac{\pi}{3} & \beta_{13-15} &= \pi \\ \beta_{16-18} &= \frac{5\pi}{3} & q_{3n} &= q_{3n+1} = -e & q_{3n+2} &= e \end{aligned}$$

and we shall take θ as a variable parameter. There are a total of 18 charges in this array. Those above the $x-y$ plane have the indices 1-9, while those below have 10-18. Indices of the form $3n$ refer to Cl^- , $3n+1$ refers to OH^- and $3n+2$ refers

to H^+ . The angular portions of the potential were calculated over a range from $\theta=0^\circ$ to 90° . The angular factors (2.4) were obtained from the tables of Tallqvist (1908). This table is not widely available, but no other table covering its range has been prepared to the knowledge of the writer. The National Bureau of Standards table (1945) covers values of M only up to and including 4.

In performing the summation (2.7), the only generally nonvanishing coefficients are ($N \leq 6$) C_2^0 , C_4^0 , C_6^0 , C_4^2 , C_6^2 , and C_6^4 , though this may be deduced directly from table 5 without explicitly performing the summation. The values of these, apart from a radial factor of the form

$$e \left(\frac{-1}{R_1^{N+1}} + \frac{-1}{R_2^{N+1}} + \frac{1}{R_3^{N+1}} \right)$$

are given in table 17. Certain general qualitative features may be noted by inspection. For $\theta=0$ (corresponding to all charges being located along the vertical axis), the symmetry is $D_{\infty h}$ (table 5). For other angles, the symmetry is generally D_{3d} , while at 90° (a planar configuration) it is D_{6h} . A particular exception occurs for $\theta=54^\circ 44'$ where the symmetry is "accidentally" O_h , belonging to the cubic point groups (section 2.4). For this value of θ , the three pairs of charge axes are mutually perpendicular and the C_2^0 term vanishes. For other values of θ (e.g., 31°), other coefficients vanish, but there is no general theoretical significance to this. It is to be noted that for $\theta < 10^\circ$, the terms responsible for departure from $D_{\infty h}$ symmetry are small, while in the vicinity of $\theta=90^\circ$ the terms causing departure from D_{6h} symmetry are large and quite sensitive to small angular variations.

Most features of even this simple problem require the specification of the radial distances of the ions, as well as the average radial integrals ($\langle r^v \rangle$) of the wave functions. On the other hand, aspects of the Zeeman effect can be understood without further knowledge of these parameters, and these aspects are most important for understanding more complicated cases. Let us assume

TABLE 17. A representative crystal field

θ	C_2^0	C_4^0	C_6^0	C_4^2	C_6^2	C_6^4
0°	9.5120	7.0898	5.8991	0	0	0
10°	9.0818	6.0491	4.1557	0.0765	0.2113	0.0001
20°	7.8430	3.3675	0.4242	.5575	1.3485	.0063
30°	5.9450	0.1662	-2.2064	1.6053	3.0365	.0619
40°	3.6168	-2.2617	-1.9088	3.0170	3.7557	.2795
45°	2.3780	-2.8802	-0.8756	3.7074	3.3393	.4953
50°	1.1392	-3.0311	.3326	4.2850	2.3851	.8007
$54^\circ 44'$	0	-2.7571	1.3109	4.6604	0.7915	1.1740
60°	-1.1890	-2.0494	1.9068	4.8160	-4.338	1.6716
70°	-3.0870	-0.0269	1.2322	4.2086	-2.5978	2.7282
80°	-4.3258	1.8852	-0.7794	2.4595	-2.3645	3.6146
90°	-4.7560	2.6587	-1.8435	0	0	3.9624

* See text.

that we have the case $\theta=0^\circ$ of the preceding table 17, i.e., $D_{\infty h}$ symmetry. We see from table 9 that the $2J+1$ states are separated into doubly degenerate pairs corresponding to $\pm M$, plus (for integral J only) a nondegenerate state $M=0$.

The application of a uniform magnetic field parallel to the z -axis will reduce the symmetry to $C_{\infty v}$ and all levels are now nondegenerate. The splitting is given by twice the diagonal matrix element $\langle M|L_z+2S_z|M\rangle B$, since the $-M$ level is depressed by this same amount and there are no nondiagonal elements, L_z+2S_z , having no dependence on ϕ . If the field is applied parallel to the y axis, the only symmetry remaining is C_{2v} , but this is referred to a horizontal axis, rather than a vertical one. The secular determinant will now involve matrix elements of L_y+2S_y . This has a ϕ dependence of the form $e^{\pm i\phi}$, and will have nonvanishing matrix elements between states differing in M by 1. Thus, of all the $\pm M$ pairs, a transverse magnetic field will have no diagonal elements and will have nondiagonal elements only in the case of $\pm \frac{1}{2}$. The magnitude of this element will be $\frac{1}{2}\sqrt{(J+\frac{1}{2})(J-\frac{1}{2}+1)}g\beta B$ and the splitting will be twice that or $(J+\frac{1}{2})g\beta B$.

If we now proceed to the planar case D_{6h} , we see from table 10 that for $J \leq \frac{5}{2}$ the states remain as in the $D_{\infty h}$ case and the Zeeman effect for these lower J values is unchanged. For $J=3$, the original pair $M=\pm 3(E_{3g,3u})$ now splits into two nondegenerate states $B_{1g,1u}$ and $B_{2g,2u}$, which are the states $\frac{1}{\sqrt{2}}|+3\rangle + \frac{1}{\sqrt{2}}|-3\rangle$ and $\frac{1}{\sqrt{2}}|+3\rangle - \frac{1}{\sqrt{2}}|-3\rangle$, respectively. This arises from the $e^{\pm 6i\phi}$ in the potential with nonzero matrix elements between terms where $\Delta M = \pm 6$. These states will show no first-order Zeeman effect, either longitudinal or transverse. For $J=4$, we see that the states coming from ± 2 and ± 4 have the same symmetry ($E_{2g,2u}$), and while we still have degenerate pairs, their properties will depend upon the solution of the associated secular determinant. One pair will be given by an expression of the form $a|\pm 4\rangle + (1-a^2)^{1/2}|\mp 2\rangle$ while the other will be of the form $(1-a^2)^{1/2}|\pm 4\rangle - a|\mp 2\rangle$. The coefficient a will be fixed by the magnitude of C_6^g . For these terms, the longitudinal Zeeman splitting will be $2[4a^2 - 2(1-a^2)] = 12a^2 - 4$ and $2[4(1-a^2) - 2a^2] = 12a^2 - 8$, respectively, while the transverse effect remains zero. This longitudinal effect thus depends explicitly on the crystal field. For $J=\frac{7}{2}$, a similar phenomenon takes place through a combination of $|\pm \frac{7}{2}\rangle$ with $|\mp \frac{5}{2}\rangle$, and the longitudinal effect will again depend explicitly on the crystal field. A major difference will occur in the transverse effect—here there are nonvanishing off-diagonal matrix elements of the form $\langle \frac{7}{2} L_y + 2S_y | \frac{5}{2} \rangle$, in the secular determinant

and there will be a transverse Zeeman splitting, also depending explicitly on the crystal field. Under still lower symmetries than D_{6h} , the Zeeman effect in odd-electron systems becomes increasingly dependent on the crystal field, since Kramers degeneracy remains in all cases, or vanishes in first order in most even-electron systems where the crystal field states generally become nondegenerate.

6.3. The Fluorescence Spectrum of the Terbium Ion

Because of the inexact knowledge of the radial parameters necessary for further work on the direct approach to a solution of a given problem, the more common procedure is to utilize the quantities $\langle r^N \rangle C_N^M$ and $\langle r^N \rangle S_N^M$ as adjustable parameters to fit as best as possible the observed data. This leaves the radial wave function integrals combined with the crystal field parameters—they are frequently left in this form since further information about either one of them separately is not readily available. We shall illustrate this procedure by some considerations of the fluorescence spectrum of $TbCl_3 \cdot 6H_2O$.

The experimentally observed fluorescence lines (Singh, J. H. U., Dissertation, 1957), observed both at $4.2^\circ K$ and $2.2^\circ K$, are given in table 18. The exciting light, unpolarized, is incident along the crystal axis (the z axis). For both temperatures, the spacing of each level from the lowest of its group is given, the average of the two temperature values, and the spacing of this (to the nearest cm^{-1}) from the mean of the group. The overall mean for each group is also given. The fluorescent light is observed on the side of the crystal opposite the incident radiation, also along the z -axis. The fluorescent lines are relatively weak and are generally observed without regard to polarization; however, a few plates were taken at $4.2^\circ K$ showing polarization (electric vector along x or y). These data were made available by private communication from Dr. Singh and are also included in table 18. In the interpretation of fluorescence spectra it should be remembered that the spectra may originate near the surface of the crystal, and that local fields, different from those in the main body of the crystal, may significantly affect the regularities expected.

The Tb ion has the configuration $4f^8$. According to Hund's rule, the ground state is a 7F multiplet with $J=6$ lowest, with J from 5 to 0 above the lowest. The fluorescence spectrum is caused by transitions from a higher level (probably 5D but this is not important for the present considerations) to various levels of the ground multiplet. Qualitatively, the agreement with experiment is excellent. The groups of lines Z, Y, X, W, V, U, T , correspond to $J=6, 5, 4, 3, 2, 1$, and 0, in that order. Within each group, there are $2J+1$ lines, indicating that the crystal field has removed all degeneracy. This is confirmed by the absence of any first order Zeeman effect.

TABLE 18. Fluorescence spectrum of $TbCl_3 \cdot 6H_2O$

Line	(4.2°K)		(2.2°K)		$\bar{\Delta\nu}$	$\overline{\Delta\nu - \Delta\nu_m}$	x y	
	ν	$\Delta\nu$	ν	$\Delta\nu$				
Z ₁	20538.81	0.00	20538.77	0.00	0.00	-167		
Z ₂	20503.66	35.15	503.72	35.05	35.10	-132		
Z ₃	498.20	40.61	498.33	40.44	40.52	-127	No	
Z ₄	461.26	77.55	461.59	77.18	77.36	-90	data	
Z ₅			454.97	83.80	83.80	-83		
Z ₆	380.70	158.11	379.70	159.07	158.59	-8		$\bar{Z} = 20372$
Z ₇	369.16	169.65	370.19	168.58	169.12	2		
Z ₈	359.49	179.32	359.27	179.50	179.41	12		
Z ₉			347.85	190.92	190.92	24		
Z ₁₀	306.14	232.67	305.73	233.04	232.86	66		
Z ₁₁	239.08	299.73	238.33	300.44	300.08	133		
Z ₁₂	193.84	344.95	192.86	345.91	345.43	178		
Z ₁₃	179.61	359.20	179.33	359.44	359.32	192		
Y ₁	18433.00	0.00	18433.04	0.00	0.00	-127	s w	
Y ₂	413.02	19.98	413.02	20.00	19.99	-107	s w	
Y ₃	400.92	32.08	400.99	32.05	32.06	-95	w s	
Y ₄	359.26	73.74	359.41	73.63	73.68	-53	s w	
Y ₅	341.00	92.00	341.09	91.95	91.98	-35	s w	$\bar{Y} = 18306$
Y ₆			333.69	99.35	99.35	-27	s w	
Y ₇	308.29	124.71	308.57	124.47	124.59	-2	s w	
Y ₈	294.96	138.04	295.90	137.14	137.59	11	s w	
Y ₉	250.87	182.13			182.13	55	s w	
Y ₁₀	124.83	308.17	124.82	308.22	308.20	181	? ?	
Y ₁₁	107.90	325.10	107.81	325.23	325.16	198	? ?	
X ₁	17193.56	0.00	17193.49	0.00	0.00	-74	w w	
X ₂			188.20	5.29	5.29	-69	w w	
X ₃	184.65	8.91	184.76	8.73	8.82	-65	s	
X ₄			135.74	57.75	57.75	-16	s w	$\bar{X} = 17120$
X ₅	131.91	61.65	131.98	61.51	61.58	-12	s w	
X ₆			104.88	88.61	88.61	15	vw	
X ₇	067.31	126.25	067.52	125.97	126.11	52	"	
X ₈			056.26	137.23	137.23	63	"	
X ₉	013.94	179.62	014.48	178.81	179.22	105	"	
W ₁	16162.72	0.00	16162.85	0.00	0.00	-98	w w	
W ₂			080.68	82.17	82.17	-15	s	
W ₃	070.26	92.46	070.33	92.52	92.49	-5	w w	
W ₄	043.44	119.28	043.44	119.41	119.34	22	w s	$\bar{W} = 16065$
W ₅	037.91	124.81	037.95	124.90	124.86	27	w w	
W ₆	032.99	129.73	032.87	129.98	129.86	32	s w	
W ₇	028.07	134.65	028.12	134.69	134.69	37	s w	
V ₁	15499.89	0.00				-145		
V ₂	430.03	69.86				-75		
V ₃	297.53	202.36				57		$\bar{V} = 15355$
V ₄	292.94	206.95				62		
V ₅	253.19	246.70	15253.17			101	vw s	
U ₁	15001.25	0.00				-23	s vw	
U ₂	14994.68	6.57				-17	vw s	$\bar{U} = 14978$
U ₃	938.41	62.84				40		
T	14691.73							$\bar{T} = 14692$

Let us consider the quantitative details. For best results, our theory requires $L-S$ coupling and a crystal field splitting small compared to the multiplet separation. We may check the validity of $L-S$ coupling by the interval rule, taking the separation of the mean of adjacent groups and dividing by the higher J value. We obtain, in cm^{-1} , the following:

$J=6$	344	$J=4$	264	$J=2$	158.5
$J=5$	237	$J=3$	237	$J=1$	286

This is not especially impressive for very good obedience to $L-S$ coupling requirements. Let us also calculate the total spread of a group, divided by the separation of the group mean from that for the next lower J . We obtain:

$J=6$	0.174	$J=4$	0.170	$J=2$	0.653
$J=5$.274	$J=3$.190	$J=1$.220

This indicates that the influence of adjacent levels may be rather significant. The irregularity here for $J=2$ is quite interesting; it is only partially due to the irregularity shown in the interval rule check, since the total spread of $J=2$ is 247 cm^{-1} while that of $J=3$ is only 179 cm^{-1} .

In beginning an analysis, we must first obtain some idea of the symmetry. From table 9, we see that the only symmetries giving no degeneracy for an even-electron system are D_{2h} , C_{2v} , C_{2h} , and C_s . The overall crystal symmetry suggests either C_{2v} or C_{2h} . From table 5, we see that D_{2h} requires the following potential coefficients: $C_2^0 C_4^0 C_6^0 C_2^2 C_4^2 C_6^2 C_4^4 C_6^4 C_6^6$. The lower symmetry C_{2v} is obtained by the inclusion of odd potential terms, which will influence the energy splittings by a very small amount (section 6.1), while C_{2h} will include even S terms. These latter will strongly influence the energy splittings. In order to reduce the number of independent parameters, we shall assume D_{2h} , which will be energetically equivalent to C_{2v} .

In deducing the field parameters, we need to know not only the energies of the levels but also the free ion M values from which they come. Where the crystal field does not remove all degeneracy, this may be partially obtained from the Zeeman splittings. For the present case, the only source of such information is the polarization data.

We begin our detailed analysis with the simplest nontrivial case $J=1$. We have three levels at -23 , -17 , and $+40$. The only potential terms of D_{2h} contributing to matrix elements within $J=1$ are C_2^0 and C_2^2 . It is a natural first guess to consider that the C_2^0 term causes a splitting of the state $|0\rangle$ to $+40$, and $|+1\rangle$ and $|-1\rangle$ to -20 , with a small splitting of the latter two to -17 and -23 by C_2^2 . This is not necessarily the case, and must be in agreement with the polarization data.

From table 14, where radiation in the z direction corresponds to $\theta=\phi=0^\circ$, $\psi=0^\circ$ for x polarization and $\psi=90^\circ$ for y polarization, we see that x -polarized radiation may be produced by either A_x , B_y , or A_{xz} , while y -polarized radiation may be produced by A_y , B_x , or A_{yz} . These are the electric dipole, magnetic dipole, or electric quadrupole terms, respectively. We shall assume that the transitions are magnetic dipole in character. We do not know the symmetry of the upper state in the fluorescent spectrum, but there are only 4 possibilities in a D_{2h} symmetry. From table 13 we deduce the allowed lower state for each possible upper state as follows:

Upper state	$B_y(U_1)$	$B_x(U_2)$
A_{1g}	B_{2g}	B_{1g}
A_{2g}	B_{1g}	B_{2g}
B_{1g}	A_{2g}	A_{1g}
B_{2g}	A_{1g}	A_{2g}

The identification of U_1 and U_2 of table 18 ideally requires that they be present in only one of the two polarizations, rather than "very weak," but the actual behavior is quite close to this ideal. We now turn to table 9, and find that the lower state 1_g yields in D_{2h} symmetry the states A_{2g} , B_{1g} , and B_{2g} . Since A_{1g} is not present in the lower group, we conclude that the upper state is either A_{1g} or A_{2g} , that the lower states at -23 and -17 are B_{1g} and B_{2g} , while the other state from $J=1$ is A_{2g} . We cannot uniquely assign B_{1g} and B_{2g} to a particular one of the -23 , -17 levels. The upper level is presumably 5D_4 and a 4g level under D_{2h} gives both A_{1g} and A_{2g} levels. Interestingly enough, the same results, without "g" and "u" subscripts, hold for C_{2v} symmetry. We now see from table 10 that B_{1g} and B_{2g} arise from the state E_{1g} of $D_{\infty h}$, i.e., they come from $M=\pm 1$ and our earlier guess is confirmed.

In order to achieve maximum simplicity in the calculations, it is in general desirable to use zero-order wave functions which belong to reps of the symmetry group. In the present case, the state $|0\rangle$ belongs to A_{2g} , but the states $|+1\rangle$ and $|-1\rangle$ are not appropriate, and we must use the (essentially real) combinations $B_{1g}: \frac{1}{\sqrt{2}}|-1\rangle + \frac{1}{\sqrt{2}}|+1\rangle$ and $B_{2g}: \frac{1}{\sqrt{2}}|-1\rangle - \frac{1}{\sqrt{2}}|+1\rangle$. We shall frequently abbreviate these as $|1+\rangle$ and $|1-\rangle$, respectively. We now introduce the perturbation $V=r^2(C_2^0 C_4^0 + C_2^2 C_4^2)$. Since the potential V is symmetric under D_{2h} and our three zero order wave functions belong to different reps, all non-diagonal matrix elements vanish, and the energies are merely $\langle 0|V|0\rangle$, $\langle 1+|V|1+\rangle$ and $\langle 1-|V|1-\rangle$. It is a general characteristic of

crystal field problems that one must be freely able to shift from complex to real forms of wave functions and vice versa, as the symmetry demands.

In the present case, we have from (2.9) $Y_2^0 = C_2^0$ and $Y_2^2 = Y_2^{-2} = \frac{1}{\sqrt{2}} C_2^2$. Hence $\langle 0|V|0\rangle =$

$(\alpha_1 || Y_2 || \alpha_1) C_{10}^{242} Y_2^0 \langle r^2 \rangle = 40 \text{ cm}^{-1}$ or, from tables 15 and 16, $(0.199469)(-0.632456) Y_2^0 \langle r^2 \rangle = 40 \text{ cm}^{-1}$, so that $Y_2^0 \langle r^2 \rangle = -317.1 \text{ cm}^{-1}$. Similarly,

$$\langle 1+|V|1+\rangle = (\alpha_1 || Y_2 || \alpha_1) \left[\frac{1}{2} (C_{10}^{242} + C_{-10}^{242}) Y_2^0 + \frac{1}{2} (C_{-12}^{242} + C_{12}^{242}) Y_2^2 \right] \langle r^2 \rangle = -17 \text{ cm}^{-1}.$$

Recalling (5.6) and using the value of Y_2^0 just computed above, we obtain $(0.199469)(0.774597) Y_2^2 \langle r^2 \rangle = +3$, or $Y_2^2 = 19.4 \text{ cm}^{-1}$. The sign of Y_2^2 is ambiguous, since the state $|1+\rangle$ could equally well have been assigned to -23 cm^{-1} . These results, while interesting, are really not too profound, since we have only two independent energies relative to the mean of the group, and we had two adjustable coefficients at our disposal.

When we proceed to the $J=2$ case, we see that the allowed levels (table 9) are (2) A_{1g}, A_{2g}, B_{1g} , and B_{2g} . In contrast to the $J=1$ case, the B_{1g} and B_{2g} are $|1-\rangle$ and $|1+\rangle$. By using table 10, we see that A_{1g} and A_{2g} arise from $|\pm 2\rangle$. In particular A_{1g} is $|2+\rangle$ while A_{2g} is $|2-\rangle$. These may most conveniently be deduced from the transformation rules of table 8, in comparison with the tesseral harmonics of table 2, keeping in mind the phase convention for odd positive M of (2.8). The only nondiagonal matrix elements in the resulting secular determinant will be $\langle 0|V|2+\rangle$, so that we have only one second degree equation to solve for the energies.

The polarization data are available only for the state at $+101 \text{ cm}^{-1}$. This has polarization like the state B_{1g} or B_{2g} of $J=1$, and hence is presumably also $|1+\rangle$ or $|1-\rangle$. We do not have any polarization data on the other levels, so we do not know which states are which. We are therefore forced to a trial-and-error process. The additional potential parameters significant for $J=2$ are C_4^0, C_4^2, C_4^4 . If we select $|1+\rangle$ as 101 cm^{-1} and any one other as $|1-\rangle$, this will fix C_4^0 and C_4^2 , so that only C_4^4 is left to fit the other three levels.

In attempting this process, the work is somewhat simplified by observing that $(\alpha_2 || Y_2 || \alpha_2)$ is very small, and that second degree terms will have only a small contribution to the matrix elements. Consequently, they were disregarded for the trial and error work, and were included later for a more exact calculation. The required matrix elements are, omitting the factor $(\alpha_2 || Y_4 || \alpha_2)$ common to all,

$$\langle 0|V|0\rangle = C_{00}^{242} Y_4^0 \langle r^4 \rangle$$

$$\begin{aligned} \langle 0|V|2+\rangle &= \frac{1}{\sqrt{2}} (C_{-2}^{242} + C_{-22}^{242}) Y_4^2 \langle r^4 \rangle \\ &= \frac{2}{\sqrt{2}} C_{-22}^{242} Y_4^2 \langle r^4 \rangle \end{aligned}$$

$$\begin{aligned} \langle 2+|V|2+\rangle &= \frac{1}{2} (C_{20}^{242} + C_{-20}^{242}) Y_4^0 \langle r^4 \rangle \\ &+ \frac{1}{2} (C_{-24}^{242} + C_{24}^{242}) Y_4^4 \langle r^4 \rangle \\ &= C_{20}^{242} Y_4^0 \langle r^4 \rangle + C_{-24}^{242} Y_4^4 \langle r^4 \rangle \end{aligned}$$

$$\langle 1+|V|1+\rangle = C_{10}^{242} Y_4^0 \langle r^4 \rangle + C_{-12}^{242} Y_4^2 \langle r^4 \rangle$$

$$\langle 1-|V|1-\rangle = C_{10}^{242} Y_4^0 \langle r^4 \rangle - C_{-12}^{242} Y_4^2 \langle r^4 \rangle$$

$$\langle 2-|V|2-\rangle = C_{20}^{242} Y_4^0 \langle r^4 \rangle - C_{-24}^{242} Y_4^4 \langle r^4 \rangle$$

These may be simplified by observing that $C_{00}^{242} : C_{10}^{242} : C_{20}^{242} = 6 : -4 : 1$ and that $C_{-22}^{242} = -\sqrt{3/8} C_{-24}^{242}$. If we introduce the following abbreviations,

$$\langle 0|V|0\rangle = V_0 \quad \langle -2|V|2+\rangle = \beta$$

$$\langle -1|V|1-\rangle = \delta$$

the matrix elements may be simplified to:

$$\langle 0|V|0\rangle = V_0$$

$$\langle 2+|V|2+\rangle = \frac{1}{6} V_0 + \beta$$

$$\langle 2-|V|2-\rangle = \frac{1}{6} V_0 - \beta$$

$$\langle 1+|V|1+\rangle = -\frac{2}{3} V_0 + \delta$$

$$\langle 0|V|2+\rangle = -\sqrt{\frac{3}{8}} (\sqrt{2}) \delta = -\frac{1}{2} \sqrt{3} \delta$$

$$\langle 1-|V|1-\rangle = -\frac{2}{3} V_0 - \delta$$

The form of these matrix elements suggests the physical situation. The C_4^0 term splits the $2J+1$ levels into one at V_0 , a pair at $-\frac{2}{3}V_0$ and a pair at $\frac{1}{6}V_0$. The first pair is then further split by

C_4^2 into states separated by 2δ . The second pair is also split into two separated by 2β through C_4^4 , but one of these interacts further (via C_4^2) with $|0\rangle$ through the nondiagonal matrix element $-\frac{1}{2}\sqrt{3}\delta$.

The trial and error fit was actually tried for all possible pairs of states as $|1+\rangle$ and $|1-\rangle$. The best fit was obtained for 101 (in agreement with the polarization data) and -75 . With this preliminary information, the second degree terms were included in the matrix elements, the fourth

degree coefficients were chosen for a least squares fit in the energy, and the following were obtained:

E (obs)	E (calc)	State
101	108.0	$ 1+\rangle$
62	65.9	$ 2-\rangle$
57	45.4	$0.820 0\rangle + 0.572 2+\rangle$
-75	-85.8	$ 1-\rangle$
-145	-133.5	$0.820 2+\rangle - 0.572 0\rangle$

$$\langle r^4 \rangle Y_4^0 = -187.8 \text{ cm}^{-1} \quad \langle r^4 \rangle Y_4^2 = 1,141 \text{ cm}^{-1}$$

$$\langle r^4 \rangle Y_4^4 = -626.2 \text{ cm}^{-1}.$$

The fit is only fair. There are three possible sources of inaccuracy in the treatment. The first of these is the possibility of deviation from the $L-S$ coupling approximation. This would appear essentially in the values of the reduced matrix elements. Purely from the empirical standpoint, they could be treated as additional parameters to provide a better fit to the data. A second appreciable source of discrepancy is the interactions with other states, principally those within the ground multiplet. The nondiagonal Wigner coefficients and Racah coefficients are both generally of the same order of magnitude as the diagonal ones, and the reduced $L-S$ matrix elements of table 15 are appropriate for all combinations within the ground multiplet. Nondiagonal matrix elements can therefore be of appreciable magnitude and, in view of the relatively small multiplet splitting, could become quite important in a more refined theory. Finally, there is also the possibility that the symmetry is such as to require the inclusion of S_N^M terms. This almost doubles the total number of parameters. This will, of course, permit an adjustment to the data, but there is no clear requirement for the inclusion of these terms. It may be noted here again that the earlier check on multiplet separation showed a distinct anomaly for the $J=2$ group.

Let us now turn our attention to $J=3$. Our allowed states are as follows:

$$A_{2z}: |0\rangle \text{ and } |2+\rangle \quad A_{1z}: |2-\rangle$$

$$B_{1z}: |3+\rangle \text{ and } |1+\rangle \quad B_{2z}: |1-\rangle \text{ and } |3-\rangle.$$

If we compute the levels from the second and fourth degree terms only, we obtain a set of levels which show a rather surprisingly similar trend to the actually observed levels, which include the effect of sixth degree terms. Unfortunately, the polarization data both here and for still higher J does not appear to follow either the rules appropriate for D_{2h} symmetry found adequate for lower J , or any other pattern easily amenable to analysis. In particular, the states given do not especially even suggest the reported polarization data.

E (obs)	E (calc)	State
37	45.1	$0.948 1+\rangle - 0.319 3+\rangle$
32	36.9	$.756 3-\rangle - .654 1-\rangle$
27	21.8	$.784 0\rangle - .620 2+\rangle$
22	8.6	$.620 0\rangle + .784 2+\rangle$
-5	-14.8	$.319 1+\rangle + .948 3+\rangle$
-15	-28.0	$ 2-\rangle$
-98	-69.6	$0.756 1-\rangle + .654 3-\rangle$

There is obviously a considerable amount of work yet to be done before the Tb spectrum is understood. Experimentally, there is need for more attention to polarization data for fixing the character of the states involved. Theoretically, there are several lines of endeavor to be pursued. The problems become numerically so tedious that machine calculation is probably required for further success. The first of these is a systematic machine calculation of nondiagonal Wigner coefficients. The published table of Simon (1954) is inadequate in range, and the extension of table 15 by desk calculations is almost out of the question. For the most part, the tables of the Racah coefficients (Simon et al., 1954) are adequate, so that the availability of Wigner coefficients would permit a study of nondiagonal contributions to the energies. Furthermore, it appears that in problems of low symmetry, such as the present one, there will be a requirement for trial-and-error calculations. These are well adapted to machine calculations, but are again, except in very simple cases, almost out of the question for desk calculators.

7. Bibliography

- 1905
Kayser, H. G. J., *Handbuch der Spectroscopie* **3**, 427 (S. Hirzel).
- 1906
Frobenius, G. and Schur, I., *Über die reellen Darstellungen der endlichen Gruppen*, *Sitzber. preuss. Akad. Wiss., Physik-math. Kl. I*, 186.
- 1908
Becquerel, J., *Recherches sur les phenomenes magneto-optiques dans les cristaux et les solutions solidifiees a la temperature de l'air liquide Radium V*, 5.
Becquerel, J. and Onnes, H. K., *The absorption spectra of the compounds of the rare earths at the temperatures obtainable with liquid hydrogen, and their change by the magnetic field*, *Koninkl. Akad. Wetenschap. (Amsterdam)* **10**, 592.
- Tallqvist, H., *Tafeln der Kugelfunctionen*, *Acta Soc. Sci. Fennicae* **33**, No. 4. (a) *Tafeln der Abgeleiteten und Zugeordneten Kugelfunctionen Erster Art*, *Acta Soc. Sci. Fennicae* **33**, No. 9.
- du Bois, H. and Elias, G. J., *Der Einfluss von Temperatur und Magnetisierung bei selektiven Absorptions- und Fluoreszenzspektren*, *Ann. Physik* **4.27**, 233.
- 1911
du Bois, H. and Elias, G. J., *Der Einfluss von Temperatur und Magnetisierung bei selektiven Absorptions- und Fluoreszenzspektren*, *Ann. Physik* **4.35**, 617.

- 1923
Schoenflies, A. Theorie der Kristallstruktur. Berlin, Gebrüder Borntraeger.
- 1925
Hund, F., Atomtheoretische Deutung des Magnetismus der Seltenen Erden, Z. Physik **33**, 855.
- 1926
Becquerel, J., Onnes, H. K., and de Haas, W. J., The absorption bands of the compounds of the rare earths, their modification by a magnetic field, and the magnetic rotation of the plane of polarisation at very low temperatures, Koninkl. Akad. Wetenschap. (Amsterdam) **29**, 264.
- 1928
Weyl, H., Gruppentheorie und Quantenmechanik (S. Hirzel).
- 1929
Bethe, H., Termaufspaltung in Kristallen, Ann. Physik **5.3**, 133. [Splitting of Terms in Crystals, Consultants Bureau, Tr.]
Brunetti, R. and Ollano, Z., L'Azione del campo elettrico sulle radiazioni degli joni di terre rare in composto chimico, Nuovo cimento **6**, 345.
Freed, S. and Spedding, F. H., Line absorption spectra in solids at low temperatures in the visible and ultraviolet regions of the spectrum, Nature **123**, 525.
Kramers, M. H. A., La rotation paramagnétique du plan de polarisation dans les cristaux uniaxes de terres rares, Koninkl. Akad. Wetenschap. (Amsterdam) **32**, 1176.
Van Vleck, J. H. and Frank, A., The effect of second order Zeeman terms on magnetic susceptibilities in the rare earth and iron groups, Phys. Rev. **34**, 1494.
- 1930
Bethe, H., Zur Theorie des Zeemaneffektes an den Salzen der seltenen Erden, Z. Physik **60**, 218.
Eckart, C., The application of group theory to the quantum dynamics of monatomic systems, Rev. Modern Phys. **2**, 305.
Kramers, H. A., Theorie generale de la rotation paramagnétique dans les cristaux, Koninkl. Akad. Wetenschap. (Amsterdam) **33**, 959.
Wigner, E., Über die elastischen Eigenschwingungen symmetrischer Systeme, Nachr. Akad. Wiss. Göttingen, Math.-physik. Kl. **2**, 133.
- 1931
Spedding, F. H., Interpretation of the spectra of rare earth crystals, Phys. Rev. **37**, 777.
Wigner, E., Gruppentheorie (F. Vieweg und Sohn).
- 1932
Penney, W. G. and Schlapp, R., Influence of crystalline fields on the spectra of salts of paramagnetic ions. I. The rare earths, especially Pr and Nd, Phys. Rev. **41**, 194.
Schlapp, R. and Penney, W. G., Influence of crystalline fields on the susceptibilities of salts of paramagnetic ions. II. The iron group, especially Ni, Cr, and Co, Phys. Rev. **42**, 666.
Wigner, E., Über die Operation der Zeitumkehr in der Quantenmechanik, Nachr. Akad. Wiss. Göttingen, Math.-physik. Kl. **5**, 546.
Spedding, F. H. and Bear, R. S., Absorption spectrum of the samarium ion in solids. I. Absorption by large single crystals of $\text{SmCl}_2 \cdot 6\text{H}_2\text{O}$, Phys. Rev. **42**, 58.
Van Vleck, J. H., Theory of electric and magnetic susceptibilities (Oxford University Press, Oxford, England).
- 1933
Spedding, F. H., Zeeman effect in solids, J. Chem. Phys. **1**, 144.
- 1935
Condon, E. U. and Shortley, G. H., Theory of atomic spectra (Cambridge University Press, England).
- 1936
Lang, R. J., The spectrum of trebly ionized cerium, Can. J. Research **14A**, 127.
- 1937
Ahlberg, J. E., Blanchard, E. R., and Lundberg, W. O., Electronic activation in crystals: The heat capacities of $\text{Sm}_2(\text{SO}_4)_3 \cdot 8\text{H}_2\text{O}$ and $\text{Nd}_2(\text{SO}_4)_3 \cdot 8\text{H}_2\text{O}$ from 3 to 40°K, J. Chem. Phys. **5**, 552.
Van Vleck, J. H., The puzzle of rare earth spectra in solids, J. Phys. Chem. **41**, 67.
- 1938
Murnaghan, F. D., Theory of group representations (Johns Hopkins Press, Baltimore, Md.).
- 1940
Opechowski, W., Sur les groupes cristallographiques "Doubles", Physica **7**, 552.
- 1941
Stratton, J. A., Electromagnetic theory (McGraw Hill Book Co., Inc., New York, N.Y.), pg. 408.
- 1942
Racah, G., Theory of complex spectra I, Phys. Rev. **61**, 186. (a) Theory of complex spectra II, Phys. Rev. **62**, 438. (b) Theory of complex spectra III, Phys. Rev. **63**, 367 (1943).
- 1945
U.S. National Bureau of Standards, Tables of associated Legendre functions (Columbia University Press, New York, N.Y.).
- 1948
Hellwege, K. H., Elektronenterme und Strahlung von Atomen in Kristallen. I. Termaufspaltung und elektrische Dipolstrahlung, Ann. Physik **6.4**, 97. (a) Elektronenterme und Strahlung von Atomen in Kristallen. II. Magnetische Dipolstrahlung, Ann. Physik **6.4**, 127. (b) Elektronenterme und Strahlung von Atomen in Kristallen. III. Elektrische Quadrupolstrahlung, Ann. Physik **6.4**, 136. (c) Elektronenterme und Strahlung von Atomen in Kristallen. IV. Symmetrieentartung und Kramerssche Entartung **6.4**, 143. (d) Elektronenterme und Strahlung von Atomen in Kristallen. V. Kubische Kristalle, Ann. Physik **6.4**, 150.
- 1949
Racah, G., Theory of complex spectra IV, Phys. Rev. **76**, 1352.
- 1952
Biedenharn, L. C., Tables of the Racah Coefficients, Oak Ridge Nat'l Lab. Rep. **1098**.
Falkoff, D. L., Colladay, G. S., and Sells, R. E., Transformation amplitudes for vector addition of angular momentum ($j3mm'j3JM$), Can. J. Phys. **30**, 253.
Klein, M. J., On a degeneracy theorem of Kramers, Am. J. Phys. **20**, 65.
Stevens, K. W. H., Matrix elements and operator equivalents connected with the magnetic properties of rare earth ions, Proc. Phys. Soc. (London) **A.55**, 209.
- 1953
Bleaney, B. and Stevens, K. W. H., Paramagnetic resonance, Repts. Prog. in Phys. **16**, 108.
Elliott, R. J. and Stevens, K. W. H., The theory of magnetic resonance experiments on salts of the rare earths, Proc. Roy. Soc. (London) **A218**, 553. (a) The magnetic properties of certain rare earth ethyl sulphates, Proc. Roy. Soc. (London) **A219**, 387.
- 1954
Simon, A., A numerical table of the Clebsch-Gordan coefficients, Oak Ridge Nat'l Rep. **1718**.
Simon, A., Vander Sluis, J. H., and Biedenharn, L. C., Tables of the Racah coefficients, Oak Ridge Nat'l Rep. **1679**.

1955

Bowers, K. D. and Owen, J., Paramagnetic resonance. II., Repts. Prog. in Phys. **18**, 304.

Judd, R. B., Operator equivalents and matrix elements for the excited states of rare earth ions, Proc. Roy. Soc. (London) **A.227**, 552. (a) The theory of the spectra of europium salts, Proc. Roy. Soc. (London) **A.228**, 120.

Mulliken, R. S. Report on notation for the spectra of polyatomic molecules, J. Chem. Phys. **23**, 1907.

1956

Dieke, G. H. and Crosswhite, H. M., Absorption spectrum and magnetic properties of ytterbium chloride, J. Opt. Soc. Am. **46**, 885.

Dieke, G. H. and Heroux, L., Absorption spectra, Zeeman effects, and magnetic properties of neodymium salts. Phys. Rev. **103**, 1227.

Melvin, M. A., Simplification in finding symmetry-adapted eigenfunctions, Rev. Mod. Phys. **28**, 18.

WASHINGTON, January 15, 1960

1957

Edmonds, A. R., Angular momentum in quantum mechanics (Princeton University Press, Princeton, N.J.).

Fick, E. and Joos, G., Kristallspektren, Handbuch der Physik **28**, 205.

Rose, M. E., Elementary theory of angular momentum (John Wiley & Sons, Inc., 440 Fourth Ave., New York, N.Y.).

Singh, Shobha, Absorption, fluorescence, and Zeeman effects of dysprosium, terbium, holmium and uranyl compounds at low temperatures, Dissertation, Johns Hopkins U.

1958

Low, W., Paramagnetic resonance spectrum of trivalent gadolinium in the cubic field of calcium fluoride, Phys. Rev. **109**, 265.

1960

Herzfeld, C. M. and Meijer, P. H. E., Group theory and crystal field theory (to be published by Academic Press, Inc., New York, New York).

THE NATIONAL BUREAU OF STANDARDS

The scope of activities of the National Bureau of Standards at its major laboratories in Washington, D.C., and Boulder, Colorado, is suggested in the following listing of the divisions and sections engaged in technical work. In general, each section carries out specialized research, development, and engineering in the field indicated by its title. A brief description of the activities, and of the resultant publications, appears on the inside of the front cover.

WASHINGTON, D.C.

Electricity and Electronics. Resistance and Reactance. Electron Devices. Electrical Instruments. Magnetic Measurements. Dielectrics. Engineering Electronics. Electronic Instrumentation. Electrochemistry.

Optics and Metrology. Photometry and Colorimetry. Optical Instruments. Photographic Technology. Length. Engineering Metrology.

Heat. Temperature Physics. Thermodynamics. Cryogenic Physics. Rheology. Molecular Kinetics. Free Radicals Research.

Atomic Physics. Spectroscopy. Radiometry. Mass Spectrometry. Solid State Physics. Electron Physics. Atomic Physics.

Radiation Physics. Neutron Physics. Radiation Theory. Radioactivity. X-ray. High Energy Radiation. Nucleonic Instrumentation. Radiological Equipment.

Chemistry. Organic Coatings. Surface Chemistry. Organic Chemistry. Analytical Chemistry. Inorganic Chemistry. Electrodeposition. Molecular Structure and Properties of Gases. Physical Chemistry. Thermochemistry. Spectrochemistry. Pure Substances.

Mechanics. Sound. Mechanical Instruments. Fluid Mechanics. Engineering Mechanics. Mass and Scale. Capacity, Density, and Fluid Meters. Combustion Controls.

Organic and Fibrous Materials. Rubber. Textiles. Paper. Leather. Testing and Specifications. Polymer Structure. Plastics. Dental Research.

Metallurgy. Thermal Metallurgy. Chemical Metallurgy. Mechanical Metallurgy. Corrosion. Metal Physics.

Mineral Products. Engineering Ceramics. Glass. Refractories. Enamels. Metals. Constitution and Microstructure.

Building Technology. Structural Engineering. Fire Protection. Air Conditioning, Heating, and Refrigeration. Floor, Roof, and Wall Coverings. Codes and Safety Standards. Heat Transfer. Concreting Materials.

Applied Mathematics. Numerical Analysis. Computation. Statistical Engineering. Mathematical Physics.

Data Processing Systems. SEAC Engineering Group. Components and Techniques. Digital Circuitry. Digital Systems. Analog Systems. Applications Engineering.

• Office of Basic Instrumentation.

• Office of Weights and Measures.

BOULDER, COLORADO

Cryogenic Engineering. Cryogenic Equipment. Cryogenic Processes. Properties of Materials. Gas Liquefaction.

Radio Propagation Physics. Upper Atmosphere Research. Ionosphere Research. Regular Prediction Services. Sun-Earth Relationships. VHF Research. Radio Warning Services. Airglow and Aurora. Radio Astronomy and Arctic Propagation.

Radio Propagation Engineering. Data Reduction Instrumentation. Radio Noise. Tropospheric Measurements. Tropospheric Analysis. Propagation-Terrain Effects. Radio-Meteorology. Lower Atmosphere Physics.

Radio Standards. High-Frequency Electrical Standards. Radio Broadcast Service. Radio and Microwave Materials. Atomic Frequency and Time Standards. Electronic Calibration Center. Microwave Circuit Standards.

Radio Communication and Systems. Low Frequency and Very Low Frequency Research. High Frequency and Very High Frequency Research. Modulation Systems. Antenna Research. Navigation Systems. Systems Analysis. Field Operations.