

**Best  
Available  
Copy**

AD-A280 279



# ATOMIC ENERGY LEVELS

As Derived From the Analyses of Optical Spectra

Volume III

42Mo-57La


72Hf-89Ac



CIRCULAR 467

BEST AVAILABLE COPY


UNITED STATES DEPARTMENT OF COMMERCE  
NATIONAL BUREAU OF STANDARDS



**COPY 4**

**LIBRARY COPY**



**94-14776**  
 *284 ps*

**94 5 17 094**

UNITED STATES DEPARTMENT OF COMMERCE, Sinclair Weeks, Secretary  
NATIONAL BUREAU OF STANDARDS, A. V. Astin, Director

# ATOMIC ENERGY LEVELS

As Derived From the Analyses of Optical Spectra

Volume III

The Spectra of Molybdenum, Technetium, Ruthenium, Rhodium, Palladium, Silver, Cadmium, Indium, Tin, Antimony, Tellurium, Iodine, Xenon, Cesium, Barium, Lanthanum—Hafnium, Tantalum, Tungsten, Rhenium, Osmium, Iridium, Platinum, Gold, Mercury, Thallium, Lead, Bismuth, Polonium, Radon, Radium, and Actinium

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Dist	Availability or Special
A-1	

BY CHARLOTTE E. MOORE



DTIC QUALITY INSPECTED 3

Circular of the National Bureau of Standards 467

Issued May 1, 1958

---

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C.  
Price \$3.00



## Preface

The present Volume, including data to December 1957, is the third of a series being prepared at the National Bureau of Standards under a project on the compilation of atomic energy levels derived from the analyses of optical spectra.

Volume I, published in June 1949, contains the energy levels of 206 spectra of the elements Hydrogen through Vanadium,  $Z=1$  through 23. Volume II, completed in August 1952, includes similar data for 152 spectra of the elements Chromium through Niobium,  $Z=24$  through 41. Volume III covers the elements Molybdenum through Lanthanum,  $Z=42$  through 57; and Hafnium through Actinium,  $Z=72$  through 89. It includes 124 spectra, and is identical in arrangement with Volumes I and II. The form of presentation is that recommended by the majority of interested scientists who received a questionnaire proposed by the National Research Council Committee on Line Spectra of the Elements, when the program was started in 1946.

The manuscript has been prepared by Charlotte E. Moore under the direction of William F. Meggers, Chief of the Spectroscopy Section of the Atomic and Radiation Physics Division. The success of this project has depended heavily upon the cordial and continuing collaboration of the many spectroscopists who have made an unusual effort to furnish data in advance of publication. Their efforts together with the unflinching cooperation of the National Research Council Committee are greatly appreciated.

Perhaps the most difficult phase of this extensive program lies ahead. The two groups of rare-earth spectra (58 Cerium through 71 Lutetium, and 90 Thorium through 100 Fermium) remain for Volume IV. At best, the preparation of this Volume within the next few years will tax the most experienced and talented workers available. These spectra are the most intricate and least known of atomic spectra. Consequently the type of collaboration we have enjoyed with qualified spectroscopists in the preparation of the first three Volumes of the "Atomic Energy Levels" series will have not only to continue but be substantially augmented if the fourth Volume is to be brought to satisfactory fruition.

A. V. ASTIN, *Director*

WASHINGTON, D. C., July 1, 1957.

230

### Contents

	Page		Page
Preface.....	II	5. The Periodic Table.....	VIII
Index to Spectra.....	IV	5.1 The chemical elements by atomic number—ionization potentials for spectra of the elements H to La and Hf to Ac (table 34).....	VIII
1. Introduction.....	VII	5.2 The chemical elements by chemical symbol (table 35).....	IX
2. Scope of the Present Tables.....	VII	5.3 The periodic system (table 36).....	IX
3. Arrangement.....	VII	5.4 Index— <i>isoelectronic</i> sequences (table 37).....	IX
4. Tables of Predicted and Observed Arrays of Terms.....	VIII	6. References.....	IX
		7. Acknowledgments.....	X

A list of additions and corrections to Volumes I and II is appended, pages 238 to 245.

### List of Tables

Table	Subject	Page	Table	Subject	Page
PREDICTED TERMS			PREDICTED TERMS—continued		
1	Mo I	XII	19	Re I	XXV
2	Te I	XIII	20	Os I	XXVI
3	Ru I	XIV	21	Ir I	XXVII
4	Rh I	XV	22	Pt I	XXVIII
5	Pd I	XVI	23	Au I	XXVIII
6	Ag I	XVI	24	Hg I	XXVIII
7	Cd I	XVII	25	Tl I	XXIX
8	In I	XVII	26	Pb I	XXIX
9	Sn I	XVIII	27	Bi I	XXX
10	Sb I	XVIII	28	Po I	XXX
11	Te I	XIX	29	At I	XXXI
12	I I	XIX	30	Rn I	XXXI
13	Xe I	XX	31	Ra I	XXXII
14	Ba I	XX	32	Ac I	XXXII
15	La I	XXI	33	References to Rare-Earth Spectra	XXXIII
**			34	Ionisation Potentials	XXXIV
16	Hf I	XXII	35	Chemical Symbols	XXXVI
17	Ta I	XXIII	36	The Periodic System	XXXVII
18	W I	XXIV	37	Index— <i>isoelectronic</i> Sequences	XXXVIII

\*\* The two groups of rare-earth spectra, Ce to Lu (E-68 to 71) and Th to Fm (E-88 to 100) will be included in Vol. IV. See references to these spectra in Table 28.

C-45166-A

### Index to Spectra

Element	Z	Spectrum	Page	Element	Z	Spectrum	Page		
Molybdenum	42	Mo I.....	1	Tin	50	Sn I.....	74		
		Mo II.....	7			Sn II.....	80		
		Mo III.....	11			Sn III.....	82		
		Mo IV.....	12			Sn IV.....	83		
		Mo V.....	13			Sn V.....	84		
		Mo VI.....	14			Sn VI.....	85		
		Mo VII.....	15			Sn XXII.....	86		
		Mo VIII.....	15			Sn XXIV.....	86		
Mo XVI.....	16	Antimony	51			Sb I.....	87		
Technetium	43					Tc I.....	17	Sb II.....	90
						Tc II.....	19	Sb III.....	92
Ruthenium	44					Ru I.....	20	Sb IV.....	93
				Ru II.....	25	Sb V.....	94		
				Ru III.....	28	Sb VI.....	95		
Rhodium	45	Rh I.....	29	Sb XXIII.....	95				
		Rh II.....	32	Tellurium	52	Te I.....	96		
		Rh III.....	34			Te II.....	98		
Palladium	46	Pd I.....	38			Te III.....	100		
		Pd II.....	41			Te IV.....	101		
		Pd III.....	44			Te V.....	102		
		Pd XVIII.....	47			Te VI.....	103		
Silver	47	Pd XX.....	47			Te VII.....	104		
		Ag I.....	48	Iodine	53	I I.....	105		
		Ag II.....	50			I II.....	108		
		Ag III.....	52			I III.....	110		
		Ag XIX.....	54			I V.....	111		
Ag XXI.....	54	I VI.....	111						
Cadmium	48	Cd I.....	55	I VII.....	112				
			Cd II.....	58	I VIII.....	112			
			Cd III.....	60	Xenon	54	Xe I.....	113	
			Cd IV.....	62			Xe II.....	118	
			Cd XX.....	63			Xe III.....	120	
			Cd XXII.....	63			Xe IV.....	123	
Indium	49	In I.....	64	Cesium	55	Cs I.....	124		
			In II.....			67	Cs II.....	128	
			In III.....			70	Cs III.....	130	
			In IV.....	71	Barium	56	Ba I.....	131	
			In V.....	72			Ba II.....	134	
			In XXI.....	73			Ba IV.....	135	
			In XXIII.....	73	Lanthanum	57	La I.....	136	
Literature References to Rare-Earth Spectra....	XXXIII	La II.....	139						
			La III.....	142					

### Index to Spectra—Continued

Element	Z	Spectrum	Page	Element	Z	Spectrum	Page
Hafnium	72	Hf I.....	143	Thallium	81	Tl I.....	202
		Hf II.....	146			Tl II.....	204
Tantalum	73	Ta I.....	149			Tl III.....	206
		Ta II.....	154			Tl IV.....	207
Tungsten	74	W I.....	156	Lead	82	Pb I.....	208
		W II.....	161			Pb II.....	211
Rhenium	75	Re I.....	164			Pb III.....	213
		Re II.....	168			Pb IV.....	215
Osmium	76	Os I.....	171			Pb V.....	217
		Os II.....	176	Bismuth	83	Bi I.....	219
Iridium	77	Ir I.....	177			Bi II.....	221
		Platinum	78			Pt I.....	181
Pt II.....	183					Bi IV.....	223
Gold	79	Au I.....	186			Bi V.....	224
		Au II.....	189			Bi VI.....	225
Mercury	80	Hg I.....	191	Polonium	84	Po I.....	227
		Hg II.....	196			Radon	86
		Hg III.....	198	Rn II.....	230		
		Hg IV.....	200	Radium	88	Ra I.....	231
Actinium	89	Ac I.....	234			Ra II.....	233
		Ac II.....	235	Actinium	89	Ac I.....	234
		Ac III.....	237			Ac II.....	235

## 1. Introduction

The present Volume is the third of a series being prepared at the National Bureau of Standards in continuation of a project started in 1946. At that time the National Research Council Committee on Line Spectra of the Elements met to discuss details regarding the preparation, in the Spectroscopy Section of this Bureau, of a critical compendium of Atomic Energy Levels. The need for

such a program was stressed because Bacher and Goudsmit did not contemplate revising their very useful book on "Atomic Energy States as Derived from the Analyses of Optical Spectra," published in 1932. The demand for these Volumes has abundantly justified the need anticipated by the Committee in formulating and supporting this extensive project.

## 2. Scope of the Present Tables

This Volume covers the elements Mo to La ( $Z=42$  to 57) and Hf to Ac ( $Z=72$  to 89), thus completing the Periodic Table except for the two groups of rare-earth elements, which will comprise Volume IV. Pending the completion of Volume IV, which is by far the most difficult of all, a current list of references to analyses of rare-earth spectra is included here in Table 33, p. xxxiii.

In spite of requests to increase the scope of the tables it has been decided to retain the general format of Volumes I and II throughout.

As the atomic number increases, the spectra become more complex. Because of configuration-interaction and breakdown of  $LS$ -coupling it is increasingly difficult to make term-designation assignments that are significant. Consequently, the present Volume has many more miscellaneous levels than are found among spectra of lighter elements. It is hoped that this compendium will provide useful data for workers interested in future studies of the general problems of configuration-interaction and coupling, even though many of the spectra are incompletely analyzed.

The important subject of hyperfine structure is handled only by general references to bibliographies on the subject. In the present Volume the publications by Walchli [1]<sup>1</sup> and by Klinkenberg [2] replace the earlier bibliographies

by Mack [3] and by Meggers [4] for many spectra. In addition, selected papers are listed for a very few spectra, particularly if they do not appear in current bibliographies summarizing hyperfine-structure data. Workers in this field will find these Volumes seriously inadequate.

Since these tables do not include the lists of observed spectral lines from which the atomic energy levels are derived, the users must consult the individual papers on the analyses of the separate spectra for wavelengths, intensities, and line classifications. This difficulty is being handled, in part, by the preparation of "An Ultraviolet Multiplet Table" [5] for selected spectra. Each Volume of "Atomic Energy Levels" is accompanied by a Section of the Ultraviolet Multiplet Table covering the same range of elements. Section 3 of this Table is now in course of preparation. For multiplet data on individual spectra in the wavelength range longer than 3000Å the "Revised Multiplet Table" [6] will suffice to some extent. A revision of this 1945 edition is in course of preparation.

Urgent requests for Grotrian Diagrams to accompany these tables have been met in part by the preparation of selected *Partial Grotrian Diagrams of Astrophysical Interest*, accompanied by tables of related lines in individual spectra of the isoelectronic sequences represented by the diagrams [7].

## 3. Arrangement

The arrangement of the data in the various columns of the table, and the notation are described in detail in Volume I, and will not be repeated here, since the same style has been followed throughout. In general, all levels of the "odd" type are printed in italics.

The letters in parentheses following the literature references have the following meaning:

IP	Ionization potential.
T	Terms.
CL	Classified lines.

GD	Grotrian diagram.
ED	Energy diagram.
ZE	Zeeman effect.
IS	Isotope shift.
hfs	Hyperfine structure.

These letters describe briefly the scope and content of the paper. If no such letters follow a reference, the paper is mentioned in the text for the spectrum in question. For example, in general no letter accompanies references to theoretical papers.

<sup>1</sup> Figures in brackets indicate literature references on p. ix.

#### 4. Tables of Predicted and Observed Arrays of Terms

As in the preceding Volumes, for complex spectra whose analyses are not seriously incomplete, arrays of observed terms follow the individual listings of atomic energy levels in the table. Similar, but more extensive, arrays of terms

predicted by theory for spectra of the different isoelectronic sequences are included in Tables 1 to 32 (pages XII to XXXII) as follows:

Table	Sequence	Table	Sequence	Table	Sequence	Table	Sequence
1	Mo I	9	Sn I	17	Ta I	25	Tl I
2	Te I	10	Sb I	18	W I	26	Pb I
3	Ru I	11	Te I	19	Re I	27	Bi I
4	Rh I	12	I I	20	Os I	28	Po I
5	Pd I	13	Xe I	21	Ir I	29	At I
6	Ag I	14	Ba I	22	Pt I	30	Rn I
7	Cd I	15	La I**	23	Au I	31	Ra I
8	In I	16	Hf I	24	Hg I	32	Ac I

Leaders indicate that predicted terms from additional electrons have been omitted.

For the Xe I and Rn I sequences the tables give predicted terms (*LS*-coupling) and predicted pairs of levels (*JL*-coupling), as has been done for the preceding spectra of the inert-gas type. The pair-coupling notation in the general form suggested by Racah [8] has been adopted as before, to take into account the departure from *LS*-coupling.

In the arrays of predicted terms the order follows in a general way that of the limit terms of the first spectrum of the sequence. Consequently, in successive Volumes the arrays are not identically arranged, since different configurations and limit-terms predominate as *Z* increases.

Predicted terms in configurations involving equivalent electrons are summarized in a number of general references to atomic spectra. [9]

#### 5. The Periodic Table

##### 5.1 The Chemical Elements by Atomic Number; Ionization Potentials (Table 34)

Requests for ionization potentials have exceeded all others for data in these Volumes. A number of those published in Volumes I and II have subsequently been revised. Ionization potentials for all spectra in Volumes I, II, and III, so far as known, are listed in Table 34, which supersedes those published earlier. Column one gives the atomic number *Z*; column two the chemical symbol of the element. Successive columns contain the ionization potentials of the atoms or ions in successive stages of ionization, *i* denoting first spectra (neutral atoms); *ii*, second spectra (singly ionized atoms); etc.

Throughout these Volumes the ionization potentials are derived by multiplying the limit in  $\text{cm}^{-1}$  (*K*) by the conversion factor 0.00012395, to express it in electron volts. This factor was recommended by Birge in 1941 [10] and adopted by the Joint Commission for Spectroscopy in 1953 "to conform to the standard book 'Atomic Energy Levels'. With the 1951 values of *h*, *c*, and *e* the value for  $\lambda_0$  is between 12396 and 12397". [11, 12]

The writer takes this opportunity to point out that she misquoted the limits for the spectra of the C I and N I sequences in Volume I. The resulting corrected ionization potentials are included both in Table 34 and in the list of corrections facing blank page 238.

Although series are not known for all the listed spectra of a given stage of ionization in the respective periods of the Periodic Table, yet it is possible to derive fairly reliable limits in cases where well established series are known for neighboring spectra. Russell has discussed this for the group of second spectra from Ca to Zn [13]. Similarly, Catalán and Velasco [14] have determined limits and ionization potentials for the first, second, and third spectra of the iron group. Catalán and Rico give similar data for the first and second spectra of the palladium group [15]. Rico has generously furnished revised values for the second spectra of this group, and for the third spectra Y III to In III, in advance of publication [16].

From a study of screening constants, Finkelnburg and Humbach have interpolated ionization potentials for the entire periodic table [17].

### 5.2 The Chemical Elements by Chemical Symbol (Table 35)

For convenience of cross reference, the chemical elements are listed in Table 35 in the alphabetical order of the chemical symbol. This Table is identical in form with Table 24 in Volume I, and Table 19 of Volume II. It includes the four additional elements, 99 Einsteinium, 100 Fermium, 101 Mendelevium, and 102 Nobelium, whose names have been adopted by the International Union of Pure and Applied Chemistry Commission on Nomenclature (Section of Inorganic Chemistry).

### 5.3 The Periodic System (Table 36)

Table 36 is identical in form with Table 25 of Volume I and Table 20 of Volume II. The general arrangement is similar to that given by Condon and Shortley on page 333 of "The Theory of Atomic Spectra" [18].

### 5.4 Index—Isoelectronic Sequences (Table 37)

This table is the index to the data in Volume III, i. e., the spectra from Mo through La and Hf through Ac, arranged similarly to the indices for Volumes I and II.

Column I gives the atomic number  $Z$  followed by the chemical symbol under the heading "Element." The remaining columns indicate the successive stages of ionization under the general heading "Spectrum," I denoting first spectra (neutral atoms), II second spectra (singly-ionized atoms), III third spectra, etc. The numbers in the body of the table indicate the page on which the data for the individual spectra may be found.

In this table, isoelectronic spectra appear on the diagonals. Alternate diagonals are printed in bold face type to emphasize the spectra of each sequence. Blanks occur for spectra in which structure is not yet known.

No isoelectronic sequences are carried into the rare-earth groups of spectra in this Volume, but they will be continued in Volume IV for the two groups of spectra of elements with  $Z > 57$  and  $Z > 89$ , respectively. The isoelectronic sequences in this part of the periodic table are, in general, very short. Those started but not completed in Volume III are as follows:

Sequence	Spectrum	Sequence	Spectrum
Ca I	Ce IV	Fr I	Th IV
Ba I	Ce III	Ra I	Th III
La I	Ce II	Ac I	Th II

## 6. References

- [1] H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL—1469, 134 pp. (1953); Suppl. II (Physics), 43 pp. (1955).
- [2] P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63-73 (April 1952).
- [3] J. E. Mack, *Rev. Mod. Phys.* **22**, No. 1, 64-76 (January 1950).
- [4] W. F. Meggers, *J. Opt. Soc. Am.* **36**, 431-448 (August 1946).
- [5] C. E. Moore, *Circ. Nat. Bur. Std.* 488, Section 1, H to V (1950); Section 2, Cr to Nb (1952); Section 3, Mo to La and Hf to Ac, in press (1958).
- [6] C. E. Moore, *Princeton Univ. Obs. Contr. No.* 20 (1945).
- [7] P. W. Merrill, *Lines of the Chemical Elements in Astronomical Spectra*, Appendix A, Carnegie Inst. Wash. Publ. 610, 103-165 (1956).
- [8] G. Racah, *Phys. Rev.* **61**, 537 (L) (1942).
- [9] H. N. Russell, *Phys. Rev.* **29**, 782 (1927); R. C. Gibbs, D. T. Wilber, and H. E. White, *Phys. Rev.* **29**, 790 (1927); R. F. Bacher and S. Goudsmit, *Atomic Energy States*, p. 10 (McGraw-Hill Book Co., Inc., New York and London, 1932); F. Hund, *Linienpektren und Periodisches System der Elemente* (Julius Springer, Berlin, 1927); C. L. B. Shudeman, *J. Franklin Inst.* **224**, 501 (1937) (Terms from equivalent  $g$ -,  $h$ -, and  $i$ -electrons).
- [10] R. T. Birge, *Rev. Mod. Phys.* **13**, No. 4, 237 (1941).
- [11] *Trans. Joint Commission for Spectroscopy*, *J. Opt. Soc. Am.* **43**, 412 (1953).
- [12] The writer used the factor 0.00012345 in preparing the multiplet tables, which explains the discrepancies between the ionization potentials quoted in the Multiplet Tables [5,6] and those used in these Volumes.
- [13] H. N. Russell, *J. Opt. Soc. Am.* **40**, 518, 1950.
- [14] M. A. Catalán y R. Velasco, *An. Real Soc. Esp. Fisica y Quimica (Madrid) [A]* **48**, 247 (1952).
- [15] M. A. Catalán y F. R. Rico, *An. Real Soc. Esp. Fisica y Quimica (Madrid) [A]* **48**, 328 (1952).
- [16] F. R. Rico, unpublished material (December 1956); M. A. Catalán y F. R. Rico, *An. Real Soc. Esp. Fisica y Quimica (Madrid) [A]* **53**, 85 (1957).
- [17] W. Finkelnburg und W. Humbach, *Naturwiss.* **43**, 35 (1955).
- [18] E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra*, (The University Press, Cambridge, England; The Macmillan Co., N. Y., corrected edition, 1951).

## 7. Acknowledgments

The present Volume could not have been prepared without unrelenting effort on the part of spectroscopists at home and abroad. At the National Bureau of Standards, W. F. Meggers, Chief of the Spectroscopy Section and Chairman of the National Research Council Committee on Line Spectra of the Elements, has given generously of his time and expert advice on numerous difficult questions, and has painstakingly supervised the work throughout. He, with the assistance of his coworkers, has contributed unpublished analyses for more spectra than any other single individual, 13 spectra in all. C. C. Kiess has likewise arranged his entire schedule to fit into this program. The unpublished data from him and his colleagues on six complex spectra, have greatly enhanced the value of this Volume. His spectra start Volume III (Mo I and Mo II) as was the case with Volume II (Cr I and Cr II). Many helpful suggestions from the standpoint of theory have been made by R. E. Trees. K. Kessler with the assistance of the computational laboratory has carried out work on several spectra with the digital computer. Much of the observing on the above programs has been done by C. H. Corliss and W. R. Bozman. All members of the Spectroscopy Section have been most willing collaborators.

At Princeton, A. G. Shenstone has arranged his entire research program to supply data for this Volume. Except for his splendid work, a number of spectra would have been seriously incomplete. The late H. N. Russell also continued to take a keen interest in the work until the time of his death.

At the Oak Ridge National Laboratory, J. R. McNally, Jr., P. M. Griffin, and others have cooperated, as have also M. Fred and F. S. Tomkins at Argonne National Laboratory.

J. E. Mack and his staff, and D. D. Laun in Wisconsin, C. J. Humphreys in Corona, Calif., and R. A. Fisher at Northwestern University have been cordial collaborators.

The response from abroad has also been gratifying. In Madrid, the late M. A. Catalán and his entire staff spared no effort in furnishing analyses especially for inclusion here. In Amsterdam, P. F. A. Klinkenberg and J. C. van den Bosch and their students have been very cooperative. G. Racah, in Jerusalem, has been a valuable consultant. R. F. Barrow and H. C. Rowlinson in England; B. Edlén in Sweden; and a group from Canada, A. M. Crooker, R. Nodwell, J. N. P. Hume, and M. F. Crawford have all provided unpublished data.

The writer is deeply indebted to these many contributors, but the users of this Volume, who will have the lasting benefit of this collaboration, are perhaps more deeply indebted. It is sincerely hoped that equally generous support will continue throughout the preparation of Volume IV, which will include the most difficult of all atomic spectra, namely those of the rare-earths.

The services of many experts are required to publish these data. Miss Sarah A. Jones, the Librarian of the Bureau, and her staff, have handled most efficiently the numerous requests for literature references. The splendid work of J. L. Mathusa and the staff in the Publications Section of the Bureau, on the details of publishing these complex tables, has attracted the attention of many users. J. E. Carpenter has been a most helpful consultant. Special thanks are due also to Mrs. Betty L. Arnold whose competence in handling the publication details has been invaluable. The personnel in the Government Printing Office have also been most cooperative. Mrs. Isabel D. Murray has provided competent technical assistance throughout the work.

The writer takes great pleasure in recording here her appreciation of all the generous assistance she has received. The most lasting pleasure comes from the willingness of so many to contribute unselfishly to these Volumes.



## **Tables**

**Predicted Terms: Mo I to La I**

**Hf I to Ac I**

**References to Rare-Earth Spectra**

**Ionization Potentials**

**Chemical Symbols**

**The Periodic System**

**Index—Isoelectronic Sequences**

TABLE I. PREDICTED TERMS OF THE MO I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3d <sup>2</sup> 4s <sup>2</sup> +	Predicted Terms										
	ns (n ≥ 5)			np (n ≥ 5)				nd (n ≥ 5)			.....
4d <sup>2</sup> 5s <sup>2</sup>	<sup>1</sup> S <sup>3</sup> S	<sup>1</sup> D <sup>3</sup> D	<sup>1</sup> F <sup>3</sup> F	<sup>1</sup> G <sup>3</sup> G	<sup>1</sup> H <sup>3</sup> H						
4d <sup>2</sup>	<sup>1</sup> S <sup>3</sup> S	<sup>1</sup> D <sup>3</sup> D	<sup>1</sup> F <sup>3</sup> F	<sup>1</sup> G <sup>3</sup> G	<sup>1</sup> H <sup>3</sup> H						
4d <sup>2</sup> ( <sup>1</sup> S)ns	<sup>1</sup> S <sup>3</sup> S					<sup>1</sup> P° <sup>3</sup> P°		<sup>1</sup> D <sup>3</sup> D			
4d <sup>2</sup> 5s( <sup>1</sup> D)ns		<sup>1</sup> D <sup>3</sup> D				<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G <sup>3</sup> G		
4d <sup>2</sup> ( <sup>1</sup> G)ns				<sup>1</sup> G <sup>3</sup> G			<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H° <sup>3</sup> H°		
4d <sup>2</sup> ( <sup>1</sup> F)ns	<sup>1</sup> P <sup>3</sup> P					<sup>1</sup> S° <sup>3</sup> S°	<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°		<sup>1</sup> P <sup>3</sup> P	<sup>1</sup> D <sup>3</sup> D
4d <sup>2</sup> ( <sup>1</sup> D)ns		<sup>1</sup> D <sup>3</sup> D				<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G <sup>3</sup> G		
4d <sup>2</sup> ( <sup>3</sup> D)ns		<sup>1</sup> D <sup>3</sup> D				<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G <sup>3</sup> G		
4d <sup>2</sup> ( <sup>1</sup> I)ns				<sup>1</sup> I <sup>3</sup> I					<sup>1</sup> G <sup>3</sup> G	<sup>1</sup> H <sup>3</sup> H	<sup>1</sup> I <sup>3</sup> I
4d <sup>2</sup> ( <sup>1</sup> F)ns			<sup>1</sup> F <sup>3</sup> F				<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H <sup>3</sup> H	
4d <sup>2</sup> 5s( <sup>1</sup> D)ns		<sup>1</sup> D <sup>3</sup> D				<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G <sup>3</sup> G		
4d <sup>2</sup> ( <sup>1</sup> F)ns			<sup>1</sup> F <sup>3</sup> F				<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H <sup>3</sup> H	
4d <sup>2</sup> 5s( <sup>1</sup> H)ns				<sup>1</sup> H <sup>3</sup> H				<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H° <sup>3</sup> H°	<sup>1</sup> I° <sup>3</sup> I°	
4d <sup>2</sup> ( <sup>1</sup> G)ns				<sup>1</sup> G <sup>3</sup> G			<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H° <sup>3</sup> H°		
4d <sup>2</sup> 5s( <sup>1</sup> P)ns	<sup>1</sup> P <sup>3</sup> P					<sup>1</sup> S° <sup>3</sup> S°	<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°		<sup>1</sup> P <sup>3</sup> P	<sup>1</sup> D <sup>3</sup> D
4d <sup>2</sup> ( <sup>1</sup> F)ns			<sup>1</sup> F <sup>3</sup> F				<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H <sup>3</sup> H	
4d <sup>2</sup> ( <sup>1</sup> H)ns				<sup>1</sup> H <sup>3</sup> H				<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H° <sup>3</sup> H°	<sup>1</sup> I° <sup>3</sup> I°	
4d <sup>2</sup> 5s( <sup>1</sup> F)ns			<sup>1</sup> F <sup>3</sup> F				<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H <sup>3</sup> H	
4d <sup>2</sup> ( <sup>1</sup> S)ns	<sup>1</sup> S <sup>3</sup> S					<sup>1</sup> P° <sup>3</sup> P°			<sup>1</sup> D <sup>3</sup> D		
4d <sup>2</sup> 5s( <sup>1</sup> G)ns				<sup>1</sup> G <sup>3</sup> G				<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H <sup>3</sup> H	<sup>1</sup> I <sup>3</sup> I
4d <sup>2</sup> 5s( <sup>1</sup> P)ns	<sup>1</sup> P <sup>3</sup> P					<sup>1</sup> S° <sup>3</sup> S°	<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°		<sup>1</sup> P <sup>3</sup> P	<sup>1</sup> D <sup>3</sup> D
4d <sup>2</sup> ( <sup>3</sup> D)ns		<sup>1</sup> D <sup>3</sup> D				<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G <sup>3</sup> G		
4d <sup>2</sup> 5s( <sup>1</sup> H)ns				<sup>1</sup> H <sup>3</sup> H				<sup>1</sup> G° <sup>3</sup> G°	<sup>1</sup> H° <sup>3</sup> H°	<sup>1</sup> I° <sup>3</sup> I°	
4d <sup>2</sup> 5s( <sup>1</sup> D)ns		<sup>1</sup> D <sup>3</sup> D				<sup>1</sup> P° <sup>3</sup> P°	<sup>1</sup> D° <sup>3</sup> D°	<sup>1</sup> F° <sup>3</sup> F°	<sup>1</sup> G <sup>3</sup> G		

TABLE 2. PREDICTED TERMS OF THE Tc I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>1</sup> +	Predicted Terms			
	n <sub>3</sub> (n ≥ 5)	n <sub>2</sub> (n ≥ 5)	n <sub>1</sub> (n ≥ 5)	n <sub>0</sub> (n ≥ 5)
4s <sup>1</sup> 5s <sup>1</sup>	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>3</sup> S)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>1</sup> D)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>3</sup> P)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>1</sup> P)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>3</sup> H)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>1</sup> G)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>3</sup> G)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>1</sup> D)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>3</sup> S)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>1</sup> D)	{ {} }	{ {} }	{ {} }	{ {} }
4s <sup>1</sup> 5s <sup>1</sup> ( <sup>3</sup> S)	{ {} }	{ {} }	{ {} }	{ {} }

TABLE 3: PREDICTED TERMS OF THE Ru I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>1</sup> +	Predicted Terms		
	n <sub>s</sub> (n ≥ 5)	n <sub>p</sub> (n ≥ 5)	n <sub>d</sub> (n ≥ 5)
4s <sup>1</sup>	{ 1s <sup>2</sup> 1p <sup>1</sup> 1G		
4s <sup>1</sup> 5s <sup>1</sup>	{ 1s <sup>2</sup> 1p <sup>1</sup> 1p <sup>1</sup> 1G 1H 1s <sup>2</sup> 1p <sup>1</sup> 1p <sup>1</sup> 1G 1G 1I		
4d <sup>1</sup> (1F) <sub>nz</sub>	{ 1p <sup>1</sup> 1F	{ 1D <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup>	{ 1P 1D 1G 1H 1P 1D 1F 1G 1H
4d <sup>1</sup> (1F) <sub>nz</sub>	{ 1p <sup>1</sup>	{ 1D <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup>	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H
4d <sup>1</sup> 5s(1D) <sub>nz</sub>	{ 1D 1D	{ 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup>	{ 1P 1D 1F 1G 1G 1S 1S 1F 1G 1G
4d <sup>1</sup> (1G) <sub>nz</sub>	{ 1G 1G	{ 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup> 1H <sup>0</sup> 1H <sup>0</sup>	{ 1P 1D 1F 1G 1G 1H 1I 1P 1D 1F 1G 1G 1H 1I
4d <sup>1</sup> (1F) <sub>s</sub>	{ 1p <sup>1</sup>	{ 1D <sup>0</sup> 1D <sup>0</sup>	{ 1P 1D 1F 1G 1G 1P 1D 1F 1G 1G
4d <sup>1</sup> (1D) <sub>nz</sub>	{ 1D 1D	{ 1p <sup>1</sup> 1p <sup>1</sup> 1F <sup>0</sup> 1F <sup>0</sup>	{ 1P 1D 1F 1G 1G 1S 1S 1F 1G 1G
4d <sup>1</sup> (1H) <sub>nz</sub>	{ 1H 1H	{ 1D <sup>0</sup> 1D <sup>0</sup> 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup> 1H <sup>0</sup> 1H <sup>0</sup>	{ 1P 1D 1F 1G 1G 1H 1I 1K 1S 1S 1F 1G 1G 1H 1I 1K
4d <sup>1</sup> 5s(1D) <sub>nz</sub>	{ 1D 1D	{ 1p <sup>1</sup> 1p <sup>1</sup> 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup>	{ 1P 1D 1F 1G 1G 1S 1S 1F 1G 1G
4d <sup>1</sup> (1F) <sub>nz</sub>	{ 1F 1F	{ 1D <sup>0</sup> 1D <sup>0</sup> 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup>	{ 1P 1D 1F 1G 1G 1S 1S 1F 1G 1G
4d <sup>1</sup> 5s(1F) <sub>nz</sub>	{ 1F 1H	{ 1D <sup>0</sup> 1D <sup>0</sup> 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup> 1H <sup>0</sup> 1H <sup>0</sup>	{ 1P 1D 1F 1G 1G 1H 1I 1K 1S 1S 1F 1G 1G 1H 1I 1K
4d <sup>1</sup> 5s(1F) <sub>nz</sub>	{ 1F 1F	{ 1D <sup>0</sup> 1D <sup>0</sup> 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup>	{ 1P 1D 1F 1G 1G 1S 1S 1F 1G 1G
4d <sup>1</sup> 5s(1G) <sub>nz</sub>	{ 1G 1G	{ 1D <sup>0</sup> 1D <sup>0</sup> 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup> 1H <sup>0</sup> 1H <sup>0</sup>	{ 1P 1D 1F 1G 1G 1H 1I 1K 1S 1S 1F 1G 1G 1H 1I 1K
4d <sup>1</sup> 5s(1H) <sub>nz</sub>	{ 1H 1H	{ 1D <sup>0</sup> 1D <sup>0</sup> 1F <sup>0</sup> 1F <sup>0</sup> 1G <sup>0</sup> 1H <sup>0</sup> 1H <sup>0</sup>	{ 1P 1D 1F 1G 1G 1H 1I 1K 1S 1S 1F 1G 1G 1H 1I 1K

TABLE 4. PREDICTED TERMS OF THE R<sub>h</sub> I ISOMERIZATION SEQUENCE

Configuration 1 <sup>h</sup> 2 <sup>h</sup> 3 <sup>h</sup> 4 <sup>h</sup> 3 <sup>h</sup> 2 <sup>h</sup> 4 <sup>h</sup> 1 <sup>h</sup> +	Predicted Terms			
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)	----
4 <sup>h</sup>	1D			
4 <sup>h</sup> 5 <sup>h</sup>	1P 1D 1D	1F 1G 1H		
4 <sup>h</sup> (1F) ns	1P 1P	1D <sup>o</sup> 1D <sup>o</sup>	1F <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1G <sup>o</sup>	1P 1D 1F 1G 1H 1H
4 <sup>h</sup> (1P) ns	1P 1P	1D <sup>o</sup> 1D <sup>o</sup>	1F <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1G <sup>o</sup>	1P 1D 1F 1G 1H 1H
4 <sup>h</sup> (1G) ns			1F <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>	1P 1D 1F 1G 1H 1H
4 <sup>h</sup> (1D) ns	1D	1P <sup>o</sup> 1P <sup>o</sup>	1F <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1G <sup>o</sup>	1P 1D 1F 1G 1H 1H
4 <sup>h</sup> (1S) ns	1S	1P <sup>o</sup> 1P <sup>o</sup>	1F <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1G <sup>o</sup>	1P 1D 1F 1G 1H 1H
4 <sup>h</sup> 5 <sup>h</sup> (1F) ns	1P 1P	1D <sup>o</sup> 1D <sup>o</sup>	1F <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1G <sup>o</sup>	1P 1D 1F 1G 1H 1H
4 <sup>h</sup> 5 <sup>h</sup> (1P) ns	1P 1P	1D <sup>o</sup> 1D <sup>o</sup>	1F <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1G <sup>o</sup>	1P 1D 1F 1G 1H 1H

TABLE 5. PREDICTED TERMS OF THE Pd I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>0</sup> +	Predicted Terms										
4d <sup>9</sup>	1S										
4d <sup>8</sup> 5s	{ 1S 1P 1D 1F 1G										
	nd (n ≥ 5)					np (n ≥ 5)					
4d <sup>9</sup> (1D)ns	{ 1D 1D					1P° 1D° 1F° 1P° 1D° 1F°					
4d <sup>8</sup> 5s(1F)ns'	{ 1F 1F					1D° 1F° 1G° 1D° 1F° 1G°					
4d <sup>8</sup> 5s(3F)ns''	{ 1F 1F					1D° 1F° 1G° 1D° 1F° 1G°					
	nd (n ≥ 5)					nf (n ≥ 4)					-----
4d <sup>9</sup> (1D)ns	{ 1S 1P 1D 1F 1G 1S 1P 1D 1F 1G					1P° 1D° 1F° 1G° 1H° 1P° 1D° 1F° 1G° 1H°					-----
4d <sup>8</sup> 5s(1F)ns'	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H					1S° 1P° 1D° 1F° 1G° 1H° 1I° 1S° 1P° 1D° 1F° 1G° 1H° 1I°					-----
4d <sup>8</sup> 5s(3F)ns''	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H					1S° 1P° 1D° 1F° 1G° 1H° 1I° 1S° 1P° 1D° 1F° 1G° 1H° 1I°					-----

TABLE 6. PREDICTED TERMS OF THE Ag I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>1</sup> 4p <sup>0</sup> +	Predicted Terms										
4d <sup>9</sup> 5s	1D										
	nd (n ≥ 5)			np (n ≥ 5)			nd (n ≥ 5)				
4d <sup>9</sup> (1S)ns	1S										
4d <sup>8</sup> 5s(1D)ns'	{ 1D 1D			1P° 1D° 1F° 1P° 1D° 1F°			1S 1P 1D 1F 1G 1S 1P 1D 1F 1G				
4d <sup>8</sup> 5s(3D)ns''	{ 1D			1P° 1D° 1F°			1S 1P 1D 1F 1G				
	nf (n ≥ 4)			ng (n ≥ 5)			-----				
4d <sup>9</sup> (1S)ns	1F°			1G			-----				
4d <sup>8</sup> 5s(1D)ns'	{ 1P° 1D° 1F° 1G° 1H° 1P° 1D° 1F° 1G° 1H°			1D 1F 1G 1H 1I 1D 1F 1G 1H 1I			-----				
4d <sup>8</sup> 5s(3D)ns''	{ 1P° 1D° 1F° 1G° 1H°			1D 1F 1G 1H 1I			-----				

TABLE 7. PREDICTED TERMS OF THE Cd : ISOLELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 3s <sup>2</sup> 3p <sup>4</sup> 4s <sup>2</sup> 4p <sup>1</sup> 4d <sup>0</sup> +	Predicted Terms		
4s <sup>2</sup> 5s <sup>2</sup>	1S		
{ 4s <sup>2</sup> 5p <sup>2</sup>	{ 1P 1D		
{ 4s <sup>2</sup> 5d <sup>2</sup>	{ 1P 1D 1F 1G		
	ns (n ≥ 5)	np (n ≥ 5)	nd (n ≥ 5)
{ 4s <sup>2</sup> 5s(1S)ns	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1P <sup>o</sup> 1D 1D	{ 1D 1D
{ 4s <sup>2</sup> 5p(1P <sup>o</sup> )ns'	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1S 1P 1D 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1P <sup>o</sup>
{ 4s <sup>2</sup> 5d(1D)ns''	{ 1D 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1P <sup>o</sup>	{ 1S 1P 1D 1F 1G
	n' (n ≥ 4)	ng (n ≥ 5)	-----
{ 4s <sup>2</sup> 5s(1S)ns	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1G 1G	-----
{ 4s <sup>2</sup> 5p(1P <sup>o</sup> )ns'	{ 1D 1F 1G 1D 1F 1G	{ 1P <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1P <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>	-----
{ 4s <sup>2</sup> 5d(1D)ns''	{ 1P <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1P <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>	{ 1D 1F 1G 1H 1I 1D 1F 1G 1H 1I	-----

TABLE 8. PREDICTED TERMS OF THE In : ISOLELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>2</sup> 4s <sup>2</sup> 4p <sup>1</sup> 4d <sup>0</sup> +	Predicted Terms			
5s <sup>2</sup> (1S) 5p	1P <sup>o</sup>			
{ 5s 5p <sup>2</sup>	{ 1S 1P 1D			
{ 5p <sup>2</sup>	{ 1P <sup>o</sup> 1D <sup>o</sup>			
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)	nf (n ≥ 6)
{ 5s <sup>2</sup> (1S) ns	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1P <sup>o</sup> 1D 1D	{ 1D 1D 1D 1D	{ 1P <sup>o</sup> 1G 1G
{ 5s 5p(1P <sup>o</sup> ) ns'	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1S 1P 1P 1D 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1D <sup>o</sup>	{ 1P <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1P <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>
{ 5s 5p(1P <sup>o</sup> ) ns''	{ 1P <sup>o</sup>	{ 1S 1P 1D 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1D <sup>o</sup>	{ 1P <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1P <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>

TABLE 9. PREDICTED TERMS OF THE  $5d^1$  ISOELECTRONIC SEQUENCE

Configuration $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 4p^6 4d^{n-1}$	Predicted Terms			
	$ns (n \geq 6)$	$np (n \geq 6)$	$nd (n \geq 5)$	$nf (n \geq 4)$
$5d^1 5p^1$	${}^1S$	${}^1D$		
$5s 5p^1$	${}^1S^o$ ${}^3S^o$	${}^1P^o$ ${}^3P^o$	${}^1D^o$ ${}^3D^o$	${}^1G^o$ ${}^3G^o$
$5s^2 5p^1(1P^o)ns$	${}^1P^o$ ${}^3P^o$	${}^1S$ ${}^3S$	${}^1D$ ${}^3D$	${}^1G$ ${}^3G$
$5s 5p^1(1P^o)ns'$	${}^1P$ ${}^3P$	${}^1S$ ${}^3S$	${}^1D$ ${}^3D$	${}^1G$ ${}^3G$

TABLE 10. PREDICTED TERMS OF THE  $5d^1$  ISOELECTRONIC SEQUENCE

Configuration $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 4p^6 4d^{n-1}$	Predicted Terms			
	$ns (n \geq 6)$	$np (n \geq 6)$	$nd (n \geq 5)$	$nf (n \geq 4)$
$5s^2 5p^1$	${}^1S^o$	${}^1P^o$ ${}^3P^o$	${}^1D^o$	
$5s 5p^1$	${}^1S$	${}^1P$ ${}^3P$	${}^1D$	
$5p^1$		${}^1P^o$		
$5s^2 5p^1(1P^o)ns$	${}^1P^o$ ${}^3P^o$	${}^1S^o$ ${}^3S^o$	${}^1D^o$ ${}^3D^o$	${}^1G^o$ ${}^3G^o$
$5s^2 5p^1(1D^o)ns'$		${}^1P^o$ ${}^3P^o$	${}^1D^o$ ${}^3D^o$	${}^1G^o$ ${}^3G^o$
$5s^2 5p^1(1S)ns''$	${}^1S$		${}^1D$	${}^1F$



TABLE 11. PREDICTED TERMS OF THE T<sub>61</sub> ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>n+1</sup>	Predicted Terms			
	ns (n ≥ 0)	np (n ≥ 0)	nd (n ≥ 0)	nf (n ≥ 0)
5s <sup>2</sup> 5p <sup>4</sup>	{ <sup>1</sup> S	<sup>3</sup> P	<sup>1</sup> D	
5s 5p <sup>5</sup>	{	<sup>3</sup> P <sup>o</sup> <sup>1</sup> P <sup>o</sup>		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>3</sup> P) <sup>ns</sup>	{ <sup>3</sup> S <sup>o</sup> <sup>1</sup> S <sup>o</sup>	<sup>3</sup> P <sup>1</sup> P	<sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>	<sup>3</sup> F <sup>1</sup> F
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>1</sup> D) <sup>ns'</sup>	{	<sup>3</sup> P <sup>1</sup> P	<sup>3</sup> S <sup>o</sup> <sup>1</sup> S <sup>o</sup>	<sup>3</sup> F <sup>1</sup> F
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>3</sup> P) <sup>ns''</sup>	{	<sup>3</sup> P <sup>1</sup> P	<sup>3</sup> S <sup>1</sup> S	<sup>3</sup> F <sup>1</sup> F

TABLE 12. PREDICTED TERMS OF THE I<sub>1</sub> ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>n+1</sup>	Predicted Terms			
	ns (n ≥ 0)	np (n ≥ 0)	nd (n ≥ 0)	nf (n ≥ 0)
5s <sup>2</sup> 5p <sup>4</sup>	<sup>3</sup> P <sup>o</sup>			
5s 5p <sup>5</sup>	<sup>3</sup> S			
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>3</sup> P) <sup>ns</sup>	{ <sup>3</sup> S <sup>o</sup> <sup>1</sup> S <sup>o</sup>	<sup>3</sup> P <sup>o</sup> <sup>1</sup> P <sup>o</sup>	<sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>	<sup>3</sup> F <sup>o</sup> <sup>1</sup> F <sup>o</sup>
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>1</sup> D) <sup>ns'</sup>	<sup>3</sup> S	<sup>3</sup> P <sup>1</sup> P	<sup>1</sup> D <sup>1</sup> D	<sup>3</sup> F <sup>1</sup> F
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>3</sup> P) <sup>ns''</sup>	{ <sup>3</sup> S <sup>1</sup> S	<sup>3</sup> P <sup>1</sup> P	<sup>1</sup> D <sup>1</sup> D	<sup>3</sup> F <sup>1</sup> F

TABLE 13. PREDICTED LEVELS OF THE Xs: ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>n+1</sup>	Predicted Terms				
5s <sup>2</sup> 5p <sup>2</sup>	1S				
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 5)	nf (n ≥ 4)	-----
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>3</sup> P <sup>o</sup> )ns	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1S 1P 1D 1S 1P 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup>	{ 1D 1F 1G 1D 1F 1G	-----
5s 5p <sup>4</sup> ( <sup>3</sup> S)ns'	{ 1S 1S	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1D 1D	{ 1F <sup>o</sup> 1F <sup>o</sup>	-----
<i>Jl</i> -Coupling Notation					
Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>n+1</sup>	Predicted Pairs				
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 5)	nf (n ≥ 4)	-----
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>3</sup> P <sub>2,0</sub> )ns	{ [1/2] <sup>o</sup>	{ [0/2] [2/2] [1/2]	{ [0/2] <sup>o</sup> [3/2] [2/2] [1/2] [2/2] <sup>o</sup>	{ [1/2] [4/2] [2/2] [3/2]	-----
5s 5p <sup>4</sup> ( <sup>3</sup> P <sub>2,0</sub> )ns'	{ [0/2] <sup>o</sup>	{ [1/2] [0/2]	{ [2/2] <sup>o</sup> [1/2] <sup>o</sup>	{ [3/2] [2/2]	-----

TABLE 14. PREDICTED TERMS OF THE Ba I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>n</sup> 5s <sup>2</sup> 5p <sup>2</sup> +	Predicted Terms				
6s <sup>2</sup>	1S				
5d <sup>2</sup>	{ 1S 1P 1D 1F 1G				
6p <sup>2</sup>	{ 1S 1P 1D				
4f <sup>2</sup>	{ 1S 1P 1L 1F 1G 1H 1I				
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 5)		
6s( <sup>3</sup> S)ns	{ 1S 1S	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1D 1D		
5d( <sup>3</sup> D)ns'	{ 1D 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup>	{ 1S 1P 1D 1F 1G 1S 1P 1D 1F 1G	-----	
6p( <sup>3</sup> P <sup>o</sup> )ns''	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1S 1P 1D 1S 1P 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup>	-----	
	nf (n ≥ 4)	ng (n ≥ 5)	-----		
6s( <sup>3</sup> S)ns	{ 1P <sup>o</sup> 1P <sup>o</sup>	{ 1G 1G	-----		
5d( <sup>3</sup> D)ns'	{ 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>	{ 1D 1F 1G 1H 1I 1D 1F 1G 1H 1I	-----		
6p( <sup>3</sup> P <sup>o</sup> )ns''	{ 1D 1F 1G 1D 1F 1G	{ 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>	-----		

TABLE 15. PREDICTED TERMS OF THE  $La$  ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>n</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Predicted Terms						
3d 6s <sup>2</sup>	1D						
5d <sup>2</sup>	$\left\{ \begin{array}{l} {}^1P \\ {}^3P \\ {}^1D \\ {}^3D \end{array} \right. \begin{array}{l} {}^3F \\ {}^1F \\ {}^3G \\ {}^1H \end{array}$						
	$ns$ ( $n \geq 6$ )			$np$ ( $n \geq 6$ )			
5d <sup>2</sup> ( <sup>3</sup> F) <sub>nz</sub>	$\left\{ \begin{array}{l} {}^3F \\ {}^1F \end{array} \right.$			$\left\{ \begin{array}{l} {}^3D^{\circ} \\ {}^1D^{\circ} \end{array} \right. \begin{array}{l} {}^3F^{\circ} \\ {}^1F^{\circ} \end{array} \begin{array}{l} {}^3G^{\circ} \\ {}^1G^{\circ} \end{array}$			
5d 6s( <sup>1</sup> D) <sub>nz</sub>	1D			1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup>			
5d 6s( <sup>3</sup> D) <sub>nz</sub>	$\left\{ \begin{array}{l} {}^3D \\ {}^1D \end{array} \right.$			$\left\{ \begin{array}{l} {}^3P^{\circ} \\ {}^1P^{\circ} \end{array} \right. \begin{array}{l} {}^3D^{\circ} \\ {}^1D^{\circ} \end{array} \begin{array}{l} {}^3F^{\circ} \\ {}^1F^{\circ} \end{array}$			
5d <sup>2</sup> ( <sup>3</sup> P) <sub>nz</sub>	$\left\{ \begin{array}{l} {}^3P \\ {}^1P \end{array} \right.$			$\left\{ \begin{array}{l} {}^3S^{\circ} \\ {}^1S^{\circ} \end{array} \right. \begin{array}{l} {}^3P^{\circ} \\ {}^1P^{\circ} \end{array} \begin{array}{l} {}^3D^{\circ} \\ {}^1D^{\circ} \end{array}$			
6s <sup>2</sup> ( <sup>1</sup> S) <sub>nz</sub>	1S			1P <sup>o</sup>			
5d <sup>2</sup> ( <sup>1</sup> G) <sub>nz</sub>	1G			1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>			
5d <sup>2</sup> ( <sup>1</sup> D) <sub>nz</sub>	1D			1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup>			
	$nd$ ( $n \geq 5$ )			$nf$ ( $n \geq 4$ )			-----
5d <sup>2</sup> ( <sup>3</sup> F) <sub>nz</sub>	$\left\{ \begin{array}{l} {}^3P \\ {}^1P \end{array} \right. \begin{array}{l} {}^3D \\ {}^1D \end{array} \begin{array}{l} {}^3F \\ {}^1F \end{array} \begin{array}{l} {}^3G \\ {}^1G \end{array} \begin{array}{l} {}^3H \\ {}^1H \end{array}$			$\left\{ \begin{array}{l} {}^3S^{\circ} \\ {}^1S^{\circ} \end{array} \right. \begin{array}{l} {}^3P^{\circ} \\ {}^1P^{\circ} \end{array} \begin{array}{l} {}^3D^{\circ} \\ {}^1D^{\circ} \end{array} \begin{array}{l} {}^3F^{\circ} \\ {}^1F^{\circ} \end{array} \begin{array}{l} {}^3G^{\circ} \\ {}^1G^{\circ} \end{array} \begin{array}{l} {}^3H^{\circ} \\ {}^1H^{\circ} \end{array} \begin{array}{l} {}^3I^{\circ} \\ {}^1I^{\circ} \end{array}$			-----
5d 6s( <sup>1</sup> D) <sub>nz</sub>	1S 1P 1D 1F 1G			1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>			-----
5d 6s( <sup>3</sup> D) <sub>nz</sub>	$\left\{ \begin{array}{l} {}^3S \\ {}^1S \end{array} \right. \begin{array}{l} {}^3P \\ {}^1P \end{array} \begin{array}{l} {}^3D \\ {}^1D \end{array} \begin{array}{l} {}^3F \\ {}^1F \end{array} \begin{array}{l} {}^3G \\ {}^1G \end{array}$			$\left\{ \begin{array}{l} {}^3P^{\circ} \\ {}^1P^{\circ} \end{array} \right. \begin{array}{l} {}^3D^{\circ} \\ {}^1D^{\circ} \end{array} \begin{array}{l} {}^3F^{\circ} \\ {}^1F^{\circ} \end{array} \begin{array}{l} {}^3G^{\circ} \\ {}^1G^{\circ} \end{array} \begin{array}{l} {}^3H^{\circ} \\ {}^1H^{\circ} \end{array}$			-----
5d <sup>2</sup> ( <sup>3</sup> P) <sub>nz</sub>	$\left\{ \begin{array}{l} {}^3P \\ {}^1P \end{array} \right. \begin{array}{l} {}^3D \\ {}^1D \end{array} \begin{array}{l} {}^3F \\ {}^1F \end{array}$			$\left\{ \begin{array}{l} {}^3D^{\circ} \\ {}^1D^{\circ} \end{array} \right. \begin{array}{l} {}^3F^{\circ} \\ {}^1F^{\circ} \end{array} \begin{array}{l} {}^3G^{\circ} \\ {}^1G^{\circ} \end{array}$			-----
6s <sup>2</sup> ( <sup>1</sup> S) <sub>nz</sub>	1D			1F <sup>o</sup>			-----
5d <sup>2</sup> ( <sup>1</sup> G) <sub>nz</sub>	1D 1F 1G 1H 1I			1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1I <sup>o</sup> 1K <sup>o</sup>			-----
5d <sup>2</sup> ( <sup>1</sup> D) <sub>nz</sub>	1S 1P 1D 1F 1G			1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>			-----

TABLE 16. PREDICTED TERMS OF THE H I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2p <sup>2</sup> 3p <sup>2</sup> 3p <sup>2</sup> 3d <sup>2</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>2</sup> 4f <sup>2</sup> 5s <sup>2</sup> 5p <sup>2</sup> +	Predicted Terms		
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)
5s <sup>2</sup> 6s <sup>2</sup>	{ 1S 1P 1D 1G 1D 1D 1F 1G 1H 1P 1P 1D 1F 1G 1H 1S 1S		
5s <sup>2</sup> 6s(1F)nz	{ 1P 1F 1P 1F	{ 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup>	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H
5s <sup>2</sup> 6s(1P)nz	{ 1P 1P	{ 1S <sup>o</sup> 1P <sup>o</sup> 1S <sup>o</sup> 1P <sup>o</sup>	{ 1P 1D 1F 1P 1D 1F
5s <sup>2</sup> 6s(1F)nz	{ 1D 1D	{ 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup>	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H
5s <sup>2</sup> 6s(1D)nz	{ 1P 1P	{ 1S <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1S <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup>	{ 1P 1D 1F 1P 1D 1F
5s <sup>2</sup> 6s(1P)nz	{ 1G 1G	{ 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>	{ 1D 1D 1F 1G 1H 1I 1D 1D 1F 1G 1H 1I
5s <sup>2</sup> 6s(1G)nz	{ 1F 1F	{ 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup>	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H
5s <sup>2</sup> (1F)nz	{ 1P 1P	{ 1S <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1S <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup>	{ 1P 1D 1F 1P 1D 1F
5s <sup>2</sup> (1P)nz	{ 1G 1G	{ 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1H <sup>o</sup>	{ 1D 1D 1F 1G 1H 1I 1D 1D 1F 1G 1H 1I
5s <sup>2</sup> (1G)nz	{ 1D 1D	{ 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup>	{ 1S 1P 1D 1F 1G 1S 1P 1D 1F 1G
5s <sup>2</sup> (1H)nz	{ 1P 1P	{ 1S <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup> 1S <sup>o</sup> 1P <sup>o</sup> 1D <sup>o</sup>	{ 1P 1D 1F 1G 1H 1I 1K 1P 1D 1F 1G 1H 1I 1K
5s <sup>2</sup> (1P)nz	{ 1F 1F	{ 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup> 1D <sup>o</sup> 1F <sup>o</sup> 1G <sup>o</sup>	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H

TABLE 17. PREDICTED TERMS OF THE Ta I ISOLECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>n</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Predicted Terms		
	$n s \ (n \geq 6)$	$n p \ (n \geq 6)$	$n d \ (n \geq 6)$
5d <sup>2</sup> 6s <sup>2</sup>	{ 4P 3D 3F 3G 3H 4S 4P 4D 4F 4G 4H 4I 4S 4P 4D 4F 4G 4H 4I		
5d <sup>2</sup> 6s(4F)nz	{ 4F 4F 4F 3F	{ 4D <sup>o</sup> 4F <sup>o</sup> 4G <sup>o</sup> 4D <sup>o</sup> 4F <sup>o</sup> 4G <sup>o</sup>	{ 4P 4D 4F 4G 4H 4P 4D 4F 4G 4H
5d <sup>2</sup> 6s(4F)nz	{ 4P 3P	{ 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup> 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup>	{ 4P 4D 4F 4G 4H 4P 4D 4F 4G 4H
5d <sup>2</sup> 6s(4F)nz	{ 4P 3P	{ 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup> 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup>	{ 4P 4D 4F 4G 4H 4P 4D 4F 4G 4H
5d <sup>2</sup> 6s(4F)nz	{ 4P 3P	{ 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup> 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup>	{ 4P 4D 4F 4G 4H 4P 4D 4F 4G 4H
5d <sup>2</sup> 6s(4G)nz	{ 4G 3G	{ 4F <sup>o</sup> 4G <sup>o</sup> 4H <sup>o</sup> 4F <sup>o</sup> 4G <sup>o</sup> 4H <sup>o</sup>	{ 4D 4F 4G 4H 4I 4D 4F 4G 4H 4I
5d <sup>2</sup> 6s(4D)nz	{ 4D 4D	{ 4P <sup>o</sup> 4D <sup>o</sup> 4F <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup> 4F <sup>o</sup>	{ 4S 4P 4D 4F 4G 4S 4P 4D 4F 4G
5d <sup>2</sup> 6s(4D)nz	{ 4D 3D	{ 4P <sup>o</sup> 4D <sup>o</sup> 4F <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup> 4F <sup>o</sup>	{ 4S 4P 4D 4F 4G 4S 4P 4D 4F 4G
5d <sup>2</sup> 6s(4H)nz	{ 4H 3H	{ 4G <sup>o</sup> 4H <sup>o</sup> 4I <sup>o</sup> 4G <sup>o</sup> 4H <sup>o</sup> 4I <sup>o</sup>	{ 4F 4G 4H 4I 4K 4F 4G 4H 4I 4K
5d <sup>2</sup> 6s(4P)nz	{ 4P 3P	{ 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup> 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup>	{ 4P 4D 4F 4G 4P 4D 4F 4G
5d <sup>2</sup> 6s(4D)nz	{ 4D 3D	{ 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup> 4S <sup>o</sup> 4P <sup>o</sup> 4D <sup>o</sup>	{ 4S 4P 4D 4F 4G 4S 4P 4D 4F 4G
5d <sup>2</sup> 6s(4F)nz	{ 4F 3F	{ 4G <sup>o</sup> 4H <sup>o</sup> 4I <sup>o</sup> 4G <sup>o</sup> 4H <sup>o</sup> 4I <sup>o</sup>	{ 4P 4D 4F 4G 4H 4I 4K 4P 4D 4F 4G 4H 4I 4K
5d <sup>2</sup> 6s(4G)nz	{ 4G 3G	{ 4F <sup>o</sup> 4G <sup>o</sup> 4H <sup>o</sup> 4F <sup>o</sup> 4G <sup>o</sup> 4H <sup>o</sup>	{ 4P 4D 4F 4G 4H 4I 4P 4D 4F 4G 4H 4I
5d <sup>2</sup> 6s(4F)nz	{ 4F 3F	{ 4D <sup>o</sup> 4F <sup>o</sup> 4G <sup>o</sup> 4D <sup>o</sup> 4F <sup>o</sup> 4G <sup>o</sup>	{ 4P 4D 4F 4G 4H 4I 4P 4D 4F 4G 4H 4I



TABLE 10. PREDICTED TERMS OF THE R<sub>1</sub> ISOELECTRONIC SEQUENCES

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>n</sup> 4f <sup>n</sup> 5s <sup>2</sup> 5p <sup>6</sup> 4	Predicted Terms		
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)
5d <sup>1</sup> 6s <sup>2</sup>	$\left\{ \begin{array}{l} 4^1P \\ 4^1D \\ 4^1D \\ 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1G \\ 4^1G \\ 4^1G \\ 4^1G \\ 4^1G \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1I \\ 4^1H \\ 4^1H \\ 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>2</sup>	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \\ 4^1D \\ 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1F \\ 4^1F \\ 4^1G \\ 4^1G \\ 4^1G \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \\ 4^1H \\ 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> 6s (5S) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1S \\ 4^1S \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$
5d <sup>3</sup> (1D) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1G \\ 4^1G \end{array} \right\}$
5d <sup>3</sup> (3P) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1F \\ 4^1F \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1G \\ 4^1G \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> (3H) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1G \\ 4^1G \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> (3G) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> (3P) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1S \\ 4^1S \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> (1D) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1G \\ 4^1G \end{array} \right\}$
5d <sup>3</sup> 6s <sup>2</sup> (1D) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1S \\ 4^1S \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> 6s (5S) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1S \\ 4^1S \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> 6s (3G) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> 6s (1P) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$
5d <sup>3</sup> 6s (1F) <sub>nz</sub>	$\left\{ \begin{array}{l} 4^1P \\ 4^1P \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1D \\ 4^1D \end{array} \right\}$	$\left\{ \begin{array}{l} 4^1H \\ 4^1H \end{array} \right\}$

TABLE 20. PREDICTED TERMS OF THE Os I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>5</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>5</sup> 4f <sup>0</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Predicted Terms		
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)
5d <sup>5</sup> 6s <sup>1</sup>	$\left\{ \begin{array}{l} {}^1D \\ {}^3P \\ {}^1S \end{array} \right\} \begin{array}{l} {}^1G \\ {}^3G \\ {}^1G \end{array} \begin{array}{l} {}^3H \\ {}^1I \end{array}$ $\left\{ \begin{array}{l} {}^3P \\ {}^1D \end{array} \right\} {}^1G$		
5d <sup>5</sup> 6s <sup>0</sup>			
5d <sup>5</sup> 6s <sup>0</sup> (D) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^1D \\ {}^3D \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1P^{\circ} \\ {}^3P^{\circ} \end{array} \right\} \begin{array}{l} {}^1D^{\circ} \\ {}^3D^{\circ} \end{array} \begin{array}{l} {}^1F^{\circ} \\ {}^3F^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array}$
5d <sup>5</sup> (F) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3P \\ {}^1P \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1D^{\circ} \\ {}^3D^{\circ} \end{array} \right\} \begin{array}{l} {}^1G^{\circ} \\ {}^3G^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array} \begin{array}{l} {}^1H \\ {}^3H \end{array}$
5d <sup>5</sup> (P) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3P \\ {}^1P \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1S^{\circ} \\ {}^3S^{\circ} \end{array} \right\} \begin{array}{l} {}^1D^{\circ} \\ {}^3D^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array} \begin{array}{l} {}^1H \\ {}^3H \end{array}$
5d <sup>5</sup> (H) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3H \\ {}^1H \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1G^{\circ} \\ {}^3G^{\circ} \end{array} \right\} \begin{array}{l} {}^1H^{\circ} \\ {}^3H^{\circ} \end{array} \begin{array}{l} {}^1I^{\circ} \\ {}^3I^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array} \begin{array}{l} {}^1H \\ {}^3H \end{array} \begin{array}{l} {}^1I \\ {}^3I \end{array} \begin{array}{l} {}^1K \\ {}^3K \end{array}$
5d <sup>5</sup> (G) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3G \\ {}^1G \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1P^{\circ} \\ {}^3P^{\circ} \end{array} \right\} \begin{array}{l} {}^1G^{\circ} \\ {}^3G^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array} \begin{array}{l} {}^1H \\ {}^3H \end{array}$
5d <sup>5</sup> (F) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3F \\ {}^1F \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1D^{\circ} \\ {}^3D^{\circ} \end{array} \right\} \begin{array}{l} {}^1G^{\circ} \\ {}^3G^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array} \begin{array}{l} {}^1H \\ {}^3H \end{array}$
5d <sup>5</sup> (D) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3D \\ {}^1D \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1P^{\circ} \\ {}^3P^{\circ} \end{array} \right\} \begin{array}{l} {}^1D^{\circ} \\ {}^3D^{\circ} \end{array} \begin{array}{l} {}^1F^{\circ} \\ {}^3F^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array}$
5d <sup>5</sup> (P) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3P \\ {}^1P \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1S^{\circ} \\ {}^3S^{\circ} \end{array} \right\} \begin{array}{l} {}^1D^{\circ} \\ {}^3D^{\circ} \end{array} \begin{array}{l} {}^1F^{\circ} \\ {}^3F^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array}$
5d <sup>5</sup> 6s <sup>0</sup> (D) <sub>ns</sub>	$\left\{ \begin{array}{l} {}^3D \\ {}^1D \end{array} \right\}$	$\left\{ \begin{array}{l} {}^1P^{\circ} \\ {}^3P^{\circ} \end{array} \right\} \begin{array}{l} {}^1D^{\circ} \\ {}^3D^{\circ} \end{array} \begin{array}{l} {}^1F^{\circ} \\ {}^3F^{\circ} \end{array}$	$\left\{ \begin{array}{l} {}^1S \\ {}^3S \end{array} \right\} \begin{array}{l} {}^1D \\ {}^3D \end{array} \begin{array}{l} {}^1F \\ {}^3F \end{array} \begin{array}{l} {}^1G \\ {}^3G \end{array}$



TABLE 21. PREDICTED TERMS OF THE Ir I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>8</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4f <sup>n</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Predicted Terms				
5d <sup>8</sup> 6s <sup>2</sup>	{ 1P 1D 1D	1F 1F 1G 1H			
5d <sup>8</sup>		1D			
		ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)	-----
5d <sup>8</sup> ( <sup>2</sup> F)nz	{ 1F 1F		1D° 1F° 1G° 1D° 1F° 1G°	1P 1D 1F 1G 1H 1P 1D 1F 1G 1H	----- -----
5d <sup>8</sup> ( <sup>2</sup> P)nz	{ 1P 1P		1S° 1P° 1D° 1S° 1P° 1D°	1P 1D 1F 1P 1D 1F	----- -----
5d <sup>8</sup> ( <sup>2</sup> G)nz		1G		1D 1F 1G 1H 1I	-----
5d <sup>8</sup> ( <sup>2</sup> D)nz		1D	1P° 1D° 1F°	1S 1P 1D 1F 1G	-----
5d <sup>7</sup> 6s( <sup>2</sup> F)nz	{ 1F 1F		1D° 1F° 1G° 1D° 1F° 1G°	1P 1D 1F 1G 1H 1P 1D 1F 1G 1H	----- -----
5d <sup>7</sup> 6s( <sup>2</sup> P)nz	{ 1P 1P		1S° 1P° 1D° 1S° 1P° 1D°	1P 1D 1F 1P 1D 1F	----- -----
5d <sup>7</sup> 6s( <sup>2</sup> F)nz	{ 1F 1F		1D° 1F° 1G° 1D° 1F° 1G°	1P 1D 1F 1G 1H 1P 1D 1F 1G 1H	----- -----
5d <sup>7</sup> 6s( <sup>2</sup> P)nz	{ 1P 1P		1S° 1P° 1D° 1S° 1P° 1D°	1P 1D 1F 1P 1D 1F	----- -----

TABLE 22. PREDICTED TERMS OF THE Pt I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>n</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Predicted Terms			
5d <sup>9</sup> 6s <sup>2</sup>	{ 1S 1P 1D 1F 1G			
5d <sup>9</sup>		1S		
		ns (n ≥ 6)	np (n ≥ 6)	-----
5d <sup>9</sup> ( <sup>2</sup> D)nz	{ 1D 1D		1P° 1D° 1F° 1P° 1D° 1F°	
5d <sup>9</sup> 6s( <sup>2</sup> F)nz'	{ 1F 1F		1D° 1F° 1G° 1D° 1F° 1G°	
5d <sup>9</sup> 6s( <sup>2</sup> P)nz''	{ 1P 1P		1S° 1P° 1D° 1S° 1P° 1D°	
5d <sup>9</sup> 6s( <sup>2</sup> F)nz'''	{ 1F 1F		1D° 1F° 1G° 1D° 1F° 1G°	
		nd (n ≥ 6)	nf (n ≥ 4)	-----
5d <sup>9</sup> ( <sup>2</sup> D)nz	{ 1S 1P 1D 1F 1G 1S 1P 1D 1F 1G		1P° 1D° 1F° 1G° 1H° 1P° 1D° 1F° 1G° 1H°	----- -----
5d <sup>9</sup> 6s( <sup>2</sup> F)nz'	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H		1S° 1P° 1D° 1F° 1G° 1H° 1I° 1S° 1P° 1D° 1F° 1G° 1H° 1I°	----- -----
5d <sup>9</sup> 6s( <sup>2</sup> P)nz''	{ 1P 1D 1F 1P 1D 1F		1D° 1F° 1G° 1D° 1F° 1G°	----- -----
5d <sup>9</sup> 6s( <sup>2</sup> F)nz'''	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H		1S° 1P° 1D° 1F° 1G° 1H° 1I° 1S° 1P° 1D° 1F° 1G° 1H° 1I°	----- -----

TABLE 23. PREDICTED TERMS OF THE Au I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Predicted Terms										
	5d <sup>9</sup> 6s <sup>2</sup>	1D									
	ns (n ≥ 6)			np (n ≥ 6)			nd (n ≥ 6)				
5d <sup>10</sup> (1S)ns	1S			1P°			1D				
5d <sup>9</sup> 6s(1D)ns'	{	1D	1P°	1D°	1F°	1S	1P	1D	1F	1G	
5d <sup>9</sup> 6s(1D)ns''		1D	1P°	1D°	1F°	1S	1P	1D	1F	1G	
	nf (n ≥ 5)					ng (n ≥ 5)					
5d <sup>10</sup> (1S)ns	1F°					1G					
5d <sup>9</sup> 6s(1D)ns'	{	1P°	1D°	1F°	1G°	1H°	1D	1F	1G	1H	1I
5d <sup>9</sup> 6s(1D)ns''		1P°	1D°	1F°	1G°	1H°	1D	1F	1G	1H	1I

TABLE 24. PREDICTED TERMS OF THE Hg I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Predicted Terms										
	5d <sup>10</sup> 6s <sup>2</sup>	1S									
5d <sup>10</sup> 6p <sup>1</sup>	{	1P		1D							
5d <sup>10</sup> 6d <sup>1</sup>		1P		1D		1F		1G			
	ns (n ≥ 6)			np (n ≥ 6)			nd (n ≥ 6)				
5d <sup>10</sup> 6s(1S)ns	{	1S			1P°			1D			
5d <sup>9</sup> 6s(1D)ns'		1D			1P°			1D°		1F°	
5d <sup>10</sup> 6p(1P°)ns''	{	1P°		1P°		1S		1P		1D	
		1P°		1S		1P		1D		1F°	
	nf (n ≥ 5)					ng (n ≥ 5)					nλ (n ≥ 6)
5d <sup>10</sup> 6s(1S)ns	{	1P°			1F°			1G			
5d <sup>9</sup> 6s(1D)ns'		1P°		1D°		1F°		1G°		1H°	
5d <sup>10</sup> 6p(1P°)ns''	{	1D		1F		1G		1H		1I	
		1D		1F		1G		1H		1I	

TABLE 25. PREDICTED TERMS OF THE Tl I ISOELECTRONIC SEQUENCE

Configuration 1s 2s 2p 3s 3p 3d 4s 4p 4d 4f 5s 5p 5d 6s 6p	Predicted Terms				
	ns (n ≥ 7)	np (n ≥ 7)	nd (n ≥ 6)	nf (n ≥ 5)	ng (n ≥ 5)
6s <sup>2</sup> (1S)6p	1P°				
6s 6p <sup>2</sup>	{ 1S 3P	1D			
6p <sup>3</sup>	{ 1S° 3P°	1D°			
6s <sup>2</sup> (1S)ns	ns	np	nd	nf	ng
6s 6p(1P°)ns'	{ 1P° 3P°	1D 3P	1D° 3P°	1F° 3F°	1G° 3G°
6s 6p(1P°)ns''	1P°	1D	1D°	1F	1G°

TABLE 26. PREDICTED TERMS OF THE Pb I ISOELECTRONIC SEQUENCE

Configuration 1s 2s 2p 3s 3p 3d 4s 4p 4d 4f 5s 5p 5d 6s 6p	Predicted Terms				
	ns (n ≥ 7)	np (n ≥ 7)	nd (n ≥ 6)	nf (n ≥ 5)	ng (n ≥ 5)
6s 6p <sup>2</sup>	{ 1S 3S°	1D			
6s 6p <sup>3</sup>	{ 1P° 3P°	1D° 3D°			
6s 6p(1P°)ns	{ 1P° 3P°	1D 3P	1D° 3P°	1F° 3F°	1G° 3G°
6s 6p(1P°)ns	{ 1P° 3P°	1D 3P	1D° 3P°	1F° 3F°	1G° 3G°

TABLE 27. PREDICTED TERMS OF THE Bi I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>n-1</sup> +	Predicted Terms			
	ns (n ≥ 7)	np (n ≥ 7)	nd (n ≥ 6)	nf (n ≥ 5)
6s <sup>2</sup> 6p <sup>2</sup>	{ <sup>1</sup> S <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup>		
6s 6p <sup>3</sup>	{ <sup>4</sup> S <sup>o</sup>	{ <sup>2</sup> P <sup>o</sup> <sup>2</sup> D <sup>o</sup>		
6p <sup>4</sup>		{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup>		
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>1</sup> P) <sup>ns</sup>	{ <sup>1</sup> P <sup>o</sup> <sup>1</sup> P <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> F <sup>o</sup>	{ <sup>3</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>3</sup> D) <sup>ns'</sup>	{ <sup>3</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> F <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> G <sup>o</sup>	{ <sup>3</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> H <sup>o</sup>
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>1</sup> S) <sup>ns''</sup>	{ <sup>1</sup> S <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup>	{ <sup>1</sup> D <sup>o</sup>	{ <sup>1</sup> F <sup>o</sup>

XXX

TABLE 28. PREDICTED TERMS OF THE Po I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>n-1</sup> +	Predicted Terms			
	ns (n ≥ 7)	np (n ≥ 7)	nd (n ≥ 6)	nf (n ≥ 5)
6s <sup>2</sup> 6p <sup>2</sup>	{ <sup>1</sup> S <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup>		
6s 6p <sup>3</sup>	{ <sup>4</sup> S <sup>o</sup>	{ <sup>2</sup> P <sup>o</sup> <sup>2</sup> D <sup>o</sup>		
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>1</sup> S) <sup>ns</sup>	{ <sup>1</sup> S <sup>o</sup> <sup>1</sup> S <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> F <sup>o</sup>	{ <sup>3</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>3</sup> D) <sup>ns'</sup>	{ <sup>3</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> F <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> G <sup>o</sup>	{ <sup>3</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> H <sup>o</sup>
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>1</sup> P) <sup>ns''</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> P <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup>	{ <sup>3</sup> P <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> F <sup>o</sup>	{ <sup>3</sup> D <sup>o</sup> <sup>1</sup> F <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> D <sup>o</sup> <sup>1</sup> H <sup>o</sup>

TABLE 29. PREDICTED TERMS OF THE At I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>10</sup>	Predicted Terms			
	6s <sup>2</sup> 6p <sup>2</sup>	6s	6p <sup>2</sup>	6d
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>1</sup> P) <sup>o</sup> ns	6s	6p <sup>2</sup> ( <sup>1</sup> P) <sup>o</sup>	6d	6f
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>3</sup> D) <sup>o</sup> ns'	6s	6p <sup>2</sup> ( <sup>3</sup> D) <sup>o</sup>	6d	6f

TABLE 30. PREDICTED TERMS OF THE Rn I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>10</sup> +	Predicted Terms			
	6s <sup>2</sup> 6p <sup>2</sup>	6s	6p <sup>2</sup>	6d
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>1</sup> P) <sup>o</sup> ns	6s	6p <sup>2</sup> ( <sup>1</sup> P) <sup>o</sup>	6d	6f
6s <sup>2</sup> 6p <sup>2</sup> ( <sup>3</sup> S) <sup>o</sup> ns'	6s	6p <sup>2</sup> ( <sup>3</sup> S) <sup>o</sup>	6d	6f

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>10</sup> 6s <sup>2</sup> +	Predicted Pairs			
	6p <sup>2</sup> ( <sup>1</sup> P <sub>1/2</sub> ) <sup>o</sup> ns	6p <sup>2</sup> ( <sup>1</sup> P <sub>1/2</sub> ) <sup>o</sup>	6d	6f
6p <sup>2</sup> ( <sup>1</sup> P <sub>3/2</sub> ) <sup>o</sup> ns'	6p <sup>2</sup> ( <sup>1</sup> P <sub>3/2</sub> ) <sup>o</sup>	6d	6f	6g

TABLE 31. PREDICTED TERMS OF THE Ra I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>1</sup> 3d <sup>2</sup> 3d <sup>3</sup> 4s <sup>1</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>2</sup> 3d <sup>3</sup> 4s <sup>2</sup> +	Predicted terms										
7s	1S										
6d	{ 1S 1P 1D 1F 1G										
7p	{ 1S 1P 1D										
	ns (n ≥ 7)			np (n ≥ 7)				nd (n ≥ 6)			
7s(8)ns	{ 1S 1S			1P° 1P°				1D 1D			
6d(7D)ns'	{ 1D 1D			1P° 1D° 1F° 1P° 1D° 1F°				1S 1P 1D 1F 1G 1S 1P 1D 1F 1G			
7p(7P°)ns''	{ 1P° 1P°			1S 1P 1D 1S 1P 1D				1P° 1D° 1F° 1P° 1D° 1F°			
	nf (n ≥ 5)			ng (n ≥ 5)				-----			
7s(8)ns	{ 1F° 1F°			1G 1G				----- -----			
6d(7D)ns'	{ 1P° 1D° 1F° 1G° 1H° 1P° 1D° 1F° 1G° 1H°			1D 1F 1G 1H 1I 1D 1F 1G 1H 1I				----- -----			
7p(7P°)ns''	{ 1D 1F 1G 1D 1F 1G			1F° 1G° 1H° 1F° 1G° 1H°				----- -----			

TABLE 32. PREDICTED TERMS OF THE Ac I ISOELECTRONIC SEQUENCE

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>1</sup> 3d <sup>2</sup> 3d <sup>3</sup> 4s <sup>1</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>2</sup> 3d <sup>3</sup> 4s <sup>2</sup> +	Predicted Terms										
6d 7s	1D										
6p	{ 1P 1D 1F 1G 1H 1P 1D 1F 1G 1H										
	ns (n ≥ 7)			np (n ≥ 7)				nd (n ≥ 6)			-----
7s(8)ns	1S			1P°				1D			-----
6d 7s(7D)ns	{ 1D 1D			1P° 1D° 1F° 1P° 1D° 1F°				1S 1P 1D 1F 1G 1S 1P 1D 1F 1G			----- -----
6d 7s(7D)ns	{ 1D			1P° 1D° 1F°				1S 1P 1D 1F 1G			-----
6d(7F)ns	{ 1F 1F			1D° 1F° 1G° 1D° 1F° 1G°				1P 1D 1F 1G 1H 1P 1D 1F 1G 1H			----- -----
6d(7F)ns	{ 1P 1P			1S 1P° 1D° 1S 1P° 1D°				1P 1D 1F 1P 1D 1F			----- -----

TABLE 33. REFERENCES TO RARE-EARTH SPECTRA  
The numbers in the table indicate the references in the bibliography below

Z	El	Spectrum				
		I	II	III	IV	VI
58	Ce		1, 2, 10, 31, †			
59	Pr	3, 12	3, 31			
60	Nd	17, 31, 32	17, 28, 31, 32			
62	Sm	9				
64	Gd	17, 30	30			
68	Er	12	12, 35			
71	Lu	8, 20	7			
90	Th	12, 31	17, 22, 31	19, 29	18, 21	
92	U	6, 15, 16, 31, 33, 34	5, 6, 22, 31, 32, 33, 34			18
94	Pu	4, 12‡, 23‡	4, 14			
95	Am	12, 13	12, 13			
96	Cm	11				

‡Paper refers to the spectra of the element, in general.

Table 33—Bibliography

- [1] W. Albertson, unpublished material (1949); (data at present in Spectroscopy Section, Nat. Bur. Std.). (T) (C L)
- [2] H. Arros, *Phys. Rev.* **89**, 94 (1954). (I S)
- [3] G. J. van den Berg, *Thesis*, Zeeman Lab. (Amsterdam) (1951). (Z E)
- [4] M. van den Berg and P. F. A. Klinkenberg, *Physica* **20**, 461 (1954).
- [5] J. C. van den Bosch, *Physica* **15**, 503 (1949). (T) (C L)
- [6] J. C. van den Bosch and G. J. van den Berg, *Physica* **15**, 329 (1949). (T) (Z E)
- [7] L. F. H. Bovey and R. W. B. Pearse, *Atomic Energy Research Establishment C/R 1976*, 19 pp. (1956). (I P) (T) (C L)
- [8] L. F. H. Bovey, E. B. M. Steers, and H. S. Wise, *Proc. Phys. Soc. London [A]* **69**, 783 (L) (1956). (C L); Infrared.
- [9] P. Brix, *Zeit. Phys.* **126**, 431 (1949). (T) (C L) (I S)
- [10] P. Brix und H. Frank, *Zeit. Phys.* **127**, 299 (1950). (I S)
- [11] J. G. Conway and R. D. McLaughlin, *J. Opt. Soc. Am.* **46**, 91 (1956). (I S)
- [12] M. Fred and F. S. Tomkins, *Argonne Symposium on Spectroscopy* (February 1956); See report by L. F. H. Bovey, *Nature* **177**, 737 (1956).†
- [13] M. Fred and F. S. Tomkins, *J. Opt. Soc. Am.* **44**, 824 (A) (1954); **47**, 1076 (1957). (I P) (T) (C L) (hfs)
- [14] P. M. Griffin and J. R. McNally, Jr., *J. Opt. Soc. Am.* **45**, 63 (L) (1955). (Z E)
- [15] C. J. Humphreys, U. S. Naval Ord. Lab., NAVORD Report 4571, 23 (1956); Observing techniques for infrared.
- [16] C. C. Kline, C. J. Humphreys, and D. D. Lann, *J. Research Nat. Bur. Std.* **57**, 57, RP 1729 (1946). (I P) (T) (C L) (Z E)
- [17] P. F. A. Klinkenberg, *Physica* **12**, 33 (1946). (T) (C L) (Z E)
- [18] P. F. A. Klinkenberg, *Physica* **16**, 185 (1950). (T) (C L) (Z E)
- [19] P. F. A. Klinkenberg, *Physica* **16**, 618 (1950). (I P) (T) (C L) (Z E) (E D)
- [20] P. F. A. Klinkenberg, *Physica* **21**, 53 (1955). (I P) (T) (C L) (G D)
- [21] P. F. A. Klinkenberg and R. J. Lang, *Physica* **15**, 774 (1949). (I P) (T) (C L) (Z E) (G D)
- [22] J. R. McNally, Jr., *Phys. Rev.* **77**, 417 (L) (1950). (T)
- [23] J. R. McNally, Jr., Oak Ridge Nat. Lab., ORNL—2154, 142 pp. (1956); Bibliography, wavelengths.
- [24] W. F. Meggers, *J. Opt. Soc. Am.* **36**, 431 (1946); Bibliography.†
- [25] W. F. Meggers, *Science* **106**, 514 (1947).†
- [26] W. F. Meggers and C. E. Moore, *J. Opt. Soc. Am.* **46**, 149 (1956).†
- [27] H. H. Nielsen, Program, *Tenth Anniversary Symposium on Molecular Structure and Spectroscopy*, The Ohio State Univ., pp. XIX, 53 (June 1956).†
- [28] G. Nöldke, *Zeit. Phys.* **143**, 274 (1955). (I S) (hfs)
- [29] G. Racah, *Physica* **16**, 651 (1950). (T); Theory.
- [30] H. N. Russell, *J. Opt. Soc. Am.* **46**, 550 (1950). (I P) (T) (C L)
- [31] PH. Schuurmanns, *Thesis*, Zeeman Lab. (Amsterdam) (1946). (T) (C L) (Z E) (hfs)
- [32] PH. Schuurmanns, *Physica* **11**, 419 (1946). (T) (C L) (Z E)
- [33] PH. Schuurmanns, J. C. van den Bosch, and N. Dijkwel, *Physica* **12**, 117 (1947). (T) (C L) (Z E)
- [34] A. R. Striganov and L. A. Korostyleva, *J. Exp. Tech. Phys., U. S. S. R.*, **2**, 277 (1956). (C L) (I S)
- [35] L. Wilets and L. C. Bradley III, *Phys. Rev.* **84**, 1055 (L) (1951). (I S)

†General references to rare-earth spectra.

‡G. Raab (*Bull. Res. Council Israel* **4A**, No. 1, 75, 1956) reports that in Ce II the fundamental level of Group I appears to be higher than the fundamental level of Group II by  $2064.65 \pm 0.04 \text{ cm}^{-1}$ .





Volume III

Spectrum

Z	Element	I	II	III	IV	V	VI	VII	VIII	XVI	XVII	XVIII	XIX	XXI	XXII	XXIII	XXIV
42	Mo	7.10	16.15	27.13	46.4	61.2	68	126	153								
43	Tc	7.28	15.26														
44	Ru	7.364	16.76	28.46													
45	Rh	7.46	18.07	31.05													
46	Pd	8.23	19.42	32.92													
47	Ag	7.574	21.48	34.83													
48	Cd	8.991	16.904	37.47	54.4												
49	In	5.735	18.96	28.08													
50	Sn	7.342	14.628	30.49	40.72	72.3											
51	Sb	8.639	16.5	25.3	44.1	56	106										
52	Te	9.01	18.6	31	38	60	72	137									
53	I	10.454	19.09						170								
54	Xe	12.127	21.2	32.1													
55	Cs	8.893	25.1														
56	Ba	5.210	10.001														
57	La**	5.61	11.42														
72	Hf	7	14.9														
73	Ta	7.88	16.2	19.17													
74	W	7.98	17.7														
75	Re	7.87	16.6														
76	Os	8.7	17														
77	Ir	9															
78	Pt	9.0	18.56														
79	Au	9.22	20.5														
80	Hg	10.43	18.751	34.2													
81	Tl	6.106	20.42	29.8	50.7												
83	Pb	7.415	15.028	31.93	42.31	66.8											
83	Bi	7.287	16.68	25.56	45.3	56.0	83.3										
84	Po	8.43															
85	At																
86	Rn	10.746															
87	Fr																
88	Ra	5.277	10.144	20?													
89	Ac	6.9	12.1														

11

TABLE 35. CHEMICAL SYMBOLS

Symbol	Element	Z	Symbol	Element	Z	Symbol	Element	Z	Symbol	Element	Z
Ac	Actinium	89	Er	Erbium	68	Mg	Magnesium	12	Rn	Radon	86
Ag	Silver	47	Es	Einsteinium	99	Mn	Manganese	25	Ru	Ruthenium	44
Al	Aluminum	13	Eu	Europium	63	Mo	Molybdenum	42	S	Sulfur	16
Am	Americium	95	F	Fluorine	9	N	Nitrogen	7	Sb	Antimony	51
Ar	Argon	18	Fe	Iron	26	Na	Sodium	11	Sc	Scandium	21
As	Arsenic	33	Fm	Fermium	100	Nb	Niobium	41	Se	Selenium	34
At	Astatine	85	Fr	Francium	87	Nd	Neodymium	60	Si	Silicon	14
Au	Gold	79	Ga	Gallium	31	Ne	Neon	10	Sm	Samarium	62
B	Boron	5	Gd	Gadolinium	64	Ni	Nickel	28	Sn	Tin	50
Ba	Barium	56	Ge	Germanium	32	No	Nobelium	102	Sr	Strontium	38
Be	Beryllium	4	H	Hydrogen	1	Np	Neptunium	93	Ta	Tantalum	73
Bi	Bismuth	83	(D	Deuterium)	1	O	Oxygen	8	Tb	Terbium	65
Bk	Berkelium	97	(T	Tritium)	1	Os	Osmium	76	Tc	Technetium	43
Br	Bromine	35	He	Helium	2	P	Phosphorus	15	Te	Tellurium	52
C	Carbon	6	Hf	Hafnium	72	Pa	Protactinium	91	Th	Thorium	90
Ca	Calcium	20	Hg	Mercury	80	Pb	Lead	82	Ti	Titanium	22
Cd	Cadmium	48	Hc	Holmium	67	Pd	Palladium	46	Tl	Thallium	81
Ce	Cerium	58	I	Iodine	53	Pm	Promethium	61	Tm	Thulium	69
Cf	Californium	98	In	Indium	49	Po	Polonium	84	U	Uranium	92
Cl	Chlorine	17	Ir	Iridium	77	Pr	Praseodymium	59	V	Vanadium	23
Cm	Curium	96	K	Potassium	19	Pt	Platinum	78	W	Wolfram	74
Co	Cobalt	27	Kr	Krypton	36	Pu	Plutonium	94	Xe	Xenon	54
Cr	Chromium	24	La	Lanthanum	57	Ra	Radium	88	Y	Yttrium	39
Cs	Cesium	55	Li	Lithium	3	Rb	Rubidium	37	Yb	Ytterbium	70
Cu	Copper	29	Lu	Lutetium	71	Re	Rhenium	75	Zn	Zinc	30
Dy	Dysprosium	66	Md	Mendelevium	101	Rh	Rhodium	45	Zr	Zirconium	40



TABLE 37. INDEX—ISOBLECTRONIC SEQUENCES  
 [The tabular entries are page numbers]

Z	Element	Spectrum																
		I	II	III	IV	V	VI	VII	VIII	XVI	XVII	XVIII	XIX	XX	XXI	XXII	XXIII	XXIV
42	Mo	1		11	12	13	14	15	16									
43	Tc	17	7															
44	Ru	20	19	28														
45	Rh	29	32	34														
46	Pd	38	41	44														
47	Ag	48	50	53	62													
48	Cd	55	58	60	71													
49	In	64	67	70	83	72												
50	Sn	74	80	82	93	84												
51	Sb	87	90	92	101	85												
52	Te	96	98	100	111	95												
53	I	105	108	110	125	103												
54	Xe	113	118	120	136	104												
55	Cs	124	128	130	143	111												
56	Ba	131	134			112												
57	La**	136	139			112												
72	Hf	143	146															
73	Ta	149	154															
74	W	156	161															
75	Re	164	168															
76	Os	171	176															
77	Ir	177																
78	Pt	181	183															
79	Au	186	189															
80	Hg	191	196	198	209													
81	Tl	202	204	206	207													
82	Pb	208	211	213	215	217												
83	Bi	219	221	222	223	224												
84	Po	227																
86	Rn	228	229															
88	Ra	231	233															
89	Ac	234	235	237														

# MOLYBDENUM

Mo I

42 electrons

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^5 5s^1 S_2$

$\alpha$   $^3S_1$ , 57200 K

I. P. 7.10 volts

The analysis is, for the most part, by Kiess, who started work on Mo I some 35 years ago, when he first became interested in Cr I and Cr II. After the discovery of the first multiplets of Mo I the analysis was postponed because of the need of further observations. About 8 years ago Kiess and Miss Harvey resumed work on a new description of the spectrum including Zeeman observations. The present line list extends from 2000 Å to 11850 Å, and includes about 7500 lines, of which about 80 percent are classified. The Zeeman observations are from spectrograms made at the Massachusetts Institute of Technology, as well as the National Bureau of Standards. Some observed  $g$ -values in the table are means of as many as 10 determinations.

Observed intersystem combinations connect the terms of different multiplicities. It is one of the few spectra in which four multiplicities are known.

In a spectrum as complex as Mo I it is particularly difficult to assign configurations. The low even configurations are discussed in the 1952 paper by Trees and Miss Harvey. The odd ones are tentatively assigned, in many cases. Configuration-interaction and the resulting perturbations complicate the problem. Since the work is still in progress some of the numbers in column three used to designate miscellaneous levels are subject to replacement by term designations. Consequently they do not run continuously.

The limit is well determined by the  $n s^1 S$  series of three members ( $n=5$  to 7).

A Monograph on Mo I is nearing completion.

## REFERENCES

- R. E. Trees and M. M. Harvey, *J. Research Nat. Bur. Std.* **66**, 397, RP 2378 (1952). (T)  
H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955). (Summary hfs)  
C. C. Kiess and M. M. Harvey, unpublished material (June 1956). (I P) (T) (C L) (Z E)

## Me I

## Me I

Config.	Desig.	J	Level	Interval	Obs. $\nu$	Config.	Desig.	J	Level	Interval	Obs. $\nu$
4d <sup>3</sup> (a <sup>3</sup> S)5s	a <sup>3</sup> S	3	0.00		1.992	4d <sup>3</sup> (a <sup>3</sup> F)5s	b <sup>3</sup> F	2	27093.44	681.04	0.890
4d <sup>3</sup> (a <sup>3</sup> S)5s	a <sup>3</sup> S	2	10768.33		1.980	3	27774.48		1.065		1.125
4d <sup>3</sup> 5s <sup>2</sup>	a <sup>3</sup> D	0	10965.97		0.000	4d <sup>3</sup> 5s <sup>2</sup>	b <sup>3</sup> G	3	27383.76	-41.84	0.913
		1	11142.83	176.86	1.490	4	27341.92		384.81		1.075
		2	11454.42	311.59	1.498	5	27726.73				1.219
		3	11858.54	404.12	1.488						
		4	12346.31	487.77	1.483	4d <sup>3</sup> 5s(a <sup>3</sup> D)5p	s <sup>3</sup> F <sup>o</sup>	0	27866.63	407.76	
4d <sup>3</sup> (a <sup>3</sup> G)5s	a <sup>3</sup> G	2	16641.08		0.331	1	28274.39		392.34		2.312
		3	16692.96	51.88	0.918	2	28666.73		181.40		1.905
		4	16747.73	54.77	1.141	3	28848.13		322.95		
		5	16784.60	36.87	1.284	4	29171.08		610.08		1.496
		6	16783.96	-0.64	1.318	5	29781.16		714.91		1.513
		6				6	30496.07				
4d <sup>3</sup> (a <sup>3</sup> P)5s	a <sup>3</sup> P	3	18229.23		1.636	4d <sup>3</sup> (a <sup>3</sup> I)5s	a <sup>3</sup> I	6	28241.03		1.015
		2	18356.53	-127.30	1.789	4d <sup>3</sup> (a <sup>3</sup> S)5p	s <sup>3</sup> P <sup>o</sup>	1	28715.29	121.31	2.500
		1	18479.65	-123.12	2.424	2	28836.60		87.06		1.831
4d <sup>3</sup> (a <sup>3</sup> D)5s	b <sup>3</sup> D	0	19969.85		-0.037	3	28923.66				1.664
		1	20130.38	160.53	1.579	4d <sup>3</sup> (a <sup>3</sup> F)5s	c <sup>3</sup> F	2	29642.97	517.66	1.02
		2	20281.03	150.65	1.537	3	30159.73		342.25		1.089
		3	20350.54	69.51	1.495	4	30501.98				1.111
		4	20157.90	-192.64	1.482	4d <sup>3</sup> (a <sup>3</sup> H)5s	b <sup>3</sup> H	4	29842.16		0.804
4d <sup>3</sup> 5s <sup>2</sup>	a <sup>3</sup> P	0	20607.57		0.000	5	29981.70		139.54		1.091
		1	22244.42	1636.85	0.977	6	30113.07		131.37		1.12
		2	22876.02	631.60	1.361	4d <sup>3</sup> 5s(a <sup>3</sup> D)5p	s <sup>3</sup> D <sup>o</sup>	1	30848.61	308.43	2.894
4d <sup>3</sup> (a <sup>3</sup> D)5s	a <sup>3</sup> D	1	20930.45		1.056	2	31155.04		499.76		2.034
		2	20950.88	20.43	1.289	3	31654.80		468.36		1.744
		3	21618.60	667.72	1.320	4	32123.18		488.75		1.658
4d <sup>3</sup> (a <sup>3</sup> G)5s	a <sup>3</sup> G	3	20947.86		0.767	5	32611.91				1.568
		4	21153.84	205.98	1.051	4d <sup>3</sup> 5s(a <sup>3</sup> D)5p	y <sup>3</sup> P <sup>o</sup>	2	31299.95	233.32	2.273
		5	21343.15	189.31	1.213	3	31633.27		379.97		1.84
4d <sup>3</sup> 5s <sup>2</sup>	a <sup>3</sup> F	4	23516.52		1.117	4	31913.24				1.73
		3	23668.12	-151.60	1.079	4d <sup>3</sup> 5s <sup>2</sup>	b <sup>3</sup> I	6	31484.60		1.028
		2	23534.46	133.66	0.667	4d <sup>3</sup> (a <sup>3</sup> G)5s	c <sup>3</sup> G	3			
4d <sup>3</sup> 5s <sup>2</sup>	c <sup>3</sup> H	4	24096.26		0.973	4	31510.57		768.45		
		5	24465.70	369.44	1.128	5	32279.02				
		6	24823.48	357.78	1.142	4d <sup>3</sup> (a <sup>3</sup> G)5s	b <sup>3</sup> G	4	32688.30		0.960
4d <sup>3</sup> 5s <sup>2</sup>	a <sup>3</sup> S	0	24472.06			4d <sup>3</sup> 5s(a <sup>3</sup> D)5p	y <sup>3</sup> P <sup>o</sup>	1	32898.81	400.28	2.550
4d <sup>3</sup>	c <sup>3</sup> D	4	25455.58		1.458	2	33299.09		656.00		1.838
		3	25707.15	-251.57	1.355	3	33955.09				1.659
		2	25794.65	-87.50	1.09	4d <sup>3</sup> (a <sup>3</sup> H)5s	a <sup>3</sup> H	5	33904.46		1.026
		1	25820.67	-26.02		4d <sup>3</sup> 5s(a <sup>3</sup> D)5p	s <sup>3</sup> F <sup>o</sup>	1	34248.43	186.28	-0.004
		0	25960.3	-159.6		2	34434.71		305.69		1.010
4d <sup>3</sup> (a <sup>3</sup> I)5s	a <sup>3</sup> I	5	25516.95		0.796	3	34740.40		429.04		1.258
		6	25548.90	31.95	1.028	4	35169.44		549.93		1.346
		7	25638.63	89.73	1.147	5	35719.37				1.395
4d <sup>3</sup> (a <sup>3</sup> S)5p	s <sup>3</sup> P <sup>o</sup>	2	25614.31		2.299	4d <sup>3</sup>	c <sup>3</sup> H	6	34810.14	-101.92	1.09
		3	25771.86	257.55	1.892	5	34912.06		-130.00		1.10
		4	26320.38	448.52	1.736	4	35042.06				0.79
4d <sup>3</sup> (a <sup>3</sup> F)5s	a <sup>3</sup> F	5	25905.58		1.394	4d <sup>3</sup> 5s(a <sup>3</sup> D)5p	s <sup>3</sup> D <sup>o</sup>	0	37128.14	164.86	0.000
		4	25997.32	-91.74	1.322	1	37293.00		286.23		1.480
		3	26189.48	-192.16	1.307	2	37679.23		389.24		1.473
		2	26335.87	-146.39	1.031	3	37988.47		454.63		1.492
		1	26283.82	52.05		4	38423.10				1.482
4d <sup>3</sup> (a <sup>3</sup> P)5s	b <sup>3</sup> P	0	26450.10			4d <sup>3</sup> 5s(b <sup>3</sup> D)5p	y <sup>3</sup> D <sup>o</sup>	0	37385.69	535.86	0.000
		1	26414.90	-35.20	1.41	1	37901.55		620.71		1.491
		2	27415.27	1000.37	1.41	2	38522.86		637.63		1.471
4d <sup>3</sup> 5s <sup>2</sup>	a <sup>3</sup> G	4	26635.82		1.014	3	39159.89		755.75		1.251
4d <sup>3</sup> (a <sup>3</sup> D)5s	b <sup>3</sup> D	3	26638.84		1.234	4	39915.64				1.463
		2	26758.70	-119.86	1.24	1 <sup>o</sup>	38626.50				1.443
		1	27362.55	-603.85	1.06						

## Me I—Continued

## Me I—Continued

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$				
4d <sup>3</sup> (a <sup>4</sup> G)5p	z <sup>4</sup> G°	2	38983.40		0.428	4d <sup>3</sup> 5s(a <sup>4</sup> H)5p	y <sup>4</sup> H°	3	43697.63	314.82	0.554				
		3	39181.48	138.08	1.070			4	44019.35	432.16	0.867				
		4	39289.63	168.15				5	44444.51	250.42					
		5	39445.91	155.58				6	44894.93	710.97					
		6	39582.19	76.96				7	45406.90						
4d <sup>3</sup> 5s(b <sup>4</sup> D)5p	z <sup>4</sup> P°	1	39358.58		2.485	4d <sup>3</sup> 5s(a <sup>4</sup> H)5p	y <sup>4</sup> G°	3	43976.68	485.07	1.075				
		2	39482.74	104.16	1.696			4	44460.63	299.11	1.170				
		3	39900.54	497.80	1.262			5	44759.74						
4d <sup>3</sup>	c <sup>4</sup> I	6	39521.09		0.982	4d <sup>3</sup> (a <sup>4</sup> D)5p	y <sup>4</sup> D°	1	44041.09	483.50	1.123				
4d <sup>3</sup> (a <sup>4</sup> G)5p	z <sup>4</sup> H°	7	39600.05					2	44584.59	396.45					
		6	39635.23	-35.18				3	44921.04						
		5	40367.34	-732.11		4d <sup>3</sup> (a <sup>4</sup> S)5d	e <sup>4</sup> D	1	44935.78	4.61					
		4	40067.67	299.67				2	44940.39	6.87	1.94				
3	39749.93	317.74		3	44947.26			9.95	1.650						
				4	44957.21			12.89	1.605						
				5	44970.10										
4d <sup>3</sup> (a <sup>4</sup> S)6s	e <sup>4</sup> S	3	39675.42		1.992	4d <sup>3</sup> (a <sup>4</sup> G)5p	z <sup>4</sup> G°	2	45398.66						
4d <sup>3</sup> 5s(b <sup>4</sup> D)5p	z <sup>4</sup> P°	0						3	45414.90	141.38	0.794				
		1	39779.89	41.54	1.346	4	45556.28	279.24	1.108						
4d <sup>3</sup> 5s(b <sup>4</sup> D)5p	z <sup>4</sup> F°	2	39939.18†		1.350	4d <sup>3</sup> 5s(b <sup>4</sup> F)5p	w <sup>4</sup> F°	5	45835.68		1.100				
		3						1	45457.86	-32.74	0.104				
		4	40843.53		1.256			2	45426.11	209.59	1.286				
								3	45634.70	234.87	1.162				
4d <sup>3</sup> 5s(b <sup>4</sup> D)5p	z <sup>4</sup> D°	1	40087.01		0.781			4	45869.67		1.334				
		2	40566.27	529.26	1.160	4d <sup>3</sup> (a <sup>4</sup> D)5p	z <sup>4</sup> F°	2	45710.01	228.51	0.890				
		3	40964.11	397.84	1.278			3	45938.62	30.69	1.110				
				4	45969.21				1.135						
2°	40840.84			4d <sup>3</sup> (a <sup>4</sup> S)5d	e <sup>4</sup> D			4	45785.69	-7.08					
4d <sup>3</sup> (a <sup>4</sup> P)5p	z <sup>4</sup> D°	4	40488.35			-340.50	1.470	3	45792.77	-7.39					
		2	40828.85			130.39	1.474	2	45800.16	-5.31					
		1	40928.46			-37.54	1.393	0	45805.47	-1.90					
		0	40736.00		0.000	0	45807.37								
4d <sup>3</sup> (a <sup>4</sup> S)6s	e <sup>4</sup> S	2	40840.28			4d <sup>3</sup> (a <sup>4</sup> P)5p	z <sup>4</sup> S°	1	45834.83		2.048				
		4d <sup>3</sup> (a <sup>4</sup> G)5p	y <sup>4</sup> F°	1	41011.74†			20.27	1.235	4d <sup>3</sup> (a <sup>4</sup> F)5p	z <sup>4</sup> D°	1	45974.34	327.44	1.078
				2	41032.01			192.40	1.053			2	46301.78	506.69	1.275
				3	41224.41			171.13	1.259			3	46808.47		
				4	41395.54			-47.83	1.343	4d <sup>3</sup> (a <sup>4</sup> F)5p	v <sup>4</sup> F°	1	46447.97	4.56	
5	41347.71				1.385	2	46462.63	0.89							
					3	46453.42	303.08	1.304							
					4	46756.50									
					5										
4d <sup>3</sup> (a <sup>4</sup> G)5p	y <sup>4</sup> F°	2	41484.54	365.43	0.781	4d <sup>3</sup> (a <sup>4</sup> F)5p	v <sup>4</sup> F°	4	46590.07		1.160				
		3	41849.97	231.62	1.084			4d <sup>3</sup> (a <sup>4</sup> D)5p	w <sup>4</sup> D°	1	46720.22				
		4	42031.59		1.225					2	46754.90				
4d <sup>3</sup> (a <sup>4</sup> D)5p	w <sup>4</sup> D°	0	42173.10	-16.96	0.000	4d <sup>3</sup> (a <sup>4</sup> I)5p	z <sup>4</sup> I°	5	46861.06	310.70	0.937				
		1	42156.14	81.27	1.438			6	47171.76	321.14	1.13				
		2	42237.41	184.99	1.442			7	47492.90		1.14				
		3	42422.40	319.42	1.425			4d <sup>3</sup> (a <sup>4</sup> D)5p	w <sup>4</sup> F°	2	46874.52	51.88	1.093		
		4	42741.82		1.403					3	46928.40				
				4											
4d <sup>3</sup> (a <sup>4</sup> G)5p	z <sup>4</sup> H°	4	42185.84	97.39	0.851	4d <sup>3</sup> (a <sup>4</sup> D)5p	w <sup>4</sup> D°	1	46895.06	215.18	1.156				
		5	42263.23	61.75	1.258			2	47110.24	74.34	1.290				
		6	42344.98					3	47184.68						
4d <sup>3</sup> (a <sup>4</sup> G)5p	z <sup>4</sup> G°	3	42970.06	226.89	0.884	4d <sup>3</sup> (a <sup>4</sup> D)5p	w <sup>4</sup> D°	3	46955.84		1.66				
		4	43196.95	748.69	1.097			10°	1	47038.81					
		5	43945.64		1.359			11°	2	47051.74		0.917			
4d <sup>3</sup> (a <sup>4</sup> D)5p	z <sup>4</sup> F°	1				4d <sup>3</sup> (a <sup>4</sup> D)5p	w <sup>4</sup> D°	2							
		2	43045.53	200.09	1.030										
		3	43245.62	284.23	1.248										
		4	43529.85	-230.77	1.365										
		5	43299.08		1.363										
4d <sup>3</sup> 5s(a <sup>4</sup> H)5p	z <sup>4</sup> I°	4	43598.54	731.69	0.779										
		5	44330.23		0.956										
		6													
		7													
		8													

Mo I—Continued

Mo I—Continued

Config.	Desig.	J	Level	Interval	Obs. $\sigma$	Config.	Desig.	J	Level	Interval	Obs. $\sigma$	
4d <sup>5</sup> 5s(b <sup>4</sup> F)5p	12°	3	47064.59		1.13	4d <sup>5</sup> 5s(b <sup>4</sup> G)5p	t <sup>4</sup> F°	1	49189.71		0.31	
	13°	3	47177.18				2	49346.90	156.59	1.20		
	y <sup>4</sup> G°	2	47186.547	124.72	0.862		3	49488.06	131.75	1.216		
		3	47311.26	98.02	1.200		4	49690.83	122.78	1.245		
		4	47409.28	249.75	1.252		5	49660.68	59.85	1.352		
		5	47659.03	45.65	1.30		28°	3	49859.38		1.348	
	14°	1	47848.15		1.164		4d <sup>5</sup> 5s(a <sup>4</sup> H)5p	y <sup>4</sup> I°	5	49350.60	49.78	0.914
	15°	2	47882.23					6	49400.38	164.75	1.108	
	16°	0	47324.26		0.00			7	49665.13		1.122	
	17°	1	47408.23		0.889		29°	4	49869.50		1.475	
18°	5	47491.68		1.112	4d <sup>5</sup> 5s(b <sup>4</sup> P)5p	w <sup>4</sup> P°	1	49484.14	227.72	2.378		
19°	2	47587.58		1.12		2	49711.86	1498.06	2.16			
20°	4	47710.81		1.056		3	51909.92		1.66			
4d <sup>5</sup> 5s(c <sup>4</sup> D)5p	v <sup>4</sup> D°	1	47779.41	305.85	0.076	4d <sup>5</sup> 5s(a <sup>4</sup> D)6s	f <sup>4</sup> D	1	49608.20		2.990	
		2	48085.26	120.69	1.044		2	49852.29	244.09	303.21		
		3	48205.95		1.37		3	50155.50	634.84	1.613		
4d <sup>5</sup> 5s(c <sup>4</sup> D)5p	u <sup>4</sup> F°	22°	3	47848.40		1.076	4	50790.34	485.35	1.611		
		1	48022.23	-195.01	0.29	4d <sup>5</sup> 5s(b <sup>4</sup> G)5p	u <sup>4</sup> F°	2	49801.41	-11.58	1.061	
		2	47827.22	201.00	1.09		3	49789.83	344.41	1.210		
		3	48022.22	163.83	1.283		4	50134.24		1.054		
		4	48122.05	355.39	1.293	5	50631.35	767.90	1.123			
5	48547.44		1.370	31°	2	49880.46		1.72				
4d <sup>5</sup> (a <sup>3</sup> I)5p	y <sup>4</sup> H°	4	47876.73	312.49	1.206	4d <sup>5</sup> 5s(c <sup>4</sup> D)5p	v <sup>4</sup> P°	3	49899.49	-1050.11	1.61	
		5	48122.22	211.04	1.195		2	51049.60	936.33	2.52		
		6	48400.26				1	50113.27		1.046		
4d <sup>5</sup> (a <sup>3</sup> F)5p	v <sup>4</sup> F°	23°	2	48038.89		1.20	4d <sup>5</sup> (a <sup>3</sup> I)5p	s <sup>4</sup> K°	8	50276.83	-319.10	0.968
		24°	6	48120.58				7	50595.93		1.046	
		2	48238.21	160.79	0.758	6		50595.93		1.046		
4d <sup>5</sup> (a <sup>3</sup> S)7s	f <sup>4</sup> S	3	48367.66		1.174	4d <sup>5</sup> (a <sup>3</sup> S)6d	g <sup>4</sup> D	5	50364.36	40.93		
		2	48238.21				4	50323.43	-48.31			
		3	48399.00				3	50371.74				
		4					2					
4d <sup>5</sup> 5s(c <sup>4</sup> D)5p	v <sup>4</sup> D°	0				4d <sup>5</sup> (a <sup>3</sup> S)6d	g <sup>4</sup> D	1				
		1	48410.88	-2.19	1.440		32°	2	50482.46		1.07	
		2	48408.69	123.41	1.496		33°	4	50625.18		1.23	
		3	48532.10	338.34	1.454		34°	3	50670.03			
4d <sup>5</sup> (a <sup>3</sup> G)5p	z <sup>4</sup> H°	4	48870.44			35°	2	50769.72				
		4	48622.08	196.89	0.926	36°	4	50770.12				
		5	48818.97	242.76	0.972	37°	5	50801.61		1.083		
4d <sup>5</sup> (a <sup>3</sup> S)7s	f <sup>4</sup> S	6	49061.73		1.108	38°	2	50925.56				
		2	48742.20			39°	4	50953.80	799.62	0.937		
4d <sup>5</sup> 5s(b <sup>4</sup> G)5p	z <sup>4</sup> G°	25°	1	48775.48			5	51753.42		1.05v		
		2	48875.84	121.40	0.60	4d <sup>5</sup> 5s(a <sup>4</sup> H)5p	w <sup>4</sup> H°	4	50953.80	799.62	0.937	
		3	48897.24	149.45	1.164		5	51753.42		1.05v		
		4	49146.69	155.29	1.25		6					
		5	49301.98	106.56	1.150		39°	1	51070.34			
		6	49408.54				40°	1	51165.07			
4d <sup>5</sup> (a <sup>3</sup> I)5p	s <sup>4</sup> H°	5	49222.21		0.970		41°	4	51234.16		1.26	
		26°	1	48955.80		0.77	42°	2	51404.83		1.675	
		27°	1	49169.31			43°	4	51585.39		1.040	



Mo I—Continued

Mo I—Continued

Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	
4d <sup>5</sup> 5s(a <sup>4</sup> D) 6s	44°	2	51875.27		1.146	Mo II(8S <sub>M</sub> )	83°	4	53855.20		1.17	
	45°	3	51898.89		0.88		84°	5	53937.80		1.14	
	46°	6	51856.30				85°	4	53994.53		1.016	
	47°	4	51853.82		1.258		86°	1	54184.34			
	48°	1	51876.03		0.87		87°	4	54292.23		1.04	
	49°	3	51824.52		1.13		88°	1	54341.00			
	50°	2	51891.30		0.56		89°	2	54342.90		1.48	
	51°	4	52025.30		1.266		90°	1	54531.12			
	52°	2	52033.09				91°	1	54571.76		0.75	
	53°	3	52135.63		0.91		92°	5	54704.54		0.911	
	54°	3	52132.01		1.34		93°	7	54738.58		1.10	
	55°	4	52404.79		1.16		94°	3	54839.36		0.31	
	56°	2	52441.17				95°	3	54889.12		0.84	
	57°	5	52441.33				96°	4	55025.75		1.050	
	58°	2	52443.57		1.12		97°	5	55139.74		0.968	
	59°	5	52465.36		1.068		98°	3	55327.60			
		<i>f</i> <sup>4</sup> D	0					99°	3	55352.31		
			1					100°	5	55387.34		1.12
			2					101°	4	55529.85		
			3					102°	4	55907.71		1.09
			4	52477.60				103°	5	55975.93		1.10
	60°	4	52579.34		1.055		4d <sup>5</sup> (a <sup>3</sup> H)5p	<i>g</i> <sup>3</sup> H°	5	56181.71		0.96
	61°	5	52667.27		1.18			104°	3	56330.79		0.03
	62°	2	52740.51				4d <sup>5</sup> (a <sup>3</sup> H)5p	<i>s</i> <sup>1</sup> I°	6	56716.58		1.00
	63°	1	52746.93					105°	6	56890.71		1.21
	64°	3	52902.26					106°	4	56945.78		
	65°	1	52941.36					107°	5	57029.31		0.98
	66°	2	52964.76					108°	3	57144.26		
	67°	4	53005.71		1.35			Limit		57260		
	68°	5	53112.20		1.40			109°	4	57818.12		
	69°	6	53124.27		1.04			110°	4	58090.10		
	70°	5	53197.79		1.324			111°	3	58192.82		
71°	3	53226.86		1.01		112°	6	58525.71		1.15		
72°	6	53290.73				113°	5	58998.62		0.88		
73°	3	53305.79				114°	3	59236.59				
74°	5	53327.30				115°	2	59249.76				
75°	4	53478.86				116°	5	59436.61		1.07		
76°	1	53433.00				117°	5	59612.36		1.09		
77°	2	53538.78		1.43		118°	5	59841.11		0.930		
78°	3	53595.18		1.18		119°	6	60192.00				
79°	7	53636.76		1.125		120°	7	61470.51				
80°	3	53643.84		1.37								
81°	3	53735.87										
82°	4	53800.43		1.49								

June 1956.

Mo I OBSERVED TERMS \*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>4</sup> +	Observed Terms			
	ns (n ≥ 5)		np (n ≥ 5)	
4d <sup>5</sup> 5s <sup>2</sup>	a <sup>1</sup> S	a <sup>3</sup> P	a <sup>1</sup> D	a <sup>3</sup> F
4d <sup>5</sup>			b <sup>1</sup> G	b <sup>3</sup> H
			c <sup>1</sup> H	c <sup>3</sup> I
4d <sup>4</sup> (a <sup>1</sup> S)ns	{ a <sup>1</sup> S, c <sup>1</sup> I, f <sup>1</sup> H a <sup>3</sup> P, e <sup>3</sup> F, f <sup>3</sup> H			
4d <sup>4</sup> 5s(a <sup>1</sup> D)ns	{ f <sup>1</sup> D f <sup>3</sup> D			
4d <sup>4</sup> (a <sup>1</sup> G)ns	{ e <sup>1</sup> G e <sup>3</sup> G			
4d <sup>4</sup> (a <sup>1</sup> P)ns	{ a <sup>3</sup> P b <sup>3</sup> P			
4d <sup>4</sup> (a <sup>1</sup> D)ns	{ b <sup>1</sup> D a <sup>3</sup> D			
4d <sup>4</sup> (a <sup>3</sup> D)ns	{ b <sup>3</sup> D			
4d <sup>4</sup> (a <sup>3</sup> I)ns	{ a <sup>3</sup> I a <sup>1</sup> I			
4d <sup>4</sup> (a <sup>3</sup> F)ns	{ a <sup>3</sup> F b <sup>3</sup> F			
4d <sup>4</sup> 5s(b <sup>1</sup> D)ns	{ z <sup>1</sup> P° z <sup>3</sup> P°			
4d <sup>4</sup> (a <sup>3</sup> F)ns	{ z <sup>1</sup> D° y <sup>1</sup> D° z <sup>3</sup> D°			
4d <sup>4</sup> 5s(a <sup>1</sup> H)ns	{ z <sup>1</sup> P° z <sup>3</sup> P°			
4d <sup>4</sup> (a <sup>1</sup> G)ns	{ z <sup>1</sup> G° z <sup>3</sup> G°			
4d <sup>4</sup> 5s(b <sup>1</sup> P)ns	{ w <sup>1</sup> P°			
4d <sup>4</sup> (a <sup>3</sup> H)ns	{ b <sup>3</sup> H a <sup>1</sup> H			
4d <sup>4</sup> 5s(b <sup>3</sup> F)ns	{ v <sup>1</sup> P° f <sup>1</sup> P° z <sup>1</sup> G° w <sup>1</sup> G°			
4d <sup>4</sup> (b <sup>1</sup> G)ns	{ v <sup>1</sup> P° v <sup>3</sup> P° u <sup>1</sup> P° u <sup>3</sup> P°			
4d <sup>4</sup> 5s(c <sup>1</sup> D)ns	{ v <sup>1</sup> P° v <sup>3</sup> P° u <sup>1</sup> P° u <sup>3</sup> P°			

\*For predicted terms in the spectra of the Mo I isoelectronic sequence, see Vol. III, Introduction.

## Mo II

(Nb I sequence; 41 electrons)

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^2 4p^6 4d^5 {}^6S_{5/2}$  $\lambda$   ${}^6S_{5/2}$  136200 K

I. P. 16.15 volts

Since 1932 no extensive work has been done on Mo II except at the National Bureau of Standards. A detailed description and analysis have been completed by Kiess especially for inclusion here, and a Monograph has been completed.

His observations extend from 2000 Å to 6000 Å, supplemented by several hundred lines to shorter waves, measured on spectrograms furnished by Boyce and by Shenstone. Altogether, about seventy percent of the known Mo II lines have been classified.

The observed  $g$ -values have been derived from Zeeman spectrograms taken at the National Bureau of Standards and at the Massachusetts Institute of Technology.

The limit is from Catalán and Rico, who have interpolated it from data on second spectra from Sr to Cd.

The prefixes  $a$ ,  $b$  of the limit terms of Mo III as given in the table are tentative. The terms of Mo III are very incompletely known, and the prefixes have been arbitrarily assigned for convenience in describing the configurations in Mo II.

The present Monograph by Kiess represents work covering a span of 35 years, culminating in the classification of more than 2600 lines. Numerous observed intersystem combinations connect the terms of different multiplicity. It is noteworthy that Kiess has found 14 of the 16 terms expected from the  $4d^5$  configuration.

## REFERENCES

M. A. Catalán y F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] 48, 323 (1952) and letter (December 1956.) (I P)

C. C. Kiess, J. Research Nat. Bur. Std. 66, 375, RP2856 (1958). (T) (C L) (Z E)

## Mo II

## Mo II

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$
$4d^5$	$a {}^6S$	$2\frac{1}{2}$	0.00			$4d^4(a {}^6D)5s$	$b {}^4D$	$0\frac{1}{2}$	24372.12	287.08	0.068
$4d^4(a {}^6D)5s$	$a {}^6D$	$0\frac{1}{2}$	11783.36	250.70	3.301			$1\frac{1}{2}$	24659.20	453.07	1.194
		$1\frac{1}{2}$	12034.06	383.22	1.847			$2\frac{1}{2}$	25112.27	220.31	1.269
		$2\frac{1}{2}$	12417.28	483.05	1.635			$3\frac{1}{2}$	25341.58		1.387
		$3\frac{1}{2}$	12900.33	560.37	1.584	$4d^5$	$a {}^4F$	$3\frac{1}{2}$	24509.30	-326.79	1.180
		$4\frac{1}{2}$	13460.70		1.543			$2\frac{1}{2}$	24836.09		1.054
$4d^5$	$a {}^4G$	$2\frac{1}{2}$	15199.25	181.31		$4d^4(a {}^4H)5s$	$a {}^4H$	$3\frac{1}{2}$	26041.18	446.97	0.798
		$3\frac{1}{2}$	15330.56	97.17				$4\frac{1}{2}$	26488.15	251.32	0.984
		$4\frac{1}{2}$	15427.73	19.24				$5\frac{1}{2}$	26739.47	374.36	1.130
		$5\frac{1}{2}$	16446.97					$6\frac{1}{2}$	27113.83		1.198
$4d^5$	$a {}^4P$	$2\frac{1}{2}$	15691.22	-7.94	1.595	$4d^5$	$a {}^2G$	$4\frac{1}{2}$	26068.60	-337.01	1.065
		$1\frac{1}{2}$	15699.16	-190.96				$3\frac{1}{2}$	26405.61		0.785
		$0\frac{1}{2}$	15890.12			$4d^4(a {}^2P)5s$	$b {}^4P$	$0\frac{1}{2}$	26603.55	1024.00	2.530
$4d^5$	$a {}^4D$	$0\frac{1}{2}$	16796.14	377.96	0.758			$1\frac{1}{2}$	27627.55	1394.57	1.700
		$1\frac{1}{2}$	17174.10	170.00	1.391			$2\frac{1}{2}$	29022.12		1.574
		$2\frac{1}{2}$	17344.10	-397.32	1.433	$4d^5$	$b {}^2F$	$3\frac{1}{2}$	27410.30	-468.59	1.148
		$3\frac{1}{2}$	16946.78		1.404			$2\frac{1}{2}$	27878.89		0.900
$4d^5$	$a {}^2D$	$2\frac{1}{2}$	22444.36	-420.00	1.082	$4d^5$	$a {}^2H$	$5\frac{1}{2}$	27627.00	-97.69	1.086
		$1\frac{1}{2}$	22864.36		0.688			$4\frac{1}{2}$	27724.69		0.987
$4d^5$	$a {}^2I$	$5\frac{1}{2}$	22980.48	267.71		$4d^4(a {}^2F)5s$	$b {}^4F$	$1\frac{1}{2}$	28883.69	-6.87	0.529
		$6\frac{1}{2}$	23248.19					$2\frac{1}{2}$	28876.82	112.14	1.035
$4d^5$	$a {}^2F$	$4\frac{1}{2}$	23832.86	-20.49	1.300			$3\frac{1}{2}$	28988.96	45.21	1.135
		$3\frac{1}{2}$	23853.35	-81.01	1.219			$4\frac{1}{2}$	29034.17		1.278
		$2\frac{1}{2}$	23934.56	-203.29	1.018	$4d^5$	$a {}^2S$	$0\frac{1}{2}$	28950.36		1.968
		$1\frac{1}{2}$	24137.65		0.514						

## Mo II—Continued

## Mo II—Continued

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$			
4d <sup>4</sup> (a <sup>4</sup> G)5e	b <sup>4</sup> G	3 $\frac{1}{2}$	30699.32	230.04	0.758	4d <sup>4</sup> (a <sup>4</sup> P)5p	y <sup>4</sup> D <sup>o</sup>	0 $\frac{1}{2}$	57319.55	821.20	0.200			
		3 $\frac{3}{4}$	30010.36		1.059			1 $\frac{1}{2}$	58140.76		1.183			
		4 $\frac{1}{4}$	30213.46		1.192			2 $\frac{1}{2}$	59347.94		1.263			
		5 $\frac{1}{4}$	30391.28		1.268			3 $\frac{1}{2}$	60702.16		1.306			
4d <sup>4</sup> (a <sup>4</sup> P)5e	a <sup>4</sup> P	0 $\frac{1}{2}$	32124.04	2295.22	0.672	4d <sup>4</sup> (a <sup>4</sup> H)5p	z <sup>4</sup> H <sup>o</sup>	3 $\frac{1}{2}$	57892.06	304.62	0.710			
		1 $\frac{1}{2}$	34419.26		1.176			4 $\frac{1}{2}$	58196.68		564.27	0.900		
4d <sup>4</sup>	b <sup>4</sup> D	2 $\frac{1}{2}$	32879.55	-206.73	1.213	4d <sup>4</sup> (a <sup>4</sup> P)5p	z <sup>4</sup> S <sup>o</sup>	0 $\frac{1}{2}$	58587.00	1.651	1.110			
		1 $\frac{1}{2}$	33086.28		0.889			6 $\frac{1}{2}$	59491.83		730.88	1.206		
4d <sup>4</sup> (a <sup>4</sup> H)5e	b <sup>4</sup> H	4 $\frac{1}{4}$	33045.37	555.70	0.983	4d <sup>4</sup> (a <sup>4</sup> F)5p	z <sup>4</sup> G <sup>o</sup>	2 $\frac{1}{2}$	59053.32	425.02	0.672			
		5 $\frac{1}{4}$	33601.07		1.057			3 $\frac{1}{2}$	59478.34		748.66	0.923		
4d <sup>4</sup> (a <sup>4</sup> G)5e	b <sup>4</sup> G	3 $\frac{1}{2}$	33146.30	108.16	0.904	4d <sup>4</sup> (a <sup>4</sup> H)5p	z <sup>4</sup> I <sup>o</sup>	4 $\frac{1}{2}$	59879.75	1244.78	1.027			
		4 $\frac{1}{2}$	33254.46		1.043			5 $\frac{1}{2}$	60294.53		723.23	0.936		
4d <sup>4</sup> (a <sup>4</sup> D)5e	c <sup>4</sup> D	3 $\frac{1}{2}$	33549.28	-320.44	1.406	4d <sup>4</sup> (a <sup>4</sup> F)5p	z <sup>4</sup> D <sup>o</sup>	1 $\frac{1}{2}$	59840.70	1151.77	0.862			
		2 $\frac{1}{2}$	33869.72		1.323			2 $\frac{1}{2}$	60992.47		1.205			
		1 $\frac{1}{2}$	33525.16		1.230			3 $\frac{1}{2}$	60135.37		837.77	1.011		
		0 $\frac{1}{2}$	33895.06		0.021			4 $\frac{1}{2}$	60973.14		1.101			
4d <sup>4</sup> (a <sup>4</sup> I)5e	b <sup>4</sup> I	6 $\frac{1}{2}$	35099.46	-306.56	1.053	4d <sup>4</sup> (a <sup>4</sup> P)5p	y <sup>4</sup> P <sup>o</sup>	0 $\frac{1}{2}$	61134.26	322.52	2.299			
		5 $\frac{1}{2}$	35406.02		1.065			1 $\frac{1}{2}$	61456.77		968.70	1.656		
4d <sup>4</sup> (a <sup>4</sup> F)5e	c <sup>4</sup> F	2 $\frac{1}{2}$	36288.80	452.50	0.913	4d <sup>4</sup> (a <sup>4</sup> P)5p	z <sup>4</sup> P <sup>o</sup>	2 $\frac{1}{2}$	62425.47	968.70	1.185			
		3 $\frac{1}{2}$	36741.30		1.065			1 $\frac{1}{2}$	61748.58		-349.47	1.220		
4d <sup>4</sup> (a <sup>4</sup> G)5e	c <sup>4</sup> G	3 $\frac{1}{2}$	37431.45	622.43	0.927	4d <sup>4</sup> (a <sup>4</sup> F)5p	z <sup>4</sup> D <sup>o</sup>	0 $\frac{1}{2}$	62096.06	-349.47	1.075			
		4 $\frac{1}{2}$	38053.88		1.105			3 $\frac{1}{2}$	62491.78		149.51	1.252		
4d <sup>4</sup> (a <sup>4</sup> D)5e	c <sup>4</sup> D	1 $\frac{1}{2}$	39243.45	669.50	0.800	4d <sup>4</sup> (a <sup>4</sup> H)5p	z <sup>4</sup> I <sup>o</sup>	2 $\frac{1}{2}$	62342.27	-208.77	1.067			
		2 $\frac{1}{2}$	39912.95		1.203			1 $\frac{1}{2}$	62551.04		-386.41	0.707		
4d <sup>4</sup> (a <sup>4</sup> D)5e	d <sup>4</sup> D	2 $\frac{1}{2}$	41421.34	-120.66	0.761	4d <sup>4</sup> (a <sup>4</sup> F)5p	z <sup>4</sup> D <sup>o</sup>	0 $\frac{1}{2}$	62937.45	-386.41	0.030			
		1 $\frac{1}{2}$	41542.00		0.761			1 <sup>o</sup>	62594.53		1.14			
4d <sup>4</sup> (a <sup>4</sup> S)5e	b <sup>4</sup> S	0 $\frac{1}{2}$	41873.66	137.32	1.993	4d <sup>4</sup> (a <sup>4</sup> H)5p	z <sup>4</sup> I <sup>o</sup>	5 $\frac{1}{2}$	62728.35	251.89	0.902			
		3 $\frac{1}{2}$	42169.30		1.873			6 $\frac{1}{2}$	62980.24		1.091			
4d <sup>4</sup>	d <sup>4</sup> G	3 $\frac{1}{2}$	42169.30	137.32	1.873	4d <sup>4</sup> (a <sup>4</sup> F)5p	y <sup>4</sup> F <sup>o</sup>	1 $\frac{1}{2}$	63002.58	389.94	0.899			
		4 $\frac{1}{2}$	42306.62		1.873			2 $\frac{1}{2}$	63392.52		-267.89	1.002		
4d <sup>4</sup> (a <sup>4</sup> F)5e	d <sup>4</sup> F	2 $\frac{1}{2}$	42925.34	66.84	1.085	4d <sup>4</sup> (a <sup>4</sup> P)5p	z <sup>4</sup> P <sup>o</sup>	3 $\frac{1}{2}$	63104.63	678.47	1.060			
		3 $\frac{1}{2}$	42992.18		1.085			4 $\frac{1}{2}$	63783.10		1.186			
4d <sup>4</sup> (b <sup>4</sup> F)5e	o <sup>4</sup> F	1 $\frac{1}{2}$	44212.16	-0.650	1.072	4d <sup>4</sup> (a <sup>4</sup> H)5p	y <sup>4</sup> G <sup>o</sup>	2 $\frac{1}{2}$	63041.47	-123.53	0.851			
		3 $\frac{1}{2}$						45853.08	1.072		3 $\frac{1}{2}$	62917.94	35.81	1.004
		4 $\frac{1}{2}$						46148.12	1.305		4 $\frac{1}{2}$	62953.75	253.68	1.214
		5 $\frac{1}{2}$						46614.14	1.375		5 $\frac{1}{2}$	63207.43	1.272	
4d <sup>4</sup> (a <sup>4</sup> T)5p	z <sup>4</sup> F <sup>o</sup>	0 $\frac{1}{2}$	47808.36	814.09	2.779	4d <sup>4</sup> (a <sup>4</sup> G)5p	y <sup>4</sup> H <sup>o</sup>	3 $\frac{1}{2}$	63298.23	199.31	0.933			
		1 $\frac{1}{2}$	48022.45		1.818			4 $\frac{1}{2}$	63497.54		642.44	1.024		
		2 $\frac{1}{2}$	48901.57		1.742			5 $\frac{1}{2}$	64139.98		974.85	1.166		
		3 $\frac{1}{2}$	49481.04		1.672			6 $\frac{1}{2}$	65114.83		1.105			
4d <sup>4</sup> (a <sup>4</sup> T)5p	z <sup>4</sup> P <sup>o</sup>	1 $\frac{1}{2}$	49040.82	567.92	2.305	4d <sup>4</sup> (a <sup>4</sup> G)5p	z <sup>4</sup> F <sup>o</sup>	2 $\frac{1}{2}$	63876.68	-517.96	0.802			
		2 $\frac{1}{2}$	49608.74		1.718			2 $\frac{1}{2}$	64394.64		1.105			
		3 $\frac{1}{2}$	49481.04		1.672			2 $\frac{1}{2}$	64394.64		0.802			
4d <sup>4</sup> (a <sup>4</sup> T)5p	z <sup>4</sup> D <sup>o</sup>	0 $\frac{1}{2}$	49949.45	242.55	2.155	4d <sup>4</sup> (a <sup>4</sup> G)5p	z <sup>4</sup> F <sup>o</sup>	1 $\frac{1}{2}$	63903.80	263.94	0.535			
		1 $\frac{1}{2}$	50182.00		1.802			2 $\frac{1}{2}$	64167.84		35.25	1.012		
		2 $\frac{1}{2}$	50577.36		1.597			3 $\frac{1}{2}$	64203.09		123.31	1.208		
		3 $\frac{1}{2}$	50908.54		1.532			4 $\frac{1}{2}$	64326.40		1.258			
4d <sup>4</sup> (a <sup>4</sup> D)5p	z <sup>4</sup> F <sup>o</sup>	1 $\frac{1}{2}$	51372.90	359.49	0.412	4d <sup>4</sup> (a <sup>4</sup> H)5p	z <sup>4</sup> H <sup>o</sup>	4 $\frac{1}{2}$	64130.22	944.49	0.934			
		2 $\frac{1}{2}$	51732.39		1.045			5 $\frac{1}{2}$	65074.71		1.112			
		3 $\frac{1}{2}$	52217.30		1.262			4d <sup>4</sup> (a <sup>4</sup> P)5p	z <sup>4</sup> S <sup>o</sup>		1 $\frac{1}{2}$	64750.66	1.926	
		4 $\frac{1}{2}$	52843.10		1.362						4d <sup>4</sup> (a <sup>4</sup> F)5p	y <sup>4</sup> G <sup>o</sup>	3 $\frac{1}{2}$	64822.22
4d <sup>4</sup> (a <sup>4</sup> D)5p	z <sup>4</sup> D <sup>o</sup>	0 $\frac{1}{2}$	54238.80	448.81	0.042	4d <sup>4</sup> (a <sup>4</sup> F)5p	y <sup>4</sup> F <sup>o</sup>	4 $\frac{1}{2}$	65094.91	842.69			1.105	
		1 $\frac{1}{2}$	54637.61		1.197			3 $\frac{1}{2}$	65260.95		-11.82	0.850		
		2 $\frac{1}{2}$	55215.85		528.24			2 $\frac{1}{2}$	65272.77		0.850			

## Mo II—Continued

## Mo II—Continued

Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	
4d <sup>a</sup> (a <sup>o</sup> G)5p	y <sup>o</sup> H <sup>o</sup>	4½	65222.53	142.07	1.175	4d <sup>a</sup> (a <sup>o</sup> S)5p	z <sup>o</sup> P <sup>o</sup>	0½	71926.30	1252.67	0.670	
		5½	65424.65					1½	73218.97			1.082
4d <sup>a</sup> (a <sup>o</sup> P)5p	y <sup>o</sup> D <sup>o</sup>	1½	65444.30	638.01	0.769	4d <sup>a</sup> (a <sup>o</sup> D)5p	w <sup>o</sup> D <sup>o</sup>	1½	72039.00	790.62	1.086	
		2½	66022.31					2½	72829.62			1.156
4d <sup>a</sup> (a <sup>o</sup> G)5p	z <sup>o</sup> G <sup>o</sup>	2½	65732.22	355.22	0.751	4d <sup>a</sup> (a <sup>o</sup> D)5p	v <sup>o</sup> F <sup>o</sup>	2½	72424.19	548.09		
		3½	66027.55					3½	73032.28			
		4½	66321.46					0½	73546.75			
		5½	66742.72						1½			74050.37
		3°	65821.24						1.209			502.62
4d <sup>a</sup> (a <sup>o</sup> D)5p	w <sup>o</sup> D <sup>o</sup>	0½	66272.65	25.79	1.075	4d <sup>a</sup> (a <sup>o</sup> F)5p	u <sup>o</sup> F <sup>o</sup>	2½	74148.50	245.04		
		1½	66329.44					3½	74421.54			
		2½	66627.28					3½	74749.55			
		3½	66716.24						2½			74928.17
4d <sup>a</sup> (a <sup>o</sup> G)5p	z <sup>o</sup> F <sup>o</sup>	3½	67321.42	-266.71	1.120	4d <sup>a</sup> (b <sup>o</sup> F)5p	w <sup>o</sup> G <sup>o</sup>	2½	74252.86	975.51		
		2½	67622.12					3½	75234.37			750.23
4d <sup>a</sup> (a <sup>o</sup> D)5p	z <sup>o</sup> P <sup>o</sup>	2½	67712.22	-728.74	1.426	4d <sup>a</sup> (b <sup>o</sup> F)5p	w <sup>o</sup> G <sup>o</sup>	4½	76524.60	1079.32		
		1½	68441.66					5½	77623.22			
		0½	69042.75					1.525	2.410			
4d <sup>a</sup> (a <sup>o</sup> G)5p	z <sup>o</sup> G <sup>o</sup>	3½	67760.30	292.17	0.921	4d <sup>a</sup> (a <sup>o</sup> F)5p	v <sup>o</sup> G <sup>o</sup>	3½	75622.65	1.140		
		4½	68022.47					4½				
4d <sup>a</sup> (a <sup>o</sup> D)5p	w <sup>o</sup> F <sup>o</sup>	1½	67821.60	194.32	1.250	4d <sup>a</sup> (b <sup>o</sup> F)5p	v <sup>o</sup> F <sup>o</sup>	1½	75810.627	856.41	-29.65	
		2½	68015.22					3½	76627.02			
		3½	68179.70					4½	76627.22			
		4½	68322.50									
4d <sup>a</sup> (a <sup>o</sup> I)5p	z <sup>o</sup> K <sup>o</sup>	6½	67822.25	1158.60	0.933	4d <sup>a</sup> (b <sup>o</sup> P)5p	v <sup>o</sup> D <sup>o</sup>	1½	75819.20			
		7½	69047.45					2½				
4d <sup>a</sup> (a <sup>o</sup> I)5p	y <sup>o</sup> I <sup>o</sup>	5½	68472.24	425.76		4d <sup>a</sup> (b <sup>o</sup> P)5p	v <sup>o</sup> D <sup>o</sup>	0½				
		6½	68908.60					1½				
4d <sup>a</sup> (a <sup>o</sup> D)5p	y <sup>o</sup> P <sup>o</sup>	0½	68645.50	257.30	0.768	4d <sup>a</sup> (a <sup>o</sup> F)5p	u <sup>o</sup> D <sup>o</sup>	1½	76206.57	118.43		
		1½	68902.20					2½	76325.00			
4d <sup>a</sup> (a <sup>o</sup> D)5p	w <sup>o</sup> F <sup>o</sup>	2½	69722.20	271.65	0.885	4d <sup>a</sup> (b <sup>o</sup> F)5p	u <sup>o</sup> G <sup>o</sup>	3½	81522.27	1029.93		
		3½	70101.55					4½	82556.20			
4d <sup>a</sup> (a <sup>o</sup> G)5p	z <sup>o</sup> H <sup>o</sup>	4½	70022.24	666.72		4d <sup>a</sup> (b <sup>o</sup> G)5p	t <sup>o</sup> G <sup>o</sup>	4½	82214.22	-559.27		
		5½	70270.57					2½	82274.15			
4d <sup>a</sup> (a <sup>o</sup> D)5p	z <sup>o</sup> D <sup>o</sup>	2½	70712.10	-24.90	0.946	4d <sup>a</sup> (b <sup>o</sup> G)5p	v <sup>o</sup> H <sup>o</sup>	4½	82224.22	531.62		
		1½	70722.20					5½	84422.50			
4d <sup>a</sup> (a <sup>o</sup> G)5p	w <sup>o</sup> G <sup>o</sup>	4½	71011.20	-182.33		4d <sup>a</sup> (b <sup>o</sup> G)5p	z <sup>o</sup> F <sup>o</sup>	3½	82222.24			
		3½	71122.22					2½				
4d <sup>a</sup> (a <sup>o</sup> I)5p	w <sup>o</sup> H <sup>o</sup>	5½	71542.22	-372.66								
		4½	71220.22			Mo III(D <sub>2</sub> )	Limit	---	122220			

February 1958.

Mo II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> +	Observed Terms	
4d <sup>2</sup>	$\left\{ \begin{array}{l} a^1G \\ a^1G \end{array} \right. \quad a^1P \quad \left\{ \begin{array}{l} a^1D \\ a^1D \\ b^1D \end{array} \right. \quad \left\{ \begin{array}{l} a^1F \\ a^1F \\ b^1F \end{array} \right. \quad \left\{ \begin{array}{l} a^1G \\ a^1G \\ b^1G \end{array} \right. \quad a^1H \quad a^1I$	
	no (n ≥ 5)	np (n ≥ 5)
4d <sup>2</sup> (e <sup>1</sup> D) <sub>ns</sub>	$\left\{ \begin{array}{l} a^1D \\ b^1D \end{array} \right.$	$\left\{ \begin{array}{l} z^1P^o \\ z^1P^o \end{array} \right. \quad \left\{ \begin{array}{l} z^1D^o \\ z^1D^o \end{array} \right. \quad \left\{ \begin{array}{l} z^1F^o \\ z^1F^o \end{array} \right.$
4d <sup>2</sup> (e <sup>1</sup> H) <sub>ns</sub>		$\left\{ \begin{array}{l} y^1G^o \\ z^1G^o \end{array} \right. \quad \left\{ \begin{array}{l} z^1H^o \\ z^1H^o \end{array} \right. \quad \left\{ \begin{array}{l} z^1I^o \\ z^1I^o \end{array} \right.$
4d <sup>2</sup> (e <sup>1</sup> F) <sub>ns</sub>	$\left\{ \begin{array}{l} b^1P \\ a^1P \end{array} \right.$	$\left\{ \begin{array}{l} z^1G^o \\ z^1G^o \end{array} \right. \quad y^1P^o \quad \left\{ \begin{array}{l} y^1D^o \\ y^1D^o \end{array} \right.$
4d <sup>2</sup> (e <sup>1</sup> F) <sub>ns</sub>	$\left\{ \begin{array}{l} b^1P \\ a^1P \end{array} \right.$	$\left\{ \begin{array}{l} z^1D^o \\ z^1D^o \end{array} \right. \quad y^1P^o \quad \left\{ \begin{array}{l} z^1G^o \\ y^1G^o \end{array} \right.$
4d <sup>2</sup> (e <sup>1</sup> G) <sub>ns</sub>		$\left\{ \begin{array}{l} z^1P^o \\ z^1P^o \end{array} \right. \quad \left\{ \begin{array}{l} z^1G^o \\ z^1G^o \end{array} \right. \quad y^1H^o$
4d <sup>2</sup> (e <sup>1</sup> G) <sub>ns</sub>		$\left\{ \begin{array}{l} z^1P^o \\ z^1P^o \end{array} \right. \quad w^1G^o \quad z^1H^o$
4d <sup>2</sup> (e <sup>1</sup> D) <sub>ns</sub>	$\left\{ \begin{array}{l} a^1D \\ d^1D \end{array} \right.$	$\left\{ \begin{array}{l} z^1P^o \\ y^1P^o \end{array} \right. \quad w^1D^o \quad w^1P^o$
4d <sup>2</sup> (e <sup>1</sup> I) <sub>ns</sub>		$w^1H^o \quad y^1I^o \quad z^1K^o$
4d <sup>2</sup> (e <sup>1</sup> D) <sub>ns</sub>	$a^1D$	$w^1P^o \quad z^1D^o \quad a^1F^o$
4d <sup>2</sup> (e <sup>1</sup> G) <sub>ns</sub>	$b^1G$	$z^1P^o$
4d <sup>2</sup> (e <sup>1</sup> F) <sub>ns</sub>	$d^1F$	$w^1D^o \quad u^1P^o \quad a^1G^o$
4d <sup>2</sup> (b <sup>1</sup> F) <sub>ns</sub>	$a^1F$	$\left\{ \begin{array}{l} y^1P^o \\ z^1P^o \end{array} \right. \quad w^1G^o$
4d <sup>2</sup> (b <sup>1</sup> F) <sub>ns</sub>		$\left\{ \begin{array}{l} y^1P^o \\ z^1P^o \end{array} \right. \quad u^1G^o$
4d <sup>2</sup> (b <sup>1</sup> F) <sub>ns</sub>		$y^1D^o$
4d <sup>2</sup> (b <sup>1</sup> G) <sub>ns</sub>		$\left\{ \begin{array}{l} z^1D^o \\ z^1D^o \end{array} \right. \quad \left\{ \begin{array}{l} z^1F^o \\ z^1F^o \end{array} \right. \quad \left\{ \begin{array}{l} z^1G^o \\ z^1G^o \end{array} \right. \quad a^1H^o$

\*For predicted terms of the Nb I isoelectronic sequence, see Vol. II, Introduction, p. XXVI.

## Mo III

(Zr : sequence; 40 electrons)

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^4 {}^4D_3$  $4d^4 {}^4D_3$ , 218000 K

I. P. 27.13 volts

The analysis is from Rico, who has classified 37 lines between 2142 Å and 2474 Å, from observations made at Madrid for the region 2200 Å to 3200 Å. These have been supplemented by spectrograms taken at Princeton covering the range 670 Å and 2200 Å.

The ground term has not been found. Combinations from this term are estimated to lie between 1250 Å and 1330 Å. Rico has classified 4 lines between 1274 Å and 1288 Å as representing transitions from  $4d^4 {}^4D_3$ , which he estimates to be approximately 1500 K above the ground state zero,  $4d^4 {}^4D_3$ . He and Catalán have interpolated the limit by comparison of the third spectra from Y to In.

All of the observed energy levels are relative to this estimated value of 1500 K entered in brackets in the table. Further analysis is necessary to establish more accurate values derived from observed combinations from the ground state zero. The observed terms have been found by comparison with similar spectra in the isoelectronic sequence and with neighboring third spectra.

Rao also reports 27 classified lines of Mo III involving the quintet terms from the  ${}^4F$  limit in Mo IV. The two papers have only a few term intervals and lines in common, chiefly in the  ${}^4F$ - ${}^4G^o$  multiplet.

## REFERENCES

V. R. Rao, Indian J. Phys. **23**, 258 (1949). (T) (C L)F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] **50**, 185 (1954). (T) (C L)M. A. Catalán y F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] **53**, 83 (1957). (I P)

## Mo III

## Mo III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^4$	$4d^4 {}^4D$	0			$4d^4 ({}^4F) 5p$	$5p {}^4D^o$	0	75083.07+z	536.27
		1					1	75599.2+z	864.8
		2					2	76164.0+z	1433.4
		3					3	77897.4+z	1226.8
		4	[1500]				4	79194.2+z	
$4d^4 ({}^4F) 5s$	$5s {}^4F$	1	22045.7+z	424.5	$4d^4 ({}^4F) 5p$	$5p {}^4F^o$	1	77361.1+z	323.8
		2	22470.2+z	609.1			2	77684.4+z	1451.2
		3	23079.3+z	772.9			3	79135.6+z	587.1
		4	23852.2+z	904.1			4	79733.7+z	247.5
		5	24756.3+z				5	79970.2+z	
$4d^4 ({}^4F) 5p$	$5p {}^4G^o$	2	78490.2+z	871.5	-----	-----	-----	-----	-----
		3	74351.7+z	1091.9	Mo IV ( ${}^4F_{10}$ )	Limit	--	218000	
		4	75443.6+z	1296.6					
		5	76740.2+z	1576.2					
		6	78316.4+z						

February 1957.

## Mo IV

(Y I sequence; 39 electrons)

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^5 \text{ } ^6F_{1/2}$  $4d^5 \text{ } ^6F_{1/2}$  374180 K

I. P. 46.4 volts

Elison has classified some 40 lines between 856.00 Å and 2140.98 Å. He has derived the limit by a linear extrapolation of isoelectronic sequence data. This value in round figures is entered in brackets in the table.

## REFERENCE

A. Y. Elison, Phys. Rev. 43, 745 (1933). (I F) (T) (C L)

## Mo IV

## Mo IV

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^5$	$4d^5 \text{ } ^6F$	1½	0.0		$4d^5 \text{ } ^6F$ 5p	$5p \text{ } ^6F^o$	1½	111756.5	
		2½	780.0	780.0			2½	112981.0	1164.5
		3½	1759.0	979.0			3½	114619.4	1698.4
		4½	2888.6	1099.6			4½	115958.7	1337.3
$4d^5 \text{ } ^6F$ 5s	$5s \text{ } ^6F$	1½	80893.4		$4d^5 \text{ } ^6F$ 5p	$5p \text{ } ^6D^o$	0½	115790.4	
		2½	81634.1	730.7			1½	116584.3	793.9
		3½	83708.3	1081.2			2½	117608.3	1018.0
		4½	84042.6	1337.3			3½	118078.5	474.2
$4d^5 \text{ } ^6F$ 5p	$5p \text{ } ^6G^o$	2½	109412.0	1923.7	Mo v ( $^6F_2$ )	Limit			
		3½	111336.6	2102.7					
		4½	113439.5	2437.1					
		5½	115876.4						[374180]

April 1951.



## Mo v

(Sr I sequence; 38 electrons)

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^9 \text{ } ^4\text{F}_7$  $4s^2 \text{ } ^4\text{F}_7$ , 493300 K

I. P. 61.2 volts

The analysis is by Trawick, who has classified 90 lines between 410.35 Å and 2159.30 Å. He has extrapolated the limit on a Moseley diagram of isoelectronic sequence data. This value of the limit, rounded off, is entered in brackets in the table.

The singlet and triplet terms are connected by observed intersystem combinations.

## REFERENCE

M. W. Trawick, Phys. Rev. 68, 233 (1935). (I P) (T) (C L)

Mo v					Mo v				
Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
$4d^9$	$4d^9 \text{ } ^4\text{F}$	2	0	1585	$4d(^4\text{D})5p$	$5p \text{ } ^4\text{P}^{\circ}$	0	157000	-444 1236
		3	1585	1			160016		
		4	3350	2			157858		
$4d^9$	$4d^9 \text{ } ^4\text{P}$	0	11165	647	$4d(^4\text{D})5d$	$5d \text{ } ^1\text{D}$	3	212020	-831 -115
		1	11812	1601			3 2	218940	
		2	12413	214671					
$4d(^4\text{D})5s$	$5s \text{ } ^1\text{D}$	2	83971	732 1724	$4d(^4\text{D})5d$	$5d \text{ } ^3\text{D}$	3	214786	215874
		1	92381				3	214671	
			2					93113	
$4d(^4\text{D})5s$	$5s \text{ } ^3\text{D}$	1	92381	732 1724	$4d(^4\text{D})5d$	$5d \text{ } ^1\text{P}$	1	215874	3162 6226
		2	93113				2 3	236714	
		3	94837					239071	
$4d(^4\text{D})5p$	$5p \text{ } ^1\text{P}^{\circ}$	3	141150	2262 1820	$4d(^4\text{D})4f$	$4f \text{ } ^1\text{F}^{\circ}$	2	241794	2357 3033
		1	147909				3 4	244296	
			2					148950	
$4d(^4\text{D})5p$	$5p \text{ } ^3\text{P}^{\circ}$	1	147909	2262 1820	$4d(^4\text{D})5d$	$5d \text{ } ^3\text{F}$	2	236714	1170 900
		2	151818				3 4	239071	
		3	153041					242104	
$4d(^4\text{D})5p$	$5p \text{ } ^3\text{D}^{\circ}$	1	148950	2262 1820	$4d(^4\text{D})4f$	$4f \text{ } ^3\text{G}^{\circ}$	3	243665	1170 900
		2	151818				4 5	244841	
		3	153041					245530	
$4d(^4\text{D})5p$	$5p \text{ } ^1\text{D}^{\circ}$	2	151760		Mo vi( $^4\text{D}_{3/2}$ )	Limit		[493300]	

April 1951.

## Mo v OBSERVED TERMS\*

Configuration $1s^2 2s^2 2p^6 3s^2$ $3p^6 3d^{10} 4s^2 4p^6 +$	Observed Terms									
$4d^9$	$4d^9 \text{ } ^4\text{P}$		$4d^9 \text{ } ^4\text{F}$							
	$ns (n \geq 5)$		$np (n \geq 5)$		$nd (n \geq 5)$		$nf (n \geq 4)$			
	$4d(^4\text{D})ns$	{	$5s \text{ } ^1\text{D}$	$5p \text{ } ^1\text{P}^{\circ}$	$5p \text{ } ^3\text{D}^{\circ}$	$5p \text{ } ^1\text{F}^{\circ}$	$5d \text{ } ^1\text{P}$	$5d \text{ } ^1\text{D}$	$5d \text{ } ^3\text{F}$	$4f \text{ } ^1\text{F}^{\circ}$
$5s \text{ } ^3\text{D}$			$5p \text{ } ^3\text{P}^{\circ}$	$5p \text{ } ^3\text{D}^{\circ}$	$5p \text{ } ^3\text{F}^{\circ}$	$5d \text{ } ^3\text{P}$	$5d \text{ } ^3\text{D}$	$5d \text{ } ^3\text{F}$	$4f \text{ } ^3\text{F}^{\circ}$	$4f \text{ } ^3\text{G}^{\circ}$

\*For predicted terms in the spectra of the Sr I isoelectronic sequence, see Vol. II Introduction, page XXIII.

## Mo VI

(Eh I sequence; 37 electrons)

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^2 D_{3/2}$  $4d^2 D_{3/2}$  549000 K

I. P. 68 volts

Sixteen lines have been classified as combinations among seven terms. The observations extend from 296 Å to 1595 Å. Trawick has published six terms and extrapolated a limit from isoelectronic sequence data. From recent series data in Zr IV combined with the series of Mo VI, Kiess has derived an improved value of the limit, which is quoted here.

Charles has recently reobserved the spectrum between 296 Å and 790 Å and confirmed three of the four terms from Trawick's list, covered by this range. The one exception is  $4f^2 F^o$ , which he revises. Trawick lists this term as follows:

Desig.	J	Level	Interval
$4f^2 F^o$	$2\frac{1}{2}$	267451	404
	$3\frac{1}{2}$	267355	

Charles also resolves the members of his  $4d^2 D - 4f^2 F^o$  group, and adds tentative values of the  $6p^2 P^o$  term.

The measurements common to the two lists are in satisfactory agreement. In the table the term values from Charles' paper are quoted, but the two terms  $5s^2 S$  and  $5d^2 D$  are from Trawick's paper.

## REFERENCES

M. W. Trawick, Phys. Rev. 46, 63 (1934). (I P) (T) (C L)

G. W. Charles, Phys. Rev. 77, 120 (1950). (T) (C L)

C. C. Kiess, unpublished material (October 1955). (I P)

## Mo VI

## Mo VI

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4p^6(^1S)4d$	$4d^2 D$	$1\frac{1}{2}$	0	2578	$4p^6(^1S)5d$	$5d^2 D$	$1\frac{1}{2}$	282837	787
		$2\frac{1}{2}$	2578				$2\frac{1}{2}$	283624	
$4p^6(^1S)5s$	$5s^2 S$	$0\frac{1}{2}$	119739		$4p^6(^1S)6s$	$6s^2 S$	$0\frac{1}{2}$	313810	
$4p^6(^1S)5p$	$5p^2 P^o$	$0\frac{1}{2}$	182404	4925	$4p^6(^1S)6p$	$6p^2 P^o$	$0\frac{1}{2}?$	336331	1408
		$1\frac{1}{2}$	187389				$1\frac{1}{2}?$	337739	
$4p^6(^1S)4f$	$4f^2 F^o$	$2\frac{1}{2}$	267045	408	-----		-----	-----	-----
		$3\frac{1}{2}$	267451		Mo VII( $^1S_0$ )	Limit	-----	[549000]	

October 1955.

## Mo VII

(Kr I sequence; 36 electrons)

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 {}^1S_0$  $4p^6 {}^1S_0$ , 1020460 K

I. P. 126 volts

The analysis is far from complete. Charles has classified 6 lines between 136 Å and 286 Å as due to transitions from the ground term to levels produced by 4d, 5d, 5s, and 6s-electrons. The level-values in the table are the rounded-off wave numbers of the observed lines. He has estimated the limit quoted above by applying a Rydberg formula to the *ns s<sub>1</sub>* levels (*n*=5, 6). The higher limit in the table has been determined by the writer by adding the interval of the ground term of Mo VIII to the lower limit.

As for Kr I the writer has introduced the *Jl*-coupling notation in the general form suggested by Racah. Charles has noted that the 4d and 5d assignments are open to some question. Consequently the pair-coupling notation for these levels may need revision.

## REFERENCES

G. Racah, Phys. Rev. 61, 537 (L) (1942).

G. W. Charles, Phys. Rev. 77, 120 (1950). (I P) (T) (C L)

## Mo VII

## Mo VII

Author	Config.	Desig.	<i>J</i>	Level	Author	Config.	Desig.	<i>J</i>	Level
$4p^6 p_0$	$4p^6$	$4p^6 {}^1S$	0	0	$6s s_1$	$4p^6 ({}^2P_{1/2}) 6s$	$6s [1\frac{1}{2}]^o$	2	
$4d d_1$	$4p^6 ({}^2P_{1/2}) 4d$	$4d [1\frac{1}{2}]^o$	2	349290	$6s s_1$	$4p^6 ({}^2P_{3/2}) 6s$	$6s' [0\frac{1}{2}]^o$	0	709450
$5s s_1$	$4p^6 ({}^2P_{1/2}) 5s$	$5s [1\frac{1}{2}]^o$	2	481290		Mo VIII ( ${}^2P_{1/2}$ )	Limit	-----	1020460
$5s s_1$	$4p^6 ({}^2P_{3/2}) 5s$	$5s' [0\frac{1}{2}]^o$	0	502990		Mo VIII ( ${}^2P_{3/2}$ )	Limit	-----	1043740
$5d d_1?$	$4p^6 ({}^2P_{1/2}) 5d$	$5d [1\frac{1}{2}]^o$	2	668280					

February 1951.

## Mo VIII

(Br I sequence; 35 electrons)

Z=42

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^5 {}^2P_{1/2}^o$  $4p^5 {}^2P_{1/2}^o$ , 1235000 K

I. P. 153 volts

The analysis is incomplete. Charles has classified 42 lines between 168 Å and 474 Å as due to transitions from the ground term to 14 higher terms.

He has extrapolated the value of the limit from isoelectronic sequence data.

## REFERENCE

G. W. Charles, Phys. Rev. 77, 120 (1950). (I P) (T) (C L)

## Mo VIII

## Mo VIII

Config.	Desig.	<i>J</i>	Level	Interval	Config.	Desig.	<i>J</i>	Level	Interval
4s <sup>2</sup> 4p <sup>2</sup>	4p <sup>2</sup> <sup>3</sup> P <sup>o</sup>	1½ 0½	0 53877	-23277	4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 4d	4d <sup>3</sup> F	3½ 2½	409395	
4s 4p <sup>3</sup>	4p <sup>3</sup> <sup>2</sup> S	0½	233830		4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> D) 4d	4d' <sup>1</sup> D	2½ 1½	423722 444240	-20618
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 4d	4d <sup>4</sup> D	3½ 2½ 1½ 0½?	301557 310419 319456	-8862 -9037	4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> D) 4d	4d' <sup>3</sup> P	1½ 0½	445430	
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 4d	4d <sup>4</sup> P	0½ 1½ 2½	363255 371345 377888	8090 6543	4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> S) 4d	4d'' <sup>1</sup> D	2½? 1½?	447888 454307	-6419
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 4d	4d <sup>4</sup> F	4½ 3½ 2½ 1½	367822 367967	-145	4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 5s	5s <sup>4</sup> P	2½ 1½? 0½	492522 503309	-10787
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 4d	4d <sup>3</sup> D	1½ 2½	376122 391179	15057	4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 5s	5s <sup>3</sup> P	1½ 0½	527428 543352	-15924
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 4d	4d <sup>3</sup> P	0½ 1½?	381102		4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> D) 5s	5s' <sup>3</sup> D	2½ 1½	586550 548916	-12366
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) 4d	4d <sup>3</sup> P	0½ 1½?	381102		4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> S) 5s	5s'' <sup>2</sup> S	0½	595062	
					Mo IX ( <sup>3</sup> P <sub>2</sub> )	Limit	-----	[1235000]	

February 1951.

## Mo VIII OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> +	Observed Terms				
4s <sup>2</sup> 4p <sup>2</sup>	4p <sup>2</sup> <sup>3</sup> P <sup>o</sup>				
4s 4p <sup>3</sup>	4p <sup>3</sup> <sup>2</sup> S				
	ns (n ≥ 5)		nd (n ≥ 4)		
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>3</sup> P) n <sub>z</sub>	{	5s <sup>4</sup> P	4d <sup>4</sup> P	4d <sup>4</sup> D	4d <sup>4</sup> F
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> D) n <sub>z</sub> '		5s <sup>3</sup> P	4d <sup>3</sup> P	4d <sup>3</sup> D	4d <sup>3</sup> F
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> D) n <sub>z</sub> '		5s' <sup>3</sup> D	4d' <sup>3</sup> P	4d' <sup>3</sup> D	
4s <sup>2</sup> 4p <sup>4</sup> ( <sup>1</sup> S) n <sub>z</sub> ''		5s'' <sup>2</sup> S		4d'' <sup>1</sup> D	

\*For predicted terms in the spectra of the Br I isoelectronic sequence, see Vol. II, Introduction, page XXI.

## Mo XVI

(Co I sequence; 27 electrons)

Z=42

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>9</sup> <sup>2</sup>D<sub>3/2</sub>3d<sup>9</sup> <sup>2</sup>D<sub>3/2</sub>

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed three lines due to the transition 3p<sup>6</sup> 3d<sup>9</sup> <sup>2</sup>D - 3p<sup>6</sup> 3d<sup>9</sup> <sup>2</sup>P<sup>o</sup>. In figure 4 of his paper on the spectra of highly-ionized atoms, the observed wave numbers are plotted against atomic number for this combination in the Co I-like spectra Rb XI to Mo XVI. For Mo XVI the wave numbers are between 1300000 and 1500000 K.

## REFERENCE

B. Edlén, *Physica* 13, No. 9, 548 (1947).

February 1950.

## TECHNETIUM

Tc I

43 electrons

Z=43

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^5 5s^2 {}^6S_{5/2}$  $\epsilon$   ${}^6S_{5/2}$  58700 K

I. P. 7.28 volts

A description of the spectrum was published by Meggers and Scribner in 1950. The next year Meggers reported the first regularities: 20 terms, 200 classified lines, and an approximate ionization potential.

The spectrum has since been reobserved by Bozman, Corliss, and Meggers. The present line list covers the range 2057.70 Å to 8917.74 Å. The analysis has been extended by Bozman especially for inclusion here. His work is still in progress; the tabular data represent only the results available when the present Volume was concluded for press. The observed  $g$ -values are, also, from Zeeman spectrograms taken at the National Bureau of Standards.

There are approximately 900 classified lines. Observed intersystem combinations connect the systems of terms of different multiplicity.

Bozman has revised the earlier value of the limit. The present limit is from the  $4d^5(a {}^6D)ns {}^6D$  series ( $n=5,6$ ). It has been derived by a Rydberg formula, with a Ritz correction  $\alpha=2.0 \times 10^{-6}$ , described in the 1951 paper. It agrees well with the value interpolated by Catalán from neighboring spectra, 58297 K.

## REFERENCES

- W. F. Meggers and B. F. Scribner, *J. Research Nat. Bur. Std.* **45**, 476, RP 2161 (1950).  
 W. F. Meggers, *J. Research Nat. Bur. Std.* **47**, 7, RP 2221 (1951). (I P) (T) (C L)  
 M. A. Catalán y F. R. Rico, *An. Real Soc. Esp. Física y Química (Madrid)* [A] **48**, 328 (1952). (I P)  
 K. G. Kessler and R. E. Treca, *Phys. Rev.* (2) **92**, 303 (1953). (hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955).  
 (Summary hfs)  
 W. R. Bozman, unpublished material (June 1957). (I P) (T) (C L) (Z E)

Tc I

Tc I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
4d <sup>5</sup> 5e	a <sup>5</sup> B	2 1/2	0.00		1.99	4d <sup>5</sup> (a <sup>5</sup> D)5p	y <sup>5</sup> P <sup>0</sup>	3 1/2 2 1/2 1 1/2	31414.07 31408.78 31503.96		1.72 1.80 2.26
4d <sup>5</sup> (a <sup>5</sup> D)5e	a <sup>5</sup> D	4 1/2 3 1/2 2 1/2 1 1/2 0 1/2	2572.89 3250.91 3700.55 4002.58 4178.72	-678.02 -449.64 -302.03 -176.14	1.60 1.59 1.65 1.86 3.32	4d <sup>5</sup> (a <sup>5</sup> D)5p	s <sup>5</sup> P <sup>0</sup>	2 1/2 1 1/2 0 1/2	31987.01 32258.07 32403.07	-331.06 -145.00	1.59 1.40 2.65
4d <sup>5</sup> (a <sup>5</sup> D)5e	a <sup>5</sup> D	3 1/2 2 1/2 1 1/2 0 1/2	10516.54 11063.08 11578.59 11890.96	-546.54 -515.51 -312.37	1.44 1.37 1.21 0.02	4d <sup>5</sup> (a <sup>5</sup> D)5p	s <sup>5</sup> D <sup>0</sup>	3 1/2 2 1/2 1 1/2 0 1/2	32690.57 32685.70 33343.17 33480.47	-465.33 -257.47 -137.30	1.39 1.39 1.20 0.02
4d <sup>5</sup> (a <sup>5</sup> P)5e	a <sup>5</sup> P	2 1/2 1 1/2 0 1/2	13252.73 14169.75 14679.92	-917.02 -510.17	1.62 1.70 2.63	4d <sup>5</sup> 5e(a <sup>5</sup> S)5p	y <sup>5</sup> P <sup>0</sup>	2 1/2 1 1/2 0 1/2	34515.90 34948.68 35276.45	-426.76 -333.79	1.61 1.74 2.66
4d <sup>5</sup> (a <sup>5</sup> F)5e	a <sup>5</sup> F	4 1/2 3 1/2 2 1/2 1 1/2	14733.14 15298.47 15624.26 15770.42	-565.33 -325.79 -146.16	1.31 1.20 1.02 0.42	4d <sup>5</sup> 5e(a <sup>5</sup> S)6e	e <sup>5</sup> S	3 1/2	37612.40		2.02
4d <sup>5</sup> (a <sup>5</sup> G)5e	a <sup>5</sup> G	5 1/2 4 1/2 3 1/2 2 1/2	16025.16 16133.99 16267.80 16415.64	-108.83 -153.81 -127.84	1.23 1.17 0.95 0.58	4d <sup>5</sup> 5e(a <sup>5</sup> S)6e	e <sup>5</sup> S	2 1/2	39598.85		2.00
4d <sup>5</sup> 5e(a <sup>5</sup> S)5p	s <sup>5</sup> P <sup>0</sup>	2 1/2 3 1/2 4 1/2	16488.71 16874.51 17582.95		445.80 1.94 1.80	4d <sup>5</sup> (a <sup>5</sup> D)6e	e <sup>5</sup> D	4 1/2 3 1/2 2 1/2 1 1/2 0 1/2	42855.22 43391.75 43796.71 44028.92 44149.51	-536.53 -404.96 -232.21 -120.59	1.56 1.57 1.65 1.87 3.30
4d <sup>5</sup> 5e(a <sup>5</sup> S)5p	s <sup>5</sup> P <sup>0</sup>	3 1/2 2 1/2 1 1/2	23265.33 23455.21 23538.40	-189.88 -133.19	1.70 1.90 2.42	4d <sup>5</sup> (a <sup>5</sup> D)6e	e <sup>5</sup> D	3 1/2 2 1/2 1 1/2 0 1/2	44102.00 44804.36 45267.84 45497.76	-702.36 -463.48 -229.92	
4d <sup>5</sup> (a <sup>5</sup> D)5p	s <sup>5</sup> D <sup>0</sup>	4 1/2 3 1/2 2 1/2 1 1/2 0 1/2	27369.78 27660.09 27940.78 28151.28 28290.69	-290.31 -280.63 -210.56 -145.41	1.53 1.57 1.64 1.85 3.30	4d <sup>5</sup> 5e(a <sup>5</sup> S)5d	e <sup>5</sup> D	1 1/2 2 1/2 3 1/2 4 1/2 5 1/2	44333.23 44337.53 44343.92 44352.66 44365.33	4.30 6.39 8.74 12.67	2.08 1.84 1.67 1.66
4d <sup>5</sup> (a <sup>5</sup> D)5p	s <sup>5</sup> F <sup>0</sup>	5 1/2 4 1/2 3 1/2 2 1/2 1 1/2 0 1/2	30067.89 30133.25 30382.08 30528.85 30630.63 30689.18	-65.96 -248.81 -146.79 -101.78 -58.49	1.50 1.41 1.38 1.29 1.07 -0.62	4d <sup>5</sup> 5e(a <sup>5</sup> S)5d	f <sup>5</sup> D	4 1/2 3 1/2 2 1/2 1 1/2 0 1/2	44919.07 44985.53 45071.02 45108.62 45189.70	-66.46 -85.49 -37.60 -81.08	1.54 1.58 1.66 1.87 3.32
4d <sup>5</sup> (a <sup>5</sup> D)5p	s <sup>5</sup> F <sup>0</sup>	4 1/2 3 1/2 2 1/2 1 1/2	31114.09 31604.99 32014.79 32253.65	-490.90 -409.80 -238.86	1.37 1.32 1.14 0.85	Tc II(7B <sub>2</sub> )	Limit		58700		

## Tc II

(Mo I sequence; 42 electrons)

Z=43

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^5 5s^1 S_2$  $a^1S_2$ , 123100 K

I. P. 15.26 volts

Meggers and Scribner first observed this spectrum in 1950, and Meggers reported the first regularities in 1951. He classified the 10 leading lines arising from the low  $^1S$  and  $^4S$  terms, and stressed the similarity between the spectra of Mn II and Tc II. In 1952, he and Catalán identified the important low  $^4D$  term from the  $3d^5$  configuration, with its leading component 3461 K above the ground term  $a^1S_2$ .

A new description of the spectrum has since been made by Bozman, Corliss, and Meggers, which covers the range 2054.468 Å to 6673.263 Å. From these observations the writer has carried the analysis somewhat further with the aid of Bureau Zeeman data from Bozman. Unfortunately, the new terms are not connected with the earlier ones because observations are lacking in the region short of 2054 Å. The writer has arbitrarily adopted 24000 K for  $a^4G_5$ , with the correction,  $x$ , to be determined later when the line list has been extended. Trees has predicted the theoretical position of this  $^4G$  term at  $22500 \pm 1000$  K.

There are about 50 classified lines, but the analysis is seriously incomplete. Intersystem combinations connect the terms in the table that need no correction,  $x$ .

No series are known, but Catalán and Rico have interpolated the limit quoted here, by comparison with neighboring second spectra in the second long period.

## REFERENCES

- W. F. Meggers and B. F. Scribner, J. Research Nat. Bur. Std. **45**, 476, RP 2161 (1950).  
 W. F. Meggers, J. Research Nat. Bur. Std. **47**, 7, RP 2221 (1951). (T) (C L)  
 W. F. Meggers and M. A. Catalán, unpublished material (1952). (T) (C L)  
 M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. Física. y Química (Madrid) [A] **48**, 334 (1952) and letter (December 1956). (I P)  
 R. E. Trees, unpublished material (1956). (T)  
 W. R. Bozman, unpublished material (1956). (Z E)  
 C. E. Moore, unpublished material (June 1957). (T) (C L)

## Tc II

## Tc II

Config.	Desig.	J	Level	Interval	Obs. $g$	Config.	Desig.	J	Level	Interval	Obs. $g$
$4d^5(^8S)5s$	$a^1S$	3	0.00		2.02	$4d^5(^4G)5p$	$z^4G^o$	2	$59034.69+x$	193.06	0.42
$4d^5$	$a^4D$	4	3461.27		1.47			3	$59227.75+x$	184.45	0.90
		3	4217.17	-755.90	1.50			4	$59412.20+x$	168.98	1.16
		2	4669.22	-452.05	1.52			5	$59581.18+x$	176.47	1.26
		1	4961.14	-291.92	1.56			6	$59757.65+x$		1.36
		0	5100.98	-139.84	0/0	$4d^5(^4G)5p$	$z^4H^o$	3	$60535.64+x$	59.14	0.54
$4d^5(^8S)5s$	$a^4S$	2	12617.20		2.04			4	$60594.78+x$	648.74	0.97
$4d^5(^4G)5s$	$a^4G$	2	$23838.39+x$	78.78	0.37			5	$61243.52+x$	472.53	1.24
		3	$23917.17+x$	61.48	0.89			6	$61718.05+x$	224.61	1.31
		4	$23978.65+x$	46.23	1.15	$4d^5(^4G)5p$	$z^4F^o$	1	$60707.68+x$	321.56	0.04
		5	$24024.88+x$	-24.88	1.30			2	$61029.24+x$	318.44	1.15
		6	$24000.00+x$		1.32			3	$61347.68+x$	71.47	1.34
$4d^5(^8S)5p$	$z^1P^o$	2	37767.21		2.36			4	$61419.15+x$	383.93	1.41
		3	38302.80	535.59	1.88			5	$61803.08+x$		
		4	39508.33	1005.58	1.78						
$4d^5(^8S)5p$	$z^3P^o$	3	43500.96		1.65	Tc III ( $^8S_{3/2}$ )	Limit		123100		
		2	43741.33	-240.37	1.86						
		1	43905.00	-163.67	2.50						

June 1957.

## RUTHENIUM

## Ru I

44 electrons

Z=44

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2 4p^6 4d^7 5s^1 {}^5F_3$  $\alpha$   ${}^5F_3$ , 59410 K

I. P. 7.364 volts

The first significant regularities in the Ru I spectrum were reported in 1925 by Meggers and Laporte and by Sommer. In 1926 relative term values confirmed by observed  $g$ -values were published by these authors. Harrison and McNally derived improved  $g$ -values from Zeeman observations made at the Massachusetts Institute of Technology, and extended the analysis in 1940. Later, McNally carried this work further. His unpublished 1941 manuscript includes 61 even and 188 odd levels, and has been made available to the National Bureau of Standards for further study of Ru I.

Kessler and Meggers reobserved the spectrum, and in 1955 published a complete new description which Kessler has used to extend the analysis. His manuscript has been furnished in advance of publication, and used for the data in the table. He has adjusted all level values to fit the 1955 observations. Most of the observed  $g$ -values in the table are from the 1940 and 1941 work of Harrison and McNally. A few additional ones are from Zeeman spectrograms taken in 1949 by Meggers with the Bitter magnet at the Massachusetts Institute of Technology, and measured by Kessler.

At present 100 even and 205 odd levels are known, and there are approximately 3250 classified lines between 2013.95 Å and 11483.91 Å. Many observed intersystem combinations connect the systems of terms of different multiplicity.

The more dubious configuration assignments are indicated by a colon in the first column of the table. Members of fragmentary terms have a designation entered in parentheses in column two, along with a number labeling the level as miscellaneous. Extensive theoretical study of Ru I by Trees confirms in detail the observed values of the low even terms, and has been an invaluable guide in making a number of the configuration assignments for additional terms.

The limit quoted above has been derived by Kessler from the two-member series  $4d^7(\alpha {}^5F)ns {}^5F$ , ( $n=5,6$ ), by means of a Rydberg formula, corrected by the factor  $\alpha = -2.43 \times 10^{-6}$  suggested by Catalán and Rico. They have obtained this Ritz correction from a study of the series in the first spectra from Rb I through Ag I. The observed limit thus corrected agrees excellently with the interpolated value 59417 obtained by Catalán and Rico.

## REFERENCES

- W. F. Meggers and O. Laporte, *J. Wash. Acad. Sci.* **16**, 143 (1926). (T) (C L) (Z E)  
 L. A. Sommer, *Zeit. Phys.* **37**, 1 (1926). (I P) (T) (C L) (Z E)  
 G. R. Harrison and J. R. McNally, Jr., *Phys. Rev.* **56**, 703 (1940). (T) (C L) (Z E)  
 J. R. McNally, Jr., *Thesis*, Mass. Inst. Tech., unpublished (June 1941). (T) (C L) (Z E)  
 M. A. Catalán y F. R. Rico, *An. Real. Soc. Esp. Física y Química (Madrid)* [A] **48**, 328 (1952). (I P)  
 K. G. Kessler and W. F. Meggers, *J. Research Nat. Bur. Std.* **55**, 97, RP 2609 (1955).  
 K. Murakawa, *J. Phys. Soc. Japan* **10**, 919 (1955). (hfs)  
 R. E. Trees, unpublished material (February 1957).  
 K. G. Kessler, unpublished material (February 1957). (I P) (T) (C L) (Z E)





## Ru I—Continued

## Ru I—Continued

Config.	Design.	J	Level	Interval	Obs. $\rho$	Config.	Design.	J	Level	Interval	Obs. $\rho$				
4d <sup>7</sup> (a <sup>2</sup> G)5p	y <sup>2</sup> F°	4	38243.38		1.107	4d <sup>7</sup> (a <sup>2</sup> H)5p	y <sup>2</sup> H°	6	43548.87		1.162				
		3	39433.70	-1190.32	0.968			5	44109.41	-560.74	1.083				
		2	40433.23	-999.53	0.889			4	44662.01	-552.60	0.928				
4d <sup>7</sup> (a <sup>2</sup> G)5p	z <sup>2</sup> H°	6	38897.50		1.174	4d <sup>7</sup> (a <sup>2</sup> H)5p	y <sup>2</sup> H°	5	43596.58			1.08			
		5	38997.09	600.41	1.048			w <sup>2</sup> C°	5	43742.81		1.272			
		4	39273.28	-976.19	0.895				4	43862.91	-120.10				
4d <sup>7</sup> (a <sup>2</sup> F)5p	z <sup>2</sup> S°	1	38587.14		1.566										
4d <sup>7</sup> (a <sup>2</sup> F)5p	y <sup>2</sup> P°	3	38708.36		1.631	4d <sup>7</sup> (a <sup>2</sup> P)5p	y <sup>2</sup> D°	2	43903.41			1.026			
		2	39008.63	-302.26	1.713			z <sup>2</sup> (C°)	6	43998.60		1.219			
		1	39773.49	-764.87	2.315										
4d <sup>7</sup> (a <sup>2</sup> G)5p	z <sup>2</sup> G°	4	39037.18		1.115	4d <sup>7</sup> (a <sup>2</sup> D)5p	z <sup>2</sup> P°	2	44234.68		1.422				
4d <sup>7</sup> (a <sup>2</sup> G)5p	y <sup>2</sup> G°	5	39450.66		1.142	4d <sup>7</sup> 5s(a <sup>2</sup> D)5p	w <sup>2</sup> D°	4	44243.49			1.473			
		4	40276.61	-825.95	1.035			3	45071.40	-827.91	1.449				
		3	40235.39	41.22	0.890			2	45790.41	-719.01	1.484				
4d <sup>7</sup> (a <sup>2</sup> P)5p	y <sup>2</sup> P°	2	39742.03		1.299			1	46191.40	-400.99	1.439				
		1	39916.54	-174.51	1.606			0	46466.35	-274.95					
		0	39894.50	22.04	0/0										
	1°(2F°)	4	40439.25		1.196		3°(2P°)	1	44301.14		1.350				
4d <sup>7</sup> (a <sup>2</sup> G)5p	z <sup>2</sup> H°	5	40816.22		1.020	4d <sup>7</sup> 5s(a <sup>2</sup> D)5p	z <sup>2</sup> F°	5	44321.81		1.308				
4d <sup>7</sup> (a <sup>2</sup> P)5p	z <sup>2</sup> D°	3	40768.15		1.159			4	44807.61	-285.80					
		2	42007.26	-1239.11	1.007			3	44800.81	-193.20					
		1	41016.65	990.61	0.895			2	45592.33	-791.52					
4d <sup>7</sup> (a <sup>2</sup> G)5p	z <sup>2</sup> F°	3	40948.65		1.137		4°	3	44441.59		0.76				
4d <sup>7</sup> (a <sup>2</sup> D)5p	z <sup>2</sup> F°	2	41182.94		0.887	4d <sup>7</sup> 5s(b <sup>2</sup> P)5p	y <sup>2</sup> S°	2	44891.40		0.383				
		3	41260.04	77.10	1.235			4d <sup>7</sup> (a <sup>2</sup> F)5p:	z <sup>2</sup> F°	3	45201.98		1.059		
		4	42346.90	1086.86	1.247					4d <sup>7</sup> (a <sup>2</sup> H)5p	y <sup>2</sup> G°	4	45364.72		0.962
4d <sup>7</sup> (a <sup>2</sup> F)6s	e <sup>2</sup> F	5	41256.40			4d <sup>7</sup> (a <sup>2</sup> F)5p:	6°	3	45475.77		1.547				
		4	43018.57	-1762.17				4d <sup>7</sup> (a <sup>2</sup> F)5p:	z <sup>2</sup> G°	4	45528.61		1.08		
		3	44176.23	-1157.66						7°	3	45549.51			
		2	43892.09	284.14						4d <sup>7</sup> (a <sup>2</sup> F)6p:	w <sup>2</sup> F°	4	45755.55	-1191.03	1.022
		1	44343.91	-451.82								3	46946.58	-301.40	0.702
4d <sup>7</sup> (a <sup>2</sup> D)5p	w <sup>2</sup> D°	3	41482.66		1.286			2	47247.98						
		2	42533.81	-1051.15	1.025			t <sup>2</sup> D°	3	45923.36		1.089			
		1	42894.42	-360.61	0.810				2	46499.70	-576.34	0.72			
4d <sup>7</sup> (a <sup>2</sup> H)5p	z <sup>2</sup> I°	7	42290.53		1.146			1	45978.14		0.44				
		6	41577.75	682.78	1.013			8°	1	46056.23		1.115			
		5	42978.28	-1400.53	0.861				9°	1	46056.23				
4d <sup>7</sup> (a <sup>2</sup> H)5p	z <sup>2</sup> G°	5	41739.30		1.197	10°	4	46067.24							
		4	42939.12	-1199.82	1.197		11°	0	46102.93		0/0				
		3	43975.79	-1036.67	0.934	12°		4	46273.20						
4d <sup>7</sup> (a <sup>2</sup> D)5p	z <sup>2</sup> D°	2	41756.15		1.182		13°	4	46400.58						
		4d <sup>7</sup> (a <sup>2</sup> F)6s	e <sup>2</sup> F	4	41825.23				4d <sup>7</sup> 5s(b <sup>2</sup> H)5p:	z <sup>2</sup> H°	5	46495.05		1.030	
				3	43115.47	-1290.24		4d <sup>7</sup> (a <sup>2</sup> D)5p:			y <sup>2</sup> P°	1	46528.26		1.05
2	44970.04			-1854.57		14°	4					46695.02			
4d <sup>7</sup> (a <sup>2</sup> H)5p	z <sup>2</sup> I°	6	42404.147				4d <sup>7</sup> 5s(b <sup>2</sup> P)5p	z <sup>2</sup> P°	3	46746.35	-57.25				
		4d <sup>7</sup> (a <sup>2</sup> P)5p	z <sup>2</sup> P°	1	42415.81				0.965	2	46803.60	-696.94			
				4d <sup>7</sup> (a <sup>2</sup> P)5p	z <sup>2</sup> S°	0			42620.80		0/0	1	47500.54		
A	4					42895.39			15°	1	46789.23				
4d <sup>7</sup> (a <sup>2</sup> D)5p	y <sup>2</sup> F°	3	42998.31		0.995										
		4d <sup>7</sup> (a <sup>2</sup> P)5p	y <sup>2</sup> S°	1	43107.52		1.533								
				2	43509.17										
1	43841.53			-332.36	1.158										
	u <sup>2</sup> D°	3			0.800										

Ru I—Continued

Ru I—Continued

Config.	Design.	J	Level	Interval	Obs. g	Config.	Design.	J	Level	Interval	Obs. g			
4d <sup>2</sup> (a <sup>2</sup> F)5d:	f <sup>2</sup> F	5	46906. 23				38°	4	48597. 45		1. 206			
		4	48542. 88	-1637. 56				39°	1	48604. 34		2. 06		
		3	48809. 28	-265. 40				40°	3	48765. 88				
		2						41°	2	48779. 15				
		1						42°	4	48853. 69		1. 23		
4d <sup>2</sup> (a <sup>2</sup> F)5d:	e <sup>2</sup> D	4	46972. 42		1. 36		43°	3	48933. 93					
		3	47188. 22	-215. 90				44°	2	49037. 35				
		2	48720. 87	-1532. 55				45°	1	49047. 61				
		1	49513. 33?	-792. 46?				46°	3	49141. 42				
		0						47°	4	49165. 05				
4d <sup>2</sup> (a <sup>2</sup> F)5d:	e <sup>2</sup> G	6	46991. 15			4d <sup>2</sup> (a <sup>2</sup> P)5d	e <sup>2</sup> P	3	49291. 06					
		5	48531. 77	-1530. 62				2	50172. 84	-881. 78				
		4						1						
		3	49552. 73					48°	3	49303. 88?				
		2						49°	1	49408. 97				
4d <sup>2</sup> (a <sup>2</sup> F)5d:	e <sup>2</sup> G	16°	47046. 54		1. 058		50°	2	49417. 50					
		5	47064. 80		1. 19			51°	4?	49447. 53				
		4	48727. 68	-1642. 88				4d <sup>2</sup> 5s(a <sup>2</sup> D)5d	e <sup>2</sup> G	2	49439. 56			
		3	49675. 97	-945. 29						C (F)	5	49592. 90		
		17°	47167. 88		1. 24					4d <sup>2</sup> 5s(a <sup>2</sup> D)5d:	f <sup>2</sup> D	5	49624. 26	
18°(D°)	0	47176. 90			4	51058. 64								
19°	4	47261. 52		1. 12	3	52729. 80	-1671. 16							
4d <sup>2</sup> (a <sup>2</sup> F)6p:	e <sup>2</sup> D°	3	47339. 32		1. 377		52°	4	49722. 18					
		2	47547. 33	-208. 01	1. 057			53°	1			49761. 61		
		1						54°	3?	49918. 28?				
		20°(D°)	2	47345. 10				1. 55	e <sup>2</sup> G°	3				
		21°	5	47485. 17						4	49949. 25			
B (F)	4	47486. 96			5	50122. 41	173. 16	1. 16						
22°	3	47526. 09		1. 134	55°	2	49970. 65							
23°	3	47635. 33		1. 06		56°	2	50027. 96						
24°	5	47642. 87					57°	1	50122. 07					
25°	3	47788. 72						58°	1	50338. 99				
26°	1	47809. 11							4d <sup>2</sup> 5s(a <sup>2</sup> D)6e	f <sup>2</sup> D	4	50350. 52		
27°	4	47817. 84			3						51364. 82	-1014. 30		
28°	3	47868. 35		1. 309	2	52001. 35					-636. 53			
29°	3	48003. 00			1									
30°	4	48109. 32			0									
4d <sup>2</sup> 5s(a <sup>2</sup> D)6e	e <sup>2</sup> D	5	48226. 23		1. 602		59°	3	50351. 93					
		4	49235. 15	-848. 82	1. 649			60°	1	50772. 05				
		3	50016. 70	-781. 55					61°	4	51360. 21		1. 094	
		2	50539. 81	-522. 11						D	3	52455. 23		
		1	50618. 88	-79. 09						62°	4	53717. 63		
4d <sup>2</sup> (a <sup>2</sup> F)5d:	e <sup>2</sup> D	34°	48405. 09		1. 14		E			4	54043. 38			
		3	48489. 66		1. 28			Ru II(F <sub>6d</sub> )		Limit	---	59410		
		2	49757. 88	-1268. 22										
		1												
		35°	3	48493. 01										
36°	5	48503. 30												
37°	3?	48570. 85												

Ru: Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>1</sup> +	Observed Terms
4d <sup>1</sup>	{ c <sup>1</sup> P b <sup>1</sup> D b <sup>1</sup> F b <sup>1</sup> G
4d <sup>2</sup> 5s <sup>1</sup>	{ d <sup>1</sup> P f <sup>1</sup> D c <sup>1</sup> F b <sup>1</sup> G b <sup>1</sup> H
	ns (n ≥ 5)
4d <sup>1</sup> (e <sup>1</sup> F)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> (e <sup>1</sup> F)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> 5s(e <sup>1</sup> D)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> (e <sup>1</sup> G)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> (e <sup>1</sup> F)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> (e <sup>1</sup> D)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> (e <sup>1</sup> H)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> 5s(e <sup>1</sup> D)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> (e <sup>1</sup> F)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> 5s(b <sup>1</sup> P)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>
4d <sup>1</sup> 5s(b <sup>1</sup> H)ns	{ s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> s <sup>1</sup> H <sup>o</sup>

\*For predicted terms in the spectra of the Ru I isoelectronic sequence, see Vol. III, Introduction.

## Ru II

(To I sequence; 43 electrons)

Z=44

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2 4p^6 4d^7 {}^4F_{3/2}$  $\epsilon {}^4F_{3/2}$  135300 K

I. P. 16.76 volts

The analysis is mostly by Shenstone, who placed Ru II on his program especially in order to provide the data for inclusion here, since the first regularities reported by Meggers and himself in 1930 had not been extended. A complete description of the spectrum, together with measurements for Zeeman patterns from the Massachusetts Institute of Technology spectrograms, was furnished him by Meggers. The Bureau observations extend from 2005.69 Å to 6662.68 Å. These were supplemented by Princeton observations in the short-wave region extending to 1055 Å.

There are about 1400 classified lines, including nearly all of the stronger lines. Observed intersystem combinations connect the terms of different multiplicity. It is noteworthy that all terms from the  $4d^7$  configuration have been found.

The limit is from Catalán and Rico who have derived it by a comparison of second spectra from Sr to Cd.

## REFERENCES

- K. G. Kessler and W. F. Meggers, *J. Research Nat. Bur. Std.*, **55**, 97, RP 2609 (1955).  
 W. F. Meggers, unpublished material (May 1955). (Z E).  
 M. A. Catalán y F. R. Rico, letter (December 1956). (I P)  
 A. G. Shenstone, unpublished material (October 1957). (I P) (T) (C L) (Z E).

## Ru II

## Ru II

Config.	Desig.	J	Level	Interval	Obs. $\nu$	Config.	Desig.	J	Level	Interval	Obs. $\nu$
$4d^7$	$\epsilon {}^4F$	4 $\frac{1}{2}$	0.0			$4d^7(\epsilon F)5s$	$b {}^4P$	3 $\frac{1}{2}$	26911.4	-2179.6	1.572
		3 $\frac{3}{2}$	1523.4	-1523.4	1 $\frac{1}{2}$			29001.0	-1398.3	1.636	
		2 $\frac{1}{2}$	2494.2	-970.8	0 $\frac{1}{2}$			30489.3		2.541	
		1 $\frac{1}{2}$	3104.7	-610.5							
$4d^7$	$\epsilon {}^4P$	3 $\frac{1}{2}$	8286.7	-230.7	1.57	$4d^7(\epsilon F)5s$	$b {}^4F$	4 $\frac{1}{2}$	27544.6	-404.2	1.200
		1 $\frac{1}{2}$	8477.4	-308.0	1.66			3 $\frac{1}{2}$	27948.8	-190.0	1.092
		0 $\frac{1}{2}$	9373.4		2.61			2 $\frac{1}{2}$	28138.8		0.940
								1 $\frac{1}{2}$	28495.3		0.415
$4d^7(\epsilon D)5s$	$\epsilon {}^4D$	4 $\frac{1}{2}$	9151.5	-990.1	1.54	$4d^7(\epsilon G)5s$	$\epsilon {}^4G$	5 $\frac{1}{2}$	29018.6	-1081.1	1.237
		3 $\frac{1}{2}$	10150.6	-701.2	1.576			4 $\frac{1}{2}$	30099.7	-339.9	1.167
		2 $\frac{1}{2}$	10851.8	-451.8	1.637			3 $\frac{1}{2}$	30439.6	146.5	1.021
		1 $\frac{1}{2}$	11303.6	-300.5	1.843			2 $\frac{1}{2}$	30293.1		0.649
		0 $\frac{1}{2}$	11604.1		2.271						
$4d^7$	$\epsilon {}^4G$	4 $\frac{1}{2}$	10860.9	-1432.5		$4d^7(\epsilon H)5s$	$b {}^4H$	5 $\frac{1}{2}$	32623.0	-63.7	1.072
		3 $\frac{1}{2}$	12293.4		4 $\frac{1}{2}$			32686.7		0.958	
$4d^7$	$\epsilon {}^4P$	1 $\frac{1}{2}$	12956.6	-1842.9		$4d^7(\epsilon D)5s$	$b {}^4D$	0 $\frac{1}{2}$	32960.9	-72.5	0.135
		0 $\frac{1}{2}$	14793.5		1 $\frac{1}{2}$			32888.4	130.6	1.200	
					2 $\frac{1}{2}$			33019.0	313.5	1.344	
$4d^7$	$\epsilon {}^4D$	2 $\frac{1}{2}$	14681.2	-2436.4	1.27	$4d^7(\epsilon F)5s$	$b {}^4P$	1 $\frac{1}{2}$	33734.9	-2360.0	1.188
		1 $\frac{1}{2}$	17017.6		0 $\frac{1}{2}$			36094.9		0.870	
$4d^7$	$\epsilon {}^4H$	5 $\frac{1}{2}$	14663.4	-1461.6		$4d^7(\epsilon F)5s$	$b {}^4F$	3 $\frac{1}{2}$	34038.4	-1260.1	1.063
		4 $\frac{1}{2}$	16125.0		2 $\frac{1}{2}$			35298.5		0.945	
$4d^7(\epsilon D)5s$	$\epsilon {}^4D$	3 $\frac{1}{2}$	19378.7	-1136.2	1.403	$4d^7$	$b {}^4D$	1 $\frac{1}{2}$	34793.8	35.9	0.965
		2 $\frac{1}{2}$	20514.9	-731.3	1.337			2 $\frac{1}{2}$	34829.2		1.102
		1 $\frac{1}{2}$	21246.2	-399.3	1.188						
		0 $\frac{1}{2}$	21645.5		0.0						
$4d^7$	$\epsilon {}^4F$	3 $\frac{1}{2}$	21557.8	731.2	0.864	$4d^7 5s^2$	$\epsilon {}^4S$	2 $\frac{1}{2}$	35857.6		
		3 $\frac{1}{2}$	22289.0		1.141						
$4d^7(\epsilon H)5s$	$\epsilon {}^4H$	6 $\frac{1}{2}$	25962.2	-157.3	1.23	$4d^7(\epsilon I)5s$	$\epsilon {}^4I$	6 $\frac{1}{2}$	35939.7	-290.1	1.06
		5 $\frac{1}{2}$	26109.5	-3.9	1.16			5 $\frac{1}{2}$	36239.8		0.90
		4 $\frac{1}{2}$	26118.4		1.14						
		3 $\frac{1}{2}$	26468.1	-349.7	0.753			$4d^7(\epsilon G)5s$	$b {}^4G$	4 $\frac{1}{2}$	36016.0
			3 $\frac{1}{2}$	36515.8		0.883					

Ru II—Continued

Ru II—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
4d <sup>9</sup> (G)5s	c <sup>3</sup> G	4 1/2	37433.1	-548.5	1.078
		3 1/2	37981.6		0.942
4d <sup>9</sup> (D)5s	c <sup>3</sup> D	2 1/2	39061.9	-780.1	1.2
		1 1/2	39712.0		0.789
4d <sup>9</sup> (S)5s	e <sup>3</sup> S	0 1/2	40226.2		2.0?
4d <sup>9</sup> (D)5s	d <sup>3</sup> D	1 1/2	42750.2	859.7	0.80
		2 1/2	43609.9		1.20
4d <sup>9</sup> (D)5p	s <sup>3</sup> D <sup>o</sup>	4 1/2	48470.9	-240.5	1.530
		3 1/2	48711.4		1.558
		2 1/2	47984.9		-572.5
		1 1/2	47708.4		-422.5
		0 1/2	47983.7		-275.8
4d <sup>9</sup> (D)5p	s <sup>3</sup> F <sup>o</sup>	5 1/2	50758.8	-87.1	1.45
		4 1/2	50845.3		1.43
		3 1/2	50882.5		-17.2
		2 1/2	51179.4		-316.9
		1 1/2	51316.6		-137.2
		0 1/2	51379.8		-63.2
4d <sup>9</sup> (D)5p	s <sup>3</sup> P <sup>o</sup>	3 1/2	51848.8	-1271.5	1.580
		2 1/2	52890.3		1.835
		1 1/2	53685.1		2.365
4d <sup>9</sup> (D)5p	s <sup>3</sup> F <sup>o</sup>	4 1/2	52964.3	-1261.1	1.246
		3 1/2	54285.4		1.262
		2 1/2	54794.8		-568.8
		1 1/2	55223.9		-429.7
4d <sup>9</sup> (D)5p	s <sup>3</sup> D <sup>o</sup>	3 1/2	53316.9	-748.2	1.40
		2 1/2	54065.1		1.347
		1 1/2	54663.8		-596.1
		0 1/2	54981.8		-318.0
4d <sup>9</sup> (D)5p	s <sup>3</sup> P <sup>o</sup>	2 1/2	55694.9	-969.7	1.571
		1 1/2	56664.6		1.712
		0 1/2	57263.3		2.648
4d <sup>9</sup> (F)5p	s <sup>3</sup> S <sup>o</sup>	1 1/2	62446.3		1.789
4d <sup>9</sup> (H)5p	s <sup>3</sup> H <sup>o</sup>	3 1/2	62626.6	380.7	0.960
		4 1/2	62907.3		1.023
		5 1/2	63307.8		400.5
		6 1/2	63588.8		280.5
4d <sup>9</sup> (H)5p	s <sup>3</sup> G <sup>o</sup>	5 1/2	63518.5	-332.8	1.24
		4 1/2	63851.3		-89.8
		3 1/2	63941.1		-84.2
		2 1/2	64025.3		0.646
4d <sup>9</sup> (P)5p	y <sup>3</sup> P <sup>o</sup>	2 1/2	63797.3	-1710.8	1.295
		1 1/2	65508.1		1.857
		0 1/2			
4d <sup>9</sup> (H)5p	s <sup>3</sup> I <sup>o</sup>	4 1/2	64385.3	1084.9	0.960
		5 1/2	65472.8		1.052
		6 1/2	66192.8		726.6
		7 1/2	66359.8?		-939.0
4d <sup>9</sup> (F)5p	y <sup>3</sup> F <sup>o</sup>	1 1/2	64921.7	-457.9	0.428
		2 1/2	64463.8		1.022
		3 1/2	64743.8		285.0
		4 1/2	65328.6		577.8
4d <sup>9</sup> (H)5p	s <sup>3</sup> G <sup>o</sup>	3 1/2	65212.0	563.3	1.032
		4 1/2	65776.3		0.994
4d <sup>9</sup> (P)5p	s <sup>3</sup> D <sup>o</sup>	2 1/2	65844.8		1.325
4d <sup>9</sup> (H)5p	s <sup>3</sup> I <sup>o</sup>	6 1/2	65546.6	-702.8	1.12
		5 1/2	66249.4		1.05
4d <sup>9</sup> (F)5p	s <sup>3</sup> F <sup>o</sup>	3 1/2	66012.3	-1086.9	1.199
		2 1/2	67099.8		0.850

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	
4d <sup>9</sup> (F)5p	y <sup>3</sup> D <sup>o</sup>	3 1/2	66332.0	-2426.0	1.243	
		2 1/2	65774.0		1.306	
		1 1/2	67900.0		874.0	
		0 1/2	68633.3		1264.7	
4d <sup>9</sup> (F)5p	z <sup>3</sup> D <sup>o</sup>	0 1/2	69097.0	-993.8	0.285	
		1 1/2	67108.2		1.187	
		2 1/2	66650.7		-452.5	
		3 1/2	68229.8		1572.5	
4d <sup>9</sup> (F)5p	y <sup>3</sup> G <sup>o</sup>	5 1/2	67601.7	426.3	1.164	
		4 1/2	67165.4		-480.7	
		3 1/2	67648.1		-300.6	
		2 1/2	67848.7			
4d <sup>9</sup> (H)5p	z <sup>3</sup> H <sup>o</sup>	5 1/2	67501.4	-619.0	1.13	
		4 1/2	68120.4		1.018	
4d <sup>9</sup> (G)5p	z <sup>3</sup> F <sup>o</sup>	4 1/2	67901.3	-1502.7	1.19	
		3 1/2	69404.0		-242.6	
		2 1/2	69648.6		26.5	
		1 1/2	69920.1			
4d <sup>9</sup> (F)5p	y <sup>3</sup> G <sup>o</sup>	3 1/2	68265.6	915.3	1.028	
		4 1/2	69180.9		1.055	
4d <sup>9</sup> (G)5p	z <sup>3</sup> G <sup>o</sup>	5 1/2	68328.3	-578.2	1.152	
		4 1/2	68906.5		-644.1	
		3 1/2	69658.6		498.0	
		2 1/2	69654.6			
4d <sup>9</sup> (P)5p	s <sup>3</sup> P <sup>o</sup>	1 1/2	68445.9	-993.6	1.16	
		0 1/2	69439.5		0.829	
4d <sup>9</sup> (G)5p	y <sup>3</sup> H <sup>o</sup>	6 1/2	69311.8	-325.6	1.196	
		5 1/2	69646.8		34.0	
		4 1/2	69612.8		-42.6	
		3 1/2	69655.4			
		1 <sup>o</sup>	1 1/2	70150.7?		1.209
4d <sup>9</sup> (F)5p	y <sup>3</sup> D <sup>o</sup>	2 1/2	70571.1	-206.4	1.157	
		1 1/2	70779.5		0.745	
4d <sup>9</sup> (P)5p	s <sup>3</sup> S <sup>o</sup>	0 1/2	71036.8		2.047	
4d <sup>9</sup> (D)5p	z <sup>3</sup> P <sup>o</sup>	2 1/2	71115.0	-751.5	1.246	
		1 1/2	71866.5		1.567	
		0 1/2	73195.5		-1329.0	
4d <sup>9</sup> (G)5p	y <sup>3</sup> H <sup>o</sup>	5 1/2	71808.9	-723.6	0.737	
		4 1/2	71926.9		0.98	
4d <sup>9</sup> (G)5p	y <sup>3</sup> F <sup>o</sup>	2 1/2	71598.6	411.9	1.177	
		3 1/2	72010.5		1.116	
4d <sup>9</sup> (D)5p	z <sup>3</sup> F <sup>o</sup>	2 1/2	72248.5		1.137	
		3 1/2				
4d <sup>9</sup> (D)5p	w <sup>3</sup> F <sup>o</sup>	1 1/2	72523.5	446.7	0.490	
		2 1/2	72970.8		0.9	
		3 1/2	73142.1		171.9	
		4 1/2	73218.8		76.2	
4d <sup>9</sup> 5s <sup>1</sup> (S)5p	y <sup>3</sup> P <sup>o</sup>	1 1/2	72760.7	148.1	1.30	
		2 1/2	72908.8		304.4	
		3 1/2	73212.2			
		2 <sup>o</sup>	1 1/2	72736.0		1.712
4d <sup>9</sup> (I)5p	s <sup>3</sup> K <sup>o</sup>	7 1/2	72903.0?	-339.0		
		6 1/2	73242.0?			
4d <sup>9</sup> (D)5p	w <sup>3</sup> D <sup>o</sup>	0 1/2	73749.5	-433.7	0.552	
		1 1/2	73315.8		1.20	
		2 1/2	73284.0		-31.8	
		3 1/2	74594.0		1310.0	

## Ru II—Continued

## Ru II—Continued

Config.	Desig.	J	Level	Interval	Obs. $\nu$	Config.	Desig.	J	Level	Interval	Obs. $\nu$
4d <sup>8</sup> ( <sup>1</sup> I)5p	z <sup>1</sup> H°	4½	73993.4	587.2	0.96	4d <sup>8</sup> 5s( <sup>4</sup> S)5p	w <sup>1</sup> P°	3½	79503.8		1.083
		5½	73960.8		1.09			2½	81781.7		
4d <sup>8</sup> ( <sup>3</sup> G)5p	z <sup>1</sup> G°	3½	73810.9	235.6	1.114			1½	82839.5	-517.8	
		4½	74045.8		0.713			0½	82650.8		
4d <sup>8</sup> ( <sup>3</sup> D)5p	z <sup>1</sup> D°	1½	74598.3	525.1	0.8	4d <sup>8</sup> ( <sup>3</sup> D)6s	e <sup>1</sup> D	4½	84510.9		
		2½	75117.4		1.2			3½			
4d <sup>8</sup> ( <sup>1</sup> G)5p	w <sup>1</sup> G°	4½	75346.3	-85.4	1.057	4d <sup>8</sup> ( <sup>3</sup> D)6s	e <sup>1</sup> D	3½	86440.4	-1083.0	
		3½	75431.7		0.911			2½	87523.4		
4d <sup>8</sup> ( <sup>3</sup> G)5p	w <sup>1</sup> H°	5½	75693.1	-726.9	1.08			1½			
		4½	76339.0		0.94			0½			
4d <sup>8</sup> ( <sup>1</sup> I)5p	y <sup>1</sup> I°	6½	75789.5	-135.9	1.08	4d <sup>8</sup> 5s( <sup>4</sup> S)6p	z <sup>1</sup> P°	3½	90165.8	-664.7	
		5½	75925.4		0.92			2½	90829.9		
4d <sup>8</sup> ( <sup>1</sup> G)5p	w <sup>1</sup> F°	3½	76977.0		1.14			1½	91503.7		
		2½									
4d <sup>8</sup> ( <sup>1</sup> D)5p	y <sup>1</sup> P°	1½	77898.7		1.33	-----					
4d <sup>8</sup> ( <sup>1</sup> D)5p	w <sup>1</sup> D°	2½	77819.0	-798.1	1.07	Ru III( <sup>4</sup> D <sub>1</sub> )	Limit	-----		135200	
		1½	78917.1		0.86						
4d <sup>8</sup> ( <sup>1</sup> D)5p	e <sup>1</sup> F°	2½	79114.9	689.1	1.0						
		3½	79804.0		1.046						

October 1957.

## Ru II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>5</sup> +	Observed Terms	
4d <sup>7</sup>	a <sup>4</sup> P a <sup>2</sup> P	a <sup>4</sup> F a <sup>2</sup> F
4d <sup>8</sup> 5s <sup>1</sup>	a <sup>4</sup> S b <sup>2</sup> D	a <sup>3</sup> G a <sup>1</sup> H
	ns (n ≥ 5)	
4d <sup>8</sup> ( <sup>3</sup> D)ns	a, e <sup>3</sup> D a, e <sup>1</sup> D	z <sup>4</sup> P° z <sup>2</sup> P°
4d <sup>8</sup> 5s( <sup>4</sup> S)ns		z <sup>1</sup> D° z <sup>3</sup> D°
4d <sup>8</sup> 5s( <sup>4</sup> S)ns		y <sup>4</sup> P° z <sup>4</sup> P° w <sup>4</sup> P°
4d <sup>8</sup> ( <sup>3</sup> H)ns		z <sup>4</sup> G° z <sup>2</sup> G°
4d <sup>8</sup> ( <sup>3</sup> H)ns		z <sup>4</sup> H° z <sup>2</sup> H°
4d <sup>8</sup> ( <sup>3</sup> H)ns		z <sup>4</sup> I° z <sup>2</sup> I°
4d <sup>8</sup> ( <sup>3</sup> P)ns	b <sup>4</sup> P b <sup>2</sup> P	z <sup>4</sup> S° z <sup>2</sup> S°
4d <sup>8</sup> ( <sup>3</sup> P)ns		y <sup>4</sup> P° z <sup>4</sup> P° z <sup>2</sup> P°
4d <sup>8</sup> ( <sup>3</sup> F)ns		y <sup>4</sup> D° z <sup>4</sup> D° z <sup>2</sup> D°
4d <sup>8</sup> ( <sup>3</sup> F)ns	b <sup>4</sup> F b <sup>2</sup> F	z <sup>4</sup> F° y <sup>4</sup> F° z <sup>4</sup> G° y <sup>4</sup> G°
4d <sup>8</sup> ( <sup>3</sup> G)ns		z <sup>4</sup> F° y <sup>4</sup> F° z <sup>4</sup> G° z <sup>2</sup> G°
4d <sup>8</sup> ( <sup>3</sup> G)ns		z <sup>4</sup> H° y <sup>4</sup> H°
4d <sup>8</sup> ( <sup>3</sup> D)ns	b <sup>4</sup> D c <sup>2</sup> D	z <sup>4</sup> P° w <sup>4</sup> D° z <sup>4</sup> D° z <sup>2</sup> D°
4d <sup>8</sup> ( <sup>1</sup> I)ns		
4d <sup>8</sup> ( <sup>1</sup> I)ns	a <sup>4</sup> I	z <sup>4</sup> H° y <sup>4</sup> I° z <sup>4</sup> K°
4d <sup>8</sup> ( <sup>1</sup> G)ns		w <sup>4</sup> F° w <sup>4</sup> G° w <sup>4</sup> H°
4d <sup>8</sup> ( <sup>1</sup> S)ns	a <sup>4</sup> S	
4d <sup>8</sup> ( <sup>1</sup> D)ns	d <sup>4</sup> D	y <sup>4</sup> P° w <sup>4</sup> D° v <sup>4</sup> F°

\*For predicted terms in the spectra of the Tc I isoelectronic sequence, see Volume III, Introduction.

## Ru III

(Mo I sequence; 42 electrons)

Z=44

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2 4p^6 4d^8 {}^1D_2$  $4d^8 {}^1D_2$ , 229600 K

I. P. 28.46 volts

The analysis is by Shenstone, who has investigated Ru III especially for inclusion here. He has observed the spectrum from 600 Å to 4004 Å and classified 40 lines. The quintet and septet systems of terms are connected by observed intersystem combinations.

He derives the limit 225900 from the  $ns$  series by means of a Ritz formula, with  $\alpha = 1.033 \times 10^{-6}$ . This value of  $\alpha$  represents  $\alpha(\text{Fe III}) \times \{\alpha(\text{Ag II})/\alpha(\text{Cu II})\}$ . Catalán and Rico have interpolated the quoted limit by comparison of the third spectra from Y to In.

## REFERENCES

A. G. Shenstone, unpublished material (June 1955). (I P) (T) (C L)

M. A. Catalán y F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] 53, 85 (1957). (I P)

## Ru III

## Ru III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^8$	$4d^8 {}^1D$	4	0.0		$4d^8({}^6S)5d$	$5d {}^1D$	1	137218.67	
		3	1158.8	-1158.8			2	137251.0	32.4
		2	1826.3	-667.5			3	137303.6	52.6
		1	2266.3	-440.0			4	137383.8	80.2
		0	2476.0	-209.7			5	137502.6	118.8
$4d^8({}^6S)5s$	$5s {}^7S$	3	27162.8		$4d^8({}^6S)6s$	$6s {}^7S$	3	137482.3	
$4d^8({}^6S)5s$	$5s {}^9S$	2	41111.7		$4d^8({}^6S)6s$	$6s {}^9S$	2	139692.3	
$4d^8({}^6S)5p$	$5p {}^1P^o$	2	76056.1		$4d^8({}^6S)5d$	$5d {}^3D$	0		
		3	76915.6	859.5			1	140345.87	141.8
		4	78673.5	1757.9			2	140487.6	-40.4
$4d^8({}^6S)5p$	$5p {}^3P^o$	3	83897.9				3	140447.2	-27.7
		2	84487.1	-489.2	4	140419.5			
		1	84882.9	-335.8					
					-----	-----			
					Ru IV ( ${}^6S_{3/2}$ )	Limit	-----	229600	

February 1957.



## RHODIUM

## Rh I

45 electrons

Z=45

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^8 5s^1 \text{ } ^4F_{3/2}$  $\alpha$   $^4F_{3/2}$  60197 K

I. P. 7.46 volts

The first spectrum of rhodium has a unique place in the history of atomic spectra. In 1901 C. P. Snyder published an array of Rh I wave numbers consisting of 19 columns and 54 rows, accounting for 476 Rh I lines. The significance of this work was not appreciated until years later when regularities were found in many spectra. It may be said fairly that this work anticipated by years the long series of investigations culminating in the Hund theory of atomic spectra. All efforts to contact this author have failed—and yet spectroscopists are deeply indebted to him for his remarkable pioneer work illustrating in detail the correct approach to the study of complex spectra.

In 1927 Sommer extended the early work on this spectrum by Meggers and Laporte. He published an extensive analysis including about 1000 classified lines between 2288.53 Å and 8425.51 Å, 136 energy levels, and numerous Zeeman observations. He derived the limit 62000 K from the  $^4F$ - and  $^2F$ -series of two members each, by using a Rydberg formula.

Subsequently Molnar and Hitchcock observed the Rh I spectrum at the Massachusetts Institute of Technology, and revised the analysis with the aid of extensive Zeeman observations made with the Bitter magnet. The data in the table are from their paper. They derived improved values for the energy levels throughout the spectrum, added 14 energy levels, and rejected 19 of Sommer's miscellaneous levels. Of these, one at 532      may be a misprint for a possible level at 53223.85 with  $J=1\frac{1}{2}$  or  $2\frac{1}{2}$ . Two others at 56898.60 ( $J=2\frac{1}{2}$ ) and 63891.07 ( $J=3\frac{1}{2}$ ), respectively, may be real, although the combinations are limited to faint lines. There are about 950 classified lines between 1987.84 Å and 8615.23 Å. The doublet and quartet systems of terms are connected by observed intersystem combinations. The following changes in notation have been introduced into the table:

Designation

Molnar and Hitchcock	Adopted AEL
$a^3D$	$a^3D$
$a^1D$	$b^1D$
$c^3F$	$c^3F$
$c^1F$	$c^1F$
$x^3P^o$	$y^3P^o$
$y^3P^o$	$z^3P^o$
$x^1F^o$	$y^1F^o$
$y^1F^o$	$z^1F^o$

## Rh I—Continued

The writer has also interchanged the configuration assignments of the two  $^2P^{\circ}$  terms having singlet limits so that the lower one, here called  $\gamma^2P^{\circ}$ , has the limit  $^1D$ , and the next higher one,  $z^2P^{\circ}$ , has the limit  $^1S$ .

Murphy has recently extended the observations between 6332.96 Å and 11021.74 Å. He has added 53 classified lines and the two levels  $b^4P_{1/2}$  and  $b^4G_{3/2}$ . The remaining levels of these terms have been furnished by the writer. A homogeneous line list and further analysis are needed. No sextet terms have been found, no terms involving  $d$ -electrons are known, and many miscellaneous levels should be assigned their proper term designations and configurations.

The limit quoted here is from Catalán and Rico. They have derived it from a study of the series in the spectra Rb I to Ag I.

## REFERENCES

- C. P. Snyder, *Astroph. J.* **14**, 179 (1901).  
 W. F. Meggers and O. Laporte, *Phys. Rev.* **28**, 653 (1926). (T) (C L)  
 L. A. Sommer, *Zeit. Phys.* **45**, 147 (1927). (I P) (T) (C L) (Z E) (E D) (G D)  
 J. P. Molnar and W. J. Hitchcock, *J. Opt. Soc. Am.* **30**, 523 (1940). (T) (C L) (Z E)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 R. J. Murphy, *J. Research Nat. Bur. Std.* **49**, 371, RP 2374 (1952). (T) (C L)  
 M. A. Catalán y F. R. Rico, *An. Real Soc. Esp. Física y Química (Madrid)* [A] **48**, 328 (1952). (I P)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955). (Summary hfs)

## Rh I

## Rh I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
$4d^8(^2F)5s$	$a^4F$	$4\frac{1}{2}$	0.00		1.329	$4d^8(^2F)5p$	$z^4F^{\circ}$	$4\frac{1}{2}$	29430.86		1.261
		$3\frac{1}{2}$	1529.97	-1529.97	1.230			$3\frac{1}{2}$	29886.34	-435.48	1.148
		$2\frac{1}{2}$	2598.03	-1068.06	1.105			$2\frac{1}{2}$	31474.60	-1608.16	0.834
		$1\frac{1}{2}$	3472.68	-874.65	0.473			$1\frac{1}{2}$	32277.43	-802.93	0.556
$4d^8$	$a^2D$	$2\frac{1}{2}$	3309.86		1.116	$4d^8(^2F)5p$	$z^2G^{\circ}$	$4\frac{1}{2}$	31613.78		1.156
		$1\frac{1}{2}$	5657.97	-2348.11	0.744			$3\frac{1}{2}$	33045.91	-1430.13	0.991
$4d^8(^2F)5s$	$a^2F$	$3\frac{1}{2}$	5690.97		1.143	$4d^8(^2F)5p$	$z^2F^{\circ}$	$3\frac{1}{2}$	32004.01		1.120
		$2\frac{1}{2}$	7791.23	-2100.26	0.947			$2\frac{1}{2}$	33946.34	-1942.33	1.011
$4d^8(^2F)5s$	$a^4P$	$2\frac{1}{2}$	9221.22		1.478	$4d^8(^2F)5p$	$z^2D^{\circ}$	$2\frac{1}{2}$	32046.32		0.999
		$1\frac{1}{2}$	10313.41	-1092.19	1.576			$1\frac{1}{2}$	33867.04	-1820.72	0.913
		$0\frac{1}{2}$	11006.05	-692.64	2.661						
$4d^8(^2F)5s$	$a^2P$	$1\frac{1}{2}$	11968.23		1.230	$4d^8(^2P)5p$	$z^4P^{\circ}$	$2\frac{1}{2}$	35333.86		1.470
		$1\frac{1}{2}$	13974.73	-2006.50	0.681			$1\frac{1}{2}$	35404.24	-70.39	1.451
		$0\frac{1}{2}$						$0\frac{1}{2}$	35669.60	-265.26	2.064
$4d^7 5s^2$	$b^4F$	$4\frac{1}{2}$	12723.07		1.317	$4d^8(^2P)5p$	$y^4D^{\circ}$	$3\frac{1}{2}$	36132.72		1.473
		$3\frac{1}{2}$	14787.87	-2064.80	1.230			$3\frac{1}{2}$	36787.48		1.346
		$2\frac{1}{2}$	16118.69	-1330.82	1.018			$2\frac{1}{2}$	36985.25	-197.77	1.150
		$1\frac{1}{2}$	16943.50	-824.81	0.416			$1\frac{1}{2}$	38038.12	-1052.87	1.134
$4d^8(^1D)5s$	$b^2D$	$2\frac{1}{2}$	13520.69		1.233	$4d^8(^2P)5p$	$y^2D^{\circ}$	$2\frac{1}{2}$	37300.60		1.400
		$1\frac{1}{2}$	14382.19	-861.50	1.034			$1\frac{1}{2}$	38474.43	-436.31	-0.160
$4d^8(^1G)5s$	$a^2G$	$4\frac{1}{2}$	16017.94		1.114	$4d^8(^2P)5p$	$z^2P^{\circ}$	$0\frac{1}{2}$	37368.62		1.342
		$3\frac{1}{2}$	16120.72	-102.78	0.900			$1\frac{1}{2}$	38210.96	842.34	1.210
$4d^7 5s^2$	$b^4P$	$2\frac{1}{2}$	23157.57			$4d^8(^2P)5p$	$y^2D^{\circ}$	$3\frac{1}{2}$	38012.75		1.436
		$1\frac{1}{2}$	23655.93	-498.36				$4\frac{1}{2}$	38668.00		
		$0\frac{1}{2}$	24686.00	-1032.07							
$4d^7 5s^2$	$b^2G$	$4\frac{1}{2}$	25820.80			$4d^8(^2P)5p$	$y^2P^{\circ}$	$2\frac{1}{2}$	38718.11		
		$3\frac{1}{2}$	27876.52	-2055.72				$1\frac{1}{2}$	39525.22	-807.17	1.494
								$0\frac{1}{2}$	38806.64		-0.072
								$5^{\circ}$	39186.76		1.323
$4d^8(^2F)5p$	$z^4D^{\circ}$	$3\frac{1}{2}$	27075.26		1.410	$4d^8(^1D)5p$	$y^2P^{\circ}$	$1\frac{1}{2}$	39231.32		1.263
		$2\frac{1}{2}$	28859.64	-1784.38	1.330			$0\frac{1}{2}$	43729.02	-4497.64	0.687
		$1\frac{1}{2}$	30397.27	-1537.63	1.165						
		$0\frac{1}{2}$	31148.68	-749.41	0.040						
$4d^8(^2F)5p$	$z^4G^{\circ}$	$5\frac{1}{2}$	29104.71		1.273	$4d^8(^2P)5p$	$y^2P^{\circ}$	$1\frac{1}{2}$	39494.00		1.180
		$4\frac{1}{2}$	28542.69	562.02	1.181			$3\frac{1}{2}$	39788.09		1.111
		$3\frac{1}{2}$	31101.75	-2559.06	1.023						
		$2\frac{1}{2}$	32243.32	-1141.57	0.937						

Rh I—Continued

Rh I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
	9°	2½	39981.30		1.045	5	4½, 5½		47862.14		
	10°	1½	40134.48		1.230	6	3½		47896.92		
4d <sup>o</sup> (D)5p	y <sup>o</sup> F°	3½ 2½	40884.83 40576.95	-292.12	1.204 1.068	7	4½		47928.07		1.15
4d <sup>o</sup> (S)5p	x <sup>o</sup> P°	1½ 0½	40603.48		1.268	24°	2½		47954.80		-0.083
4d <sup>o</sup> (P)5p	z <sup>o</sup> S°	0½	40604.58		1.870	8	2½		48239.04		1.20
4d <sup>o</sup> (P)5p	z <sup>o</sup> S°	1½	40900.00		1.232	25°	2½		48797.74		1.066
	11°	2½	41445.67		1.983	26°	3½		48811.47		0.928
4d <sup>o</sup> (F)6e	e <sup>o</sup> F	4½ 3½ 2½ 1½	41880.86 44444.52 45471.24 45811.72	-2583.66 -1026.72 -340.48	1.31 1.22	27°	2½		48713.30		
	12°	4½	41953.07		1.127	9	2½		50042.87		
4d <sup>o</sup> (F)6e	e <sup>o</sup> F	3½ 2½	42292.51 44474.68	-2182.17		10	1½, 2½		50111.87		1.36
	13°	4½	42325.46		1.113	28°	2½		50146.60		
4d <sup>o</sup> (D)5p	z <sup>o</sup> D°	1½ 2½	42431.48 43421.40	989.92	1.346 1.008	11	3½, 4½		50190.80		1.00
	14°	3½	42495.48		1.231	12	5½		50199.18		1.12
	15°	4½	43042.31		1.119	13	2½		50207.73		
	16°	3½	43047.54		1.139	14	3½		50233.05		1.12
4d <sup>o</sup> (G)5p	z <sup>o</sup> H°	5½ 4½	43070.04		1.097	15	3½, 4½		50277.62		1.15
4d <sup>o</sup> (G)5p	z <sup>o</sup> F°	2½ 3½	43777.07 43904.73	127.66	0.957 1.069	16	3½		50284.72		0.97
	17°	3½	44588.12		1.260	17	1½		50290.06		
	18°	1½	44620.63		0.520	18	2½		50408.21		1.02
	19°	2½	44786.67		0.654	19	1½		50451.97		
4d <sup>o</sup> (G)5p	y <sup>o</sup> G°	3½ 4½	45177.63 45315.54	137.91	0.948 1.106	20	2½		50721.44		
	20°	2½	45683.41		1.237	21	3½		51324.35		0.74
	21°	1½	46280.09		1.089	22	1½, 2½		51356.22		
	22°	4½	46511.10		1.115	23	2½, 3½		51419.45		
	23°	0½	46762.86		0.024	24	2½, 3½		51477.77		
1	3½		47713.63		1.38	30°	2½, 3½		51608.71		
2	5½		47793.54		1.12	25	1½		51636.54		0.92
3	2½ ?		47832.35		1.33	26	0½, 1½		52005.37		0.960
4	4½		47857.39		1.25	27	1½		52413.28		
						28	1½, 2½		52473.55		
						32°	1½, 2½		53599.28		
						27	1½		56087.60		
						28	1½, 2½		57610.48		
						Rh u(F <sub>4</sub> )	Limit	-----	60197		

February 1955.

## Rh I Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>2</sup> +	Observed Terms		
4d <sup>2</sup>	a <sup>3</sup> D		
4d <sup>2</sup> 5s <sup>2</sup>	b <sup>3</sup> P	b <sup>3</sup> F	b <sup>3</sup> G
	ns (n ≥ 5)		np (n ≥ 5)
4d <sup>2</sup> ( <sup>3</sup> F)ns	a <sup>3</sup> P, a <sup>3</sup> F		s <sup>3</sup> D°, s <sup>3</sup> F°, s <sup>3</sup> G°
4d <sup>2</sup> ( <sup>1</sup> P)ns	a <sup>3</sup> P, a <sup>3</sup> F		s <sup>3</sup> D°, s <sup>3</sup> F°, s <sup>3</sup> G°
4d <sup>2</sup> ( <sup>1</sup> D)ns	b <sup>3</sup> D		y <sup>3</sup> P°, s <sup>3</sup> D°, y <sup>3</sup> F°
4d <sup>2</sup> ( <sup>1</sup> G)ns		a <sup>3</sup> G	s <sup>3</sup> F°, y <sup>3</sup> G°, s <sup>3</sup> H°
4d <sup>2</sup> ( <sup>1</sup> S)ns			s <sup>3</sup> P°

\*For predicted terms of the Rh I isoelectronic sequence, see Vol. III, Introduction.

## Rh II

(Ru I sequence; 44 electrons)

Z=45

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>7</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>8</sup> <sup>3</sup>F<sub>4</sub>

a<sup>3</sup>F<sub>4</sub> 145800 K

I. P. 18.07 volts

The analysis is by Sancho, who has revised and extended the earlier work on Rh II by Livingood and Shenstone, and by Hitchcock. From spectrograms made at Princeton by Catalán, he has prepared a new line list extending from 1100 Å to 4800 Å. There are 494 classified lines. Observed intersystem combinations connect the systems of terms having different multiplicity.

The limit has been interpolated by Catalán and Rico from a study of the ns series of the spectra Sr II to Cd II.

The following observed g-values have been furnished by van Kleef:

Desig.	J	Obs. g	Desig.	J	Obs. g
a <sup>3</sup> F	5	1.37	s <sup>3</sup> F°	5	1.43
	4	1.31		4	1.38
				3	1.32
			s <sup>3</sup> D°	4	1.38

## REFERENCES

- J. J. Livingood and A. G. Shenstone, unpublished material; see R. F. Bacher and S. Goudamit, *Atomic Energy States*, p. 387 (McGraw-Hill Book Co., Inc., New York, N. Y., and London, 1932). (T)  
 W. J. Hitchcock, unpublished material (March 1952). (T) (C L)  
 Th. A. M. van Kleef, unpublished material (September 1956). (Z E)  
 F. J. Sancho, unpublished material (November 1956). (T) (C L)  
 M. A. Catalán y F. R. Rico, letter (December 1956). (I F)

## Rh II

## Rh II

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
4d <sup>2</sup>	a <sup>3</sup> F	4	0.0	-2401.3	4d <sup>2</sup>	a <sup>3</sup> G	4	14855.4	
		3	2401.3	-1179.4					
		2	3580.7						
4d <sup>2</sup>	a <sup>3</sup> D	2	8164.4		4d <sup>2</sup> (a <sup>3</sup> F)5s	a <sup>3</sup> F	5	16884.8	-1655.6
4d <sup>2</sup>	a <sup>3</sup> P	0	10760.8	-245.8			4	18540.4	-1252.0
		1	10515.0	1128.7			3	19792.4	-854.5
		2	11643.7				2	20646.9	-533.1
							1	21180.0	

## Rh II—Continued

## Rh II—Continued

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval								
$4d^7(a^1F)5s$	$b^1F$	4	25376.9	-2062.5	$4d^7(a^1G)5p$	$s^1H^{\circ}$	6	78112.1	1460.1								
		3	27439.4	-1395.2			5	70643.0	-1517.4								
		2	28834.6				4	78180.4									
$4d^7(a^1F)5s$	$a^1P$	3	27801.4	-330.0	$4d^7(a^1P)5p$	$s^1P^{\circ}$	2	78128.5	-1517.8								
		2	28131.4	-941.6			1	78646.3									
		1	29073.0				0										
$4d^7(a^1G)5s$	$a^1G$	5	31730.5	-874.2	$4d^7(a^1P)5p$	$s^1S^{\circ}$	1	78331.5									
		4	32604.7	-1240.7			$y^1D^{\circ}$	1	78780.7	-190.0							
		3	33845.4					2	78570.7	618.0							
0			3	78185.7													
$4d^7(a^1P)5s$	$b^1P$	2	34243.0	-483.1	$4d^7(a^1G)5p$	$s^1G^{\circ}$	4	78870.6									
		1	34726.1	-285.9			$4d^7(a^1H)5p$	$x^1G^{\circ}$	5	78444.3	-2060.2						
		0	35012.0						4	76504.5	-1090.5						
			3	77595.0													
$4d^7(a^1H)5s$	$a^1H$	6	35787.2	-1007.3	$4d^7(a^1P)5p$	$y^1P^{\circ}$	2	78780.1	-817.7								
		5	36794.5	-2148.7			1	74537.8									
		4	38943.2				0										
$4d^7(a^1P)5s$	$c^1P$	2	36366.3	-343.7	$4d^7(a^1G)5p$	$s^1F^{\circ}$	3	74364.4									
		1	36710.0				$4d^7(a^1G)5p$	$y^1G^{\circ}$	5	74509.8	-460.6						
		0							4	74979.4	-51.1						
			3	75030.5													
$4d^7(a^1G)5s$	$b^1G$	4	36486.4		$4d^7(a^1G)5p$	$s^1H^{\circ}$	5	74836.8									
		$4d^7(a^1D)5s$	$a^1D$	3			36938.9	-1683.4	$4d^7(a^1P)5p$	$x^1D^{\circ}$	3	76691.1	-875.5				
				2			38622.3	-65.2			2	76586.6	-1066.4				
1	38687.5				1	77633.0											
$4d^7(a^1H)5s$	$a^1H$	5	41045.3		$4d^7(a^1H)5p$	$s^1I^{\circ}$	7	75905.4									
		$4d^7(a^1D)5s$	$b^1D$	2			42872.3		6	76538.8							
				1					5								
$4d^7(a^1F)5p$	$s^1F^{\circ}$	5	57080.8	473.5	$4d^7(a^1D)5p$	$w^1D^{\circ}$	3	76032.4	25.1								
		4	56547.3	-1811.2			2	76007.3	-134.2								
		3	58358.5	-1340.1			1	76141.5									
		2	59698.6	-874.6			$4d^7(a^1D)5p$	$x^1F^{\circ}$	4	77842.8	-1316.9						
		1	60573.2						3	79159.7	858.8						
0			2	78503.9													
$4d^7(a^1F)5p$	$s^1G^{\circ}$	6	59702.4	-27.0	$4d^7(a^1H)5p$	$s^1I^{\circ}$			6	78297.6							
		5	59789.4	-1443.7					$4d^7(a^1H)5p$	$y^1H^{\circ}$	6	78735.4	-663.9				
		4	61173.1	-766.7			5	79399.3			-854.7						
		3	61939.8	-348.5			4	80254.0									
		2	62288.3				$4d^7(a^1F)5p$	$y^1S^{\circ}$			1	79044.2					
$4d^7(a^1F)5p$	$s^1G^{\circ}$	5	62194.4	-1765.1	$4d^7(a^1P)5p$	$s^1P^{\circ}$					1	79547.6					
		4	63959.5	-1361.7					$4d^7(a^1P)5p$	$s^1D^{\circ}$	2	79717.5					
		3	65321.2								$4d^7(a^1D)5p$	$y^1D^{\circ}$	2	79938.8			
		$4d^7(a^1F)5p$	$s^1F^{\circ}$	4									62326.1	-1128.8	$4d^7(a^1H)5p$	$y^1G^{\circ}$	4
				3			63454.9	-1515.5					$4d^7(a^1D)5p$	$y^1F^{\circ}$			3
2	64970.4				$4d^7(a^1D)5p$	$x^1P^{\circ}$	2	81955.0									-1296.4
$4d^7(a^1F)5p$	$s^1S^{\circ}$			2			63261.4		1	83251.4							
				1					0								
		$4d^7(a^1F)5p$	$s^1D^{\circ}$	3			64819.4	-1259.4	$4d^7(a^1H)5p$	$y^1H^{\circ}$	5	82830.7					
				2			66078.8	-690.9			$4d^7(a^1D)5p$	$y^1P^{\circ}$	1	83176.2			
				1	66769.7		$4d^7(a^1F)5p$	$s^1D^{\circ}$					3	87394.0	-536.1		
$4d^7(a^1P)5p$	$s^1P^{\circ}$			1	69022.9	17.5							2	87930.1	102.6		
				2	69040.4	114.7							1	87827.5			
		3	69155.1														
		$4d^7(a^1G)5p$	$y^1F^{\circ}$	4	70038.3	-2285.4			$Rh III(F_{04})$	Limit	-----	145800					
				3	72303.7	-2848.7											
2	75152.4																
$4d^7(a^1F)5p$	$y^1D^{\circ}$	0	70199.2	343.0													
		1	70542.2	816.4													
		2	71358.6	349.1													
		3	71707.7	-702.0													
		4	71005.7														

Rh II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> +	Observed Terms			
	4d <sup>8</sup>	{ a <sup>3</sup> P a <sup>1</sup> D a <sup>3</sup> F a <sup>1</sup> G		
	nd (n ≥ 5)		np (n ≥ 5)	
4d <sup>7</sup> (a <sup>3</sup> F)ns	{ a <sup>3</sup> F b <sup>3</sup> F		z <sup>3</sup> D° z <sup>1</sup> D°	y <sup>3</sup> F° y <sup>1</sup> F°
4d <sup>7</sup> (a <sup>1</sup> P)ns	{ a <sup>3</sup> P a <sup>1</sup> P		x <sup>3</sup> S° y <sup>3</sup> S°	z <sup>1</sup> P° y <sup>1</sup> P°
4d <sup>7</sup> (a <sup>3</sup> G)ns	{ a <sup>3</sup> G b <sup>3</sup> G			y <sup>3</sup> F° y <sup>1</sup> F°
4d <sup>7</sup> (a <sup>1</sup> P)ns	{ b <sup>3</sup> P		x <sup>3</sup> S° z <sup>1</sup> P°	z <sup>1</sup> D° y <sup>1</sup> D°
4d <sup>7</sup> (a <sup>3</sup> H)ns	{ a <sup>3</sup> H a <sup>1</sup> H			z <sup>3</sup> G° y <sup>3</sup> G°
4d <sup>7</sup> (a <sup>1</sup> D)ns	{ a <sup>3</sup> D b <sup>3</sup> D			y <sup>3</sup> H° y <sup>1</sup> H°
4d <sup>7</sup> (a <sup>3</sup> F)ns				z <sup>3</sup> I° z <sup>1</sup> I°

\*For predicted terms in the spectra of the Ru I isoelectronic sequence, see Volume III, Introduction.

Rh III

(Tc I sequence; 43 electrons)

Z=45

1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>7</sup> <sup>4</sup>F<sub>6,4</sub>

a<sup>4</sup>F<sub>6,4</sub> 250500 K

I. P. 31.05 volts

The analysis has been carried out by Catalán, Sales, and Shenstone especially for inclusion here. They have observed the spectrum from 687A to 4800 A, and classified 854 lines. Observed intersystem combinations connect the systems of terms having different multiplicity.

The limit has been interpolated by comparison of third spectra from Y to In, and furnished by Catalán and Rico in advance of publication.

REFERENCES

- M. A. Catalán, M. Sales and A. G. Shenstone, unpublished material (October 1955). (I P) (T) (C L)  
 M. A. Catalán y F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] 53, 85 (1957). (I P)

Rh III

Rh III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
4d <sup>7</sup>	a <sup>4</sup> F	4½	0.0	-2147.8	4d <sup>7</sup>	a <sup>3</sup> H	5½	19490.2	-2502.1
		3½	2147.8	4½			21992.3		
		2½	3485.7						
		1½	4322.0						
4d <sup>7</sup>	a <sup>4</sup> P	2½	11062.3	65.2	4d <sup>7</sup>	a <sup>3</sup> D	2½	19528.5	-2518.1
		1½	10997.1	-1472.7			1½	22046.6	
		0½	12469.8						
4d <sup>7</sup>	a <sup>3</sup> G	4½	14044.0	-1212.8	4d <sup>8</sup> ( <sup>3</sup> D)5s	a <sup>4</sup> D	4½	43022.0	-1372.4
		3½	15256.8				3½	44394.4	
4d <sup>7</sup>	a <sup>3</sup> P	1½	16870.7	-1433.0			2½	45278.2	-598.4
		0½					1½	45876.6	-350.5
							0½	46227.1	

## Rh III—Continued

## Rh III—Continued

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
4d <sup>7</sup>	b <sup>2</sup> D	1½ 2½	49183. 4 50442. 3	1258. 9	4d <sup>6</sup> ( <sup>2</sup> D)5p	s <sup>2</sup> D <sup>o</sup>	3¼ 2¼ 1¼ 0¼	100845. 9 101857. 9 102855. 1 103827. 3	-1012. 0 -877. 2 -292. 3
4d <sup>6</sup> ( <sup>2</sup> D)5s	a <sup>2</sup> D	3¼ 2¼ 1¼ 0¼	54632. 2 56125. 7 57012. 5 57531. 3	-1493. 5 -885. 8 -518. 8	4d <sup>6</sup> ( <sup>2</sup> P)5s <sup>2</sup>	c <sup>2</sup> P	0¼ 1¼ 2¼	101445. 1 101890. 8 102224. 0	445. 7 333. 2
4d <sup>6</sup> ( <sup>2</sup> P)5s	b <sup>2</sup> P	2¼ 1¼ 0¼	62717. 1 64273. 7 68793. 1	-3556. 6 -2519. 4	4d <sup>6</sup> ( <sup>2</sup> F)5s <sup>2</sup>	c <sup>2</sup> F	1¼ 2¼ 3¼ 4¼	102194. 2 102302. 2 102410. 4 102468. 0	8. 0 208. 2 57. 6
4d <sup>6</sup> ( <sup>2</sup> H)5s	a <sup>2</sup> H	6¼ 5¼ 4¼ 3¼	64250. 0 64882. 8 65217. 9 65483. 6	-632. 8 -335. 1 -265. 7	4d <sup>6</sup> ( <sup>2</sup> D)5p	s <sup>2</sup> P <sup>o</sup>	2¼ 1¼ 0¼	104183. 4 105493. 1 106856. 9	-1309. 7 -742. 8
4d <sup>6</sup> ( <sup>2</sup> F)5s	b <sup>2</sup> F	4¼ 3¼ 2¼ 1¼	66104. 4 66684. 6 67025. 7 67476. 8	-580. 2 -341. 1 -451. 1	4d <sup>6</sup> ( <sup>2</sup> F)5s <sup>2</sup>	d <sup>2</sup> F	2¼ 3¼	106665. 1 107009. 4	344. 3
4d <sup>6</sup> ( <sup>2</sup> G)5s	a <sup>2</sup> G	5¼ 4¼ 3¼ 2¼	70561. 3 72245. 9 73036. 2 73278. 3	-1684. 6 -790. 3 -242. 1	4d <sup>6</sup> ( <sup>2</sup> P)5s <sup>2</sup>	c <sup>2</sup> P	0¼ 1¼	106726. 5 107813. 0	1086. 5
4d <sup>6</sup> ( <sup>2</sup> H)5s	b <sup>2</sup> H	5¼ 4¼	72580. 2 73192. 0	-611. 8	4d <sup>6</sup> ( <sup>2</sup> G)5s <sup>2</sup>	d <sup>2</sup> G	4¼ 3¼	111897. 5 112923. 5	-1026. 0
4d <sup>6</sup> ( <sup>2</sup> P)5s	b <sup>2</sup> P	1¼ 0¼	72837. 7 74909. 6	-2071. 9	4d <sup>6</sup> ( <sup>2</sup> P)5p	s <sup>2</sup> S <sup>o</sup>	1¼	113245. 4	
4d <sup>6</sup> ( <sup>2</sup> F)5s	b <sup>2</sup> F	3¼ 2¼	76752. 3 77708. 6	-956. 3	4d <sup>6</sup> ( <sup>2</sup> P)5p	y <sup>2</sup> P <sup>o</sup>	2¼ 1¼ 0¼	113201. 1 114353. 0	-2671. 9
4d <sup>6</sup> ( <sup>2</sup> G)5s	b <sup>2</sup> G	4¼ 3¼	79304. 7 80600. 4	-1295. 7	4d <sup>6</sup> ( <sup>2</sup> H)5p	s <sup>2</sup> G <sup>o</sup>	5¼ 4¼ 3¼ 2¼	113312. 1 113965. 1 114311. 5 114558. 8	-653. 0 -346. 4 -246. 7
4d <sup>6</sup> ( <sup>2</sup> D)5s	b <sup>2</sup> D	0¼ 1¼ 2¼ 3¼	81589. 5 81713. 6 81893. 5 82156. 5	124. 1 179. 9 263. 0	4d <sup>6</sup> ( <sup>2</sup> H)5p	s <sup>2</sup> H <sup>o</sup>	6¼ 5¼ 4¼ 3¼	113893. 7 114280. 3 114522. 1 115491. 7	-386. 6 -541. 8 -669. 6
4d <sup>6</sup> ( <sup>2</sup> I)5s	a <sup>2</sup> I	6¼ 5¼	82095. 7 82414. 5	-314. 8	4d <sup>6</sup> ( <sup>2</sup> D)5s <sup>2</sup>	e <sup>2</sup> D	2¼ 1¼	115013. 2 116104. 0	-1090. 8
4d <sup>6</sup> ( <sup>2</sup> G)5s	c <sup>2</sup> G	4¼ 3¼	83128. 7 83367. 3	-238. 6	4d <sup>6</sup> ( <sup>2</sup> H)5p	s <sup>2</sup> I <sup>o</sup>	7¼ 6¼ 5¼ 4¼	115590. 0 116209. 0 116344. 0 116064. 8	-619. 0 -135. 0 279. 2
4d <sup>6</sup> ( <sup>2</sup> D)5s	c <sup>2</sup> D	1¼ 2¼	84004. 5 86377. 2	372. 7	4d <sup>6</sup> ( <sup>2</sup> P)5p	y <sup>2</sup> D <sup>o</sup>	3¼ 2¼ 1¼ 0¼	115769. 4 119712. 3 120568. 4 120858. 7	-3942. 9 -856. 1 -290. 3
4d <sup>6</sup> ( <sup>2</sup> S)5s	a <sup>2</sup> S	0¼	88151. 0		4d <sup>6</sup> ( <sup>2</sup> P)5p	z <sup>2</sup> D <sup>o</sup>	2¼ 1¼	115991. 6 118078. 0	-2086. 4
4d <sup>6</sup> ( <sup>2</sup> D)5s	d <sup>2</sup> D	2¼ 1¼	88248. 7 88526. 3	-277. 6	4d <sup>6</sup> ( <sup>2</sup> F)5p	y <sup>2</sup> F <sup>o</sup>	1¼ 2¼ 3¼ 4¼	116011. 1 116322. 8 116407. 1 116364. 4	311. 7 84. 3 457. 1
4d <sup>6</sup> ( <sup>2</sup> D)5p	z <sup>2</sup> D <sup>o</sup>	4¼ 3¼ 2¼ 1¼ 0¼	91818. 7 92031. 6 92285. 5 92412. 8 92809. 3	-202. 9 -804. 3 -586. 9 -396. 5	4d <sup>6</sup> ( <sup>2</sup> H)5p	s <sup>2</sup> G <sup>o</sup>	4¼ 3¼	116295. 7 116965. 0	-669. 3
4d <sup>6</sup> ( <sup>2</sup> D)5p	z <sup>2</sup> L <sup>o</sup>	5¼ 4¼ 3¼ 2¼ 1¼ 0¼	97805. 8 97567. 3 97378. 4 97630. 0 97768. 9 97814. 3	38. 5 188. 3 -251. 6 -138. 9 -45. 4	4d <sup>6</sup> ( <sup>2</sup> F)5p	z <sup>2</sup> D <sup>o</sup>	3¼ 2¼ 1¼ 0¼	116990. 8 118247. 7 118988. 9 119418. 9	-1256. 9 -741. 2 -430. 0
4d <sup>6</sup> ( <sup>2</sup> F)5s	c <sup>2</sup> F	3¼ 2¼	97867. 8 98213. 3	-345. 5	4d <sup>6</sup> ( <sup>2</sup> F)5p	y <sup>2</sup> G <sup>o</sup>	5¼ 4¼ 3¼ 2¼	117275. 0 117596. 4 117919. 1 118129. 9	-321. 4 -322. 7 -210. 8
4d <sup>6</sup> ( <sup>2</sup> D)5p	z <sup>2</sup> P <sup>o</sup>	3¼ 2¼ 1¼	99048. 9 100941. 8 102273. 6	-1894. 9 -1331. 8	4d <sup>6</sup> ( <sup>2</sup> S)5s <sup>2</sup>	b <sup>2</sup> S	0¼	118210. 27	
4d <sup>6</sup> ( <sup>2</sup> D)5p	z <sup>2</sup> F <sup>o</sup>	4¼ 3¼ 2¼ 1¼	100767. 3 102446. 8 103513. 9 103920. 2	-1689. 5 -867. 1 -666. 3	4d <sup>6</sup> ( <sup>2</sup> H)5p	s <sup>2</sup> I <sup>o</sup>	6¼ 5¼	118277. 6 118955. 4	-675. 8
					4d <sup>6</sup> ( <sup>2</sup> F)5p	z <sup>2</sup> F <sup>o</sup>	3¼ 2¼	118850. 2 119481. 2	-631. 0

Rh III—Continued

Rh III—Continued

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
4d <sup>9</sup> ( <sup>9</sup> H)5p	s <sup>1</sup> H°	5½ 4½	119734.0 120225.0	-1151.0	4d <sup>9</sup> ( <sup>1</sup> D)5p	w <sup>1</sup> D°	1½ 2½	132362.9 132622.8	268.9
4d <sup>9</sup> ( <sup>9</sup> G)5p	s <sup>1</sup> G°	5½ 4½ 3½ 2½	119241.7 120052.7 121022.0 121225.7	-211.0 -1002.3 -763.7	4d <sup>9</sup> ( <sup>1</sup> D)5p	w <sup>1</sup> P°	0½ 1½	132214.5 132752.0	544.5
4d <sup>9</sup> ( <sup>9</sup> F)5p	y <sup>1</sup> G°	4½ 3½	119222.0 120202.6	-902.6	4d <sup>9</sup> ( <sup>1</sup> F)5p	v <sup>1</sup> D°	2½ 1½	142222.5 142222.6	-337.1
4d <sup>9</sup> ( <sup>9</sup> F)5p	s <sup>1</sup> P°	1½ 0½	120214.3 122211.4	-1227.1	4d <sup>9</sup> ( <sup>1</sup> F)5p	v <sup>1</sup> G°	3½ 4½	142222.6 144224.3	392.7
4d <sup>9</sup> ( <sup>9</sup> G)5p	s <sup>1</sup> F°	4½ 3½ 2½ 1½	122221.5 122227.4 122222.2 122222.2	-1002.9 -423.4	4d <sup>9</sup> ( <sup>1</sup> F)5p	u <sup>1</sup> F°	3½ 2½	142222.0 147221.3	-532.3
4d <sup>9</sup> ( <sup>9</sup> G)5p	y <sup>1</sup> H°	6½ 5½ 4½ 3½	122222.2 122222.2 122222.0 122407.3	-222.3 -122.8 -44.3	4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	w <sup>1</sup> P°	0½ 1½ 2½	142222.2 142222.2 147222.2	222.7 722.3
4d <sup>9</sup> ( <sup>9</sup> F)5p	y <sup>1</sup> D°	2½ 1½	124402.1 124277.6	-574.5	4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	v <sup>1</sup> D°	0½ 1½ 2½ 3½	142222.1 147222.4 147741.2 147222.2	322.3 422.2 222.3
4d <sup>9</sup> ( <sup>9</sup> F)5p	s <sup>1</sup> S°	0½	122222.2		4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	w <sup>1</sup> G°	2½ 3½ 4½ 5½	142222.1 142222.3 142222.1 142222.2	222.2 214.2 242.7
4d <sup>9</sup> ( <sup>9</sup> G)5p	y <sup>1</sup> F°	2½ 3½	122240.7 127242.2	922.1	4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	y <sup>1</sup> S°	1½	142724.2	
4d <sup>9</sup> ( <sup>9</sup> D)5p	s <sup>1</sup> P°	0½ 1½ 2½	122442.1 122222.2 127147.2	322.4 324.3	4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	u <sup>1</sup> D°	0½ 1½ 2½ 3½	151242.2 151242.2 152222.2 152222.2	122.2 472.7 -52.4
4d <sup>9</sup> ( <sup>9</sup> G)5p	y <sup>1</sup> H°	5½ 4½	122222.2 127242.2	-1422.2	4d <sup>9</sup> ( <sup>1</sup> F)5p <sup>1</sup>	u <sup>1</sup> D°	1½ 2½	152222.4 152222.2	322.4
4d <sup>9</sup> ( <sup>9</sup> D)5p	w <sup>1</sup> D°	0½ 1½ 2½ 3½	127222.2 122222.2 122222.2 122222.2	222.1 222.2 112.3	4d <sup>9</sup> ( <sup>1</sup> D)6s	e <sup>1</sup> D	4½ 3½ 2½ 1½ 0½	152722.0?	
4d <sup>9</sup> ( <sup>9</sup> D)5p	w <sup>1</sup> F°	1½ 2½ 3½ 4½	122222.4 122222.2 122122.2 122222.2	122.1 117.7 422.7	4d <sup>9</sup> ( <sup>1</sup> F)5p <sup>1</sup>	u <sup>1</sup> G°	4½ 3½	152222.2 152222.2	-322.2
4d <sup>9</sup> ( <sup>1</sup> I)5p	s <sup>1</sup> K°	6½ 7½	122141.0 122222.2	1422.2	4d <sup>9</sup> ( <sup>1</sup> F)5p <sup>1</sup>	v <sup>1</sup> F°	1½ 2½ 3½ 4½	152401.2 152222.2 152222.2 152222.2	22.0 -4.4 422.0
4d <sup>9</sup> ( <sup>9</sup> G)5p	s <sup>1</sup> G°	4½ 3½	122422.2 122222.2	-22.2	4d <sup>9</sup> ( <sup>1</sup> F)5p <sup>1</sup>	t <sup>1</sup> F°	2½ 3½	154222.2 152222.2	522.0
4d <sup>9</sup> ( <sup>1</sup> I)5p	s <sup>1</sup> H°	4½ 5½	122402.4 122222.2	522.0	4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	t <sup>1</sup> D°	1½ 2½	154222.2 152222.2	222.2
4d <sup>9</sup> ( <sup>1</sup> G)5p	w <sup>1</sup> G°	4½ 3½	122122.4 122222.1	-122.2	4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	v <sup>1</sup> P°	0½ 1½	152222.2 152222.0	344.4
4d <sup>9</sup> ( <sup>1</sup> G)5p	s <sup>1</sup> F°	3½ 2½	122222.2 122222.2	-222.2	4d <sup>9</sup> ( <sup>1</sup> P)5p <sup>1</sup>	y <sup>1</sup> S°	0½	152222.2	
4d <sup>9</sup> ( <sup>9</sup> D)5p	y <sup>1</sup> P°	0½ 1½	122222.2 122222.2	372.2	4d <sup>9</sup> ( <sup>1</sup> G)5p <sup>1</sup>	s <sup>1</sup> F°	3½ 2½	122222.2 122222.2	-522.2
4d <sup>9</sup> ( <sup>1</sup> G)5p	w <sup>1</sup> H°	5½ 4½	122222.2 122222.2	-222.2	4d <sup>9</sup> ( <sup>1</sup> G)5p <sup>1</sup>	t <sup>1</sup> G°	4½ 3½	122222.2 122222.2	-1122.4
4d <sup>9</sup> ( <sup>9</sup> D)5p	s <sup>1</sup> D°	1½ 2½	122222.2 122222.2	422.2	4d <sup>9</sup> ( <sup>1</sup> G)5p <sup>1</sup>	v <sup>1</sup> H°	5½ 4½	122222.2 122222.2	-222.2
4d <sup>9</sup> ( <sup>1</sup> I)5p	y <sup>1</sup> I°	6½ 5½	122222.2 122222.2	-222.2	4d <sup>9</sup> ( <sup>1</sup> D)5p <sup>1</sup>	s <sup>1</sup> D°	2½ 1½	122222.2 122222.2	-222.2
4d <sup>9</sup> ( <sup>9</sup> D)5p	w <sup>1</sup> F°	3½ 2½	122222.2 122222.2	-122.2	4d <sup>9</sup> ( <sup>1</sup> D)5p <sup>1</sup>	u <sup>1</sup> P°	1½ 0½	122222.2 122222.2	-222.2
4d <sup>9</sup> ( <sup>1</sup> S)5p	s <sup>1</sup> P°	1½ 0½	122222.2 122222.2	-222.2	4d <sup>9</sup> ( <sup>1</sup> D)5p <sup>1</sup>	r <sup>1</sup> F°	3½ 2½	122222.2 122222.2	-222.2
4d <sup>9</sup> ( <sup>1</sup> D)5p	v <sup>1</sup> F°	2½ 3½	122222.2 122222.2	422.0					
					Rh rv( <sup>1</sup> D <sub>1</sub> )	Limit		222222.0	



Rh III OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>1</sup> +	Observed Terms					
	a <sup>1</sup> P a <sup>3</sup> P	a <sup>1</sup> D a <sup>3</sup> D	a <sup>1</sup> F a <sup>3</sup> F	a <sup>1</sup> G a <sup>3</sup> G	a <sup>1</sup> H a <sup>3</sup> H	a <sup>1</sup> I a <sup>3</sup> I
4d <sup>7</sup>						
4d <sup>6</sup> (1D)ns		a <sup>1</sup> D a <sup>3</sup> D				
4d <sup>6</sup> (3F)ns	b <sup>1</sup> P b <sup>3</sup> P					
4d <sup>6</sup> (3H)ns				a <sup>1</sup> H b <sup>3</sup> H		
4d <sup>6</sup> (3F)ns			b <sup>1</sup> F b <sup>3</sup> F			
4d <sup>6</sup> (3G)ns				a <sup>1</sup> G b <sup>3</sup> G		
4d <sup>6</sup> (1D)ns		b <sup>1</sup> D c <sup>3</sup> D				
4d <sup>6</sup> (1I)ns					a <sup>1</sup> I	
4d <sup>6</sup> (1G)ns	a <sup>1</sup> S			c <sup>1</sup> D		
4d <sup>6</sup> (3S)ns						
4d <sup>6</sup> (1D)ns		d <sup>1</sup> D				
4d <sup>6</sup> (3F)ns			c <sup>1</sup> F			
4d <sup>6</sup> (3P)ns <sup>†</sup>	c <sup>1</sup> P c <sup>3</sup> P					
4d <sup>6</sup> (3F)ns <sup>†</sup>			c <sup>1</sup> F d <sup>3</sup> F			
4d <sup>6</sup> (1G)ns <sup>†</sup>				d <sup>1</sup> G		
4d <sup>6</sup> (1D)ns <sup>†</sup>	b <sup>1</sup> S	e <sup>1</sup> D				
4d <sup>6</sup> (3S)ns <sup>†</sup>						

\*For predicted terms in the spectra of the Tc I isoelectronic sequence, see Vol. III, Introduction.  
 †This entry refers to the higher limit term of the given type, for this configuration.

## PALLADIUM

## Pd I

46 electrons

Z=46

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4s$  $4d^{10} 1S_0$ , 67296.0 K

I. P. 8.33 volts

The analysis is from Shenstone, who, in 1930, revised and extended the earlier work by McLennan and Smith, Bechert and Catalán, Meggers and Laporte, and others. With the aid of recent ultraviolet observations he has carried it still further, and furnished the additional levels especially for inclusion here. There are 390 classified lines between 1723.65 Å and 9234.02 Å. In the 1930 term table he gives observed  $g$ -values based on all available material. More recent Zeeman data from the Massachusetts Institute of Technology have been published by Lindsley and Rosen, who also derived more accurate values for some of the energy levels. In the table, the two-place entries for energy levels are from their paper. The first column in the table (labeled "S") gives Shenstone's notation.

In his 1953 paper, Shenstone discusses forbidden transitions in the Pd I spectrum. He explains the level previously called  $K_1$  (54885.7), as resulting from "forced" dipole radiation.

The singlet and triplet systems of terms are connected by observed intersystem combinations.

Mack has noted that the sequence Pd I—Sn V shows a gradual transition from  $LS$ - to  $jj$ -coupling for levels from the  $d^9 s$  and  $d^9 p$  configurations. By analogy with Cu II, the writer has prepared the following table giving the  $JL$ -coupling notation in the form suggested by Racah.

Author	Config.	Desig.	J	Author	Config.	Desig.	J
$nd^9 1D_2$	$4d^9(^1D_{20})nd$	$nd$ [2½]	3	$nd^9 1S_0$	$4d^9(^1D_{20})nd$	$nd$ [0½]	1
$nd^9 1D_2$			2	$nd^9 1G_2$	"	$nd$ [4½]	5
$nd^9 1D_1$	$4d^9(^1D_{10})nd$	$nd'$ [1½]	1	$nd^9 1G_1$	"	$nd$ [4½]	4
$nd^9 1D_1$			2	$nd^9 1F_2$	"	$nd$ [1½]	2
$np^5 1P_1$	$4d^9(^1D_{20})np$	$np$ [1½] <sup>*</sup>	2	$nd^9 1F_1$	"	$nd$ [1½]	1
$np^5 1P_1$			1	$nd^9 1D_2$	"	$nd$ [2½]	3
$np^5 1F_1$	"	$np$ [3½] <sup>*</sup>	3	$nd^9 1D_1$	"	$nd$ [2½]	2
$np^5 1F_1$			4	$nd^9 1F_2$	"	$nd$ [3½]	3
$np^5 1D_1$	"	$np$ [2½] <sup>*</sup>	3	$nd^9 1F_1$	"	$nd$ [3½]	4
$np^5 1D_1$			2	$nd^9 1F_0$	"	$nd$ [0½]	0
$np^5 1P_1$	$4d^9(^1D_{20})np$	$np'$ [0½] <sup>*</sup>	0	$nd^9 1P_1$	$4d^9(^1D_{20})nd$	$nd'$ [0½]	1
$np^5 1P_1$			1	$nd^9 1G_2$	"	$nd'$ [3½]	3
$np^5 1F_1$	"	$np'$ [2½] <sup>*</sup>	2	$nd^9 1G_1$	"	$nd'$ [3½]	4
$np^5 1F_1$			3	$nd^9 1D_1$	"	$nd'$ [1½]	1
$np^5 1D_1$	"	$np'$ [1½] <sup>*</sup>	1	$nd^9 1D_2$	"	$nd'$ [1½]	2
$np^5 1D_1$			2	$nd^9 1F_2$	"	$nd'$ [2½]	2
				$nd^9 1F_1$	"	$nd'$ [2½]	3
				$nd^9 1S_0$	"	$nd'$ [0½]	0



Pd I—Continued

Pd I—Continued

S	Config.	Design.	J	Level	Interval	Obs. <i>g</i>	S	Config.	Design.	J	Level	Interval	Obs. <i>g</i>
26 <sub>1</sub>	4d <sup>9</sup> 5s( ) 5p		2	60069. 5			8s <sup>2</sup> D <sub>3</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 8s	8s <sup>2</sup> D	2	61736. 2		
27 <sub>1</sub>	4d <sup>9</sup> 5s( ) 5p		2	60224. 4						1			
6d <sup>4</sup> G <sub>1</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d	6d <sup>4</sup> G	1	60224. 8			s <sup>1</sup> F <sub>1</sub>	4d <sup>9</sup> 5s( <sup>2</sup> F) 5p	5p' <sup>1</sup> F°	3	62316. 2		1. 06
28 <sub>1,2</sub>	4d <sup>9</sup> 5s( ) 5p		2, 4	60310. 3			s <sup>1</sup> D <sub>1</sub>	4d <sup>9</sup> 5s( <sup>2</sup> F) 5p	5p' <sup>1</sup> D°	2	62626. 2		
6d <sup>4</sup> G <sub>3</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d	6d <sup>4</sup> G	5	60316. 4	-2. 0		31°	4d <sup>9</sup> 5s( ) 5p		27	62824. 4†		
6d <sup>4</sup> G <sub>2</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d		4	60318. 4	-3534. 6		32 <sub>1</sub>	4d <sup>9</sup> 5s( ) 5p		3	62476. 4		
6d <sup>4</sup> G <sub>1</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d		3	62853. 0			33 <sub>1,2</sub>	4d <sup>9</sup> 5s( ) 5p		1, 2	62662. 3†		
30°	4d <sup>9</sup> ( ) 4f†			60317. 1†			9s <sup>2</sup> D <sub>3</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 9s	9s <sup>2</sup> D	2	63671. 7†		
6d <sup>2</sup> F <sub>3</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d	6d <sup>2</sup> F	2	60322. 8						1			
			0	60324. 6						2			
6d <sup>2</sup> D <sub>3</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d	6d <sup>2</sup> D	3	60370. 4	-26. 0		6d <sup>1</sup> G <sub>4</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d	6d <sup>1</sup> G	4	62872. 7		
6d <sup>2</sup> D <sub>1</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d		2	60396. 4	-3492. 9		6d <sup>1</sup> F <sub>3</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d	6d <sup>1</sup> F	3	62939. 8		
6d <sup>2</sup> D <sub>1</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d		1	62866. 3			34 <sub>1,2</sub>	4d <sup>9</sup> 5s( ) 5p		1, 2	62961. 5†		
6d <sup>2</sup> F <sub>4</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d	6d <sup>2</sup> F	4	60404. 0	6. 1		35 <sub>1,2</sub>	4d <sup>9</sup> 5s( ) 5p		1, 2	64266. 0†		
6d <sup>2</sup> F <sub>3</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d		3	60397. 9	-3589. 5								
6d <sup>2</sup> F <sub>2</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6d		2	62937. 4									
29 <sub>1</sub>	4d <sup>9</sup> 5s( ) 5p		2	60655. 2									
s <sup>1</sup> G <sub>4</sub>	4d <sup>9</sup> 5s( <sup>2</sup> F) 5p	5p' <sup>1</sup> G°	4	60722. 7		1. 20†		Pd II( <sup>2</sup> D <sub>3/2</sub> )	Limit	---	67226. 6		
18 <sub>1,2</sub>	4d <sup>9</sup> 5s( ) 5p		1, 2	60722. 5				Pd II( <sup>2</sup> D <sub>1/2</sub> )	Limit	---	70775. 0		
7s <sup>1</sup> D <sub>2</sub>	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 7s	7s <sup>1</sup> D	2	61638. 6									

February 1955.

Pd I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>9</sup> 4s <sup>1</sup> 4p <sup>4</sup> +	Observed Terms	
4d <sup>9</sup>	4d <sup>9</sup> <sup>1</sup> S	
4d <sup>9</sup> 5s		5s <sup>2</sup> F
	ns (n ≥ 5)	np (n ≥ 5)
4d <sup>9</sup> ( <sup>2</sup> D) ns	{ 5-9s <sup>2</sup> D 5-7s <sup>1</sup> D	5, 6p <sup>2</sup> P° 5, 6p <sup>1</sup> D° 5, 6p <sup>2</sup> F° 5p <sup>1</sup> P° 5p <sup>1</sup> D° 5, 6p <sup>1</sup> F°
4d <sup>9</sup> 5s( <sup>2</sup> F) ns'		5p' <sup>1</sup> D° 5p' <sup>1</sup> F° 5p' <sup>1</sup> G°
	nd (n ≥ 5)	
4d <sup>9</sup> ( <sup>2</sup> D) ns	{ 5, 6d <sup>4</sup> S 5, 6d <sup>2</sup> P 5, 6d <sup>1</sup> D 5, 6d <sup>2</sup> F 5, 6d <sup>2</sup> G 5d <sup>1</sup> S 5d <sup>1</sup> P 5d <sup>1</sup> D 5, 6d <sup>1</sup> F 5, 6d <sup>1</sup> G	

\*For predicted terms in the spectra of the Pd I isoelectronic sequence, see Vol. III, Introduction.

## Pd II

(Rh I sequence; 45 electrons)

Z=46

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^9 {}^1D_{3/2}$  $4d^9 {}^1D_{3/2}$  156700 K

I. P. 19.42 volts

In 1928 Shenstone revised and extended the earlier work on this spectrum, and published 356 classified lines in the interval 1115.16 Å to 4156.95 Å. This list is from observations by himself and others. Those in the vacuum region are by H. E. White, who located the ground term.

Blair used a Schüller tube to observe the spectrum more completely, in order to extend the analysis of terms from the  $4d^9 6s$  and  $4d^9 5d$  configurations. His list covers the range from 2124.33 Å to 3382.57 Å, and includes 257 classified lines. About 60 lines are common to both published lists. Blair improved the values of eleven higher levels given by Shenstone, and added 35 new levels. He revised the levels at 112872.7 K and 114940.2 K, designated by Shenstone  $5d {}^4G_5$  and  $6s {}^4P_3$ , respectively, retaining the latter level as a miscellaneous one labeled  $sd 3_{3/2}$ , and rejecting the former one.

The  $g$ -values in the table are from the 1941 paper with the exception of four entries for high levels, which are from Shenstone. Three place  $g$ -values are, in general, from resolved patterns observed at the Massachusetts Institute of Technology.

Shenstone determined the approximate limit 160605 K from the  $5,6s$  'F' series. This was recently revised by Catalán and Rico from a study of the series in second spectra from Sr II to Cd II. The revised value is quoted here. Blair comments that the ' $F_{6s}$ ', ' $F_{5s}$ ' levels "appear to be approaching the same limit as would be expected with the inverted type of convergence pointed out by Shenstone . . ." This type of convergence is discussed under Ni I in the present volumes.

Observed intersystem combinations connect the doublet and quartet systems of terms.

## REFERENCES

- A. G. Shenstone, *Nature* 123, 727 (1928).  
 A. G. Shenstone, *Phys. Rev.* 33, 30 (1928). (I P) (T) (C L) (Z E)  
 H. A. Blair, *Phys. Rev.* 34, 173 (1930). (T) (C L)  
 C. H. Lindaley and N. Rosen, *J. Opt. Soc. Am.* 31, 531 (1941). (Z E)  
 M. A. Catalán y F. R. Rico, *An. Real Soc. Esp. de Física y Química (Madrid)* [A] 48, 328 (1952) and letter (December 1956). (I P)

Pd II

Pd II

Authors	Config.	Design.	J	Level	Interval	Obs. p	Authors	Config.	Design.	J	Level	Interval	Obs. p
4d <sup>2</sup> D <sub>2</sub> 4d <sup>2</sup> D <sub>1</sub>	4d <sup>2</sup>	4d <sup>2</sup> D	3 1/2 1 1/2	0 3539	-3539		bp <sup>2</sup> D <sub>1</sub> bp <sup>2</sup> D <sub>2</sub> bp <sup>2</sup> D <sub>3</sub> bp <sup>2</sup> D <sub>4</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> D°	0 1/2 1 1/2 2 1/2 3 1/2	81806.0 82333.7 82460.0 83066.3	528.7 118.3 606.3	0.049 1.17 1.233 1.22
5s <sup>2</sup> F <sub>1</sub> 5s <sup>2</sup> F <sub>2</sub> 5s <sup>2</sup> F <sub>3</sub> 5s <sup>2</sup> F <sub>4</sub>	4d <sup>2</sup> (F)5s	5s <sup>2</sup> F	4 1/2 3 1/2 2 1/2 1 1/2	25081.0 27094.3 28927.0 29945.6	-2013.3 -1832.7 -1018.6	1.330 1.248 1.031 0.424	kp <sup>2</sup> H <sub>1</sub> kp <sup>2</sup> H <sub>2</sub>	4d <sup>2</sup> (G)5p	5p <sup>2</sup> H°	4 1/2 5 1/2	83789.7 85592.9	1803.2	0.91 1.09
5s <sup>2</sup> F <sub>1</sub> 5s <sup>2</sup> F <sub>2</sub>	4d <sup>2</sup> (F)5s	5s <sup>2</sup> F	3 1/2 2 1/2	32277.9 34422.0	-2144.1	1.148 1.006	cp <sup>2</sup> D <sub>1</sub> cp <sup>2</sup> D <sub>2</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> D°	2 1/2 1 1/2	83802.6 84572.3	-569.8	1.19 0.834
5s <sup>2</sup> P <sub>1</sub> 5s <sup>2</sup> P <sub>2</sub> 5s <sup>2</sup> P <sub>3</sub>	4d <sup>2</sup> (P)5s	5s <sup>2</sup> P	2 1/2 1 1/2 0 1/2	36281.5 37365.0 38502.4	-1083.5 -1137.4	1.323 1.402 2.666	cp <sup>2</sup> S <sub>1</sub> cp <sup>2</sup> P <sub>1</sub> cp <sup>2</sup> P <sub>2</sub>	4d <sup>2</sup> (P)5p	5p <sup>2</sup> S° 5p <sup>2</sup> P°	0 1/2 1 1/2 0 1/2	85070.7 85151.0 86038.1	-2887.1	1.839 1.411 0.70
5s <sup>2</sup> D <sub>1</sub> 5s <sup>2</sup> D <sub>2</sub>	4d <sup>2</sup> (D)5s	5s <sup>2</sup> D	1 1/2 2 1/2	39571.1 41198.3	1627.2	1.165 1.239	kp <sup>2</sup> F <sub>1</sub> kp <sup>2</sup> F <sub>2</sub>	4d <sup>2</sup> (G)5p	5p <sup>2</sup> F°	3 1/2 2 1/2	86043.1 87120.4	-1077.3	1.09 0.927
5s <sup>2</sup> P <sub>1</sub> 5s <sup>2</sup> P <sub>2</sub>	4d <sup>2</sup> (P)5s	5s <sup>2</sup> P	1 1/2 0 1/2	43647.8 43939.8	-292.0	1.220 0.681	bp <sup>2</sup> G <sub>1</sub> kp <sup>2</sup> G <sub>1</sub> kp <sup>2</sup> G <sub>2</sub>	4d <sup>2</sup> (P)5p	5p <sup>2</sup> S° 5p <sup>2</sup> G°	1 1/2 3 1/2 4 1/2	86279.6 86622.5 86622.4	299.9	2.00 0.89 1.11
5s <sup>2</sup> G <sub>1</sub> 5s <sup>2</sup> G <sub>2</sub>	4d <sup>2</sup> (G)5s	5s <sup>2</sup> G	4 1/2 3 1/2	44505.7 44613.9	-108.2	1.099 0.886	6s <sup>2</sup> F <sub>1</sub> 6s <sup>2</sup> F <sub>2</sub> 6s <sup>2</sup> F <sub>3</sub> 6s <sup>2</sup> F <sub>4</sub>	4d <sup>2</sup> (F)6s	6s <sup>2</sup> F	4 1/2 3 1/2 2 1/2 1 1/2	104616.0 105308.2 108147.0 109330.0	-692.2 -2838.8 -1183.0	
ap <sup>2</sup> D <sub>1</sub> ap <sup>2</sup> D <sub>2</sub> ap <sup>2</sup> D <sub>3</sub> ap <sup>2</sup> D <sub>4</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> D°	3 1/2 2 1/2 1 1/2 0 1/2	65847.0 67962.2 70119.2 71178.9	-2715.2 -2157.6 -1059.1	1.40 1.328 1.168 0.038	6s <sup>2</sup> F <sub>1</sub> 6s <sup>2</sup> F <sub>2</sub> 6s <sup>2</sup> F <sub>3</sub>	4d <sup>2</sup> (F)6s	6s <sup>2</sup> F	3 1/2 2 1/2	108146.0 109928.5	-1782.5	
ap <sup>2</sup> G <sub>1</sub> ap <sup>2</sup> G <sub>2</sub> ap <sup>2</sup> G <sub>3</sub> ap <sup>2</sup> G <sub>4</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> G°	5 1/2 4 1/2 3 1/2 2 1/2	69611.5 67299.0 69610.2 71069.5	1312.5 -2311.2 -1459.3	1.28 1.17 1.069 0.718	5d <sup>2</sup> D <sub>1</sub> 5d <sup>2</sup> D <sub>2</sub>	4d <sup>2</sup> (F)5d	5d <sup>2</sup> D	3 1/2 2 1/2 1 1/2 0 1/2	110204.5 110621.9	-617.4	
ap <sup>2</sup> F <sub>1</sub> ap <sup>2</sup> F <sub>2</sub> ap <sup>2</sup> F <sub>3</sub> ap <sup>2</sup> F <sub>4</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> F°	4 1/2 3 1/2 2 1/2 1 1/2	69878.4 71244.7 73112.4 78348.0	-1366.3 -1867.7 764.4	1.28 1.11 1.047 0.724	5d <sup>2</sup> F <sub>1</sub> 5d <sup>2</sup> F <sub>2</sub> 5d <sup>2</sup> F <sub>3</sub> 5d <sup>2</sup> F <sub>4</sub>	4d <sup>2</sup> (F)5d	5d <sup>2</sup> F	4 1/2 3 1/2 2 1/2 1 1/2	110886.6 111090.3 114228.2 115449.7	-203.7 -3137.9 -1221.5	
ap <sup>2</sup> G <sub>1</sub> ap <sup>2</sup> G <sub>2</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> G°	4 1/2 3 1/2	72285.0 74318.7	-2033.7	1.15 0.99	ed <sup>2</sup> G <sub>1</sub> ed <sup>2</sup> G <sub>2</sub> ed <sup>2</sup> G <sub>3</sub> ed <sup>2</sup> G <sub>4</sub>	4d <sup>2</sup> (F)5d	5d <sup>2</sup> G	5 1/2 4 1/2 3 1/2 2 1/2	110910.3 111247.0 113835.7	-336.7 -2588.7	
ap <sup>2</sup> D <sub>1</sub> ap <sup>2</sup> D <sub>2</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> D°	2 1/2 1 1/2	72732.6 75098.0	-2365.4	1.020 0.920	ed <sup>2</sup> H <sub>1</sub> ed <sup>2</sup> H <sub>2</sub>	4d <sup>2</sup> (F)5d	5d <sup>2</sup> H	6 1/2 5 1/2 4 1/2 3 1/2	110986.8 110624.5	362.3	
ap <sup>2</sup> F <sub>1</sub> ap <sup>2</sup> F <sub>2</sub>	4d <sup>2</sup> (F)5p	5p <sup>2</sup> F°	3 1/2 2 1/2	73327.5 75753.6	-2426.1	1.09 1.115	1 <sub>1/2</sub>			3 1/2	114001.4		
bp <sup>2</sup> P <sub>1</sub> bp <sup>2</sup> P <sub>2</sub> bp <sup>2</sup> P <sub>3</sub>	4d <sup>2</sup> (P)5p	5p <sup>2</sup> P°	2 1/2 1 1/2 0 1/2	76767.4 76764.5 76807.4	12.9 -52.9	1.34 1.317 2.266	ed <sup>2</sup> F <sub>1</sub> ed <sup>2</sup> F <sub>2</sub> ed <sup>2</sup> F <sub>3</sub>	4d <sup>2</sup> (F)5d	5d <sup>2</sup> F?	3 1/2 2 1/2	114241.0		
cp <sup>2</sup> F <sub>1</sub> cp <sup>2</sup> F <sub>2</sub>	4d <sup>2</sup> (D)5p	5p <sup>2</sup> F°	2 1/2 3 1/2	78765.3 79708.0	942.7	1.056 1.25	ed <sup>2</sup> 2 <sub>1/2</sub> ed <sup>2</sup> 3 <sub>1/2</sub>	4d <sup>2</sup> (F)5d		2 1/2 2 1/2	114647.6 114940.2		
cp <sup>2</sup> P <sub>1</sub> cp <sup>2</sup> P <sub>2</sub>	4d <sup>2</sup> (D)5p	5p <sup>2</sup> P°	0 1/2 1 1/2	79612.4 80956.1	1343.7	1.09 1.130	ed <sup>2</sup> 4 <sub>1/2</sub>	4d <sup>2</sup> (F)5d		0 1/2	115426.9		
cp <sup>2</sup> D <sub>1</sub> cp <sup>2</sup> D <sub>2</sub>	4d <sup>2</sup> (D)5p	5p <sup>2</sup> D°	1 1/2 2 1/2	81557.2 82057.0	499.8	1.186 1.20							

Pd II—Continued

Authors	Config.	Design.	J	Level	Interval	Obs. <i>g</i>
ed 5 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> F)5d		3½	115551.5		
ed 6 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> F)5d		3½	115771.2		
6s <sup>1</sup> P <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> F)6s	6s' <sup>1</sup> P	2½	116511.6	-1633.0	
6s <sup>1</sup> P <sub>3/2</sub>			1½	118144.6	-773.8	
6s <sup>1</sup> P <sub>3/2</sub>			0½	118918.4		
ed 7 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> F)5d		2½	117149.7		
ed 8 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> F)5d		1½	117332.1		
ed 9 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> F)5d		2½	118648.5		
6s <sup>1</sup> D <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> D)6s	6s'' <sup>1</sup> D	1½	118781.7	612.9	
6s <sup>1</sup> D <sub>3/2</sub>			2½	119394.6		
6s <sup>1</sup> F <sub>3/2</sub>	4d <sup>9</sup> ( <sup>1</sup> F)6s	6s' <sup>1</sup> F	0½	119080.6	663.0	
6s <sup>1</sup> F <sub>5/2</sub>			1½	119743.6		
6s <sup>1</sup> G <sub>3/2</sub>	4d <sup>9</sup> ( <sup>1</sup> G)6s	6s''' <sup>1</sup> G	4½	122706.0	-12.5	
6s <sup>1</sup> G <sub>5/2</sub>			3½	122718.5		
fd 10 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> D)5d		1½	123755.0		
fd 11 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> D)5d		1½	123866.1		
fd 12 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> D)5d		2½	124149.6		
fd 13 <sub>1/2</sub>	4d <sup>9</sup> ( <sup>1</sup> D)5d		2½	124280.3		
fd 14	4d <sup>9</sup> ( <sup>1</sup> D)5d			124930.0		
gd <sup>1</sup> D	4d <sup>9</sup> ( <sup>1</sup> G)5d	5d''' <sup>1</sup> D	2½	12885.2		
			1½			
gd <sup>1</sup> F <sub>3/2</sub>	4d <sup>9</sup> ( <sup>1</sup> G)5d	5d''' <sup>1</sup> F	3½	128623.6	-40.9	
gd <sup>1</sup> F <sub>5/2</sub>			2½	128664.5		
gd <sup>1</sup> G <sub>3/2</sub>	4d <sup>9</sup> ( <sup>1</sup> G)5d	5d''' <sup>1</sup> G	4½	129309.3	-73.2	
gd <sup>1</sup> G <sub>5/2</sub>			3½	129382.5		
	Pd III ( <sup>1</sup> F <sub>2</sub> )	Limit		156700		

December 1956.

Pd II OBSERVED TERMS\*

Configuration	Observed Terms
1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3d <sup>9</sup> 4s <sup>1</sup> 4p <sup>1</sup> +	
4d <sup>9</sup>	4d <sup>9</sup> <sup>1</sup> D
4d <sup>9</sup> ( <sup>1</sup> F)ns	ns (n ≥ 5)
4d <sup>9</sup> ( <sup>1</sup> F)ns'	{ 5, 6s <sup>1</sup> F 5, 6s <sup>1</sup> F
4d <sup>9</sup> ( <sup>1</sup> D)ns''	{ 5, 6s <sup>1</sup> P 5, 6s <sup>1</sup> P
4d <sup>9</sup> ( <sup>1</sup> G)ns'''	5, 6s <sup>1</sup> <sup>1</sup> D
	5, 6s <sup>1</sup> <sup>1</sup> G
	np (n ≥ 5)
	5p <sup>1</sup> D° 5p <sup>1</sup> F° 5p <sup>1</sup> G° 5p <sup>1</sup> D° 5p <sup>1</sup> F° 5p <sup>1</sup> G° 5p' <sup>1</sup> G° 5p' <sup>1</sup> P° 5p' <sup>1</sup> D° 5p' <sup>1</sup> G° 5p' <sup>1</sup> P° 5p' <sup>1</sup> D° 5p'' <sup>1</sup> P° 5p'' <sup>1</sup> D° 5p'' <sup>1</sup> F° 5p''' <sup>1</sup> F° 5p''' <sup>1</sup> G° 5p''' <sup>1</sup> H°
	nd (n ≥ 5)
	5d <sup>1</sup> D 5d <sup>1</sup> F 5d <sup>1</sup> G 5d <sup>1</sup> H 5d <sup>1</sup> D 5d <sup>1</sup> F 5d <sup>1</sup> F 5d''' <sup>1</sup> D 5d''' <sup>1</sup> F 5d''' <sup>1</sup> G

\* For predicted terms in the spectrum of the Rh I isoelectronic sequence, see Vol. III, Introduction.

## Pd III

(Ru I sequence; 44 electrons)

Z=46

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^8 \text{ } ^3F_4$  $a \text{ } ^3F_4$ , 265600 K

I. P. 32.92 volts

The extensive analysis is by Shenstone, who has carried it out especially for inclusion here. His observations extend from 695A to 2950A. He has classified 800 lines and derived the limit 265900 from the  $ns \text{ } ^3F$  series, ( $n=5,6$ ) by means of a Ritz formula, with an assumed value of  $\alpha=4 \times 10^{-6}$ . The quoted value is from Catalán and Rico, who have obtained it by comparison of the third spectra from Y to In. Observed inter-system combinations connect the terms of different multiplicities.

Shenstone's investigation of this spectrum is so complete that only three levels "from the expected configurations" remain to be found. He emphasizes the fact that "the assignment of terms to ions is in many cases doubtful" and he feels that the levels should not be grouped into terms in the table. On account of the increased inaccuracy of all levels lower than  $a \text{ } ^3F_4$ , he suggests, also, that this level be adopted as zero.

The writer has rearranged his data to conform to the general format adopted for these Volumes. The users are urged to bear in mind the reservations made by Shenstone regarding this format and to consult him for further details.

## REFERENCES

- A. G. Shenstone, unpublished material (January 1956). (I P) (T) (C L)  
 M. A. Catalán y F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] **53**, 85 (1957). (I P)



## Pd III

## Pd III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval		
4d <sup>8</sup>	a <sup>3</sup> F	4	0.0		4d <sup>7</sup> ( <sup>3</sup> P)5p	z <sup>3</sup> G°	2	118388.6			
		3	3229.7	-3229.7			4d <sup>7</sup> ( <sup>3</sup> F)5p	z <sup>3</sup> F°	4	118449.3	-1352.6
		2	4687.3	-1457.6					3	118801.9	-1936.1
			2	118738.0							
4d <sup>8</sup>	a <sup>1</sup> D	2	10230.5		4d <sup>7</sup> ( <sup>3</sup> F)5p	z <sup>3</sup> G°	5	118508.9	-2379.7		
4d <sup>8</sup>	a <sup>3</sup> P	0	13699.1		4d <sup>7</sup> ( <sup>3</sup> F)5p	z <sup>3</sup> D°	4	114886.6	-1632.3		
		1	13470.3	-228.8			3	116518.9			
		2	14634.3	1164.0							
4d <sup>8</sup>	a <sup>1</sup> G	4	17880.4		4d <sup>7</sup> ( <sup>3</sup> F)5p	z <sup>3</sup> D°	3	115736.6	-1053.7		
4d <sup>7</sup> ( <sup>3</sup> F)5s	a <sup>3</sup> F	5	52915.9	-2172.9	4d <sup>7</sup> ( <sup>3</sup> P)5p	z <sup>3</sup> P°	2	118790.8	-776.3		
		4	55088.8	-1652.7			1	117566.6			
		3	56741.5	-1103.5			1	119227.9	815.2		
		2	57845.0	-682.3			2	120045.1	179.3		
		1	58527.3				3	120222.4			
4d <sup>7</sup> ( <sup>3</sup> F)5s	b <sup>3</sup> F	4	62560.9	-2694.5	4d <sup>7</sup> ( <sup>3</sup> G)5p	z <sup>3</sup> H°	4	122771.8	-2083.9		
		3	65255.4	-1824.0			5	120687.9	2641.4		
		2	67079.4				6	123329.3			
4d <sup>7</sup> ( <sup>3</sup> P)5s	a <sup>3</sup> P	3	65708.0	-80.3	4d <sup>7</sup> ( <sup>3</sup> G)5p	y <sup>3</sup> F°	4	121161.5	-3075.4		
		2	65788.3	-1363.1			3	124236.9	-3141.7		
		1	67151.4				2	127378.6			
4d <sup>7</sup> ( <sup>3</sup> G)5s	a <sup>3</sup> G	5	69985.8	-1061.4	4d <sup>7</sup> ( <sup>3</sup> P)5p	y <sup>3</sup> D°	0	121652.6	81.8		
		4	71047.2	-1738.9			1	121794.4	1258.7		
		3	72786.1				2	122893.1	510.3		
4d <sup>7</sup> ( <sup>3</sup> P)5s	b <sup>3</sup> P	2	72745.0	-257.6	4d <sup>7</sup> ( <sup>3</sup> P)5p	z <sup>3</sup> P°?	3	123503.4	-1603.6		
		1	73002.6	-1278.6			4	121899.8			
		0	74281.2				2	123313.1	-2164.3		
4d <sup>7</sup> ( <sup>3</sup> H)5s	a <sup>3</sup> H	6	74673.3	-1294.3	4d <sup>7</sup> ( <sup>3</sup> P)5p	z <sup>3</sup> S°	1	123779.0			
		5	75967.6	-2613.5			4d <sup>7</sup> ( <sup>3</sup> G)5p	z <sup>3</sup> G°	4	124431.9	
		4	78581.1						4d <sup>7</sup> ( <sup>3</sup> P)5p	y <sup>3</sup> D°	3
			2	124439.5	-536.6						
			1	124976.1							
4d <sup>7</sup> ( <sup>3</sup> G)5s	b <sup>3</sup> G	4	75403.0		4d <sup>7</sup> ( <sup>3</sup> H)5p	y <sup>3</sup> G°	5	124760.4	-3224.7		
							4	127985.1	-1697.4		
							3	129682.5			
4d <sup>7</sup> ( <sup>3</sup> P)5s	c <sup>3</sup> P	2	75455.0	-600.8	4d <sup>7</sup> ( <sup>3</sup> H)5p	z <sup>3</sup> I°	7	127545.8	2382.7		
		1	76055.8	-2676.7			6	125163.1	-2743.5		
		0	78732.57				5	127906.6			
4d <sup>7</sup> ( <sup>3</sup> D)5s	a <sup>3</sup> D	3	76231.4	-1938.4	4d <sup>7</sup> ( <sup>3</sup> G)5p	z <sup>3</sup> G°	5	125989.9	-1524.1		
		2	78169.8	49.8			4	126914.0	-103.5		
		1	78120.0				3	127017.6			
4d <sup>7</sup> ( <sup>3</sup> H)5s	a <sup>3</sup> H	5	80805.1		4d <sup>7</sup> ( <sup>3</sup> G)5p	z <sup>3</sup> F°	3	125894.3			
							4d <sup>7</sup> ( <sup>3</sup> P)5p	y <sup>3</sup> P°	2	126078.5	-1071.0
									1	127149.5	
			0								
4d <sup>7</sup> ( <sup>3</sup> P)5s	a <sup>3</sup> P	1	82620.3		4d <sup>7</sup> ( <sup>3</sup> G)5p	z <sup>3</sup> H°	5	127187.1			
							4d <sup>7</sup> ( <sup>3</sup> D)5p	z <sup>3</sup> D°	3	127657.9	-1140.0
									2	128797.9	-1043.5
			1	129841.4							
4d <sup>7</sup> ( <sup>3</sup> D)5s	b <sup>3</sup> D	2	83204.3		4d <sup>7</sup> ( <sup>3</sup> P)5p	w <sup>3</sup> D°	3	128038.4	-175.9		
							2	128214.3	-286.8		
							1	128501.1			
4d <sup>7</sup> ( <sup>3</sup> F)5s	c <sup>3</sup> F	2	85420.7	409.7	4d <sup>7</sup> ( <sup>3</sup> D)5p	z <sup>3</sup> F°	4	130476.3	-1568.4		
		3	85830.4	964.8			3	130044.7	1423.7		
		4	86795.2				2	130621.0			
4d <sup>7</sup> ( <sup>3</sup> F)5s	a <sup>3</sup> F	3	90684.3		4d <sup>7</sup> ( <sup>3</sup> H)5p	z <sup>3</sup> I°	6	130959.3			
							4d <sup>7</sup> ( <sup>3</sup> P)5p	z <sup>3</sup> S°	0	131482.87	
4d <sup>7</sup> ( <sup>3</sup> D)5s <sup>†</sup>	b <sup>3</sup> D	1	103529.47	20.2							
		2	103549.6	869.5							
		3	104419.1								
4d <sup>7</sup> ( <sup>3</sup> F)5p	z <sup>3</sup> F°	5	106145.4	726.7							
		4	104418.7	-2474.8							
		3	106893.5	-1748.2							
		2	108641.7	-1138.4							
		1	109730.1								
4d <sup>7</sup> ( <sup>3</sup> F)5p	z <sup>3</sup> D°	4	108115.7	-1608.8							
		3	109724.5	-1216.5							
		2	110911.0	-768.4							
		1	111709.4	-137.8							
		0	111847.2								
4d <sup>7</sup> ( <sup>3</sup> F)5p	z <sup>3</sup> G°	6	109015.4	-80.9							
		5	109076.3	-1672.7							
		4	110749.0	-882.2							
		3	111631.2	-171.7							
		2	111802.9								

## Pd III—Continued

## Pd III—Continued

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval		
4d <sup>7</sup> ( <sup>2</sup> H)5p	y <sup>2</sup> H°	6	131550.6	-749.9 -1149.7	4d <sup>7</sup> ( <sup>2</sup> F)5p	z <sup>1</sup> G°	4	141018.4	-936.4 117.1		
		5	132300.6				4d <sup>7</sup> ( <sup>2</sup> F)5p	v <sup>1</sup> D°		3	141283.7
		4	133450.8							2	142220.1†
4d <sup>7</sup> ( <sup>2</sup> F)5p	y <sup>2</sup> S°	1	132181.8	4d <sup>7</sup> ( <sup>2</sup> D)5p <sup>1</sup>	w <sup>2</sup> P°	2	153229.6	-637.5 -822.5			
4d <sup>7</sup> ( <sup>2</sup> F)5p	z <sup>1</sup> P°	1	132716.7			1	154527.0				
4d <sup>7</sup> ( <sup>2</sup> D)5p	z <sup>1</sup> D°	2	132742.9			0	155389.5				
4d <sup>7</sup> ( <sup>2</sup> F)5p	y <sup>1</sup> D°	2	132951.1	4d <sup>7</sup> ( <sup>2</sup> D)5p <sup>1</sup>	u <sup>1</sup> D°	3	156576.8	-499.2 -1455.1			
4d <sup>7</sup> ( <sup>2</sup> H)5p	y <sup>1</sup> G°	4	134638.8			2					
4d <sup>7</sup> ( <sup>2</sup> F)5p	w <sup>1</sup> G°	3	135159.3			1					
4d <sup>7</sup> ( <sup>2</sup> D)5p	z <sup>2</sup> P°	2	135218.8	3889.8 2825.1	4d <sup>7</sup> ( <sup>1</sup> F)6s	e <sup>1</sup> F	5	169831.1	-499.2 -1455.1		
		1	137488.9				4	170330.3			
		0	137740.5†				3	171785.4			
4d <sup>7</sup> ( <sup>2</sup> D)5p	y <sup>1</sup> F°	3	135356.8	4d <sup>7</sup> ( <sup>1</sup> F)6s	e <sup>2</sup> F	4	173635.6	-1937.5 -1385.1			
4d <sup>7</sup> ( <sup>2</sup> D)5p	y <sup>1</sup> P°	1	136782.3			3					
4d <sup>7</sup> ( <sup>2</sup> H)5p	y <sup>1</sup> H°	5	136847.4			2					
4d <sup>7</sup> ( <sup>2</sup> F)5p	x <sup>1</sup> F°	3	137180.6	4d <sup>8</sup> 5s( <sup>2</sup> D)5p	y <sup>2</sup> F°	5	193222.5	-1937.5 -1385.1			
4d <sup>7</sup> ( <sup>2</sup> F)5p	w <sup>2</sup> F°	2	138415.5			4	195180.0				
4d <sup>7</sup> ( <sup>2</sup> F)5p		3	140473.3			3	196545.1				
4d <sup>7</sup> ( <sup>2</sup> F)5p	z <sup>1</sup> D°	2	140563.5	2057.8 1506.8	-----	-----	-----	265600	-----		
		4	141980.1								
		4d <sup>7</sup> ( <sup>2</sup> F)5p								4	141980.1
4d <sup>7</sup> ( <sup>2</sup> F)5p	x <sup>1</sup> D°	2	140563.5								

February 1957.

## Pd III OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> +	Observed Terms			
4d <sup>8</sup>	{ a <sup>2</sup> P a <sup>1</sup> D a <sup>2</sup> F a <sup>1</sup> G			
		ns (n ≥ 5)	np (n ≥ 5)	
4d <sup>7</sup> ( <sup>1</sup> F)nz	{ a, c <sup>2</sup> F b, e <sup>2</sup> F		z <sup>2</sup> D° z <sup>1</sup> D°	z <sup>2</sup> F° z <sup>1</sup> F°
4d <sup>7</sup> ( <sup>2</sup> F)nz	{ a <sup>2</sup> P c <sup>2</sup> P		z <sup>2</sup> S° y <sup>2</sup> S°	z <sup>2</sup> P° y <sup>2</sup> P°
4d <sup>7</sup> ( <sup>2</sup> G)nz	{ a <sup>2</sup> G b <sup>1</sup> G			y <sup>2</sup> D° y <sup>1</sup> D°
4d <sup>7</sup> ( <sup>2</sup> F)nz	{ b <sup>2</sup> P a <sup>1</sup> P		z <sup>2</sup> S° z <sup>1</sup> S°	z <sup>2</sup> P° z <sup>1</sup> P°
4d <sup>7</sup> ( <sup>2</sup> H)nz	{ a <sup>2</sup> H a <sup>1</sup> H		w <sup>2</sup> D° y <sup>1</sup> D°	z <sup>2</sup> G° z <sup>1</sup> G°
4d <sup>7</sup> ( <sup>2</sup> D)nz	{ a <sup>2</sup> D b <sup>1</sup> D			z <sup>2</sup> H° z <sup>1</sup> H°
4d <sup>7</sup> ( <sup>2</sup> F)nz	{ c <sup>2</sup> F a <sup>1</sup> F			y <sup>2</sup> G° y <sup>1</sup> G°
4d <sup>7</sup> ( <sup>2</sup> D)nz <sup>1</sup>	b <sup>2</sup> D			y <sup>2</sup> H° y <sup>1</sup> H°
4d <sup>8</sup> 5s( <sup>2</sup> D)nz				z <sup>2</sup> I° z <sup>1</sup> I°
				x <sup>2</sup> P° y <sup>1</sup> P°
				z <sup>2</sup> D° z <sup>1</sup> D°
				z <sup>2</sup> F° y <sup>1</sup> F°
				w <sup>2</sup> D° z <sup>1</sup> D°
				w <sup>2</sup> F° z <sup>1</sup> F°
				w <sup>2</sup> G° z <sup>1</sup> G°
				w <sup>2</sup> P° u <sup>1</sup> D°
				y <sup>2</sup> F°

\*For predicted terms in the spectra of the Ru I isoelectronic sequence, see Vol. III, Introduction.

†This entry denotes the higher of the two <sup>1</sup>D limit terms for this configuration.

## Pd XVIII

(Cu I sequence; 29 electrons)

Z=46

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 S_{1/2}$  $4s^2 S_{1/2}$ 

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed the doublets due to the transitions  $4s-5p$ ,  $4p-5d$ ,  $4d-5f$ , and  $4p-5s$ , from Pd XVIII to In XXI. In figure 3 of his paper on the spectra of highly-ionized atoms, the lines are indicated on the photograph of vacuum spark spectra of Pd observed from 40 Å to 80 Å.

By analogy with Cu I the writer has assumed the ground state indicated above.

## REFERENCE

B. Edlén, *Physica* 13, No. 9, 549 (1947).

March 1953.

## Pd XX

(Co I sequence; 27 electrons)

Z=46

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^9 D_{3/2}$  $3d^9 D_{3/2}$ 

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed three lines due to the transition  $3p^6 3d^9 D-3p^6 3d^{10} P^o$ , in the region between 40 Å and 80 Å. In figure 4 of his paper on the spectra of highly-ionized atoms, the observed wave numbers are plotted against atomic number for this combination in the Co I-like spectra Pd XX to Sn XXIV. For Pd XX the wave numbers are between 1500000 and 1800000 K.

## REFERENCE

B. Edlén, *Physica* 13, No. 9, 548 (1947).

March 1953.

## SILVER

## Ag I

47 electrons

Z=47

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 \text{ } ^1\text{S}_{0,1}$  $5s^2 \text{ } ^1\text{S}_{0,1}$ , 61106.50 K

I. P. 7.574 volts

The early work on this spectrum by Blair and others has been revised and extended by Shenstone. The analysis is from Shenstone's 1940 paper, supplemented by Rasmussen's extension of the  $^1\text{P}^\circ$ ,  $^1\text{D}$ , and  $^1\text{F}^\circ$  series. For the terms common to the two papers, values of the latter have been used in the present compilation. Shenstone's notation is given in column 1 of the table.

About 190 lines have been classified in the range between 1507.37 Å and 39951 Å. The doublet and quartet systems of terms are connected by observed intersystem combinations.

Shenstone has determined the limit from the  $^1\text{S}$  series, by using an extended Ritz formula. For 10 levels having as limit the  $^1\text{D}$  term in Ag II, he has assigned the  $J$ -value of the limit, as is indicated in the configuration column.

The following  $g$ -values have been determined from Zeeman observations of five Ag I lines that appear as impurities on spectrograms measured at the National Bureau of Standards:

Desig.	$J$	Level	Obs. $g$
$5s \text{ } ^1\text{S}$	$0\frac{1}{2}$	0.00	1.998
$5p \text{ } ^1\text{P}^\circ$	$0\frac{1}{2}$	29552.05	0.666
	$1\frac{1}{2}$	30472.71	1.330
$6s \text{ } ^1\text{S}$	$0\frac{1}{2}$	42556.15	1.995
$5d \text{ } ^1\text{D}$	$1\frac{1}{2}$	48744.00	0.801

## REFERENCES

- A. G. Shenstone, *Phys. Rev.* **57**, 894 (1940). (I P) (T) (C L)  
 E. Rasmussen, *Det. Kgl. Danske Videnskab., Math.-fys. Medd.* (Copenhagen) **18**, 24 (1940). (T) (C L)  
 A. G. Shenstone, *Phys. Rev.* **72**, 1273 (L) (1947). (T)  
 J. E. Mack, *Rev. Mod. Phys.* **23**, No. 1, 64 (1950). (Summary hfs)  
 P. F. A. Klippenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs).  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955). (Summary hfs)

Ag I

Ag I

Author	Config.	Desig.	J	Level	Interval	Author	Config.	Desig.	J	Level	Interval
5e S <sub>14</sub>	4d <sup>0</sup> (S)5e	5e S	0%	0.00		12e S <sub>14</sub>	4d <sup>0</sup> (S)12e	12e S	0%	59574.56	
5p F <sub>14</sub>	4d <sup>0</sup> (S)5p	5p P°	0%	88652.06	920.66	11d D <sub>14</sub>	4d <sup>0</sup> (S)11d	11d D	1%	59751.15	0.36
5p F <sub>14</sub>			1%	80472.71		11d D <sub>14</sub>			2%	59751.51	
5e D <sub>14</sub>	4d <sup>0</sup> 5e?	5e D	2%	30242.28	4471.90	12d D	4d <sup>0</sup> (S)12d	12d D	1%, 2%	60007.6	
5e D <sub>14</sub>			1%	34714.16		Ag II(S <sub>6</sub> )	Limit			61106.50	
6e S <sub>14</sub>	4d <sup>0</sup> (S)6e	6e S	0%	42556.15		5e 5p D <sub>14</sub>	4d <sup>0</sup> 5e(D)5p	5p' D°	3%	61775.1	-408.5
6p F <sub>14</sub>	4d <sup>0</sup> (S)6p	6p P°	0%	48297.38	203.39	5e 5p D <sub>14</sub>			2%	62181.6	-1349.1
6p F <sub>14</sub>			1%	48600.77		5e 5p D <sub>14</sub>			1%	62630.7	-1045
5d D <sub>14</sub>	4d <sup>0</sup> (S)5d	5d D	1%	48744.00	20.22	5e 5p D <sub>14</sub>	4d <sup>0</sup> 5e(D)5p	5p' P°	2%	62953.7	2703.9
5d D <sub>14</sub>			2%	48764.22		5e 5p P <sub>14</sub>			3%	65637.6	
7e S <sub>14</sub>	4d <sup>0</sup> (S)7e	7e S	0%	51886.98		5e 5p P	4d <sup>0</sup> 5e(D)5p	5p' P°	0%, 1%	65988	
7p F <sub>14</sub>	4d <sup>0</sup> (S)7p	7p P°	0%	54040.99	80.17	5e 5p D <sub>14</sub>	4d <sup>0</sup> 5e(D)5p	5p' D°	1%	66340	632
7p F <sub>14</sub>			1%	54121.16		5e 5p D <sub>14</sub>			2%	66972.0	
6d D <sub>14</sub>	4d <sup>0</sup> (S)6d	6d D	1%	54203.13	10.47	1°	4d <sup>0</sup> 5e(D)5p	5p'' P°	3%	72334	
6d D <sub>14</sub>			2%	54213.60		2°	4d <sup>0</sup> 5e(D)5p	5p'' P°	1%	73470	
4f F	4d <sup>0</sup> (S)4f	4f P°	2%, 3%	54204.73		3°	4d <sup>0</sup> 5e(D)5p	5p'' D°	2%	73587	
8e S <sub>14</sub>	4d <sup>0</sup> (S)8e	8e S	0%	55581.29		5e 6e D <sub>14</sub>	4d <sup>0</sup> 5e(D)6e	6e' D	3%	79412.9	-751
5e 5p P <sub>14</sub>	4d <sup>0</sup> 5e(D)5p	5p' P°	2%	56223.3	-2281.4	5e 6e D <sub>14</sub>			2%	80164	-1102
5e 5p P <sub>14</sub>			1%	58504.7	-2032.8	5e 6e D <sub>14</sub>			1%	81268	-2717
5e 5p P <sub>14</sub>			0%	60637.6		5e 6e D <sub>14</sub>			0%	83982.8	
8p P <sub>14</sub>	4d <sup>0</sup> (S)8p	8p P°	0%	56620.72	39.85	e D <sub>14</sub>	4d <sup>0</sup> 5e(D)6e	6e' D	2%	82070	-2524
8p P <sub>14</sub>			1%	56660.67		e D <sub>14</sub>			1%	84594	
	4d <sup>0</sup> (S)5f	5f P°	2%, 3%	56691.40		f D <sub>14</sub>	4d <sup>0</sup> 5e(D)6e	6e'' D	2%	86484	
7d D <sub>14</sub>	4d <sup>0</sup> (S)7d	7d D	1%	56699.79	5.75				1%		
7d D <sub>14</sub>			2%	56705.54							
5g G	4d <sup>0</sup> (S)5g	5g G	3%, 4%	56711.1		11 <sub>14, 14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	11		86896	
9e S <sub>14</sub>	4d <sup>0</sup> (S)9e	9e S	0%	57425.11		12 <sub>14, 14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	12		87122.0	
	4d <sup>0</sup> (S)9p	9p P°	0%	58006.80	21.90	13 <sub>14, 14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	13		87144	
9p P <sub>14</sub>			1%	58027.10		14 <sub>14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	14	2%	87222	
8d D <sub>14</sub>	4d <sup>0</sup> (S)8d	8d D	1%	58050.01	3.47	15 <sub>14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	15	3%	87312.0	
8d D <sub>14</sub>			2%	58063.48		16 <sub>14, 14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	16		87356.6	
	4d <sup>0</sup> (S)6f	6f P°	2%, 3%	58065.66		17	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	17	3%?	88594	
10e S <sub>14</sub>	4d <sup>0</sup> (S)10e	10e S	0%	58478.13		18 <sub>14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	18	2%	88738	
5e 5p F <sub>14</sub>	4d <sup>0</sup> 5e(D)5p	5p' F°	4%	58901.9	112.2	19 <sub>14, 14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )5d	19		88816	
5e 5p F <sub>14</sub>			3%	58729.7	-531.4	5e 7e D <sub>14</sub>	4d <sup>0</sup> 5e(D <sub>2</sub> )7e	7e' D	3%	90366	
5e 5p F <sub>14</sub>			2%	59321.1	-1920.8				2%		
5e 5p F <sub>14</sub>			1%	61241.9					1%		
	4d <sup>0</sup> (S)10p	10p P°	0%	58834.26	15.58				0%		
			1%	58849.83							
9d D <sub>14</sub>	4d <sup>0</sup> (S)9d	9d D	1%	58862.46	2.20						
9d D <sub>14</sub>			2%	58864.66							
11e S <sub>14</sub>	4d <sup>0</sup> (S)11e	11e S	0%	59135.99		20 <sub>14, 14</sub>	4d <sup>0</sup> 5e?	20		90612	
10d D <sub>14</sub>	4d <sup>0</sup> (S)10d	10d D	1%	59388.97	1.75						
10d D <sub>14</sub>			2%	59390.72							

## Ag I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> +	Observed Terms					
	4d <sup>9</sup> 5s <sup>1</sup>	5s <sup>1</sup> <sup>2</sup> D				
ns (n ≥ 5)		np (n ≥ 5)		nd (n ≥ 5)	nf (n ≥ 4)	ng (n ≥ 5)
4d <sup>9</sup> ( <sup>2</sup> S)ns		5-12s <sup>2</sup> S	5-10p <sup>2</sup> P°		5-12d <sup>2</sup> D	4-6f <sup>2</sup> F°
4d <sup>9</sup> 5s( <sup>2</sup> D)ns'	{	6, 7s' <sup>4</sup> D	5p' <sup>4</sup> P°	5p' <sup>4</sup> D°	5p' <sup>4</sup> F°	
6s' <sup>2</sup> D		5p' <sup>2</sup> P°	5p' <sup>2</sup> D°	5p' <sup>2</sup> F°		
4d <sup>9</sup> 5s( <sup>1</sup> D)ns''		6s'' <sup>2</sup> D	5p'' <sup>2</sup> P°	5p'' <sup>2</sup> D°	5p'' <sup>2</sup> F°	

\*For predicted terms of the Ag I isoelectronic sequence, see Vol. III, Introduction.

## Ag II

(Pd I sequence; 46 electrons)

Z=47

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> <sup>1</sup>S<sub>0</sub>

4d<sup>9</sup> <sup>1</sup>S<sub>2</sub>, 173300 K

I. P. 21.48 volts

More than 380 lines of Ag II have been classified in the range from 728.73 Å to 9052.69 Å. The terms are from the papers by Gilbert and by Rasmussen, who have extended the early work of Shenstone, Blair, Duffendack and Thomson, Beals, and others.

Catalán and Rico have adjusted the value of the limit from a study of the spectra Sr II to Cd II. Their value is quoted here. Gilbert has derived the ionization limit 173274.0 K from the 4d<sup>n</sup> ns <sup>2</sup>D series (n=6 to 8) by means of a Ritz formula. To quote, "of the 4d<sup>n</sup> ns <sup>2</sup>D series the <sup>2</sup>D<sub>1</sub> and <sup>2</sup>D<sub>3</sub> series converge to the 4d<sup>n</sup> <sup>2</sup>D<sub>3/2</sub> state of Ag III, and the <sup>2</sup>D<sub>1</sub> and <sup>2</sup>D<sub>3</sub> series converge to 4d<sup>n</sup> <sup>2</sup>D<sub>3/2</sub>". From these two limits he determines the interval of the ground term <sup>2</sup>D of Ag III as 4607 cm<sup>-1</sup>. In the table the J-values of the limits are indicated.

The pair-coupling notation, introduced first in these Volumes for the inert gases to denote the departure from LS-coupling, applies also to spectra of the Pd I sequence. In the text of Pd I a small auxiliary table giving this notation, may be found.

The three-place g-values in the table are from unpublished Zeeman patterns of 37 lines measured by W. F. Meggers, C. C. Kiees, and C. J. Humphreys at the National Bureau of Standards. These Ag II lines appeared as impurities in various spectra, observed mostly at the Massachusetts Institute of Technology. The weakest determinations are indicated by a colon. Two-place entries are from the paper by Shenstone and Blair, who derived g-values from unresolved Zeeman patterns of Ag II from their formula for blended patterns.

Observed intersystem combinations connect the systems of terms of different multiplicities. The ground term has been connected with the rest by Shenstone from White's observations of three lines, subsequently confirmed by Menzies, as follows:

White		Menzies		Desig.
A (vac)	K	A (vac)	K	
1107.05	90330.2	1107.01	90333.4	4d <sup>9</sup> <sup>1</sup> S <sub>2</sub> -5p <sup>2</sup> D <sub>1</sub>
1112.46	89890.9	1112.35	89892.8	4d <sup>9</sup> <sup>1</sup> S <sub>2</sub> -5p <sup>1</sup> F <sub>1</sub>
1196.87	83621.1	1196.76	83623.8	4d <sup>9</sup> <sup>1</sup> S <sub>2</sub> -5p <sup>1</sup> F <sub>1</sub>

## Ag II—Continued

## REFERENCES

- A. G. Shenstone, *Phys. Rev.* **31**, 317 (1933). (I P) (T) (C L)  
 A. C. Mearns, *Proc. Roy. Soc. (London)* [A] **123**, 134 (1929). (T) (C L)  
 A. G. Shenstone and H. A. Blair, *Phil. Mag.* **3**, 770 (1929). (S E)  
 H. A. Blair, *Phys. Rev.* **36**, 173 (1930). (T) (C L)  
 O. S. Duffendack and K. Thomson, *Phys. Rev.* **69**, 100 (1922). (T) (C L)  
 W. P. Gilbert, *Phys. Rev.* **47**, 847 (1935). (I P) (T) (C L)  
 E. Rasmussen, *Kgl. Danske Videnskab., Math.-fys. Medd.* (Copenhagen) **18**, No. 5, 10 (1940). (T) (C L) (I S)  
 M. A. Catalán y F. R. Risco, *An. Real Soc. Esp. Física y Química* (Madrid) [A] **48**, 328 (1952) and letter (December 1955). (I P)

## Ag II

## Ag II

Config.	Desig.	J	Level	Interval	Obs. $g$	Config.	Desig.	J	Level	Interval	Obs. $g$
4d <sup>10</sup>	4d <sup>10</sup> 1S	0	0.0			4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p	6p <sup>1</sup> P <sup>o</sup>	2	130781.01	-2116.63	
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5e	5e <sup>1</sup> D	3	39163.9	-1577.1	1.318	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p		1	132837.64	-3328.18	
		2	40741.0	-2997.7	1.119			0	136166.33		
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5e		1	43738.7		0.488	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d	5d <sup>1</sup> P	1	130756.0		1.51
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5e	5e <sup>1</sup> D	2	46045.7		1.028	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d	5d <sup>1</sup> G	4	131510.5		1.065
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5p	5p <sup>3</sup> P <sup>o</sup>	2	80173.90	-3449.16	1.464	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d	5d <sup>1</sup> D	2	131783.9		1.147
		1	85681.36	-2514.66	1.410	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d	5d <sup>1</sup> F	3	132191.8		1.095
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5p		0	86136.03			4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p	6p <sup>3</sup> F <sup>o</sup>	4	132485.50	411.97	
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5p	5p <sup>3</sup> F <sup>o</sup>	4	83665.41	1497.85	1.249		3	138013.55	-4501.38		
		3	82167.56	-4716.39	1.059	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )6p		2	136514.91		
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5p		2	86833.95		0.889		1 <sup>o</sup>	4	132684.46		
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5p	5p <sup>1</sup> D <sup>o</sup>	3	86456.54	1259.93	1.240	4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p	6p <sup>1</sup> D <sup>o</sup>	3	135017.94	101.65	
		2	85196.61	-5134.13	0.851		2	132916.29	-4305.14		
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5p		1	90330.74		0.52	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )6p		1	137221.43		
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5p	5p <sup>1</sup> F <sup>o</sup>	3	89130.64		1.076		2 <sup>o</sup>	2	133137.81		
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5p	5p <sup>1</sup> P <sup>o</sup>	1	89891.40		1.008		3 <sup>o</sup>	2	135589.4		
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5p	5p <sup>1</sup> D <sup>o</sup>	2	90883.70		1.099		4 <sup>o</sup>	3	134339.04		
4d <sup>9</sup> 5e <sup>2</sup>	5e <sup>2</sup> <sup>3</sup> F	4	93928.66	-4029.15		4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d	5d <sup>1</sup> S	0	134449.4		
		3	97957.81	-1644.09			5 <sup>o</sup>	1	134795.04		
		2	99601.90				6 <sup>o</sup>	2	135411.22		
4d <sup>9</sup> 5e <sup>2</sup>	5e <sup>2</sup> <sup>3</sup> F	2	105258.18	-3676.16		4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )6p	6p <sup>1</sup> P <sup>o</sup>	1	136828.51		
		1	108934.34			4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )6p	6p <sup>1</sup> F <sup>o</sup>	3	137172.63		
		0				4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )6p	6p <sup>1</sup> D <sup>o</sup>	2	137464.20		
4d <sup>9</sup> 5e <sup>2</sup>	5e <sup>2</sup> 1S	0	109123.59				7 <sup>o</sup>	3	138633.3		
4d <sup>9</sup> 5e <sup>2</sup>	5e <sup>2</sup> 1D	2	110769.41			4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s	7s <sup>3</sup> D	3	144473.9	-145.1	
4d <sup>9</sup> 5e <sup>2</sup>	5e <sup>2</sup> 1G	4	113598.12				2	144619.0	-4459.4		
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6e	6e <sup>3</sup> D	3	120529.45	-377.68	1.293:	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )7s	7s <sup>1</sup> D	2	149179.1		
		2	120907.13	-4215.25	1.195:	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )7s	7s <sup>3</sup> D	3	155085.3	-71.5	
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6e		1	125122.38		0.53	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )8s	8s <sup>3</sup> D	2	155156.8	-4534.0	
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )6e	6e <sup>1</sup> D	2	125400.89		1.11	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )8s		1	159690.8		
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5d	5d <sup>3</sup> S	1	125568.9		1.84	4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )8s	8s <sup>1</sup> D	2	159738.8		
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5d	5d <sup>3</sup> G	5	126660.5	-12.6	1.03						
		4	126673.1	-4573.2	0.757						
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d		3	131246.3								
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5d	5d <sup>3</sup> P	2	126760.2	-3.5	1.334						
		1	126763.7	-1765.1	0.90						
"		0	128528.8								
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5d	5d <sup>3</sup> D	3	127205.0	-312.0	1.288:						
		2	127517.0	-3983.5	1.006:						
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d		1	131500.5		0.74						
4d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )5d	5d <sup>3</sup> F	4	127601.8	117.3	1.218:	Ag III( <sup>2</sup> D <sub>3/2</sub> )	Limit	-----	173300		
		3	127484.5	-4664.4	0.958	Ag III( <sup>2</sup> D <sub>1/2</sub> )	Limit	-----	177862		
4d <sup>9</sup> ( <sup>2</sup> D <sub>1/2</sub> )5d		2	132148.9		0.71						

March 1952.

## Ag II Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> +	Observed Terms				
4s <sup>2</sup>	4s <sup>2</sup> <sup>1</sup> S				
4s <sup>2</sup> 5s	{ 5s <sup>2</sup> <sup>1</sup> S	5s <sup>2</sup> <sup>3</sup> P	5s <sup>2</sup> <sup>1</sup> D	5s <sup>2</sup> <sup>3</sup> F	5s <sup>2</sup> <sup>1</sup> G
	ns (n ≥ 5)			np (n ≥ 5)	
4s <sup>2</sup> ( <sup>1</sup> D) <sub>ns</sub>	{	5-8s <sup>3</sup> D	5-8s <sup>1</sup> D	5, 6p <sup>3</sup> P°	5, 6p <sup>1</sup> D°
				5, 6p <sup>3</sup> F°	5, 6p <sup>1</sup> F°
	nd (n ≥ 5)				
4s <sup>2</sup> ( <sup>1</sup> D) <sub>ns</sub>	{ 5d <sup>1</sup> S	5d <sup>3</sup> P	5d <sup>1</sup> D	5d <sup>3</sup> F	5d <sup>1</sup> G
	{ 5d <sup>1</sup> S	5d <sup>3</sup> P	5d <sup>1</sup> D	5d <sup>3</sup> F	5d <sup>1</sup> G

\*For predicted terms in the spectra of the Pd I isoelectronic sequence, see Vol. III, Introduction.

## Ag III

(Rh I sequence; 45 electrons)

Z=47

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>9</sup> <sup>2</sup>D<sub>3/2</sub>

4d<sup>9</sup> <sup>2</sup>D<sub>3/2</sub>, 289900 K

I. P. 34.82 volts

The analysis is by Gilbert, who has observed the spectrum from 500 Å to 3000 Å, thereby revising and greatly extending the earlier work by B. V. R. Rao. Gilbert has published 257 classified lines between 709.80 Å and 3013.81 Å. The limit is from Catalán and Rico, who have interpolated it by comparison of third spectra from Y to In.

Observed intersystem combinations connect the doublet and quartet systems of terms.

## REFERENCES

- B. V. R. Rao, Proc. Indian Acad. Sci. **1A**, 23 (1934). (I P) (T) (C L)  
 W. P. Gilbert, Phys. Rev. **48**, 328 (1935). (I P) (T) (C L)  
 M. A. Catalán y F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] **53**, 85 (1957). (I P)



## Ag III

## Ag III

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval																																																																																																																																																																																				
4d	4d <sup>2</sup> D	3/2	0	-4007	4d(2F)5p	5p <sup>2</sup> G°	4/2	185850	-2479																																																																																																																																																																																				
		1/2	4007				3/2	187720		4d(2F)5s	5s <sup>2</sup> F	4/2	68250	-2514	4d(2F)5p	5p <sup>2</sup> F°	3/2	186738	-2072	3/2	65764	2/2	188804	3/2	68145			1/2	60851			4d(2F)5s	5s <sup>2</sup> F	3/2	71801	-2243	4d(2P)5p	5p' <sup>2</sup> P°	0/2	189143	794	2/2	73934	1/2	189937	4d(2P)5s	5s' <sup>2</sup> P	2/2	76406	-1007	4d(2D)5p	5p'' <sup>2</sup> F°	2/2	191875	1760	1/2	77413	3/2	192635	0/2	793267			4d(2D)5s	5s'' <sup>2</sup> D	1/2	80131	2100	4d(2D)5p	5p'' <sup>2</sup> P°	0/2	193457	1488	2/2	82231	1/2	194955	4d(2P)5s	5s' <sup>2</sup> P	1/2	85182	-2295	4d(2D)5p	5p'' <sup>2</sup> D°	1/2	196356	406	0/2	87477	2/2	197688	4d(2G)5s	5s''' <sup>2</sup> G	4/2	85599	-128	4d(2P)5p	5p' <sup>2</sup> D°	0/2	198808	123	3/2	85727	1/2	19981	4d(2F)5p	5p <sup>2</sup> D°	3/2	115412	-3731	4d(2G)5p	5p''' <sup>2</sup> H°	4/2	198809	3133	2/2	119143	5/2	19948	1/2	129014			0/2	123408			4d(2F)5p	5p <sup>2</sup> G°	5/2	120359	2428	4d(2P)5p	5p' <sup>2</sup> D°	2/2	198849	-473	4/2	117891	1/2	199322	3/2	121068			2/2	122522			4d(2F)5p	5p <sup>2</sup> F°	4/2	122300	-1331	4d(2P)5p	5p' <sup>2</sup> P°	1/2	140078	-3703	3/2	123451	0/2	145781	2/2	126808			1/2	125987			4d(2F)5p	5p <sup>2</sup> D°	2/2	125096	-2775	4d(2G)5p	5p''' <sup>2</sup> F°	3/2	140881	-1284	1/2	127870	2/2	142165				
4d(2F)5s	5s <sup>2</sup> F	4/2	68250	-2514	4d(2F)5p	5p <sup>2</sup> F°	3/2	186738	-2072																																																																																																																																																																																				
		3/2	65764				2/2	188804																																																																																																																																																																																					
		3/2	68145																																																																																																																																																																																										
		1/2	60851																																																																																																																																																																																										
4d(2F)5s	5s <sup>2</sup> F	3/2	71801	-2243	4d(2P)5p	5p' <sup>2</sup> P°	0/2	189143	794																																																																																																																																																																																				
		2/2	73934				1/2	189937																																																																																																																																																																																					
4d(2P)5s	5s' <sup>2</sup> P	2/2	76406	-1007	4d(2D)5p	5p'' <sup>2</sup> F°	2/2	191875	1760																																																																																																																																																																																				
		1/2	77413				3/2	192635																																																																																																																																																																																					
		0/2	793267																																																																																																																																																																																										
4d(2D)5s	5s'' <sup>2</sup> D	1/2	80131	2100	4d(2D)5p	5p'' <sup>2</sup> P°	0/2	193457	1488																																																																																																																																																																																				
		2/2	82231				1/2	194955																																																																																																																																																																																					
4d(2P)5s	5s' <sup>2</sup> P	1/2	85182	-2295	4d(2D)5p	5p'' <sup>2</sup> D°	1/2	196356	406																																																																																																																																																																																				
		0/2	87477				2/2	197688																																																																																																																																																																																					
4d(2G)5s	5s''' <sup>2</sup> G	4/2	85599	-128	4d(2P)5p	5p' <sup>2</sup> D°	0/2	198808	123																																																																																																																																																																																				
		3/2	85727				1/2	19981																																																																																																																																																																																					
4d(2F)5p	5p <sup>2</sup> D°	3/2	115412	-3731	4d(2G)5p	5p''' <sup>2</sup> H°	4/2	198809	3133																																																																																																																																																																																				
		2/2	119143				5/2	19948																																																																																																																																																																																					
		1/2	129014																																																																																																																																																																																										
		0/2	123408																																																																																																																																																																																										
4d(2F)5p	5p <sup>2</sup> G°	5/2	120359	2428	4d(2P)5p	5p' <sup>2</sup> D°	2/2	198849	-473																																																																																																																																																																																				
		4/2	117891				1/2	199322																																																																																																																																																																																					
		3/2	121068																																																																																																																																																																																										
		2/2	122522																																																																																																																																																																																										
4d(2F)5p	5p <sup>2</sup> F°	4/2	122300	-1331	4d(2P)5p	5p' <sup>2</sup> P°	1/2	140078	-3703																																																																																																																																																																																				
		3/2	123451				0/2	145781																																																																																																																																																																																					
		2/2	126808																																																																																																																																																																																										
		1/2	125987																																																																																																																																																																																										
4d(2F)5p	5p <sup>2</sup> D°	2/2	125096	-2775	4d(2G)5p	5p''' <sup>2</sup> F°	3/2	140881	-1284																																																																																																																																																																																				
		1/2	127870				2/2	142165																																																																																																																																																																																					
					Ag IV(2F <sub>4</sub> )	Limit		220900																																																																																																																																																																																					

February 1957.

## Ag III OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> +	Observed Terms	
4d	4d <sup>2</sup> D	
	$n_s$ ( $n \geq 5$ )	$n_p$ ( $n \geq 5$ )
4d(2F) $n_z$	{ 5s <sup>2</sup> F 5s <sup>2</sup> F	5p <sup>2</sup> D° 5p <sup>2</sup> F° 5p <sup>2</sup> G° 5p <sup>2</sup> D° 5p <sup>2</sup> F° 5p <sup>2</sup> G°
4d(2P) $n_z'$	{ 5s' <sup>2</sup> P 5s' <sup>2</sup> P	5p' <sup>2</sup> P° 5p' <sup>2</sup> D° 5p' <sup>2</sup> P° 5p' <sup>2</sup> D°
4d(2D) $n_z''$	5s'' <sup>2</sup> D	5p'' <sup>2</sup> P° 5p'' <sup>2</sup> D° 5p'' <sup>2</sup> F°
4d(2G) $n_z'''$	5s''' <sup>2</sup> G	5p''' <sup>2</sup> F° 5p''' <sup>2</sup> H°

\*For predicted terms in the spectra of the Rh I isoelectronic sequence, see Vol. III, Introduction.

## Ag XIX

(Cu I sequence; 29 electrons)

Z=47

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 S_{0,1}$  $4s^2 S_{0,1}$ 

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed the doublets due to the transitions  $4s-5p$ ,  $4p-5d$ ,  $4d-5f$ , and  $4p-5s$ , from Pd XVIII to In XXI. In figure 3 of his paper on the spectra of highly-ionized atoms, the lines are indicated on the photograph of vacuum spark spectra of Ag observed from 40 Å to 80 Å.

By analogy with Cu I the writer has assumed the ground state indicated above.

## REFERENCE

B. Edlén, *Physica* 12, No. 9, 549 (1947).

March 1948.

## Ag XXI

(Co I sequence; 27 electrons)

Z=47

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^9 D_{3,1}$  $3d^9 D_{3,1}$ 

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed three lines due to the transition  $3p^4 3d^9 D-3p^4 3d^{10} P^o$ , in the region between 40 Å and 80 Å. In figure 4 of his paper on the spectra of highly-ionized atoms, the observed wave numbers are plotted against atomic number for this combination in the Co I-like spectra Pd XX to Sn XXIV. For Ag XXI the wave numbers are between 1500000 and 1900000 K.

## REFERENCE

B. Edlén, *Physica* 12, No. 9, 548 (1947).

March 1948.

## CADMIUM

Cd I

48 electrons

 $Z=48$ Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 1S_0$  $5s^2 1S_0$ , 72538.8 K

I. P. 8.991 volts

More than 200 lines in the spectrum have been classified. The observations extend from 682.85 Å to nearly 39087 Å, but the existing lists of wavelengths are not homogeneous. The accuracy of the present data could be greatly improved with modern observations, particularly in the infrared.

The early analysis was by Paschen and Götze and by Fowler, who found the leading series. The limit, and most of the terms in the table are from Fowler's Report. Ruark added the terms: 12 to 15s  $1S$ ; 15, 16s  $3S$ ; 9 to 11, 13 to 15d  $1D$ ; 17 to 19d  $3D$ , and 5p  $3P$ . Shenstone and Pittenger revised the values of the terms 7 to 12d  $3D$ . Beutler observed in absorption 26 lines between 682 Å and 1022 Å, which he classifies as transitions from the ground term to terms above the ionization limit having the configurations  $4d^9 5s^2 (3D)np$  and  $4d^9 5s^2 (3D)nf$ . The  $J$ -values of the limit term in Cd II,  $3D$ , are from this paper. For these levels the values of the observed wave numbers are entered in the table. For triplet terms only the component with  $J=1$  is indicated, since only the combinations with the ground term ( $J=0$ ) are to be expected.

Observed intersystem combinations connect the singlet and triplet systems of terms.

The entries in the table given to more than one decimal are from interferometer measurements of 18 lines, of which 14 are from the paper by Meggers and Burns, 3 from Kietzler, Boreman, and Burns, and 1 from Ignatieff. The level 5p  $3P_2$  is from Garton and Rajaratnam, who have observed wide diffuse features at 2212 Å and 2270 Å which they designate as combinations of this level with 5p  $3P_1^o$  and 5p  $3P_2^o$ , respectively.

Observations of forbidden transitions and of hyperfine structure are discussed in numerous papers. The Paschen-Back effect of the three lines at 4678 Å, 4799 Å, and 5085 Å, 5p  $3P^o$ —6s  $3S$ , are described in detail by Satô. Green and Gray give a similar report for six lines between 3403 Å and 3614 Å, 5p  $3P_2^o$ —5d  $1D$ .

Cd I occupies a unique position among spectra in that it furnishes the primary standard of wavelength. The line 6438.4696 Å, 5p  $1P_1^o$ —5d  $1D_2$ , is the fundamental wavelength on which all other standards are based. The specifications for the production of this primary standard are defined in the 1925 report of Commission 14 of the International Astronomical Union—the Commission on Standards of Wave-Length. The summary of the directly measured wavelengths "of the Cadmium Red Line in terms of the International Metre (1 Å. =  $1 \times 10^{-10}$  m)" is published in the 1950 report of this Commission.

## Cd I—Continued

## REFERENCES

- A. Ignatieff, *Ann. der Phys.* [4] **43**, 1117 (1914).  
 W. F. Meggers and K. Burns, *Sci. Papers Bur. Std.* **18**, 185, No. 441 (1922).  
 F. Paschen und R. Götze, *Seriengesetze der Linienspektren*, p. 110 (Julius Springer, Berlin, 1922). (T) (C L)  
 A. Fowler, *Report on Series in Line Spectra*, p. 142 (Fleetway Press, London, 1922). (I P) (T) (C L)  
*Trans. Intern. Astron. Union* **3**, 40 (1925).  
 A. E. Ruark, *J. Opt. Soc. Am.* **11**, 199 (1925). (T) (C L)  
 H. Beutler, *Zeit. Phys.* **87**, 19 (1933). (T) (C L)  
 J. B. Green and D. E. Gray, *Phys. Rev.* **45**, 273 (1933). (Z E)  
 C. W. Hetsler, R. W. Boreman, and K. Burns, *Phys. Rev.* **48**, 656 (1935).  
 S. Satô, *Sci. Reports Tohoku Univ. (Sendai)* [1] **25**, 686 (1936). (Z E)  
 A. G. Shenstone and J. T. Pittenger, *J. Opt. Soc. Am.* **39**, 220 (1949). (T)  
*Trans. Intern. Astron. Union* **7**, 146 (1950).  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab. ORNL-1469, Suppl. II, **39** (1955).  
 (Summary hfs)  
 W. R. S. Garton and A. Rajaratnam, *Proc. Phys. Soc. (London)* [A] **68**, 1107 (1955). (T) (C L)

Cd I

Cd I

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
4d <sup>10</sup> 5s <sup>2</sup>	5s <sup>2</sup> 1S	0	0.0		4d <sup>10</sup> 5s(2S)7p	7p 1P°	1	66494.8	
4d <sup>10</sup> 5s(2S)5p	5p 1P°	0	30114.017 30656.130 31888.996	542.113 1170.866	4d <sup>10</sup> 5s(2S)4f	4f 1F°	2, 3, 4	65581.7	
4d <sup>10</sup> 5s(2S)5p	5p 1P°	1	43698.474		4d <sup>10</sup> 5s(2S)8s	8s 1S	1	66680.920	
4d <sup>10</sup> 5s(2S)6s	6s 1S	1	51484.013		4d <sup>10</sup> 5s(2S)8s	8s 1S	0	66904.7	
4d <sup>10</sup> 5s(2S)6s	6s 1S	0	53310.16		4d <sup>10</sup> 5s(2S)8p	8p 1P°	0 1 2	67829.6 67848.1 67875.8	12.5 33.1
4d <sup>10</sup> 5s(2S)6p	6p 1P°	0	58390.9 58461.6 58635.7	70.7 174.1	4d <sup>10</sup> 5s(2S)7d	7d 1D	2	67837.1	
4d <sup>10</sup> 5s(2S)5d	5d 1D	2	59219.824		4d <sup>10</sup> 5s(2S)7d	7d 1D	1 2 3	67990.1 67992.9 67997.5	2.8 4.6
4d <sup>10</sup> 5s(2S)5d	5d 1D	1	59485.793 59497.895 59518.017	12.102 18.122	4d <sup>10</sup> 5s(2S)8p	8p 1P°	1	68065.4	
4d <sup>10</sup> 5s(2S)6p	6p 1P°	1	59905.6		4d <sup>10</sup> 5s(2S)5f	5f 1F°	2, 3, 4	68093.7	
4d <sup>10</sup> 5s(2S)7s	7s 1S	1	62563.461		4d <sup>10</sup> 5s(2S)9s	9s 1S	1	68682.2	
4d <sup>10</sup> 5s(2S)7s	7s 1S	0	63086.7		4d <sup>10</sup> 5s(2S)9s	9s 1S	0	68799.6	
4d <sup>10</sup> 5s(2S)7p	7p 1P°	0	64995.9 65021.3 65098.8	25.4 71.5	4d <sup>10</sup> 5s(2S)8d	8d 1D	2	69292.5	
4d <sup>10</sup> 5s(2S)6d	6d 1D	2	65134.871		4d <sup>10</sup> 5s(2S)9p	9p 1P°	0 1 2	69314.5 69331.4 69340.8	6.9 18.8
4d <sup>10</sup> 5s(2S)6d	6d 1D	1	65353.5 65359.3 65367.257	5.8 8.0	4d <sup>10</sup> 5s(2S)8d	8d 1D	1 2 3	69400.5 69402.8 69405.5	2.3 2.7

## Cd I—Continued

## Cd I—Continued

Config.	Design.	<i>J</i>	Level	Interval	Config.	Design.	<i>J</i>	Level	Interval
4d <sup>10</sup> 5s(°S)9p	9p 1P°	1	69435. 7		4d <sup>10</sup> 5s(°S)16d	16d 1D	1, 2, 3	71972. 1	
4d <sup>10</sup> 5s(°S)10e	10e 1S	1	69905. 9		4d <sup>10</sup> 5s(°S)17d	17d 1D	1, 2, 3	72046. 7	
4d <sup>10</sup> 5s(°S)10e	10e 1S	0	69873. 1		4d <sup>10</sup> 5s(°S)18d	18d 1D	1, 2, 3	72108. 2	
4d <sup>10</sup> 5s(°S)9d	9d 1D	2	70175. 9		4d <sup>10</sup> 5s(°S)19d	19d 1D	1, 2, 3	72155. 7	
4d <sup>10</sup> 5s(°S)10p	10p 1P°	0, 1, 2	70907. 3		Cd II(°S <sub>ns</sub> )	Limit	-----	72538. 8	
4d <sup>10</sup> 5s(°S)9d	9d 1D	1	70244. 2	1. 3	4d <sup>10</sup> 5p <sup>1</sup>	5p <sup>1</sup> 1P	0	73996. 2	749. 2
		2	70245. 5	1. 6			1	74745. 4	1115
		3	70247. 1				2	75860	
4d <sup>10</sup> 5s(°S)10p	10p 1P°	1	70962. 6		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )5p	5p'' 1P°	1	97800	
4d <sup>10</sup> 5s(°S)11a	11a 1S	1	70501. 2		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )5p	5p'' 1P°	1	103480	
4d <sup>10</sup> 5s(°S)11e	11e 1S	0	70543. 2		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )5p	5p'' 1D°	1	104351	
4d <sup>10</sup> 5s(°S)10d	10d 1D	2	70741. 9		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )6p	6p'' 1P°	1	127464	
4d <sup>10</sup> 5s(°S)10d	10d 1D	1	70788. 3	0. 7	4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )6p	6p'' 1P°	1	132998	
		2	70789. 0	1. 1	4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )6p	6p'' 1D°	1	133044	
		3	70790. 1		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )7p	7p'' 1P°	1	134196	
4d <sup>10</sup> 5s(°S)11p	11p 1P°	1	70800. 0		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )4f7	4f'' 1P°?	1	135097	
4d <sup>10</sup> 5s(°S)12e	12e 1S	1	70962. 0		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )8p	8p'' 1P°	1	137027	
4d <sup>10</sup> 5s(°S)12e	12e 1S	0	70992. 1		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )9p	9p'' 1P°	1	138524	
4d <sup>10</sup> 5s(°S)11d	11d 1D	2	71119. 5		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )6f7	6f'' 1P°?	1	138679	
4d <sup>10</sup> 5s(°S)12p	12p 1P°	1	71167. 9		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )10p	10p'' 1P°	1	139417	
4d <sup>10</sup> 5s(°S)11d	11d 1D	1	71159. 6	1. 0	4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )7p	7p'' 1P°	1	139652	
		2	71160. 6		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )11p	11p'' 1P°	1	139987	
		3	71160. 6		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )12p	12p'' 1P°	1	140410	
4d <sup>10</sup> 5s(°S)13e	13e 1S	1	71281. 8		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )13p?	13p'' 1P°?	1	140655	
4d <sup>10</sup> 5s(°S)13e	13e 1S	0	71299. 0		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )4f7	4f'' 1P°	1	140655	
4d <sup>10</sup> 5s(°S)12d	12d 1D	1	71424. 1	0. 9	4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )8p	8p'' 1P°	1	142613	
		2	71425. 0		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )5f7	5f'' 1P°?	1	143903	
		3	71425. 0		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )9p	9p'' 1P°	1	144154	
4d <sup>10</sup> 5s(°S)14e	14e 1S	1	71515. 6		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )6f	6f'' 1P°?	1	144531	
4d <sup>10</sup> 5s(°S)14e	14e 1S	0	71528. 0		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )10p	10p'' 1P°	1	145034	
4d <sup>10</sup> 5s(°S)13d	13d 1D	2	71596. 3		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )7f7	7f'' 1P°?	1	145314	
4d <sup>10</sup> 5s(°S)13d	13d 1D	1, 2, 3	71618. 5		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )11p	11p'' 1P°	1	145603	
4d <sup>10</sup> 5s(°S)15e	15e 1S	1	71689. 6		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )12p	12p'' 1P°	1	145993	
4d <sup>10</sup> 5s(°S)15e	15e 1S	0	71692. 4		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )13p	13p'' 1P°	1	146281	
4d <sup>10</sup> 5s(°S)14d	14d 1D	2	71748. 9		4d <sup>10</sup> 5s <sup>1</sup> (°D <sub>11d</sub> )14p	14p'' 1P°	1	146446	
4d <sup>10</sup> 5s(°S)14d	14d 1D	1, 2, 3	71767. 2						
4d <sup>10</sup> 5s(°S)16e	16e 1S	1	71818. 6						
4d <sup>10</sup> 5s(°S)15d	15d 1D	2	71869. 2						
4d <sup>10</sup> 5s(°S)15d	15d 1D	1, 2, 3	71880. 7						

August 1955.

## Cd I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> +	Observed Terms			
4d <sup>10</sup> 5s <sup>2</sup>	5s <sup>2</sup> <sup>1</sup> S			
4d <sup>10</sup> 5p <sup>2</sup>	5p <sup>2</sup> <sup>3</sup> P			
	ns (n ≥ 6)	np (n ≥ 5)	nd (n ≥ 5)	nℓ (n ≥ 4)
4d <sup>10</sup> 5s( <sup>3</sup> S)ns	{ 6-16s <sup>3</sup> S 6-15s <sup>1</sup> S	5-10p <sup>3</sup> P° 5-12p <sup>1</sup> P°	5-10d <sup>3</sup> D 5-11, 13-15d <sup>1</sup> D	4, 6f <sup>3</sup> F°
4d <sup>10</sup> 5s( <sup>1</sup> D)ns''		5-14p'' <sup>3</sup> P°    5-6p'' <sup>3</sup> D° 5-13p'' <sup>1</sup> P°		4, 6f'' <sup>3</sup> F°? 4-7f'' <sup>1</sup> F°?

\*For predicted terms in the spectra of the Cd I isoelectronic sequence, see Vol. III, Introduction.

## Cd II

(Ag I sequence; 47 electrons)

Z=48

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s <sup>2</sup>S<sub>1/2</sub>

5s <sup>2</sup>S<sub>1/2</sub> 136374.74 K

I. P. 16.904 volts

The analysis is from the paper by Shenstone and Pittenger, who have revised and extended the earlier work by Takahashi and by von Salis. In the range from 761 Å to 8066 Å there are 256 classified lines. Shenstone's notation is given in the first column of the table. He has redetermined the limit from the <sup>2</sup>S and <sup>2</sup>D series; and stresses the irregularity of both the <sup>2</sup>P° and <sup>2</sup>F° series, as was pointed out by von Salis.

Observed intersystem combinations connect the doublet and quartet systems of terms. The combination 4f <sup>2</sup>F° - 5g <sup>2</sup>G has recently been observed by C. H. Corliss and R. J. Murphy at 9670.84 Å and 9658.75 Å. These observations furnish the value of 5g <sup>2</sup>G in the table, and fit well into the ng <sup>2</sup>G series (n=6 to 11) reported earlier.

## REFERENCES

- G. v. Salis, Ann. der Phys. [4] 76, 153 (1924). (I P) (T) (C L)  
 Y. Takahashi, Ann. der Phys. [5] 3, 27 (1929). (I P) (T) (C L)  
 A. G. Shenstone and J. T. Pittenger, J. Opt. Soc. Am. 30, 220 (1949). (I P) (T) (C L)  
 J. E. Mack, Rev. Mod. Phys. 22, 64 (1950). (Summary hfs)  
 C. H. Corliss and R. J. Murphy, unpublished material (March 1953). (T) (C L)

Cd II

Cd II

Author	Config.	Desig.	J	Level	Interval	Author	Config.	Desig.	J	Level	Interval
5e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)5e	5e <sup>2</sup> S	0 1/2	0. 00		10e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)10e	10e <sup>2</sup> S	0 1/2	127152. 89	
5p <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)5p	5p <sup>2</sup> F°	0 1/2	44136. 08	2482. 47	7f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)7f	7f <sup>2</sup> F°	3 1/2	127242. 09	-33. 95
5p <sup>2</sup> F <sub>1/2</sub>			1 1/2	46818. 55		7f <sup>2</sup> F <sub>3/2</sub>			2 1/2	127283. 04	
a <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> 5e	5e <sup>2</sup> D	2 1/2	69258. 91	-5634. 75	7g <sup>2</sup> G	4d <sup>10</sup> ( <sup>1</sup> S)7g	7g <sup>2</sup> G	3 1/2	127396. 62	
a <sup>2</sup> D <sub>1/2</sub>			1 1/2	74893. 66		9d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)9d	9d <sup>2</sup> D	1 1/2		
6e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)6e	6e <sup>2</sup> S	0 1/2	82990. 66		9d <sup>2</sup> D <sub>1/2</sub>		2 1/2	127712. 28	14. 85	
5d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)5d	5d <sup>2</sup> D	1 1/2	89689. 25	154. 53	10p <sup>2</sup> P <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)10p	10p <sup>2</sup> P°	1 1/2	127996. 73	-79. 6
5d <sup>2</sup> D <sub>1/2</sub>			2 1/2	89843. 78		10p <sup>2</sup> P <sub>3/2</sub>			0 1/2	128076. 3	
6p <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)6p	6p <sup>2</sup> F°	0 1/2	94710. 41	673. 22	y <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> 5e( <sup>1</sup> D)5p	5p'' <sup>2</sup> F°	3 1/2	129189. 5	-2528. 7
6p <sup>2</sup> F <sub>1/2</sub>			1 1/2	95383. 63		y <sup>2</sup> F <sub>1/2</sub>			2 1/2	131718. 3	
s <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> 5e( <sup>1</sup> D)5p	5p' <sup>2</sup> F°	2 1/2	108189. 30	-3251. 66	11e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)11e	11e <sup>2</sup> S	0 1/2	129343. 05	
s <sup>2</sup> F <sub>1/2</sub>			1 1/2	109440. 86		y <sup>2</sup> P <sub>1/2</sub>	4d <sup>10</sup> 5e( <sup>1</sup> D)5p	5p'' <sup>2</sup> P°	1 1/2	129344. 86	
s <sup>2</sup> F <sub>3/2</sub>			0 1/2	112196. 50					0 1/2		
7e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)7e	7e <sup>2</sup> S	0 1/2	107300. 88		8f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)8f	8f <sup>2</sup> F°	2 1/2	129419. 38	267. 22
4f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)4f	4f <sup>2</sup> F°	2 1/2	108419. 47	12. 96	8f <sup>2</sup> F <sub>1/2</sub>		3 1/2	129886. 60		
4f <sup>2</sup> F <sub>1/2</sub>			3 1/2	108432. 43		8g <sup>2</sup> G	4d <sup>10</sup> ( <sup>1</sup> S)8g	8g <sup>2</sup> G	3 1/2	129502. 43	
s <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> 5e( <sup>1</sup> D)5p	5p' <sup>2</sup> F°	4 1/2	109736. 82	-431. 29	10d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)10d	10d <sup>2</sup> D	1 1/2		129708. 50
s <sup>2</sup> F <sub>1/2</sub>			3 1/2	110168. 11		10d <sup>2</sup> D <sub>1/2</sub>			2 1/2	129718. 66	
s <sup>2</sup> F <sub>3/2</sub>			1 1/2	112785. 19		11p <sup>2</sup> P°	4d <sup>10</sup> ( <sup>1</sup> S)11p	11p <sup>2</sup> P°	0 1/2	129956. 3	260. 3
6d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)6d	6d <sup>2</sup> D	1 1/2	110174. 10	73. 50	11p <sup>2</sup> P <sub>1/2</sub>		1 1/2	130216. 6		
6d <sup>2</sup> D <sub>1/2</sub>			2 1/2	110247. 60		12e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)12e	12e <sup>2</sup> S	0 1/2	130835. 86	
7p <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)7p	7p <sup>2</sup> F°	0 1/2	112361. 06	129. 33	9f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)9f	9f <sup>2</sup> F°	2 1/2	130878. 61	47. 21
7p <sup>2</sup> F <sub>1/2</sub>			1 1/2	112490. 39		9f <sup>2</sup> F <sub>1/2</sub>			3 1/2	130925. 82	
s <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> 5e( <sup>1</sup> D)5p	5p' <sup>2</sup> D°	3 1/2	114793. 91?	-158. 39	9g <sup>2</sup> G	4d <sup>10</sup> ( <sup>1</sup> S)9g	9g <sup>2</sup> G	3 1/2	130946. 09	
s <sup>2</sup> D <sub>1/2</sub>			2 1/2	114962. 30		11d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)11d	11d <sup>2</sup> D	1 1/2		
s <sup>2</sup> D <sub>3/2</sub>			1 1/2	116285. 71		11d <sup>2</sup> D <sub>1/2</sub>			2 1/2	131100. 14	
s <sup>2</sup> D <sub>1/2</sub>			0 1/2	117988. 79	-1763. 08	12p <sup>2</sup> P <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)12p	12p <sup>2</sup> P°	0 1/2	131261. 1	113. 6
s <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> 5e( <sup>1</sup> D)5p	5p' <sup>2</sup> F°	2 1/2	116963. 70		12p <sup>2</sup> P <sub>1/2</sub>			1 1/2	131374. 7	
s <sup>2</sup> F <sub>1/2</sub>			3 1/2	119373. 19	2419. 49	13e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)13e	13e <sup>2</sup> S	0 1/2	131899. 07	-8. 15
8e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)8e	8e <sup>2</sup> S	0 1/2	118040. 65	10f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)10f	10f <sup>2</sup> F°	3 1/2	131953. 72?		
5f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)5f	5f <sup>2</sup> F°	3 1/2	118444. 36	-101. 41	10f <sup>2</sup> F <sub>1/2</sub>		2 1/2	131961. 87?		
5f <sup>2</sup> F <sub>1/2</sub>			2 1/2	118545. 77		10g <sup>2</sup> G	4d <sup>10</sup> ( <sup>1</sup> S)10g	10g <sup>2</sup> G	3 1/2	131978. 47	
s <sup>2</sup> P <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)5g	5g <sup>2</sup> G	3 1/2	118769. 95	-237. 88	12d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)12d	12d <sup>2</sup> D	1 1/2		132086. 77
s <sup>2</sup> P <sub>3/2</sub>			4 1/2			119066. 11	12d <sup>2</sup> D <sub>1/2</sub>			2 1/2	132092. 18
7d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)7d	7d <sup>2</sup> D	1 1/2	119522. 72	39. 27	13p <sup>2</sup> P <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)13p	13p <sup>2</sup> P°	0 1/2	132280. 4?	
7d <sup>2</sup> D <sub>1/2</sub>			2 1/2	119561. 99		11f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)11f	11f <sup>2</sup> F°	2 1/2	132707. 91?	
s <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> 5e( <sup>1</sup> D)5p	5p' <sup>2</sup> D°	1 1/2	120135. 22	1146. 42	11g <sup>2</sup> G	4d <sup>10</sup> ( <sup>1</sup> S)11g	11g <sup>2</sup> G	3 1/2		132742. 26
s <sup>2</sup> D <sub>1/2</sub>			2 1/2	121231. 64		13d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)13d	13d <sup>2</sup> D	1 1/2	132824. 25	
8p <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)8p	8p <sup>2</sup> F°	0 1/2	120618. 46	92. 72	13d <sup>2</sup> D <sub>1/2</sub>		2 1/2	132828. 29	4. 04	
8p <sup>2</sup> F <sub>1/2</sub>			1 1/2	120711. 18		4d <sup>10</sup> ( <sup>1</sup> S)14d	14d <sup>2</sup> D	1 1/2	133389. 76		
9e <sup>2</sup> S <sub>1/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)9e	9e <sup>2</sup> S	0 1/2	123751. 48				2 1/2			
6f <sup>2</sup> F <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)6f	6f <sup>2</sup> F°	3 1/2	123972. 10	-16. 29						
6f <sup>2</sup> F <sub>1/2</sub>			2 1/2	123988. 39							
6g <sup>2</sup> G	4d <sup>10</sup> ( <sup>1</sup> S)6g	6g <sup>2</sup> G	3 1/2	124151. 26							
			4 1/2								
8d <sup>2</sup> D <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)8d	8d <sup>2</sup> D	1 1/2	124613. 30	23. 20						
8d <sup>2</sup> D <sub>1/2</sub>			2 1/2	124636. 50							
9p <sup>2</sup> P <sub>3/2</sub>	4d <sup>10</sup> ( <sup>1</sup> S)9p	9p <sup>2</sup> P°	0 1/2	125223. 06	31. 43						
9p <sup>2</sup> P <sub>1/2</sub>			1 1/2	125254. 49							
						Cd III( <sup>1</sup> S <sub>0</sub> )	Limit			136374. 74	

March 1953.

## Cd II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>2</sup> +	Observed Terms				
	5s <sup>2</sup> 5D				
4d <sup>n</sup> 5s <sup>2</sup>	ns (n ≥ 5)	np (n ≥ 5)	nd (n ≥ 5)	n'f (n' ≥ 4)	ng (n ≥ 5)
4d <sup>10</sup> ( <sup>1</sup> S)ns	5-13s <sup>2</sup> S	5-13p <sup>2</sup> <sup>3</sup> P°	5-14d <sup>2</sup> D	4-11f <sup>2</sup> F°	5-11g <sup>2</sup> G
4d <sup>9</sup> 5s( <sup>3</sup> D)ns'	{	5p <sup>2</sup> <sup>1</sup> P°	5p <sup>2</sup> <sup>3</sup> D°	5p <sup>2</sup> <sup>1</sup> F°	
4d <sup>9</sup> 5s( <sup>1</sup> D)ns''		5p <sup>2</sup> <sup>3</sup> P°	5p <sup>2</sup> <sup>1</sup> F°		

\*For predicted terms in the spectra of the Ag I isoelectronic sequence, see Vol. III, Introduction.

## Cd III

(Pd I sequence; 46 electrons)

Z=48

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> <sup>1</sup>S<sub>0</sub>

4d<sup>10</sup> <sup>1</sup>S<sub>0</sub>, 302300 K

I. P. 37.47 volts

The analysis is by Shenstone and Pittenger, who have extended the early work by Gibbs and White and by McLennan, McLay, and Crawford, as well as that by Mazumder. The number of classified lines now totals 138; and the observations extend from 677 Å to 3035 Å. Inter-system combinations connecting the singlet and triplet systems of terms, have been observed.

Mack has noted that the sequence Pd I—Sn V shows a gradual transition from *LS*- to *jj*-coupling for levels from the d<sup>n</sup>s and d<sup>n</sup>p configurations. In Cd III Shenstone indicates the *J*-value of the limit for levels from the d<sup>n</sup>s and d<sup>n</sup>d configurations, as is done in *Jl*-coupling. In the text for Pd I the writer has included a table giving the *Jl*-coupling notation for the related pairs of levels in the form of notation suggested by Racah.

The limit from the 4d<sup>n</sup> ns series (n=5, 6) is 304690. From the s-electron series in Cd I and Cd II Shenstone predicted "a reasonable value for the Ritz correction in the Cd III series," and thus obtained a more reliable determination of the limit than that derived from a Rydberg series of only two members. The quoted value is from Catalán and Rico who have derived it by comparison of the third spectra from Y to In.

## REFERENCES

- J. E. Mack, *Phys. Rev.* **24**, 17 (1929).  
 A. G. Shenstone and J. T. Pittenger, *J. Opt. Soc. Am.* **36**, 221 (1949). (I F) (T) (C L)  
 M. A. Catalán y F. R. Rico, *An. Real Soc. Esp. Física y Química (Madrid)* [A] **53**, 85 (1957). (I F)



Cd III

Cd III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
4d <sup>10</sup>	4d <sup>10</sup> 1S	0	0.0		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1S	1	202966.1	
4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5e	5e 1D	3	80454.3	-1900.3	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )6e	6e 1D	3	204421.1	-457.0
		2	82354.6	-3364.9			2	204908.1	-5812.4
4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5e		1	86219.5		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )6e		1	210220.5	
4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5e	5e 1D	2	88871.8		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1G	5	205119.47	-177.3
4d <sup>9</sup> ( <sup>1</sup> D)5p	5p 1P°	3	133813.3	-4941.1	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d		4	205296.7	-5540.8
		1	138754.4	-3356.4	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d		3	210637.5	
		0	142110.8		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1P	2	206407.0	
4d <sup>9</sup> ( <sup>1</sup> D)5p	5p 1F°	4	139031.8	2816.9			1		
		3	139214.3	-6540.5	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1D	3	206087.7	-716.6
		2	142754.8				2	206804.3	-5248.5
4d <sup>9</sup> ( <sup>1</sup> D)5p	5p 1D°	3	142322.8	2463.4	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d		1	212049.8	
		2	140429.4	-7195.4	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1F	4	206967.8	357.5
		1	147624.8		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d		3	206610.3	-5314.5
4d <sup>9</sup> ( <sup>1</sup> D)5p	5p 1P°	1	148076.1		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d		2	211924.8	
4d <sup>9</sup> ( <sup>1</sup> D)5p	5p 1F°	3	148090.6		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1P	1	210147.1	
4d <sup>9</sup> ( <sup>1</sup> D)5p	5p 1D°	2	148461.3		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )6e	6e 1D	2	210588.0	
4d <sup>9</sup> 5e	5e 1F	4	179022.4	-5071.9	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1G	4	211518.5	
		3	184094.3	-1704.2	4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1D	2	212425.0	
		2	185798.5		4d <sup>9</sup> ( <sup>1</sup> D <sub>3/2</sub> )5d	5d 1F	3	212708.2	
4d <sup>9</sup> 5e	5e 1P	2	191762.3	-4261.4	4d <sup>9</sup> 5e	5e 1S	0	213107.7	
		1	196123.7	-184.5					
		0	196258.2						
4d <sup>9</sup> 5e	5e 1D	2	198431.0						
4d <sup>9</sup> 5e	5e 1G	4	200932.8		Cd IV ( <sup>1</sup> D <sub>3/2</sub> )	Limit		302300	

February 1957.

Cd III OBSERVED TERMS\*

Configuration 1e 2e 2p 3e 3p 3d <sup>10</sup> 4e 4p <sup>+</sup>	Observed Terms				
4d <sup>10</sup>	4d <sup>10</sup> 1S				
4d <sup>9</sup> 5e	5e 1S	5e 1P	5e 1D	5e 1F	5e 1G
	ns (n ≥ 5)		np (n ≥ 5)		
4d <sup>9</sup> ( <sup>1</sup> D)nz	{ 5, 6e 1D 5, 6e 1D		5p 1P°	5p 1D°	5p 1F°
	nd (n ≥ 5)				
4d <sup>9</sup> ( <sup>1</sup> D)nz	5d 1S	5d 1P	5d 1D	5d 1F	5d 1G

\*For predicted terms in the spectra of the Pd I isoelectronic sequence, see Vol. III, Introduction.

## Cd IV

(Rh 1 sequence; 45 electrons)

Z=48

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^9 \ ^1D_{3/2}$  $4d^9 \ ^1D_{3/2}$ 

K

I. P.

volts

The analysis is from Green, who has classified 185 lines between 489.49 Å and 1929.70 Å. Observed intersystem combinations connect the doublet and quartet systems of terms. No series are known in Cd IV.

## REFERENCE

M. Green, Phys. Rev. **60**, 117 (1941). (T) (C L)

## Cd IV

## Cd IV

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^9$	$4d^9 \ ^1D$	$2\frac{1}{2}$ $1\frac{1}{2}$	0 5812	-5812	$4d^9(^1P)5p$	$5p' \ ^1P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$ $2\frac{1}{2}$	189404 180411 190894	1007 283
$4d^9(^1F)5s$	$5s \ ^1F$	$4\frac{1}{2}$ $3\frac{1}{2}$ $2\frac{1}{2}$ $1\frac{1}{2}$	106704 111734 114758 116125	-3080 -3024 -1367	$4d^9(^1D)5p$	$5p'' \ ^1F^o$	$2\frac{1}{2}$ $3\frac{1}{2}$	181953 184367	2414
$4d^9(^1F)5s$	$5s \ ^1F$	$3\frac{1}{2}$ $2\frac{1}{2}$	118333 120557	-2224	$4d^9(^1D)5p$	$5p'' \ ^1P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	184218 185548	1330
$4d^9(^1G)5s$	$5s''' \ ^1G$	$4\frac{1}{2}$ $3\frac{1}{2}$	133860 134010	-150	$4d^9(^1D)5p$	$5p'' \ ^1D^o$	$2\frac{1}{2}$ $1\frac{1}{2}$	186255 1864857	-230
$4d^9(^1F)5p$	$5p \ ^1D^o$	$3\frac{1}{2}$ $2\frac{1}{2}$ $1\frac{1}{2}$ $0\frac{1}{2}$	178387 177258 180848 182804	-4865 -3616 -1998	$4d^9(^1P)5p$	$5p' \ ^1D^o$	$0\frac{1}{2}$ $1\frac{1}{2}$ $2\frac{1}{2}$ $3\frac{1}{2}$	186570 188290 188338 187508	1850 118 -830
$4d^9(^1F)5p$	$5p \ ^1G^o$	$5\frac{1}{2}$ $4\frac{1}{2}$ $3\frac{1}{2}$ $2\frac{1}{2}$	179178 175319 178575 180818	3854 -4056 -1437	$4d^9(^1G)5p$	$5p''' \ ^1H^o$	$4\frac{1}{2}$ $5\frac{1}{2}$	186590 201293	4733
$4d^9(^1F)5p$	$5p \ ^1F^o$	$4\frac{1}{2}$ $3\frac{1}{2}$ $2\frac{1}{2}$ $1\frac{1}{2}$	181678 182973 186307 189176	-1301 -3334 4131	$4d^9(^1P)5p$	$5p' \ ^1D^o$	$2\frac{1}{2}$ $1\frac{1}{2}$	186585 200743	-1218
$4d^9(^1F)5p$	$5p \ ^1F^o$	$4\frac{1}{2}$ $3\frac{1}{2}$ $2\frac{1}{2}$ $1\frac{1}{2}$	181678 182973 186307 189176	-1301 -3334 4131	$4d^9(^1P)5p$	$5p' \ ^1P^o$	$1\frac{1}{2}$ $0\frac{1}{2}$	201987 206408	-4435
$4d^9(^1F)5p$	$5p \ ^1D^o$	$2\frac{1}{2}$ $1\frac{1}{2}$	184301 187813	-2912	$4d^9(^1P)5p$	$5p' \ ^1S^o$	$0\frac{1}{2}$	202655	
$4d^9(^1F)5p$	$5p \ ^1G^o$	$4\frac{1}{2}$ $3\frac{1}{2}$	185368 188147	-2781	$4d^9(^1G)5p$	$5p''' \ ^1F^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	202838 204196	-1360
$4d^9(^1F)5p$	$5p \ ^1F^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	187168 188290	-1128	$4d^9(^1P)5p$	$5p' \ ^1S^o$	$1\frac{1}{2}$	204294	

February 1966.

## Cd IV Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>n</sup> 4s <sup>1</sup> 4p <sup>1</sup> +	Observed Terms		
4d <sup>1</sup>	4d <sup>1</sup> D		
	ns (n ≥ 5)	np (n ≥ 5)	
4d <sup>1</sup> (F) <sub>ns</sub>	{ 5s <sup>1</sup> F 5s <sup>1</sup> F	5p <sup>1</sup> D°	5p <sup>1</sup> F°
4d <sup>1</sup> (F) <sub>ns'</sub>		5p <sup>1</sup> D°	5p <sup>1</sup> F°
4d <sup>1</sup> (D) <sub>ns''</sub>		5p' <sup>1</sup> G°	5p' <sup>1</sup> G°
4d <sup>1</sup> (G) <sub>ns'''</sub>		5p' <sup>1</sup> G°	5p' <sup>1</sup> G°
		5p'' <sup>1</sup> P°	5p'' <sup>1</sup> P°
		5p'' <sup>1</sup> D°	5p'' <sup>1</sup> D°
		5p''' <sup>1</sup> F°	5p''' <sup>1</sup> F°
			5p''' <sup>1</sup> H°

\*For predicted terms in the spectra of the Rh I isoelectronic sequence, see Vol. III, Introduction.

## Cd XX

(Cu I sequence; 29 electrons)

Z=48

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>1</sup> S<sub>1/2</sub>

4s<sup>1</sup> S<sub>1/2</sub>

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed the doublets due to the transitions 4s-5p, 4p-5d, 4d-5f, and 4p-5s, from Pd XVIII to In XXI. In figure 3 of his paper on the spectra of highly-ionized atoms, the lines are indicated on the photograph of vacuum spark spectra of Cd observed from 40 Å to 80 Å.

By analogy with Cu I the writer has assumed the ground state indicated above.

## REFERENCE

B. Edlén, *Physica* 13, No. 9, 549 (1947).

March 1953.

## Cd XXII

(Co I sequence; 27 electrons)

Z=48

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>9</sup> D<sub>5/2</sub>

3d<sup>9</sup> D<sub>5/2</sub>

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed two lines due to the transition 3p<sup>6</sup> 3d<sup>9</sup> D-3p<sup>6</sup> 3d<sup>10</sup> P°, in the region between 40 Å and 80 Å. In figure 4 of his paper on the spectra of highly-ionized atoms, the observed wave numbers are plotted against atomic number for this combination in the Co I-like spectra Pd XX to Sn XXIV. For Cd XXII the wave numbers are between 1600000 K and 2000000 K.

## REFERENCE

B. Edlén, *Physica* 13, No. 9, 548 (1947).

March 1953.

## INDIUM

In I

49 electrons

Z=49

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^2 P_{1/2}^0$  $5p^2 P_{1/2}^0$  46669.93 K

I. P. 5.785 volts

Approximately 145 In I lines in the interval 1169 Å to 12912 Å have been classified. The most extensive term list is given in Paschen's 1938 paper, which includes the  $5p^2 {}^4P$  term, reported by Sawyer and Lang and by Lansing, evaluated from measurements of hyperfine-structure components.

Paschen's level called "A", which he designates as part of the  $6s^2 {}^4P$  term, has been included in the table with the writer's tentative assignment of  $J=2\frac{1}{2}$ .

Edlén has pointed out that the term for which Paschen (1938) suggested the designation  $5s 5p^2 {}^2D$  "must be  $4f {}^2F^0$ , as it was originally identified in earlier papers".

The terms  $5f {}^2F^0$  and  $6f {}^2F^0$  have been added from infrared observations by Meggers and Murphy in the interval 6847 Å to 12912 Å. Their paper contains an excellent summary of the work on this spectrum. The term  $5p^2 {}^2P$  has been published by Garton and also by Clearman; and the terms  $5p^2 {}^2S$ ,  $6p^2 {}^2S$  are given in Garton's 1950 paper. The writer has received from Garton in advance of publication his manuscript giving revised values of these terms, and also the series terms  $np^2 {}^2S$  ( $n=7, 8$ ),  $ns {}^2S$  ( $n=16$  to 24), and  $nd {}^2D$  ( $n=16$  to 30). He has observed the new series members in absorption, and from the  ${}^2S$  series derives the limit  $46670.5 \pm 0.7$  K.

Brackets in the table denote estimated values extrapolated from the series. Paschen's value of the limit, based on well-observed series, is quoted in the table. He derived it from an empirical formula of the Hicks type.

Observed intersystem combinations connect the quartet terms with the doublets.

## REFERENCES

- R. A. Sawyer and R. J. Lang, *Phys. Rev.* **34**, 719 (1929). (T) (C L)  
 W. D. Lansing, *Phys. Rev.* **34**, 506 (1929). (C L)  
 J. Okubo and S. Satô, *Reports Tohoku Imp. Univ. (Sendai)* [1] Prof. K. Honda Anniversary Vol. p. 8 (1936). (Z E) (hfs)  
 F. Paschen, *Ann. der Phys.* [5] **33**, 148 (1938). (I P) (T) (C L) (hfs)  
 W. R. S. Garton, *Nature* **168**, 150, 1950; *Proc. Phys. Soc. London [A]* **64**, 509 (1951). (T) (C L)  
 H. E. Clearman, *J. Opt. Soc. Am.* **43**, 374 (1952). (T) (C L)  
 W. F. Meggers and R. J. Murphy, *J. Research Nat. Bur. Std.* **48**, 334, RP 2320 (1953). (I P) (T) (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 W. R. S. Garton, *Proc. Phys. Soc. London [A]* **67**, 864 (1954). (I P) (T) (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955). (Summary hfs)  
 B. Edlén, letter (February 1958). (T)

In I

In I

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
5 <sup>a</sup> (18)5p	5p 1P°	0% 1%	0.00 2912.59	2312.58	5 <sup>a</sup> (18)11d	11d 1D	1% 2%	45211.51 45221.08	9.57
5 <sup>a</sup> (18)6e	6e 1S	0%	24372.87		5 <sup>a</sup> (18)9f	9f 1F°	3% 2%	45308.31 45303.39	-0.08
5 <sup>a</sup> (18)6p	6p 1P°	0% 1%	31216.61 32114.79	298.18	5 <sup>a</sup> (18)13e	13e 1S	0%	45394.18	
5 <sup>a</sup> (18)8d	8d 1D	1% 2%	33992.12 33915.42	23.30	5 <sup>a</sup> (18)12d	12d 1D	1% 2%	45493.98 45502.04	8.06
5 <sup>a</sup> 5p <sup>a</sup>	5p <sup>a</sup> 1P	0% 1% 2%	34977.66 36020.80 37451.90	1043.14 1481.10	5 <sup>a</sup> (18)10f	10f 1F°	3% 2%	45568.89 45564.04	-0.15
5 <sup>a</sup> (18)7e	7e 1S	0%	36301.69		5 <sup>a</sup> (18)14e	14e 1S	0%	45630.44	
5 <sup>a</sup> (18)7p	7p 1P°	0% 1%	38961.35 38978.84	111.49	5 <sup>a</sup> (18)13d	13d 1D	1% 2%	45702.08 45709.02	6.94
5 <sup>a</sup> (18)6d	6d 1D	1% 2%	39043.45 39068.37	49.89	5 <sup>a</sup> (18)11f	11f 1F°	2% 3%	45766.50 45766.56	0.06
5 <sup>a</sup> (18)4f	4f 1F°	2% 3%	39707.89 39707.96	0.67	5 <sup>a</sup> (18)15e	15e 1S	0%	45806.88	
5 <sup>a</sup> (18)8e	8e 1S	0%	40636.71		5 <sup>a</sup> (18)14d	14d 1D	1% 2%	45859.30 45865.32	6.02
5 <sup>a</sup> (18)8p	8p 1P°	0% 1%	41887.10 41881.44	54.34	5 <sup>a</sup> (18)12f	12f 1F°	2% 3%	45908.98	
5 <sup>a</sup> (18)7d	7d 1D	1% 2%	41836.28 41861.57	25.34	5 <sup>a</sup> (18)16e	16e 1S	0%	45941.9	
5 <sup>a</sup> (18)5f	5f 1F°	2% 3%	42290.0		5 <sup>a</sup> (18)15d	15d 1D	1% 2%	45982.2 45986.69	4.5
5 <sup>a</sup> (18)9e	9e 1S	0%	42718.55		5 <sup>a</sup> (18)17e	17e 1S	0%	46047.4	
5 <sup>a</sup> (18)8d	8d 1D	1% 2%	43335.63 43354.62	18.99	5 <sup>a</sup> (18)16d	16d 1D	1% 2%	46078.6	
5 <sup>a</sup> (18)9p	9p 1P°	0% 1%	43399.09 43399.53	30.44	5 <sup>a</sup> (18)18e	18e 1S	0%	46131.6	
5 <sup>a</sup> (18)6f	6f 1F°	2% 3%	43584.4		5 <sup>a</sup> (18)17d	17d 1D	1% 2%	46156.1	
5 <sup>a</sup> (18)10e	10e 1S	0%	43680.83		5 <sup>a</sup> (18)19e	19e 1S	0%	46200.0	
5 <sup>a</sup> (18)9d	9d 1D	1% 2%	44234.26 44249.37	15.11	5 <sup>a</sup> (18)18d	18d 1D	1% 2%	46219.9	
5 <sup>a</sup> (18)10p	10p 1P°	0% 1%	44375.17 44394.13	18.96	5 <sup>a</sup> (18)20e	20e 1S	0%	46255.6	
5 <sup>a</sup> (18)7f	7f 1F°	2% 3%	44408.80 44408.88	0.08	5 <sup>a</sup> (18)19d	19d 1D	1% 2%	46272.3	
5 <sup>a</sup> (18)11e	11e 1S	0%	44595.86		5 <sup>a</sup> (18)21e	21e 1S	0%	46302.1	
5 <sup>a</sup> (18)10d	10d 1D	1% 2%	44815.06 44827.02	11.96	5 <sup>a</sup> (18)20d	20d 1D	1% 2%	46316.0	
5 <sup>a</sup> (18)11p	11p 1P°	0% 1%	[44853.28] 44865.79	[12.56]	5 <sup>a</sup> (18)21d	21d 1D	1% 2%	46352.7	
5 <sup>a</sup> (18)8f	8f 1F°	3% 2%	44988.70 44988.71	-0.01	5 <sup>a</sup> (18)23e	23e 1S	0%	46374.9	
5 <sup>a</sup> (18)12e	12e 1S	0%	45067.19		5 <sup>a</sup> (18)22d	22d 1D	1% 2%	46384.3	

## In I—Continued

## In I—Continued

Config.	Desig.	<i>J</i>	Level	Interval	Config.	Desig.	<i>J</i>	Level	Interval
5s <sup>2</sup> ( <sup>1</sup> S)24s	24s <sup>1</sup> S	0½	46402.6		5s <sup>2</sup> ( <sup>1</sup> S)30d	30d <sup>1</sup> D	1½ 2½	46525.5	
5s <sup>2</sup> ( <sup>1</sup> S)23d	23d <sup>1</sup> D	1½ 2½	46410.6						
5s <sup>2</sup> ( <sup>1</sup> S)24d	24d <sup>1</sup> D	1½ 2½	46434.5		In II( <sup>1</sup> S <sub>0</sub> )	Limit		46669.93	
5s <sup>2</sup> ( <sup>1</sup> S)25d	25d <sup>1</sup> D	1½ 2½	46455.2		5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> S	0½	59118	905
5s <sup>2</sup> ( <sup>1</sup> S)26d	26d <sup>1</sup> D	1½ 2½	46472.5		5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> P	0½ 1½	59657 60652	
5s <sup>2</sup> ( <sup>1</sup> S)27d	27d <sup>1</sup> D	1½ 2½	46488.1		5s 5p( <sup>1</sup> P°)6s	6s' <sup>1</sup> P°	0½ 1½ 2½	68321.7	
5s <sup>2</sup> ( <sup>1</sup> S)28d	28d <sup>1</sup> D	1½ 2½	46501.7		5s 5p( <sup>1</sup> P°)6p	6p' <sup>1</sup> S	0½	77997	
5s <sup>2</sup> ( <sup>1</sup> S)29d	29d <sup>1</sup> D	1½ 2½	46514.2		5d 5p( <sup>1</sup> P°)7p	7p' <sup>1</sup> S	0½	84651	
					5s 5p( <sup>1</sup> P°)8p	8p' <sup>1</sup> S	0½	87700	

February 1958.

## In I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>2</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>10</sup> +	Observed Terms			
5s <sup>2</sup> ( <sup>1</sup> S)5p	5p <sup>1</sup> P°			
5s 5p <sup>2</sup>	{	5p <sup>2</sup> <sup>1</sup> S	5p <sup>2</sup> <sup>1</sup> P	5p <sup>2</sup> <sup>1</sup> P
		n <sub>s</sub> (n ≥ 6)	n <sub>p</sub> (n ≥ 6)	n <sub>d</sub> (n ≥ 5)
5s <sup>2</sup> ( <sup>1</sup> S)ns		6-24s <sup>1</sup> S	6-11p <sup>1</sup> P°	5-30d <sup>1</sup> D
5s 5p( <sup>1</sup> P°)ns'	{	6s' <sup>1</sup> P°	6-8p' <sup>1</sup> S	4-12f <sup>1</sup> F°

\*For predicted terms in the spectra of the In I isoelectronic sequences, see Vol. III, Introduction.

## In II

(Cd I sequence; 48 electrons)

Z=49

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 \text{ } ^1S_0$  $5s^2 \text{ } ^1S_0$ , 163195 K

I. P. 18.86 volts

The analysis is from Paschen and Campbell, who have revised and extended the earlier work by Lang and Sawyer, and by others. Approximately 570 lines are classified between 680 Å and 9246 Å. Many lines exhibit conspicuous hyperfine structure which Paschen and Campbell discuss in detail along with their analysis of the fine structure. Each member of the G-series consists of one pair of double levels called  $n^+G_{3,4}$  and  $n^-G_{3,4}$  in the paper on analysis. In the table the writer has labeled the  $n^+G_{3,4}$  pair as  $^{1,2}G_{3,4}$  and the  $n^-G_{3,4}$  pair as  $^3G_{3,4}$  in accordance with the suggestion of Trees. By analogy, the  $^+H$  and  $^-H$  series are handled similarly in the table. There is evidently a mixing of  $J$ -values among the four levels of each series member.

Paschen and Campbell list absolute term values based on the H-series as derived by means of a Rydberg formula without a Ritz correction. Their limit, quoted above, is 18 K lower than the earlier value, by Lang and Sawyer.

Observed inter-system combinations connect the singlet and triplet systems of terms.

## REFERENCES

- R. J. Lang and R. A. Sawyer, Zeit. Phys. 71, 453 (1931). (I P) (T) (C L)  
 F. Paschen und J. S. Campbell, Ann. der Phys. [5] 81, 29 (1933). (I P) (T) (C L) (hfs)  
 J. E. Mack, Rev. Mod. Phys. 22, No. 1, 64 (1950). (Summary hfs)  
 R. E. Trees, unpublished material (August 1953).

## In II

## In II

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$5s^2$	$5s^2 \text{ } ^1S$	0	0		$5s(5S)7s$	$7s \text{ } ^3S$	1	121437. 71	
$5s(5S)5p$	$5p \text{ } ^3P^o$	0	48875	1074	$5s(5S)7s$	$7s \text{ } ^1S$	0	123368. 04	
		1	43549	2478	$5s(5S)4f$	$4f \text{ } ^3F^o$	2	123657. 85	5. 19
		2	45827				3	123643. 04	16. 68
$5s(5S)5p$	$5p \text{ } ^1P^o$	1	63033. 81				4	123659. 72	
$5s(5S)6s$	$6s \text{ } ^3S$	1	93919. 03		$5s(5S)4f$	$4f \text{ } ^1F^o$	3	123694. 07	
$5s(5S)6s$	$6s \text{ } ^1S$	0	97025. 36		$5s(5S)6d$	$6d \text{ } ^3D$	1	124737. 90	33. 97
$5p^2$	$5p^2 \text{ } ^1D$	2	97623. 61				2	124771. 87	53. 43
$5p^2$	$5p^2 \text{ } ^3P$	0	101603. 3	1641. 2			3	124825. 30	
		1	102244. 47	2315. 72	$5s(5S)6d$	$6d \text{ } ^1D$	2	126666. 16	
		2	105560. 19		$5s(5S)7p$	$7p \text{ } ^3P^o$	0	126928. 03	62. 14
$5s(5S)5d$	$5d \text{ } ^3D$	1	102083. 63	86. 00			1	126990. 17	259. 19
		2	102169. 63	133. 60	$5s(5S)7p$	$7p \text{ } ^1P^o$	1	127568. 53	
		3	102308. 23		$5s(5S)8s$	$8s \text{ } ^3S$	1	133067. 67	
$5s(5S)6p$	$6p \text{ } ^3P^o$	0	107657. 87	179. 30	$5s(5S)8s$	$8s \text{ } ^1S$	0	133549. 12	
		1	107837. 17	588. 35	$5s(5S)5f$	$5f \text{ } ^3F^o$	2	133935. 63	4. 00
		2	108425. 52				3	133939. 63	16. 08
$5s(5S)6p$	$6p \text{ } ^1P^o$	1	108775. 39				4	133955. 71	
$5s(5S)5d$	$5d \text{ } ^1D$	2	113880. 07						
$5p^2$	$5p^2 \text{ } ^1S$	0	121284. 71						

## In II—Continued

## In II—Continued

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
5e(8)8f	8f 1F°	3	133979.84		5e(8)8g	8g 2G 1.2G	3, 4 4, 5	145295.24 145298.81	3.57
5e(8)8g	8g 2G 1.2G	3, 4 4, 5	134507.18 134511.01	3.83	5e(8)8h	8h 2H°	4, 5 5, 6	145331.51 145335.05	3.54
5e(8)7d	7d 2D	1 2 3	134721.67 134739.19 134767.07	17.52 27.88	5e(8)10d	10d 2D	1 2 3	145378.26 145382.99 145390.20	4.73 7.21
5e(8)7d	7d 2D	2	135396.63		5e(8)10d	10d 2D	2	145494.76	
5e(8)8p	8p 2P°	0 1 2	135856.79 135894.5	137.8	5e(8)11p	11p 2P°	0 1 2	145650.93	
5e(8)8p	8p 2P°	1	136092.10		5e(8)11p	11p 2P°	1	145678.91	
5e(8)9e	9e 2S	1	139132.27		5e(8)12a	12a 2S	1	146532.37	
5e(8)9e	9e 2S	0	139332.53		5e(8)12a	12a 2S	0	146598.23	
5e(8)6f	6f 2F°	2 3 4	139544.50 139547.83 139552.41	3.33 14.58	5e(8)9f	9f 2F°	2 3 4	146609.33 146607.43 146619.81	-1.90 12.38
5e(8)6f	6f 2F°	3	139581.23		5e(8)9f	9f 2F°	3	146633.49	
5e(8)6g	6g 2G 1.2G	3, 4 4, 5	139916.70 139920.44	3.74	5e(8)9g	9g 2G 1.2G	3, 4 4, 5	146745.44 146749.04	3.60
5e(8)8d	8d 2D	1 2 3	140077.42 140087.88 140104.34	10.46 16.46	5e(8)9h	9h 2H°	4, 5 5, 6	146764.61 146768.15	3.54
5e(8)8d	8d 2D	2	140403.60		5e(8)11d	11d 2D	1 2 3	146807.73 146810.4 146815.83	2.7 5.4
5e(8)9p	9p 2P°	0 1 2	140729.75 140817.6	87.8	5e(8)11d	11d 2D	2	146885.43	
5e(8)9p	9p 2P°	1	140840.13		5e(8)13a	13a 2S	1	147629.77	
5e(8)10e	10e 2S	1	142703.52		5e(8)10f	10f 2F°	2 3 4	147677.75 147675.63 147687.14	-2.12 11.51
5e(8)10e	10e 2S	0	142852.13		5e(8)13a	13a 2S	0	147676.63	
5e(8)7f	7f 2F°	2 3 4	142922.96 142925.87 142938.34	2.41 12.97	5e(8)10f	10f 2F°	3	147699.67	
5e(8)7f	7f 2F°	3	142954.91		5e(8)10g	10g 2G 1.2G	3, 4 4, 5	147782.28 147785.81	3.53
5e(8)7g	7g 2G 1.2G	3, 4 4, 5	143178.79 143182.34	3.55	5e(8)10h	10h 2H° 1.2H°	4, 5 5, 6	147796.52 147800.08	3.54
5e(8)9d	9d 2D	1 2 3	143292.63 143300.48 143311.02	6.85 10.54	5e(8)12d	12d 2D	1 2 3	147828.63 147831.50 147835.53	2.87 4.03
5e(8)9d	9d 2D	2	143479.10		5e(8)12d	12d 2D	2	147884.13	
5e(8)10p	10p 2P°	0 1 2	143701.70 143736.87	34.1	5e(8)14a	14a 2S	1	148436.04	
5e(8)10p	10p 2P°	1	143758.23		5e(8)11f	11f 2F°	2 3 4	148467.23 148463.68 148475.19	-3.61 11.57
5e(8)11a	11a 2S	1	144985.21		5e(8)14a	14a 2S	0	148470.73	
5e(8)11a	11a 2S	0	145081.50		5e(8)11f	11f 2F°	3	148486.65	
5e(8)8f	8f 2F°	2 3 4	145110.98 145118.70 145124.50	1.72 11.80	5e(8)11g	11g 2G 1.2G	3, 4 4, 5	148549.25 148552.80	3.55
5e(8)8f	8f 2F°	3	145139.69		5e(8)11h	11h 2H° 1.2H°	4, 5 5, 6	148660.11 148663.65	3.54



## In II—Continued

## In II—Continued

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
5s(7S)13d	13d <sup>1</sup> D	1	148584. 8	2. 8	5s(7S)15d	15d <sup>1</sup> D	2	149633. 23	
		2	148587. 6	3. 3			2	149633. 23	
		3	148590. 9				1	149692. 67	
5s(7S)13d	13d <sup>1</sup> D	2	149626. 13		5s(7S)17a	17a <sup>3</sup> S	1	149692. 67	
5s(7S)15e	15e <sup>3</sup> S	1	149046. 06		5s(7S)17a	17a <sup>1</sup> S	0	[149006. 8]	
5s(7S)12f	12f <sup>3</sup> F <sup>o</sup>	2	149065. 89	-3. 14 11. 17	5s(7S)14g	14g <sup>1</sup> G	3, 4	149945. 47	3. 54
		3	149068. 78		4, 5	149949. 01			
		4	149073. 89		5s(7S)14h	14h <sup>3</sup> H <sup>o</sup>	4, 5 5, 6	149951. 19 149954. 70	
5s(7S)15e	15e <sup>1</sup> S	0	149072. 46		5s(7S)16d	16d <sup>1</sup> D	1	149963. 4	3. 3 0. 8
5s(7S)12f	12f <sup>1</sup> F <sup>o</sup>	3	149084. 04		2	149966. 7			
3	149084. 04		3	149967. 5					
5s(7S)12g	12g <sup>3</sup> G	3, 4 4, 5	149132. 24 149135. 78	3. 54	5s(7S)16d	16d <sup>1</sup> D	2	[149983. 88]	
5s(7S)12h	12h <sup>3</sup> H <sup>o</sup>	4, 5 5, 6	149140. 90 149144. 44	3. 54	5s(7S)18e	18e <sup>3</sup> S	1	[150193. 12]	
5s(7S)14d	14d <sup>1</sup> D	1	149160. 5	3. 0 1. 9	5s(7S)17d	17d <sup>1</sup> D	1, 2 3	150253. 2 150253. 97	0. 8
		2	149163. 5		2	150267. 05			
		3	149165. 4		5s(7S)19e	19e <sup>3</sup> S	1	[150438. 41]	
5s(7S)14d	14d <sup>1</sup> D	2	[149192. 0]		5s(7S)18d	18d <sup>1</sup> D	1 2, 3	150486. 3 150488. 6	2. 3
5s(7S)16e	16e <sup>3</sup> S	1	149519. 15		5s(7S)18d	18d <sup>1</sup> D	2	[150499. 27]	
5s(7S)16e	16e <sup>1</sup> S	0	149539. 30		5s(7S)19d	19d <sup>1</sup> D	1 2 3	150683. 0	
5s(7S)13g	13g <sup>3</sup> G <sup>1,3</sup> G	3, 4	149585. 71	3. 54					
		4, 5	149589. 25		In III(7S <sub>0</sub> )	Limit		152195	
5s(7S)13h	13h <sup>3</sup> H <sup>o</sup>	4, 5	149598. 78	3. 54					
		5, 6	149599. 89						
		3	149609. 4		0. 5				
5s(7S)15d	15d <sup>1</sup> D	2	149609. 86	2. 8					
		3	149612. 7						
		3	149612. 7						

August 1953.

## In II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 3p <sup>2</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>10</sup> +	Observed Terms			
5s <sup>2</sup>	5s <sup>2</sup> <sup>1</sup> S			
5p <sup>2</sup>	{	5p <sup>2</sup> <sup>1</sup> S	5p <sup>2</sup> <sup>3</sup> P	
			5p <sup>2</sup> <sup>1</sup> D	
5s(7S)nz	{	ns (n ≥ 6)	np (n ≥ 5)	nd (n ≥ 5)
		6-17a <sup>3</sup> S 6-16a <sup>1</sup> S	5-11p <sup>3</sup> P <sup>o</sup> 5-11p <sup>1</sup> P <sup>o</sup>	5-19d <sup>3</sup> D 5-13, 15, 17d <sup>1</sup> D
		nf (n ≥ 4)	ng (n ≥ 5)	nh (n ≥ 6)
5s(7S)nz	{	4-12f <sup>3</sup> F <sup>o</sup> 4-12f <sup>1</sup> F <sup>o</sup>	5-14g <sup>1,3</sup> G	8-14h <sup>1,3</sup> H <sup>o</sup>

\*For predicted terms in the spectra of the Cd I isoelectronic sequence, see Vol. III, Introduction.

## In III

(Ag I sequence; 47 electrons)

Z=49

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 S_{1/2}$  $5s^2 S_{1/2}$ , 226100 K

I. P. 28.03 volts

The analysis is from Nodwell, who has furnished his tentative results in advance of publication, especially for inclusion here. He has classified 56 lines in the interval 685.31 Å to 6197.72 Å. The  $5p'$   $^1P^o$  term requires further confirmation.

The limit is from Catalán and Rico, who have derived it by comparison of the third spectra from Y to In.

## REFERENCES

R. Nodwell, unpublished material (December 1955). (I P) (T) (C L)

M. A. Catalán y F. R. Rico, An. Real Soc. Esp. Física y Química (Madrid) [A] 53, 85 (1957). (I P)

## In III

## In III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^{10}(^1S)5s$	$5s^2 S$	$0\frac{1}{2}$	0. 0		$4d^{10}(^1S)6d$	$6d^2 D$	$1\frac{1}{2}$ $2\frac{1}{2}$	170531. 8 170715. 3	183. 5
$4d^{10}(^1S)5p$	$5p^2 P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	57181. 1 61523. 2	4342. 1	$4d^{10}(^1S)7p$	$7p^2 P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	180551. 87 181150. 87	599. 3
$4d^{10} 5s^2$	$5s^2 ^2D$	$2\frac{1}{2}$ $1\frac{1}{2}$	115568. 7 122414. 8	-6846. 1	$4d^{10}(^1S)5g$	$5g^2 G$	$3\frac{1}{2}, 4\frac{1}{2}$	186522. 0	
$4d^{10}(^1S)6s$	$6s^2 S$	$0\frac{1}{2}$	126874. 2		$4d^{10}(^1S)8s$	$8s^2 S$	$0\frac{1}{2}$	189369. 6	
$4d^{10}(^1S)5d$	$5d^2 D$	$1\frac{1}{2}$ $2\frac{1}{2}$	128452. 8 128742. 5	289. 7	$4d^{10}(^1S)7d$	$7d^2 D$	$1\frac{1}{2}$ $2\frac{1}{2}$	190033. 5 190131. 9	98. 4
$4d^{10}(^1S)6p$	$6p^2 P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	144583. 6 145922. 6	1339. 0	$4d^{10}(^1S)6g$	$6g^2 G$	$3\frac{1}{2}, 4\frac{1}{2}$	198647. 9	
$4d^{10} 5s(^2D)5p$	$5p^2 ^1P^o$	$2\frac{1}{2}$ $1\frac{1}{2}$ $0\frac{1}{2}$	1588287 1676227 171768. 47	-8900 -4140	$4d^{10}(^1S)9s$	$9s^2 S$	$0\frac{1}{2}$	201149. 07	
$4d^{10}(^1S)4f$	$4f^2 ^1F^o$	$2\frac{1}{2}$ $3\frac{1}{2}$	161968. 8 161976. 8	8. 0	$4d^{10}(^1S)7g$	$7g^2 G$	$3\frac{1}{2}, 4\frac{1}{2}$	205961. 3	
$4d^{10}(^1S)7s$	$7s^2 S$	$0\frac{1}{2}$	169429. 8		In iv( $^1S_0$ )	Limit	-----	226100	

February 1957.

## In IV

(Pd I sequence; 46 electrons)

Z=49

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^9 \text{ } ^1\text{S}_0$  $4d^9 \text{ } ^1\text{S}_0$ , 439000 K

I. P. 54.4 volts

The analysis is by Gibbs and White, who have classified 36 lines between 472.48 Å and 1725.91 Å. Observed combinations connect the singlet and triplet systems of terms.

The writer has interpolated the limit quoted here, from the isoelectronic spectra Pd I through Sn v. This gives an ionization potential 3.5 volts less than the one quoted by Kruger and Shoupp, which in turn is from a Moseley diagram by Gibbs and White.

Crooker and Nodwell have recently observed the spectrum with a vacuum-spark source and will doubtless be able to revise and extend the present analysis.

## REFERENCES

- R. C. Gibbs and H. E. White, *Phys. Rev.* **31**, 776 (1928). (I P) (T) (C L)  
 P. G. Kruger and W. E. Shoupp, *Phys. Rev.* **46**, 124 (1934). (I P)  
 A. M. Crooker and R. Nodwell, letter (December 1955).

## In IV

## In IV

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^9$	$4d^9 \text{ } ^1\text{S}$	0	0		$4d^9(^1D)5p$	$5p \text{ } ^3D^\circ$	3 2 1	<del>205953</del> <del>202129</del> <del>211650</del>	3824 -9521
$4d^9(^1D_{3/2})5s$	$5s \text{ } ^3D$	3	128785	-2196					
		2	130981						
$4d^9(^1D_{3/2})5s$		1	135893	-4912	$4d^9(^1D)5p$	$5p \text{ } ^1P^\circ$	1	208708	
$4d^9(^1D_{3/2})5s$	$5s \text{ } ^1D$	2	138764		$4d^9(^1D)5p$	$5p \text{ } ^1F^\circ$	3	208886	
$4d^9(^1D)5p$	$5p \text{ } ^3P^\circ$	2	194004	-6658	$4d^9(^1D)5p$	$5p \text{ } ^1D^\circ$	2	212785	
		1	200682	-4395					
		0	205057		-----	-----	-----	-----	
$4d^9(^1D)5p$	$5p \text{ } ^1F^\circ$	4	201158	4452	In $v(^1D_{3/2})$	Limit	----	439000	
		3	196708	-8651					
		2	205357						

January 1956.

## In V

(Rh sequence; 45 electrons)

Z=49

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^4 4d^2 \text{ } ^1D_{3/2}$  $4d^2 \text{ } ^1D_{3/2}$ 

K

I. P.

volts

The spectrum is incompletely known. The analysis is from Green, who has classified 43 lines between 368.67 Å and 423.16 Å. Observed intersystem combinations connect the doublet and quartet systems of terms. No series have been found.

## REFERENCE

M. Green, Phys. Rev. **68**, 117 (1941). (T) (C L)

In V

In V

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval																																																																																																																										
$4d^2$	$^1D$	$3\frac{1}{2}$	0	-7165	$4d^2(^1D)5p$	$5p'' \text{ } ^1F^{\circ}$	$2\frac{1}{2}$	258598	2662																																																																																																																										
		$1\frac{1}{2}$	7165				$3\frac{1}{2}$	261390		$4d^2(^1F)5p$	$^1D^{\circ}$	$3\frac{1}{2}$	236317	-4734	$4d^2(^1D)5p$	$5p'' \text{ } ^1P^{\circ}$	$0\frac{1}{2}$	261230	852	$2\frac{1}{2}$	241061	$1\frac{1}{2}$	262082	$1\frac{1}{2}$	246142	$5p'' \text{ } ^1D^{\circ}$	$2\frac{1}{2}$	262971	$0\frac{1}{2}$		$1\frac{1}{2}$	263567	$4d^2(^1F)5p$	$^1G^{\circ}$	$5\frac{1}{2}$		-585	$4d^2(^1P)5p$	$5p' \text{ } ^1D^{\circ}$	$3\frac{1}{2}$	264124	-1947	$4\frac{1}{2}$		$2\frac{1}{2}$	266071	$3\frac{1}{2}$		$1\frac{1}{2}$	265908	163	$2\frac{1}{2}$	245128	$0\frac{1}{2}$		$4d^2(^1F)5p$	$^1F^{\circ}$	$4\frac{1}{2}$		-4176	$4d^2(^1P)5p$	$5p' \text{ } ^1D^{\circ}$	$1\frac{1}{2}$	268226	623	$3\frac{1}{2}$	248515	$2\frac{1}{2}$	268849	$2\frac{1}{2}$	252691	5978		$1\frac{1}{2}$	246713			$4d^2(^1F)5p$	$^1D^{\circ}$	$2\frac{1}{2}$	249644	-3004	$4d^2(^1P)5p$	$5p' \text{ } ^1P^{\circ}$	$1\frac{1}{2}$	270197	-5108	$1\frac{1}{2}$	252648	$0\frac{1}{2}$	275305	$4d^2(^1F)5p$	$^1F^{\circ}$	$3\frac{1}{2}$		-165	$4d^2(^1P)5p$	$5p' \text{ } ^1S^{\circ}$	$0\frac{1}{2}$	270850		$2\frac{1}{2}$	253878	$3\frac{1}{2}$	271245	$4d^2(^1F)5p$	$^1G^{\circ}$	$4\frac{1}{2}$			$4d^2(^1G)5p$	$5p''' \text{ } ^1F^{\circ}$	$2\frac{1}{2}$	272585	-1320	$3\frac{1}{2}$	254803			$4d^2(^1P)5p$	$^1P^{\circ}$	$2\frac{1}{2}$	257129		$4d^2(^1P)5p$	$5p' \text{ } ^1S^{\circ}$	$1\frac{1}{2}$	273867		$1\frac{1}{2}$	257294
$4d^2(^1F)5p$	$^1D^{\circ}$	$3\frac{1}{2}$	236317	-4734	$4d^2(^1D)5p$	$5p'' \text{ } ^1P^{\circ}$	$0\frac{1}{2}$	261230	852																																																																																																																										
		$2\frac{1}{2}$	241061				$1\frac{1}{2}$	262082																																																																																																																											
		$1\frac{1}{2}$	246142				$5p'' \text{ } ^1D^{\circ}$	$2\frac{1}{2}$				262971																																																																																																																							
		$0\frac{1}{2}$						$1\frac{1}{2}$		263567																																																																																																																									
$4d^2(^1F)5p$	$^1G^{\circ}$	$5\frac{1}{2}$		-585	$4d^2(^1P)5p$	$5p' \text{ } ^1D^{\circ}$	$3\frac{1}{2}$	264124	-1947																																																																																																																										
		$4\frac{1}{2}$					$2\frac{1}{2}$	266071																																																																																																																											
		$3\frac{1}{2}$					$1\frac{1}{2}$	265908		163																																																																																																																									
		$2\frac{1}{2}$	245128				$0\frac{1}{2}$																																																																																																																												
$4d^2(^1F)5p$	$^1F^{\circ}$	$4\frac{1}{2}$		-4176	$4d^2(^1P)5p$	$5p' \text{ } ^1D^{\circ}$	$1\frac{1}{2}$	268226	623																																																																																																																										
		$3\frac{1}{2}$	248515				$2\frac{1}{2}$	268849																																																																																																																											
		$2\frac{1}{2}$	252691				5978																																																																																																																												
		$1\frac{1}{2}$	246713																																																																																																																																
$4d^2(^1F)5p$	$^1D^{\circ}$	$2\frac{1}{2}$	249644	-3004	$4d^2(^1P)5p$	$5p' \text{ } ^1P^{\circ}$	$1\frac{1}{2}$	270197	-5108																																																																																																																										
		$1\frac{1}{2}$	252648				$0\frac{1}{2}$	275305																																																																																																																											
$4d^2(^1F)5p$	$^1F^{\circ}$	$3\frac{1}{2}$		-165	$4d^2(^1P)5p$	$5p' \text{ } ^1S^{\circ}$	$0\frac{1}{2}$	270850																																																																																																																											
		$2\frac{1}{2}$	253878				$3\frac{1}{2}$	271245																																																																																																																											
$4d^2(^1F)5p$	$^1G^{\circ}$	$4\frac{1}{2}$			$4d^2(^1G)5p$	$5p''' \text{ } ^1F^{\circ}$	$2\frac{1}{2}$	272585	-1320																																																																																																																										
		$3\frac{1}{2}$	254803																																																																																																																																
$4d^2(^1P)5p$	$^1P^{\circ}$	$2\frac{1}{2}$	257129		$4d^2(^1P)5p$	$5p' \text{ } ^1S^{\circ}$	$1\frac{1}{2}$	273867																																																																																																																											
		$1\frac{1}{2}$	257294																																																																																																																																
		$0\frac{1}{2}$																																																																																																																																	

February 1956.

## In v OBSERVED TERMS\*

Configuration $1s^2 2s^2 2p^6 3s^2$ $3p^6 3d^{10} 4s^2 4p^6 +$	Observed Terms
$4d^0$	$4d^0 D$
	$np (n \geq 5)$
$4d^1(^4F)ns$	$\left\{ \begin{array}{lll} 5p \ ^4D^0 & 5p \ ^4F^0 & 5p \ ^4G^0 \\ 5p \ ^4D^0 & 5p \ ^4F^0 & 5p \ ^4G^0 \end{array} \right.$
$4d^1(^4P)ns'$	$\left\{ \begin{array}{lll} 5p' \ ^4S^0 & 5p' \ ^4P^0 & 5p' \ ^4D^0 \\ 5p' \ ^4S^0 & 5p' \ ^4P^0 & 5p' \ ^4D^0 \end{array} \right.$
$4d^1(^1D)ns''$	$5p'' \ ^1P^0 \quad 5p'' \ ^1D^0 \quad 5p'' \ ^1F^0$
$4d^1(^1G)ns'''$	$5p''' \ ^1P^0$

\*For predicted terms in the spectra of the Rh I isoelectronic sequence, see Vol. III, Introduction.

## In XXI

(Cu I sequence; 29 electrons)

Z=49

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 S_{0g}$

$4s^2 S_{0g}$

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed the doublets due to the transitions  $4s-5p$ ,  $4p-5d$ ,  $4d-5f$ , and  $4p-5s$ , from Pd XVIII to In XXI. In figure 3 of his paper on the spectra of highly-ionized atoms, the lines are indicated on the photograph of vacuum spark spectra of In observed from 40 Å to 80 Å.

By analogy with Cu I the writer has assumed the ground state indicated above.

## REFERENCE

B. Edlén, *Physica* 12, No. 9, 549 (1947).

March 1953.

## In XXIII

(Co I sequence; 27 electrons)

Z=49

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 D_{3g}$

$3d^8 D_{3g}$

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed two lines due to the transition  $3p^6 3d^8 D-3p^6 3d^{10} P^0$ , in the region between 40 Å and 80 Å. In figure 4 of his paper on the spectra of highly-ionized atoms, the observed wave numbers are plotted against atomic number for this combination in the Co I-like spectra Pd XX to Sn XXIV. For In XXIII the wave numbers are between 1700000 K and 2000000 K.

## REFERENCE

B. Edlén, *Physica* 12, No. 9, 548 (1947).

March 1953.

## TIN

## Sn I

50 electrons

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^2 \ ^3P_1$  $5p^2 \ ^3P_1$ , 59231.8 K

I. P. 7.342 volts

In 1940 Meggers published an extensive analysis of Sn I in which he revised the earlier work by Back, Green and Loring, and others. His observations extend from 2000 Å to 13000 Å, and are supplemented by corrected radiometric observations of Randall and Wright between 12788 Å and 24738 Å. Spectrograms taken by Shenstone cover the short-wave interval 1400 Å to 2200 Å. Meggers lists 378 lines between 1697.59 Å and 24738 Å, of which 80 percent are classified. Observed intersystem combinations connect the singlet and triplet systems of terms.

Garton has observed the Sn I spectrum in absorption from 1300 Å to 1800 Å. He reports the discovery of about 50 new atomic lines and states that analysis of this extension of Sn I is in progress.

Barrow and Rowlinson have recently extended the ultraviolet observations of Sn I and reported a number of series lines above the ionization limit. All but nine levels in the table higher than 59453 are from their list. Owing to the difficulty of fitting these new series on to the known series, the writer submitted these data to Shenstone with the hope that he might be able to interpret them. This he has been able to do by means of a new spectrogram in the Schumann region connecting the separate regions of the spectrum. The present analysis is from his manuscript, prepared especially for inclusion here. He has made some revisions to Meggers' analysis and classified about one hundred additional lines. The limits quoted in the table are based chiefly on the series by Barrow and Rowlinson, the strongest ones being  $nd \ ^1P_1^o$  ( $n=8$  to 19) and  $nd \ ^3P_1^o$  ( $n=8$  to 27). Meggers derived the limit 59155.0K from the  $ns \ ^3P^o$  series ( $n=6$  to 15) and  $ns \ ^1P^o$  series ( $n=6$  to 8), by means of a Ritz formula.

The ground term of Sn II,  $5p \ ^3P^o$ , has an interval of 4251K. As a consequence of this wide interval, the levels of a given configuration in Sn I having this term as limit, divide themselves into two widely separated groups, the lower of which has as limit  $5p \ ^3P_{3/2}^o$ , and the higher has the limit  $5p \ ^3P_{1/2}^o$ . The  $J$ -values of the limit are quoted in the table so that the related pairs of levels can be distinguished. The  $J$ -coupling notation can be worked out from the auxiliary table that precedes the references below.

Back, and Green and Loring have observed the Zeeman effect for Sn I lines. Their results are in good agreement. The  $g$ -values in the table are chiefly from Back's paper. Some  $g$ -values of Sn I lines, measured as impurity lines by W. F. Meggers, C. C. Kiess, and C. J. Humphreys on spectrograms at the National Bureau of Standards, have been averaged with those of Back. This applies only to the levels of the  $5p^2$ - and  $6s$ -configurations.

In these Volumes the general format is to arrange the levels by terms. Shenstone emphasizes strongly that in the case of Sn I this is particularly misleading. In compliance with his urgent request, an exception is being made for Sn I, and the levels are repeated in a Supplementary Table in numerical order over the range in which overlapping occurs.

## Sm I—Continued

Author	Config.	Desig.	J	Author	Config.	Desig.	J
no <sup>1</sup> P <sub>1</sub> <sup>1</sup> P <sub>1</sub>	5p( <sup>1</sup> P <sub>1/2</sub> )ns	ns [0%] <sup>a</sup>	0 1	nd <sup>1</sup> F <sub>1</sub> <sup>1</sup> D <sub>1</sub>	5p( <sup>1</sup> P <sub>1/2</sub> )nd	nd [1%] <sup>a</sup>	2 1
no <sup>1</sup> P <sub>1</sub> <sup>1</sup> P <sub>1</sub>	5p( <sup>1</sup> P <sub>1/2</sub> )ns	ns' [1%] <sup>a</sup>	2 1	nd <sup>1</sup> D <sub>1</sub> <sup>1</sup> D <sub>1</sub>	"	nd [3%] <sup>a</sup>	2 3
np <sup>1</sup> P <sub>1</sub> <sup>1</sup> P <sub>1</sub>	5p( <sup>1</sup> P <sub>1/2</sub> )np	np [0%]	1 0	nd <sup>1</sup> D <sub>1</sub> <sup>1</sup> F <sub>1</sub>	5p( <sup>1</sup> P <sub>1/2</sub> )nd	nd' [2%] <sup>a</sup>	2 3
np <sup>1</sup> D <sub>1</sub> <sup>1</sup> D <sub>1</sub>	"	np [1%]	2 1	nd <sup>1</sup> P <sub>1</sub> <sup>1</sup> P <sub>1</sub>	"	nd' [1%] <sup>a</sup>	2 1
np <sup>1</sup> P <sub>1</sub> <sup>1</sup> S <sub>1/2</sub>	5p( <sup>1</sup> P <sub>1/2</sub> )np	np' [0%]	1 0	nd <sup>1</sup> P <sub>1</sub> <sup>1</sup> P <sub>1</sub>	"	nd' [0%] <sup>a</sup>	0 1
np <sup>1</sup> D <sub>1</sub> <sup>1</sup> P <sub>1</sub>	"	np' [2%]	3 2	nd <sup>1</sup> F <sub>1</sub> <sup>1</sup> F <sub>1</sub>	"	nd' [3%] <sup>a</sup>	4 3
np <sup>1</sup> S <sub>1/2</sub> <sup>1</sup> D <sub>1</sub>	"	np' [1%]	1 2				

## REFERENCES

- J. B. Green and R. A. Loring, *Phys. Rev.* **30**, 574 (1927). (I P) (T) (C L) (ZE)  
 E. Back, *Zeit. Phys.* **43**, 309 (1927). (T) (C L) (Z E)  
 H. M. Randall and N. Wright, *Phys. Rev.* **36**, 457 (1931). (I P) (T) (C L)  
 W. F. Meggers, *J. Research Nat. Bur. Std.* **24**, 153, RP 1275 (1940). (I P) (T) (C L)  
 W. R. S. Garton, *Nature* **168**, 690 (1950); *Proc. Phys. Soc. London [A]* **64**, 591 (1951). (T)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 R. F. Barrow and H. C. Rowlinson, unpublished material (Nov. 1953). (T) (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955).  
 (Summary hfs)  
 A. G. Shenstone, unpublished material (June 1956). (I P) (T) (C L)

## Sm I

## Sm I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> P	0	0.0	1691.8	0/0	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p	6p <sup>1</sup> P	1	46603.4		
		1	1691.8	1785.9	1.502	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p	6p <sup>1</sup> S	0	46936.37		
		2	3427.7		1.452	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 5d	5d <sup>1</sup> D°	2	47145.7		0.941
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> D	2	8613.0		1.062	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p	6p <sup>1</sup> S	1	47805.7		
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> S	0	17162.6		0/0	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p	6p <sup>1</sup> D	2	48189.7		
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6s	6s <sup>1</sup> P°	0	34640.8	273.4	0/0	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 7s	7s <sup>1</sup> P°	0	48216.2		
		1	34914.2	3714.6	1.280	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 7s		1	48222.1	5.9	1.316
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6s		2	38628.8		1.501	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 7s		2	52415.7	4193.6	
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6s	6s <sup>1</sup> P°	1	39257.1		1.121	5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 5d	5d <sup>1</sup> P°	2	48669.6	-312.4	1.406
5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> S°	2	39925.5			"	1	48922.0	-505.2	1.229	
		0	43430.0	-1087.7		5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 5d	5d <sup>1</sup> F°	3	49893.8		1.043
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p	6p <sup>1</sup> P	1	43242.3	4992.9		5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 5d	5d <sup>1</sup> P°	1	50125.9		1.066
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p		2	47235.2			5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 7p	7p <sup>1</sup> P	0	51374.8	-619.0	
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p	6p <sup>1</sup> D	1	43368.5	-129.8		5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 7p		1	50755.8	4431.1	
		2	43238.7	3768.1		5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 7p		2	55186.9		
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6p		3	47006.8			5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 6d	6d <sup>1</sup> D°	1	51475.1	-464.6	0.863
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 5d	5d <sup>1</sup> F°	2	43683.0	3905.0	0.865	"	2	51010.5	744.4		
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 5d		3	47488.0		1.246	"	3	51754.9			
		4				5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 7p	7p <sup>1</sup> D	1	51113.3	57.5	
5s <sup>2</sup> 5p( <sup>1</sup> P <sub>1/2</sub> ) 5d	5d <sup>1</sup> D°	1	44508.8	-364.2	0.635		2	51170.8	3969.1		
		2	44144.6	421.5	1.131		3	55139.9			
"		3	44576.1		1.167						





## Sn I—Continued

## Sn I—Continued

Config.	Desig.	J	Level	Interval	Obs. #	Config.	Desig.	J	Level	Interval	Obs. #
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 5f		37	59046.6			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 14d	14d <sup>1</sup> P°	1	62651.8		
<hr/>						5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 14d	14d <sup>1</sup> P°	1	62716.4		
Sn II ( <sup>2</sup> F <sub>3/2</sub> )	Limit	----	59231.8			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 16e	16e <sup>1</sup> P°	1	62745.9		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9e	9e <sup>1</sup> P°	1	59453.0			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 15d	15d <sup>1</sup> D°	2	62774.6		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>1</sup> P°	1	60051.4			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 15d	15d <sup>1</sup> P°	1	62776.2		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>1</sup> P°	1	60397.0			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 15d	15d <sup>1</sup> P°	1	62827.5		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9d	9d <sup>1</sup> P°	1	60643.4			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 17e	17e <sup>1</sup> P°	1	62852.6		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9d	9d <sup>1</sup> P°	1	61212.2			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 16d	16d <sup>1</sup> D°	2	62871.2		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9d	9d <sup>1</sup> F°	3	61247.5			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 16d	16d <sup>1</sup> P°	1	62878.1		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10d	10d <sup>1</sup> D°	2	61534.2			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 16d	16d <sup>1</sup> P°	1	62919.6		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10d	10d <sup>1</sup> P°	1	61556.3			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 17d	17d <sup>1</sup> D°	2	62954.2		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10d	10d <sup>1</sup> P°	1	61744.2			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 17d	17d <sup>1</sup> P°	1	62956.4		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10d	10d <sup>1</sup> F°	3	61762.0			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 17d	17d <sup>1</sup> P°	1	62962.5		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12e	12e <sup>1</sup> P°	1	61859.5			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 18d	18d <sup>1</sup> D°	2	63012.1		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 11d	11d <sup>1</sup> D°	2	61963.0			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 18d	18d <sup>1</sup> P°	1	63025.9		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 11d	11d <sup>1</sup> P°	1	61973.2			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 18d	18d <sup>1</sup> P°	1	63061.3		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 11d	11d <sup>1</sup> P°	1	62110.6			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 19d	19d <sup>1</sup> D°	2	63072.6		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 11d	11d <sup>1</sup> F°	3	62127			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 19d	19d <sup>1</sup> P°	1	63080.3		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12e	12e <sup>1</sup> P°	1	62191.5			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 19d	19d <sup>1</sup> P°	1	63104.2		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d <sup>1</sup> D°	2	62263.9			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 20d	20d <sup>1</sup> P°	1	63122.5		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d <sup>1</sup> P°	1	62267.3			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 21d	21d <sup>1</sup> P°	1	63160.4		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d <sup>1</sup> P°	1	62373.3			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 22d	22d <sup>1</sup> P°	1	63191.6		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d <sup>1</sup> F°	3	62385			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 23d	23d <sup>1</sup> P°	1	63220.7		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 14e	14e <sup>1</sup> P°	1	62431.0			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 24d	24d <sup>1</sup> P°	1	63244.3		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 13d	13d <sup>1</sup> D°	2	62483.1			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 25d	25d <sup>1</sup> P°	1	63267.9		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 13d	13d <sup>1</sup> P°	1	62484.0			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 26d	26d <sup>1</sup> P°	1	63283.5		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 13d	13d <sup>1</sup> P°	1	62564.5			5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 27d	27d <sup>1</sup> P°	1	63297.5		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 15e	15e <sup>1</sup> P°	1	62609.4								
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 14d	14d <sup>1</sup> D°	2	62647.2			Sn II ( <sup>2</sup> F <sub>3/2</sub> )	Limit	----	63483.2		

July 1956.

Sm I—SUPPLEMENTARY TABLE

Config.	Desig.	J	Level	Config.	Desig.	J	Level
5s <sup>2</sup> 5p <sup>3</sup>	5p <sup>3</sup> <sup>1</sup> P	0	0.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>1</sup> F°	2	51160.2
"	5p <sup>3</sup> <sup>3</sup> P	1	1691.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>3</sup> D	2	51170.8
"	5p <sup>3</sup> <sup>1</sup> P	2	3427.7	"	7p <sup>1</sup> P	0	51374.8
"	5p <sup>3</sup> <sup>3</sup> D	2	8613.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>3</sup> D°	1	51475.1
"	5p <sup>3</sup> <sup>3</sup> S	0	17162.6	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>1</sup> D°	3	51754.9
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6s	6s <sup>1</sup> P°	0	34640.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 4f		2?	52249.0
"	6s <sup>3</sup> P°	1	34914.2	"		2	52263.8
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6s	6s <sup>3</sup> P°	2	39693.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>3</sup> S	0	52265.3
"	6s <sup>1</sup> P°	1	39857.1	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7s	7s <sup>3</sup> P°	2	52415.7
5s 5p <sup>3</sup>	5p <sup>3</sup> <sup>3</sup> S°	2	39695.5	"	7s <sup>1</sup> P°	1	52706.8
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6p	6p <sup>3</sup> P	1	42342.3	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8s	8s <sup>3</sup> P°	1	53090.6
"	6p <sup>1</sup> D	2	42338.7	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7d	7d <sup>3</sup> D°	1	53592.2
"	6p <sup>3</sup> D	1	42368.5	"	7d <sup>1</sup> D°	2	53631.4
"	6p <sup>1</sup> P	0	42430.0	"	7d <sup>3</sup> D°	3	53826.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 5d	5d <sup>3</sup> F°	2	43683.0	"	7d <sup>1</sup> F°	2	54311.7
"	5d <sup>1</sup> D°	2	44144.6	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>3</sup> P°	2	54653.4
"	5d <sup>3</sup> D°	1	44508.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>3</sup> D°?	1	54712.8
"	5d <sup>1</sup> D°	3	44576.1	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 5f		3?	54762.6
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6p	6p <sup>1</sup> P	1	46603.4	"		2?	54767.7
"	6p <sup>3</sup> S	0	46936.3?	"		2?	54771.8
"	6p <sup>1</sup> D	3	47006.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>1</sup> D°	3	54830.5
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 5d	5d <sup>1</sup> D°	2	47145.7	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>1</sup> P	1	54990.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6p	6p <sup>3</sup> P	2	47235.2	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>3</sup> P°	1	55074.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 5d	5d <sup>3</sup> F°	3	47458.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>1</sup> D°	2	55130.9?
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6p	6p <sup>3</sup> S	1	47805.7	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>3</sup> D	3	55139.9
"	6p <sup>1</sup> D	2	48189.7	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 9s	9s <sup>3</sup> P°	1	55156.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7s	7s <sup>3</sup> P°	0	48216.2	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>3</sup> P	2	55186.9
"	7s <sup>1</sup> P°	1	48222.1	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>1</sup> D°	2	55226.1
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 5d	5d <sup>3</sup> P°	2	48689.6	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>3</sup> S	1	55373.8
"	5d <sup>1</sup> P°	1	48982.0	"	7p <sup>1</sup> D	2	55500.6
"	5d <sup>3</sup> P°	0	49487.2	5s 5p <sup>3</sup>	5p <sup>3</sup> <sup>3</sup> S°	1	55687.4
"	5d <sup>1</sup> F°	3	49893.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>3</sup> F°	3	55739.7
"	5d <sup>1</sup> P°	1	50125.9	"	6d <sup>3</sup> P°	0	55782.4
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>3</sup> P	1	50755.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>3</sup> F°	2	55803.2
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>1</sup> D°	2	51010.5	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>3</sup> P°	1	56244.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7p	7p <sup>1</sup> D	1	51113.3	"	6d <sup>1</sup> F°	3	56298.8

Sn I—SUPPLEMENTARY TABLE—Continued

Config.	Desig.	<i>J</i>	Level	Config.	Desig.	<i>J</i>	Level
		1 or 2	56357.4	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d <sup>2</sup> D°	2	58100.9
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 10s	10s <sup>1</sup> P°	1	56389.9	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 14s	14s <sup>1</sup> P°	1	58143.6
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 4f		3?	56396.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 15s	15s <sup>1</sup> P°	1	58324.3
"		3?	56400.7	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 16s	16s <sup>1</sup> P°	1	58523.3
"		2	56486.5	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 15d	15d <sup>1</sup> D°	1	58595.0
"		1?	56488.5	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 17s	17s <sup>1</sup> P°	1	58607.5
5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>2</sup> D°	2	56544.8	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 16d	16d <sup>1</sup> D°	1	58679.6
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 4f		3?	56550.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 18s	18s <sup>1</sup> P°	1	58690.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 4f	<sup>1</sup> D	1	56632.9	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 17d	17d <sup>1</sup> D°	1	58747.8
5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>2</sup> D°	1	56658.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 19s	19s <sup>1</sup> P°	1	58758.4
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7d	7d <sup>1</sup> F°?	3	56839.4	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7d	7d <sup>1</sup> P°	1	58970.3
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 11s	11s <sup>1</sup> P°	1	57004.5	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 5f		3?	59046.6
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 9d	9d <sup>1</sup> D°	2	57105.7	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7d	7d <sup>1</sup> F°	3	59190.3
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8s	8s <sup>1</sup> P°	2	57149.8				
5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>2</sup> D°	3	57181.6	Sn II( <sup>2</sup> P <sub>1/2</sub> )	Limit		59231.3
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8s	8s <sup>1</sup> P°	1	57288.3	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 9s	9s <sup>1</sup> P°	2	59375.9
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 10d	10d <sup>1</sup> D°	2	57533.7	"	9s <sup>1</sup> P°	1	59453.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 12s	12s <sup>1</sup> P°	1	57563.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>1</sup> P°	1	60051.4
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 11d	11d <sup>1</sup> D°	3	57847.0?	"	8d <sup>1</sup> P°	1	60397.0
"	11d <sup>1</sup> D°	2	57856.3	"	8d <sup>1</sup> F°	3	60450.0
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 13s	13s <sup>1</sup> P°	1	57899.0	5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 10s	10s <sup>1</sup> P°	2	60598.8
5s <sup>2</sup> 5p( <sup>2</sup> P <sub>1/2</sub> ) 7d	7d <sup>1</sup> P°	1	57934.7				

See remaining levels on page 77.

Sn I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>2</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>10</sup> +	Observed Terms				
	5s <sup>2</sup> 5p <sup>2</sup>	{	5p <sup>2</sup> <sup>1</sup> S	5p <sup>2</sup> <sup>1</sup> P	5p <sup>2</sup> <sup>1</sup> D
5s 5p <sup>2</sup>	{	5p <sup>2</sup> <sup>3</sup> S°		5p <sup>2</sup> <sup>3</sup> D°	
		ns (n ≥ 6)		np (n ≥ 6)	
5s <sup>2</sup> 5p( <sup>2</sup> P°) ns	{	6-19s <sup>1</sup> P°	6-9, 12-17s <sup>1</sup> P°	6, 7p <sup>1</sup> S	6, 7p <sup>1</sup> P
				6, 7p <sup>1</sup> S	6, 7p <sup>1</sup> P
		nd (n ≥ 5)			
5s <sup>2</sup> 5p( <sup>2</sup> P°) ns	{	5-27d <sup>1</sup> P°	5-12, 15-17d <sup>1</sup> D°	5-12d <sup>1</sup> F°	
		5-19d <sup>1</sup> P°	5, 6, 10-19d <sup>1</sup> D°	5-7d <sup>1</sup> F°	

\*For predicted terms in the spectra of the Sn I isoelectronic sequence, see Vol. III, Introduction.

## Sn II

(In 1 sequence; 49 electrons)

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4 4d^{10} 5s^2 5p^2 P_{1/2}^o$  $5p^2 P_{1/2}^o$  118017.9 K

I. P. 14.628 volts

In 1938 McCormick and Sawyer published an extension and revision of the earlier work on Sn II. Shenstone has recently reobserved the spectrum between 600 Å and 2500 Å, revised a number of their term values, carried the analysis further, and confirmed the 1938 value of the limit from a Ritz series calculation. There are approximately 145 classified lines in the interval 899 Å to 7904 Å. The doublet and quartet systems of terms are connected by observed intersystem combinations.

In the table, the energy-level values given to one decimal are from Shenstone, who furnished them in advance of publication for inclusion here. Previously identified levels not confirmed by his new observations because of the limited range, are given without a decimal.

Green and Loring observed the Zeeman effect for 8 Sn II lines. The  $g$ -values in the last column of the table have been derived by the writer from their observed patterns.

Shenstone comments that "the analysis is far from complete since it requires new observations with the hollow cathode, over the whole range of wavelengths". This is indicated by the miscellaneous levels in the table. He notes that those labeled 1°, 2° and 3° have the configuration  $5s 5p(^2P^o)6s$  or  $5s 5p(^2P^o)5d$ .

## REFERENCES

- J. B. Green and R. A. Loring, *Phys. Rev.* **30**, 574 (1927). (C L) (Z E)  
 W. W. McCormick and R. A. Sawyer, *Phys. Rev.* **54**, 71 (1938). (I P) (T) (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, 44 (1953). (Summary hfs)  
 W. R. S. Garton, *Proc. Phys. Soc. London [A]* **67**, 864 (1954). (T) (C L)  
 A. G. Shenstone, unpublished material (June 1955). (I P) (T) (C L)

## Sn II

## Sn II

Config.	Desig.	$J$	Level	Interval	Config.	Desig.	$J$	Level	Interval
$5s^2(^1S)5p$	$5p^2 ^1P^o$	0½	0.0	4251.4	$5s 5p^2$	$5p^2 ^2S ?$	0½	80206.1	1262.7
		1½	4251.4				1½	80455.3	
$5s 5p^2$	$5p^2 ^1P$	0½	46464.2	1903.8	$5s 5p^2$	$5p^2 ^2P ?$	1½	81718.0	1262.7
		1½	48368.0	2362.0			0½	86280.2	
		2½	50730.0				1½	89886	
$5s^2(^1S)6s$	$6s ^2S$	0½	56885.9		$5s^2(^1S)4f$	$4f ^2F^o$	3½	89886	-6
$5s 5p^2?$	$5p^2 ^2D$	1½	58843.8	619.6	$5s^2(^1S)6d$	$6d ^2D$	2½	89886	111.4
		2½	59463.4				1½	90242.1	
$5s^2(^1S)5d?$	$5d ^2D$	1½	71405.6	642.0	$5s^2(^1S)7p$	$7p ^2P^o$	2½	90353.5	363
		2½	72047.6				0½	91905	
$5s^2(^1S)6p$	$6p ^2P^o$	0½	71494.3	883.0			1½	92266	
		1½	72377.3						

## Sn II—Continued

## Sn II—Continued

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
5s <sup>2</sup> ( <sup>1</sup> S)8s	8s <sup>1</sup> S	0½	98492. 1		5s <sup>2</sup> ( <sup>1</sup> S)9g	9g <sup>1</sup> G	2½ 4½	112670	
5s <sup>2</sup> ( <sup>1</sup> S)8f	8f <sup>1</sup> F°	2½ 3½	98659 98663	-4	5s <sup>2</sup> ( <sup>1</sup> S)11d	11d <sup>1</sup> D	1½ 2½		112697. 8
5s <sup>2</sup> ( <sup>1</sup> S)7d	7d <sup>1</sup> D	1½ 2½	100284. 6 100339. 2	54. 6	5s <sup>2</sup> ( <sup>1</sup> S)10g	10g <sup>1</sup> G	2½ 4½	113607	
5s <sup>2</sup> ( <sup>1</sup> S)8p	8p <sup>1</sup> P°	0½ 1½	101198 101387	191		2° ( <sup>1</sup> P°?)	1½?		113819. 0
5s <sup>2</sup> ( <sup>1</sup> S)9s	9s <sup>1</sup> S	0½	104689. 1		5s <sup>2</sup> ( <sup>1</sup> S)11g	11g <sup>1</sup> G	2½ 4½	114374	
5s <sup>2</sup> ( <sup>1</sup> S)6f	6f <sup>1</sup> F°	2½ 3½	105387						
5s <sup>2</sup> ( <sup>1</sup> S)8d	8d <sup>1</sup> D	1½ 2½	105732. 6 105764. 8	31. 7					
5s <sup>2</sup> ( <sup>1</sup> S)6g	6g <sup>1</sup> G	3½ 4½	105735		Sn III( <sup>1</sup> S <sub>0</sub> )	Limit		119917. 6	
5s <sup>2</sup> ( <sup>1</sup> S)9p	9p <sup>1</sup> P°	0½ 1½		106368			3°	{ 0½ or 1½	119980. 9?
5s <sup>2</sup> ( <sup>1</sup> S)10s	10s <sup>1</sup> S	0½	108354. 1			4°	{ 0½ or 1½	120063. 8	
5s <sup>2</sup> ( <sup>1</sup> S)7g	7g <sup>1</sup> G	3½ 4½	109000			5°	{ 1½ or 2½	120253. 6	
5s <sup>2</sup> ( <sup>1</sup> S)9d	9d <sup>1</sup> D	1½ 2½		109007. 3 109025. 9	18. 6		6° ( <sup>1</sup> D°?)	3½?	122491. 6
	1° ( <sup>1</sup> P°?)	0½?	109223. 4?		5s 5p( <sup>1</sup> P°)5d	7° ( <sup>1</sup> P°?)	1½?	123156. 9	
5s 5p( <sup>1</sup> P°)6s	6s' <sup>1</sup> P°	0½ 1½ 2½	109455. 5 110772. 8 114345. 7	1222. 7 2467. 5	5s 5p( <sup>1</sup> P°)5d	8° ( <sup>1</sup> P°?)	2½?	124246. 4?	
5s <sup>2</sup> ( <sup>1</sup> S)11s	11s <sup>1</sup> S	0½	110695. 2		5s 5p( <sup>1</sup> P°)5d	9° ( <sup>1</sup> D°?)	1½?	124687. 7	
5s <sup>2</sup> ( <sup>1</sup> S)8g	8g <sup>1</sup> G	3½ 4½	111118		5s 5p( <sup>1</sup> P°)5d	10° ( <sup>1</sup> D°?)	1½?	128168. 0	
5s <sup>2</sup> ( <sup>1</sup> S)10d	10d <sup>1</sup> D	1½ 2½		111128. 8 111141. 3	12. 5	5s 5p( <sup>1</sup> P°)5d	11°	{ 1½ or 2½	128708. 3

July 1955.

## Sn II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> +	Observed Terms		
5s 5p <sup>2</sup>	{ 5p <sup>2</sup> <sup>1</sup> S	5p <sup>2</sup> <sup>3</sup> P 5p <sup>2</sup> <sup>1</sup> P	5p <sup>2</sup> <sup>1</sup> D
	ns (n ≥ 6)	np (n ≥ 5)	nd (n ≥ 5)
5s <sup>2</sup> ( <sup>1</sup> S)ns 5s 5p( <sup>1</sup> P°)ns'	6-11s <sup>1</sup> S	5-9p <sup>1</sup> P°	5-11d <sup>1</sup> D
	6s' <sup>1</sup> P°		
	nf (n ≥ 4)	ng (n ≥ 5)	
5s <sup>2</sup> ( <sup>1</sup> S)ns	4-6f <sup>1</sup> F°	6-11g <sup>1</sup> G	

\*For predicted terms in the spectra of the In I isoelectronic sequence, see Vol. III, Introduction.

## Sn III

(Cd I sequence; 48 electrons)

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 \text{ } ^1S_0$  $5s^2 \text{ } ^1S_0$ , 246020.0 K

I. P. 30.49 volts

Shenstone has revised and extended the early analysis by Gibbs and Vieweg and others, and furnished the tabular data in advance of publication. There are now 125 classified lines between 614 Å and 3963 Å in his line list based on new observations of Sn III. He has derived the limit from the  $\alpha s \text{ } ^1S$  series by means of a Ritz formula. Observed intersystem combinations connect the singlet and triplet systems of terms.

## REFERENCE

A. G. Shenstone, unpublished material (June 1955). (I P) (T) (C L)

## Sn III

## Sn III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$5s^2$	$5s^2 \text{ } ^1S$	0	0.0						
$5s \text{ } ^3S$	$5p \text{ } ^3P^o$	0	53548.0	1648.4	$5s \text{ } ^3S$	$6d \text{ } ^3D$	1	187915.7	80.6
		1	55190.4	4032.2			2	187996.3	132.7
		2	59228.6				3	188129.0	
$5s \text{ } ^1S$	$5p \text{ } ^1P^o$	1	79911.3		$5s \text{ } ^1S$	$6d \text{ } ^1D$	2	189689.9	
$5p^2$	$5p^2 \text{ } ^3P$	0	127308.5	2811.1	$5s \text{ } ^3S$	$5g \text{ } ^3G$	3	206110.3	0.0
		1	130119.6	4447.6			4	206110.3	23.0
		2	134567.2				5	206133.3	
$5p^2$	$5p^2 \text{ } ^1D$	2	128204.7		$5s \text{ } ^1S$	$5g \text{ } ^1G$	4	206125.7	
$5s \text{ } ^3S$	$6s \text{ } ^3S$	1	139637.9		$5s \text{ } ^3S$	$8s \text{ } ^3S$	1	207956.9	
$5s \text{ } ^1S$	$5d \text{ } ^3D$	1	141321.5	204.9	$5s \text{ } ^1S$	$7d \text{ } ^3D$	1	208647.4	47.5
		2	141526.4	311.7			2	208694.9	80.1
		3	141838.1				3	208775.0	
$5s \text{ } ^3S$	$6s \text{ } ^1S$	0	143590.8		$5p \text{ } ^3P^o$	$5d$	2	208731.6	
$5s \text{ } ^1S$	$5d \text{ } ^1D$	2	154115.5		$5s \text{ } ^3S$	$7d \text{ } ^1D$	2	209255.8?	
$5s \text{ } ^3S$	$6p \text{ } ^3P^o$	0	159959.8	275.6	$5p \text{ } ^3P^o$	$5d$	2 or 3	213840.7?	
		1	160815.4	1223.0	$5p \text{ } ^1P^o$	$5d$	1	215613.0?	
		2	161438.4		$5p \text{ } ^3P^o$	$5d$	1 or 2	215856.4	
$5s \text{ } ^1S$	$6p \text{ } ^1P^o$	1	162725.1		$5p \text{ } ^3P^o$	$5d$	0		
$5s \text{ } ^3S$	$4f \text{ } ^3F^o$	2	179308.8	36.2			1	216752.0	867.4
		3	179343.0	98.1			2	217619.4	
		4	179441.1		$5s \text{ } ^1S$	$9s \text{ } ^3S$	1	219504.3?	
$5s \text{ } ^1S$	$4f \text{ } ^1F^o$	3	179702.2						
$5s \text{ } ^3S$	$7s \text{ } ^3S$	1	186688.3						
$5s \text{ } ^1S$	$7s \text{ } ^1S$	0	187399.5?						
					-----	-----	-----	-----	-----
					Sn IV ( $^3S_{01}$ )	Limit	-----	246020.0	

July 1955.

## Sm IV

(Ag I sequence; 47 electrons)

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 S_{1/2}$  $5s^2 S_{1/2}$  328550.0 K

I. F. 40.72 volts

The analysis is by Shenstone, who has revised and extended the early work of Lang and others especially for inclusion here. He has reobserved the spectrum and has a total of 45 classified lines between 595 Å and 4216 Å. His discovery of the  $4d^9 5s^2 {}^2D$  term is of special interest.

His value of the limit is based on a Ritz series calculation for the  $ns^2 S$  series.

## REFERENCE

A. G. Shenstone, unpublished material (June 1955). (I F) (T) (C L)

## Sm IV

## Sm IV

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^{10}(S)5s$	$5s^2 S$	$0\frac{1}{2}$	0.0		$4d^{10}(S)7s$	$7s^2 S$	$0\frac{1}{2}$	237615.7	
$4d^{10}(S)5p$	$5p^2 P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	69663.9 70072.3	6508.4	$4d^{10}(S)5g$	$5g^2 G$	$4\frac{1}{2}$ $3\frac{1}{2}$	258282.3 258282.7	-0.4
$4d^{10}(S)5d$	$5d^2 D$	$1\frac{1}{2}$ $2\frac{1}{2}$	165304.7 165410.8	106.1	$4d^9 5s({}^1D)5p$	$5p'' {}^2F^o?$	$2\frac{1}{2}$ $3\frac{1}{2}$	259876.3	
$4d^9 5s^2$	$5s^2 {}^2D$	$2\frac{1}{2}$ $1\frac{1}{2}$	169233.6 177889.07	-8655.4	$4d^9 5s({}^1D)5p$	$5p'' {}^2D^o?$	$1\frac{1}{2}$ $2\frac{1}{2}$	260398.5	
$4d^{10}(S)6s$	$6s^2 S$	$0\frac{1}{2}$	174133.8		$4d^9 5s({}^1D)5p$	$5p'' {}^2P^o?$	$0\frac{1}{2}$ $1\frac{1}{2}$	263718.5	
$4d^{10}(S)6p$	$6p^2 P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	197850.9 200030.3	2179.9					
$4d^{10}(S)4f$	$4f^2 F^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	210257.7 210313.3	-60.5					
$4d^{10}(S)6d$	$6d^2 D$	$1\frac{1}{2}$ $2\frac{1}{2}$	224795.7 235127.7	332.0					
					$Sn \nu({}^1S_0)$	Limit		328550	

July 1955.

(Pd 1 sequence; 46 electrons)

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4 4d^{10} 5s_0$  $4d^{10} 5s_0$ , 583000 K

I. P. 72.3 volts

The analysis is by Shenstone, who has revised and extended the earlier work by Gibbs and White, and others. The new observations extend from 824A to 1535A and include 58 classified lines.

Shenstone has determined the limit from the  $ns$ -series by means of a Ritz formula with an assumed value of  $\alpha=0.312 \times 10^{-6}$ .

Observed intersystem combinations connect the singlet and triplet systems of terms.

## REFERENCES

- R. C. Gibbs and H. E. White, Proc. Nat. Acad. Sci. 14, 245, 550 (1928). (T) (C L)  
 A. G. Shenstone, unpublished material (June 1955). (I P) (T) (C L)

Sn v					Sn v				
Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
$4d^{10}$	$4d^{10} 5s_0$	0	0.0		$4d^9(^3D)5p$	$5p \ ^1F^0$	3	879844.7	
$4d^9(^3D)5s$	$5s \ ^3D$	3	182578.8	-2479.2	$4d^9(^3D)5p$	$5p \ ^1D^0$	2	888018.7	
		2	185058.0		$4d^9(^3D)5d$	1	2 or 3	874811.6?	
		1	191202.0	-6144.0	$4d^9(^3D)5d$	2	1 or 2	379050.0?	
$4d^9(^3D)5s$	$5s \ ^1D$	2	193229.8		$4d^9(^3D)6s$	$6s \ ^3D$	3	380266.5	-1136.9
$4d^9(^3D)5p$	$5p \ ^3P^0$	2	258372.1	-8623.9		2		381403.4	-7881.6
		1	267896.0			1		389285.0	
		0			$4d^9(^3D)5d$	3	3 or 4	381226.3	
$4d^9(^3D)5p$	$5p \ ^1F^0$	4	268731.1	6414.7	$4d^9(^3D)5d$	4	3 or 4	386584.7	
		3	262316.4		$4d^9(^3D)6s$	$6s \ ^1D$	2	390233.4?	
		2	273336.0	-11018.6					
$4d^9(^3D)5p$	$5p \ ^3D^0$	3	274576.8	5280.8					
		2	289096.0						
		1	281169.1	-12073.1					
$4d^9(^3D)5p$	$5p \ ^1P^0$	1	276589.4		Sn vi( $^3D_{3/2}$ )	Limit		583000	

July 1955.



## Sn VI

(Rh I sequence; 45 electrons)

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^9 \ ^1D_{3/2}$  $4d^9 \ ^1D_{3/2}$ 

K

I. P.

volts

The spectrum is very incompletely analyzed. The listed terms are from Green, who has classified 38 lines between 289.21 Å and 326.43 Å, from a study of the isoelectronic sequence. Observed combinations connect the doublet and quartet systems of terms. No series have been found in Sn VI. The spectrum should be reobserved.

## REFERENCE

M. Green, Phys. Rev. **66**, 117 (1941). (T) (C L)

## Sn VI

## Sn VI

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^9$	$4d^9 \ ^1D$	$2\frac{1}{2}$ $1\frac{1}{2}$	0 8715	-8715	$4d^9(^1D)5p$	$5p'' \ ^1F^{\circ}$	$2\frac{1}{2}$ $3\frac{1}{2}$	331356 333898	2512
$4d^9(^3F)5p$	$5p \ ^3G^{\circ}$	$5\frac{1}{2}$ $4\frac{1}{2}$ $3\frac{1}{2}$ $3\frac{1}{2}$	315000		$4d^9(^1D)5p$	$5p'' \ ^3P^{\circ}$	$0\frac{1}{2}$ $1\frac{1}{2}$	334183 334348	165
$4d^9(^3F)5p$	$5p \ ^3F^{\circ}$	$4\frac{1}{2}$ $3\frac{1}{2}$ $2\frac{1}{2}$ $1\frac{1}{2}$	319857 324885 317581	-5068 7344	$4d^9(^1D)5p$	$5p'' \ ^1D^{\circ}$	$2\frac{1}{2}$ $1\frac{1}{2}$	335638 336590	-892
$4d^9(^3F)5p$	$5p \ ^1D^{\circ}$	$2\frac{1}{2}$ $1\frac{1}{2}$	320718		$4d^9(^3P)5p$	$5p' \ ^1D^{\circ}$	$3\frac{1}{2}$ $2\frac{1}{2}$ $1\frac{1}{2}$ $0\frac{1}{2}$	336678 339859 339443	-3181 416
$4d^9(^3F)5p$	$5p \ ^3F^{\circ}$	$3\frac{1}{2}$ $2\frac{1}{2}$	325277 326712	-1435	$4d^9(^3P)5p$	$5p' \ ^3D^{\circ}$	$1\frac{1}{2}$ $2\frac{1}{2}$	341789 342861	1132
$4d^9(^3F)5p$	$5p \ ^3G^{\circ}$	$4\frac{1}{2}$ $3\frac{1}{2}$	327322		$4d^9(^3P)5p$	$5p' \ ^3P^{\circ}$	$1\frac{1}{2}$ $0\frac{1}{2}$	344400 350245	-5845
$4d^9(^3P)5p$	$5p' \ ^3P^{\circ}$	$2\frac{1}{2}$ $1\frac{1}{2}$ $0\frac{1}{2}$	329131 330098 328907	-967 1291	$4d^9(^3P)5p$	$5p' \ ^3S^{\circ}$	$0\frac{1}{2}$	344883	
					$4d^9(^1G)5p$	$5p''' \ ^1F^{\circ}$	$3\frac{1}{2}$ $2\frac{1}{2}$	345770 347021	-1251

February 1956.

## Sn VI OBSERVED TERMS\*

Configuration $1s^2 2s^2 2p^6 3s^2$ $3p^6 3d^{10} 4s^2 4p^6 +$	Observed Terms
$4d^9$	$4d^9 \ ^1D$
	$np \ (n \geq 5)$
$4d^9(^3F)ns$	$\left\{ \begin{array}{l} 5p \ ^3D^{\circ} \quad 5p \ ^3F^{\circ} \quad 5p \ ^3G^{\circ} \\ 5p \ ^3F^{\circ} \quad 5p \ ^3G^{\circ} \end{array} \right.$
$4d^9(^3P)ns'$	$\left\{ \begin{array}{l} 5p' \ ^3S^{\circ} \quad 5p' \ ^3P^{\circ} \quad 5p' \ ^3D^{\circ} \\ 5p' \ ^3P^{\circ} \quad 5p' \ ^3D^{\circ} \end{array} \right.$
$4d^9(^1D)ns''$	$5p'' \ ^1P^{\circ} \quad 5p'' \ ^3D^{\circ} \quad 5p'' \ ^1F^{\circ}$
$4d^9(^1G)ns'''$	$5p''' \ ^1F^{\circ}$

\*For predicted terms in the spectra of the Rh I isoelectronic sequence, see Vol. III, Introduction.

## Sn XXIII

(Cu I sequence; 29 electrons)

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 S_{1/2}$  $4s^2 S_{1/2}$ 

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed the leading doublets from the  $4s-5p$  and  $4p-5d$  configurations from Pd xviii to In xxi. In figure 3 of his paper on the spectra of highly-ionized atoms, the lines are indicated on the photograph of vacuum-spark spectra of these elements from 40 A to 80 A. He states that the doublets have been traced in this sequence, as far as Sb xxiii.

By analogy with Cu I the writer has assumed the ground state indicated above.

## REFERENCE

B. Edlén, *Physica* 12, No. 9, 549 (1947).

March 1953.

## Sn XXIV

(Co I sequence; 27 electrons)

Z=50

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^7 D_{3/2}$  $3d^7 D_{3/2}$ 

K

I. P.

volts

This spectrum has not been analyzed, but Edlén has observed two lines due to the transition  $3p^6 3d^7 D-3p^6 3d^{10} P^o$ , in the region between 40 A and 80 A. In figure 4 of his paper on the spectra of highly-ionized atoms, the observed wave numbers are plotted against atomic number for this combination in the Co I-like spectra Pd xx to Sn xxiv. For Sn xxiv the wave numbers are between 1800000 K and 2100000 K.

## REFERENCE

B. Edlén, *Physica* 12, No. 9, 548 (1947).

March 1953.

## ANTIMONY

Sb I

51 electrons

Z=51

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3 \text{ } ^3\text{S}_{1,1}$  $5p^3 \text{ } ^3\text{S}_{1,1}$  69700 K

I. P. 8.639 volts

The analysis is by Meggers and Humphreys, whose line list extends from 1388.91 Å to 12466.75 Å. They observed the region from 2000 Å to 13000 Å. For the short-wave region, 1300 Å to 2000 Å, Shenstone made the spectrograms at Princeton. Of the 466 Sb I lines recorded, seventy-four percent have been classified. The observed  $g$ -values are from Zeeman patterns observed by Löwenthal in 1929. The paper by Meggers and Humphreys includes an extensive history and bibliography covering the earlier work on Sb I.

Observed intersystem combinations connect the doublet and quartet systems of terms. "Since it is not possible to assign definite  $L$ -values" to all of the levels, many have been numbered and listed as miscellaneous. The numbers assigned by the authors are entered in column one of the table.

The limit is from the  $ns \text{ } ^3\text{P}(n=6-9)$  and  $ns \text{ } ^1\text{P}(n=6, 7)$  series, and has been derived by a Ritz formula.

## REFERENCES

- W. F. Meggers and C. J. Humphreys, *J. Research Nat. Bur. Std.* **28**, 463, RP 1464 (1942). (I P) (T) (C L) (Z E)  
 P. F. A. Klippenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955). (Summary hfs)



Sb I—Continued

Sb I—Continued

Authors	Config.	Design.	J	Level	Interval	Obs. $\rho$	Authors	Config.	Design.	J	Level	Interval	Obs. $\rho$
29*			3%	66088.0			55				66045.8		
46			0%	66354.0			31*			0%	66069.9		
47			2%	66536.0			56			1%	66141.5		
48			3%	66743.7			57				66349.0		
49			1%	66838.6				5s <sup>2</sup> 5p <sup>2</sup> (P <sub>1</sub> ) 6s	9s 'P	0%	66503	2506	
30*			2%	66837.1				5s <sup>2</sup> 5p <sup>2</sup> (P <sub>2</sub> ) 6s		1%	71089		
50			3%	66967.3			58			2%	66991.0		
51			1%	67052.6				Sb II (P <sub>2</sub> )	Limit	---	69769		
52			0%	67437.7			59			0%	66968.3		
53				67794.6			60	5s <sup>2</sup> 5p <sup>2</sup> (D <sub>2</sub> ) 7s	7s 'D	1%	70880.3		
54			1%	67840.3						2%			

February 1955.

Sb I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> +	Observed Terms
5s <sup>2</sup> 5p <sup>2</sup>	{ 5p <sup>2</sup> 'S <sup>o</sup> 5p <sup>2</sup> 'P <sup>o</sup> 5p <sup>2</sup> 'D <sup>o</sup>
	ns (n ≥ 6)
5s <sup>2</sup> 5p <sup>2</sup> (P) ns	{ 6-9s 'P 6,7s 'P
5s <sup>2</sup> 5p <sup>2</sup> (D) ns'	6,7s' 'D
5s <sup>2</sup> 5p <sup>2</sup> (S) ns''	6s'' 'S

\* For predicted terms in the spectra of the Sb I isoelectronic sequence, see Vol. III, Introduction.

## Sb II

(Sn I sequence; 50 electrons)

Z=51

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3 \ ^3P_1$  $5p^3 \ ^3P_1$ , 133337.5 K

I. P. 16.5 volts

This spectrum should be observed over the entire range. A complete analysis based on homogeneous observations is needed. The leading multiplets have been found by Lang and Vestine. Two other papers on analysis, one by Krishnamurty, and one by Murakawa and Suwa extend the work, but the three term lists are discordant, and it is difficult to interpret existing data in such a way as to provide a satisfactory array of energy levels. The writer has prepared a line list from the wavelengths published by Lang and Vestine (691 Å to 7343 Å), and unpublished measurements by W. F. Meggers (1272 Å to 8742 Å). From this list she has made a multiplet array of all published levels, and revised all level values, but the lists are not in good agreement and the tolerance between observed and calculated wave numbers is inexcusably large.

With intensities and intervals as a guide, she has attempted to reallocate the levels whose published designations appear subject to question, into a term array that compares not too unfavorably with that of Sn I and with the second spectra of nitrogen, phosphorus, and arsenic.

Approximately 220 lines are classified. Eleven levels have been rejected as unreal; two odd levels  $68410.8$  ( $J=1$  or  $2$ ), and  $70160.9$  ( $J=1$  or  $2$ ) have also been omitted from the table, but may be real. It should be emphasized that most of the designation assignments of the higher levels are extremely tentative and require confirmation; in fact, some of the levels listed may be spurious.

Yamanouchi has made a theoretical study of 10 levels of Sb II from the list by Murakawa and Suwa. Eight are unchanged in the present list, but for two, namely, the level at 90353.8 K published as  $^3P_1$  and one at 86051.9 K as  $^1P_1$ , the writer has changed the respective designations to  $7p \ ^3D_1$  and  $6p \ ^3P_1$ . These changes are admittedly tentative.

Lang adopts the limit 150000 K derived from two singlet series of two members each,  $6s \ ^1P^{\circ} - np \ ^1D$  and  $6s \ ^1P^{\circ} - np \ ^1S$  ( $n=5, 6$ ). This limit gives an ionization potential of 19 volts. The extrapolated value by Finkelnburg derived from a study of screening constants,  $16.7 \pm 0.5$ , is probably more reliable. The limit quoted above is from the paper by Murakawa and Suwa.

## REFERENCES

- R. J. Lang and E. H. Vestine, *Phys. Rev.* **43**, 233 (1932). (I P) (T) (C L)  
 S. G. Krishnamurty, *Indian J. Phys.* **10**, 83 (1936). (T) (C L)  
 K. Murakawa and S. Suwa, *Reports Inst. Sci. Tech. Tokyo Univ.* **1**, 121 (1947). (I P) (T) (C L)  
 Y. Yamanouchi, *Reports Inst. Sci. Tech. Tokyo Univ.* **3**, 96 (1948). (T)  
 W. Finkelnburg and F. Stern, *Phys. Rev.* **77**, 303 (1950). (I P)

Sb II

Sb II

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
5e <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 'P	0	0.0	3055.0	5e <sup>2</sup> 5p('P <sup>2</sup> )6p	6p 'B	1 or 2	92542.0	
		1	3055.0	2904.0			1	92000.9	
		2	5659.0	2904.0			2	92542.2	
5e <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 'D	2	12799.8		5e <sup>2</sup> 5p('P <sup>2</sup> )6p	6p 'D	2	92542.2	
5e <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 'B	0	29006.0		5e <sup>2</sup> 5p('P <sup>2</sup> )6p	6p 'B	0	92311.1	
5e 5p <sup>2</sup>	5p <sup>2</sup> 'D°	1	69291.7	210.9	5e <sup>2</sup> 5p('P <sup>2</sup> )7p	7p 'D	2	95464.1	
		2	69292.8	1285.6	5e <sup>2</sup> 5p('P <sup>2</sup> )7p	7p 'B	0	96910.9	
		3	67322.2		5e <sup>2</sup> 5p('P <sup>2</sup> )7s	7s 'P°	0		
5e <sup>2</sup> 5p('P <sup>2</sup> )6s	6s 'P°	0	69139.9	292.2	5e <sup>2</sup> 5p('P <sup>2</sup> )7s	7s 'P°	1	97939.4	7082.5
		1	69455.1	5739.4	5e <sup>2</sup> 5p('P <sup>2</sup> )7s	7s 'P°	2	104724.9	
		2	75874.5		5e <sup>2</sup> 5p('P <sup>2</sup> )8p	8p 'D	1		
5e <sup>2</sup> 5p('P <sup>2</sup> )5d	5d 'D°	1			5e <sup>2</sup> 5p('P <sup>2</sup> )8p	8p 'D	2	105262.2	
		2	79339.9		5e <sup>2</sup> 5p('P <sup>2</sup> )8p	8p 'D	3		
		3			5e <sup>2</sup> 5p('P <sup>2</sup> )4f	4f 'F	2	105591.1	-120.9
5e <sup>2</sup> 5p('P <sup>2</sup> )6s	6s 'P°	1	75899.9		5e <sup>2</sup> 5p('P <sup>2</sup> )4f	4f 'F	3	105470.2	
		2	76992.1		5e <sup>2</sup> 5p('P <sup>2</sup> )4f	4f 'F	4		
		3	77159.4	-447.3	5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'F°	2	106915.2	
5e 5p <sup>2</sup>	5p <sup>2</sup> 'P°	1			5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'F°	3		
		0			5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'F°	4		
		2	78391.2		5e <sup>2</sup> 5p('P <sup>2</sup> )7s	7s 'P°	1	107463.3	
5e 5p <sup>2</sup>	5p <sup>2</sup> 'D°	1 or 2	82221.8		5e <sup>2</sup> 5p('P <sup>2</sup> )8p	8p 'D	2	111266.2	
		1	83825.8	2257.8	5e <sup>2</sup> 5p('P <sup>2</sup> )8p	8p 'D	1 or 2	111679.6	
		2	86182.6	5399.6	5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'D°	1		
5e <sup>2</sup> 5p('P <sup>2</sup> )6p	6p 'D	3	91552.2		5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'D°	2	111684.7	
		1	84316.9		5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'D°	3		
		0	86494.5	-442.6	5e 5p <sup>2</sup>	5p <sup>2</sup> 'B°	1	111671.1	
5e <sup>2</sup> 5p('P <sup>2</sup> )6p	6p 'P	1	86051.9	5666.3	5e <sup>2</sup> 5p('P <sup>2</sup> )6s	6s 'P°	0		
		2	91718.2		5e <sup>2</sup> 5p('P <sup>2</sup> )6s	6s 'P°	1	112356.9	
		1	87831.0		5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'P°	0		
5e <sup>2</sup> 5p('P <sup>2</sup> )6p	6p 'P?	1 or 2	89562.6		5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'P°	1	112404.1	
		2	89626.2		5e <sup>2</sup> 5p('P <sup>2</sup> )6d	6d 'P°	2		
		1	90322.2		5e <sup>2</sup> 5p('P <sup>2</sup> )8s	8s 'P°	1	112753.0	
5e <sup>2</sup> 5p('P <sup>2</sup> )5d	5d 'P°	1	90322.2		5e 5p <sup>2</sup>	5p <sup>2</sup> 'P°	1	113904.5	
5e <sup>2</sup> 5p('P <sup>2</sup> )7p	7p 'D	1	90352.8	292.5	5e <sup>2</sup> 5p('P <sup>2</sup> )7d	7d 'P°	1	122517.0?	
		2	90647.2						
		3							
5e <sup>2</sup> 5p('P <sup>2</sup> )7p	7p 'P	1	90608.4						
		0							
		1	90896.9						
5e <sup>2</sup> 5p('P <sup>2</sup> )7p	7p 'P	2							
		1							
		2							
5e <sup>2</sup> 5p('P <sup>2</sup> )5d	5d 'P°	0	91390.2		Sb III('P <sup>2</sup> )	Limit		122327.5	
		1							
		2							

## Sb III

(In I sequence; 49 electrons)

Z=51

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3 \text{ } ^3\text{P}_{0,1}$  $5p \text{ } ^3\text{P}_{0,1}$ , 294248 K

I. P. 25.3 volts

In 1930 Lang published a list of empirical term values of Sb III, based on an absolute value of  $4f \text{ } ^3\text{F}^{\circ} = 64000$  K, arbitrarily chosen by comparison with the run of the terms in the In I and Sn II spectra. In 1932 he and Vestine investigated the Sb II spectrum and in the course of the work re-examined the Sb III lines. As a result they rejected 5 of the previously published Sb III terms.

In 1947 Murakawa and Suwa revised and extended the earlier analysis, from observations made with a condensed hollow-cathode discharge in a neon atmosphere, as the source. They retained two levels rejected by Lang and Vestine, namely, the 'P levels 54365 K and 57960 K. Nine terms are common to the two lists; and the two levels 92948 K and 93417 K appear with different designation assignments. The limit and tabular data quoted here are from this later paper.

There are 49 classified lines between 691 Å and 6926 Å. The spectrum needs further study.

## REFERENCES

- R. J. Lang, *Phys. Rev.* **35**, 445, 1930. (I P) (T) (C L)  
 R. J. Lang and E. H. Vestine, *Phys. Rev.* **42**, 241 (1932). (T)  
 J. S. Badami, *Nature* **136**, 836 (1935). (hfs)  
 K. Murakawa and S. Suwa, *Reports Inst. Sci. Tech. Tokyo Univ.* **1**, 121 (1947). (I P) (T) (C L)

## Sb III

## Sb III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$5s^2(4s)5p$	$5p \text{ } ^3\text{P}^{\circ}$	$0\frac{1}{2}$ $1\frac{1}{2}$	0 6576	6576	$5s^2(4s)5p$	$6p \text{ } ^3\text{P}^{\circ}$	$0\frac{1}{2}$ $1\frac{1}{2}$	114780.5 116388.5	1668.0
$5s 5p^2$	$5p^2 \text{ } ^4\text{P}$	$0\frac{1}{2}$ $1\frac{1}{2}$ $2\frac{1}{2}$	54365 57960 64356	3595 6396	$5s^2(4s)4f$	$4f \text{ } ^3\text{F}^{\circ}$	$3\frac{1}{2}$ $2\frac{1}{2}$	136815.6 136869.6	-56.0
$5s 5p^2$	$5p^2 \text{ } ^3\text{D}$	$1\frac{1}{2}$ $2\frac{1}{2}$	76522.8 77791.9	1269.1	$5s^2(4s)7s$	$7s \text{ } ^3\text{S}$	$0\frac{1}{2}$	143126.8	
$5s 5p^2$	$5p^2 \text{ } ^3\text{P}$	$0\frac{1}{2}$ $1\frac{1}{2}$	92948.9 101953.5	9004.6	$5s^2(4s)6d$	$6d \text{ } ^3\text{D}$	$1\frac{1}{2}$ $2\frac{1}{2}$	144682.6 144918.6	236.0
$5s^2(4s)6s$	$6s \text{ } ^3\text{S}$	$0\frac{1}{2}$	93417.8		$5s^2(4s)8s$	$8s \text{ } ^3\text{S}$	$0\frac{1}{2}$	163388.2	
$5s^2(4s)5d$	$5d \text{ } ^3\text{D}$	$1\frac{1}{2}$ $2\frac{1}{2}$	96819.5 100385.4	1565.9	$5s^2(4s)5g$	$5g \text{ } ^3\text{G}$	$3\frac{1}{2}, 4\frac{1}{2}$	164302.0	
$5s 5p^2$	$5p^2 \text{ } ^3\text{S}$	$0\frac{1}{2}$	100010?		$5s^2(4s)6g$	$6g \text{ } ^3\text{G}$	$3\frac{1}{2}, 4\frac{1}{2}$	176539.7	
					Sb IV(4s)	Limit		294248	

January 1954.



## Sb IV

(Cd : sequence; 48 electrons)

Z=51

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 1S_0$  $5s^2 1S_0$ , 356156 K

I. P. 44.1 volts

The analysis is seriously incomplete and needs revision from more precise wavelengths. The terms in the table are from Badami, who extended the ultraviolet work by Gibbs and Vieweg and by Green and Lang. He suggested that the configurations of the two low  $^1D$  terms as given below be interchanged. This has not been done since  $5p^2 1D$  is lower than  $5s(^3S)5d 1D$  in related spectra of the sequence. His complete array of combinations contains a total of 74 classified lines between 456 Å and 3922 Å.

Observed intersystem combinations connect the singlet and triplet systems of terms.

The limit is from Gibbs and Vieweg, who report some second series members and list absolute term values that fit an extrapolation along the isoelectronic sequence.

## REFERENCES

- J. B. Green and R. J. Lang, Proc. Nat. Acad. Sci. 14, 706 (1928). (C L)  
 E. C. Gibbs and A. M. Vieweg, Phys. Rev. 34, 400 (1929). (I P) (T) (C L)  
 J. S. Badami, Proc. Phys. Soc. London 43, 538 (1931). (I P) (T) (C L)  
 J. E. Mack, Rev. Mod. Phys. 22, No. 1, 64 (1950). (Summary hfs)

## Sb IV

## Sb IV

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$5s^2$	$5s^2 1S$	0	0		$5s(^3S)6p$	$6p 1P^o$	1	219038	
$5s(^3S)5p$	$5p 1P^o$	0	64435	2265	$5s(^3S)4f$	$4f 1F^o$	2	227069	85
		1	68700	5960			3	227144	160
		2	72560				4	227304	
$5s(^3S)5p$	$5p 1P^o$	1	95952		$5s(^3S)4f$	$4f 1F^o$	3	229598	
$5p^2$	$5p^2 1P$	0	152076	4312	$5s(^3S)6d$	$6d 1D$	1	254683	143
		1	156388	7136			2	254826	253
		2	163524				3	255079	
$5p^2$	$5p^2 1D$	2	155956		$5s(^3S)6d$	$6d 1D$	2	255558	
$5s(^3S)5d$	$5d 1D$	1	178914	350	$5s(^3S)7s$	$7s 1S$	1	257771	
		2	179264	356	$5s(^3S)7s$	$7s 1S$	0	262912	
		3	179620						
$5s(^3S)5d$	$5d 1D$	2	182801						
$5s(^3S)6s$	$6s 1S$	1	188622		Sb v( $^3S_{02}$ )	Limit		356156	
$5s(^3S)6p$	$6p 1P^o$	0	215396	350					
		1	215758	2066					
		2	217808						

August 1953.

Sb IV Observed Terms\*

1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 2s <sup>2</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>2</sup> +	Observed Terms			
5s <sup>2</sup> 5p <sup>2</sup>	5s <sup>2</sup> <sup>1</sup> S			
	{ 5p <sup>2</sup> <sup>3</sup> P 5p <sup>2</sup> <sup>1</sup> D			
	ns (n ≥ 6)	np (n ≥ 5)	nd (n ≥ 5)	n'f (n' ≥ 4)
5s( <sup>2</sup> S)ns	{ 6, 7s <sup>2</sup> S 7s <sup>1</sup> S	5, 6p <sup>3</sup> P <sup>o</sup> 5, 6p <sup>1</sup> P <sup>o</sup>	5, 6d <sup>3</sup> D 5, 6d <sup>1</sup> D	4/ <sup>3</sup> F <sup>o</sup> 4/ <sup>1</sup> F <sup>o</sup>

\*For predicted terms in the spectra of the Cd I isoelectronic sequence, see Vol. III, Introduction.

Sb V

(Ag I sequence; 47 electrons)

Z=51

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s <sup>2</sup>S<sub>1/2</sub>

5s <sup>2</sup>S<sub>1/2</sub> 449300 K

I. P. 56 volts

This spectrum is very incompletely known. The analysis is from Badami, who has revised the early work of Lang on the basis of observations by Gibbs, Vieweg, and Gartlein, added 6p <sup>3</sup>P<sup>o</sup>, and rejected Lang's 5f <sup>3</sup>F<sup>o</sup> term. There are 12 classified lines in the range 699.22 Å to 3362.94 Å.

The limit is from Lang, who derived absolute term values "by treating the f-orbits as hydrogenic — and obtaining thus an estimate of the F term value." His terms fit a Moseley diagram of the isoelectronic sequence Ag I to Sb V. This limit may be considerably in error. Further observations are needed to extend the series.

REFERENCES

- R. J. Lang, Proc. Nat. Acad. Sci. 13, 341 (1927). (I P) (T) (C L)  
 R. C. Gibbs, A. M. Vieweg, and C. W. Gartlein, Phys. Rev. 34, 406 (1929).  
 J. S. Badami, Proc. Phys. Soc. (London) 42, 541 (1931). (T) (C L)

Sb V

Sb V

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
4d <sup>10</sup> ( <sup>1</sup> S)5s	5s <sup>2</sup> S	0½	0		4d <sup>10</sup> ( <sup>1</sup> S)6s	6s <sup>2</sup> S	0½	224653	3199
4d <sup>10</sup> ( <sup>1</sup> S)5p	5p <sup>3</sup> P <sup>o</sup>	0½ 1½	81566 90553	8987	4d <sup>10</sup> ( <sup>1</sup> S)6p	6p <sup>3</sup> P <sup>o</sup>	0½ 1½	254380 257579	
4d <sup>10</sup> ( <sup>1</sup> S)5d	5d <sup>3</sup> D	1½ 2½	201927 203043	1116	Sb VI( <sup>1</sup> S <sub>0</sub> )	Limit	-----	449300	

March 1953.

## Sb VI

(Pd I sequence; 46 electrons)

Z=51

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} {}^1S_0$  $4d^{10} {}^1S_0$ , 868140 K

I. P. 108 volts

Schoepfle classified 31 lines in the interval 883.45 Å to 1331.59 Å, but did not observe the resonance lines. The lines were classified from a study of a Moseley diagram of the sequence. Later, Kruger and Shoupp reported three lines in the region between 279 Å and 291 Å as combinations from the ground term, and revised Schoepfle's value of  $5p {}^1P^0$ . They give the limit quoted above and have also based their work on a study of a Moseley diagram. The quoted ionization potential may be in error by several volts. The tabular entries are from the earlier paper supplemented by these later additions and corrections. The writer has rounded off the values, and entered the  $J$ -value of the parent term by analogy with Pd I.

The singlet and triplet systems of terms are connected by observed intersystem combinations.

The level values of these terms published by L. and E. Bloch in 1937 derived from their own measurements differ considerably from those in the table, but there is agreement on the resonance lines at 284 Å and 279 Å representing combinations of the ground term with  $5p {}^1P^0$  and  $5p {}^1D_1^0$ , respectively.

## REFERENCES

- G. K. Schoepfle, *Phys. Rev.* **43**, 742 (1933). (I P) (T) (C L)  
 P. G. Kruger and W. E. Shoupp, *Phys. Rev.* **66**, 124 (1934). (I P) (T) (C L)  
 L. Bloch et E. Bloch, *J. Phys. Rad.* [7] **8**, 224 (1937). (T) (C L)

## Sb VI

## Sb VI

Config.	Desig.	$J$	Level	Interval	Config.	Desig.	$J$	Level	Interval
$4d^{10}$	$4d^{10} {}^1S$	0	0		$4d^9({}^2D_{3/2})5p$	$5p {}^1D^0$	3	348570	6770
$4d^9({}^2D_{3/2})5s$	$5s {}^1D$	2	243220	-2770			2	348300	-14630
$4d^9({}^2D_{1/2})5s$		3	245990	-7220	$4d^9({}^2D_{1/2})5p$		1	357480	
$4d^9({}^2D_{3/2})5s$		1	253210		$4d^9({}^2D_{3/2})5p$	$5p {}^1P^0$	1	351010	
$4d^9({}^2D_{1/2})5s$	$5s {}^1D$	2	256360		$4d^9({}^2D_{1/2})5p$	$5p {}^1F^0$	3	355300	
$4d^9({}^2D_{3/2})5p$	$5p {}^1P^0$	2	351460	-11070	$4d^9({}^2D_{3/2})5p$	$5p {}^1D^0$	2	359170	
$4d^9({}^2D_{1/2})5p$		1	348530	-6270					
$4d^9({}^2D_{3/2})5p$		0	348300						
$4d^9({}^2D_{1/2})5p$	$5p {}^1F^0$	4	348260	8320	Sb VII( ${}^2D_{3/2}$ )	Limit		[868140]	
$4d^9({}^2D_{3/2})5p$		3	354930	-3230					
$4d^9({}^2D_{1/2})5p$		2	348160						

March 1953.

## Sb XXIII

(Cu I sequence; 29 electrons)

Z=51

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s {}^2S_{1/2}$  $4s {}^2S_{1/2}$  K

I. P. volts

This spectrum has not been analyzed, but Edlén has observed the leading doublets from the  $4s-5p$  and  $4p-5d$  configurations from Pd XVIII to In XXI. In figure 3 of his paper on the spectra of highly-ionized atoms, the lines are indicated on the photograph of vacuum-spark spectra of these elements from 40 Å to 80 Å. He states that the doublets have been traced in this sequence as far as Sb XXIII.

By analogy with Cu I the writer has assumed the ground state indicated above.

## REFERENCE

- B. Edlén, *Physica* **13**, No. 9, 548 (1947).

March 1953

## TELLURIUM

Te I

52 electrons

Z=52

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4 4d^{10} 5s^2 5p^4 \ ^3P_1$

$5p^4 \ ^3P_1$ , 72667 K

I. P. 9.01 volts

The analysis is from Ruedy, who has classified 62 lines between 5083.0A and 11084.5A from observations he made by using an electrodeless discharge tube as the source, thus revising and extending the early work by McLennan and others.

The writer has tentatively assigned  $J$ -values to the members of the  $7p \ ^3P$  term on the basis of the intensities of the combination with  $6s \ ^4S_1^o$ . The  $^4D^o$  terms are listed in the table as unresolved, although the author assigns  $J=2$  to the observed levels of this series.

In 1929, McLennan and Crawford predicted the positions of the two terms  $5p^4 \ ^1D_2$  and  $5p^4 \ ^1S_0$  by extrapolation from the first spectra of the oxygen group. These "forbidden" transitions have since been confirmed by laboratory observations, and are described in detail in the 1938 reference below. They are attributed to magnetic dipole and electric quadrupole radiation. The predicted positions of these terms are quoted in the table.

Bartelt revised and extended Ruedy's analysis in 1934, on the basis of observations by McLennan and his associates, and listed 42 classified lines between 1645A and 3175.15A. Further confirmation of his results based on more precise observations is needed.

Ruedy has derived the limit from the  $6p \ ^4P_1 - ns \ ^4S_1^o$  series ( $n=9, 10, 11$ ), by means of a Hicks formula.

The Zeeman effect in the spectra of tellurium is being investigated by Vander Sluis and Griffin at the Oak Ridge National Laboratory. Their extension of the Te I analysis is in progress but will not be completed in time for inclusion here. This spectrum is in serious need of further study, particularly in the short wave region.

### REFERENCES

- J. C. McLennan and M. F. Crawford, *Nature* **124**, 874 (1929). (T)  
J. E. Ruedy, *Phys. Rev.* **41**, 588 (1932). (I P) (T) (C L)  
O. Bartelt, *Zeit. Phys.* **88**, 522 (1934). (T) (C L)  
H. Niewodniczanski and F. Lipifski, *Nature* **142**, 1160 (1938). (C L)  
P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
J. B. Green and R. A. Loring, *Phys. Rev.* **90**, 80 (1953). (Z E)  
H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469 Suppl. II, 29 (1955). (Summary hfs).  
P. M. Griffin, letter (January 1957).

Te I

Te I

Config.	Desig.	J	Level	Interval	Obs. $g$	Config.	Desig.	J	Level	Interval	Obs. $g$		
$5p^4$	$5p^4 \ ^3P$	2	0	-4751 44	1.49	$5p^3(4S^0)7d$	$7d \ ^3D^0$	1 to 3	68914				
		1	4751					$5p^3(4S^0)9e$	$9e \ ^3S^0$	2	68964		
		0	4707										
$5p^4$	$5p^4 \ ^1D$	2	10559			$5p^3(4S^0)8d$	$8d \ ^1D^0$	0 to 4	69145				
$5p^4$	$5p^4 \ ^3S$	0	23199			$5p^3(4S^0)8d$	$8d \ ^3D^0$	1 to 3	69577				
$5p^3(4S^0)6e$	$6e \ ^3S^0$	2	44853		1.98	$5p^3(4S^0)10e$	$10e \ ^3S^0$	2	69670				
$5p^3(4S^0)6e$	$6e \ ^3P^0$	1	46653		1.90	$5p^3(4S^0)9d$	$9d \ ^3D^0$	0 to 4	70137				
$5p^3(4S^0)6p$	$6p \ ^3P$	1	54162	39 336		$5p^3(4S^0)9d$	$9d \ ^3D^0$	1 to 3	70398				
		2	54201					$5p^3(4S^0)11e$	$11e \ ^3S^0$	2	70468		
		3	54537										
$5p^3(4S^0)6p$	$6p \ ^3P$	2	55813	138 -171		$5p^3(4S^0)10d$	$10d \ ^3D^0$	0 to 4	70767				
		1	55675					$5p^3(4S^0)10d$	$10d \ ^3D^0$	1 to 3	70930		
		0	55846										
$5p^3(4S^0)7p$	$7p \ ^3P$	1	63557	113 251		$5p^3(4S^0)11d$	$11d \ ^3D^0$	0 to 4	71184				
		2	63670					$5p^3(4S^0)11d$	$11d \ ^3D^0$	1 to 3	71297		
		3	63921										
$5p^3(4S^0)7p$	$7p \ ^3P$	2	63922	-167 107		$5p^3(4S^0)12d$	$12d \ ^3D^0$	1 to 3	71557				
		1	64089					$5p^3(4S^0)13d$	$13d \ ^3D^0$	1 to 3	71749		
		0	63982										
$5p^3(4S^0)6d$	$6d \ ^3D^0$	1 to 3	65695										
$5p^3(4S^0)8e$	$8e \ ^3S^0$	2	65933										
$5p^3(4S^0)7d$	$7d \ ^3D^0$	0 to 4	67437										
						Te II( $4S^0_{1/2}$ )	Limit		72667				

January 1957.

## Te I OBSERVED TERMS\*

Configuration $1s^2 2s^2 2p^6 3s^2$ $3p^2 3d^0 4s^2$ $4p^2 4d^0 5s^2 +$	Observed Terms		
$5p^4$	{ $5p^4 \ ^3S$ $5p^4 \ ^3P$ $5p^4 \ ^1D$		
	$ns (n \geq 6)$	$np (n \geq 6)$	$nd (n \geq 5)$
$5p^3(4S^0)ns$	{ $6, 8$ to $11s \ ^3S^0$ $6e \ ^3S^0$		
	$6, 7p \ ^3P$ $6, 7p \ ^3P$	$7$ to $11d \ ^3D^0$ $6$ to $13d \ ^3D^0$	

\*For predicted terms in the spectra of the Te I isoelectronic sequence, see Vol. III, Introduction.

## Te II

(Sb I sequence; 51 electrons)

Z=52

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3 \text{ } ^4\text{S}_{1/2}$  $5p^3 \text{ } ^4\text{S}_{1/2}$   $156000 \pm 3000$  KI. P.  $18.6 \pm 0.4$  volts

The analysis is by Handrup and Mack, who have revised and extended the earlier work on this spectrum especially for inclusion here. Their investigation is still in progress. It has been carried to its present state only with the aid of the excellent Zeeman effect data furnished by Vander Sluis and Griffin of the Oak Ridge National Laboratory, and of ultraviolet line measurements from plates made by S. Avellón of Lund University. The  $g$ -values quoted here supplement and partly supersede those published in the 1951 reference.

There are 29 odd and 38 even levels listed in the table, of which 4, listed with question marks, need further confirmation. All odd levels are in italics. The observations extend from 157 Å to 8972 Å, and about 300 lines have been classified.

The zeros after the decimal point in the level 71193.00 have been arbitrarily adopted by the authors because of the inaccuracy of the far ultraviolet wave numbers on which the combinations from the ground term are based.

The configurations  $5s 5p^4$  and  $5s^2 5p^2 ({}^1\text{D } {}^1\text{S})6p$  are almost completely known. The authors point out, however, that on account of the effects of configuration-interaction and intermediate coupling, in many cases the assignment of configurations and  $LS$ -designations has little meaning. For example, at least the two levels 76301 and 78448 contribute to  $5p^4 \text{ } ^4\text{P}_{0,2}$ . The limit terms  ${}^3\text{P}$ ,  ${}^1\text{D}$ ,  ${}^1\text{S}$  are, also, not indicated in the table for most levels of the  $5s^2 5p^2 n\ell$  configurations; and a number of configurations are listed for groups of levels. Ross and Murakawa have suggested some approximate quantum number assignments.

No series have been found in Te II, but from the spacing of successive groups of levels Handrup and Mack estimate the series limit quoted above. Their ionization potential agrees well with the value  $18.8 \pm 0.5$  volts interpolated by Finkelburg and Humbach from a study of screening constants.

## REFERENCES

- J. E. Mack, K. Murakawa, J. S. Ross, F. A. Pick, and J. C. van den Bosch, Phys. Rev. **83**, 654 (L) (1951). (T) (Z E) (hfs)
- J. S. Ross and K. Murakawa, Phys. Rev. **85**, 559 (1952). (T) (C L) (hfs) (I S)
- J. B. Green and R. A. Loring, Phys. Rev. **90**, 80 (1953). (C L) (Z E)
- W. Finkelburg und W. Humbach, Naturwiss. **42**, 35 (1955). (I P)
- M. Bernardus Handrup and J. E. Mack, unpublished material (December 1956); See Bull. Am. Phys. Soc. [II] **2**, 199 (A) (1957). (I P) (T) (C L) (Z E)

Te II

Te II

Config.	Design.	J	Level	Interval	Obs. g	Config.	Design.	J	Level	Interval	Obs. g
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 5S°	1½	0					2½	111947. 17		1. 19
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 1D°	1½ 2½	10889 12483	2201		5s <sup>2</sup> 5p <sup>2</sup> (1D) 6p		1½	112272. 87		1. 09
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 1P°	0½ 1½	20548 24033	3487				2½	112549. 29		1. 06
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 1P°	2½	71193. 00	-3700. 85	1. 59	5s <sup>2</sup> 5p <sup>2</sup> 7e 5s <sup>2</sup> 5p <sup>2</sup> 6d		0½	112824. 36		2. 16
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 1P°	1½	74893. 85		1. 71	5s <sup>2</sup> 5p <sup>2</sup> (1D) 6p		0½	114069. 12		0. 77
		0½	76301. 30		2. 63	5s <sup>2</sup> 5p <sup>2</sup> 7e 5s <sup>2</sup> 5p <sup>2</sup> 6d		1½	115700. 94		0. 91
		0½	78448. 63		2. 29	5s <sup>2</sup> 5p <sup>2</sup> (1D) 6p		1½	116102. 19		1. 29
		1½	81895. 89		1. 12			2½	116837. 76?		1. 20
		1½	82743. 83		1. 45	5s <sup>2</sup> 5p <sup>2</sup> 7e		0½?	117192. 78		
		0½	83577. 90		1. 06	5s <sup>2</sup> 5p <sup>2</sup> 6d		1½	117264. 36		0. 98
		2½	85049. 79		1. 09			1½	117340. 09		1. 60
5s <sup>2</sup> 5p <sup>2</sup>		1½	85160. 15		0. 88			2½	117686. 31		0. 85
5s <sup>2</sup> 5p <sup>2</sup> 6e		2½	85592. 36		1. 50			3½	117859. 52		1. 13
5s <sup>2</sup> 5p <sup>2</sup> 5d		1½	86760. 28		0. 94			0½?	118009. 36		
		2½	87404. 98		1. 25			0½	118325. 30		1. 17
		2½	88925. 17		1. 06?			2½	120617. 57		1. 51
		1½	88961. 52		1. 17	5s <sup>2</sup> 5p <sup>2</sup> 7e		2½	121064. 23		
		1½	92192. 22		1. 13	5s <sup>2</sup> 5p <sup>2</sup> 6d		0½	121106. 68		0. 52
		2½	92793. 80		1. 24			1½	121174. 38		1. 23
5s <sup>2</sup> 5p <sup>2</sup> (1P) 6p		0½	93979. 34		0. 79			1½	121519. 36		1. 20
5s <sup>2</sup> 5p <sup>2</sup>		2½	94860. 95		1. 28			2½	122197. 27		1. 07
5s <sup>2</sup> 5p <sup>2</sup> 6e		1½	95208. 88		0. 98	5s <sup>2</sup> 5p <sup>2</sup> 7p		2½	122616. 98?		1. 06
5s <sup>2</sup> 5p <sup>2</sup> 5d		1½	96145. 17		1. 32	5s <sup>2</sup> 5p <sup>2</sup> 4f		2½	122888. 03		
		0½	97780. 47		1. 26			3½	123655. 00		1. 24
		1½	99585. 08		1. 26	5s <sup>2</sup> 5p <sup>2</sup> 7e		0½	124082. 49		
		2½	100112. 52		1. 40	5s <sup>2</sup> 5p <sup>2</sup> 6d		2½	124646. 81		1. 32
		1½	101221. 44		1. 31			3½	125967. 61		1. 15
		0½	101371. 26		2. 32	5s <sup>2</sup> 5p <sup>2</sup> 7p 5s <sup>2</sup> 5p <sup>2</sup> 4f		3½	126164. 69?		
		2½	102324. 95		1. 31			0½	126212. 47		2. 03
		3½	103106. 32		1. 38	5s <sup>2</sup> 5p <sup>2</sup> 7e		1½	126517. 39		0. 98
		1½	103956. 32		1. 74	5s <sup>2</sup> 5p <sup>2</sup> 6d					
		1½	105006. 55		1. 21						
		2½	105583. 48		1. 29	Te m(1P <sub>0</sub> )	Limit		150000		
		0½	106119. 73		0. 88						

January 1967.

## Te III

(Sn I sequence; 50 electrons)

Z=52

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3 \text{ } ^3\text{P}_0$  $5p^3 \text{ } ^3\text{P}_0$ , 247000 K

I. P. 31 volts

This spectrum is in serious need of further study. Krishnamurty and Rao have classified approximately 200 lines between 612 Å and 6977 Å. R. J. Lang furnished them with an unpublished list of lines in the region between 600 Å and 2000 Å, to supplement their observations.

The limit is derived from the two-member series  $np \text{ } ^3\text{P}$ , namely,  $6s \text{ } ^3\text{P}_2^o - 5p^3 \text{ } ^3\text{P}$ , and  $6s \text{ } ^3\text{P}_2^o - 6p \text{ } ^3\text{P}_2$ , and "a very large error is possible when the limit is obtained solely from the above series in which the first member lies in the extreme ultra-violet." A rounded-off value is quoted in the table.

Observed intersystem combinations connect the singlet and triplet systems of terms.

The observed  $g$ -values in the table are from the paper by Green and Loring.

The writer has revised several of the published term designations to conform more nearly to the related spectra Sn I and Sb II, and to give better agreement with the observed  $g$ -values. Some  $J$ -values of the limit term have been added in column one of the table, and more levels are entered as miscellaneous than in the published papers. The more dubious levels have been omitted here pending confirmation. The tabular data are tentative, but should serve as a guide in the search for more regularities.

## REFERENCES

- S. G. Krishnamurty, Proc. Roy. Soc. London [A] 151, 178 (1935). (I P) (T) (C L)  
 S. G. Krishnamurty and K. R. Rao, Proc. Roy. Soc. London [A] 158, 562 (1937). (T) (C L)  
 J. B. Green and R. A. Loring, Phys. Rev. 89, 82 (1953). (Z E)

Te III

Te III

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$
$5s^2 5p^3$	$5p^3 \text{ } ^3\text{P}$	0	0			$5s^2 5p(^3\text{P}^o)5d$	$^3\text{F}^o$	2	104708.8		
		1	4751	4751				3	108305.6	1.07	
		2	8165	3414							
$5s^2 5p^3$	$5p^3 \text{ } ^1\text{D}$	2	17358			$5s^2 5p(^3\text{P}_{1/2})6s$	$6s \text{ } ^3\text{P}^o$	0	107482.1	255.6	1.37
					1			107717.7	7698.3	1.28	
$5s 5p^3$		1	88885.7			$5s^2 5p(^3\text{P}_{1/2})6s$		2	115410.0		
					2			109006.1			
$5s 5p^3$		1	92300.9			$5s^2 5p(^3\text{P}_{1/2})6s$	$6s \text{ } ^1\text{P}^o$	1	114209.9		1.00
"	$^3\text{P}^o$	2	95024.6			$5s^2 5p(^3\text{P}^o)5d$	$5d \text{ } ^3\text{P}^o$	0			
"	$^3\text{P}^o$	1	96574.4			$5s^2 5p(^3\text{P}_{1/2})5d$		1	117789.4	-1077.5	1.26
					2			118711.9	1.34		
"	$^1\text{D}^o$	2	100463.4		1.13	$5s^2 5p(^3\text{P}^o)5d$	$5d \text{ } ^1\text{D}^o$	1	115739.4	6770.7	
					2			122510.1	-1613.4	1.35	
		1	101149.0		3			120896.7			
						1 or 2		122121.2			



## Te III—Continued

## Te III—Continued

Config.	Design.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Design.	<i>J</i>	Level	Interval	Obs. <i>g</i>
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 5d	5d 1D°	2	184783.3			5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d	6d 1P°	2	168797.8		
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 5d	5d 1P°	1	187183.6			5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d		3	163796.4	988.6	
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 5d	5d 1F°	3	187841.8			5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d		4	168543.7?	4817.3?	
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p	6p 1D	1	128617.9	3711.2	0.68	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d	6d 1D°	1	163384.9	3286.5	
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p		2	132329.1	7620.6	1.16	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d		2	168611.4	4450.5	
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p		3	139949.7		1.28	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d		3	171061.9		
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p	6p 1P	1	138289.7		1.02	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d	6d 1F°	3	168071.9		
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p	6p 1P	0	132262.4	-145.7		5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 7s	7s 1P°	0	170683.4		
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p		1	132116.7	7547.8	1.24	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 7s		1	178410.9	7827.5	
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p		2	139664.5		1.43	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d		2	171869.4		
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p	6p 3S	1	141803.0		1.76	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d	6d 1D°	2	171888.8		
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6p	6p 1D	2	142982.0		1.14	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d		1 or 2	173569.8		
5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 6d	6d 1P°	0	161198.6	8819.6	1.43	5s <sup>2</sup> 5p(5P <sub>1/2</sub> ) 7s	7s 1P°	1	174497.5		
"		1	170018.2								
		2									
						Te IV(5P <sub>1/2</sub> )	Limit		247000		

October 1953.

## Te IV

(In I sequence; 49 electrons)

Z=52

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s<sup>2</sup> 5p<sup>2</sup> P<sub>0,1/2</sub><sup>o</sup>5p<sup>2</sup> P<sub>0,1/2</sub><sup>o</sup> 305091 K

I. P. 38 volts

The analysis is from Rao, who has classified 27 lines between 749 Å and 3585 Å. An error in the published value of 7s 3S is corrected here. His value of the limit is the mean derived from a Rydberg formula for the 3S and 3D series of two members each. From isoelectronic sequence data he estimates that the error in the absolute term values "probably does not exceed about 3000 cm<sup>-1</sup>". Further study of the spectrum is needed.

The *g*-values are from the observed Zeeman pattern of the line 3585.27 Å—the only Te IV line observed by Green and Loring.

## REFERENCES

- K. R. Rao, Proc. Roy. Soc. London [A] **133**, 220 (1931). (I P) (T) (C L)  
 J. B. Green and R. A. Loring, Phys. Rev. **90**, 80 (1953). (Z E)

## Te IV

Config.	Design.	<i>J</i>	Level	Interval	Obs. <i>g</i>
5s <sup>2</sup> ( <sup>1</sup> S)5p	5p <sup>1</sup> P <sup>o</sup>	0½	0	9222	
		1½	9222		
5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>3</sup> D	1½	92772	2039	
		2½	94811		
5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> S	0½	109536		
5s 5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> P	0½	119009	947	
		1½	119056		
5s <sup>2</sup> ( <sup>1</sup> S)5d	5d <sup>3</sup> D	1½	127445. 0	771. 5	
		2½	128216. 5		
5s <sup>2</sup> ( <sup>1</sup> S)6s	6s <sup>1</sup> S	0½	133456. 0		2. 02
5s <sup>2</sup> ( <sup>1</sup> S)6p	6p <sup>1</sup> P <sup>o</sup>	0½	161359. 4	2619. 6	0. 66
		1½	163959. 0		
5s <sup>2</sup> ( <sup>1</sup> S)7s	7s <sup>1</sup> S	0½	205343. 6		
5s <sup>2</sup> ( <sup>1</sup> S)6d	6d <sup>3</sup> D	1½	202941. 4	411. 4	
		2½	203352. 8		
Te v( <sup>1</sup> S <sub>0</sub> )	Limit		365091		

August 1953.

## Te v

(Cd I sequence; 48 electrons)

Z=52

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s<sup>2</sup> <sup>1</sup>S<sub>0</sub>5s<sup>2</sup> <sup>1</sup>S<sub>0</sub> 486244 K

I. P. 60 volts

The spectrum is incompletely known, but Gibbs and Vieweg have classified 23 lines between 603 Å and 1549 Å. Their absolute term values have been extrapolated on a Moseley diagram of the isoelectronic sequence. Observed intersystem combinations connect the singlet and triplet systems of terms.

L. and E. Bloch have observed five additional lines between 358 Å and 402 Å, from which they derive the three following levels:

6p <sup>3</sup> P <sub>1</sub> <sup>o</sup>	274454
6p <sup>1</sup> P <sub>1</sub> <sup>o</sup>	278738
7s <sup>1</sup> S <sub>1</sub>	334459

These levels require further confirmation.

From a study of screening constants Finkelnburg and Humbach have interpolated an ionization potential of 66 volts.

## REFERENCES

- R. C. Gibbs and A. M. Vieweg, Phys. Rev. **34**, 400 (1929). (I P) (T) (C L)  
 L. et E. Bloch, J. Phys. Rad. [7] **6**, 441 (1935). (T) (C L)  
 W. Finkelnburg und W. Humbach, Naturwiss. **42**, 35 (1955). (I P)

## Te v

Config.	Desig.	<i>J</i>	Level	Interval
5s <sup>2</sup>	5s <sup>2</sup> <sup>1</sup> S	0	0	
5s(5s)5p	5p <sup>3</sup> P <sup>o</sup>	0	75109	2914
		1	79083	7974
		2	85997	
5s(5s)5p	5p <sup>1</sup> P <sup>o</sup>	1	111707	
5p <sup>2</sup>	5p <sup>2</sup> <sup>3</sup> P	0	176248	6171
		1	182419	10173
		2	192592	
5p <sup>2</sup>	5p <sup>2</sup> <sup>1</sup> D	2	182797	
5s(5s)5d	5d <sup>3</sup> D	1	215612	525
		2	216137	852
		3	216989	
5s(5s)5d	5d <sup>1</sup> D	2	221498	
5s(5s)6s	6s <sup>2</sup> S	1	240637	
Te vi(5s <sub>1/2</sub> )	Limit	-----	486244	

August 1955.

## Te vi

(Ag I sequence; 47 electrons)

*Z* = 52Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s <sup>2</sup>S<sub>1/2</sub>5s <sup>2</sup>S<sub>1/2</sub> 583500 K

I. P. 72 volts

This spectrum is incompletely analyzed. The terms are from K. R. Rao, who has classified 10 lines in the range between 540.24 Å and 1313.90 Å. The limit is a mean value of that given by the *ns* <sup>2</sup>S series (*n* = 5, 6) derived by a Rydberg formula, and of the one given by an extrapolation along the isoelectronic sequence based on Moseley's law. The writer has rounded off the value of this mean. Further observations are needed to lengthen the series and correct the present inaccuracy in the limit.

## REFERENCE

K. R. Rao, Proc. Royal Soc. London [A] 123, 220 (1931). (I P) (T) (C L)

## Te vi

## Te vi

Config.	Desig.	<i>J</i>	Level	Interval	Config.	Desig.	<i>J</i>	Level	Interval
4d <sup>10</sup> (5s)5s	5s <sup>2</sup> S	0½	0		4d <sup>10</sup> (5s)6p	6p <sup>3</sup> P <sup>o</sup>	0½ 1½	314190 318589	4399
4d <sup>10</sup> (5s)5p	5p <sup>3</sup> P <sup>o</sup>	0½ 1½	93338 106150	11814					
4d <sup>10</sup> (5s)5d	5d <sup>3</sup> D	1½ 2½	239081 239725	1644	Te vii(5s)	Limit	-----	583500	
4d <sup>10</sup> (5s)6s	6s <sup>2</sup> S	0½	278439						

March 1953.

## Te VII

(Pd I sequence; 46 electrons)

Z=52

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s$ , $4d^{10} 5s$ , 1106300 K

I. P. 137 volts

In 1933 Schoepfle classified 24 lines between 784.09 Å and 1123.36 Å, but did not observe the resonance lines. The following year Kruger and Shoupp reported three lines in the region from 227 Å to 236 Å as combinations from the ground term, and revised the earlier value of  $5p^1P^o$ . Their limit, quoted above, and all the terms are based on a study of a Moseley diagram for the isoelectronic sequence. The quoted ionization potential may be in error by several volts. The terms in the table are from the 1933 paper supplemented by these later additions and corrections. The writer has rounded off the values, and assigned the *J*-value of the parent term by analogy with Pd I.

The singlet and triplet systems of terms are connected by observed intersystem combinations.

The level values of these same terms published by L. and E. Bloch in 1937 as prepared from their own measurements differ considerably from those adopted here, but there is agreement on the two resonance lines at 232 Å and 227 Å representing combinations from the ground term to  $5p^1P^o$  and  $5p^1D^o$ , respectively.

## REFERENCES

- G. K. Schoepfle, *Phys. Rev.* **43**, 742 (1933). (I P) (T) (C L)  
 P. G. Kruger and W. E. Shoupp, *Phys. Rev.* **46**, 124 (1934). (I P) (T) (C L)  
 L. Bloch et E. Bloch, *J. Phys. Rad.* [7] **8**, 224 (1937). (T) (C L)

## Te VII

## Te VII

Config.	Desig.	<i>J</i>	Level	Interval	Config.	Desig.	<i>J</i>	Level	Interval
$4d^{10}$	$4d^{10} 5s$	0	0		$4d^9(^3D_{30})5p$	$5p^1D^o$	3	489910	8300
$4d^9(^3D_{30})5s$	$5s^1D$	3	309000	-3090	$4d^9(^3D_{10})5p$		2	481810	-17350
$4d^9(^3D_{10})5s$		2	312090	-8280	$4d^9(^3D_{10})5p$		1	486900	
$4d^9(^3D_{10})5s$		1	320370		$4d^9(^3D_{10})5p$	$5p^1P^o$	1	490410	
$4d^9(^3D_{10})5s$	$5s^1D$	2	323620		$4d^9(^3D_{10})5p$	$5p^1F^o$	3	496580	
$4d^9(^3D_{30})5p$	$5p^1P^o$	2	409310	-13560	$4d^9(^3D_{10})5p$	$5p^1D^o$	2	440870	
$4d^9(^3D_{10})5p$		1	499570	-7180					
$4d^9(^3D_{10})5p$		0	490060						
$4d^9(^3D_{30})5p$	$5p^1F^o$	4	482940	10300	Te VIII( $^3D_{30}$ )	Limit		1106300	
$4d^9(^3D_{10})5p$		3	419840	-15510					
$4d^9(^3D_{10})5p$		2	488160						

March 1953.

## IODINE

II

53 electrons

Z=53

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^5 {}^2P_{3/2}$  $5p^5 {}^2P_{3/2}$  84340 K

I. P. 10.454 volts

The analysis is by Kiess and Corliss, who have reobserved the spectrum, revised, and extended the earlier work especially for inclusion here. The new observations extend from 1200 Å to 23000 Å. The Vacuum Spectrograph of the Bureau was used for the short-wave region. For wavelengths longer than 12000 Å observations were made by Plyler and by Humphreys, who used PbS detectors with an electrodeless discharge as source.

From 2000 Å to 13000 Å the lines were measured relative to the International Secondary Standards of Iron. Precise energy-level values were derived by means of the 400 lines classified in this range. The values were then used to calculate wavelengths suitable as standards in the regions shorter than 2000 Å and longer than 12000 Å, i. e., to correct the wavelength scales at the extreme ends of the observed range. The final line list thus has internally consistent standards over the entire range. The total number of classified lines is approximately 625. Observed intersystem combinations connect the systems of terms having different multiplicities.

The observed  $g$ -values are from Zeeman spectrograms taken in a magnetic field of about 36000 oersteds, with the Weiss magnet of the National Bureau of Standards.

The limit is excellently determined by means of a Ritz formula, from three series:  $ns {}^4P_{3/2}(n=6 \text{ to } 11)$ ,  $np {}^4P_{3/2}(n=6 \text{ to } 9)$ , and  $nd {}^4F_{3/2}(n=5 \text{ to } 9)$ .

## REFERENCES

- S. F. Evans, Proc. Roy. Soc. London [A] **133**, 417 (1931). (I P) (T) (C L)  
 K. Murakawa, Zeit. Phys. **100**, 171 (1938). (I P) (T) (C L)  
 P. F. A. Klinkenberg, Rev. Mod. Phys. **24**, No. 2, 63 (1952). (Summary hfs)  
 F. E. Ebbach and R. A. Fisher, J. Opt. Soc. Am. **44**, 868 (1954). (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 29 (1955). (Summary hfs)  
 R. Onaka, Phys. Rev. (2) **106**, 1178 (1957). (hfs)  
 C. C. Kiess and C. H. Corliss, unpublished material (October 1957). (I P) (T) (C L) (Z E)

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
5 <sup>st</sup> 5 <sup>pt</sup>	5 <sup>pt</sup> 2 <sup>nd</sup>	1 1/2 0%	0.00 7002.15	-7002.15	0.673	5 <sup>st</sup> 5 <sup>pt</sup> (CP)7 <sup>p</sup>	7 <sup>p</sup> 4 <sup>D</sup>	3 1/2 2 1/2 1 1/2 0%	75194.10 82033.80 81782.98	-8839.70 310.83	1.42 1.39
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>a</sup>	6 <sup>a</sup> 4 <sup>P</sup>	2 1/2 1 1/2 0%	54833.46 61819.81 60099.27	-7186.35 923.54	1.576 1.618 2.561	5 <sup>st</sup> 5 <sup>pt</sup> (CP)5 <sup>d</sup>	5 <sup>d</sup> 4 <sup>D</sup>	3 1/2	75511.13		1.320
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>a</sup>	6 <sup>a</sup> 2 <sup>P</sup>	1 1/2 0%	50002.88 63184.76	-7083.88	1.385 0.799	5 <sup>st</sup> 5 <sup>pt</sup> (CP)5 <sup>d</sup>	5 <sup>d</sup> 4 <sup>F</sup>	4 1/2	75704.08		1.338
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>p</sup>	6 <sup>p</sup> 4 <sup>P</sup>	2 1/2 1 1/2 0%	64908.34 67082.12 65358.98	-2155.78 1208.16	1.524 1.415 1.555	5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 15	0%	75714.44		0.80
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>p</sup>	6 <sup>p</sup> 4 <sup>S</sup>	1 1/2	64900.01		1.619		nd 16	1 1/2	75823.98		1.41
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>p</sup>	6 <sup>p</sup> 2 <sup>D</sup>	2 1/2 1 1/2	66844.49 78307.81	-7162.72	1.217 1.316		nd 17	2 1/2	75908.91		1.14
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>p</sup>	6 <sup>p</sup> 4 <sup>D</sup>	3 1/2 2 1/2 1 1/2 0%	65670.00 79529.17 71976.74 73387.18	-6859.17 552.43 -1410.44	1.420 1.370 1.317 1.137	5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 18	2 1/2	76106.57		
5 <sup>st</sup> 5 <sup>pt</sup> (D)6 <sup>a</sup>	6 <sup>a</sup> 2 <sup>D</sup>	2 1/2 1 1/2	66020.64 66355.21	-334.57	1.258 0.828		nd 19	1 1/2	76136.43		1.04
5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 1	1 1/2	67298.45			5 <sup>st</sup> 5 <sup>pt</sup> (CP)4 <sup>f</sup>	nd 20	0%	76746.96		
	nd 2	2 1/2	68549.77				nd 21	2 1/2	76903.28		
5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 3	2 1/2	68588.00				4 <sup>f</sup> 1°	2 1/2	77307.47		
	nd 4	1 1/2	68612.02				4 <sup>f</sup> 2°	3 1/2	77313.00		
	nd 5	2 1/2	70151.32				4 <sup>f</sup> 3°	3 1/2	77390.66		
5 <sup>st</sup> 5 <sup>pt</sup> (S)6 <sup>a</sup>	6 <sup>a</sup> 2 <sup>S</sup>	0%	70354.93		1.96		4 <sup>f</sup> 4°	1 1/2	77366.76		
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>p</sup>	6 <sup>p</sup> 2 <sup>P</sup>	0 1/2 1 1/2	71501.58 73064.56	1553.04	1.239 1.329		4 <sup>f</sup> 5°	2 1/2	77359.68		
5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>p</sup>	6 <sup>p</sup> 2 <sup>S</sup>	0%	71813.97		1.377		4 <sup>f</sup> 6°	1 1/2	77406.86		
5 <sup>st</sup> 5 <sup>pt</sup> (CP)7 <sup>a</sup>	7 <sup>a</sup> 4 <sup>P</sup>	2 1/2 1 1/2 0%	71903.44 78991.16 78415.66	-6987.72 475.50	1.59	5 <sup>st</sup> 5 <sup>pt</sup> (CP)8 <sup>a</sup>	8 <sup>a</sup> 4 <sup>P</sup>	2 1/2	77450.76		1.563
5 <sup>st</sup> 5 <sup>pt</sup> (CP)7 <sup>a</sup>	7 <sup>a</sup> 2 <sup>P</sup>	1 1/2 0%	72294.83 79285.26	-6990.43		5 <sup>st</sup> 5 <sup>pt</sup> (CP)8 <sup>a</sup>	8 <sup>a</sup> 2 <sup>P</sup>	1 1/2	77555.77		1.35
5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 6	1 1/2	72876.75			5 <sup>st</sup> 5 <sup>pt</sup> (D)6 <sup>p</sup>	6 <sup>p</sup> 2 <sup>P</sup>	1 1/2	78416.56		1.39
	nd 7	1 1/2	73477.25			5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 2°	1 1/2	78592.76		1.00
	nd 8	1 1/2	73639.28			5 <sup>st</sup> 5 <sup>pt</sup> (CP)8 <sup>p</sup>	8 <sup>p</sup> 4 <sup>P</sup>	2 1/2 1 1/2 0%	78732.31 79824.78 79903.11	-492.47 21.67	1.454 1.30
5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 9	2 1/2	73846.96			5 <sup>st</sup> 5 <sup>pt</sup> (CP)7 <sup>p</sup>	7 <sup>p</sup> 2 <sup>P</sup>	0 1/2 1 1/2	78780.04 82451.20	3651.16	1.479
	nd 10	3 1/2	73977.68		1.21	5 <sup>st</sup> 5 <sup>pt</sup> (CP)8 <sup>p</sup>	8 <sup>p</sup> 4 <sup>S</sup>	1 1/2	78816.61		1.71
	nd 11	1 1/2	74587.40		0.78	5 <sup>st</sup> 5 <sup>pt</sup> (CP)8 <sup>p</sup>	8 <sup>p</sup> 2 <sup>D</sup>	2 1/2	78854.86		1.11
	nd 12	1 1/2	74625.56			5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>d</sup>	6 <sup>d</sup> 4 <sup>D</sup>	3 1/2	78904.05		1.36
	nd 13	2 1/2	74823.48			5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>d</sup>	6 <sup>d</sup> 4 <sup>P</sup>	2 1/2	78943.37		1.41
5 <sup>st</sup> 5 <sup>pt</sup> (CP)7 <sup>p</sup>	7 <sup>p</sup> 4 <sup>P</sup>	2 1/2 1 1/2 0%	74965.77 75621.41 75303.13	-655.64 318.28	1.472 1.483 1.53	5 <sup>st</sup> 5 <sup>pt</sup> (CP)8 <sup>p</sup>	8 <sup>p</sup> 4 <sup>D</sup>	3 1/2	79003.70		1.03
5 <sup>st</sup> 5 <sup>pt</sup> (CP)7 <sup>p</sup>	7 <sup>p</sup> 4 <sup>S</sup>	1 1/2	75049.57		1.506	5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 22	1 1/2	79030.99		1.23
5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 14	2 1/2	75177.26		1.26	5 <sup>st</sup> 5 <sup>pt</sup> (CP)6 <sup>d</sup>	6 <sup>d</sup> 4 <sup>F</sup>	4 1/2	79055.01		1.332
5 <sup>st</sup> 5 <sup>pt</sup> (CP)7 <sup>p</sup>	7 <sup>p</sup> 2 <sup>D</sup>	2 1/2 1 1/2	75191.37 82424.18	-7232.81	1.24 1.27	5 <sup>st</sup> 5 <sup>pt</sup> ( )nd	nd 23	3 1/2	79150.55		
							nd 24	1 1/2	79395.92		1.42
						5 <sup>st</sup> 5 <sup>pt</sup> (CP)8 <sup>p</sup>	8 <sup>p</sup> 2 <sup>P</sup>	0 1/2	79418.49		1.17
							5 <sup>f</sup> 1°	1 1/2	79701.73		1.02
							5 <sup>f</sup> 2°	0 1/2	79835.03		
						5 <sup>st</sup> 5 <sup>pt</sup> (CP)5 <sup>f</sup>	5 <sup>f</sup> 3°	2 1/2	79840.23		
							5 <sup>f</sup> 4°	3 1/2	79844.58		
							5 <sup>f</sup> 5°	1 1/2	79847.88		
									79881.74		

## I I—Continued

## I I—Continued

Config.	Design.	J	Level	Interval	Obs. g	Config.	Design.	J	Level	Interval	Obs. g
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)9s	9s <sup>2</sup> P	2½	79914. 68		1. 56	5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)10s	10s <sup>2</sup> P	2½	81252. 48		1. 53
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)9s	9s <sup>1</sup> P	1½	79947. 20								
5s <sup>2</sup> 5p <sup>4</sup> ( )np	np <sup>3</sup> °	1½	80069. 94		1. 50	5s <sup>2</sup> 5p <sup>4</sup> ( )nd	nd 29	2½	81371. 2		
	np <sup>4</sup> °	2½	80185. 87		1. 05		nd 30	0½	81551. 9		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)9p	9p <sup>4</sup> P°	2½	80224. 45				nd 31	1½	81642. 25		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)7d	7d <sup>4</sup> D	3½	80676. 17		1. 40	5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)7p	7p <sup>2</sup> S°	0½	81608. 82		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)7d	7d <sup>4</sup> P	2½	80680. 12		1. 37	5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)8d	8d <sup>4</sup> D	3½	81689. 02		
5s <sup>2</sup> 5p <sup>4</sup> ( )nd	nd 26	1½	80690. 98			5s <sup>2</sup> 5p <sup>4</sup> ( )np	np <sup>5</sup> °	3½	81693. 55		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)9p	9p <sup>2</sup> S°	1½	80720. 85			5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)8d	8d <sup>4</sup> F	4½	81760. 58		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)7d	7d <sup>4</sup> F	4½	80772. 35			5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)7f	7f <sup>1</sup> °	2½	82026. 20		
5s <sup>2</sup> 5p <sup>4</sup> ( )nd	nd 27	1½	80782. 49		1. 46		7f <sup>2</sup> °	3½	82030. 35		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)9p	9p <sup>4</sup> D°	2½	80797. 95			5s 5p <sup>4</sup>	5p <sup>4</sup> <sup>2</sup> S	0½	82028. 61		
5s <sup>2</sup> 5p <sup>4</sup> ( )nd	nd 28	3½	80869. 05			5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)10p	10p <sup>4</sup> D°	3½	82035. 00		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)9p	9p <sup>4</sup> D°	3½	80945. 44		1. 32	5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)11s	11s <sup>4</sup> P	2½	82074. 10		
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)6f	6f <sup>1</sup> °	2½	81205. 59			5s <sup>2</sup> 5p <sup>4</sup> ( )np	np <sup>6</sup> °	2½	82214. 04		
	6f <sup>2</sup> °	3½	81207. 32			5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)9d	9d <sup>4</sup> F	4½	82385. 44		
	6f <sup>3</sup> °	1½	81216. 75			5s <sup>2</sup> 5p <sup>4</sup> ( )nd	nd 32	2½	82452. 83		
	6f <sup>4</sup> °	2½	81219. 30			5s <sup>2</sup> 5p <sup>4</sup> ( )np	np <sup>7</sup> °	1½	82515. 84		
						I II( <sup>2</sup> P <sub>2</sub> )	Limit	-----	84340		

October 1957.

## I I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> +	Observed Terms		
5s <sup>2</sup> 5p <sup>4</sup>	5p <sup>4</sup> <sup>2</sup> P°		
5s 5p <sup>4</sup>	5p <sup>4</sup> <sup>2</sup> S		
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 5)
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>2</sup> P)ns	{ 6 to 11s <sup>4</sup> P 6 to 9s <sup>2</sup> P	6 to 9p <sup>4</sup> S° 6 to 9p <sup>4</sup> P° 6 to 10p <sup>4</sup> D° 6, 7p <sup>2</sup> S° 6 to 8p <sup>2</sup> P° 6 to 9p <sup>2</sup> D°	6, 7d <sup>4</sup> P 5 to 8d <sup>4</sup> D 5 to 9d <sup>4</sup> F
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>1</sup> D)ns'	6s' <sup>2</sup> D	6p' <sup>2</sup> P°	
5s <sup>2</sup> 5p <sup>4</sup> ( <sup>1</sup> S)ns''	6s'' <sup>2</sup> S		

\*For predicted terms in the spectra of the I I isoelectronic sequence, see Vol. III, Introduction.

## I II

(Te I sequence; 52 electrons)

Z=53

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^4 \ ^3P_1$  $5p^4 \ ^3P_1$ , 154050 K

I. P. 19.09 volts

The analysis is chiefly by Lacroute who reported in 1935 that more than 1200 lines were known. He published Zeeman data for many lines and observed several series. He utilized the measurements of L. Bloch and E. Bloch in the region from about 2200 Å to 798 Å.

Murakawa in studying the hyperfine structure of I II also reported regularities in the spectrum, and in 1938 revised the series and extended the analysis. Lacroute carried the work further and published in 1939 the term list that has been used for the analysis reported here. Ten levels have been omitted as extremely dubious.

A new description of I II is in course of preparation by Kiess, Corliss, and Martin at the National Bureau of Standards. The available observations extend from 2023.100 Å to 9046.27 Å and include Zeeman spectrograms taken with metal halide lamps as the source. The writer has revised the known term values from these new measurements by starting with  $6p \ ^3P_1$  equal to 100410.00K. These values are given to two decimal places in the table. A complete new description of I II in the region short of 2000 Å is in progress at the Bureau. Several excellent spectrograms extending from 800 Å to 2500 Å have just been obtained. With these observations, additions and corrections to the tabular data will be made.

There are approximately 500 classified lines. Observed intersystem combinations connect the systems of terms of different multiplicities.

Martin has added some new levels, extended the known series, and derived the limit quoted above, from the  $ns \ ^3S^{\circ}$  series ( $n=6$  to 9), by means of an extended Ritz formula.

Most of the observed  $g$ -values in the table given to two decimal places are from Lacroute. All three-place entries are new values derived from the Bureau Zeeman data.

## REFERENCES

- P. Lacroute, Ann. de Phys. (11) **3**, 5 (1935). (I P) (T) (C L) (Z E)  
 K. Murakawa, Zeit. Phys. **109**, 162 (1938). (I P) (T) (C L) (hfs)  
 P. Lacroute, Ann. d'Astroph. **2**, 318 (1939). (I P) (T) (C L) (Z E)  
 C. C. Kiess, C. H. Corliss, and W. C. Martin, unpublished material (December 1957). (I P) (T) (C L) (Z E)

## I II

## I II

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$		
$5s^2 5p^4$	$5p^4 \ ^3P$	2	0	-7090 639		$5s^2 5p^4(^3D^{\circ})6s$	$6s' \ ^3D^{\circ}$	1	92140.04	1558.74 2959.26	0.685		
		1	7090					2	93498.78		1.16		
		0	6451					3	96658.04		1.36		
$5s^2 5p^4$	$5p^4 \ ^1D$	2	13731			$5s^2 5p^4$	$5p^4 \ ^1P^{\circ}$	1	95963.31		1.05		
$5s^2 5p^4$	$5p^4 \ ^1S$	0	32629			$5s^2 5p^4(^3D^{\circ})5d$	$5d' \ ^3D^{\circ}$	1		2859.76	1.26		
$5s^2 5p^4(^3S^{\circ})6s$	$6s \ ^3S^{\circ}$	2	81040.12	1.86	2			97090.73					
		1	81915.20		3			99950.49					
		0	84229.62										
$5s^2 5p^4$	$5p^4 \ ^3P^{\circ}$	2	81915.20	-2314.42	1.54	$5s^2 5p^4(^3D^{\circ})6s$	$6s' \ ^1D^{\circ}$	2	97708.27		0.996		
		1	84229.62	-1161.98	1.530			$5s^2 5p^4(^3D^{\circ})5d$	$5d' \ ^3F^{\circ}$	2		1201.78	1.24?
		0	85391.60		0/0					3	98183.18		
$5s^2 5p^4(^3S^{\circ})6s$	$6s \ ^3S^{\circ}$	1	84850.39		1.753	4	99384.96						
		$5s^2 5p^4(^3S^{\circ})5d$	$5d \ ^3D^{\circ}$	4	86043.66	308.27	1.480	$5s^2 5p^4(^3S^{\circ})6p$	$6p \ ^3P$	1	99226.99	107.54 1075.47	2.309
				3	85735.39	-436.89	1.457			2	99334.53		1.714
2	86172.28			-1569.19	1.434	3	100410.00			1.622			
$5s^2 5p^4(^3S^{\circ})5d$	$5d \ ^3D^{\circ}$	1	87741.47	-2670.93	1.472	$5s^2 5p^4(^3D^{\circ})5d$	$5d' \ ^1G^{\circ}$	4	101165.25				
		0	90412.40		0/0			$5s^2 5p^4(^3S^{\circ})6p$	$6p \ ^3P$	2	102620.89	969.30 -1359.88	1.501
		3	93013.39	2507.63	1.35					1	101651.59		1.520
2	90806.78	-4326.99	1.138	0	103011.47	0/0							
		1	94832.75		0.653								



I II—Continued

I II—Continued

Config.	Design.	J	Level	Interval	Obs. g	Config.	Design.	J	Level	Interval	Obs. g
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)6 <sup>a</sup>	6 <sup>a</sup> ' <sup>a</sup> P°	0 1 2	108071 107519. 18	1448	1. 39	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> S	1	128570. 61		1. 78
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)6 <sup>a</sup>	6 <sup>a</sup> ' <sup>a</sup> P°	1	107628. 58			5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> g	1	129780. 20		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)5 <sup>d</sup>	5 <sup>d</sup> ' <sup>a</sup> P°	0 1 2	108544. 98		1. 45	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>d</sup> or 5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>e</sup>	k° l°	1 2	130622. 68 130624. 98		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)5 <sup>d</sup>	5 <sup>d</sup> ' R°	1	109303. 77			5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> P	0 1 2	131452. 20		1. 26
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> P	1	110014. 15		0. 915	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)8 <sup>a</sup>	8 <sup>a</sup> <sup>a</sup> S°	2	132844. 61		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)5 <sup>d</sup>	5 <sup>d</sup> ' Y°	2	110486. 48		1. 36	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)8 <sup>a</sup>	8 <sup>a</sup> <sup>a</sup> S°	1	133304. 47		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> F	2 3 4	111305. 50 115275. 29 115361. 39	3969. 79 86. 10	0. 915 1. 212 1. 29	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>d</sup> or 5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>a</sup>	a° b° c°	3 3 3	133557. 94 133962. 93 134031. 30		1. 23 1. 16 0. 81
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> D	3 2 1	112426. 46 112186. 91 113820. 24	239. 55 -1633. 33	1. 12 1. 158 0. 985	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>d</sup>	6 <sup>d</sup> ' <sup>a</sup> S°	0	134242. 57		0/0
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> F	3	114164. 65		1. 13	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)7 <sup>d</sup>	7 <sup>d</sup> <sup>a</sup> D°	4 3 2	134509. 75 134449. 09	60. 66	
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)5 <sup>d</sup>	5 <sup>d</sup> ' O°	2	114480. 00					1 0	134554. 39		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> P	0 1 2	114643. 21 116092. 24 115715. 90	1449. 03 -376. 34	0/0 1. 39 1. 425	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>d</sup>	6 <sup>d</sup> ' <sup>a</sup> D°	1 2 3	134967. 58 135592. 96	624. 78	
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)5 <sup>d</sup>	5 <sup>d</sup> ' Z°	2	117942. 70		1. 38?	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>p</sup>	7 <sup>p</sup> ' <sup>a</sup> P	1	135350. 02		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)7 <sup>a</sup>	7 <sup>a</sup> <sup>a</sup> S°	2	118614. 91		1. 894	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>p</sup>	7 <sup>p</sup> ' <sup>a</sup> F	2 3 4	135691. 17 139698. 67 139507. 13	4007. 6 -191. 54	
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> D	2	119090. 24		1. 11			3	136104. 20 136415. 92 136871. 44	-311. 72 -455. 52	
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)7 <sup>a</sup>	7 <sup>a</sup> <sup>a</sup> S°	1	119896. 14			5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>p</sup>	7 <sup>p</sup> ' <sup>a</sup> D	1	136534. 34		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)5 <sup>d</sup>	5 <sup>d</sup> ' <sup>a</sup> P°	0 1 2	121083. 11		1. 443	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>p</sup>	7 <sup>p</sup> ' <sup>a</sup> D	2 1	136708. 04		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)5 <sup>d</sup>	5 <sup>d</sup> ' <sup>a</sup> D°	3 2 1	121202. 92 126146. 16?	-4943. 23	1. 17 1. 18	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>p</sup> or 5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)5 <sup>f</sup>		3	136740. 86		1. 21
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)6 <sup>d</sup>	6 <sup>d</sup> <sup>a</sup> D°	0 1 2 3 4	121709. 05 122250. 90 121896. 02 121785. 70 121833. 41	541. 85 -424. 88 -40. 32 47. 71	0/0 1. 466 1. 453 1. 424 1. 45	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>d</sup> or 5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>a</sup>	e° f° g°	4 3 2	137512. 54 138323. 94		1. 24 1. 27
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)5 <sup>d</sup>	5 <sup>d</sup> ' U°	3	123587. 33			5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>d</sup>	6 <sup>d</sup> ' <sup>a</sup> P°	0 1 2	137021. 77 138371. 5?		0/0
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)5 <sup>f</sup>	5 <sup>f</sup> <sup>a</sup> F	5 4 3 2 1	124749. 64 124691. 29 124719. 30 124790. 59 124842. 28	58. 35 -28. 01 -71. 29 -58. 67		5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>p</sup>	7 <sup>p</sup> ' <sup>a</sup> F	3	139225. 07		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)7 <sup>p</sup>	7 <sup>p</sup> <sup>a</sup> P	1 2 3	124958. 36 125091. 79 125490. 94	133. 43 399. 15		5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)6 <sup>d</sup> or 5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>a</sup>	A° i°	2 2	139559. 17 139641. 52		1. 09 1. 04
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)5 <sup>f</sup>	5 <sup>f</sup> <sup>a</sup> F	2 3 4	125229. 61 125234. 37 125434. 51	4. 76 250. 14	0. 83	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>p</sup>	7 <sup>p</sup> ' <sup>a</sup> P	2 1 0	139918. 97 140044. 25	-125. 28	
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)6 <sup>d</sup>	6 <sup>d</sup> <sup>a</sup> D°	3 2 1	125295. 68 125254. 88 125346. 42	40. 80 -91. 54	1. 28 1. 095 0. 674	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)9 <sup>a</sup>	9 <sup>a</sup> <sup>a</sup> S°	2	139965. 27		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> P°)6 <sup>p</sup>	6 <sup>p</sup> ' <sup>a</sup> D	1 2 3	125532. 05 129438. 75	3906. 70	0. 83 1. 32	5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> S°)9 <sup>a</sup>	9 <sup>a</sup> <sup>a</sup> S°	1 2	140133. 05 140196. 75		
5 <sup>a</sup> 5 <sup>p</sup> ( <sup>a</sup> D°)7 <sup>a</sup>	7 <sup>a</sup> ' <sup>a</sup> D°	1 2 3	127897		0. 89				154050		

I III (<sup>a</sup>S<sub>12</sub>)

Limit

I II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>n+1</sup>	Observed Terms			n' (n ≥ 4)
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)	
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> 1S	5p <sup>2</sup> 1P	5p <sup>2</sup> 1D	
5s 5p <sup>3</sup>	{ 5p <sup>3</sup> 1P <sup>o</sup> 5p <sup>3</sup> 1P <sup>o</sup>			
5s <sup>2</sup> 5p <sup>2</sup> (S <sup>2</sup> ) 1S	{ 6, 7p <sup>2</sup> 1S <sup>o</sup> 6, 7p <sup>2</sup> 1S <sup>o</sup>	6, 7p <sup>2</sup> 1P 6p <sup>2</sup> 1P	6, 7p <sup>2</sup> 1D 6, 7p <sup>2</sup> 1D	6, 7p <sup>2</sup> 1D <sup>o</sup> 6, 6d <sup>2</sup> 1D <sup>o</sup>
5s <sup>2</sup> 5p <sup>2</sup> (D <sup>2</sup> ) 1S'	{ 6, 7p <sup>2</sup> 1D <sup>o</sup> 6p <sup>2</sup> 1D <sup>o</sup>	6, 7p <sup>2</sup> 1P 6, 7p <sup>2</sup> 1P	6, 7p <sup>2</sup> 1D 6, 7p <sup>2</sup> 1D	6d <sup>2</sup> 1S <sup>o</sup> 6, 6d <sup>2</sup> 1P <sup>o</sup> 6, 6d <sup>2</sup> 1D <sup>o</sup> 6d <sup>2</sup> 1P <sup>o</sup> 6d <sup>2</sup> 1D <sup>o</sup>
5s <sup>2</sup> 5p <sup>2</sup> (P <sup>2</sup> ) 1S''	{ 6p <sup>2</sup> 1P <sup>o</sup> 6p <sup>2</sup> 1P <sup>o</sup>	6p <sup>2</sup> 1S 6p <sup>2</sup> 1P	6p <sup>2</sup> 1D	6d <sup>2</sup> 1P <sup>o</sup> 6d <sup>2</sup> 1D <sup>o</sup>

\*For predicted terms in the spectrum of the Te I isoelectronic sequence, see Vol. III, Introduction.

I III

(Sb I sequence; 51 electrons)

Z=53

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>2</sup> 4d<sup>10</sup> 5s<sup>2</sup> 5p<sup>2</sup> 5S<sub>1/2</sub>

5p<sup>2</sup> 5S<sub>1/2</sub> K

I. P. volts

Seth has reported 20 levels, based on measurements by L. and E. Bloch, which he assigns to 6s, 6p, and 6d electrons. In the 1949 reference below it is stated that a few additional levels, and possibly two series members are known. Spectrograms of I III covering a long spectral range are needed to confirm and extend the present work on analysis.

REFERENCES

J. B. Seth, Proc. Indian Acad. Sci. 1A, 593 (1955). (T) (C L)  
S. G. Krishnamurty and T. V. Parthasarathy, Nature 164, 407 (1949).

October 1953.

## I IV

(Sn I sequence; 50 electrons)

Z=53

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3 {}^3P_0$  $5p^3 {}^3P_0$ 

K

I. P.

volts

This spectrum has not been adequately investigated. Krishnamurty has reported 68 lines between 2224.43 Å and 3546.90 Å, half of which he has classified as combinations among 9 even levels and 10 odd levels. By a comparison of isoelectronic data he assigns some term designations and assumes an arbitrary absolute value of 192000.0 K for  $6s {}^3P_2^o$ . Accurate measurements in the short-wave region are needed to establish the ground term and extend the study. Lacking these, the writer has made a crude extrapolation that places  $6s {}^3P_2^o$  roughly 154000 K above the ground state zero. She estimates the value of the limit as approximately 345000 K, and the ionization potential as roughly  $43 \pm$  volts. The error in all these estimates is probably large.

## REFERENCE

S. G. Krishnamurty, Proc. Phys. Soc. London **48**, 277 (1936). (T) (C L)

August 1953.

## I VI

(Cd I sequence; 48 electrons)

Z=53

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 {}^1S_0$  $5s^2 {}^1S_0$ 

K

I. P.

volts

Krishnamurty and Fernando have utilized measurements by L. and E. Bloch to work out an array of 6 multiplets from 17 lines in the range 483 Å to 919 Å. They suggest combinations giving intervals for the  $5p {}^3P^o$  term that are not inconsistent with those extrapolated along the sequence. Confirmation is needed.

## REFERENCE

S. G. Krishnamurty and I. Fernando, Indian J. Phys. **23**, 172 (1949). (C L)

August 1953.

## I VII

(Ag I sequence; 47 electrons)

Z=53

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 4d^{10} 4s^2 4p^4 4d^{10} 5s^2 5s_{1/2}$  $5s^2 5s_{1/2}$ 

K

I. P. volts

Little is known about this spectrum. Five lines measured by L. Bloch, E. Bloch, and N. Felici have been attributed to I VII and classified by Fernando. The terms have been worked out by the writer from these lines.

## REFERENCES

- L. Bloch, E. Bloch, et N. Felici, J. Phys. Rad. [7] 8, No. 9, 355 (1937).  
I. Fernando, Current Sci. 17, 362 (1948). (C L)

## I VII

## I VII

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$4d^{10}(^1S)5s$	$5s^2 S$	0½	0		$4d^{10}(^1S)5d$	$5d^2 D$	1½ 2½	274020 276250	2230
$4d^{10}(^1S)5p$	$5p^2 P^o$	0½ 1½	104960 119960	15000					

March 1953.

## I VIII

(Pd I sequence; 46 electrons)

Z=53

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 4d^{10} 4s^2 4p^4 4d^{10} ^1S_0$  $4d^{10} ^1S_0$ , 1370370 K

I. P. 170 volts

Kruger and Shoupp have classified three lines between 190 Å and 196 Å as combinations with the ground term, and determined the limit from a study of the Moseley diagram of the isoelectronic sequence. The writer has rounded off their term values, and assigned the J-value of the parent term by analogy with Pd I. Two of the lines have been classified independently in the 1937 paper listed below. The quoted ionization potential may be in error by several volts.

## REFERENCES

- P. G. Kruger and W. E. Shoupp, Phys. Rev. 46, 124 (1934). (I P) (T) (C L)  
L. Bloch, E. Bloch et N. Felici, J. Phys. Rad. [7] 8, 362 (1937). (C L)

## I VIII

## I VIII

Config.	Desig.	J	Level	Config.	Desig.	J	Level
$4d^{10}$	$4d^{10} ^1S$	0	0				
$4d^{10}(^3D_{3/2})5p$	$5p^2 P^o$	2 1 0	508770	$4d^{10}(^3D_{3/2})5p$	$5p^2 D^o$	3 2 1	585880
$4d^{10}(^3D_{3/2})5p$	$5p^2 P^o$	1	515060	$I_{IX}(^3D_{3/2})$	Limit		[1370370]

March 1953.

## XENON

## Xe I

54 electrons

Z=54

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 \ ^1S_0$  $5p^6 \ ^1S_0$ , 97834.4 K

I. P. 12.127 volts

The observations of Xe I extend from 926 Å to 21048 Å. Those in the infrared region beyond the photographic limit are listed in the 1949 and 1952 references by Sittner and Peck, and by Humphreys and Kostkowski, respectively. The ultraviolet observations have been extended by Beutler to 926 Å, i. e. beyond the limit, 1192 Å, reached by Boyce.

The energy levels have been taken from an unpublished manuscript kindly furnished by Edlén, who has made a critical study of the published papers on Xe I. Most of the levels are listed in the extensive term array given by Humphreys and Meggers in 1933. Three-place entries are from interferometric measurements. Predicted values, furnished by Edlén, are in brackets.

The observed  $g$ -values are from the 1941 paper, by Green, Hurlburt, and Bowman. Pogány has also published a shorter list of  $g$ -values.

The limit given by Boyce, 97834.4 K, has been used for the present compilation. The other levels are relative to the absolute value of  $6s(1/2)_2^2 = 30766.353$ , adopted by Humphreys and Meggers.

White has discussed the effect of autoionization on the appearance of higher series members having the limit  $^2P_{3/2}^o$ . Beutler has observed a number of these lines in absorption. The levels  $ns'[0X]^\circ$ ,  $n=8$  to 12 and  $nd'[1X]^\circ$ ,  $n=6$  to 18, are from his paper. The separation of the two limits  $^2P_{1/2}^o - ^2P_{3/2}^o$  as observed by Edlén (1944) in Xe II is 10537.01 K.

The Paschen notation is entered in the first column of the table. The letters X, Y, T, Z, U, V, and W, adopted when configurations involving  $f$ -electrons were found, are also included in this column.

As for the other inert gases, Edlén has suggested a pair-coupling notation to take into account the departure from  $LS$ -coupling. By analogy with the other inert gas spectra, and by taking into account the general run of the observed  $g$ -values, the writer has assigned the following provisional  $LS$ -designations, although they have little, if any, significance in most cases.

Paschen	Desig.	Paschen	Desig.	Paschen	Desig.
$(n-5)s_0$	$ns \ ^1P_1$	$3-7d_0$	$5-9d \ ^1P_1$	$3s_1'''$	$5d \ ^1D_2$
$(n-5)s_1$	$ns \ ^1P_1$	$3-7d_1$	$5-9d \ ^1P_1$	$3s_2'''$	$5d \ ^1D_2$
$(n-5)s_2$	$ns \ ^1P_1$	$3-7d_2$	$5-9d \ ^1P_1$	$3s_3'''$	$5d \ ^1D_2$
$(n-5)s_3$	$ns \ ^1P_1$	$3-7d_3$	$5-9d \ ^1P_1$	$3s_4'''$	$5d \ ^1D_2$
		$3-9d_0$	$5-11d \ ^1F_3$		
$2-4p_{10}$	$6-8p \ ^1S_1$	$3-7d_1'$	$5-9d \ ^1F_3$	$4-6X$	$4-6f \ ^1D_2$
$2-4p_0$	$6-8p \ ^1D_2$	$3-7d_1$	$5-9d \ ^1F_3$	$4-6Y$	$4-6f \ ^1D_2$
$2-5p_0$	$6-9p \ ^1D_2$	$3-7d_2$	$5-9d \ ^1F_3$	$4-6U$	$4-6f \ ^1D_2$
$2-4p_1$	$6-8p \ ^1D_2$	$3-7d_2$	$5-9d \ ^1F_3$	$4-6V$	$4-6f \ ^1D_2$
$2-5p_1$	$6-9p \ ^1D_2$				
$2-5p_2$	$6-9p \ ^1D_2$				
$2-5p_3$	$6-9p \ ^1D_2$				

Xe I—Continued

The *Jl*-coupling notation in the general form suggested by Racah is here adopted, and the arrangement is similar to that used for Ne I, Ar I, and Kr I. Because of the wide interval of the parent term, the high levels observed by Beutler might more appropriately be entered after the first ionization limit, as has been suggested by Edlén. In the general form of presentation adopted in these Volumes for noble gas spectra they stand out so conspicuously, however, that no exception in arrangement has been made for Xe I.

REFERENCES

C. J. Humphreys, *Bur. Std. J. Research* **3**, 1041, RP 245 (1930). (T) (C L)  
 H. E. White, *Phys. Rev.* **33**, 1786, 2016 (1931). (E D)  
 E. Rasmussen, *Dissertation*, p. 50 (Copenhagen, 1932). (T) (C L)  
 C. J. Humphreys and W. F. Meggers, *Bur. Std. J. Research* **10**, 139, RP 521 (1933). (I P) (T) (C L)  
 H. Beutler, *Zeit. Phys.* **83**, 177 (1935). (T) (C L)  
 B. Pogány, *Zee-man Verhandeligen*, p. 226 (Martinus Nijhoff 1935). (Z E)  
 W. F. Meggers, *J. Research Nat. Bur. Std.* **14**, 495, RP 781 (1935). (C L)  
 J. C. Boyce, *Phys. Rev.* **49**, 730 (1936). (I P) (C L)  
 J. B. Green, E. H. Huriburt, and D. W. Bowman, *Phys. Rev.* **59**, 72 (1941). (Z E)  
 G. Racah, *Phys. Rev.* **61**, 537 (L) (1942).  
 B. Edlén, *Ark. Mat. Astron. Fys. (Stockholm)* **29A**, No. 32 p. 1 (1943). (C L)  
 J. B. Green, *Phys. Rev.* **64**, 161 (1943). (Z E)  
 B. Edlén, *Phys. Rev.* **65**, 248 (1944).  
 B. Edlén, unpublished material (November 1946). (T)  
 W. R. Sittner and E. R. Peck, *J. Opt. Soc. Am.* **30**, 474 (1949). (C L)  
 C. J. Humphreys and H. J. Kostkowski, *J. Research Nat. Bur. Std.* **49**, 73 RP 2345 (1952). (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 71 (1952). (Summary hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 30 (1955). (Summary hfs)  
 C. J. Humphreys and E. Paul, Jr., U. S. Naval Ord. Lab., NAVORD Report 4589, 25 (1956). (C L)

Xe I

Xe I

Paschen	Config.	Desig.	J	Level	Obs. g	Paschen	Config.	Desig.	J	Level	Obs. g
P <sub>0</sub>	5p <sup>6</sup>	5p <sup>6</sup> 1S	0	0.0		3s <sup>1</sup> <sub>1</sub> <sup>''''</sup> 3s <sup>1</sup> <sub>1</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )5d	5d' [1½]°	2 1	91447.99 93618.75	1.274
1s <sub>1</sub> 1s <sub>2</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )6s	6s [1½]°	2 1	87068.047 89045.693	1.500 1.204	3s <sup>1</sup> <sub>1</sub> <sup>''</sup> 3s <sup>1</sup> <sub>1</sub>	"	5d' [2½]°	2 3	91153.16 91747.070	1.126
1s <sub>1</sub> 1s <sub>2</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )6s	6s' [0½]°	0 1	76197.292 77185.560	0/0 1.321	2s <sub>1</sub> 2s <sub>2</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )7s	7s [1½]°	2 1	85189.31 85440.53	
2p <sub>10</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )6p	6p [0½]	1	77269.649	1.852	2s <sub>1</sub> 2s <sub>2</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )7s	7s' [0½]°	0 1	96721.48 96801.093	1.308
2p <sub>1</sub> 2p <sub>2</sub>	"	6p [2½]	2 3	78120.303 78403.562	1.106 1.336	3p <sub>10</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )7p	7p [0½]	1	87927.652	1.728
2p <sub>1</sub> 2p <sub>2</sub>	"	6p [1½]	1 2	78956.538 79212.970	1.022 1.379	3p <sub>1</sub> 3p <sub>2</sub>	"	7p [2½]	2 3	88352.201 88469.732	1.123 1.348
2p <sub>1</sub>	"	6p [0½]	0	80119.474	0/0	3p <sub>1</sub> 3p <sub>2</sub>	"	7p [1½]	1 2	88745.081 88687.020	0.903 1.347
2p <sub>1</sub> 2p <sub>2</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )6p	6p' [1½]	1 2	88379.647 89162.880	0.790 1.195	3p <sub>1</sub> 3p <sub>2</sub>	"	7p [0½]	0	88842.781	0/0
2p <sub>1</sub> 2p <sub>2</sub>	"	6p' [0½]	1 0	89279.233 89860.538	1.552 0/0	4d <sub>1</sub> 4d <sub>2</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )6d	6d [0½]°	0 1	86491.52 86550.28	
3d <sub>1</sub> 3d <sub>2</sub>	5p <sup>6</sup> ( <sup>2</sup> P <sub>1/2</sub> )5d	5d [0½]°	0 1	79771.798 79987.16	1.395	4d <sub>1</sub> 4d <sub>2</sub>	"	6d [3½]°	4 3	88912.195 89025.39	
3d <sub>1</sub> 3d <sub>2</sub>	"	5d [3½]°	4 3	80197.16 80970.93		4d <sub>1</sub> 4d <sub>2</sub>	"	6d [1½]°	2 1	88708.96 90032.65	
3d <sub>1</sub> 3d <sub>2</sub>	"	5d [1½]°	2 1	80323.28 83890.47	1.376	4d <sub>1</sub> 4d <sub>2</sub>	"	6d [2½]°	2 3	89243.75 89535.05	
3d <sub>1</sub> 3d <sub>2</sub>	"	5d [2½]°	2 3	81929.04 82430.78							

Xe I—Continued

Xe I—Continued

Paschen	Config.	Design.	J	Level	Obs. g	Paschen	Config.	Design.	J	Level	Obs. g
4e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )6d	6d' [1½] <sup>o</sup>	2 1	100418		5p <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )9p	9p [0½]	0	94286. 01	0/0
3e <sub>i</sub> 3e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )8e	8e [1½] <sup>o</sup>	2 1	90805. 045 90852. 839	1. 465 1. 182	6d <sub>i</sub> 6d <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )8d	8d [0½] <sup>o</sup>	0 1	94184. 889 94222. 522	0/0 1. 190
3e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )8e	8e' [0½] <sup>o</sup>	0 1	101486		6d <sub>i</sub> 6d <sub>i</sub>	"	8d [3½] <sup>o</sup>	4 3	94222. 825 94290. 735	1. 236 1. 076
4X 4Y	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )4f	4f [1½]	1 2	90840. 300 90849. 968	0. 504 1. 11	6d <sub>i</sub> 6d <sub>i</sub>	"	8d [1½] <sup>o</sup>	2 1	94222. 168 94235. 24	1. 303 0. 914
4T 4Z	"	4f [4½]	5 4	90861. 18 90862. 03		6d <sub>i</sub> 6d <sub>i</sub>	"	8d [2½] <sup>o</sup>	2 3	94339. 249 94370. 504	0. 987 1. 246
4U 4V	"	4f [2½]	3 2	90907. 614 90910. 583	1. 18 0. 86	6e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )8d	8d' [1½] <sup>o</sup>	2 1	106029	
4W	"	4f [3½]	3 4	90944. 55 90944. 65		6X 6Y	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )6f	6f [1½]	1 2	94735. 411 94737. 654	0. 50 1. 09
4p <sub>10</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )8p	8p [0½]	1	92153. 809	1. 801	6T 6Z	"	6f [4½]	5 4	94744. 73 94745. 17	
4p <sub>9</sub> 4p <sub>8</sub>	"	8p [2½]	2 3	92221. 884 92265. 470	1. 103 1. 272	6U 6V	"	6f [2½]	3 2	94758. 58 94760. 415	1. 17 0. 87
4p <sub>7</sub> 4p <sub>6</sub>	"	8p [1½]	1 2	92333. 588 92371. 442	1. 036 1. 395	6W	"	6f [3½]	3 4	94769. 90 94769. 99	
4p <sub>5</sub>	"	8p [0½]	0	92555. 659	0/0	5e <sub>i</sub> 5e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )10e	10e [1½] <sup>o</sup>	2 1	94760. 431 94787. 803	1. 512 1. 164
5d <sub>3</sub> 5d <sub>2</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )7d	7d [0½] <sup>o</sup>	0 1	92260. 449 92128. 795	0/0 1. 273	5e <sub>2</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )10e	10e' [0½] <sup>o</sup>	0 1	[106316]	
5d <sub>i</sub> 5d <sub>i</sub>	"	7d [3½] <sup>o</sup>	4 3	92445. 434 92648. 655	1. 217 1. 026	6p <sub>10</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )10p	10p [0½]	1	95154. 88	
5d <sub>2</sub> 5d <sub>1</sub>	"	7d [1½] <sup>o</sup>	2 1	92722. 043 92714. 555	1. 196 0. 819	6p <sub>9</sub> 6p <sub>8</sub>	"	10p [2½]	2 3	95182. 17 95197. 04	
5d <sub>i</sub> 5d <sub>i</sub>	"	7d [2½] <sup>o</sup>	2 3	92879. 032 92734. 107	1. 073 1. 263	6p <sub>7</sub> 6p <sub>6</sub>	"	10p [1½]	1 2	95216. 97 95230. 10	
5e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )7d	7d' [1½] <sup>o</sup>	2 1	103419		6p <sub>5</sub>	"	10p [0½]	0	95226. 57	
5X 5Y	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )5f	5f [1½]	1 2	93363. 199 93366. 764	0. 50 1. 10	7d <sub>3</sub> 7d <sub>2</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )9d	9d [0½] <sup>o</sup>	0 1	95180. 107 95222. 213	0/0 1. 217
5T 5Z	"	5f [4½]	5 4	93378. 01 93378. 70		7d <sub>i</sub> 7d <sub>i</sub>	"	9d [3½] <sup>o</sup>	4 3	95250. 456 95283. 522	1. 237 1. 078
5U 5V	"	5f [2½]	3 2	93401. 96 93404. 484	1. 17 0. 87	7d <sub>2</sub> 7d <sub>1</sub>	"	9d [1½] <sup>o</sup>	2 1	95274. 239 95422. 22	1. 298 0. 899
5W	"	5f [3½]	3 4	93421. 30 93421. 40		7d <sub>i</sub> 7d <sub>i</sub>	"	9d [2½] <sup>o</sup>	2 3	95313. 212 95335. 225	0. 980 1. 225
4e <sub>i</sub> 4e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )9e	9e [1½] <sup>o</sup>	2 1	93398. 758 93422. 615	1. 496 1. 154	7e <sub>i</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )9d	9d' [1½] <sup>o</sup>	2 1	105965	
4e <sub>2</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )9e	9e' [0½] <sup>o</sup>	0 1	103954		7X 7Y	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )7f	7f [1½]	1 2	95561. 582 95563. 079	
5p <sub>10</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )9p	9p [0½]	1	94067. 46		7T 7Z	"	7f [4½]	5 4	95567. 65 95567. 99	
5p <sub>9</sub> 5p <sub>8</sub>	"	9p [2½]	2 3	94110. 66 94135. 04	1. 307	7U 7V	"	7f [2½]	3 2	95576. 44 95577. 71	
5p <sub>7</sub> 5p <sub>6</sub>	"	9p [1½]	1 2	94169. 37 94190. 75	1. 386	7W	"	7f [3½]	3 4	95583. 59 95583. 67	

Xo I—Continued

Xo I—Continued

Paschen	Config.	Design.	J	Level	Obs. g	Paschen	Config.	Design.	J	Level	Obs. g
6a <sub>1</sub> 6a <sub>2</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )11s	11s [1½] <sup>o</sup>	2 1	95579. 469 95591. 48	1. 188	9X 9Y	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )9f	9f [1½]	1 2	96463. 14 96463. 78	
6a <sub>1</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )11s	11s' [0½] <sup>o</sup>	0 1	106137		9T 9Z	"	9f [4½]	5 4	96466. 14 96466. 23	
7p <sub>10</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )11p	11p [0½]	1	[95831. 3]		9U 9V	"	9f [2½]	3 2	96470. 24 96470. 90	
7p <sub>9</sub> 7p <sub>8</sub>	"	11p [3½]	2 3	[95849. 1] 95858. 09		9W	"	9f [3½]	3, 4	96473. 66	
7p <sub>7</sub> 7p <sub>6</sub>	"	11p [1½]	1 2	95871. 30 95879. 91		8a <sub>1</sub> 8a <sub>2</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )13s	13s [1½] <sup>o</sup>	2 1	96479. 64 96481. 13	
7p <sub>5</sub>	"	11p [0½]	0	95916. 04		9p <sub>10</sub> 9p <sub>9</sub> 9p <sub>8</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )13p	13p [2½]	2 3 1	96607. 28 96617. 41	
8d <sub>10</sub> 8d <sub>9</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )10d	10d [0½] <sup>o</sup>	0 1	95896. 76 95913. 388	1. 081	9p <sub>7</sub> 9p <sub>6</sub>	"	13p [1½]	1 2	96617. 41	
8d <sub>8</sub> 8d <sub>7</sub>	"	10d [3½] <sup>o</sup>	4 3	95898. 708 95918. 585		9p <sub>5</sub>	"	13p [0½]	0	96634. 60	
8d <sub>6</sub> 8d <sub>5</sub>	"	10d [1½] <sup>o</sup>	2 1	95906. 108 96046. 88		10d <sub>10</sub> 10d <sub>9</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )12d	12d [0½] <sup>o</sup>	0 1	[96609. 5] 96616. 73	
8d <sub>4</sub> 8d <sub>3</sub>	"	10d [2½] <sup>o</sup>	2 3	95938. 31 95947. 183		10d <sub>8</sub> 10d <sub>7</sub>	"	12d [3½] <sup>o</sup>	4 3	96623. 31 96632. 14	
8a <sub>1</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )10d	10d' [1½] <sup>o</sup>	2 1	106548		10d <sub>6</sub> 10d <sub>5</sub>	"	12d [1½] <sup>o</sup>	2 1	96627. 59 96634. 90	
8X 8Y	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )8f	8f [1½]	1 2	96096. 76 96097. 82		10d <sub>4</sub> 10d <sub>3</sub>	"	12d [2½] <sup>o</sup>	2 3	96641. 85 96649. 67	
8T 8Z	"	8f [4½]	5 4	96101. 11 96101. 26		10a <sub>1</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )12d	12d' [1½] <sup>o</sup>	2 1	107816	
8U 8V	"	8f [2½]	3 2	96106. 82 96107. 87		10X 10Y	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )10f	10f [1½]	1 2	96725. 0	
8W	"	8f [3½]	3 4	96111. 68 96111. 75		10T 10Z	"	10f [4½]	5 4	96726. 98 96727. 07	
7a <sub>1</sub> 7a <sub>2</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )12s	12s [1½] <sup>o</sup>	2 1	96109. 73 96125. 88		10U 10V	"	10f [2½]	3 2	96730. 00 [96730. 5]	
7a <sub>3</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )12s	12s' [0½] <sup>o</sup>	0 1	[106658]		10W	"	10f [3½]	3	[96732. 5]	
8p <sub>10</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )12p	12p [0½]	1	[96290. 1]		9a <sub>1</sub> 9a <sub>2</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )14s	14s [1½] <sup>o</sup>	2 1	96731. 83 [96737. 9]	
8p <sub>9</sub> 8p <sub>8</sub>	"	12p [2½]	2 3	[96292. 6] 96299. 46		10p <sub>10</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )14p	14p [2½]	2 3	96830. 65	
8p <sub>7</sub> 8p <sub>6</sub>	"	12p [1½]	1 2	96307. 37 96313. 40		10p <sub>9</sub>	"	14p [1½]	1 2	96837. 65	
8p <sub>5</sub>	"	12p [0½]	0	96338. 33		11d <sub>10</sub> 11d <sub>9</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )13d	13d [3½] <sup>o</sup>	4 3	96848. 64 96849. 10	
9d <sub>10</sub> 9d <sub>9</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )11d	11d [0½] <sup>o</sup>	0 1	96306. 13 96316. 67	1. 082	11d <sub>8</sub> 11d <sub>7</sub>	"	13d [2½] <sup>o</sup>	2 3	96856. 08 96862. 08	
9d <sub>8</sub> 9d <sub>7</sub>	"	11d [3½] <sup>o</sup>	4 3	96322. 06 96334. 987		11a <sub>1</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )13d	13d' [1½] <sup>o</sup>	2 1	[107434]	
9d <sub>6</sub> 9d <sub>5</sub>	"	11d [1½] <sup>o</sup>	2 1	96329. 13 96424. 88		11T 11Z	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )11f	11f [4½]	5 4	96919. 72	
9d <sub>4</sub> 9d <sub>3</sub>	"	11d [2½] <sup>o</sup>	1 3	96348. 54 96359. 07		11U	"	11f [2½]	3 2	96922. 02	
9a <sub>1</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )11d	11d' [1½] <sup>o</sup>	2 1	106939							



Xe I—Continued

Xe I—Continued

Paschen	Config.	Desig.	J	Level	Obs. $\nu$	Paschen	Config.	Desig.	J	Level	Obs. $\nu$
10 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )15s	15s [1½] <sup>o</sup>	2 1	[99999. 6]		13 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )15d	15d' [1½] <sup>o</sup>	2 1	107708	
11 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )15p	15p [2½]	2 3	99999. 2		14 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )16d	16d' [1½] <sup>o</sup>	2 1	107808	
12 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )14d	14d [3½] <sup>o</sup>	4 3	97007. 41		15 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )17d	17d' [1½] <sup>o</sup>	2 1	107870	
12 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )14d	14d' [1½] <sup>o</sup>	2 1	107588		16 <sub>o</sub>	5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )18d	18d' [1½] <sup>o</sup>	2 1	107938	
	Xe II( <sup>2</sup> P <sub>1/2</sub> )	Limit	-----	97884. 4			Xe II( <sup>2</sup> P <sub>1/2</sub> )	Limit	-----	108371. 4	

October 1956.

Xe I OBSERVED LEVELS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> +	Observed Terms							
	5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> <sup>3</sup> S						
	ns (n ≥ 6)	np (n ≥ 6)			nd (n ≥ 5)			n' (n ≥ 4)
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sup>o</sup> )ns	{ 6-14s <sup>1</sup> P <sup>o</sup> 6-9, 11s <sup>1</sup> P <sup>o</sup>	6-8p <sup>3</sup> S 6p <sup>3</sup> S	6-9p <sup>1</sup> P 6p <sup>1</sup> P	6-9p <sup>1</sup> D 6-9p <sup>1</sup> D	5-9d <sup>3</sup> P <sup>o</sup> 5-9d <sup>1</sup> P <sup>o</sup>	5d <sup>1</sup> D <sup>o</sup> 5d <sup>1</sup> D <sup>o</sup>	5-11d <sup>3</sup> F <sup>o</sup> 5-9d <sup>1</sup> F <sup>o</sup>	4-6f <sup>1</sup> D 4-6f <sup>1</sup> D
Jl-Coupling Notation								
	Observed Pairs							
	ns (n ≥ 6)	np (n ≥ 6)			nd (n ≥ 5)			n' (n ≥ 4)
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> )ns	6-14s [1½] <sup>o</sup>	6-13p [0½] 6-15p [2½] 6-14p [1½]			5-12d [0½] <sup>o</sup> 5-14d [3½] <sup>o</sup> 5-12d [1½] <sup>o</sup> 5-13d [2½] <sup>o</sup>			4-9f [1½] 4-11f [4½] 4-11f [2½] 4-9f [3½]
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>2</sup> P <sub>3/2</sub> )ns'	6-9, 11s' [0½] <sup>o</sup>	6p' [1½] 6p' [0½]			5d' [2½] <sup>o</sup> 5-12, 14-18d' [1½] <sup>o</sup>			

\*For predicted levels in the spectra of the Xe I isoelectronic sequence, see Vol. III, Introduction.

## Xe II

(I I sequence; 53 electrons)

Z=54

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 \text{ } ^1\text{P}_{11}$  $5p^6 \text{ } ^1\text{P}_{11}$ , 171068.4 K

I. P. 21.2 volts

The analysis is from Humphreys' 1939 paper with the revisions suggested in the 1939 paper on Zeeman Effect by Humphreys, Meggers, and de Bruin. More than 660 lines have been classified, including 20 in the ultraviolet from 740.406 Å to 1244.756 Å reported by Boyce, in addition to Humphreys' list, which extends from 2230.79 Å to 10220.8 Å.

Edlén attributes a line listed as Xe I at 9487.76 Å to the forbidden transition giving the interval of the ground term of Xe II, namely, 10537.01 K. He states that the line is due to a magnetic-dipole transition and that the transition probability is  $21 \text{ sec}^{-1}$ .

The limit is "the approximate Rydberg limit of the s-electron series", the absolute term values being based on the  $\text{ } ^1\text{P}_{11}$  series.

The g-values in the table are from the paper by Humphreys, Meggers, and de Bruin, and from Angenetter. For the most part, they are average values of the various observers.

Observed intersystem combinations connect the doublet and quartet systems of terms.

## REFERENCES

- C. J. Humphreys, T. L. deBruin, and W. F. Meggers, Bur. Std. J. Research **6**, 287, RP 275 (1931). (I P)  
(T) (C L)  
J. C. Boyce, Phys. Rev. **48**, 730 (1936). (C L)  
C. J. Humphreys, J. Research Nat. Bur. Std. **33**, 19, RP 1164 (1939). (I P) (T) (C L)  
C. J. Humphreys, W. F. Meggers, and T. L. de Bruin, J. Research Nat. Bur. Std. **33**, 683, RP 1264 (1939).  
(T) (C L) (Z E)  
H. Angenetter, Zeit. Phys. **115**, 309 (1940). (Z E)  
B. Edlén, Phys. Rev. **68**, 249 (1944). (C L)  
A. G. Shenstone, Proc. Roy. Soc. [A] **219**, 419 (1953).

## Xe II

## Xe II

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g		
$5s^2 5p^6$	$5p^6 \text{ } ^1\text{P}^o$	$1\frac{1}{2}$	0.0	-10537.3		$5s^2 5p^6 \text{ } ^1\text{P}^o 5d$	5d $\text{ } ^1\text{P}$	$2\frac{1}{2}$	106475.17	-906.58	0.67		
		$0\frac{1}{2}$	10537.3					$1\frac{1}{2}$	107381.75			475.64	1.79
$5s^2 5p^6$	$5p^6 \text{ } ^3\text{S}$	$0\frac{1}{2}$	90873.83		2.02	$5s^2 5p^6 \text{ } ^3\text{P}^o 5d$	5d $\text{ } ^3\text{D}$	$1\frac{1}{2}$	107904.50	1658.65	1.20		
		$1\frac{1}{2}$	93068.40	-1995.94	1.56			$2\frac{1}{2}$	109563.15			1.33	
		$0\frac{1}{2}$	95064.34	-8093.17	1.38								
$5s^2 5p^6 \text{ } ^3\text{P}^o 6s$	6s $\text{ } ^3\text{P}$	$2\frac{1}{2}$	93068.40		1.56	$5s^2 5p^6 \text{ } ^3\text{D}^o 6s$	6s' $\text{ } ^3\text{D}$	$2\frac{1}{2}$	108007.25	-4917.45	1.22		
		$1\frac{1}{2}$	95064.34		1.38			$1\frac{1}{2}$	112924.70			0.95	
		$0\frac{1}{2}$	101157.51		2.43								
$5s^2 5p^6 \text{ } ^3\text{P}^o 5d$	5d $\text{ } ^3\text{D}$	$3\frac{1}{2}$	95437.60	40.92	1.38	$5s^2 5p^6 \text{ } ^3\text{D}^o 5d$	5d' $\text{ } ^3\text{D}$	$1\frac{1}{2}$	111326.93	1376.65	1.24		
		$2\frac{1}{2}$	95396.68	-636.76	1.34			$2\frac{1}{2}$	112703.58			1.13	
		$1\frac{1}{2}$	96033.44	-824.70	1.18								
		$0\frac{1}{2}$	96858.14		0.50								
$5s^2 5p^6 \text{ } ^3\text{P}^o 5d$	5d $\text{ } ^3\text{F}$	$4\frac{1}{2}$	99404.90	582.24	1.31	$5s^2 5p^6 \text{ } ^3\text{F}^o 6p$	6p $\text{ } ^3\text{P}^o$	$2\frac{1}{2}$	111958.84	166.76	1.47		
		$3\frac{1}{2}$	98822.66	-2712.99	1.07			$1\frac{1}{2}$	111798.08			1.61	
		$2\frac{1}{2}$	101535.65					$0\frac{1}{2}$	113672.85			-1890.77	1.50
		$1\frac{1}{2}$											
$5s^2 5p^6 \text{ } ^3\text{P}^o 6s$	6s $\text{ } ^3\text{P}$	$1\frac{1}{2}$	102799.06	-1451.00	1.59	$5s^2 5p^6 \text{ } ^3\text{F}^o 6p$	6p $\text{ } ^3\text{D}^o$	$3\frac{1}{2}$	115705.53	193.04	1.40		
		$0\frac{1}{2}$	104250.06		0.56			$2\frac{1}{2}$	115512.89			-3270.82	1.28
								$1\frac{1}{2}$	116783.11			-3631.76	1.37
$5s^2 5p^6 \text{ } ^3\text{P}^o 5d$	5d $\text{ } ^3\text{P}$	$1\frac{1}{2}$	105313.33	-634.21	1.15	$5s^2 5p^6 \text{ } ^3\text{D}^o 5d$	5d' $\text{ } ^3\text{F}$	$2\frac{1}{2}$	114751.08	162.88	1.10		
		$0\frac{1}{2}$	105947.54		1.36			$3\frac{1}{2}$	114913.96				

Xe II—Continued

Xe II—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
5s 5p <sup>2</sup> ( <sup>3</sup> S)5d	5d'' <sup>1</sup> D	3/4	110085.50	-5216.68			5°	1/4	139128.84		
		1/4	124302.18					1/4	139182.98		
5s 5p <sup>2</sup> ( <sup>3</sup> F)6p	6p'' <sup>3</sup> S°	0%	121179.89		2.08		7°	1/4	139498.63		
5s 5p <sup>2</sup> ( <sup>3</sup> F)6p	6p'' <sup>3</sup> S°	1/4	121628.30		1.28		9°	1/4	139646.1		
5s 5p <sup>2</sup> ( <sup>3</sup> F)6p	6p'' <sup>1</sup> D°	3/4	123112.50	-1177.00	1.39		11°	2/4	139646.1		
		1/4	124289.50		1.32			13°	0%		
5s 5p <sup>2</sup> ( <sup>3</sup> F)6p	6p'' <sup>1</sup> P°	1/4	123254.61	-1316.41	1.35		15°	2/4	140030.11		1.08
		0%	124571.02		1.16			17°	0%		
5s 5p <sup>2</sup> ( <sup>3</sup> S)6s	6s'' <sup>3</sup> S	0%	124070.09			5s 5p <sup>2</sup> ( <sup>3</sup> S)6p	6p'' <sup>1</sup> P°	1/4	140210.22		
5s 5p <sup>2</sup> ( <sup>1</sup> D)5d	5d'' <sup>1</sup> P	0%	127010.7	516.7			19°	0%	140335.12		
		1/4	127537.4					1/4	140883.4		
5s 5p <sup>2</sup> ( <sup>1</sup> D)6p	6p'' <sup>1</sup> F°	2/4	128257.19	1196.84	0.92	5s 5p <sup>2</sup> ( <sup>3</sup> F)6d	6d'' <sup>3</sup> P	0%	144140.2	3256.8	2.25
5s 5p <sup>2</sup> ( <sup>1</sup> D)5d	5d'' <sup>3</sup> S	0%	129248.5		1.15			1/4	143399.4	-740.8	0.83
		3/4	130094.03					2/4	143399.4		1.19
5s 5p <sup>2</sup> ( <sup>1</sup> D)6p	6p'' <sup>1</sup> P°	1/4	129687.37	-2072.72	1.40	5s 5p <sup>2</sup> ( <sup>3</sup> F)7s	7s'' <sup>1</sup> P	1/4	140987.0		
5s 5p <sup>2</sup> ( <sup>1</sup> D)6p	6p'' <sup>1</sup> D°	0%	129741.09		0.71			0%	142930.0	-547.8	1.59
		1/4	131223.78	272.92	0.90	5s 5p <sup>2</sup> ( <sup>3</sup> F)6d	6d'' <sup>1</sup> D	1/4	142382.2		1.05
5s 5p <sup>2</sup> ( <sup>3</sup> F)7s	7s'' <sup>3</sup> P	2/4	132112.7		1.58			0%	142930.0		
		1/4	132189.2	-670.5	1.40	5s 5p <sup>2</sup> ( <sup>3</sup> F)6d	6d'' <sup>3</sup> D	2/4	144385.0	-1555.1	1.21
5s 5p <sup>2</sup> ( <sup>3</sup> F)7s	7s'' <sup>1</sup> P	0%	132000.9	-1871.7	2.08			1/4	145940.1		1.04
		2	0%	132916.9			5s 5p <sup>2</sup> ( <sup>3</sup> F)6d	6d'' <sup>1</sup> P	1/4	145222.7	
4	0%				133454.2					5s 5p <sup>2</sup> ( <sup>1</sup> D)7s	7s'' <sup>1</sup> D
		6	2/4	133915.1				14	2/4		
5s 5p <sup>2</sup> ( <sup>3</sup> F)6d	6d'' <sup>1</sup> D				3/4	135507.2	-39.8			1.39	5s 5p <sup>2</sup> ( <sup>1</sup> D)6d'
		2/4	135547.0	-161.2	1.33	2/4	149697.3				
5s 5p <sup>2</sup> ( <sup>3</sup> F)6d	6d'' <sup>1</sup> F	1/4	135708.2		1.33	5s 5p <sup>2</sup> ( <sup>1</sup> D)6d'	6d'' <sup>1</sup> P	1/4	148374.9		
		0%						3/4	148958.4		
16	1/4	136509.25			1.68	5s 5p <sup>2</sup> ( <sup>1</sup> D)6d'	6d'' <sup>3</sup> D	2/4	149281.0		
			18	0%	136554.17				0.94	37°	2/4
8	2/4	136569.6						5s 5p <sup>2</sup> ( <sup>1</sup> D)6d	6d'' <sup>1</sup> F		
			5s 5p <sup>2</sup> ( <sup>3</sup> F)6d	6d'' <sup>3</sup> F	4/4	136597.7	-2496.4			1.14	39°
3/4	136604.1	-546.2			1.14	41°	1/4	157222.1			
10	2/4	136726.5							Xe m( <sup>3</sup> P <sub>2</sub> )	Limit	-----
			5s 5p <sup>2</sup> ( <sup>1</sup> D)4f	4f'' <sup>3</sup> G°	3/4	137312.21					
5s 5p <sup>2</sup> ( <sup>1</sup> D)4f	4f'' <sup>1</sup> F°	2/4			137329.18	104.57	1.06				
		12	1/4	138062.8		1.00					
20	0%				138726.4		1.86				
		3°	3/4	139005.4							

## Xe II OBSERVED TERMS \*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>n</sup> +	Observed Terms				
5s <sup>2</sup> 5p <sup>2</sup>	5p <sup>2</sup> <sup>3</sup> P <sup>o</sup>				
5s 5p <sup>3</sup>	5p <sup>3</sup> <sup>4</sup> S				
	ns (n ≥ 6)		np (n ≥ 6)		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>3</sup> P)nx	{	6, 7s <sup>4</sup> P	6p <sup>4</sup> S <sup>o</sup>	6p <sup>4</sup> P <sup>o</sup>	6p <sup>4</sup> D <sup>o</sup>
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>1</sup> D)nx'		6, 7s <sup>2</sup> P	6p <sup>2</sup> S <sup>o</sup>	6p <sup>2</sup> P <sup>o</sup>	6p <sup>2</sup> D <sup>o</sup>
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>3</sup> S)nx''		6, 7s' <sup>2</sup> D	6p' <sup>2</sup> P <sup>o</sup>	6p' <sup>2</sup> D <sup>o</sup>	6p' <sup>2</sup> F <sup>o</sup>
	nd (n ≥ 5)		nf (n ≥ 4)		
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>3</sup> P)nx	{	5, 6d <sup>4</sup> P	5, 6d <sup>4</sup> D	5, 6d <sup>4</sup> F	
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>1</sup> D)nx'		5d' <sup>2</sup> S	5d' <sup>2</sup> P	5, 6d' <sup>2</sup> D	5, 6d' <sup>2</sup> F
5s <sup>2</sup> 5p <sup>2</sup> ( <sup>3</sup> S)nx''			5d'' <sup>2</sup> D		4f' <sup>2</sup> F <sup>o</sup> 4f' <sup>2</sup> G <sup>o</sup>

\* For predicted terms in the spectra of the I I isoelectronic sequence, see Vol. III, Introduction.

## Xe III

(Te I sequence; 52 electrons)

Z=54

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s<sup>2</sup> 5p<sup>4</sup> <sup>3</sup>P<sub>1</sub>

5p<sup>4</sup> <sup>3</sup>P<sub>1</sub>, 259089 K

I. P. 32.1 volts

The terms are from the 1936 paper by Humphreys, with the revisions introduced by Humphreys, Meggers, and de Bruin from a study of the Zeeman effect, reported in 1939. Humphreys lists nearly 300 classified lines between 2235.35 Å and 7653.8 Å. Boyce has observed the Xe III spectrum in the extreme ultraviolet, and in collaboration with Humphreys has classified 128 lines between 627.393 Å and 1978.702 Å. "The most intense lines of the (<sup>4</sup>S) family have been classified by de Bruin". Excepting one level, 6d <sup>4</sup>D<sup>o</sup>, the terms found by de Bruin are incorporated in the present analysis.

Observed intersystem combinations connect the singlet and triplet systems of terms.

"The absolute term values are arrived at from an estimation of the limit of the 5s<sup>2</sup> 5p<sup>2</sup>(<sup>3</sup>S)nd <sup>4</sup>D<sup>o</sup> series", (n=5, 6).

Edlén has classified two lines whose wave numbers agree with the ground-level separations of terms from 5s<sup>2</sup> 5p<sup>4</sup>, giving the following intervals:

$$5p^4 \ ^3P_2 - 5p^4 \ ^3P_1 = 9794.6 \text{ K}$$

$$5p^4 \ ^3P_2 - 5p^4 \ ^1D_2 = 17098.97 \text{ K.}$$

He attributes these forbidden lines to magnetic-dipole transitions. His value of the interval <sup>3</sup>P<sub>2</sub> - <sup>3</sup>P<sub>1</sub> is quoted in the table.

For miscellaneous levels the designated published numbers have been retained in column two, with the running electron added. The last six levels have "nx" substituted for the running electron, since it is not known whether this electron is 6d or 7s.

## Xe III—Continued

## REFERENCES

J. C. Boyce, *Phys. Rev.* **49**, 730 (1936). (I P) (C L)C. J. Humphreys, *J. Research Nat. Bur. Std.* **16**, 639, RP 898 (1936). (I P) (T) (C L)C. J. Humphreys, W. F. Meggers, and T.-L. de Bruin, *J. Research Nat. Bur. Std.* **33**, 683, RP 1264 (1939). (T) (C L) (Z E)B. Edlén, *Phys. Rev.* **65**, 248 (1944). (C L)

## Xe III

## Xe III

Config.	Desig.	J	Level	Interval	Obs.g	Config.	Desig.	J	Level	Interval	Obs.g				
5s <sup>2</sup> 5p <sup>4</sup>	5p <sup>4</sup> <sup>1</sup> P	2	0	-9794.6 1664		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6s	6s' <sup>1</sup> D°	1	133234.88	1433.21 3990.79	0.38 1.18 1.33				
		1	9795					2	134667.49						
		0	8131					3	136658.28						
5s <sup>2</sup> 5p <sup>4</sup>	5p <sup>4</sup> <sup>1</sup> D	2	17100			5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' 41°	2	136367.66		0.90				
		5s <sup>2</sup> 5p <sup>4</sup>	5p <sup>4</sup> <sup>3</sup> S	0	37398					5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' <sup>3</sup> D°	1	138146.81	3918.88 1092.05	0.50 1.12 1.22
5s 5p <sup>5</sup>	5p <sup>5</sup> <sup>3</sup> P°	2	98263	-5306 -5890		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' <sup>3</sup> S°	1	140731.23				1.56		
		1	103589									2	142064.49		
0	109469					3	143156.54		0.31						
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)5d	5d' <sup>4</sup> D°	4		-250.88 -593.51 -244.01		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6s	6s' <sup>1</sup> D°	2	143048.37						
		3	111605.48							5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' <sup>1</sup> D°	2	145300.39		2.28
		2	111856.34									1	146781.71	180.89 2099.00	1.70 1.57
		1	112449.85									2	148962.60		
		0	112893.86							3	149061.60				
5s 5p <sup>5</sup>	5p <sup>5</sup> <sup>1</sup> P°	1	119026.28			5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 13°	1	147797.64						
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)5d	5d' <sup>3</sup> D°	3	121229.85	-3461.87 2768.69				5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 15°	2	148370.28				
		2	124691.72							5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 17°	3	148413.09		
		1	121923.03					5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 19°			2	150404.88		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)6s	6s <sup>4</sup> S°	2	121476.00		1.95	5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 17°	3	148413.09						
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)6s	6s <sup>3</sup> S°	1	125617.20		1.77	5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)6p	6p <sup>3</sup> P	0	152808.36	-2507.07 1756.65	0.00 1.56 1.50				
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' <sup>3</sup> F°	4		-2229.44		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 19°	1	150301.29						
		3	126120.07									2	152057.94		
		2	123349.51												
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' <sup>1</sup> F°	3	127722.29			5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)6s	6s'' <sup>3</sup> P°	0	150505.45	977.15 2410.81	0.00 1.47				
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' <sup>3</sup> G°	3	130174.00	1986.06		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)6s	6s'' <sup>1</sup> P°	1	151422.60						
		4	132180.06									2	153893.41		
		5													
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)5d	5d' <sup>1</sup> G°	4	132438.59			5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 23°	2	158928.29						

Xe III—Continued

Xe III—Continued

Config.	Design.	J	Level	Interval	Obs.g	Config.	Design.	J	Level	Interval	Obs.g
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6p	6p' <sup>2</sup> D	1	158997. 10	3263. 00 4114. 12	1. 17 1. 22	5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)6p	6p'' 28	1	182134. 41		
		2	162290. 10								
		3	166374. 22								
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 25°	1	159333. 39			5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)7s	7s °S°	2	182337. 31		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6p	6p' <sup>2</sup> F	2	160691. 50	1903. 43 3960. 02	0. 84 1. 08 1. 28	5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)6p	6p'' <sup>2</sup> P	2 1 0	183220. 15		
		3	162594. 93								
		4	166554. 95								
		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)5d	5d'' 27°								
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6p	6p' <sup>1</sup> F	3	164438. 80	1. 09		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)6p	6p'' 34	1	184570. 60		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6p	6p' 4	1	164511. 82	0. 42		5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)7s	7s °S°	1	188023. 28		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6p	6p' <sup>2</sup> P	0	166295. 67	1790. 48 -1205. 94	1. 30 1. 38	5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)ns	ns' 29°	2	195897. 68		
		1	169086. 15								
		2	166880. 21								
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)4f?	4f' 6	4	166355. 52	1. 31		"	ns' 33°	2 or 3	196261. 30		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)6p	6p' <sup>1</sup> D	2	171990. 00	1. 08		"	ns' 35°	2 or 3	196537. 90		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)6p	6p'' <sup>1</sup> D	3	173946. 80	-4009. 27 2724. 72	1. 18 0. 65	"	ns' 37°	2 or 3	196808. 67		
		2	177956. 07								
		1	175231. 35								
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)6p	6p'' 26	1	178029. 58	1. 51		"	ns' 39°	3	200060. 37		
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)6d	6d <sup>1</sup> D°	4	182122. 97	-341. 27 -18. 51 -68. 38 -289. 66		Xe IV( <sup>4</sup> S <sub>1/2</sub> )	Limit	-----	259669		
		3	182464. 24								
		2	182432. 76								
		1	182551. 13								
		0	182840. 79								

October 1953.

Xe III OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> +	Observed Terms		
5s <sup>2</sup> 5p <sup>4</sup>	{ 5p <sup>4</sup> <sup>1</sup> S	5p <sup>4</sup> <sup>2</sup> P	5p <sup>4</sup> <sup>1</sup> D
5s 5p <sup>5</sup>	{ 5p <sup>5</sup> <sup>2</sup> P°	5p <sup>5</sup> <sup>1</sup> P°	
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 5)
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>4</sup> S°)ns	6, 7s <sup>4</sup> S° 6, 7s <sup>2</sup> S°	6p <sup>2</sup> P 6p <sup>2</sup> P	5, 6d <sup>1</sup> D° 5d <sup>2</sup> D°
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> D°)ns'		6p' <sup>2</sup> P 6p' <sup>1</sup> D 6p' <sup>2</sup> F 6p' <sup>1</sup> D 6p' <sup>1</sup> F	5d' <sup>2</sup> S° 5d' <sup>2</sup> D° 5d' <sup>2</sup> F° 5d' <sup>2</sup> G° 5d' <sup>1</sup> D° 5d' <sup>1</sup> F° 5d' <sup>1</sup> G°
5s <sup>2</sup> 5p <sup>3</sup> ( <sup>2</sup> P°)ns''	6s'' <sup>2</sup> P° 6s'' <sup>1</sup> P°	6p'' <sup>2</sup> P 6p'' <sup>1</sup> D	

\*For predicted terms in the spectra of the Te I isoelectronic sequence, see Vol. III, Introduction.

## Xe IV

(Sb 1 sequence; 51 electrons)

Z=54

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3 \text{ } ^4\text{S}_{1/2}$  $5p^3 \text{ } ^4\text{S}_{1/2}$ 

K

I. P.

volts

The analysis is by Humphreys, who has furnished the data in the table in advance of publication. His work is based on observations made by Boyce. There are 83 classified lines in the interval 558 Å to 1249 Å.

Further observations are needed to extend the analysis. In cases where the observed combinations do not determine the  $J$ -value, the two possible choices are indicated in column three. No series are known in this spectrum.

## REFERENCES

J. C. Boyce, unpublished material (1936).

C. J. Humphreys, unpublished material (October 1953). (T) (C L)

## Xe IV

## Xe IV

Config.	Desig.	$J$	Level	Interval	Config.	Desig.	$J$	Level	Interval
$5s^2 5p^3$	$5p^3 \text{ } ^4\text{S}^\circ$	$1\frac{1}{2}$	0				$0\frac{1}{2}$	154532	
$5s^2 5p^3$	$5p^3 \text{ } ^2\text{D}^\circ$	$1\frac{1}{2}$ $2\frac{1}{2}$	13268 17512	4244			$0\frac{1}{2}$	157205?	
$5s^2 5p^3$	$5p^3 \text{ } ^2\text{P}^\circ$	$0\frac{1}{2}$ $1\frac{1}{2}$	28036? 41837	13847			$0\frac{1}{2}$	158610	
		$1\frac{1}{2}, 2\frac{1}{2}$	99664				$1\frac{1}{2}, 2\frac{1}{2}$	159643	
		$1\frac{1}{2}, 2\frac{1}{2}$	106924				$0\frac{1}{2}$	161435	
		$0\frac{1}{2}$	109255				$1\frac{1}{2}$	163596	
		$1\frac{1}{2}$	121929				$0\frac{1}{2}$	165281?	
		$1\frac{1}{2}, 2\frac{1}{2}$	125475				$1\frac{1}{2}$	165995	
		$1\frac{1}{2}$	133027				$0\frac{1}{2}$	167208	
		$1\frac{1}{2}, 2\frac{1}{2}$	134981				$1\frac{1}{2}, 2\frac{1}{2}$	170132	
		$1\frac{1}{2}, 2\frac{1}{2}$	136495				$1\frac{1}{2}, 2\frac{1}{2}$	170491	
		$1\frac{1}{2}$	145107				$1\frac{1}{2}$	173221	
		$1\frac{1}{2}, 2\frac{1}{2}$	146205				$1\frac{1}{2}, 2\frac{1}{2}$	176043	
		$1\frac{1}{2}, 2\frac{1}{2}$	148684				$1\frac{1}{2}, 2\frac{1}{2}$	179001	
							$1\frac{1}{2}, 2\frac{1}{2}$	186052	

January 1956.

## CESIUM

Cs I

55 electrons

Z=55

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 6s^2 S_{1/2}$  $6s^2 S_{1/2}$  31406.71 K

I. P. 3.893 volts

A number of references have been used to derive the best available level values from the separate sets of wavelength measurements, starting with the ground state zero. For the principal series and the limit quoted above, the results of Kratz are quoted. He has observed this series in absorption to  $n=73$ . McNally and his associates have also carried the observations of this series in absorption to  $n=62$ , and give the limit as 31406.32 K. For the  $^2D$  series Mack has furnished his unpublished observations of the forbidden lines  $6s^2 S - nd^2 D$  ( $n=9$  to 21), for inclusion here. The work of Meissner and Weinmann has provided data for some  $^2D$ ,  $^2S$  terms and for the  $^2F^\circ$  series. Meggers' observations furnish revised values for  $6p^2 P^\circ$  and  $6d^2 D$ .

The well-known series from Fowler and Paschen-Götze are given by Bacher and Goudsmit. Their terms that still need to be re-evaluated from more accurate observations have been rounded off in the table, namely,  $11, 12s^2 S$ ;  $5, 6g^2 G$ ; and  $6h^2 H^\circ$ . New measurements in the near infrared furnished by Fisher in advance of publication, give the connection of the  $5d^2 D$  and  $nj^2 F^\circ$  terms with the rest of the terms, and the quoted value of  $7s^2 S$ .

Beutler and Guggenheimer have classified 53 ultraviolet lines observed in absorption, between 653.60 Å and 1007.5 Å, as transitions from the ground term to levels above the ionization limit. The high limits are from the Cs II analysis. These authors have classified four groups of lines called A, B, C, D, having the respective limits in Cs II  $^2P_{3/2}^\circ$ ,  $^2P_{1/2}^\circ$ ,  $^2P_{3/2}^\circ$ , and  $^2P_{1/2}^\circ$ . The value of the limit they use for group C ( $^2P_{3/2}^\circ$ ), 150870K, has since been replaced in Cs II by one at 153772 K. The writer has, therefore, attempted to adjust the series for group C by assuming that their first series members  $5d$  and  $7s$  are correct as given, and adopting the best available lines that fit the revised limit. Consequently, their level at 142577 K is now labeled  $6d^2$  instead of  $8s$ ; their level at 139694 K, formerly called  $6d$ , has been rejected; and the  $8s$  level is tentatively placed at 144501 K. The miscellaneous levels, which are the wave numbers of the observed ultraviolet lines, have been assigned numbers by the writer. The double entries of  $J$  for these levels (unlike those for unresolved terms) indicate that the existing data are insufficient to determine which value of  $J$  is correct.

The ground state  $6s^2 S_{1/2}$  has a hyperfine structure with a separation of about 0.30 K observed by Kratz from  $740$  to  $72p^2 P^\circ$ . Mack has also observed it in the  $6s^2 S - nd^2 D$  observations from  $n=9$  to 15.

More than 200 lines are classified between 653.60 Å and  $7.4 \mu$ .



## Cs I—Continued

## REFERENCES

- K. W. Meissner, *Ann. der Phys.* [4] **65**, 378 (1921). (T) (C L)  
 A. Fowler, *Report on Series in Line Spectra*, p. 106 (Fleetway Press, London, 1922). (I P) (T) (C L)  
 F. Paschen und R. Götze, *Seriengesetze der Linienspektren*, p. 63 (Julius Springer, Berlin, 1922). (I P) (T) (C L)  
 R. F. Baehar and S. Goudsmit, *Atomic Energy States*, p. 171 (McGraw-Hill Book Co., Inc., New York and London, 1932). (T)  
 W. F. Meggers, *Bur. Std. J. Research* **10**, 674, RP 558 (1933). (C L)  
 H. Beutler und K. Guggenheimer, *Zeit. Phys.* **88**, 25 (1934). (T) (C L)  
 H. Kopfermann und H. Krüger, *Zeit. Phys.* **102**, 527 (1936). (hfs)  
 K. W. Meissner und W. Weinmann, *Ann. der Phys.* [5] **39**, 758 (1937). (T) (C L)  
 N. Tai-Zé and C. Shang-Yi, *Phys. Rev.* **54**, 1045 (1938). (Pressure shift)  
 S. Millman and P. Kusch, *Phys. Rev.* **69**, 438 (1940). (hfs)  
 J. R. McNally, Jr., J. P. Molnar, W. J. Hitchcock, and N. F. Oliver, *J. Opt. Soc. Am.* **30**, 57 (1940). (I F) (T) (C L)  
 D. Harting and P. F. A. Klinkenberg, *Physica* **14**, 669 (1949). (Z E)  
 H. R. Kratz, *Phys. Rev.* **78**, 1844 (1949). (I P) (T) (C L)  
 J. E. Mack, unpublished material (1949). (T) (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 R. A. Fisher, unpublished material (February 1954). (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 30 (1955). (Summary hfs)

## Cs I

## Cs I

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
5p <sup>4</sup> (S)6s	6s <sup>2</sup> S	0½	0.00		5p <sup>4</sup> (S)10s	10s <sup>2</sup> S	0½	28300.28	
5p <sup>4</sup> (S)6p	6p <sup>2</sup> P <sup>o</sup>	0½ 1½	11178.24 11732.35	554.11	5p <sup>4</sup> (S)6f	6f <sup>2</sup> F <sup>o</sup>	3½ 2½	28329.660 28329.768	-0.102
5p <sup>4</sup> (S)5d	5d <sup>2</sup> D	1½ 2½	14492.490 14597.08	97.59	5p <sup>4</sup> (S)6g	6g <sup>2</sup> G	3½, 4½	28347	
5p <sup>4</sup> (S)7s	7s <sup>2</sup> S	0½	18535.51		5p <sup>4</sup> (S)6h	6h <sup>2</sup> H <sup>o</sup>	4½, 5½	28356	
5p <sup>4</sup> (S)7p	7p <sup>2</sup> P <sup>o</sup>	0½ 1½	21765.65 21946.66	181.01	5p <sup>4</sup> (S)10p	10p <sup>2</sup> P <sup>o</sup>	0½ 1½	28787.09 28755.83	26.84
5p <sup>4</sup> (S)6d	6d <sup>2</sup> D	1½ 2½	22568.89 22631.83	42.94	5p <sup>4</sup> (S)9d	9d <sup>2</sup> D	1½ 2½	28828.90 28836.06	7.16
5p <sup>4</sup> (S)8s	8s <sup>2</sup> S	0½	24317.17		5p <sup>4</sup> (S)11s	11s <sup>2</sup> S	0½	29130	
5p <sup>4</sup> (S)4f	4f <sup>2</sup> F <sup>o</sup>	3½ 2½	24472.287 24472.463	-0.176	5p <sup>4</sup> (S)7f	7f <sup>2</sup> F <sup>o</sup>	3½ 2½	29142.156 29142.225	-0.069
5p <sup>4</sup> (S)8p	8p <sup>2</sup> P <sup>o</sup>	0½ 1½	25709.14 25791.78	82.64	5p <sup>4</sup> (S)11p	11p <sup>2</sup> P <sup>o</sup>	0½ 1½	29403.68 29421.10	17.42
5p <sup>4</sup> (S)7d	7d <sup>2</sup> D	1½ 2½	26047.86 26068.83	20.97	5p <sup>4</sup> (S)10d	10d <sup>2</sup> D	1½ 2½	29468.54 29473.22	4.68
5p <sup>4</sup> (S)9s	9s <sup>2</sup> S	0½	26910.68		5p <sup>4</sup> (S)12s	12s <sup>2</sup> S	0½	29666	
5p <sup>4</sup> (S)5f	5f <sup>2</sup> F <sup>o</sup>	3½ 2½	26971.415 26971.562	-0.147	5p <sup>4</sup> (S)8f	8f <sup>2</sup> F <sup>o</sup>	3½ 2½	29678.955 29678.979	-0.044
5p <sup>4</sup> (S)5g	5g <sup>2</sup> G	3½, 4½	27010		5p <sup>4</sup> (S)12p	12p <sup>2</sup> P <sup>o</sup>	0½ 1½	29852.85 29864.72	11.87
5p <sup>4</sup> (S)9p	9p <sup>2</sup> P <sup>o</sup>	0½ 1½	27637.29 27681.96	44.67	5p <sup>4</sup> (S)11d	11d <sup>2</sup> D	1½ 2½	29896.64 29899.89	3.25
5p <sup>4</sup> (S)8d	8d <sup>2</sup> D	1½ 2½	27811.25 27822.94	11.69	5p <sup>4</sup> (S)9f	9f <sup>2</sup> F <sup>o</sup>	3½ 2½	30042.515 30042.540	-0.025

## Cs I—Continued

## Cs I—Continued

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
5p <sup>a</sup> (1S)13p	13p <sup>a</sup> P°	0½ 1½	30166. 00 30174. 51	8. 51	5p <sup>a</sup> (1S)24p	24p <sup>a</sup> P°	0½, 1½	31144. 13	
5p <sup>a</sup> (1S)12d	12d <sup>a</sup> D	1½ 2½	30197. 02 30199. 35	2. 33	5p <sup>a</sup> (1S)25p	25p <sup>a</sup> P°	0½, 1½	31168. 04	
5p <sup>a</sup> (1S)10f	10f <sup>a</sup> F°	3½ 2½	30302. 383 30302. 390	-0. 007	5p <sup>a</sup> (1S)26p	26p <sup>a</sup> P°	0½, 1½	31188. 87	
5p <sup>a</sup> (1S)14p	14p <sup>a</sup> P°	0½ 1½	30395. 16 30399. 49	6. 33	5p <sup>a</sup> (1S)27p	27p <sup>a</sup> P°	0½, 1½	31207. 07	
5p <sup>a</sup> (1S)13d	13d <sup>a</sup> D	1½ 2½	30416. 06 30417. 76	1. 70	5p <sup>a</sup> (1S)28p	28p <sup>a</sup> P°	0½, 1½	31223. 09	
5p <sup>a</sup> (1S)11f	11f <sup>a</sup> F°	3½, 2½	30494. 59		5p <sup>a</sup> (1S)29p	29p <sup>a</sup> P°	0½, 1½	31237. 24	
5p <sup>a</sup> (1S)15p	15p <sup>a</sup> P°	0½ 1½	30565. 87 30567. 98	4. 71	5p <sup>a</sup> (1S)30p	30p <sup>a</sup> P°	0½, 1½	31249. 81	
5p <sup>a</sup> (1S)14d	14d <sup>a</sup> D	1½ 2½	30580. 46 30581. 79	1. 33	5p <sup>a</sup> (1S)31p	31p <sup>a</sup> P°	0½, 1½	31261. 05	
5p <sup>a</sup> (1S)12f	12f <sup>a</sup> F°	3½, 2½	30640. 93		5p <sup>a</sup> (1S)32p	32p <sup>a</sup> P°	0½, 1½	31271. 13	
5p <sup>a</sup> (1S)16p	16p <sup>a</sup> P°	0½ 1½	30685. 76 30687. 52	3. 76	5p <sup>a</sup> (1S)33p	33p <sup>a</sup> P°	0½, 1½	31280. 17	
5p <sup>a</sup> (1S)15d	15d <sup>a</sup> D	1½ 2½	30707. 11 30707. 96	0. 85	5p <sup>a</sup> (1S)34p	34p <sup>a</sup> P°	0½, 1½	31288. 36	
5p <sup>a</sup> (1S)17p	17p <sup>a</sup> P°	0½ 1½	30796. 20 30799. 15	2. 95	5p <sup>a</sup> (1S)35p	35p <sup>a</sup> P°	0½, 1½	31295. 76	
5p <sup>a</sup> (1S)16d	16d <sup>a</sup> D	1½ 2½	30806. 59 30807. 2	0. 6	5p <sup>a</sup> (1S)36p	36p <sup>a</sup> P°	0½, 1½	31302. 51	
5p <sup>a</sup> (1S)18p	18p <sup>a</sup> P°	0½ 1½	30878. 07 30880. 41	2. 34	5p <sup>a</sup> (1S)37p	37p <sup>a</sup> P°	0½, 1½	31308. 61	
5p <sup>a</sup> (1S)17d	17d <sup>a</sup> D	1½, 2½	30886. 7		5p <sup>a</sup> (1S)38p	38p <sup>a</sup> P°	0½, 1½	31314. 21	
5p <sup>a</sup> (1S)19p	19p <sup>a</sup> P°	0½ 1½	30944. 49 30946. 45	1. 94	5p <sup>a</sup> (1S)39p	39p <sup>a</sup> P°	0½, 1½	31319. 39	
5p <sup>a</sup> (1S)18d	18d <sup>a</sup> D	1½, 2½	30951. 5		5p <sup>a</sup> (1S)40p	40p <sup>a</sup> P°	0½, 1½	31324. 11	
5p <sup>a</sup> (1S)20p	20p <sup>a</sup> P°	0½ 1½	30999. 15 31000. 74	1. 59	5p <sup>a</sup> (1S)41p	41p <sup>a</sup> P°	0½, 1½	31328. 47	
5p <sup>a</sup> (1S)19d	19d <sup>a</sup> D	1½, 2½	31005. 0		5p <sup>a</sup> (1S)42p	42p <sup>a</sup> P°	0½, 1½	31332. 47	
5p <sup>a</sup> (1S)21p	21p <sup>a</sup> P°	0½ 1½	31044. 63 31046. 00	1. 37	5p <sup>a</sup> (1S)43p	43p <sup>a</sup> P°	0½, 1½	31336. 19	
5p <sup>a</sup> (1S)20d	20d <sup>a</sup> D	1½, 2½	31049. 5		5p <sup>a</sup> (1S)44p	44p <sup>a</sup> P°	0½, 1½	31339. 67	
5p <sup>a</sup> (1S)22p	22p <sup>a</sup> P°	0½, 1½	31084. 08		5p <sup>a</sup> (1S)45p	45p <sup>a</sup> P°	0½, 1½	31342. 84	
5p <sup>a</sup> (1S)21d	21d <sup>a</sup> D	1½, 2½	31086. 7		5p <sup>a</sup> (1S)46p	46p <sup>a</sup> P°	0½, 1½	31345. 80	
5p <sup>a</sup> (1S)23p	23p <sup>a</sup> P°	0½, 1½	31116. 40		5p <sup>a</sup> (1S)47p	47p <sup>a</sup> P°	0½, 1½	31348. 58	
					5p <sup>a</sup> (1S)48p	48p <sup>a</sup> P°	0½, 1½	31351. 18	
					5p <sup>a</sup> (1S)49p	49p <sup>a</sup> P°	0½, 1½	31353. 57	
					5p <sup>a</sup> (1S)50p	50p <sup>a</sup> P°	0½, 1½	31355. 85	
					5p <sup>a</sup> (1S)51p	51p <sup>a</sup> P°	0½, 1½	31357. 96	
					5p <sup>a</sup> (1S)52p	52p <sup>a</sup> P°	0½, 1½	31359. 95	
					5p <sup>a</sup> (1S)53p	53p <sup>a</sup> P°	0½, 1½	31361. 80	

## Cs I—Continued

## Cs I—Continued

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
5p <sup>2</sup> ( <sup>1</sup> S)54p	54p <sup>1</sup> P°	0½, 1½	31363. 60		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )7s	7s <sup>1</sup> P°	1½	123498	-707
5p <sup>2</sup> ( <sup>1</sup> S)55p	55p <sup>1</sup> P°	0½, 1½	31365. 23		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )7s		0½	124205	
5p <sup>2</sup> ( <sup>1</sup> S)56p	56p <sup>1</sup> P°	0½, 1½	31366. 78		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d 7°	1½	124804	
5p <sup>2</sup> ( <sup>1</sup> S)57p	57p <sup>1</sup> P°	0½, 1½	31368. 31		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )7s	7s <sup>1</sup> P°	1½	125096	
5p <sup>2</sup> ( <sup>1</sup> S)58p	58p <sup>1</sup> P°	0½, 1½	31369. 69		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )6d	6d 8°	0½, 1½	126203	
5p <sup>2</sup> ( <sup>1</sup> S)59p	59p <sup>1</sup> P°	0½, 1½	31370. 96		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )6d	6d 9°	0½, 1½	126488	
5p <sup>2</sup> ( <sup>1</sup> S)60p	60p <sup>1</sup> P°	0½, 1½	31372. 26		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )6d	6d 10°	0½, 1½	127426	
5p <sup>2</sup> ( <sup>1</sup> S)61p	61p <sup>1</sup> P°	0½, 1½	31373. 43		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )6d	6d 11°	0½, 1½	127868	
5p <sup>2</sup> ( <sup>1</sup> S)62p	62p <sup>1</sup> P°	0½, 1½	31374. 56		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )6d	6d 12°	0½, 1½	127973	
5p <sup>2</sup> ( <sup>1</sup> S)63p	63p <sup>1</sup> P°	0½, 1½	31375. 62		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d' <sup>1</sup> F°	1½	130329	
5p <sup>2</sup> ( <sup>1</sup> S)64p	64p <sup>1</sup> P°	0½, 1½	31376. 62		5p <sup>2</sup> 6s( <sup>1</sup> P <sub>1</sub> )5d	5d' <sup>1</sup> F°	1½	130527	
5p <sup>2</sup> ( <sup>1</sup> S)65p	65p <sup>1</sup> P°	0½, 1½	31377. 59		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )8s	8s <sup>1</sup> P°	0½	131211	
5p <sup>2</sup> ( <sup>1</sup> S)66p	66p <sup>1</sup> P°	0½, 1½	31378. 56		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )6d	6d 13°	0½, 1½	131256	
5p <sup>2</sup> ( <sup>1</sup> S)67p	67p <sup>1</sup> P°	0½, 1½	31379. 46		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )8s	8s <sup>1</sup> P°	1½	131504	
5p <sup>2</sup> ( <sup>1</sup> S)68p	68p <sup>1</sup> P°	0½, 1½	31380. 29		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )7d	7d 14°	0½, 1½	131826	
5p <sup>2</sup> ( <sup>1</sup> S)69p	69p <sup>1</sup> P°	0½, 1½	31381. 08		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )7d	7d 15°	0½, 1½	132550	
5p <sup>2</sup> ( <sup>1</sup> S)70p	70p <sup>1</sup> P°	0½, 1½	31381. 87		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )7d	7d 16°	0½, 1½	132706	
5p <sup>2</sup> ( <sup>1</sup> S)71p	71p <sup>1</sup> P°	0½, 1½	31382. 67		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )9s?	9s <sup>1</sup> P°	1½	133564	
5p <sup>2</sup> ( <sup>1</sup> S)72p	72p <sup>1</sup> P°	0½, 1½	31383. 23		5p <sup>2</sup> 6s( <sup>1</sup> P <sub>1</sub> )5d	5d' 17°	1½	133906	
5p <sup>2</sup> ( <sup>1</sup> S)73p	73p <sup>1</sup> P°	0½, 1½	31383. 65		5p <sup>2</sup> 6s( <sup>1</sup> P <sub>1</sub> )5d	5d' 18°	0½	134212	
					5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )8d	8d 19°	} 0½, 1½	134614	
					5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )9s?	9s 20°			
Cs II( <sup>1</sup> S <sub>0</sub> )	Limit		31406. 71		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )8d	8d 21°	0½, 1½	135014	
5p <sup>2</sup> ( <sup>3</sup> P <sub>1</sub> )6s <sup>2</sup>	6s <sup>2</sup> <sup>1</sup> P°	1½	99259	-9810	5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )7s	7s' <sup>1</sup> P°	0½	135164	
5p <sup>2</sup> ( <sup>1</sup> P <sub>1</sub> )6s <sup>2</sup>		0½	109069		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )10s	10s <sup>1</sup> P°	1½?	135322	
5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d 1°	1½	113491		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )12s?	12s <sup>1</sup> P°	1½	} 136855	
5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d 2°	0½	114560		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )11s?	11s 22°	0½, 1½		
5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d 3°	0½	117539		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )13s	13s <sup>1</sup> P°	1½	137297	
5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d 4°	1½	117707		5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )12s?	12s 23°	0½, 1½	137541	
5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d 5°	1½	118574		5p <sup>2</sup> 6s( <sup>1</sup> P <sub>1</sub> )7s	7s' <sup>1</sup> P°	0½	137794	1009
5p <sup>2</sup> 6s( <sup>3</sup> P <sub>1</sub> )5d	5d 6°	0½	122653				1½	138803	
					Cs II( <sup>3</sup> P <sub>1</sub> )	Limit		138799	

## Cs I—Continued

## Cs I—Continued

Config.	Design.	<i>J</i>	Level	Interval	Config.	Design.	<i>J</i>	Level	Interval
Cs II( <sup>1</sup> Pt)	Limit	-----	139312		5p <sup>6</sup> 6s( <sup>1</sup> Pt) 9s	9s' 1P°	0%	149390	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 6d?	6d' 1P°	0%	148577		5p <sup>6</sup> 6s( <sup>1</sup> Pt) 8d	8d' 3P°	0%, 1%	150404	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 6d	6d' 24°	0%, 1%			5p <sup>6</sup> 6s( <sup>1</sup> Pt) 9d	9d' 31°	0%, 1%	151551	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 6d	6d' 25°	0%, 1%	148991		5p <sup>6</sup> 6s( <sup>1</sup> Pt) 9d	9d' 32°	0%, 1%	151744	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 6d	6d' 26°	0%, 1%	148877		5p <sup>6</sup> 6s( <sup>1</sup> Pt) 11s	11s' 1P°	0%	151995	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 8s?	8s' 1P°	0%	144501		5p <sup>6</sup> 6s( <sup>1</sup> Pt) 10d	10d' 33°	0%, 1%	152998	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 8s	8s' 1P°	0%?	146309		5p <sup>6</sup> 6s( <sup>1</sup> Pt) 11d	11d' 34°	0%, 1%	152722	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 7d	7d' 27°	0%, 1%	147922		5p <sup>6</sup> 6s( <sup>1</sup> Pt) 12d?	12d' 35°	0%, 1%	152999	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 7d	7d' 28°	0%, 1%	147994		Cs II( <sup>1</sup> Pt)	Limit	-----	153772	
5p <sup>6</sup> 6s( <sup>1</sup> Pt) 7d	7d' 29°	0%, 1%	148225		Cs II( <sup>1</sup> Pt)	Limit	-----	154273	

February 1955.

## Cs II

(Xe I sequence; 54 electrons)

*Z* = 55Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s<sup>2</sup> 5p<sup>6</sup> <sup>1</sup>S<sub>0</sub>5p<sup>6</sup> <sup>1</sup>S<sub>0</sub> 202263 K

I. P. 25.1 volts

The analysis of Cs II was started by Sommer in 1924 and has since been revised and extended by Sawyer and his associates. There are nearly 350 classified lines, and the observations extend from 564.25 Å to 6955.519 Å.

The levels listed in the table are from the 1942 paper, which contains an array of the observed combinations. From the descriptions of the lines given in the various papers, including that in the 1942 paper on "The Hyperfine Structure of Cs II", and, by analogy with Xe I, the writer has tentatively assigned the *Jl*-coupling notation to levels from the 6, 7s, 5, 6d, and 6, 7p configurations. These assignments need further confirmation and are far from definitive. Trees has pointed out that configuration-interaction is prominent in this spectrum; consequently, further assignments of the pair-coupling notation are very difficult. The numbers used by Wheatley and Sawyer to label the many levels have, therefore, been retained in column 1 of the table.

The limit is from Boyd and Sawyer, who derived it by means of a Rydberg formula from the two-member series 5p<sup>6</sup> <sup>1</sup>S<sub>0</sub> - *ns*'[0%]<sub>11</sub>, *n* = 6, 7. The higher member of the limit term has been obtained by adding the interval of the ground term of Cs III, 13884 K, to the limit quoted above.

## REFERENCES

- O. Laporte, G. R. Miller, and R. A. Sawyer, *Phys. Rev.* **39**, 458 (1932). (I P) (T) (C L)  
 J. Olthoff and R. A. Sawyer, *Phys. Rev.* **43**, 766 (1932). (T) (C L)  
 M. A. Wheatley and R. A. Sawyer, *Phys. Rev.* **61**, 591 (1942). (T) (C L)  
 R. A. Boyd and R. A. Sawyer, *Phys. Rev.* **61**, 601 (1942). (I P) (hfs)  
 J. E. Mack, *Rev. Mod. Phys.* **22**, 64 (1950). (Summary hfs)

## Cs II

## Cs II

Authors	Config.	Desig.	J	Level	Interval	Authors	Config.	Desig.	J	Level	Interval
	5p <sup>4</sup>	5p <sup>4</sup> 1B	0	0.00		23f			1	166392. 31	
3f 5f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )6s	6s [1½] <sup>o</sup>	2 1	107392. 33 107906. 01		24f			2	168717. 91	
13f 14f	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )6s	6s' [0½] <sup>o</sup>	0 1	122365. 51 122966. 03		25f			0	162352. 92	
4f 2f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )5d	5d [0½] <sup>o</sup>	0 1	107563. 14 106222. 77		26f			3	162322. 96	
6f 7f	"	5d [1½] <sup>o</sup>	2 1	108304. 19 110945. 18		29f			2	164444. 82	
8f 11f	"	5d [3½] <sup>o</sup>	4 3	112239. 53 105949. 74		30f			1	164466. 70	
9f 10f	"	5d [2½] <sup>o</sup>	2 3	112796. 08 115716. 61		31f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )7d,		1	164866. 77	
11f 15f	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )5d	5d' [1½] <sup>o</sup>	2 1	119665. 41 123056. 44		32f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )8s,		1	165813. 70	
12f	"	5d' [2½] <sup>o</sup>	2 3	120404. 87		33f	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )6d		1	165843. 10	
1	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )6p	6p [0½]	1	126518. 54		34f			2	165890. 08	
2 <sub>s</sub> 3 <sub>s</sub>	"	6p [2½]	2 3	128089. 83 129107. 65		35f			3	166117. 72	
4 <sub>s</sub> 5 <sub>s</sub>	"	6p [1½]	1 2	129989. 72 130766. 00		36f			2	166131. 11	
6 <sub>s</sub>	"	6p [0½]	0	133153. 54		37f			3	166600. 74	
7 <sub>s</sub> 9 <sub>s</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )6p	6p' [1½]	1 2	141555. 59 143394. 19		38f			2	166687. 09	
8 <sub>s</sub> 10 <sub>s</sub>	"	6p' [0½]	1 0	143352. 12 144523. 45		39f			2	166961. 57	
16f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )7s	7s [1½] <sup>o</sup>	2 1	149212. 25		40f			3	167015. 97	
27f 28f	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )7s	7s' [0½] <sup>o</sup>	0 1	163024. 20 163180. 20		41f			1	167434. 91	
17f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )6d	6d [0½] <sup>o</sup>	0 1	149606. 33		42f			0	168721. 56	
19f 18f	"	6d [1½] <sup>o</sup>	2 1	152791. 49 152172. 11		43f			1	169213. 40	
22f	"	6d [3½] <sup>o</sup> ?	4 3	153678. 17		44f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )8d,		2	169528. 87	
20f 21f	"	6d [2½] <sup>o</sup> ?	2 3	153302. 27 153556. 54		45f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )9s,		1	170363. 90	
11 <sub>s</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )7p	7p [0½]	1	155965. 45		46f	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )7d,		2	170504. 22	
12 <sub>s</sub> 13 <sub>s</sub>	"	7p [2½]	2 3	156115. 18 156513. 16		47f	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )8s		3	172544. 21	
15 <sub>s</sub> 14 <sub>s</sub>	"	7p [1½]	1 2	157062. 46 156757. 30		48f			3	172226. 52	
16 <sub>s</sub>	"	7p [0½]	0	157743. 28		49f			0	172244. 21	
17 <sub>s</sub> 19 <sub>s</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>3/2</sub> )7p	7p' [1½]	1 2	166096. 39 168525. 71		50f			1	173227. 00	
18 <sub>s</sub>	"	7p' [0½]	1 0	167453. 71		51f			1	175951. 42	
						52f	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )10s	10s [1½] <sup>o</sup>	2 1	177243. 50	
						27 <sub>s</sub>			2	188791. 09	
						26 <sub>s</sub>			2	191002. 05	
						25 <sub>s</sub>			3	191103. 04	
						24 <sub>s</sub>	5p <sup>4</sup> ( <sup>2</sup> P <sub>1/2</sub> )nf		2	192475. 51	
						23 <sub>s</sub>			3	192530. 62	
						21 <sub>s</sub>			1	192675. 63	
						22 <sub>s</sub>			3	197642. 3	
						20 <sub>s</sub>			3	200406. 90	
							Cs III( <sup>2</sup> P <sub>1/2</sub> )	Limit		202263	
							Cs III( <sup>2</sup> P <sub>3/2</sub> )	Limit		216147	

January 1954.

## Cs III

(I I sequence; 53 electrons)

Z=55

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 {}^3P_{1,1}$  $5p^6 {}^3P_{1,1}$ 

K

I. P.

volts

This spectrum needs further study. Fitzgerald and Sawyer have classified 17 lines between 529.7 Å and 877.9 Å as combinations of the ground term with higher energy levels. By analogy with Xe II the writer has labeled the third level in the table as  $5p^6 {}^3S$ . The remaining "even" levels are probably from the  $5s^2 5p^4 5d$  and  $5s^2 5p^4 6s$  configurations.

From a study of screening constants Finkelnburg and Humbach have extrapolated an ionization potential of  $34.6 \pm 0.7$  volts.

## REFERENCES

- M. A. Fitzgerald and R. A. Sawyer, *Phys. Rev.* **46**, 576 (1934). (T) (C L)  
 W. Finkelnburg und W. Humbach, *Naturwiss.* **42**, 35 (1955). (I P)

## Cs III

Config.	Design.	J	Level	Interval
$5s^2 5p^4$	$5p^4 {}^1P^o$	$1\frac{1}{2}$	0	-13884
		$0\frac{1}{2}$	13884	
$5s^2 5p^4$	$5p^4 {}^3S$	$0\frac{1}{2}$	127786	
			136146	
			138453	
			145655	
			155040	
			167872	
			181758	
			182546	
	188794			

January 1955.

## BARIUM

## Ba I

56 electrons

Z=56

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 6s^2 {}^1S_0$  $6s^2 {}^1S_0$ , 42932.4 K

I. P. 5.210 volts

The Ba I spectrum, like those of Ca I and Sr I, has contributed an important share in the development of the theory of atomic spectra. Russell and Saunders in 1925 included Ba I in their discussion of terms involving two excited electrons in addition to the "regular" series. White discusses these terms further in his paper on "Auto-Ionization in the Alkaline Earth Metals and the Inert Gases."

The "regular" terms have been taken from Saunders, Paschen-Götze, and Fowler, with the revisions suggested by Shenstone and Russell in 1932. The limit is also from this later paper, which discusses in detail the perturbed series of Ba I. Whitelaw, Langstroth, and others have also investigated perturbations in this spectrum.

Sullivan and Burns have improved a number of the energy-level values from their accurate interferometric measurements of Ba I. With these as a start, the writer has prepared a line list and multiplet array, and revised the decimals of the remaining values for inclusion here. In the long-wave region she has utilized Meggers' 1933 data supplemented by his 1935 unpublished observations. Russell's unpublished material (1933) has also been placed at her disposal for this work. In this revised list there are approximately 390 classified lines between 2473 Å and 30934 Å. Observed intersystem combinations connect the singlet and triplet systems of terms.

The writer has derived the  $g$ -values in the table from the Zeeman observations given by B. E. Moore. The measurements published by Moore ( $\Delta\lambda/\lambda^2$ ) have been divided by the factor 1.09 throughout, a correction factor adopted to reduce them to the theoretical Landé patterns for a few very well-known lines. For unresolved patterns the formulas for blends given by Shenstone and Blair, as discussed by Russell (1930), have been used.

A monograph on Ba I is needed. The existing data are not homogeneous, even though the spectrum is well analyzed. The complete analysis is given in the 1955 Bureau reference below.

## REFERENCES

- B. E. Moore, *Ann. der Phys.* [4] **25**, 309 (1908). (Z E)  
 F. A. Saunders, *Astroph. J.* **51**, 23 (1920). (I P) (T) (C L)  
 F. Paschen und R. Götze, *Seriengesetze der Linienspektren* p. 90 (Julius Springer, Berlin, 1922). (I P) (T) (C L)  
 A. Fowler, *Report on Series in Line Spectra*, p. 133 (Fleetway Press, London, 1922). (I P) (T) (C L)  
 H. N. Russell and F. A. Saunders, *Astroph. J.* **61**, 38 (1925). (I P) (T) (C L)  
 H. N. Russell, *Phys. Rev.* **36**, 1590 (1930).  
 H. E. White, *Phys. Rev.* **33**, 2016 (1931).  
 A. G. Shenstone and H. N. Russell, *Phys. Rev.* **39**, 415 (1932). (I P) (T) (C L)  
 N. G. Whitelaw, *Phys. Rev.* **44**, 544 (1933).  
 G. O. Langstroth, *Proc. Roy. Soc. London [A]* **142**, 286 (1933).  
 W. F. Meggers, *Bur. Std. J. Research* **10**, 680, RP 558 (1933). (T) (C L)  
 W. F. Meggers, unpublished material (1935).  
 F. Sullivan and K. Burns, *Science Studies* **9**, No. 3, 7 (1941). (T) (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 H. N. Russell and C. E. Moore, *J. Research Nat. Bur. Std.* **55**, 299, RP2633 (1955). (I P) (T) (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 30 (1955). (Summary hfs)





Ba I—Continued

Ba I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
6s(6S)10p	10p 1P°	1	39308.74			6s(6S)12p	12p 1P°	1	40481.83		
6s(6S)9d	9d 1D	2	39334.94			6s(6S)9f	9f 1F°	3	40614.15		
6s(6S)11s	11s 1S	1	39624.60			6s(6S)9f	9f 1F°	2	40614.15	7.64	
6s(6S)7f	7f 1F°	2	39678.08	2.56		6s(6S)9f	9f 1F°	3	40681.80	5.98	
		3	39680.69	2.40				4	40687.78		
6s(6S)7f	7f 1F°	3	39705.11			6s(6S)10f	10f 1F°	2	40895.87	3.42	
5d(6D)4f	4f 1F°?	2	39785.31			6s(6S)10f	10f 1F°	3	40898.69	-1.62	
		3				6s(6S)10f	10f 1F°	4	40897.07		
		4				6s(6S)11f	11f 1F°	4	41098.50	-0.38	
6s(6S)10d	10d 1D	1	39892.19	2.34		6s(6S)11f	11f 1F°	3	41098.88	-3.37	
		2	39894.53	10.63		6s(6S)11f	11f 1F°	2	41100.85		
		3	39905.15			6s(6S)12f	12f 1F°	4	41247.58	0.54	
		1 or 2	39905.45?			6s(6S)12f	12f 1F°	3	41247.04	-3.52	
6s(6S)11p	11p 1P°	1	39984.75			6s(6S)12f	12f 1F°	2	41250.58		
6s(6S)8f	8f 1F°	2	40838.91	2.46		6s(6S)13f	13f 1F°	4	41364.65	0.29	
		3	40841.37	2.65		6s(6S)13f	13f 1F°	3	41364.38		
		4	40844.08			6s(6S)13f	13f 1F°	2	41364.38		
6s(6S)8f	8f 1F°	3	40857.63			6s(6S)14f	14f 1F°	4, 3, 2	41456.78		
6s(6S)11d	11d 1D	1	40380.32			6s(6S)15f	15f 1F°	4, 3, 2	41530.05		
		2	40382.68	2.86							
		3	40382.68			Ba II(6S <sub>1/2</sub> )	Limit		42832.4		

December 1955.

Ba I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms
6s <sup>2</sup>	6s <sup>2</sup> 1S
5d <sup>2</sup>	{ 5d <sup>2</sup> 1P 5d <sup>2</sup> 1D
6p <sup>2</sup>	{ 6p <sup>2</sup> 1S 6p <sup>2</sup> 1P 6p <sup>2</sup> 1D
	ns (n ≥ 7)
6s(6S)ns	{ 7-11s 1S 7-9s 1S
5d(6D)ns'	{ 7s' 1D 7s' 1D
	np (n ≥ 6)
6s(6S)ns	{ 6, 7p 1P° 6-12p 1P°
5d(6D)ns'	{ 6, 7p' 1P° 6, 7p' 1D° 6, 7p' 1F° 6, 7p' 1P° 6, 7p' 1D° 6, 7p' 1F°
	nd (n ≥ 5)
6s(6S)ns	{ 5-11d 1D 5-9d 1D
5d(6D)ns'	{ 4-15f 1F° 4-9f 1F° 6d' 1S 6d' 1P 6d' 1D 6d' 1F 6d' 1G 6d' 1G 4f' 1F°?

\*For predicted terms in the spectra of the Ba I isoelectronic sequence, see Volume III, Introduction.

## Ba II

(Ca I sequence; 55 electrons)

Z=56

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 6s^2 S_{0,4}$  $6s^2 S_{0,4}$ , 80686.87 K

I. P. 10.001 volts

Although the early work on this spectrum has been revised and extended, a monograph on Ba II is needed. The series are well known, but the observed wavelengths are heterogeneous and do not give precise level-values throughout.

The analysis is from Rasmussen, supplemented by additional series members reported by Saunders, Schneider, and Buckingham. Interferometric measurements have been made for a few lines of Ba II. Sullivan and Burns have utilized their observations to derive improved values for six terms. With these as a start the writer has adjusted the decimals of other terms from the wavelengths given by Rasmussen. Rounded off entries in the table are from ultra-violet measurements in the 1934 reference.

The limit is from the  $^2G$  series reported by Rasmussen. He notes that the  $^2F$  series exhibits perturbations such as are well known in Al II.

Approximately 85 lines are classified between 1397 Å and 14000 Å.

The observed  $g$ -values entered in the table for  $6s^2 S_{0,4}$  and  $6p^2 P_{0,4}$  have been determined at the National Bureau of Standards from Zeeman spectrograms on which the line at 4934 Å appeared as an impurity. The other  $g$ -values have been derived by the writer from observations of the lines at 5854 Å and 6141 Å, reported by Back; and from those given by Moore.

## REFERENCES

- B. E. Moore, *Ann. der Phys.* [4] **25**, 314 (1908). (Z E)  
 A. Fowler, *Report on Series in Line Spectra* p. 137 (Fleetway Press, London, 1922). (T) (C L)  
 E. Back, *Ann. der Phys.* [4] **70**, 360, 363 (1923). (Z E)  
 E. Rasmussen, *Zeit. Phys.* **83**, 404 (1933). (I P) (T) (C L)  
 F. A. Saunders, E. G. Schneider, and E. Buckingham, *Proc. Nat. Acad. Sci.* **20**, 291 (1934). (T) (C L)  
 F. Sullivan and K. Burns, *Sci. Studies* **9**, 7 (1941). (T) (C L)  
 J. E. Mack, *Rev. Mod. Phys.* **22**, No. 1, 64 (1950). (Summary hfs)

## Ba II

## Ba II

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
$5p^4(^1S)6s$	$6s^2 S$	0½	0.00		1.974	$5p^4(^1S)7d$	$7d^2 D$	1½ 2½	59800.31 59894.98	94.67	
$5p^4(^1S)5d$	$5d^2 D$	1½ 2½	4873.850 5674.824	800.974	0.79 1.12	$5p^4(^1S)8p$	$8p^2 P^o$	0½ 1½	61336 61636	300	
$5p^4(^1S)6p$	$6p^2 P^o$	0½ 1½	80261.568 81962.428	1690.860	0.672 1.32	$5p^4(^1S)5g$	$5g^2 G$	{ 3½ 4½ }	63026.53		
$5p^4(^1S)7s$	$7s^2 S$	0½	42355.182		1.98	$5p^4(^1S)6f$	$6f^2 F^o$	2½ 3½	64596.31 64697.08	100.77	
$5p^4(^1S)6d$	$6d^2 D$	1½ 2½	45949.496 46154.89	205.39	0.79 1.18	$5p^4(^1S)9s$	$9s^2 S$	0½	65683.41		
$5p^4(^1S)4f$	$4f^2 F^o$	2½ 3½	48258.590 48483.29	224.70		$5p^4(^1S)8d$	$8d^2 D$	1½ 2½	66673.80 66725.41	51.61	
$5p^4(^1S)7p$	$7p^2 P^o$	0½ 1½	49390.05 50011.22	621.17		$5p^4(^1S)6g$	$6g^2 G$	{ 3½ 4½ }	68425.87		
$5p^4(^1S)5f$	$5f^2 F^o$	2½ 3½	57390.93 57631.64	240.71		$5p^4(^1S)7f$	$7f^2 F^o$	2½ 3½	69211.70 69260.46	48.76	
$5p^4(^1S)8s$	$8s^2 S$	0½	58025.18			$5p^4(^1S)10s$	$10s^2 S$	0½	70014.72		

## Ba II—Continued

## Ba II—Continued

Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>
5p <sup>4</sup> ( <sup>1</sup> S)9d	9d <sup>1</sup> D	1½ 2½	70620. 40 70651. 87	31. 47		5p <sup>4</sup> ( <sup>1</sup> S)9g	9g <sup>1</sup> G	{ 3½ 4½	75244. 09		
5p <sup>4</sup> ( <sup>1</sup> S)7g	7g <sup>1</sup> G	{ 3½ 4½	71682. 38			5p <sup>4</sup> ( <sup>1</sup> S)10f	10f <sup>1</sup> F°	2½ 3½	75438 75451	13	
5p <sup>4</sup> ( <sup>1</sup> S)8f	8f <sup>1</sup> F°	2½ 3½	72142. 72 72170. 19	27. 47		5p <sup>4</sup> ( <sup>1</sup> S)12d	12d <sup>1</sup> D	1½ 2½	75945. 85		
5p <sup>4</sup> ( <sup>1</sup> S)11s	11s <sup>1</sup> S	0½	72705. 33			5p <sup>4</sup> ( <sup>1</sup> S)10g	10g <sup>1</sup> G	{ 3½ 4½	76279. 64		
5p <sup>4</sup> ( <sup>1</sup> S)10d	10d <sup>1</sup> D	1½ 2½	73101. 90 73122. 23	20. 33		5p <sup>4</sup> ( <sup>1</sup> S)11f	11f <sup>1</sup> F°	2½ 3½	76413 76429	16	
5p <sup>4</sup> ( <sup>1</sup> S)8g	8g <sup>1</sup> G	{ 3½ 4½	73795. 84			5p <sup>4</sup> ( <sup>1</sup> S)11g	11g <sup>1</sup> G	{ 3½ 4½	77046		
5p <sup>4</sup> ( <sup>1</sup> S)9f	9f <sup>1</sup> F°	2½ 3½	74091. 00 74108. 92	17. 92		5p <sup>4</sup> ( <sup>1</sup> S)12g	12g <sup>1</sup> G	{ 3½ 4½	77628		
5p <sup>4</sup> ( <sup>1</sup> S)12s	12s <sup>1</sup> S	0½	74491. 51								
5p <sup>4</sup> ( <sup>1</sup> S)11d	11d <sup>1</sup> D	1½ 2½	74765. 73 74779. 66	13. 93		Ba III( <sup>1</sup> S <sub>0</sub> )	Limit	-----	80486. 87		

February 1954.

## Ba IV

(I I sequence; 53 electrons)

Z=56

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 5s<sup>2</sup> 5p<sup>5</sup> <sup>2</sup>P<sub>1/2</sub><sup>o</sup>5p<sup>5</sup> <sup>2</sup>P<sub>1/2</sub><sup>o</sup>

K

I. P.

volts

This spectrum is very incompletely analyzed. Fitzgerald and Sawyer have reported the 22 levels in the table. These levels account for 53 classified lines; 13 between 570.1 Å and 740.0 Å, and 40 between 2220.15 Å and 4591.72 Å. The writer has labeled the third level, 5p<sup>5</sup> <sup>2</sup>S, by analogy with Xe II. The authors state that the levels in the middle group are probably from the 5s<sup>2</sup> 5p<sup>4</sup> 5d and 5s<sup>2</sup> 5p<sup>4</sup> 6s configurations, whereas the higher odd levels, 196000 K to 198000 K, are probably from 5s<sup>2</sup> 5p<sup>4</sup> 6p.

## REFERENCE

M. A. Fitzgerald and R. A. Sawyer, Phys. Rev. **46**, 576 (1934). (T) (C L)

## Ba IV

## Ba IV

Config.	Desig.	<i>J</i>	Level	Interval	Config.	Desig.	<i>J</i>	Level	Interval
5s <sup>2</sup> 5p <sup>4</sup>	5p <sup>4</sup> <sup>2</sup> P°	1½ 0½	0 17830	-17830			2½	158998	
5s <sup>2</sup> 5p <sup>4</sup>	5p <sup>4</sup> <sup>2</sup> S	0½	152963				2½	163254	
		2½	154481				1½	175431	
		3½	154631				2½	175926	
		0½ or 1½	156738				2½	196549	
		3½	157656				2½	197504	
		0½	157741				3½	197698	
		2½	158170				1½	197711	
		2½	158444				1½	198005	
		3½	158750				2½	198758	
							2½	198889	

January 1954.

## LANTHANUM

## La I

57 electrons

Z=57

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 5d 6s^2 {}^2D_{1/2}$  $\epsilon {}^2D_{1/2}$  45298 KI. P.  $5.61 \pm 0.03$  volts

The analysis is from Russell and Meggers and is based on Meggers' observations of Zeeman effects and wavelengths, which cover the range 2647.13 Å to 10954.6 Å. There are 540 classified lines. Observed intersystem combinations connect the doublet and quartet systems of terms.

The analysis is well confirmed by the 1932 Zeeman observations, but the earlier measurements have since been largely supplanted by further study at the Massachusetts Institute of Technology. All but eight of the  $g$ -values in the table are from the 1945 reference.

The limit is from three series of two members each, namely,  $6s$ ,  $7s$   ${}^4F$ ,  $6s$ ,  $7s$   ${}^2F$ , and  $6s^2$   ${}^3D$ ,  $7s$   ${}^4D$ . By comparison with Ca I, Sr I, and Ba I, the authors introduced a correction to the differences of the Rydberg denominators amounting to  $\Delta n^* = 1.060$  for the  ${}^4F$  and  ${}^2F$  series, and  $\Delta n^* = 1.010$  for the  ${}^3D$ ,  ${}^4D$  series.

The authors point out the significant fact that the electron configurations of the lanthanum atom anticipate "the atom building process which accounts for the rare-earth elements."

## REFERENCES

- W. F. Meggers, *Bur. Std. J. Research* **9**, 239, RP 468 (1932). (Z E)  
 H. N. Russell and W. F. Meggers, *Bur. Std. J. Research* **9**, 625, RP 497 (1932). (I P) (T) (C L) (G D) (Z E)  
 G. R. Harrison, N. Rosen, and J. R. McNally, Jr., *J. Opt. Soc. Am.* **35**, 658 (1945). (Z E)  
 R. A. Fisher and F. Eshbach, unpublished material (March 1954). (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 30 (1955). (Summary hfs)

## La I

## La I

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$	
$5d 6s^2$	$\alpha {}^2D$	$1\frac{1}{2}$	0.00	1053.20	0.798	$5d 6s(\alpha {}^2D)6p$	$z {}^4F^{\circ}$	$1\frac{1}{2}$	15260.36	370.72	0.93	
		$2\frac{1}{2}$	1053.20		1.198			$2\frac{1}{2}$	15651.08		1388.47	1.096
$5d^2(\alpha {}^2F)6s$	$\alpha {}^4F$	$1\frac{1}{2}$	2668.20	341.81	0.405	$5d 6s(\alpha {}^2D)6p$	$z {}^4D^{\circ}$	$0\frac{1}{2}$	14095.70	613.26	794.71	1.23
		$2\frac{1}{2}$	3010.01		1.030			$1\frac{1}{2}$	14708.96			
		$3\frac{1}{2}$	3494.58		1.239			$2\frac{1}{2}$	15505.67			
		$4\frac{1}{2}$	4121.61		1.338			$3\frac{1}{2}$	18099.28			
$5d^2(\alpha {}^2F)6s$	$\alpha {}^2F$	$2\frac{1}{2}$	7011.90	1040.25	0.893	$5d 6s(\alpha {}^2D)6p$	$z {}^2D^{\circ}$	$2\frac{1}{2}$	14804.10	-227.55	1.06	
		$3\frac{1}{2}$	8052.15		1.137			$1\frac{1}{2}$	15051.65		0.917	
$5d^2(\alpha {}^2F)6s$	$\alpha {}^4P$	$0\frac{1}{2}$	7231.36	259.10	2.648	$5d 6s(\alpha {}^2D)6p$	$z {}^2F^{\circ}$	$2\frac{1}{2}$	15196.80	1341.64	0.906	
		$1\frac{1}{2}$	7490.46		1.704			$3\frac{1}{2}$	16538.44		1.19	
		$2\frac{1}{2}$	7679.94		1.498							
$5d^2(\delta {}^2D)6s$	$\delta {}^2D$	$1\frac{1}{2}$	8446.03	737.74	0.942	$6s^2(\alpha {}^2S)6p$	$z {}^2P^{\circ}$	$0\frac{1}{2}$	15219.90	1060.39	1.329	
		$2\frac{1}{2}$	9183.77		1.250			$1\frac{1}{2}$	16280.80			
$5d^2(\alpha {}^2F)6s$	$\alpha {}^4P$	$0\frac{1}{2}$	9044.21	675.23	0.690	$5d 6s(\alpha {}^2D)6p$	$y {}^2F^{\circ}$	$2\frac{1}{2}$	16856.88	1053.36	0.810	
		$1\frac{1}{2}$	9719.44		1.23			$3\frac{1}{2}$	17910.18		1.086	
$5d^2(\alpha {}^2G)6s$	$\alpha {}^2G$	$4\frac{1}{2}$	9919.94	-41.02	1.12	$5d 6s(\alpha {}^2D)6p$	$z {}^4P^{\circ}$	$0\frac{1}{2}$	17567.56	229.74	1.175	
		$3\frac{1}{2}$	9960.96		0.91			$1\frac{1}{2}$	17797.30			
								$2\frac{1}{2}$	18157.00			

La I—Continued

La I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g		
5d <sup>2</sup> (a <sup>1</sup> F)6p	z <sup>1</sup> G°	2½	17947. 16		1. 063	5d <sup>2</sup> (b <sup>1</sup> D)6p	w <sup>1</sup> P°	0¼					
		3½	18603. 96	656. 79	1. 051			1½	27226. 27	1. 335			
		4½	19189. 34	525. 39	1. 173			5d <sup>2</sup> (b <sup>1</sup> D)6p	u <sup>1</sup> F°	2¼	27393. 00	646. 54	0. 868
		5½	20117. 40	988. 06	1. 200					3½	28039. 64	1. 13	
5d 6s(a <sup>1</sup> D)6p	y <sup>1</sup> D°	1¼	18172. 39		0. 802	4f	t <sup>1</sup> F°	2¼	27669. 38	873. 72	0. 87		
		2¼	19379. 44	1207. 05	1. 186			3½	28543. 10	1. 00			
5d 6s(a <sup>1</sup> D)6p	y <sup>1</sup> P°	1¼	20019. 00		1. 020	5d <sup>2</sup> (b <sup>1</sup> D)6p	w <sup>1</sup> D°	1¼	27968. 63	537. 86	0. 836		
		0¼	20197. 38	-178. 38	0. 60			2¼	28506. 39	1. 20			
5d <sup>2</sup> (a <sup>1</sup> F)6p	y <sup>1</sup> F°	1¼	20083. 08		0. 731	5d <sup>2</sup> (a <sup>1</sup> P)4f	w <sup>1</sup> D°	0¼	28893. 47	306. 06	0. 00		
		2¼	20358. 30	255. 28	1. 013			1¼	29199. 63	1. 15			
		3½	20763. 31	425. 01	1. 178			2¼	29502. 17	1. 263			
		4½	21364. 06	620. 75	1. 278			3½	29894. 91	1. 352			
5d <sup>2</sup> (a <sup>1</sup> F)6p	z <sup>1</sup> F°	2¼	20972. 22		0. 894	4f	e <sup>1</sup> D°	1¼	28971. 82	803. 75	0. 71		
		3½	21662. 61	690. 39	0. 995			2¼	29776. 67	1. 253			
5d <sup>2</sup> (a <sup>1</sup> F)6p	y <sup>1</sup> D°	0¼	22246. 64		-0. 227	33	6°	0¼	29461. 33				
		1¼	22439. 37	192. 73	1. 192			34?	29594. 81				
		2¼	22804. 26	364. 89	1. 362					e <sup>1</sup> F	1¼	29874. 89	479. 43
		3½	23303. 31	499. 05	1. 183			2¼	30354. 32		705. 37		
5d <sup>2</sup> (a <sup>1</sup> P)6p	z <sup>1</sup> S°	0¼	23260. 90		1. 89	3½	31059. 69	864. 21	0. 93				
		2°	23466. 85		1. 12	4¼	31923. 90						
5d <sup>2</sup> (a <sup>1</sup> P)6p	z <sup>1</sup> D°	0¼	23528. 38		0. 152	7°	1¼	29936. 73		1. 492			
		1¼	23704. 76	176. 38	1. 135			8°	1¼	30417. 47		1. 533	
		2¼	24048. 06	341. 30	1. 272					e <sup>1</sup> F°	2¼	30788. 40	176. 42
		3½	25063. 42	1037. 36	1. 379			3½	30964. 82		1. 16		
5d 6s(a <sup>1</sup> D)4f	w <sup>1</sup> F°	2¼	23876. 00		0. 957	9°	2¼	30896. 88		1. 424			
		3½	24409. 70	534. 70	1. 17			e <sup>1</sup> F	2¼	31119. 08	989. 50		
		4°	24173. 86		0. 72				3½	32108. 58			
		e <sup>1</sup> F°?	2¼	24507. 89				1. 159	e <sup>1</sup> D	0¼			
3½	25378. 46	870. 57	1. 13	1¼									
5d <sup>2</sup> (a <sup>1</sup> P)6p	z <sup>1</sup> S°	1¼	24639. 27		1. 781	2¼	31287. 65						
		3½	25378. 46		1. 13	3½							
5d <sup>2</sup> (a <sup>1</sup> F)6p	z <sup>1</sup> D°	1¼	24762. 62		0. 855	4f	r <sup>1</sup> F°	2¼	31477. 16	663. 44	0. 92		
		2¼	25218. 25	455. 63	1. 244			3½	32140. 60	1. 16			
5d <sup>2</sup> (a <sup>1</sup> F)4f	z <sup>1</sup> F°	1¼	24910. 39		0. 724	5d <sup>2</sup> (a <sup>1</sup> P)6p	u <sup>1</sup> D°	1¼	31761. 68	741. 12	0. 80		
		2¼	24984. 33	73. 94	1. 063			2¼	32492. 80	1. 145			
		3½	25380. 33	396. 00	1. 228			e <sup>1</sup> P°	0¼	32290. 26	913. 95	0. 70	
		4½	25997. 27	616. 94	1. 319				1¼	33204. 20	1. 27		
5d <sup>2</sup> (a <sup>1</sup> G)6p	z <sup>1</sup> H°	4¼	25089. 50		0. 94	5d <sup>2</sup> (a <sup>1</sup> G)4f	y <sup>1</sup> H°	4¼	32410. 76	107. 36	0. 92		
		5½	25874. 68	785. 18	1. 11			5½	32518. 12	1. 11			
5d 6s(a <sup>1</sup> D)6p	z <sup>1</sup> P°	0¼	25453. 92		0. 985	35			33286. 50				
		1¼	25960. 39	496. 47	1. 433			10°	36722. 38				
5d <sup>2</sup> (a <sup>1</sup> P)6p	y <sup>1</sup> P°	0¼	25616. 90		2. 29	11°	3½?		37731. 90		1. 187		
		1¼	25643. 08	26. 12	1. 600			12°?	39597. 58				
		2¼	26338. 90	695. 88	1. 524					13°	39631. 27		
5d <sup>2</sup> (a <sup>1</sup> F)4f	y <sup>1</sup> G°	2¼	27022. 60		0. 57	14°			40322. 46				
		3½	27455. 34	432. 74	0. 976			15°	40343. 40				
		4½	28089. 18	633. 84	1. 159								
		5½	28743. 21	654. 03	1. 28								
5d <sup>2</sup> (a <sup>1</sup> G)6p	y <sup>1</sup> G°	3¼	27132. 50		0. 94	-----			45293				
		4¼	27619. 69	487. 19	1. 12	-----							
						La II(a <sup>1</sup> F <sub>2</sub> )	Limit						

La I Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>0</sup> 4p <sup>0</sup> 4d <sup>0</sup> 5s <sup>0</sup> 5p <sup>0</sup> +	Observed Terms		
	ns (n ≥ 6)	np (n ≥ 6)	n'(n' ≥ 4)
5d 6s <sup>2</sup>	a <sup>3</sup> D		
5d <sup>2</sup> (a <sup>3</sup> F) <sub>ns</sub>	a, e <sup>3</sup> F a, e <sup>3</sup> F	y <sup>1</sup> D <sup>o</sup> y <sup>3</sup> F <sup>o</sup> z <sup>1</sup> D <sup>o</sup> z <sup>3</sup> F <sup>o</sup>	z <sup>1</sup> F <sup>o</sup> y <sup>1</sup> G <sup>o</sup>
5d 6s(a <sup>1</sup> D) <sub>ns</sub>		y <sup>3</sup> P <sup>o</sup> z <sup>1</sup> D <sup>o</sup> z <sup>3</sup> F <sup>o</sup>	
5d 6s(a <sup>3</sup> D) <sub>ns</sub>	e <sup>1</sup> D	z <sup>1</sup> D <sup>o</sup> z <sup>3</sup> F <sup>o</sup> y <sup>1</sup> D <sup>o</sup> y <sup>3</sup> F <sup>o</sup>	w <sup>1</sup> F <sup>o</sup>
5d <sup>2</sup> (a <sup>3</sup> P) <sub>ns</sub>	a <sup>3</sup> P a <sup>3</sup> P	z <sup>1</sup> S <sup>o</sup> y <sup>1</sup> P <sup>o</sup> z <sup>1</sup> D <sup>o</sup> z <sup>3</sup> S <sup>o</sup> z <sup>3</sup> P <sup>o</sup> u <sup>1</sup> D <sup>o</sup>	w <sup>1</sup> D <sup>o</sup>
6s <sup>2</sup> (a <sup>1</sup> S) <sub>ns</sub>		s <sup>1</sup> P <sup>o</sup> ?	
5d <sup>2</sup> (a <sup>1</sup> G) <sub>ns</sub>	a <sup>3</sup> G	w <sup>1</sup> P <sup>o</sup> w <sup>1</sup> D <sup>o</sup> e <sup>3</sup> F <sup>o</sup> u <sup>3</sup> F <sup>o</sup>	y <sup>3</sup> G <sup>o</sup> z <sup>3</sup> H <sup>o</sup>
5d <sup>2</sup> (b <sup>1</sup> D) <sub>ns</sub>	b <sup>1</sup> D		y <sup>3</sup> H <sup>o</sup>

\*For predicted terms in the spectra of the La I isoelectronic sequence, see Vol. III, Introduction.

## La II

(Ba I sequence; 56 electrons)

 $Z=57$ Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 5d^2 \text{ } ^3\text{F}_2$  $a \text{ } ^3\text{F}_2$ ; 92240 KI. P.  $11.43 \pm 0.07$  volts

Meggers' observations from 2142.81 Å to 10954.6 Å have resulted in the classification of 728 lines. The extensive analysis by Russell and Meggers is of particular interest because the study of La I, La II, and La III "shows that the atom building process which accounts for the rare-earth elements is actually anticipated in the electron configurations of the lanthanum atom." In addition, La II "is the most completely developed example of a 2-electron spectrum which has yet been investigated;" at least ten types of configurations are known "and almost all of the terms arising from each."

The Zeeman observations available in 1932 were extremely useful in assigning the term designations, but have since been improved and extended at the Massachusetts Institute of Technology. The  $g$ -values in the table are from the 1945 paper.

The dubious  $^1\text{D}_2$  term at 64706.767 has been rejected.

The limit has been derived from the  $ns \text{ } ^1\text{D}$  series ( $n=6,7$ ) having the limit  $5d \text{ } ^1\text{D}$  in La III, and the  $nd \text{ } ^3\text{D}$  series ( $n=5,6$ ) having the limit  $6s \text{ } ^3\text{S}$ . The term  $a \text{ } ^3\text{D}$  is used in both, since  $a \text{ } ^1\text{D}$  and  $a \text{ } ^3\text{D}$ , belong to both the  $5d(^3\text{D})6s$  and the  $6s(^3\text{S})5d$  configurations. In determining the limit the authors have assumed the difference in the Rydberg denominators,  $\Delta n^* = 1.05$  for a running  $s$ -electron and 1.15 for a running  $d$ -electron, by analogy with Ca I, Sr I, and Ba I.

The authors ascribe two triads  $^3\text{P}^\circ\text{D}^\circ\text{F}^\circ$  and  $^1\text{P}^\circ\text{D}^\circ\text{F}^\circ$  to "6p 5d" electrons. In assigning limits these electrons could belong with either of two limits, namely,  $6p(^2\text{P}^\circ)5d$  or  $5d(^2\text{D})6p$ . Since the  $^3\text{D}$  limit term is the ground term in La III and the  $^2\text{P}^\circ$  limit is 42014 K above the ground term, the triads are here entered with the lower limit.

Observed intersystem combinations connect the triplet and singlet systems of terms.

## REFERENCES

- W. F. Meggers and K. Burns, J. Opt. Soc. Am. and Rev. Sci. Instr. **14**, 449 (1927). (hfs)  
 W. F. Meggers, Bur. Std. J. Research **9**, 239, RP 468 (1932). (Z E)  
 H. N. Russell and W. F. Meggers, Bur. Std. J. Research **9**, 625, RP 497 (1932). (I P) (T) (C L) (G D) (Z E)  
 G. R. Harrison, N. Rosen, and J. R. McNally, Jr., J. Opt. Soc. Am. **35**, 658 (1945). (T) (Z E)  
 K. Murakawa, J. Phys. Soc. Japan **9**, 391 (1954). (hfs)

## La II

## La II

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$		
$5d^2$	$a \text{ } ^3\text{F}$	2	0.00		0.721	$5d^2$	$a \text{ } ^1\text{G}$	4	7473.32		1.000		
		3	1016.10	1016.10	1.083			$5d^2$	$b \text{ } ^1\text{D}$	2	10094.86		1.005
		4	1970.70	954.60	1.248					$6s(^3\text{S})4f$	$z \text{ } ^3\text{F}^\circ$	2	14147.98
$5d(^2\text{D})6s$	$a \text{ } ^1\text{D}$	2	1394.46		0.977	3	14375.17	1.056					
		3	3250.35	696.45 658.75	0.498 1.133 1.334	4	15698.74	1.247					
$5d^2$	$a \text{ } ^3\text{D}$	1	1895.15		0.498	$6s(^3\text{S})4f$	$z \text{ } ^1\text{F}^\circ$	3	15773.77		1.017		
		2	2591.60		1.133			$5d(^2\text{D})4f$	$z \text{ } ^1\text{G}^\circ$	4	16599.17		0.969
		3	3250.35		1.334					$5d(^2\text{D})4f$	$y \text{ } ^3\text{F}^\circ$	2	17211.93
$5d^2$	$a \text{ } ^3\text{P}$	0	5249.70		0/0	3	18235.56	1.086					
		1	5718.12	468.42	1.497	4	19214.54	1.232					
$6s^2$	$a \text{ } ^1\text{S}$	0	7394.57		0/0								

## La II—Continued

## La II—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g						
5d(°D)4f	z 'H°	4	17885. 68	754. 79	0. 846	5d(°D)6d	f 'G	3	52857. 88	475. 49	0. 861						
		5	18580. 41	1169. 21	1. 017			4	53333. 37			1. 086					
		6	19749. 68		1. 1787			5	54434. 65			1. 21					
5d(°D)4f	z 'D°	2	18895. 41		0. 923	5d(°D)6d	e 'P	1	53302. 56		1. 335						
5d(°D)4f	z 'G°	3	20408. 88	928. 78	0. 757	5d(°D)6d	f 'F	2	53885. 24	954. 80	0. 751						
		4	21331. 60	951. 30	1. 049			3	54840. 04			1. 088					
		5	22282. 90		1. 197			4	55321. 35			1. 136					
5d(°D)4f	z 'D°	1	21441. 73	664. 29	0. 542	5d(°D)6d	e 'S	1	54365. 80		1. 455						
		2	22108. 08	431. 28	1. 167			5d(°D)6d	e 'S	0	54793. 82						
		3	22837. 30		1. 288					5d(°D)6d	e 'P	0	54964. 197	266. 14	1. 552		
5d(°D)4f	z 'P°	1	22705. 15	21. 45	1. 431	5d(°D)6d	e 'P	1	55230. 33	806. 27	1. 203						
		2	23248. 93	541. 78	1. 459			2	56036. 60								
		0	22683. 70		0/0			4f <sup>2</sup>	e 'H			4	55107. 25	874. 84	0. 883		
5d(°D)6p	y 'D°	2	24462. 68		0. 887	5d(°D)6d	g 'D	2	55184. 05		1. 183						
5d(°D)4f	y 'F°	3	24522. 70		1. 034	5d(°D)6d	f 'G	4	56035. 70		1. 027						
5d(°D)6p	y 'D°	1	25973. 37	1414. 74	0. 782			1°	3	57364. 18		1. 07					
		2	27388. 11	927. 14	1. 168				4f <sup>2</sup>	g 'F	2	57399. 58	518. 92	0. 675			
		3	28315. 25		1. 308	3	57918. 50				1. 085						
5d(°D)6p	z 'F°	2	28414. 01	423. 65	0. 825	4f <sup>2</sup>	g 'F	4	58259. 41	340. 91	1. 196						
		3	28537. 68	1727. 74	1. 088			2°	4			58748. 90					
		4	28586. 40		1. 245			4f <sup>2</sup>	g 'G			4	59527. 60		1. 046		
5d(°D)4f	z 'P°	1	27423. 91		0. 876	4f <sup>2</sup>	3°	2, 3	59612. 64								
5d(°D)6p	y 'P°	0	27545. 85	608. 70	0/0			4f <sup>2</sup>	h 'D	2	59900. 08		1. 035				
		1	28154. 55	1343. 50	1. 267					6p <sup>2</sup>	f 'P	0	60094. 84	1033. 99	0/0		
5d(°D)4f	z 'H°	5	28525. 71		1. 004	6p <sup>2</sup>	f 'P	1	61128. 83	1377. 53	1. 528						
		5d(°D)6p	y 'P°	1	30353. 33				1. 074			2	62506. 36	1. 416			
				6s(°S)6p	z 'P°			0	31785. 22			375. 17	0/0	4°	4	60744. 17	1. 25
5d(°D)6p	z 'F°	1	32160. 99	1043. 42	1. 492	4°	5°	3, 4	61017. 68								
		2	32204. 41		1. 494			6°	3, 4	61614. 46		0. 977					
		3	32201. 05		1. 005					6p(°P°)4f	e 'F	2	35787. 53	1167. 12	0. 719		
6p(°P°)4f	e 'G	3	35452. 66	1720. 13	0. 876	6p(°P°)4f	e 'I	3	36954. 65	835. 92	1. 061						
		4	37172. 79	1845. 95	1. 127			4f <sup>2</sup>	g 'P			0	63463. 95	239. 23	0/0		
		5	39018. 74		1. 21							1	63703. 18			1. 471	
6p(°P°)4f	e 'F	2	35787. 53	1167. 12	0. 719	4f <sup>2</sup>	g 'P	2	64278. 92	575. 74	1. 414						
		3	36954. 65	835. 92	1. 113			7°	4			63598. 87		1. 205			
		4	37790. 57		1. 113							6s(°S)6d	h 'D	1	64361. 28	168. 62	0. 506
6p(°P°)4f	e 'F	3	37209. 71		0. 944	6s(°S)6d	h 'D	2	64529. 90	162. 69	1. 217						
		6p(°P°)4f	e 'D	1	38534. 11			-312. 62	0. 497			8°	3	64411. 17		0. 97	
				2	38221. 49			1181. 06	1. 071					6p <sup>2</sup>	f 'S	0	66591. 91
6p(°P°)4f	e 'D	3	39402. 55		1. 274	6p(°P°)7p?	i 'D	1		69233. 90	1. 414						
		6p(°P°)4f	e 'G	4	39221. 65				1. 059			2	3	69233. 90	1. 205		
				6p(°P°)4f	e 'D			2	40457. 71								1. 036
6s(°S)6p	z 'P°	1	45692. 17		0. 999	4f <sup>2</sup>	g 'S	0	69505. 06								
5d(°D)7s	f 'D	1	49733. 13	151. 22	0. 500							La III(°D <sub>14</sub> )	Limit	-----	92240		
		2	49884. 35	1344. 22	1. 117												
		3	51228. 57		1. 315												
5d(°D)7s	f 'D	2	51523. 86		1. 036												
5d(°D)6d	f 'F	3	52137. 67		0. 987												
5d(°D)6d	g 'D	1	52169. 66	565. 15	0. 621												
		2	52734. 81	954. 75	1. 154												
		3	53689. 56		1. 218												

March 1954.



## La II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>n</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>n</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms	
5d <sup>1</sup>	{ a <sup>3</sup> P    b <sup>1</sup> D    a <sup>3</sup> F    a <sup>1</sup> G	
6d <sup>1</sup>	{ a <sup>3</sup> S	
6p <sup>1</sup>	{ f <sup>1</sup> S    f <sup>3</sup> P    i <sup>1</sup> D	
4f <sup>1</sup>	{ g <sup>1</sup> S    g <sup>3</sup> P    h <sup>1</sup> D    g <sup>3</sup> F    g <sup>1</sup> G    e <sup>3</sup> H    e <sup>1</sup> I	
	ns (n ≥ 6)	np (n ≥ 6)
5d(2D)ns	{ a, f <sup>3</sup> D a, f <sup>1</sup> D	y <sup>3</sup> P°    y <sup>3</sup> D°    z <sup>3</sup> F° y <sup>1</sup> P°    y <sup>1</sup> D°    z <sup>1</sup> F°
6s(2S)ns'	{	z <sup>3</sup> P° z <sup>1</sup> P°
6p(2P°)ns''	{	i <sup>1</sup> D?
	nd (n ≥ 5)	nf (n ≥ 4)
5d(2D)ns	{ e <sup>3</sup> S    e <sup>3</sup> P    g <sup>3</sup> D    f <sup>3</sup> F    f <sup>3</sup> G e <sup>1</sup> S    e <sup>1</sup> P    g <sup>1</sup> D    f <sup>1</sup> F    f <sup>1</sup> G	z <sup>3</sup> P°    z <sup>3</sup> D°    y <sup>3</sup> F°    z <sup>3</sup> G°    z <sup>3</sup> H° z <sup>1</sup> P°    z <sup>1</sup> D°    y <sup>1</sup> F°    z <sup>1</sup> G°    z <sup>1</sup> H°
6s(2S)ns'	{                    h <sup>3</sup> D	z <sup>3</sup> F° z <sup>1</sup> F°
6p(2P°)ns''	{	e <sup>3</sup> D    e <sup>3</sup> F    e <sup>3</sup> G e <sup>1</sup> D    e <sup>1</sup> F    e <sup>1</sup> G

\*For predicted terms in the spectra of the Ba I isoelectronic sequence, see Vol. III, Introduction.

## La III

(Ca I sequence; 55 electrons)

Z=57

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6 5d^1 {}^2D_{3/2}$  $5d {}^2D_{3/2}$  154630 K

I. P. 19.17 volts

The analysis is from Russell and Meggers, who have extended the earlier work on La III. Lang has added the  $5f {}^2F^\circ$  and  $8s {}^2S$  terms from his observations in the short-wave region. Fifteen lines between 1081.61 Å and 3517.14 Å have been classified. The limit is from the  $6, 7s {}^2S$  series. In deriving it, Russell and Meggers have assumed that  $\Delta n^* = 1.050$ , by comparison with the run of Rydberg denominators in related spectra from K I thru Ba II.

Gibbs and Schoepfle have reported that from additional observations between 500 Å and 2100 Å they have extended the  $ns {}^2S$  and  $nd {}^2D$  series to six members, and located more  ${}^2P^\circ$  and possibly  ${}^2F^\circ$  and  ${}^2G$  terms. They also confirm the above limit. Unfortunately, their new terms are not available for inclusion here.

## REFERENCES

- H. N. Russell and W. F. Meggers, Bur. Std. J. Research **9**, 625, RP 497 (1932). (I P) (T) (C L) (Z E)  
 R. J. Lang, Canadian J. Research [A] **13**, 4 (1935). (C L)  
 R. C. Gibbs and G. K. Schoepfle, Phys. Rev. **45**, 747 (A) (1934).  
 J. E. Mack, Rev. Mod. Phys. **32**, No. 1, 64 (1950). (Summary hfs)

## La III

Config.	Desig.	J	Level	Interval	Obs. g
$5p^6({}^1S)5d$	$5d {}^2D$	$1\frac{1}{2}$	0.00	1603.23	
		$2\frac{1}{2}$	1603.23		
$5p^6({}^1S)6s$	$6s {}^2S$	$0\frac{1}{2}$	13590.76		2.10
$5p^6({}^1S)6p$	$6p {}^2P^\circ$	$0\frac{1}{2}$	42014.92	3095.72	0.63
		$1\frac{1}{2}$	45110.64		
$5p^6({}^1S)7s$	$7s {}^2S$	$0\frac{1}{2}$	82345.0		
$5p^6({}^1S)6d$	$6d {}^2D$	$1\frac{1}{2}$	82378.75	433.76	
		$2\frac{1}{2}$	82812.51		
$5p^6({}^1S)5f$	$5f {}^2F^\circ$	$2\frac{1}{2}$	92454.6	80.0	
		$3\frac{1}{2}$	92534.6		
$5p^6({}^1S)8s$	$8s {}^2S$	$0\frac{1}{2}$	113496		
La IV ( ${}^1S_0$ )	Limit	-----	154630		

April 1954.

## HAFNIUM

## Hf I

72 electrons

Z=72

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^2 6s^2 {}^3F_3$  $6s^2 {}^3F_3$ 

K

I. P. volts

The analysis is by Meggers, who has made a supreme effort to submit his results for inclusion here just as press copy for this Volume is being concluded. The work is not completed. He plans to make further observations in the shortwave region and to continue the analysis with the aid of electronic computers.

The present line list extends from 1917.73 Å to 12043.08 Å, and is based on recent observations made at the National Bureau of Standards, by using microwave-excited quartz tubes containing hafnium iodide, as light source. The Zeeman Observations (2600 Å to 5000 Å) are from spectrograms taken at the Massachusetts Institute of Technology, supplemented by observations (to 8700 Å) at the National Bureau of Standards. At present there are approximately 2000 classified lines.

Meggers has pointed out the unique character of the Hf I energy levels. The two sets of levels "odd" and "even" overlap throughout their entire range. This dovetailing of the two sets is so pronounced that it causes confusion if all levels are arranged in numerical order. Consequently, in the table the two sets are listed separately, with the "even" levels preceding the "odd" levels.

The large number of "even" levels for which no term designations can as yet be assigned suggests that the configurations  $5d^4$ ,  $5d^2 6s 6d$ ,  $5d^2 6s 7s$ , and possibly  $5d^3 6p^2$  are all represented.

No series can be established with certainty from the existing data. By comparison with neighboring spectra it appears likely that the ionization potential may be near 7 volts, thus placing the limit near 56500 K.

## REFERENCES

- W. F. Meggers and B. F. Scribner, *Bur. Std. J. Research* **4**, 169, RP139 (1930). (T) (C L)  
 W. F. Meggers, *J. Opt. Soc. Am.* **44**, 348 (A) (1954). (Z E)  
 D. R. Speck and F. A. Jenkins, *Phys. Rev.* (2) **101**, 1831 (L) (1956). (hfs)  
 W. F. Meggers, unpublished material (June 1957). (T) (C L) (Z E)

## Hf I

## Hf I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
$5d^3 6s^2$	$a {}^3F$	2	0.00		0.695	$5d^3 6s^2$	$a {}^1G$	4	10532.54		1.008
		3	2356.68	2356.68	1.083						
		4	4567.64	2210.96	1.240						
$5d^3 6s^2$	$a {}^3P$	0	5521.78		0.000	$5d^3 (b {}^4F) 6s$	$a {}^3F$	1	14092.28	648.40	0.00
		1	6572.55	1050.77	1.500			2	14740.68	932.65	1.00
		2	8983.75	2411.20	1.300			3	15873.33	1093.27	1.25
				4	16766.60			1134.68	1.36		
				5	17901.28				1.40		
$5d^3 6s^2$	$a {}^1D$	2	5638.62		1.165	$5d^3 (b {}^4P) 6s$	$a {}^3P$	1	20784.87	123.55	
								2	20908.42	1290.66	1.74
								3	22199.08		

Hf I—Continued

Hf I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
5P(b-F)6e	b-F	2	23327. 71	-447. 47 372. 57	0. 90			3	42360. 90		1. 15
		3	23380. 24					4	42670. 81		1. 24
		4	23352. 81								
5P( )6e		5	24085. 14					2	42719. 58		1. 19
		2	25084. 14					1	42782. 55		1. 53
		3	25281. 81					2	43016. 47		
		3	25678. 60					3	43489. 28		
		3	26715. 37					1	43561. 76		1. 47
		1	26918. 14					4	43838. 30		1. 30
		4	27074. 50					5	44131. 99		1. 40
		2	28200. 54					2	44375. 42		1. 35
		1	28527. 97					4	44777. 50		1. 25
		2	30146. 40					3	45082. 28		
		3	31054. 64			1. 09		2	45254. 91		
		2	31119. 20			0. 89		1	45270. 13		
		2	31619. 97			1. 02		3	45364. 84		
		1	32091. 59			1. 16		1	45474. 60		1. 41
		1	34991. 64			0. 50		3	45649. 36		1. 18
		2	35115. 28			1. 11		2	45728. 62		
		3	36523. 84			1. 26		3	45994. 60		
		2	37038. 83			1. 06?		2	46056. 21		
		1	37336. 20			0. 48		2	46183. 08		
		2	38289. 38			1. 09		3	46280. 26		
		3	39127. 61			1. 31		4	46282. 34		
		1	39286. 19			0. 22		2	46437. 72		
		1	39546. 34			0. 81		2	46860. 75		
		2	39788. 89			1. 02		2	46961. 96		
		1	39790. 70			0. 58		3	46966. 61		
		0	39848. 87					4	46992. 38		
		2	40130. 23			1. 21		2	47304. 97		
2	40513. 55			1. 11		3	47345. 93				
1	40618. 62			1. 50		4	47606. 21				
2	40636. 26			1. 07?		2	47702. 41				
3	40882. 06			1. 23		1	47874. 44				
2	41211. 03			1. 45		3	47938. 44				
2	41298. 36			0. 83		4	48022. 20				
3	41475. 84			1. 27?		3	48049. 43				
3	41739. 36			1. 33		1	48129. 57				
3	42061. 59					3	48319. 05				
1	42111. 70					4	48356. 90				
4	42175. 58			1. 30?		2	48407. 27				
4	42283. 48			1. 34		2	48594. 84				

## Hf I—Continued

## Hf I—Continued

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$
		3	48598.99			5d <sup>6</sup> 6s(b <sup>3</sup> D)6p	z <sup>1</sup> D <sup>o</sup>	2	31610.80		1.00
		2	48708.40					2	31945.84		
		3	48809.77			5d <sup>6</sup> 6s(a <sup>3</sup> F)6p	z <sup>3</sup> F <sup>o</sup>	1	31958.25	1169.23	2.080
		3	48869.43					2	33121.48	17.64	1.345
		4	49005.35					3	33159.18		1.442
		5	49540.65					3	32553.30		0.950
		4	49623.40					1	33157.87		0.775
		3	49791.86					2	33558.15		0.822
		4	50393.21					3	33949.28		1.09
		5	51408.65					4	33994.71		1.04
								2	33994.86		0.93
5d <sup>6</sup> 6s(b <sup>3</sup> D)6p	z <sup>1</sup> D <sup>o</sup>	1	14017.83	2145.53	0.55			1	34596.45		0.972
		2	16163.36	2218.13	1.17			4	34806.89		0.997
		3	18381.49		1.29			3	34877.04		0.960
5d <sup>6</sup> 6s(b <sup>3</sup> D)6p	z <sup>3</sup> F <sup>o</sup>	2	14435.13	106.55	0.89			2	34947.95		1.16
		3	14541.68	3683.31	1.08			1	35884.33		1.055
		4	18824.99		1.24			3	35453.83		1.002
5d <sup>6</sup> 6s(a <sup>3</sup> F)6p	z <sup>3</sup> G <sup>o</sup>	2	18011.05	1281.64	0.40			5	35993.70		
		3	19292.69	1667.40	0.95			4	36075.02		1.07
		4	20960.09	1941.25	1.16			3	36237.34		1.042
		5	22901.34		1.25			5	36419.26		
		6						3	36609.85		1.255
5d <sup>6</sup> 6s(b <sup>3</sup> D)6p	z <sup>3</sup> P <sup>o</sup>	0	18270.11	-126.74	0.00			2	36772.85		1.156
		1	18143.37	1647.93	1.43			4	36850.03		1.09
		2	19791.30		1.41			1	36949.29		0.53
5d <sup>6</sup> 6s(a <sup>3</sup> F)6p	z <sup>3</sup> F <sup>o</sup>	1	21738.70	711.86	0.04			4	37269.82		
		2	22450.56	998.05	1.00			1	38151.71		
		3	23448.61	1336.62	1.20			2	38325.47		1.154
		4	24785.23	1417.30	1.33			3	38407.80		1.29
		5	26202.53		1.36			5	38845.44		
5d <sup>6</sup> 6s(b <sup>3</sup> D)6p	z <sup>1</sup> F <sup>o</sup>	3	23644.74		1.075			4	38987.83		1.14
5d <sup>6</sup> 6s(a <sup>3</sup> F)6p	z <sup>3</sup> D <sup>o</sup>	0	24966.78	227.69	0.00			3	39081.86		
		1	25194.47	439.74	1.45			3	39193.91		1.29
		2	25634.21	671.57	1.42			2	39435.20		1.367
		3	26305.78	1210.63	1.400			1	39704.41		0.625
		4	27516.41		1.475			2	40194.46		0.510
5d <sup>6</sup> 6s(b <sup>3</sup> D)6p	z <sup>1</sup> P <sup>o</sup>	1	26465.93		1.000			2	40267.35		1.08
5d <sup>6</sup> 6s(a <sup>3</sup> F)6p	y <sup>3</sup> F <sup>o</sup>	2	27149.64	504.68	1.00			1	40704.12		1.15
		3	27654.32	3078.90	1.153			3	40767.44		1.22
		4	30733.22		1.28			2, 37	40957.54		
5d <sup>6</sup> 6s(a <sup>3</sup> P)6p	y <sup>3</sup> P <sup>o</sup>	0	28526.19	-992.44	0.00			1	41193.97		
		1	27533.75	1867.69	1.266			3	41422.43		
		2	29401.44		1.75						
5d <sup>6</sup> 6s(a <sup>3</sup> P)6p	z <sup>3</sup> S <sup>o</sup>	1	28047.91		1.94						
5d <sup>6</sup> 6s(a <sup>3</sup> F)6p	y <sup>3</sup> D <sup>o</sup>	1	28790.25	-522.86	0.96						
		2	28267.39	1729.42	1.145						
		3	29996.81		1.23						
5d <sup>6</sup> 6s(a <sup>3</sup> F)6p	z <sup>3</sup> G <sup>o</sup>	3	28583.69	662.96	0.95						
		4	29246.65	1730.34	1.077						
		5	30976.99		1.21						
5d <sup>6</sup> 6s(a <sup>3</sup> P)6p	y <sup>3</sup> D <sup>o</sup>	0									
		1									
		2	29752.84	1589.67	1.50						
		3	31342.51	600.80	1.39						
		4	31945.31		1.41						

Hf I—Continued

Hf I—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
		2	41768.67		0.927			3	45112.22		
		3	41824.58					2	45138.88		
		1	42036.37		1.23			3	45455.15		
		2	42302.12		0.877			1	45799.48		
		1	42485.50		0.86			2	46170.12		
		4	42871.02		1.15			3	46775.98?		
		3	43104.71		1.09			2	46981.20		
		2	43517.85					3	47590.09		
		1	43786.58		0.17			4	48047.90		
		1	43844.63					2	48951.97		
		1	44257.54								

June 1957.

Hf I OBSERVED TERMS\*

1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms		
5d <sup>2</sup> 6s <sup>2</sup>	{ a <sup>3</sup> P a <sup>1</sup> D a <sup>3</sup> F a <sup>1</sup> G		
	ns (n ≥ 6)	np (n ≥ 6)	
5d <sup>2</sup> 6s(a <sup>3</sup> F)nx	{	z <sup>3</sup> D <sup>o</sup> y <sup>3</sup> D <sup>o</sup>	z <sup>3</sup> F <sup>o</sup> y <sup>3</sup> F <sup>o</sup> z <sup>3</sup> G <sup>o</sup> y <sup>3</sup> G <sup>o</sup>
5d <sup>2</sup> 6s(a <sup>3</sup> P)nx	{	z <sup>3</sup> P <sup>o</sup> y <sup>3</sup> D <sup>o</sup>	
5d <sup>2</sup> 6s(b <sup>3</sup> D)nx	{	z <sup>3</sup> P <sup>o</sup> z <sup>3</sup> P <sup>o</sup>	z <sup>3</sup> D <sup>o</sup> z <sup>3</sup> D <sup>o</sup> z <sup>3</sup> F <sup>o</sup> z <sup>3</sup> F <sup>o</sup>
5d <sup>2</sup> 6s(a <sup>3</sup> P)nx	{	z <sup>3</sup> S <sup>o</sup> y <sup>3</sup> P <sup>o</sup>	
5d <sup>2</sup> (b <sup>3</sup> F)nx	{ a <sup>3</sup> F b <sup>3</sup> F		
5d <sup>2</sup> (b <sup>3</sup> P)nx	a <sup>3</sup> P		

\*For predicted terms in the spectra of the Hf I isoelectronic sequence, see Vol. III, Introduction.

Hf II

(Lu I sequence; 71 electrons)

Z=72

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d 6s<sup>2</sup> <sup>2</sup>D<sub>11/2</sub>

a<sup>3</sup>D<sub>11/2</sub> 120000 K

I. P. 14.9 volts

The analysis is by Meggers, who has reobserved the spectrum, revised the level values, and extended the 1934 analysis to include approximately 950 classified lines. The new observations extend from 2005.36 A to 8804.35 A. Observed intersystem combinations connect the terms of different multiplicity.

## Hf II—Continued

The observed  $g$ -values are from spectrograms taken by Meggers in 1949 at the Massachusetts Institute of Technology, through the courtesy of G. R. Harrison.

In 1954 Gehatia reported a theoretical study of the low even configurations of Hf II. The agreement between his theoretical and the experimental energy level- and  $g$ -values is excellent. His predictions have enabled Meggers to find the energy levels  $c^2D_{1/2}$  and  $b^2P_{3/2}$  listed in the table, and will doubtless serve to locate the high  $5d^2^2D$  term, when new observations of Hf II in the vacuum region become available.

Although Hf II belongs in the Lu I isoelectronic sequence so far as atomic number is concerned, it does not have the structure of a rare-earth spectrum, to which  $f$ -electrons contribute. Meggers points out the "remarkable parallelism between the Hf II and La I spectra." From this analogy, together with the assumption that  $e^2D$  and  $e^4F$  form series with  $a^2D$  and  $a^4F$ , respectively, he derives the approximate value of the limit quoted here. He plans to extend the observations to shorter waves in the near future, with the hope of improving the series.

## REFERENCES

- W. F. Meggers and B. F. Scribner, J. Research Nat. Bur. Std. **13**, 625, RP 732 (1934). (I P) (T) (C L) (Z E) (G D)  
 M. Gehatia, Phys. Rev. **94**, 618 (1954). (T)  
 W. F. Meggers, unpublished material (January 1957). (T) (C L) (Z E)

## Hf II

## Hf II

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$			
$5d\ 6s^2$	$a^2D$	$1\frac{1}{2}$	0.00	3050.88	0.787	$5d^2$	$a^2H$	$5\frac{1}{2}$	30941.97	-935.77	1.10			
		$2\frac{1}{2}$	3050.88		1.173			$4\frac{1}{2}$	31877.74		1.05			
$5d^2(a^2F)6s$	$a^4F$	$1\frac{1}{2}$	3644.65	1260.20	0.425	$5d\ 6s(a^2D)6p$	$z^2P^\circ$	$0\frac{1}{2}$	33136.20	3237.22	0.217			
		$2\frac{1}{2}$	4904.85		1.052			$1\frac{1}{2}$	36373.42		1.323			
		$3\frac{1}{2}$	6344.34		1.236			$5d\ 6s(a^2D)6p$	$z^2D^\circ$		$2\frac{1}{2}$	33180.92	1.036	
		$4\frac{1}{2}$	8361.76		1.328						$1\frac{1}{2}$	34123.93	0.800	
$5d^2(a^2P)6s$	$a^4P$	$0\frac{1}{2}$	11951.70	969.24	2.598	$5d^2(a^2F)6p$	$z^4G^\circ$	$2\frac{1}{2}$	34942.36	3556.17	0.981			
		$1\frac{1}{2}$	12920.94		1.664			$3\frac{1}{2}$	38498.53		1.044			
		$2\frac{1}{2}$	13485.56		1.410			$4\frac{1}{2}$	42591.09		1.168			
$5d^2(a^2F)6s$	$a^4F$	$2\frac{1}{2}$	12070.46	3013.80	0.964	$5d^2$	$b^2P$	$0\frac{1}{2}$	35146.73	2177.49	0.74			
		$3\frac{1}{2}$	15084.26		1.10			$1\frac{1}{2}$	37324.22		1.28			
$5d^2(b^2D)6s$	$b^2D$	$1\frac{1}{2}$	14359.42	3009.45	1.034	$5d^2$	$b^2F$	$2\frac{1}{2}$	37398.44	41.22	1.14			
		$2\frac{1}{2}$	17368.87		1.273			$3\frac{1}{2}$	37439.66		1.14			
$5d^2(a^2P)6s$	$a^2P$	$0\frac{1}{2}$	15254.29	2576.05	0.737	$5d\ 6s(a^2D)6p$	$y^2D^\circ$	$1\frac{1}{2}$	37885.90	3875.34	0.780			
		$1\frac{1}{2}$	17830.34		1.122			$2\frac{1}{2}$	41761.24		1.077			
$5d^2(a^2G)6s$	$a^2G$	$4\frac{1}{2}$	17389.06	-321.66	1.14	$5d\ 6s(a^2D)6p$	$z^4P^\circ$	$0\frac{1}{2}$	38398.56	827.90	2.596			
		$3\frac{1}{2}$	17710.72		0.940			$1\frac{1}{2}$	39228.46		1.615			
$5d^2$	$b^4F$	$1\frac{1}{2}$	18897.64	1237.30	0.446	$5d\ 6s(a^2D)6p$	$z^2F^\circ$	$2\frac{1}{2}$	38578.63	2828.23	0.860			
		$2\frac{1}{2}$	20134.94		1.030			$3\frac{1}{2}$	41406.86		1.105			
		$3\frac{1}{2}$	21637.97		1.221			$5d^2(a^2F)6p$	$y^4F^\circ$		$1\frac{1}{2}$	42518.10	1162.65	
		$4\frac{1}{2}$	23145.57		1.309						$2\frac{1}{2}$	43680.75	0.966	
$5d^2$	$b^4P$	$0\frac{1}{2}$	26996.51	288.62	2.58	$5d\ 6s(a^2D)6p$	$y^2F^\circ$	$2\frac{1}{2}$	44399.96	1724.93	1.171			
		$1\frac{1}{2}$	27285.13		1.643			$4\frac{1}{2}$	46124.89		1.223			
		$2\frac{1}{2}$	28547.05		1.573			$5d\ 6s(a^2D)6p$	$y^2P^\circ$		$1\frac{1}{2}$	42770.56	1.220	
$3\frac{1}{2}$	28458.16	-353.33	$0\frac{1}{2}$	43044.26	0.570									
$5d\ 6s(a^2D)6p$	$z^4F^\circ$	$1\frac{1}{2}$	28068.79	1336.33	0.512	$5d\ 6s(a^2D)6p$	$y^2F^\circ$	$2\frac{1}{2}$	43900.56	790.16	1.00			
		$2\frac{1}{2}$	29405.12		1.076			$3\frac{1}{2}$	44690.72		1.142			
		$3\frac{1}{2}$	33778.24		4371.12			1.252	$5d^2(a^2F)6p$		$y^4D^\circ$	$0\frac{1}{2}$	45643.25	1031.11
		$4\frac{1}{2}$	33185.67		4409.43			1.33				$1\frac{1}{2}$	46674.56	1.194
$5d^2$	$b^2G$	$4\frac{1}{2}$	28104.83	-353.33	0.997	$5d\ 6s(a^2D)6p$	$y^2F^\circ$	$2\frac{1}{2}$	43900.56	790.16	1.00			
		$3\frac{1}{2}$	28458.16		0.897			$3\frac{1}{2}$	44690.72		1.142			
		$1\frac{1}{2}$	29160.04		2624.12			0.486	$5d^2(a^2P)6p$		$z^2S^\circ$	$0\frac{1}{2}$	46495.37	1.680
		$1\frac{1}{2}$	31784.16		2570.97			1.216				$5d^2(a^2F)6p$	$z^2G^\circ$	$3\frac{1}{2}$
$2\frac{1}{2}$	34355.13	2527.36	1.320	$4\frac{1}{2}$	49840.47	1.153								
$3\frac{1}{2}$	36882.49	1.380												
$5d^2$	$c^2D$	$1\frac{1}{2}$	30594.63	2183.53	0.91									
		$2\frac{1}{2}$	32778.16		1.21									

Hf II—Continued

Hf II—Continued

Config.	Design.	J	Level	Interval	Obs. $\rho$	Config.	Design.	J	Level	Interval	Obs. $\rho$
5d <sup>2</sup> (b <sup>1</sup> D)6p	z <sup>1</sup> D°	1½ 2½	47973.58 49005.64	1032.06	1.11 1.246	5d <sup>2</sup> (a <sup>1</sup> G)6p	v <sup>1</sup> F°	2½ 3½	68774.9 64448.1	1667.2	1.05 1.15
5d <sup>2</sup> (a <sup>1</sup> F)6p	z <sup>1</sup> S°	1½	47996.48				1	2½	65643.9		
	y <sup>1</sup> P°	0½ 1½ 2½	50419.49		1.725	5d <sup>2</sup> (a <sup>1</sup> F)7p	2(e <sup>1</sup> F?)	1½	69310.2		
							4	2½	69483.4		
5d <sup>2</sup> (a <sup>1</sup> F)6p	z <sup>1</sup> F°	2½ 3½	50910.23 52339.98	1429.70	0.88 1.162	5d 6s(a <sup>1</sup> D)7s	5(e <sup>1</sup> D?)	1½	69963.3		
5d <sup>2</sup> (a <sup>1</sup> F)6p	w <sup>1</sup> D°	1½ 2½	51133.59 52702.59	1569.00	1.074 1.26	5d <sup>2</sup> (a <sup>1</sup> F)7p	3(e <sup>1</sup> F?)	2½	70666.0		
							7	1½	72159.1		
5d <sup>2</sup> (a <sup>1</sup> P)6p	z <sup>1</sup> D°	0½ 1½ 2½ 3½	52096.4 52717.46 54893.53 58448.21	621.1 1976.07 3754.68	0.722 1.43 1.276 1.14	5d 6s(a <sup>1</sup> D)7s	6(e <sup>1</sup> D?)	2½	73665.1		
5d <sup>2</sup> (b <sup>1</sup> D)6p	w <sup>1</sup> F°	2½ 3½	52204.7 53227.27	1022.6	1.132		9	1½	75158.2		
5d <sup>2</sup> (b <sup>1</sup> D)6p	z <sup>1</sup> P°	0½ 1½	52847.8 55077.0	2229.4	1.28		10	2½	75801.1		
5d <sup>2</sup> (a <sup>1</sup> P)6p	z <sup>1</sup> P°	0½ 1½ 2½	53330.5 53494.10 55080.35	163.6 1566.25	2.394 1.544 1.168		11	1½	76016.4		
							12	2½	76585.0		
5d <sup>2</sup> (a <sup>1</sup> G)6p	z <sup>1</sup> H°	4½ 5½	53546.17 56571.61	3025.44	1.02 1.09		13	1½	77163.1		
							14	3½	77314.4		
5d <sup>2</sup> (a <sup>1</sup> G)6p	y <sup>1</sup> G°	3½ 4½	53972.48 56638.80	1666.34	1.175 1.06	5d <sup>2</sup> (a <sup>1</sup> G)7s	16(e <sup>1</sup> G?)	3½	79804.4		
5d <sup>2</sup> (a <sup>1</sup> P)6p	w <sup>1</sup> P°	1½ 0½	56345.74 57477.6	-1131.9	1.02 0.72	5d <sup>2</sup> (a <sup>1</sup> G)7s	17(e <sup>1</sup> G?)	4½	81050.9		
	v <sup>1</sup> D°	1½ 2½	58117.4		1.22		18	4½	82100.5		
5d <sup>2</sup> (a <sup>1</sup> P)6p	w <sup>1</sup> D°	1½ 2½	61318.3 64664.1	3345.8			19	4½	83129.8		
						Hf III(2F <sub>2</sub> )	Limit	-----	129000		

January 1957.

Hf II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms					
	5d 6s <sup>2</sup>	a <sup>1</sup> D				
5d <sup>2</sup>	{ b <sup>1</sup> P b <sup>3</sup> P	c <sup>1</sup> D	b <sup>1</sup> F b <sup>3</sup> F	b <sup>1</sup> G	a <sup>1</sup> H	
	ns (n ≥ 6)			np (n ≥ 6)		
5d <sup>2</sup> (a <sup>1</sup> F)ns	{ a, e <sup>1</sup> F a <sup>3</sup> F			y <sup>1</sup> D° w <sup>1</sup> D°	y <sup>1</sup> F° z <sup>1</sup> F°	z <sup>1</sup> G° z <sup>3</sup> G°
5d 6s(a <sup>1</sup> D)ns	{ e <sup>1</sup> D			z <sup>1</sup> P° z <sup>3</sup> P°	z <sup>1</sup> D° y <sup>1</sup> D°	z <sup>1</sup> F° z <sup>3</sup> F°
5d <sup>2</sup> (a <sup>1</sup> P)ns	{ a <sup>1</sup> P a <sup>3</sup> P			z <sup>1</sup> S° z <sup>3</sup> S°	z <sup>1</sup> P° w <sup>1</sup> P°	z <sup>1</sup> D° u <sup>1</sup> D°
5d 6s(a <sup>1</sup> D)ns				y <sup>1</sup> P°	z <sup>1</sup> D°	y <sup>1</sup> F°
5d <sup>2</sup> (a <sup>1</sup> G)ns			a, e <sup>1</sup> G			v <sup>1</sup> F° y <sup>1</sup> G°
5d <sup>2</sup> (b <sup>1</sup> D)ns	b <sup>1</sup> D			z <sup>1</sup> P°	z <sup>1</sup> D°	w <sup>1</sup> F°

\*For predicted terms in the spectra of the La I isoelectronic sequence, see Vol. III, Introduction.



## TANTALUM

## Ta I

73 electrons

Z = 73

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^3 6s^2 \text{ } ^4F_{1/2}$  $6s^2 \text{ } ^4F_{1/2}$  63600 K

I. P. 7.88 volts

The first regularities in Ta I were reported by C. C. Kiess and Mrs. H. K. Kiess in 1933. Subsequently, at the Zeeman Laboratory, Miss van den Berg, Klinkenberg, and van den Bosch extended the study, and reported in 1952 that 1262 lines had been classified. In the 1956 reference below some theoretical aspects of this work are discussed.

The present analysis is by C. C. Kiess and Mrs. H. K. Kiess, who have continued their study of Ta I over a period of nearly 25 years, and furnished their unpublished data especially for inclusion here. Their observations extend from 1953.08 Å to 12139.58 Å. The total number of classified lines now exceeds 2300. Observed intersystem combinations connect the systems of terms of different multiplicity.

They have not yet completed their analysis, and some designation assignments are tentative, as indicated by a question mark in column two of the table. Many of the levels are listed as miscellaneous owing to the departure from LS-coupling. During much of the work, R. E. Trees has generously assisted by providing the predicted positions of even terms derived from his theoretical calculations.

The observed  $g$ -values given to three decimal places in the table are from spectrograms taken by Harrison and his colleagues at the Massachusetts Institute of Technology. All Zeeman measurements used for the present work have been made by C. C. Kiess. The resulting observed  $g$ -values have been averaged for 46 even levels and 170 odd levels, and the work is still in progress. This wealth of reliable Zeeman data for Ta I exceeds that for any spectrum known to the writer. For example, the tabular  $g$ -values for  $\alpha \text{ } ^4F_{1/2}$  ( $0.447 \pm 0.002$ ), and for  $\alpha \text{ } ^4F_{3/2}$  ( $1.031 \pm 0.004$ ) each represent means of 38 determinations. For the remaining low even levels the number of individual observations used for the final means fall into the following groups:

No. observations	30 to 34	20 to 28	10 to 15	1 to 9
No. $g$ -values	4	8	5	4

These Zeeman data should provide ideal material for a detailed study of  $g$ -sharing and coupling in a complex spectrum.

Many of the Ta I lines exhibit wide hyperfine structure. Means of the hyperfine structure components, weighted by intensity, have been used for such lines in the present work.

The limit is from the  $ns \text{ } ^4F$  series ( $n=6$  to 8), and has been determined by means of a Rydberg formula.

## REFERENCES

- C. C. Kiess and H. K. Kiess, *Bur. Std. J. Research* **11**, 277, RP 589 (1933). (T)  
 P. F. A. Klinkenberg, G. J. van den Berg, and J. C. van den Bosch, *Physica* **16**, 861 (1950). (T) (C L) (Z E).  
 C. C. Kiess and H. K. Kiess, *J. Opt. Soc. Am.* **41**, 867 (A) (1951). (I P)  
 G. J. van den Berg, *Dissertation*, Univ. Amsterdam, 76 pp (1951). (I P) (T) (C L) (Z E)  
 P. F. A. Klinkenberg, G. J. van den Berg, and J. C. van den Bosch, *Physica* **17**, 167 (L) (1951). (I P) (T)  
 G. J. van den Berg, P. F. A. Klinkenberg, and J. C. van den Bosch, *Physica* **18**, 221 (1952). (I P) (T) (C L) (Z E)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 T. Kamei, *Phys. Rev.* **99**, 789 (1955). (hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, ORNL-1469, Suppl. II, 31 (1955). (Summary hfs)  
 TH. A. M. van Kleef and P. F. A. Klinkenberg, *Phys. Rev.* **101**, 489 (L) (1956).  
 C. C. Kiess and H. K. Kiess, unpublished material (July 1957). (I P) (T) (C L) (Z E)

Ta I

Ta I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g		
5d <sup>3</sup> 6s <sup>2</sup>	a <sup>4</sup> F	1½	0.00	2010.10	0.447	5d <sup>3</sup> 6s(a <sup>4</sup> F)6p	z <sup>4</sup> F°	5½	23141.47				
		2½	2010.10	1953.82	1.031			0¼	<del>23555.41</del>			888.01	-0.320
		3½	3963.92	1657.12	1.218			1½	<del>21843.48</del>			937.70	1.126
		4½	5621.04		1.272			2½	<del>25181.18</del>			1404.81	1.239
5d <sup>3</sup> 6s <sup>2</sup>	a <sup>4</sup> P	0¼	6049.42	19.49	2.454		?	3½	<del>26585.93</del>	1147.50	1.356		
		1½	6068.91	3184.52	1.527			4½	<del>27753.43</del>	2627.79	1.390		
		2½	9253.43		1.580			5½	<del>30361.28</del>		1.334		
5d <sup>3</sup> 6s <sup>2</sup>	a <sup>4</sup> G	3½	9705.38	984.94	0.912	5d <sup>3</sup> ( <sup>4</sup> F)6s	b <sup>4</sup> F	1½					
		4½	10690.32		1.063			2½	23512.34			1.048	
5d <sup>3</sup> (a <sup>4</sup> D)6s	a <sup>4</sup> D	0¼	9758.97	216.84	3.020	5d <sup>3</sup> 6s(a <sup>4</sup> F)6p	z <sup>4</sup> D°	0¼	<del>24516.69</del>	222.34	2.888		
		1½	9975.81	1267.82	1.542			1½	<del>24759.03</del>	2055.73	1.620		
		2½	11243.63	991.13	1.641			2½	<del>26794.76</del>	985.86	1.416		
		3½	12234.76	1116.69	1.578			3½	<del>27780.68</del>	986.03	1.374		
5d <sup>3</sup> 6s <sup>2</sup>	a <sup>4</sup> P	1½	10950.22	-841.91	1.407		?	4½	<del>28766.65</del>		1.337		
		0¼	11792.13		1.116			0¼	24698.70				
5d <sup>3</sup>	a <sup>4</sup> S	2½	11796.14		2.134			3½	24917.90				
5d <sup>3</sup> 6s <sup>2</sup>	a <sup>4</sup> D	2½	12865.97	-3037.80	1.214			4½	24934.58				
		1½	15903.77		1.199			4½	<del>25185.89</del>				
		2½	14875.707					2½	25224.06				
5d <sup>3</sup> 6s <sup>2</sup>	a <sup>4</sup> H	5½	15114.14	-276.87	1.021			4½	25376.41				
		4½	18391.01		1.014			1½	25478.30				
5d <sup>3</sup> 6s <sup>2</sup>	a <sup>4</sup> F	2½	17224.47	158.65	0.882	5d <sup>3</sup> 6s(a <sup>4</sup> F)6p	y <sup>4</sup> D°	0¼	<del>25512.63</del>	851.06	0.028		
		3½	17383.12		1.125			1½	<del>26363.69</del>	1770.19	1.393		
5d <sup>3</sup> 6s(a <sup>4</sup> F)6p	z <sup>4</sup> G°	1½	<del>17384.65</del>	1793.80	0.851	5d <sup>3</sup> 6s(a <sup>4</sup> P)6p	z <sup>4</sup> P° ?	2½	<del>26219.62</del>	-370.41	1.338		
		2½	<del>19178.45</del>	1381.81	1.194			1½	<del>26590.03</del>	-276.02	1.576		
		3½	<del>20560.26</del>	2121.45	1.231			0¼	<del>26866.05</del>		2.650		
		4½	<del>22081.71</del>	2327.12	1.302			4½	26678.47		1.223		
		5½	<del>25008.83</del>					3½	<del>26960.46</del>		1.606		
		6½						2½	27167.82		1.606		
5d <sup>3</sup> 6s <sup>2</sup> (a <sup>4</sup> F)6p	z <sup>4</sup> D°	2½	17993.74		0.732			5½	27783.07		1.351		
		0¼	<del>18504.78</del>	1153.06	0.172			3½	<del>28182.60</del>		1.115		
		1½	<del>19657.78</del>	1509.83	1.018			5½	28285.99		1.356		
		2½	<del>21167.61</del>	2759.07	1.120			4½	<del>28766.65</del>		1.337		
5d <sup>3</sup> 6s <sup>2</sup> (a <sup>4</sup> P)6p	z <sup>4</sup> S°	3½	<del>23926.68</del>		1.326			2½	<del>28862.01</del>		1.247		
		0¼	<del>20340.39</del>		1.956			2½	<del>29343.46</del>		0.814		
		3½	20646.54	506.79	0.818			3½	<del>29722.95</del>		1.217		
5d <sup>3</sup> (b <sup>4</sup> H)6s	a <sup>4</sup> H	4½	21153.33	1275.23	1.089	5d <sup>3</sup> 6s(a <sup>4</sup> P)6p	y <sup>4</sup> D°	0¼	<del>29902.27</del>	762.39	2.994		
		5½	22428.56	1086.30	1.159			1½	<del>30684.66</del>	1822.09	1.812		
		6½	23514.86		1.217			2½	<del>32486.75</del>	1607.91	1.687		
		5½	26022.74		1.260			3½	<del>34094.66</del>		1.515		
5d <sup>3</sup> 6s <sup>2</sup> (a <sup>4</sup> P)6p	z <sup>4</sup> D°	1½	<del>20772.32</del>	1275.13	0.812			3½	30015.61		1.186		
		2½	<del>22047.45</del>		1.179			4½	<del>30021.20</del>		1.186		
5d <sup>3</sup> (a <sup>4</sup> D)6s	a <sup>4</sup> D	2½	21091.53					3½	30590.95		1.043		
		0¼	22235.97	-854.96	1.020	5d <sup>3</sup> 6s(a <sup>4</sup> P)6p	z <sup>4</sup> S°	1½	30894.67		1.808		
		1½	21381.01	3165.19	1.294								
		2½	24546.20	2028.82	1.42								
3½	26575.02												
5d <sup>3</sup> (b <sup>4</sup> G)6s	a <sup>4</sup> G	2½	21622.92	1138.29	0.894								
		3½	22761.21	1151.68	1.008								
		4½	23912.89	2109.85	1.185								
		5½	26022.74		1.260								
5d <sup>3</sup> 6s(a <sup>4</sup> F)6p	z <sup>4</sup> F°	1½	<del>21855.07</del>	1508.02	0.666								
		2½	<del>23363.09</del>	1618.76	1.078								
		3½	<del>24981.85</del>	944.49	1.235								
		4½	<del>25928.34</del>		1.292								
		3½	<del>22380.53</del>		1.060								
		2½	22434.37										
1½	22842.84		0.894										

Ta I—Continued

Ta I—Continued

Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	
5d <sup>3</sup> 6s(a <sup>3</sup> F)6p	z <sup>3</sup> P°	1½	31088.34	895.08 1236.33	2.108	5d <sup>3</sup> 6s(a <sup>3</sup> G)6p	y <sup>4</sup> G° ?	2½	36825.97	1853.08 2007.37	0.825	
		2½	31981.42		1.684			3½	38679.06		1.055	
		3½	33197.75		1.568			4½	40688.42		1.208	
5d <sup>3</sup> 6s(a <sup>3</sup> F)6p	z <sup>3</sup> P°	2½	31428.05					5½				
		3½	31428.05					3½	37143.26			
		0½	31500.99		0.758			2½	37145.60		1.501	
		1½						0½	37461.46		1.257	
		4½	31530.02		1.086			1½	37523.54		0.973	
		1½	31553.89		0.888			3½	37681.25		1.242	
		3½	31600.95		1.089			2½	37630.09		1.211	
		3½	32132.38		0.846			1½	37760.67		0.518	
		1½	32214.94		0.728		5d <sup>3</sup> 6s(a <sup>3</sup> H)6p	z <sup>4</sup> I° ? ?	4½	37942.84	1525.79 1443.20	0.878
		5½	33070.28	1.349		5½			39488.63	1.040		
		2½	33184.97	1.273		6½			40911.83			
		4½	33497.16	1.031		7½						
		4½	33615.49		1.067		5d <sup>4</sup> (a <sup>4</sup> D)6p	y <sup>4</sup> F°	3½	38253.39		1.112
		2½	34001.16		1.140				0½	38507.54	1080.27	0.730
		1½	34078.42		1.089				1½	39587.81	1993.17	1.083
		2½	34792.22		1.483				2½	41580.98	1596.39	1.356
		3½	34799.71		1.210				3½	43177.37	1629.27	1.417
		4½	35217.94		1.582				4½	44808.64	1032.11	1.280
		1½	35242.94		1.534				5½	46838.75		
		4½	35497.65		0.940				1½	38545.70		1.359
1½	35721.0		1.010		2½	38753.75				0.952		
3½	35746.18		1.101		0½	38845.25				0.460		
5½	35813.47		1.200		1½	38994.33				1.027		
2½	35876.47		1.490		2½	39059.52				1.054		
2½	35925.09				1½	39253.07				1.128		
5d <sup>3</sup> 6s(a <sup>3</sup> H)6p	z <sup>4</sup> G° ?	2½	36013.97	785.91 13.44 1697.06	0.985			4½	39422.40		1.143	
		3½	36799.88		1.133		0½	39490.14		0.560		
		4½	38813.32		1.088		3½	39526.7				
		5½	40510.38		1.264		3½	39641.24		1.090		
		2½	36069.2				2½	39688.20		0.930		
5d <sup>3</sup> 6s(a <sup>3</sup> H)6p	z <sup>4</sup> H° ? ?	3½	36159.13	917.58 1117.51 1166.46	0.852			2½	39786.52		1.065	
		4½	37076.71		1.046		3½	39936.13		1.295		
		5½	38194.22				1½	40230.01		1.232		
		6½	39360.68		1.061		0½	40244.74		1.056		
		0½	36345.8			0.536		3½	40333.03		1.159	
5d <sup>3</sup> 6s(a <sup>3</sup> G)6p	y <sup>4</sup> F° ?	1½	36530.06	1867.93	0.716			1½	40539.2		0.956	
		2½	38447.99		1.045		3½	40755.90		1.134		
		3½					2½	40851.61		1.265		
		4½										
		4½	36831.15		1.211							

Ta I—Continued

Ta I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g		
5d <sup>2</sup> 6s(a <sup>2</sup> F)7s	e <sup>2</sup> F	3½	40981.79		1.245	5d <sup>2</sup> (a <sup>2</sup> D)6p	z <sup>2</sup> D <sup>o</sup>	0½	44146.16		0.714		
		2½	41010.07		1.227			3½	44165.58		1.067		
		0½	41151.26		-0.657			4½	44173.68		1.115		
		1½	41594.83	442.57	0.996			1½	44360.19		1.309		
		2½	42501.59	908.76	1.250			2½	44386.40		1.179		
		3½	43982.43	1480.84	1.433			4½	44402.56		1.146		
		4½	45636.83	1654.40	1.411			1½	44483.96		1.034		
		5½	47319.57	1682.74	1.434			0½	44518.10	728.12	2.378		
		0½	41179.9		0.434			1½	45248.22	9.76	1.85		
		1½	41197.67		1.335			2½	45855.33	1153.81	1.662		
		3½	41856.56		1.533			3½	46409.79	1390.02	1.532		
		2½?	41584.94					4½	47799.81				
		1½	41692.64		1.183			3½	44669.23		1.682		
		2½	41879.23		1.199			3½	44893.40		1.227		
		0½	41902.98		0.745			5½	45057.34		1.056		
		3½	42017.95		1.107			3½	45114.71		1.337		
		1½	42178.76?					5d <sup>2</sup> (a <sup>2</sup> D)6p	y <sup>2</sup> P <sup>o</sup>	1½	45241.73		1.245
		4½	42247.00		1.268					2½	45714.68	472.95	
		1½	42408.16		1.227					3½	47621.88	1907.20	
		3½	42751.72		1.170					1½	45276.39		
1½	42778.70		1.321	3½	45648.26		1.264						
2½	42844.73		1.409	1½	45723.58		1.112						
3½	42953.1		1.046?	3½	45841.6								
2½	42982.8		1.172	3½?	46060.6								
4½	43090.22		1.222	2½	46116.93		1.254						
1½	43167.07		1.135	3½	46172.44		1.186						
5d <sup>2</sup> (a <sup>2</sup> D)6p	y <sup>2</sup> P <sup>o</sup>	5½	43185.09		1.219	1½	46176.76		0.896				
		2½	43239.22		0.973	5½	46552.13		1.266				
		4½	43391.71			1½	46887.79		0.737				
		0½	43478.20		2.18	2½	46933.38						
		1½	44689.31	1211.11	1.809	1½	46974.03		1.871				
		2½	46323.31	1634.00	1.601	2½	46981.89						
		4½	43493.24?			3½	47013.02						
		2½	43496.65		1.22	5½	47339.48						
		3½	43533.3		1.215	2½	47397.06						
		2½	43550.78		0.945	3½	47625.01						
5d <sup>2</sup> 6s(a <sup>2</sup> F)7s	e <sup>2</sup> F	1½	43818.63		1.642	3½	47931.16						
		2½	43880.84		1.284	3½	48069.61						
		1½	43964.50			2½	48290.45						
		2½	44461.60	497.10	0.988	2½	48369.35						
		3½	47817.16	3355.56		4½	48468.96						
		4½	50509.7	2692.5		4½	49149.30						
		1½	44095.95										
		5½	44141.31		1.122								

Ta I—Continued

Ta I—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
		3½	49551. 07			5d <sup>9</sup> 6s(a <sup>4</sup> F)8s	f <sup>4</sup> F	1½	54440. 57		0. 502
		3½	49599. 81					2½			
		3½	49888. 57					3½			
		2½	49981. 38					4½			
		4½	50014. 07					2½	54481. 09		
		1½	50145. 55					4½	55232. 59		
		4½	50439. 3					2½	56050. 2		
		1½	50877. 75					1½	58391. 76		
		3½	51418. 1					1½	58583. 88		
		3½	51988. 5					0½	61551. 17		
		4½	52723. 73					2½	61737. 0		
		1½	52907. 8					1½	61789. 26		
						Ta II( <sup>4</sup> F <sub>1</sub> )	Limit		63600		

July 1957.

Ta I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms	
5d <sup>9</sup> 6s	{ a <sup>4</sup> P a <sup>4</sup> P a <sup>3</sup> D a <sup>4</sup> F a <sup>4</sup> F a <sup>3</sup> G a <sup>3</sup> H	
5d <sup>9</sup>	a <sup>4</sup> S	
	ns (n ≥ 6)	np (n ≥ 6)
5d <sup>9</sup> 6s(a <sup>4</sup> F)nz	{ a, f <sup>4</sup> F	z <sup>4</sup> D° z <sup>4</sup> F° z <sup>4</sup> G° y <sup>4</sup> D° z <sup>4</sup> F°
5d <sup>9</sup> 6s(a <sup>3</sup> F)nz		z <sup>4</sup> D°
5d <sup>9</sup> 6s(a <sup>3</sup> P)nz		z <sup>3</sup> S° z <sup>3</sup> P° z <sup>3</sup> D° z <sup>3</sup> P° z <sup>3</sup> D°
5d <sup>9</sup> 6s(a <sup>3</sup> P)nz		z <sup>3</sup> S° z <sup>3</sup> P° y <sup>3</sup> D°
5d <sup>9</sup> 6s(a <sup>3</sup> G)nz		y <sup>3</sup> F° y <sup>3</sup> G°
5d <sup>9</sup> (a <sup>3</sup> D)nz	{ a <sup>3</sup> D a <sup>3</sup> D	y <sup>3</sup> P° z <sup>3</sup> D° y <sup>3</sup> ° y <sup>3</sup> P°
5d <sup>9</sup> 6s(a <sup>3</sup> H)nz		z <sup>3</sup> G° z <sup>3</sup> H° z <sup>3</sup> I°
5d <sup>9</sup> (b <sup>3</sup> H)nz	a <sup>3</sup> H	
5d <sup>9</sup> (b <sup>3</sup> G)nz	a <sup>3</sup> G	
5d <sup>9</sup> ( <sup>3</sup> F)nz	b <sup>3</sup> F	

\*For predicted terms of the Ta I isoelectronic sequence, see Vol. III, Introduction.

## Ta II

(Hf I sequence; 72 electrons)

Z=73

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^3 6s^2 F_1$  $a^5F_1$ 

K

I. P.

volts

A preliminary list of 115 levels and  $g$ -values for Ta II was published in 1950; seven more levels were added in 1951. The present analysis is by Kiess, who has revised and greatly extended the early work especially for inclusion here. His observations range from 2000 Å to 5300 Å and include some 2400 lines, of which about 1000 are classified. Observed intersystem combinations connect the systems of terms having different multiplicity.

The three-place  $g$ -values in the table are from Zeeman spectrograms taken at the Massachusetts Institute of Technology; two-place values are from Bureau spectrograms.

In his recent work Kiess has greatly benefited by the theoretical investigations of Trees. In particular, one level at 16424 K listed in the 1950 paper has been rejected, and one at 23294 K has been assigned a revised  $J$ -value, on the basis of Trees' work.

No series are known for Ta II. From a study of screening constants Finkelnburg and Humbach interpolate an ionization potential of  $16.2 \pm 0.5$  volts. This places the limit near 130700 K.

## REFERENCES

- C. C. Kiess, G. R. Harrison, and W. J. Hitchcock, *J. Research Nat. Bur. Std.* **44**, 245, RP 2075 (1950). (T) (Z E).  
 G. J. van den Berg, *Dissertation* pp 26, 38 (Univ. Amsterdam 1951). (T) (Z E)  
 B. M. Brown and D. H. Tombouljian, *Phys. Rev.* **88**, 1158 (1952); **91**, 1580 (1953). (hfs)  
 R. E. Trees, W. F. Cahill, and P. Rabinowitz, *J. Research Nat. Bur. Std.* **55**, 335, RP 2639 (1955).  
 W. Finkelnburg und W. Humbach, *Naturwiss.* **43**, 35 (1955). (I P)  
 R. E. Trees, unpublished material (April 1957).  
 C. C. Kiess, unpublished material (April 1957). (T) (C L) (Z E)

Ta II

Ta II

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$				
$5d^3(^4F)6s$	$a^4F$	1	0.00		0.000	$5d^3 6s^2$	$a^4G$	4	14205.53						
		2	1031.33	1031.33	1.008			$5d^3(^3D)6s$	$a^3D$	1	14627.75	2540.75	0.850		
		3	2642.19	1610.86	1.250					2	17168.50	1385.33	1.211		
		4	4415.70	1773.51	1.350					3	18553.83		1.33		
		5	6186.72	1771.02	1.33					$5d^3(^3H)6s$	$a^3H$	4	15851.12		1.16
$5d^3 6s^2$	$a^3F$	2	3180.04	3651.31	0.750	5									
		3	6331.35	2914.98	1.098	6									
		4	9746.33		1.225	$5d^3(^3P)6s$	$b^3P$	0							
		$5d^3 6s^2$	$a^3P$	0	4124.77			1205.89	0.000			1	17375.11	1125.49	1.171
				1	5330.66			327.34	1.550	2	18500.60		1.462		
2	5658.00				1.340			$5d^3(^3F)6s$	$c^3F$	2	22928.62	691.74	0.700		
$5d^3(^4F)6s$	$b^4F$			2	9690.46					4890.54	1.063	3	23620.36	-537.66	1.076
				3	14581.00	3912.67	1.004			4	23082.70		1.026		
		4	18493.67		1.23	$5d^3(^3D)6s$	$b^3D$			2	23294.70		1.120		
		$5d^3(^4P)6s$	$a^4P$	1	10713.30					1162.16	2.374	$5d^3(^3P)6s$	$c^3P$	0	23381.19
				2	11875.46			560.39	1.48	1					
3	12435.85				1.614			2							
$5d^3(^3G)6s$	$a^3G$			3	11767.14			938.18	0.915	$5d^3(^3P)6s$	$a^3P$			1	23406.14
				4	12705.32	125.62	1.021	$5d^3$	$b^3H$					4	24432.78
		5	12830.94		1.280	5	25413.95						1.052		
		$5d^3$	$a^3D$	0	12600.37	874.53	0.000					6			
				1	13475.40	1019.50	1.510					$5d^3(^3F)6s$	$a^3F$	3	24869.61
2	14494.90			1231.16	1.472										
3	15726.06			1505.16	1.476										
4	17231.22				1.23										
5															
$5d^3 6s^2$	$a^3D$	2	13560.25		1.111										

Ta I—Continued

Ta II—Continued

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$	
5d <sup>o</sup>	b <sup>o</sup> G	3	26829.09		0.855			5	46295.07			
		4										
		5										
		2	29256.87		0.522			2 <sup>o</sup> D	2	46387.28		1.31
		1	33706.50		0.291			4	46645.81		1.210	
		2	33715.15		0.800			3	46831.39		1.204	
		3	36112.97					2	46850.69		1.097	
		2	36177.12		0.947			3	47169.18		1.093	
		3	36763.73		1.169			4	47220.92		1.198	
		1	36987.73		0.685			2	47514.61		1.292	
		2 <sup>o</sup> G	2	37230.80		0.64		1	47596.02			
		2 <sup>o</sup> F	2	38515.55		1.026		0	47800.95			
	1 <sup>o</sup> F	1	38535.26		0.472		3	47825.41		1.198		
	3 <sup>o</sup> G	3	38962.38		1.001		0	48064.50				
	3 <sup>o</sup> F	3	39295.83		1.140		2	48166.44				
	4 <sup>o</sup> G	4	39743.64		1.222		2	48223.11		1.448		
	0	40023.76					4	48470.35		1.268		
	2	40233.53		1.158			2	48666.55		1.137		
	1	40304.78		1.232			1	48776.36		0.750		
	2	41145.00		1.161			3	48962.66		1.275		
	1	41355.06		1.893			2	49080.44		1.33		
	3	41554.52		1.204			3	49536.28		1.318		
	5	41709.02					2	49592.90		1.63		
	4	41775.29		0.72			3	49646.60		1.045		
	4 <sup>o</sup> F	4	42122.91		1.265		1	49827.03		2.023		
	2	42153.32		1.210			4	49937.82				
	3	42959.59		1.117			3	51073.88		1.156		
	2	43064.95		1.061			2	51197.32		1.363		
	3	43544.44		1.086			1	51326.37		1.052		
	1	43553.87		1.50			2	51534.21		1.317		
	4	44005.16					3	52121.22		1.088		
	1	44206.19		0.255			1	52155.76		1.618		
	2	44259.20		1.251			3	52520.41		1.403		
	3	44430.38		0.950			1	52824.53		0.930		
	0 <sup>o</sup> D	0	44434.70		0.00		3	53465.72		1.226		
	5 <sup>o</sup> F	5	44585.33		1.245		2	53644.22		1.339		
	4	44826.00		1.264			2	54533.78		1.035		
	3	44835.23		1.34			3	54848.81		0.935		
	1 <sup>o</sup> D	1	45233.85		1.461		3	55122.36		1.250		
	2	45446.91		1.150			2	55543.11		1.46		
	1	46174.63		1.361			2	56018.75		0.991		
	4	46226.97		1.405			4	56142.49		1.26		
						4	56752.94		1.22			

April 1957.

## TUNGSTEN

W I

74 electrons

Z=74

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^4 6s^2 {}^3D_3$  $\alpha$   ${}^3D_3$ , 64400 K

I. P. 7.98 volts

The spectrum needs further critical study. The analysis is chiefly from Laporte and Mack, who have reported 300 levels with 201 "tentative"  $g$ -values, and 2378 classified lines between 2008.64 Å and 10477.97 Å. Their observational material was taken mostly from the literature, supplemented by Mack's observations made at Princeton for the region short of 2500 Å. The earlier work by Catalán and Poggio and by Laun was revised and extended by these authors.

Later, the spectrum was observed by Kiess at the National Bureau of Standards in the region longer than 2000 Å. Short of this limit, observations were made by Boyce with the Carnegie Institution Spectrograph. Laun measured these spectrograms and prepared a complete line list extending from 1783.734 Å to 7727.14 Å. The earlier long-wave Bureau measurements used by Mack and Laporte are quoted in Laun's revised line list to extend the observations to 10477.97 Å.

Laun has revised the published level values, changed some  $J$ -values, and added more than 50 new levels. His unpublished material has been furnished for inclusion here. The writer has tentatively retained 4 of the 16 published levels rejected by Laun. There are now approximately 3700 classified lines.

Mack and Laporte assigned a number of configurations and term designations to the known levels. Murakawa has confirmed some of the configuration assignments from a study of the isotope displacement effect, worked out from observations of the hyperfine structure of W I. He attributes some levels, also, to the  $5d^3 6s^2 6p$  configuration.

The term and configuration assignments in the table are tentative. The writer has consulted Racah, and included some of his very provisional rearrangements of known levels, although they are far from definitive. She has also made a few other changes on the basis of the present known combinations. Observations indicate that the line at 4008.753 Å ( $\alpha$   ${}^3S_1 - s$   ${}^3P_2$ ) is the *voie ultime*. From this evidence the  $s$   ${}^3P^{\circ}$  term in the table belongs predominantly to the configuration  $5d^3(\alpha$   ${}^3S)6p$ , instead of  $5d^3 6s(\alpha$   ${}^3D)6p$ , as published. The  ${}^3P^{\circ}$  term from the latter configuration is apparently not yet known.

Mack feels that grouping of levels into terms is not justifiable in W I because of the departure from  $LS$ -coupling. The writer has, however, retained the arrangement by terms among lower levels, in spite of considerable overlapping. This plan conforms to the general format of these Volumes and facilitates a ready comparison of the relative positions of the terms, the term intervals, and the observed  $g$ -values among similar spectra. She urges the users to bear in mind the reservations regarding this format that are described in the 1943 paper, and to consult this paper for further details.

The observed  $g$ -values are quoted from the 1943 paper by Laporte and Mack, pending further study of Zeeman spectrograms taken at the Massachusetts Institute of Technology. There are more than 80 given only to tenths; these are urgently in need of revision based on present observations.

The limit is from Laporte and Mack. It is derived by a Rydberg formula from the  $5d^3 6s(\alpha$   ${}^3D)ns$   ${}^3D_3$  series ( $n=6,7$ ) corrected by  $-2.8$  percent. This Ritz correction from other spectra may need further revision when more series are known for neighboring spectra.



## W I—Continued

## REFERENCES

- D. D. Laun, *J. Research Nat. Bur. Std.* **21**, 207, RP 1125 (1938). (T)  
 O. Laporte and J. E. Mack, *Phys. Rev.* **63**, 246 (1943). (I P) (T) (C L)  
 K. Murakawa, *J. Phys. Soc. Japan* **8**, 215 (1953). (hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 31 (1955).  
 (Summary hfs)  
 D. D. Laun, unpublished material (June 1955). (T) (C L)  
 G. Racah, letter, (March 1956).

W I

W I

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
5d <sup>4</sup> 6s <sup>2</sup>	a <sup>3</sup> D	0	0.00		0/0	5d <sup>4</sup> 6s(a <sup>3</sup> D) 6p	s <sup>3</sup> F <sup>o</sup>	1	25983.60	1678.92	0.54
		1	1670.30	1670.30	1.51			2	27662.52	1476.59	1.21
		2	3325.53	1655.23	1.48			3	29139.11	2293.80	1.06
		3	4830.00	1504.47	1.50			4	31432.91	1937.12	1.32
		4	6219.33	1389.33	1.49			5	33370.03		1.39
5d <sup>4</sup> (a <sup>3</sup> S) 6s	a <sup>3</sup> S	3	2951.29		1.98	5d <sup>4</sup> (a <sup>3</sup> S) 6p	s <sup>3</sup> P <sup>o</sup>	2	26229.77	1258.34	1.84
5d <sup>4</sup> 6s <sup>2</sup>	a <sup>3</sup> P	0	9528.07		0/0			3	27488.11	401.57	1.72
		1	13307.06	3778.99	1.32			4	27389.68		1.71
		2	19253.58	5946.52	1.18			2	26367.28		0.87
5d <sup>4</sup> 6s <sup>2</sup>	a <sup>3</sup> H	4	12161.95		0.99	5d <sup>4</sup> 6s(a <sup>3</sup> D) 6p	s <sup>3</sup> D <sup>o</sup>	0	26629.46	1149.04	0/0
		5	15069.94	2907.99	1.05			1	27778.50	1417.34	1.25
		6	17008.50	1938.56	1.4			2	29195.84	717.01	1.28
5d <sup>4</sup> 6s <sup>2</sup>	a <sup>3</sup> G	3	13348.54		0.92			3	29912.85	2915.27	1.31
		4	16431.28	3082.74	1.02			4	32228.12		1.7
		5	19826.04	3394.76	1.20			5	27849.84		
5d <sup>4</sup> 6s <sup>2</sup>	a <sup>3</sup> F	2	13777.70		1.09	5d <sup>4</sup> 6s(a <sup>3</sup> D) 6p	s <sup>3</sup> P <sup>o</sup>	1	28198.90	1194.50	1.83
		3	17701.14	3923.44	1.02			2	29395.40	1193.24	1.64
		4	17107.02	-594.12	1.19			3	30586.64		
5d <sup>4</sup> 6s <sup>2</sup>	a <sup>3</sup> D	3	15459.99		1.17			5	28233.44		
		2	14976.21	483.78	1.06			6	28392.72		
		1	18082.80	-3106.59	0.7						
5d <sup>4</sup> (a <sup>3</sup> G) 6s	a <sup>3</sup> G	2	18116.84		1.08			1	30683.52		1.39
		3	18974.47	857.63	1.06			1	31325.42		0.86
		4	19256.23	281.76	1.20			2	31817.63		1.52
		5	19535.04	278.81	1.21			3	32238.02		1.3
		6	19648.56	113.52	1.32						
5d <sup>4</sup> (a <sup>3</sup> S) 6s	a <sup>3</sup> S	2	18280.48		1.43			0	32386.51		
5d <sup>4</sup> 6s(a <sup>3</sup> D) 6p	s <sup>3</sup> F <sup>o</sup>	0	19389.43		0/0			3	32957.58		1.43
		1	20064.30	674.87	1.54			2	33141.38		1.51
		2	21448.76	1384.46	1.48			2	33944.06		
		3	23047.31	1598.55	1.53			4	34121.63		1.5
		4	24765.39	1716.08	1.50			3	34228.60		
		5	26676.48	1913.09	1.46			1	34342.41		1.56
		6	28643.06?	2966.58?				3	34354.08		0.71
	a <sup>3</sup> P?	3	19827.67		1.28			2	34485.86		0.82
		2	20983.06	-1155.39				4	34632.58		0.89
		1	20427.81	555.25	2.1			1	34719.33		0.15
		0	20174.28					4	35116.78		1.2
5d <sup>4</sup> 6s(a <sup>3</sup> D) 6p	s <sup>3</sup> D <sup>o</sup>	1	21453.90		2.51			2	35311.54		1.0
		2	23964.67	2510.77	1.93			3	35499.15		1.0
		3	26189.20	2224.53	1.80						
		4	28797.24	2608.04	1.61						
		5	29773.34	976.10	1.55						
	<sup>3</sup> D?	4	22476.68		1.48						
		0	22773.84								
	<sup>3</sup> F?	5	22852.84		1.2						
	<sup>3</sup> D?	3	23930.10		1.4						

W I—Continued

W I—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
		2	38731. 94		1. 5			2	41104. 59		1. 5
		3	38943. 17		1. 4			0	41187. 38		0/0
		4	39082. 28		1. 24			6	41171. 44		
		1	39190. 48		1. 62			4	41198. 16		1. 23
		5	39275. 10		1. 27			6	41417. 48		1. 23
		0	39583. 29					3	41499. 43		1. 11
		2	39873. 70		1. 50			2	41533. 19		1. 06
		3	39874. 96		1. 50			3	41694. 34		1. 28
		2	39904. 14		1. 57			2	41734. 13		1. 1
		4	37146. 34		1. 1			4	41871. 94		1. 11
		5	37309. 13		1. 25			0	41965. 81		
		2	37466. 30		1. 28			2	41978. 61		0. 8
		3	37674. 08		1. 13			6	42239. 04		
		1	37773. 96					3	42251. 51		1. 32
		4	38001. 12		1. 1			1	42262. 30		1. 5
		3	38053. 05		1. 11			2	42449. 60		
		6	38203. 12					4	42450. 24		
		3	38206. 38					3	42514. 14		1. 22
		4	38259. 40					1	42573. 47		1. 3
		1	38355. 81					3	42601. 19		1. 12
		0	38576. 14					5	42826. 00		1. 11
		4	38743. 44					4	42910. 74		1. 18
		2	39030. 25					5	43034. 10		
		1	39183. 20		1. 01			0	43053. 86		0/0
		5	39361. 01		1. 13			1	43217. 32		1. 3
		5	39614. 05		1. 20			2	43227. 66		1. 3
		1	39636. 62		1. 44			4	43251. 00		1. 14
		3	39646. 41		1. 46			7	43411. 50		1. 20
		2	39707. 02		1. 00	5d <sup>4</sup> 6e(a'D)7e	e'D	1	43451. 98	1467. 86	2. 83
		4	39719. 96		1. 17			2	44919. 84	1576. 78	1. 9
		2	40011. 50		1. 0			3	46496. 62	1478. 92	1. 74
		4	40233. 97		1. 53			4	47975. 54	1379. 14	1. 68
		3	40269. 35		1. 03			5	49354. 68		1. 7
		1	40411. 12		1. 58			3	43478. 58		1. 3
		5	40476. 42		1. 04			2	43514. 67		0. 9
		4	40583. 07					4	43720. 87		
		3	40965. 85		0. 96			5	43741. 37		1. 09
		1	40770. 73		1. 28			3	43850. 84		1. 17
		2	40888. 39		1. 26			1	43892. 63		1. 05
		5	40911. 97		1. 03			5	43924. 25		1. 2
		3	40922. 83		1. 32			2	43976. 22		1. 15
								4	43985. 41		1. 24
								3	44081. 00		1. 2

## W I—Continued

## W I—Continued

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$
5d <sup>4</sup> 6a (a <sup>4</sup> D)7a	e <sup>4</sup> D	1	44353. 43		1. 03			4	47360. 19		
		2	44367. 50		1. 1			3	47361. 77		1. 3
		6	44390. 43		1. 23			2	47443. 53		
		3	44447. 03		1. 33			3	47433. 70		1. 29
		5	44543. 76		1. 3			6	47541. 55		1. 23
		2	44596. 33		1. 11			3	47593. 44		1. 2
		1	44737. 31		1. 1			4	47639. 33		1. 4
		6	44923. 90		1. 23			5	47850. 86		
		4	44940. 57		1. 20			4	47963. 63		1. 23
		7	44970. 73		1. 20			5	48133. 39		1. 2
		3	45014. 54		1. 3			3	48170. 67		
		2	45019. 03					2	48244. 29		1. 4
		4	45116. 77		1. 2			7	48350. 70		
		0	45225. 22			1233. 03	0/0	2	48318. 35		1. 4
		1	46453. 30			1620. 02		3	48326. 43		
		2	48073. 32			1577. 72	1. 55	1	48339. 96		
		3	49656. 04			1467. 10	1. 66	4	48676. 03		1. 20
		4	51123. 14				1. 4	6	48834. 63		
		4	45263. 55				1. 1	1	48733. 10		
		1	45373. 99					3	49073. 14		1. 26
		2	45433. 37				0. 63	4	49147. 95		
		5	45451. 63				1. 16	2	49151. 94		
		3	45551. 33				1. 30	5	49187. 93		1. 25
		3	45677. 69				1. 24	2	49370. 16		
		1	45760. 30					3	49417. 93		
		5	45789. 14				1. 19	1	49443. 63		
		4	45969. 10				1. 36	3	49514. 34		1. 3
		2	45903. 43				0. 5	2	49517. 96		
		3	46063. 03				1. 46	4	49636. 57		
		1	46104. 64					2	49699. 57		1. 9
		6	46106. 31					4	49733. 53		
		1	46231. 66					4	49733. 53		
		2	46337. 75				0. 8	0	49733. 53		
		3	46335. 43				1. 4	3	49866. 03		
		5	46506. 37				1. 33	5	50137. 55		1. 11
4	46625. 05				1. 14	3	50135. 70				
6	46673. 31				1. 13	4	50234. 63		1. 03		
2	46806. 45				1. 1	6	50429. 13?				
5	46854. 30				1. 21	2	50434. 54				
4	46931. 34				1. 0	1	50533. 54				
2	47073. 40				1. 3	2	50713. 90				
1	47255. 55					3	50800. 44		1. 0		
2	47337. 79				0. 9						

W 1—Continued

W 1—Continued

Config.	Desig.	J	Level	Interval	Obs. #	Config.	Desig.	J	Level	Interval	Obs. #
		5	50806.07					4	54118.78		
		4	50894.09					5	54310.80		
		4	50908.47		1. 3			1	54419.65		
		1	50949.07					3	54556.43		
		6	50981.90					6	54733.38		
		3	51078.18					2	54859.15		
		2	51182.41					6	54895.96		
		5	51290.75					4	54911.61		
		3	51800.53					1	54941.01		
		2	51806.90					5	55009.23		
		2	51893.35					2	55032.67		
		1	51763.44					4	55043.34		
		4	51856.10					2	55084.01		
		6	51978.63					4	55333.12		1. 45
		3	52015.28					3	55389.30		
		4	52059.78					5	55455.32		
		2	52064.02					5	55492.22		
		5	52081.09					3	55546.06		
		2	52152.62					2	55619.65		
		2	52183.20					7	55690.82		
		3	52255.90					5	55795.48		
		2	52284.76					2	55835.14		
		5	52395.52					1	55859.35		
		4	52436.44					4	55887.24		
		2	52503.41					4	55955.36		
		5	52774.14					5	55987.90		
		6	52855.98					3	56108.52		
		3	52943.48					4	56174.63		
		4	52992.89					4	56255.65		
		1	53042.02					5	56290.48		
		4	53112.29					3	56484.33		
		5	53194.25		1. 3			4	56502.10		
		6	53222.40					6	56596.61		
		4	53232.41					5	56573.54		
		3	53245.53					3	56717.16		
		3	53390.42					4	56831.67		
		2	53669.22					6	56832.90		
		1	53847.77					5	57143.42		
		4	53842.62					6	57143.92		
		5	53992.72					5	57560.81		
		2	53959.22					6	57702.30		

## W I—Continued

## W I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
		4	57803. 667					5	59203. 63		
		6	57919. 17					6	59410. 53		
		3	58091. 54					2	59422. 00		
		5	58179. 39					5	59673. 32		
		3	58206. 03					2	59999. 12		
		5	58562. 70					4	60385. 09		
		3	58644. 10					5	60741. 46		
		4	58777. 80					5	62154. 50		
		5	58903. 98								
		6	59128. 32			W II( <sup>3</sup> D <sub>3/2</sub> )	Limit		64400		
		4	59171. 71								

March 1956.

## W I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms	
5d <sup>1</sup> 6s <sup>2</sup>	{ a <sup>3</sup> P    a <sup>3</sup> D a <sup>1</sup> D    a <sup>3</sup> F    a <sup>1</sup> G    a <sup>3</sup> H	
	ns (n ≥ 6)	np (n ≥ 6)
5d <sup>2</sup> (a <sup>3</sup> S)nz	{ a <sup>3</sup> S a <sup>1</sup> S	s <sup>1</sup> P <sup>o</sup>
5d <sup>2</sup> (a <sup>1</sup> G)nz	a <sup>1</sup> G	
5d <sup>1</sup> 6s(a <sup>1</sup> D)nz	{ a <sup>1</sup> D a <sup>3</sup> D	s <sup>1</sup> P <sup>o</sup> s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>3</sup> D <sup>o</sup> s <sup>3</sup> F <sup>o</sup>

\*For predicted terms in the spectra of the W I isoelectronic sequence, see Vol. III, Introduction.

## W II

(Ta I sequence; 73 electrons)

Z=74

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>1</sup> 6s<sup>2</sup> <sup>1</sup>D<sub>3/2</sub>a<sup>1</sup>D<sub>3/2</sub>

K

I. P.

volts

In 1938 a preliminary analysis was published by Laun. Later Kiess reobserved the spectrum and Zeeman effect, at the National Bureau of Standards, in the region longer than 2000 Å. Short of 2000 Å, observations were made by Boyce with the Carnegie Spectrograph. Laun has measured these spectrograms and extended his earlier analysis especially for inclusion here. There are more than 1,250 classified lines between 1756.684 Å and 6219.77 Å.

Mack and Mrs. Taschek have added seven new levels and the three-place g-values listed in the table. Their Zeeman data have been taken from observations made with the Bitter magnet, at the Massachusetts Institute of Technology.

## W II—Continued

The authors have a decided preference for arrangement by levels throughout the spectrum because the departure from *LS*-coupling makes it difficult to assign term designations except for the lowest terms. The writer has retained the listing by terms for the lower levels in conformance with the general format adopted for these Volumes. A ready comparison of term intervals and observed *g*-values can thus be made with similar spectra.

The higher levels are listed in increasing numerical order. By analogy with Mo II, and from an examination of the observed *g*-values, Kees and the writer have assigned tentative designations to a number of these miscellaneous levels. These should be confirmed by further study. They are consistent with theoretical calculations made by Trees.

No series have been detected in W II. From a study of screening constants, Finkelnburg and Humbach interpolate an ionization potential of  $17.7 \pm 0.5$  volts.

## REFERENCES

- D. D. Laun, J. Research Nat. Bur. Std. **21**, 207, RP 1125 (1938). (T) (C L)  
 W. Finkelnburg and W. Humbach, Naturwiss. **42**, 35 (1955). (I P)  
 D. D. Laun, unpublished material (March 1956). (T) (C L)  
 J. E. Mack and I. M. Taschek, unpublished material (April 1956). (T) (Z E)

## W II

## W II

Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>		
5d <sup>4</sup> ( <sup>3</sup> D)6s	a <sup>3</sup> D	0½	0.00		3.186	5d <sup>5</sup>	<sup>4</sup> F	3½	22194.08		1.119		
		1½	1518.78	1518.78	1.839								
		2½	3172.52	1653.74	1.639			5d <sup>4</sup> ( <sup>3</sup> F)6s	<sup>3</sup> P	1½	22503.06		1.22
		3½	4716.32	1543.80	1.563								
	4½	6147.16	1430.84	1.522	5d <sup>4</sup> ( <sup>3</sup> D)6s	<sup>1</sup> D	3½	23046.80					
5d <sup>5</sup>	a <sup>3</sup> S	2½	7420.43		1.913	5d <sup>5</sup>	<sup>4</sup> F	4½	23234.87		1.249		
5d <sup>4</sup> ( <sup>3</sup> F)6s	a <sup>3</sup> F	1½	8711.26	2589.82	0.624	5d <sup>5</sup>	<sup>1</sup> D	2½	23450.50		1.297		
		2½	11301.08	2110.88	1.084	5d <sup>5</sup>	<sup>1</sup> D	3½	23803.84				
		3½	13411.96	1445.26	1.186								
		4½	14857.22		1.234								
5d <sup>4</sup> ( <sup>3</sup> P)6s	a <sup>3</sup> P	0½	8832.66	1759.86	2.383	5d <sup>5</sup>	<sup>3</sup> I	5½	23955.40		1.10		
		1½	10592.52	2841.58	1.471	5d <sup>5</sup>	<sup>4</sup> F	1½	24991.56				
		2½	13434.10		1.526	5d <sup>5</sup>	<sup>3</sup> P	1½	25169.87		1.64		
5d <sup>4</sup> ( <sup>3</sup> D)6s	a <sup>3</sup> D	0½	13173.38	1460.98	0.455	5d <sup>4</sup> ( <sup>3</sup> H)6s	<sup>3</sup> H	4½	26158.74				
		1½	14634.36	333.46	1.183	5d <sup>5</sup>	<sup>3</sup> P	2½	26227.00				
		2½	14967.82	179.20	1.013								
		3½	15147.02		0.872								
5d <sup>4</sup> ( <sup>3</sup> G)6s	a <sup>3</sup> G	2½	16234.84	354.83	0.995	5d <sup>4</sup> ( <sup>3</sup> H)6s	<sup>3</sup> H	5½	26929.34				
		3½	16589.67	-36.53	1.153	5d <sup>4</sup> ( <sup>3</sup> D)6p	<sup>3</sup> F°	0½	36185.35		0.678		
		4½	16553.14	883.88	1.137								
		5½	17437.02		1.181								
5d <sup>4</sup> ( <sup>3</sup> H)6s	<sup>3</sup> H	3½	18000.70		1.098	5d <sup>4</sup> ( <sup>3</sup> F)6p	<sup>3</sup> S°	0½	38576.39		1.614		
5d <sup>5</sup>	<sup>3</sup> D	1½	18990.96		0.90	5d <sup>4</sup> ( <sup>3</sup> D)6p	<sup>3</sup> F°	1½	39129.41		1.147		
5d <sup>5</sup>	<sup>3</sup> G	4½	19070.68		1.102			2½	39836.81		0.889		
5d <sup>5</sup>	<sup>3</sup> G	2½	19276.52		0.997	5d <sup>4</sup> ( <sup>3</sup> D)6p	<sup>3</sup> F°	2½	40049.45		1.292		
5d <sup>4</sup> ( <sup>3</sup> F)6s	<sup>3</sup> P	0½	19404.08		0.64			1½	42298.80		1.498		
5d <sup>4</sup> ( <sup>3</sup> H)6s	<sup>3</sup> H	6½	19442.54					3½	42390.87		1.161		
5d <sup>5</sup>	<sup>3</sup> D	2½	19637.38		1.102			2½	44354.82		1.390		
5d <sup>5</sup>	<sup>3</sup> G	3½	20039.74		1.117			0½	44455.18		-0.217		
5d <sup>5</sup>	<sup>3</sup> D	1½	20455.93					4½	44758.10		1.270		
5d <sup>5</sup>	<sup>3</sup> G	5½	20524.35		1.197	5d <sup>4</sup> ( <sup>3</sup> D)6p	<sup>3</sup> F°	3½	44877.18		1.277		
5d <sup>4</sup> ( <sup>3</sup> H)6s	<sup>3</sup> H	4½	20780.38		1.065			1½	44911.63		1.221		
5d <sup>5</sup>	<sup>3</sup> F	2½	22139.97		1.06			0½	45457.02		0.519		

## W II—Continued

## W II—Continued

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$			
5d <sup>+</sup> (D)6p	°F°	1½	45553.70		1.033	5d <sup>+</sup> (H)6p	°H°	2½	54375.89		0.623			
		2½	46175.48		1.452			0½	54408.60	1.141				
		3½	46355.40		1.236			3½	54488.57					
		4½	46493.43		1.311			2½	54704.61					
		0½	46825.87					5½	54858.58					
		1½	47179.94		1.007			2½	55168.30					
		2½	47413.33		1.111			4½	55392.37			1.061		
		1½	47533.64					1½	55488.01					
		2½	48284.48		1.366			1½	56084.30			1.021		
		5½	48332.73					5½	56376.45					
		3½	48830.70		1.006			5d <sup>+</sup> (H)6p	°H°			6½	56439.58	
		1½	48982.86									3½	56612.74	1.22
		3½	49124.52		1.499							3½	56768.61	1.147
		0½	49154.50									2½	56874.99	0.815
		4½	49181.04		1.409							1½	56932.27	
		2½	49242.10		1.510							4½	57089.46	
		2½	50292.33		1.334							3½	57729.92	1.184
1½	50430.25			2½	57856.70	1.36								
4½	50863.05		1.194	1½	58007.60									
3½	51045.25			2½	58336.98									
1½	51264.40		1.58	3½	58709.56									
2½	51432.03		1.301	1½	58747.94									
5d <sup>+</sup> (D)6p	°F°	5½	51495.00		1.054	5½	58891.74			1.144				
		3½	51863.03		0.937	3½	59276.81			1.102				
		2½	52087.02			4½	59399.34			1.179				
		3½	52275.22		1.297	3½	59869.14			1.125				
		0½	52355.11		0.981	3½	59933.66							
		4½	52567.15			5½	60212.84	1.130						
		0½	52593.72		1.14	3½	60256.45							
		3½	52901.74		1.374	4½	60278.71							
		2½	53113.52		1.262	2½	60656.51							
		1½	53322.71		1.357	4½	61055.80							
		3½	53332.07		0.968	5½	61240.81	1.120						
		4½	53369.97		1.086	3½	61320.29							
		1½	53422.22		0.976	4½	61360.55							
		°P°?	0½	53440.17		2.038	5½	61589.46	1.149					
			2½	54022.24			6½	61602.16						
			4½	54056.54		1.122	2½	62333.20						
		°P°?	1½	54127.20		1.606	4½	64516.37						
5½	54222.06													

April 1956.

## RHENIUM

## Re I

75 electrons

Z=75

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4 4d^{10} 4f^{14} 5s^2 5p^4 5d^5 6s^2 \text{ } ^6S_{7/2}$  $6s^2 \text{ } ^6S_{7/2}$ , 63530 K

I. P. 7.87 volts

The Re I spectrum furnishes one of the finest examples of international collaboration of any spectrum on the "Atomic Energy Levels" program. The present analysis represents the work of American, Dutch, and Spanish spectroscopists. Klinkenberg, Meggers, Velasco, and Catalán have completed their study of Re I especially for inclusion here.

The observational material consists of a new description of the spectrum supplied by Meggers, including some 4200 lines. This was supplemented by measurements of Velasco from spectrograms covering the vacuum region, taken by Catalán at Princeton University. The present line list extends from about 1700 Å to 11788.9 Å and includes 2764 classified lines arising from permitted transitions among 71 even and 211 odd levels.

The first measurements of hyperfine structure in Re I were made by Meggers. In his 1931 paper he states that about 25 percent of the 2,000 or more lines . . . show hyperfine structure of 2 to 6 components.

Observed  $g$ -values are known for 75 percent of the energy levels. These are from Zeeman spectrograms taken by Meggers at the Massachusetts Institute of Technology, through the courtesy of G. R. Harrison. The patterns were measured by Meggers, and these data were given to Catalán and Velasco who, in turn, derived the  $g$ -values.

The present interpretation is the result of all three groups, and greatly extends the earlier work on Re I by these authors. Observed intersystem combinations connect the systems of terms of different multiplicity. For a number of terms in the table the  $LS$ -designation assignments are based on theoretical work by Trees. He points out that the hyperfine structure exhibited by the  $a^6S$  term probably results from the influence of the  $5d^5(^6S)6s^2 \text{ } ^6S$  term. His work is still in progress.

The limit is derived from two triplet terms, interpreted as combinations of  $6p^4P^o$  with  $7s^2 \text{ } ^6S$  and  $8s^2 \text{ } ^6S$ , as reported in 1931. No improved series have been found from the present work.

## REFERENCES

- W. F. Meggers, *Bur. Std. J. Research* **6**, 1027, RP 322 (1931). (I P) (T) (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 W. F. Meggers, *J. Research Nat. Bur. Std.* **49**, 187, RP 2355 (1952). (hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 31 (1955). (Summary hfs)  
 R. E. Trees, unpublished material (November 1956).  
 P. F. A. Klinkenberg, W. F. Meggers, R. Velasco, and M. A. Catalán, *J. Research Nat. Bur. Std.* **59**, 319, RP 2804 (1957). (I P) (T) (C L) (Z E)



Re I

Re I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
5d <sup>8</sup> 6s <sup>2</sup>	a *G	2½	0.00		1.950	5d <sup>8</sup> (G) 6s	b *G	2½	33281.65		
5d <sup>8</sup> 6s <sup>2</sup>	a *P	2½	11583.96		1.278		3½	33823.66		542.01	
		1½	13826.12	-2242.16	1.485		4½	34194.27		370.61	
		0½	15165.89	-1339.77	2.368		5½	33317.57		-876.70	
5d <sup>8</sup> (D) 6s	a *D	4½	11754.52		1.545	5d <sup>8</sup> 6s(a *D) 6p	y *P°	1½	33589.18		2.335
		3½	14216.86	-2462.34	1.567		2½				
		2½	15770.42	-1553.56	1.309		3½				
		1½	16827.51	-557.09	1.706	5d <sup>8</sup> 6s(a *S) 6p	z *P°	2½	35267.94		1.385
		0½	17238.80	-910.79	2.521		1½	37915.87	-2647.93	1.495	
5d <sup>8</sup> 6s <sup>2</sup>	a *G	2½	14621.46		1.151		0½	37797.95	117.92	2.620	
		3½	15058.19	436.73	1.153	5d <sup>8</sup> 6s(a *G) 6p	z *G°	1½	37063.65		0.626
		4½	16619.28	1561.09	1.175		2½	37381.41	317.76	0.990	
		5½	16307.15	-312.13	1.242		3½	38580.74	1139.33	1.216	
5d <sup>8</sup> 6s <sup>2</sup>	a *D	3½	17330.82		1.255		4½	38994.73	473.99	1.25	
		2½	19457.89	-2127.07	1.361		5½	39916.39	921.66	1.314	
		1½	20481.73	-1023.84	1.451		6½	40493.54	576.95	1.364	
		0½	19757.91	723.82	0.983	5d <sup>8</sup> 6s(a *D) 6p	z *F°	0½	37381.41		0.745
5d <sup>8</sup> 6s(a *S) 6p	z *P°	2½	18950.1		2.274		1½	39926.75	2545.34	1.335	
		3½	20447.8	1497.7	1.926		2½	39064.92	-861.83	1.335	
		4½	23631.82	3184.0	1.768		3½	37765.65	-1299.27	1.335	
5d <sup>8</sup> 6s <sup>2</sup>	a *F	3½	21775.40		1.135		4½	39655.99	1890.34	1.444	
		2½	24425.40	-2650.00	1.067		5½	39196.74	-459.25	1.405	
5d <sup>8</sup> 6s <sup>2</sup>	a *G	4½	22160.04		1.198		2½	37697.68		1.219	
		3½	24724.22	-2564.18	1.03		2½	39670.52		1.107	
5d <sup>8</sup> 6s <sup>2</sup>	a *D	1½	22422.83		0.781	5d <sup>8</sup> 6s(a *S) 6p	z *P°	3½	39344.75		1.223
		2½	23154.81	731.98	1.189		2½	43569.36	-3724.61	1.504	
							1½	43702.20	-132.84	1.962	
5d <sup>8</sup> 6s <sup>2</sup>	a *I	5½	23956.00		0.995		1½	40808.85		1.351	
		6½	26348.96	2392.96	1.100		0½	40810.17			
5d <sup>8</sup> 6s <sup>2</sup>	a *F	1½	26131.57		0.650		2½	40821.83		1.126	
		2½	28030.32	1898.75	1.12						
		3½	28542.13	511.81							
		4½	27514.31	-1027.82	1.13	5d <sup>8</sup> 6s(a *P) 6p	y *D°	4½			
5d <sup>8</sup> (P) 6s	b *P	2½	26661.43		1.32		3½	40946.53		1.534	
		1½	30526.60	-3865.17			2½	41313.02	-366.49	1.278	
		0½					1½	41557.08	-244.06	1.655	
5d <sup>8</sup> (H) 6s	a *H	6½	27130.14		1.208		3½	41163.91		1.325	
		5½	27243.88	-113.74	1.18		4½	41453.18		1.372	
		4½	27161.35	82.53			3½	41843.85		1.190	
		3½	28809.87	-1648.52	0.93						
5d <sup>8</sup> (D) 6s	b *D	3½	27141.13		1.34	5d <sup>8</sup> 6s(a *P) 6p	z *D°	0½	41991.56		0.61
		2½	29900.38	-2659.25	1.17		1½	42254.19	262.63	1.578	
		1½	27827.65	1972.73	0.888		2½	43044.02	789.83	1.449	
		0½	30131.57	-2303.92			3½	43407.87	363.85	1.368	
5d <sup>8</sup> 6s <sup>2</sup>	a *S	0½	27384.80				4½	42140.06		1.249	
5d <sup>8</sup> 6s(a *S) 6p	z *P°	3½	28889.72		1.709	5d <sup>8</sup> 6s(a *S) 7s	e *S	3½	42598.27		1.957
		2½	28354.18	35.54	1.871		1½	43341.85		0.975	
		1½	28961.55	-107.38	2.333		1½?	43409.11			
5d <sup>8</sup> 6s <sup>2</sup>	a *H	5½	30559.91		1.07		4½	43453.30		1.336	
		4½	31399.30	-839.39			3½	43815.01		1.348	
5d <sup>8</sup> (F) 6s	b *F	4½	30645.33				2½	43949.98		1.385	
		3½	31186.08		1.17		5½	43996.30		1.26	
		2½	31460.62	-274.54			2½	44054.21			
		1½					1½	44148.45		1.573	
5d <sup>8</sup> 6s <sup>2</sup>	b *F	3½	31982.99				4½	44224.58		1.244	
		2½	32435.14	-452.15			2½	44308.73		1.223	
5d <sup>8</sup> 6s(a *D) 6p	z *D°	0½	32443.61		1.762		0½	44416.33		0.15	
		1½	32591.63	148.02	1.500						
		2½	33408.73	817.10	1.454						
		3½	34520.25	1111.52	1.440						
		4½	35923.02	1402.77							

Re I—Continued

Re I—Continued

Config.	Desig.	J	Level	Interval	Obs. #	Config.	Desig.	J	Level	Interval	Obs. #			
5d <sup>8</sup> 6s(a <sup>1</sup> S)7s	e <sup>1</sup> S	2½	44703.37		1.866			1½	49850.08		1.268			
		1½	44780.09		1.330			5½	49886.07		1.075			
		3½	44901.15		1.253			2½	49840.96					
		2½	44902.60					2½	49873.11		1.344			
		3½	44946.18		0.905			4½	49888.61		1.058			
5d <sup>8</sup> ( <sup>1</sup> D)6p	y <sup>1</sup> F <sup>o</sup>	5½	45088.68	-1291.62 -1558.30 -1090.22 -644.45 -192.28	1.40	5d <sup>8</sup> 6s(a <sup>1</sup> S)6d	e <sup>1</sup> D	3½	49883.88		1.175			
		4½	45374.85		1.341			1½	49895.57		1.989			
		3½	47888.55		1.31			2½	50110.88		1.765			
		2½	49088.77		1.495			2½	50158.87		1.422			
		1½	49887.85		1.27			4½	50198.94		1.345			
		0½	49889.55		0.410			1½	50883.88		1.234			
		1½	45181.81					3½	50810.70		1.170			
		2½	45338.03					1½	50332.60	8.02 18.61 36.49 68.62	2½	50340.62		2.027
		4½	45343.57					1.131	3½		50359.23		1.820	
		3½	45468.83					1.300	4½		50395.72		1.690	
		0½	45817.86					2.197	5½		50464.34		1.629	
		1½	45876.34					1.384	4½		50401.01		1.055	
	5½	45904.85		1.175	2½		50463.80		1.144					
	3½	45937.18		1.298	1½		50571.08		0.935					
	2½	46118.84		1.405	5½		50663.51		1.220					
	1½	46141.11		0.716	3½		50889.00		1.240					
	3½	46358.99		1.271	2½		50934.18		1.385					
	2½	46509.40		1.371	2½		50973.17		1.489					
	2½	46649.48		1.334	5d <sup>8</sup> 6s(a <sup>1</sup> S)6d		e <sup>1</sup> D	4½	50994.15	5.68 -47.09 4.83 -19.14	4½	50988.47		1.548
	1½	46733.38		1.858				3½	51035.56			1.558		
	1½	47004.34		1.285				2½	51030.73			1.546		
	2½	47101.61		0.893				1½	51049.87					
	1½	47178.1						0½	51049.87					
	0½	47178.9		2.215				4½	51027.96					
	4½	47305.73		1.106				1½	51183.00			1.456		
	3½	47358.36		1.151				4½	51193.49					
	6½	47506.79		1.26				3½	51229.64			1.231		
	0½	47664.74		1.50				2½	51477.15			0.97		
3½	47689.01		1.196	4½		51486.89			1.20					
1½	47703.78		1.443	1½		51578.05			0.92					
1½	47779.91		1.362	2½	51590.08		1.02							
5½	47859.83		1.20	3½	51647.88		1.139							
2½	47899.88		1.217	1½	51874.43		1.29							
2½	47970.88		1.169	5½	51945.94									
3½	48184.80		1.252	3½	51955.08		1.200							
5½	48569.63		1.27	4½	52001.37		1.357							
3½	48786.35		1.53	6½	52128.35		1.17							
3½	48857.60		1.24	2½	52218.11		1.208							
3½	49087.85		1.499	6½	52227.80									
4½	49170.78		1.135	1½	52278.88		1.422							
								3½	52573.18					

Re I—Continued

Re I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g	
5d <sup>4</sup> 6s <sup>2</sup> (a <sup>4</sup> S) 8s	f <sup>8</sup> S	0%	52489. 45		1. 815	5d <sup>4</sup> 6s <sup>2</sup> (a <sup>4</sup> D) 7s	f <sup>4</sup> D	4%	56411. 82			
		2%	52472. 31					0%	56451. 87		1. 287	
		3%	52610. 01					6%	56523. 41			1. 25
		4%	52741. 30					3%	56699. 41			
		3%	52857. 44		1. 20			2%	56822. 75			
		2%	52881. 34		1. 05			1%	56866. 06			0. 479
		1%	52951. 59		1. 14			0%	57073. 31			0. 94
		1%	52954. 46					3%	57090. 29			
		2%	53058. 74		1. 184			3%	57172. 70			1. 19
		0%	53125. 19		1. 145			4%	57290. 86			1. 19
		0%	53228. 28		1. 372			3%	57371. 75			
		1%	53293. 72		1. 323			5%	57391. 44			1. 30
		4%	53320. 95		1. 38			6%	57452. 83			
		3%	53335. 94		1. 290			2%	57524. 31			1. 40
		4%	53379. 35		1. 09			4%	57656. 85			1. 19
		2%	53389. 73		1. 19			0%	57665. 43			
		3%	53392. 00					1%	59268. 79	1608. 36		
		5%	53467. 57		1. 20			2%	59268. 79	-1208. 15		
		3%	53738. 46		1. 203			3%	59368. 90	308. 26		
		1%	53842. 53		1. 169			1%	58050. 58			1. 111
		0%	53928. 90		1. 113			3%	58260. 03			1. 43
		5%	53948. 73		1. 13			4%	58279. 31			
		3%	54018. 19		0. 723			2%	58332. 42			
		3%	54086. 69		1. 22			1%	58442. 51			1. 32
		3%	54177. 25		1. 270			0%	58541. 88			
		2%	54228. 18		1. 290			5%	58619. 32			1. 165
		2%	54409. 61		1. 069			1%	59127. 30			1. 20
		1%	54467. 40		1. 361			2%	59171. 73			
3%	54513. 84			1%	59223. 45							
5%	54554. 61			4%	59394. 29							
6%	54729. 52			5%	59412. 46			1. 15				
4%	54813. 12		1. 185	3%	59784. 67			1. 100				
2%	54823. 02		1. 182	3%	59926. 91							
0%	54928. 85		1. 314	2%	60737. 28			1. 22				
5%	55223. 85		1. 149	2%	60837. 10							
3%	55454. 42		1. 17	5%	62142. 97							
6%	55776. 51		1. 056	3%	62350. 13							
3%	55901. 20		1. 22	3%	62428. 51			1. 17				
1%	55902. 01		0. 447	5%	63267. 63			1. 135				
4%	55912. 42		1. 06									
4%	56307. 97											
						Re II(7E <sub>2</sub> )	Limit		63530			
									63772. 40			
									63934. 76		1. 07	

## Re I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms						
5d <sup>9</sup> 6s <sup>2</sup>	a <sup>4</sup> S	a <sup>4</sup> P	a <sup>4</sup> D	a <sup>4</sup> F	a <sup>4</sup> G	a <sup>4</sup> H	a <sup>4</sup> I
	a <sup>4</sup> S		a <sup>4</sup> D	a <sup>4</sup> F	a <sup>4</sup> G		
			b <sup>4</sup> F				
	ns (n ≥ 6)			np (n ≥ 6)		nd (n ≥ 6)	
5d <sup>9</sup> 6s(a <sup>4</sup> S)nz	c, f <sup>4</sup> S						a <sup>4</sup> D
	a <sup>4</sup> S						a <sup>4</sup> D
5d <sup>9</sup> ( <sup>4</sup> D)nz			a <sup>4</sup> D			y <sup>4</sup> F <sup>o</sup>	
			b <sup>4</sup> D				
5d <sup>9</sup> 6s(a <sup>4</sup> S)nz							
5d <sup>9</sup> 6s(a <sup>4</sup> G)nz							
5d <sup>9</sup> 6s(a <sup>4</sup> P)nz							
5d <sup>9</sup> ( <sup>4</sup> P)nz		b <sup>4</sup> P					
5d <sup>9</sup> ( <sup>4</sup> H)nz							a <sup>4</sup> H
5d <sup>9</sup> ( <sup>4</sup> F)nz				b <sup>4</sup> F			
5d <sup>9</sup> ( <sup>4</sup> G)nz					b <sup>4</sup> G		
5d <sup>9</sup> 6s(a <sup>4</sup> D)nz			f <sup>4</sup> D				
						y <sup>4</sup> P <sup>o</sup>	
						z <sup>4</sup> D <sup>o</sup>	
						z <sup>4</sup> F <sup>o</sup>	

\*For predicted terms in the spectra of the Re I isoelectronic sequence, see Volume III, Introduction.

## Re II

(W I sequence; 74 electrons)

Z=75

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>9</sup> 6s<sup>2</sup> <sup>7</sup>S<sub>2</sub>

6s<sup>2</sup> <sup>7</sup>S<sub>2</sub>

K

I. P.

volts

In 1952, Meggers published a description of the Re II spectrum that included some 1800 lines in the range 2000.47 Å to 6026.60 Å. Later, Catalán extended the observations in the short-wave region to 1500 Å, with the vacuum spectrograph at Princeton University; and Sales provided measurements in this region.

On the basis of this new description, the present analysis has been carried out by Meggers, Catalán, and Sales. Nine "odd" levels have also been contributed by Tech.

The observed *g*-values are from Zeeman spectrograms taken by Meggers with the Bitter magnet at the Massachusetts Institute of Technology, through the courtesy of G. R. Harrison.

Approximately 1000 lines are classified, but the assignment of term designations has not been attempted in many cases because of the marked departure from *LS*-coupling and the effect of configuration-interaction. The absence of the low 5d<sup>9</sup> <sup>4</sup>D term is conspicuous, but persistent search has failed to reveal this term.

No series are known in Re II. From a study of screening constants Finkelnburg and Humbach have extrapolated an ionization potential of 16.6 ± 0.5 volts, which places the limit near 133900 K.

## REFERENCES

- W. F. Meggers, J. Research Nat. Bur. Std. 49, 187, RP 2355 (1952).  
 J. L. Tech, unpublished material (August 1955). (T) (C L)  
 W. Finkelnburg und W. Humbach, Naturwiss. 42, 35 (1955). (I P)  
 W. F. Meggers, M. A. Catalán, and M. Sales, unpublished material (November 1954), (October 1955), (December 1957). (T) (C L) (Z E)

## Re II

## Re II

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$
5d <sup>2</sup> ( <sup>3</sup> S)6s	a <sup>3</sup> S	3	0.0		2.05	5d <sup>2</sup> ( <sup>3</sup> S)6p	z <sup>1</sup> P <sup>o</sup>	2	43937.7		2.19
5d <sup>2</sup> 6s	a <sup>1</sup> D	0	13777.3	1046.7	0/0			3	45147.6	1209.8	1.79
		1	14824.0	-471.8	1.50			4	50680.9	5533.4	
		2	14352.2	578.3	1.53	5d <sup>2</sup> ( <sup>3</sup> S)6p	z <sup>3</sup> P <sup>o</sup>	1	49439.3	3237.7	2.42
		3	14930.5	-47.9	1.48			2	52677.0	1125.1	1.82
		4	14882.6		1.48			3	53808.1		1.66
5d <sup>2</sup> ( <sup>3</sup> S)6s	a <sup>3</sup> S	2	17223.5		1.71	5d <sup>2</sup> ( <sup>1</sup> F)6p	z <sup>1</sup> D <sup>o</sup>	0	54488.0	1904.1	1.291
5d <sup>2</sup> ( <sup>1</sup> G)6s	a <sup>1</sup> G	2	18845.8	293.9	0.64			1	56378.1	766.5	1.80
		3	19139.7	1323.5	1.12			2	57138.6	2191.3	1.145
		4	20463.2	513.2	1.15			3	58329.9	3529.1	1.27
		5	20976.4	1254.7	1.26	5d <sup>2</sup> ( <sup>1</sup> F)6p	v <sup>1</sup> P <sup>o</sup>	3	62859.0		
		6	22031.1		1.24			2	55059.8	-91.6	1.81
5d <sup>2</sup> ( <sup>1</sup> F)6s	a <sup>1</sup> F	3	21629.1	-1711.7	1.26			1	55150.8		2.27
		2	23340.8	-381.6	1.67	5d <sup>2</sup> ( <sup>1</sup> F)6p	z <sup>3</sup> F <sup>o</sup>	1			1.083
		1	23722.4		1.903			2	55444.0	1606.2	1.135
		2	22544.7		1.20			3	57050.8	2545.9	1.177
		1	23146.2		1.10			4	59696.1	4686.2	1.31
		5	23381.2		1.19	5d <sup>2</sup> ( <sup>1</sup> F)6p	z <sup>1</sup> G <sup>o</sup> ?	2			
		4	23893.5		1.275			3	57429.4	1852.4	1.056
		3	24763.1		1.41			4	59281.8	1645.0	1.625
5d <sup>2</sup> ( <sup>1</sup> G)6s	a <sup>1</sup> G?	3	25321.2	916.1	0.90			5	60928.8		1.18
		4	26237.3	-249.8	1.12	5d <sup>2</sup> ( <sup>1</sup> F)6p	z <sup>1</sup> P <sup>o</sup>	0			
		5	25987.5		1.22			1			
		1	26864.3		1.46			2	59388.3		1.520
		2	26767.6		1.09			3	59665.8		1.19
		3	27627.7		1.23			1	60487.8		1.128
		3	27746.0		1.22			3	60525.0		
		4	28095.3		1.22			2	62057.3		1.17
		2	29077.3					1	617.8		1.52
		1	29427.2		1.368			3	6114.9		1.243
		4	29638.9		1.292			3	63043.3		1.34
		2	29728.1		0/0			4	63187.7		1.27
		0	29773.2		1.398			2	63464.8		1.50
		3	30224.8		1.26			4	64407.8		1.21
		3	30717.5		1.188			2	64411.4		0.812
		4	30982.5		1.42			3	64948.8		1.36
		2	31012.8		1.08			3	64986.9		
		4	32206.5		1.114			1	65011.0		0.796
		3	32257.8		1.15			3	65572.3		1.28
		5	32344.7		1.06			2	65674.0		
		2	32875.8		1.54			3	66100.3		1.40
		4	32957.3		1.188			3	66248.1		
		4	33169.3					4	66371.8		1.38
		3	33989.8		1.23			4	66476.8		
		2	34937.2		1.21			5	66514.6		1.18
		5	36063.5					5	66536.7		
		3	36993.4					2	66730.5		1.274
		3	37318.5					2	66782.6		
		2	37382.0		1.071			5	67049.1		1.29

Re II—Continued

Re II—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
		1	67876.1		1.292			2	71396.8		
		3	67914.0		1.194			2	71732.0		1.40
		4	67922.1		1.15			2	71942.0		
		5	68252.5		1.16			2	72134.8		
		3	68354.8					2	72536.8		
		2	68322.4					3	72246.6		
		3	68322.0					3	72552.9		
		4	68376.7					4	74064.4		
		3	69142.5		1.053			3	74070.8		
		2	69170.0		1.30			2	74322.8		
		4	69222.6		1.35			3	74422.2		
		3	69302.8		1.677			3	74632.0		
		3	69722.2					3	75117.0		
		4	69776.3		1.25			2	75227.1		
		4	69792.2					4	76137.1		
		1	70137.8		1.637			3	76327.2		
		3	70811.0		1.17			3	77190.7		1.11
		1	71022.7					3	77222.4		
		4	71122.5		1.19			3	77275.9		

December 1957.

Re II Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>2</sup> +	Observed Terms	
5d <sup>2</sup> 6s <sup>2</sup>	a <sup>1</sup> D	
	ns (n ≥ 6)	np (n ≥ 6)
5d <sup>2</sup> ( <sup>1</sup> S)ns	{ a <sup>1</sup> S a <sup>1</sup> G a <sup>1</sup> G? a <sup>1</sup> P	s <sup>1</sup> P <sup>o</sup> s <sup>1</sup> P <sup>o</sup>
5d <sup>2</sup> ( <sup>3</sup> G)ns		y <sup>1</sup> P <sup>o</sup> s <sup>1</sup> P <sup>o</sup>
5d <sup>2</sup> ( <sup>1</sup> F)ns		
5d <sup>2</sup> ( <sup>3</sup> F)ns		s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> F <sup>o</sup> s <sup>1</sup> G <sup>o</sup> ?

\*For predicted terms in the spectra of the W I isoelectronic sequence, see Volume III, Introduction.

## OSMIUM

## Os I

76 electrons

Z=76

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^8 6s^2 {}^1D_2$  $\alpha$   ${}^1D_2$ , 70450 K

I. P. 8.7 volts

As early as 1900, "long before any theory of complex spectra had been developed", Snyder noted regularities among 55 Os I lines, but he never published his results. In 1934 Albertson extended the analyses to include 137 levels and more than 1050 classified lines. In 1938 he reported that he had reobserved the spectrum at the Massachusetts Institute of Technology with high dispersion and recorded more than 4500 lines, of which 2169 were classified as combinations among 234 energy levels. He derived the limit from the two-member series  $6s^2 {}^1D_2$ ,  $7s {}^1D_2$  by means of a Rydberg formula.

In 1949 van den Bosch made more complete observations at the Massachusetts Institute of Technology, including Zeeman spectrograms. He has, also, had access to Albertson's unpublished material and, in collaboration with van Kleef, has extended the number of energy levels to 257, and furnished the present list in advance of publication especially for inclusion here. All the tabular  $g$ -values are from van den Bosch. His present line list covers the range 2000 Å to 8645 Å and includes approximately 6000 lines, of which 2387 are classified. Owing to the departure from  $LS$ -coupling and to the effect of configuration-interaction most of the levels have not been assigned  $LS$ -term designations.

## REFERENCES

- W. Albertson, *Phys. Rev.* **45**, 304 (1934). (I P) (T) (C L)  
 W. Albertson, *Phys. Rev.* **53**, 940 (A) (1938), and unpublished material. (T) (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **34**, No. 2, 71 (1962). (Summary hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 31 (1955). (Summary hfs)  
 J. C. van den Bosch and Th. A. M. van Kleef, unpublished material (August 1956). (T) (C L) (Z E)

Os I

Os I

Config.	Design.	J	Level	Interval	Obs. g	Config.	Design.	J	Level	Interval	Obs. g					
5d <sup>5</sup> 6s <sup>2</sup>	a <sup>1</sup> D	4	0.00		1.44			5	24291.97		0.80					
		3	4159.32	-4159.32	1.47			2	25069.71							
		2	2740.49	1418.83	1.44			3	25593.94							
		1	5766.14	-3025.65	1.47			4	25601.55							
		0	6092.79	-326.65				2	27954.38	2.04						
5d <sup>7</sup> ( <sup>1</sup> F) 6s	a <sup>1</sup> F	5	5143.92	-3598.91	1.38			3	28139.52		1.24					
		4	8742.83	-2635.17	1.31			4	28601.55							
		3	11378.00	1212.02	1.26			2	28139.52	1.24						
		2	10165.98	-2854.09	1.45			3	28351.77	1.59						
		1	13020.07		0.31			2	28371.68	1.74						
5d <sup>7</sup> ( <sup>1</sup> F) 6s	a <sup>1</sup> F	4	11030.58	-3060.79	1.16	5d <sup>5</sup> 6s(a <sup>1</sup> D) 6p	s <sup>1</sup> P <sup>o</sup>	4	28411.95		-39.91	1.59				
		3	14091.37	1316.99	1.10			3	28371.68	1.74						
		2	12774.38		1.00			2								
		2	13364.83		0.94			2	28411.95							
		2	14848.05		1.08			2	28809.97							
5d <sup>5</sup> 6s <sup>2</sup>	a <sup>1</sup> H	5	14338.99	-509.06	1.13	5d <sup>5</sup> 6s(a <sup>1</sup> D) 6p	s <sup>1</sup> F <sup>o</sup>	6	29099.41		1.48					
		6	14852.33	513.34	1.14			5								
		2	15222.57		1.61			4								
		3	15390.76		1.54			3								
		2	16212.41		1.74			2								
5d <sup>7</sup> ( <sup>1</sup> P) 6s	a <sup>1</sup> P	3	15390.76		1.54			1			1.40					
		2						0								
		1						3	29381.65							
		1	16212.41		1.74			4	29394.30							
		1	17667.34		1.40			2	30056.79							
5d <sup>7</sup> ( <sup>1</sup> P) 6s	a <sup>1</sup> P	2						2	30078.88		1.48					
		1	18301.40					5	30279.85	1.47						
		3	18901.94		1.20			1	30524.98	1.557						
		0	19048.91					4	30591.45	1.54						
		4	19108.87		1.03			2	32457.44	1.84						
		2	19410.66		0.97			4	32684.61	1.47						
		4	19893.07		1.06			3	33124.48	1.63						
		1	21033.45		0.88			3	34125.40	1.44						
		3	21123.66		0.94			5	34365.33	1.42						
		2	21303.36		1.26			4	34803.88	1.37						
		1	22563.65					2	35090.50	1.57						
		5d <sup>5</sup> 6s(a <sup>1</sup> D) 6p	s <sup>1</sup> D <sup>o</sup>	5	23482.90			847.21	1.55				3	35616.98		1.51
				4	22615.69			-2397.24	1.62				1	35919.55	1.29	
3	25012.93			-262.49	1.73	2	36345.81	1.15								
2	25875.48				2.04	2	36634.54	1.36								
1						3	36806.27	1.48								
2	23317.60				1.27											
4	23322.66				1.04											
3	23984.58				1.07											



## Os I—Continued

## Os I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
		8	36817. 94		1. 28			4	48746. 57		1. 32
		4	36886. 39		1. 41			3	49011. 58		1. 05
		1	36976. 51		1. 31			5	49401. 88		1. 22
		3	37808. 58		1. 26			2	49437. 13		1. 00
		4	37908. 77		1. 45			3	49515. 75		1. 21
		2	37981. 71		1. 45			2	49610. 81		1. 05
		4	38180. 09		1. 24			4	49754. 64		1. 05
		1	38244. 90		1. 55			4	49863. 69		1. 20
		3	38364. 16		1. 23			5	49876. 18		1. 13
		2	38390. 75		1. 41			2	44075. 34		1. 16
		3	38486. 01		1. 61			3	44144. 14		1. 13
		1	38613. 44		0. 29			1	44474. 88		1. 01
		2	38741. 19		1. 60			2	44668. 97		1. 31
		1	38875. 97		1. 90			1	44789. 93		1. 01
		3	39338. 95		1. 18			6	44839. 09		1. 19
		5	39406. 98		1. 18			3	44869. 71		1. 35
		2	39493. 80		0. 93			4	44899. 67		1. 22
		2	39674. 89		1. 91			5	44921. 18		1. 27
		4	40087. 01		1. 36			6	45315. 88		1. 30
		6	40390. 44		1. 30			4	45388. 69		1. 07
		4	40561. 92		1. 32			2	45503. 45		1. 32
		1	40497. 48		2. 18			3	45561. 99		1. 00
		2	40883. 05		0. 82			5	45758. 63		1. 23
		6	41023. 22		1. 23			3	45774. 55		1. 12
		5	41225. 02		1. 27			2	46169. 54		1. 19
		3	41331. 98		1. 08			1	46208. 36		0. 62
		4	41725. 58		1. 04			4	46263. 46		1. 12
		3	41875. 93		1. 30			3	46327. 99		1. 11
		1	42299. 83		1. 00			2	46406. 90		1. 15
		4	42310. 25		1. 22			1	46515. 45		1. 19
		3	42316. 68		1. 03			3	46776. 29		1. 25
		1	42422. 36		1. 86			2	46812. 87		1. 15
		0	42522. 75								
		2	42653. 36		1. 41			1 or 2	46947. 12		

Os I—Continued

Os I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
5d <sup>6</sup> 6s(a <sup>4</sup> D)7s	e <sup>4</sup> D	3	47057. 29		1. 21	5d <sup>6</sup> 6s(a <sup>4</sup> D)7s	e <sup>4</sup> D	4	50274. 91		1. 12
		4	47152. 42		1. 12			4	50223. 52		1. 12
		5	47192. 73		1. 57			2	50311. 71		1. 15
		4	45757. 44	-1522. 71	1. 50			3	50322. 27		1. 22
		3	51122. 12	-2400. 00	1. 71			5	50577. 19		1. 19
		2						3	50453. 20		
		1						4	50503. 53		1. 19
		5	47200. 25		1. 21			3	50530. 59		
		1	47477. 05		1. 14			1	50522. 22		1. 06
		2	47532. 52		0. 99			6	50630. 22		1. 13
		3	47614. 24		1. 12			1	50670. 23		1. 19
		3	47227. 22		1. 11			5	50902. 27		1. 19
		3	47245. 24					4	50937. 24		1. 18
		4	47253. 53		1. 17			2	50922. 77		1. 12
		2	47234. 29		1. 21			4	51022. 49	-1263. 32	1. 59
		3	47257. 27					3	52401. 21	252. 12	
		4	48121. 22					2	52142. 29		1. 25
		3	48202. 52	1. 14				1			
		1	48322. 02					0			
		1	48402. 22	0. 92				3	51042. 72		1. 25
		4	48522. 27	1. 07				0	51122. 22		
		1	48542. 42					1	51217. 74		0. 66
		4	48752. 10	1. 16				3	51222. 22		1. 27
		2	48772. 42	1. 15				2	51211. 24		0. 90
		3	48274. 22	1. 22				4	51222. 22		1. 07
		2	48222. 22					2	51542. 22		1. 50
		1	49052. 22	0. 22				2	51252. 22		1. 09
		2	49112. 10	1. 10				3	51752. 22		1. 13
		3	49122. 11	1. 22				6	52042. 22		1. 17
		3	49202. 02					3	52122. 11		1. 02
3	49421. 12			2	52122. 02		1. 21				
4	49421. 22	1. 20		2	52227. 72						
5	49524. 22			1	52274. 22		1. 14				
2	49214. 24	1. 10		4	52420. 22		1. 11				
3	49220. 42			3	52522. 21						
2	49272. 27	0. 92		3	52204. 12		0. 99				
3	49222. 24			3	52277. 22						
3	49247. 02	1. 19		4	52712. 24		1. 13				
1	50007. 22	1. 42		6	52201. 22		1. 13				
3	50122. 27	1. 16		3	52227. 22		1. 24				

## Os I—Continued

## Os I—Continued

Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Desig.	<i>J</i>	Level	Interval	Obs. <i>g</i>
		2	59075. 24		1. 20			2	56092. 46		1. 29
		4	59131. 25		1. 19			5	56222. 41		
		3	59338. 28		1. 39			4	56729. 16		
		2	59424. 05					1	57122. 30		
		4	59447. 29		1. 13			3	57334. 23		
		1	59454. 11					3	57826. 90		
		6	59578. 09		1. 10			4	58098. 19		
		2	59594. 54					3	58688. 20		
		2	59647. 07					4	58768. 76		
		3	59758. 43					3	58905. 20		
		3	59886. 85					3	58957. 24		
		2	54049. 89		0. 96			3	59074. 97		
		2	54064. 40					2	59105. 95		
		1	54902. 45		1. 57			2	59142. 16		
		6	54928. 11		1. 21			2	59170. 43		
		1	54424. 27		0. 59			2	59229. 84		
		3	54676. 96		1. 82			3	59282. 85		
		2	54753. 20					3	59556. 04		
		1	54800. 48		1. 46			3	59582. 97		
	1 or 2		54847. 38					2	59602. 19		
	2		54867. 69					3	59750. 04		
	3		55388. 87		1. 78			4	60649. 49		
	5		55402. 47		1. 40						
	4		55410. 00			Os II( <sup>6</sup> D <sub>5/2</sub> )	Limit		70450		

October 1956.

## Os I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms		
5d <sup>9</sup> 6s <sup>2</sup>	{	a <sup>4</sup> D	a <sup>3</sup> H
		ns (n ≥ 6)	
5d <sup>9</sup> 6s(a <sup>4</sup> D)nx	{	e <sup>3</sup> D e <sup>3</sup> D	s <sup>3</sup> P <sup>o</sup> s <sup>3</sup> D <sup>o</sup> s <sup>3</sup> F <sup>o</sup>
5d <sup>7</sup> ( <sup>4</sup> F)nx	{	a <sup>4</sup> F a <sup>3</sup> F	
5d <sup>7</sup> ( <sup>4</sup> P)nx	{	a <sup>4</sup> P a <sup>3</sup> P	

\*For predicted terms in the spectra of the Os I isoelectronic sequence, see Vol. III, Introduction.

## Os II

(Re 1 sequence; 75 electrons)

Z=76

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^6 6s^0 D_{5/2}$  $a^0 D_{5/2}$  K

I. P. volts

The analysis is from van den Bosch who has furnished the tabular data in advance of publication, especially for inclusion here. His observations were made at the Massachusetts Institute of Technology, and include Zeeman spectrograms taken with the Bitter magnet. There are about 250 observed lines between 2280 Å and 4550 Å, of which 73 are classified.

No series are known. From a study of screening constants Finkelnburg and Humbach interpolate an ionization potential of  $17 \pm 1$  volts.

Most of the levels have no *LS*-designations because of the departure from *LS*-coupling and the effect of configuration-interaction.

## REFERENCES

W. Finkelnburg und W. Humbach, *Naturwiss.* **42**, 35 (1955). (I P)

J. C. van den Bosch, unpublished material (August 1956). (T) (C L) (Z E)

## Os II

## Os II

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
$5d^6(^0D)6s$	$a^0 D$	$4\frac{1}{2}$	0.00		1.52	$5d^6(^0D)6p$	$s^0 D^0$	$3\frac{1}{2}$	24980.73		1.18
		$3\frac{1}{2}$	3593.15	-3593.15	1.57			$2\frac{1}{2}$	25452.13		1.09
		$2\frac{1}{2}$	3928.94	-335.79	1.62			$2\frac{1}{2}$	37321.53		2.22
		$1\frac{1}{2}$	5592.05	-1663.11	1.79			$2\frac{1}{2}$	39389.49		1.85
		$0\frac{1}{2}$	6636.57	-1044.52	3.17			$4\frac{1}{2}$	44315.40	513.04	1.44
		$2\frac{1}{2}$	7891.93		1.70			$3\frac{1}{2}$	43902.36	-2571.15	1.56
		$3\frac{1}{2}$	11459.90		1.27			$2\frac{1}{2}$	46373.51	216.32	1.61
		$2\frac{1}{2}$	11654.08		1.55			$1\frac{1}{2}$	46157.19	-1970.89	1.74
		$1\frac{1}{2}$	13136.61		1.40			$0\frac{1}{2}$	48128.08		2.96
		$3\frac{1}{2}$	13203.88		1.25			$2\frac{1}{2}$	48798.70		1.39
		$2\frac{1}{2}$	13414.80		1.34			$3\frac{1}{2}$	49149.39		1.51
		$4\frac{1}{2}$	15605.58		1.14			$4\frac{1}{2}$	51951.61		1.40
		$3\frac{1}{2}$	17242.26		0.99			$3\frac{1}{2}$	52206.48		1.42
		$1\frac{1}{2}$	17424.39		0.97			$3\frac{1}{2}$	54379.27		1.34
		$2\frac{1}{2}$	17569.40		1.35			$2\frac{1}{2}$	54445.19		1.45
		$5\frac{1}{2}$	17688.64		1.20			$2\frac{1}{2}$	55538.65		1.09
		$2\frac{1}{2}$	19590.91		0.81			$1\frac{1}{2}$	55635.92		0.70
		$3\frac{1}{2}$	19985.93		1.10			$3\frac{1}{2}$	56791.35		1.28
		$4\frac{1}{2}$	21590.81		1.11			$3\frac{1}{2}$	57402.59		1.29
		$2\frac{1}{2}$	24465.66		1.27						

October 1956.

## IRIDIUM

Ir I

77 electrons

Z=77

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4 4d^{10} 4f^{14} 5s^2 5p^4 5d^7 6s^2$   $^4F_{9/2}$  $\alpha$   $^4F_{9/2}$  75000 K

I. P. 9 volts

The first regularities in Ir I were discovered by Snyder in 1900, who worked out an array for 240 lines, but never published it. Albertson extended this work in 1932, and again in 1938, when he reported that 1937 lines out of some 3100 were classified as combinations among 214 energy levels. He determined the approximate value of the limit by assuming "a simple series relationship" between  $\alpha$   $^4F$  and  $\epsilon$   $^4F$ .

A new description of the spectrum including Zeeman observations has been made recently by van Kleef, who has revised and extended the earlier work. He has also utilized Zeeman spectrograms taken by van den Bosch in 1949 at the Massachusetts Institute of Technology. His present line list extends from 2000.70 Å to 8592.601 Å. Out of a total of some 4000 lines he reports that approximately 2000 have been classified. He has furnished the present data in advance of publication, especially for inclusion here. There are 59 even and 163 odd levels known, but most of them have not been assigned *LS*-designations because of the departure from *LS*-coupling and the effect of configuration-interaction.

## REFERENCES

- W. Albertson, *Phys. Rev.* **43**, 443 (1932). (T) (C L)  
 W. Albertson, *Phys. Rev.* **53**, 940 (A) (1938); **54**, 183 (1938). (I P) (T) (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 70 (1952). (Summary hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 31 (1955). (Summary hfs)  
 Th. A. M. van Kleef, unpublished material (September 1956). (T) (C L) (Z E)

Ir I

Ir I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g	
5d <sup>7</sup> 6s <sup>2</sup>	a <sup>4</sup> F	4½	0.00		1.30			2½	39324.57		1.22	
		3½	6323.91	-6323.91	1.21			2½	39805.97		1.34	
		2½	5784.62	590.29	1.20			5½	39940.87		1.32	
		1½	4078.94	1705.68	1.12			3½	40291.19		1.25	
5d <sup>7</sup> ( <sup>2</sup> F)6s	b <sup>4</sup> F	4½	2834.98	-4271.63	1.33			4½	40389.83		1.23	
		3½	7106.61	-2770.93	1.23			1½	40684.73		1.22	
		2½	9877.54	-1953.55	1.17			3½	40710.78		1.33	
		1½	11831.09		0.54			4½	41118.71		1.37	
5d <sup>7</sup> ( <sup>2</sup> F)6s	a <sup>2</sup> P	1½	10578.68	-1927.00	0.97			0½	41210.33		2.39	
		0½	12505.68		1.22			2½	41522.22		1.32	
5d <sup>7</sup> ( <sup>2</sup> F)6s	a <sup>2</sup> F	2½	12218.47	869.43	1.13			2½	41817.15		1.12	
		3½	13087.90		1.17			3½	43178.15		1.12	
5d <sup>7</sup> ( <sup>2</sup> F)6s	a <sup>2</sup> P	2½	12951.67	-3613.68	1.49			1½	43200.89		0.98	
		1½	16565.35	-115.85	1.39			3½	43592.21		1.45	
		0½	16681.20		2.61			1½	44569.85		1.49	
5d <sup>7</sup> 6s <sup>2</sup>	a <sup>2</sup> G	4½	13939.80	-3839.44	1.10			2½	44596.77		1.26	
		3½	17779.24		0.92			4½	44642.67		1.34	
5d <sup>7</sup> 6s <sup>2</sup>	b <sup>2</sup> P	2½	16103.32	-2443.72	1.34			4½	44652.43		1.25	
		1½	18547.04	-1689.66	1.39			1½	44785.44		1.53	
		0½	20236.70		2.08			3½	45111.68		1.08	
5d <sup>7</sup> ( <sup>1</sup> D)6s	a <sup>2</sup> D	2½	19060.62	-3049.62	1.00			2½	45185.95		1.11	
		1½	22110.24		1.00			1½	45259.14		1.27	
5d <sup>7</sup> 6s <sup>2</sup>	a <sup>2</sup> H	5½	19593.25	-3912.66	1.09			0½	45415.26		1.93	
		4½	23505.91		1.04			0½	45503.15		0.32	
5d <sup>7</sup> 6s <sup>2</sup>	b <sup>2</sup> D	2½	23310.36	-2919.12	1.20			2½	45570.89		1.03	
		1½	26229.48		0.94			3½	45895.85		1.16	
5d <sup>7</sup> 6s( <sup>2</sup> F)6p	s <sup>2</sup> D <sup>o</sup>	4½	26307.50	-4222.16	1.48			5½	45957.33		1.15	
		3½	30529.66	-2535.17	1.44			2½	46093.84		1.21	
		2½	33064.83	601.25	1.50			4½	46220.32		1.23	
		1½	32463.58	-3184.36	1.58			4½	46371.64		1.36	
		0½	35647.94		1.30			1½	46471.84		1.21	
5d <sup>7</sup> ( <sup>1</sup> G)6s	b <sup>2</sup> G	3½	26365.16	1548.68	0.92			1½	46618.13		1.05	
		4½	27913.84		1.03			3½	46879.08		1.18	
5d <sup>7</sup>	c <sup>2</sup> D	2½	26404.17	-1565.88	1.16			2½	47011.09		1.13	
		1½	27970.05		0.85			3½	47165.12		1.23	
5d <sup>7</sup> 6s( <sup>2</sup> F)6p	s <sup>2</sup> F <sup>o</sup>	5½	28452.32	-4061.11	1.40			0½	47203.81		2.31	
		4½	32513.43	-1361.00	1.36							
		3½	33874.43	-1045.40	1.38							
		2½	34919.83	-1470.32	1.34							
		1½	36390.15	-1730.79	1.14							
		0½	38120.94		0.66							
5d <sup>7</sup> 6s( <sup>2</sup> F)6p	s <sup>2</sup> G <sup>o</sup>	6½	32830.78	-1349.70	1.37							
		5½	34180.42	-900.32	1.24							
		4½	35080.80	-329.83	1.27							
		3½	35410.63	-129.71	1.32							
		2½	35540.34	-2152.41	1.59							
		1½	37692.75		0.79							
5d <sup>7</sup> 6s( <sup>2</sup> F)6p	s <sup>2</sup> D <sup>o</sup>	3½	37515.31	69.18	1.38							
		2½	37446.13		1.33							
		1½										
		0½										
5d <sup>7</sup> 6s( <sup>2</sup> F)6p	s <sup>2</sup> F <sup>o</sup>	4½	37871.69	-286.55	1.25							
		3½	38153.24		1.27							
		2½										
		1½										
		4½	38229.75		1.22							
		2½	38352.13		1.39							
		1½	38484.74		1.44							
		3½	38562.05		1.35							
0½	39229.22		1.58									

## Ir I—Continued

## Ir I—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
		4½	47806. 57		1. 08			2½	52327. 33		1. 17
		2½	47537. 29		1. 21			0½	52388. 38		1. 91
		3½	47548. 69		1. 25			2½	52806. 16		1. 16
		1½	47824. 93		1. 46			1½	52806. 57		1. 36
		5½	47858. 47		1. 30			2½	53552. 93		1. 20
		2½	48206. 57		1. 27			4½	53642. 06		1. 13
		4½	48299. 24		1. 20			3½	53686. 99		1. 12
		1½	48440. 83		1. 24			3½	53771. 90		1. 25
		3½	48448. 65		1. 05			5½	54061. 29		1. 14
		3½	48629. 22		1. 19			2½	54119. 23		0. 97
		1½	48801. 91		1. 31			3½	54140. 83		1. 10
		2½	49146. 44		1. 22			4½	54263. 79		1. 16
		4½	49158. 61		1. 26			1½	54319. 92		1. 23
		1½	49342. 51		1. 22			3½	54566. 06		0. 95
		0½	49446. 25		0. 67			2½	54639. 31		0. 98
		2½	49621. 32		1. 36			4½	54667. 54		1. 04
		5½	49719. 17		1. 31			3½	54711. 10		1. 05
		2½	49779. 37		1. 35			4½	54892. 63		1. 36
		3½	49823. 54		1. 22			3½	54894. 82		1. 06
		4½	50050. 69		1. 15			1½	54985. 38		1. 31
		1½	50100. 71		1. 36			4½	55035. 94		1. 10
		2½	50169. 88		1. 00			5½	55114. 11		1. 12
		4½	50434. 46		1. 09			1½	55160. 88		1. 17
		0½	50445. 02		1. 16			2½	55303. 80		1. 50
		1½	50564. 12		1. 19			3½	55382. 00		
		3½	50580. 39		0. 95			1½	55497. 33		1. 16
		5½	50606. 38		1. 04			3½	55619. 39		1. 16
		3½	51107. 94		1. 15			0½	55635. 36		1. 16
		1½	51166. 54		1. 21			1½	55885. 44		
5d <sup>7</sup> 6s(4F) 7s	e 4F	5½	51175. 94	-1332. 95	1. 42			2½	55913. 57		1. 17
		4½	52508. 89		1. 38			4½	55979. 05		1. 16
		3½						1½	56016. 60		1. 05
		2½						0½	56128. 34		0. 65
		1½						2½	56264. 76		1. 22
		0½						3½	56371. 08		0. 96
		2½	51427. 15		1. 12			4½	56416. 08		1. 38
		4½	51470. 74		1. 22			1½	56538. 45		1. 23
		2½	51814. 75		1. 33			1½	56788. 22		1. 21
		3½	51852. 38		1. 24			3½	56792. 60		1. 27
		0½	51983. 92		1. 42			3½	57042. 80		1. 15
		2½	52051. 75		1. 25			3½	57186. 64		0. 99
		3½	52134. 11		1. 32			1½	57247. 57		1. 39
		3½	52224. 37		1. 30						
		5½	52266. 22		0. 98						
		1½	52303. 65		-0. 10						

Ir I—Continued

Ir I—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
		3½	57697. 48		0. 98			1½	60662. 18		0. 94
		1½	57737. 04		1. 40			1½	60663. 13		1. 13
		4½	57774. 99		1. 14			2½	60781. 97		1. 00
		2½	58048. 08		1. 00			1½	60794. 40		0. 94
		2½	58117. 84		1. 36			3½	61074. 71		1. 13
		3½	58470. 79					1½	61431. 49		
		4½	58570. 87		0. 92			1½	61431. 37		0. 45
		3½	58625. 10		1. 03			3½	62018. 84		1. 35
		1½	59155. 48		0. 88			4½	62079. 88		
		2½	59354. 02		1. 01			3½	62164. 91		1. 35
		5½	59525. 87		1. 28			5½	62208. 67		1. 20
		1½	59564. 11		1. 19			4½	62220. 91		1. 39
		0½	59618. 93		2. 01			3½	62377. 72		1. 23
		3½	59647. 72		1. 32			0½	62425. 80		
		4½	59785. 87					3½	62757. 18		1. 11
		2½	59824. 53					4½	62835. 40		1. 28
		3½	59878. 02		1. 30			1½	63391. 27		
		4½	60012. 25		1. 38			4½	63469. 72		1. 27
		2½	60062. 45		1. 35			3½	63976. 18		1. 41
		3½	60963. 04					3½	64618. 16		
		2½	60411. 48								
		3½	60643. 36								
						Ir I (P)	Limit		75000		

October 1956.

Ir I OBSERVED TERMS\*

Configuration 1 <sup>st</sup> 2 <sup>nd</sup> 3 <sup>rd</sup> 4 <sup>th</sup> 3 <sup>rd</sup> 3 <sup>rd</sup> 4 <sup>th</sup> 4 <sup>th</sup> 4 <sup>th</sup> 4 <sup>th</sup> 5 <sup>th</sup> 5 <sup>th</sup> +	Observed Terms	
5d <sup>7</sup> 6s <sup>2</sup>	{ b <sup>4</sup> P	a <sup>4</sup> F
5d <sup>8</sup>	{ b <sup>3</sup> D	a <sup>3</sup> G a <sup>3</sup> H
	{ c <sup>3</sup> D	
	ns (n ≥ 6)	np (n ≥ 6)
5d <sup>6</sup> ( <sup>4</sup> F)ns	{ b <sup>4</sup> F	
	{ a <sup>4</sup> F	
5d <sup>6</sup> ( <sup>2</sup> F)ns	{ a <sup>4</sup> P	
	{ a <sup>2</sup> P	
5d <sup>6</sup> ( <sup>3</sup> D)ns	{ a <sup>3</sup> D	
5d <sup>6</sup> ( <sup>1</sup> G)ns	{ b <sup>3</sup> G	
5d <sup>7</sup> 6s( <sup>4</sup> F)ns	{ c <sup>4</sup> F	s <sup>4</sup> D° s <sup>4</sup> F° s <sup>4</sup> G°
		s <sup>4</sup> D° s <sup>4</sup> F°

\*For predicted terms in the spectra of the Ir I isoelectronic sequence, see Vol. III, Introduction.



## PLATINUM

Pt I

78 electrons

Z=78

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^9 6s^1 D_3$  $6s^1 D_3$ , 72300 K

I. P. 9.0 volts

"A fairly complete wave-number system for Pt was sent to Rowland by Snyder in October, 1900", according to Meggers and Laporte, who interpreted the early regularities among the low levels when they observed Pt I in absorption in 1926. Haussmann studied the Zeeman effect of 173 lines and reported 72 levels in 1927. Livingood revised and extended Haussmann's work in 1929, and his paper is quoted for most of the level values in the table. Except for terms arising from the  $5d^9 ns$  and  $5d^{10}$  configurations, the assignment of term designations on the basis of *LS*-coupling is extremely tentative, and the spectrum needs further study.

No homogeneous line list exists, although the observations extend from 1928.85 Å to 10757.78 Å. More than 440 lines have been classified. Kessler and Meggers have recently observed the Pt I spectrum from 6648.32 Å to 10757.78 Å. In an attempt to classify these lines in the long-wave region the writer prepared a complete line list and combination array; added 5 new "odd" levels; and rejected 2, namely, 46007.9 and 46793.9. A number of strong lines in the newly observed region still remain unclassified, however. Further observations are needed to extend the analysis. Livingood's notation is in the first column of the table. The new levels can be detected from the absence of an entry in this column.

The observed *g*-values are from Livingood. These are also inadequate for further study of the spectrum.

Goble pointed out in 1935 that in the Pt I sequence *jj*-coupling applies more nearly than *LS*-coupling, to the levels having the configuration  $5d^9(^2D)6p$ . In *LS*-coupling this configuration gives rise to the terms  $^1S(P^0D^0F^0)$ . In *jj*-coupling the same number of energy levels and the same resultant *J*-values occur, but each level is defined by two *j*-values, one for the *d*-electron ( $1\frac{1}{2}$  or  $2\frac{1}{2}$ ), and one for the *p*-electron ( $0\frac{1}{2}$  or  $1\frac{1}{2}$ ). By analogy with Au II and Hg III, Shenstone has suggested the *j*-values given in the table for 8 levels of this group. Of the remaining "odd" levels, three with *J*=2 and one with *J*=1 belong to the  $5d^9 6p$  configuration. The rest are doubtless from  $5d^9 6s 6p$ ; but configuration-interaction and the lack of adequate observational material make it impossible to assign with assurance *LS*-designations with limit terms. Consequently, the levels are listed in numerical order in the table. A table giving the transformation from *jj*- to *LS*-coupling may be found in the book by Condon and Shortley, p. 294.

The limit is from the  $6, 7s^1 D_3$  series, which represents the removal of an *s*-electron from the  $d^9 s$  configuration. It has been derived by a Rydberg formula, but corrected to allow for the percentage error known to apply in the first long period, in cases where the more accurate Ritz formula could be compared with the Rydberg formula for series of three members.

## REFERENCES

- W. F. Meggers and O. Laporte, *Phys. Rev.* **28**, 642 (1926). (T) (C L)  
 A. C. Haussmann, *Astroph. J.* **66**, 333 (1927). (T) (C L) (Z E)  
 J. J. Livingood, *Phys. Rev.* **34**, 185 (1929). (I P) (T) (C L) (Z E)  
 A. T. Goble, *Phys. Rev.* **48**, 346 (1935).  
 E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra*, p. 294 (Cambridge University Press, London, 1951).  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 K. G. Kessler, W. F. Meggers, and C. E. Moore, *J. Research Nat. Bur. Std.* **53**, 225, RP 2538 (1954). (C L)  
 A. G. Shenstone, unpublished material (September 1954).  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab. ORNL-1469, Suppl. II, 31 (1955). (Summary hfs)

Pt I

Pt I

L	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	L	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
e <sup>1</sup> D <sub>2</sub>	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6s	6s <sup>1</sup> D	3	0.0	-775.9	1.01	28i	5d <sup>2</sup> 6s 6p		4	48965.1		1.34
e <sup>1</sup> D <sub>2</sub>			2	775.9			29i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		1	47740.6		1.43
e <sup>1</sup> F <sub>4</sub>	5d <sup>2</sup> 6s <sup>2</sup>		4	823.7		1.25	30i	5d <sup>2</sup> 6s 6p		4	48361.9		1.25
e <sup>1</sup> G <sub>4</sub>	5d <sup>2</sup>	5d <sup>2</sup> <sup>1</sup> S	0	6140.0			31i			2	48535.6		1.02
e <sup>1</sup> F <sub>2</sub>	5d <sup>2</sup> 6s <sup>2</sup>		2	6567.5		1.12	32i	5d <sup>2</sup> 6s 6p		3	48779.3		1.22
e <sup>1</sup> F <sub>1</sub>	5d <sup>2</sup> 6s <sup>2</sup>		3	10116.8			33i	5d <sup>2</sup> 6s 6p		3	49886.1		1.19
e <sup>1</sup> D <sub>1</sub>	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6s	6s <sup>1</sup> D	1	10132.0			34i			1	49544.5		1.24
e <sup>1</sup> D <sub>2</sub>	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6s	6s <sup>1</sup> D	2	13496.3		1.17	35i			2	49880.8		1.12
e <sup>1</sup> F <sub>3</sub>	5d <sup>2</sup> 6s <sup>2</sup>		2	15501.8		0.92	36i			1	50055.3		0.87
e <sup>1</sup> P <sub>1</sub>	5d <sup>2</sup> 6s <sup>2</sup>		1	18566.5			50i	5d <sup>2</sup> 6s 6p		3	51097.5		1.21
e <sup>1</sup> G <sub>2</sub>	5d <sup>2</sup> 6s <sup>2</sup>		4	21967.1		1+?	37i			2	51286.9		1.13
b <sup>1</sup> D <sub>2</sub>	5d <sup>2</sup> 6s <sup>2</sup>		2	26638.6		0.97	38i			2	51545.5		1.25
s <sup>1</sup> D <sub>1</sub>	5d <sup>2</sup> 6s( <sup>1</sup> F)6p		4	30157.0		1.46	39i			2	51752.3		1.34
1i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>3/2</sub>		2	38620.0		1.39	40i			1	52071.6		1.22
2i	5d <sup>2</sup> 6s 6p		5	33880.5		1.32	e <sup>1</sup> D <sub>2</sub>	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s	7s <sup>1</sup> D	3	52379.3	-287.9	1.32
3i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>3/2</sub>		3	34122.1		1.21	e <sup>1</sup> D <sub>2</sub>	"	2	52667.2	1.04		
4i	5d <sup>2</sup> 6s 6p		3	35321.7		1.33	41i			2	52708.3		1.46
	5d <sup>2</sup> 6s 6p		4	36296.4			42i			1	53019.2		1.08
s <sup>1</sup> G <sub>1</sub>	5d <sup>2</sup> 6s( <sup>1</sup> F)6p		6	36781.6		1.33	43i			2	53953.3		1.32
5i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>3/2</sub>		1	36844.7		1.09		5d <sup>2</sup> 6s 6p		3	54011.1		
6i			2	37342.1		1.15	44i	5d <sup>2</sup> 6s 6p		3	54839.2		1.21
7i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		4	37590.7		1.25	45i,3			2?	55216.8		0.96
8i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		3	37769.0		1.17		5d <sup>2</sup> 6s 6p		3	55536.0		
s <sup>1</sup> F <sub>1</sub>	5d <sup>2</sup> 6s( <sup>1</sup> F)6p		5	38536.2		1.30	e <sup>1</sup> F <sub>3</sub>	5d <sup>2</sup> 6s( <sup>1</sup> F)7s		5	55640.7		1.41
10i			2	38815.9		0.88	D <sub>4</sub>			4	56784.4		1.27
11i	5d <sup>2</sup> 6s 6p		4	40194.2		1.21				3	57506.4?		
12i			2	40516.3		1.38	Q <sub>2</sub>			3	59731.5		1.3
13i			2	40787.9		1.20	E <sub>1</sub>			3	59751.2		
14i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		0	40873.5			F <sub>1</sub>			3	59764.3		1.27
15i	5d <sup>2</sup> 6s 6p		3	40970.1		1.12	G <sub>1</sub>			1	59782.8		1.07
16i			1	41202.7		0.92	H <sub>1</sub>			3	59872.1		1.23
17i	5d <sup>2</sup> 6s 6p		3	42660.2		1.19	I <sub>4</sub>			4	59882.4		1.17
18i			1	43187.8		1.39	J <sub>1</sub>			2	59908.1		1.02
19i			1	43945.7		1.21	e <sup>1</sup> D <sub>1</sub>	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s	7s <sup>1</sup> D	1	60357.8		0.52
20i	5d <sup>2</sup> 6s 6p		4	44432.7		1.20	e <sup>1</sup> D <sub>2</sub>	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s	7s <sup>1</sup> D	2	60640.6		1.08
21i			2	44444.4		1.21	M <sub>1</sub>			3	60790.4		1.07
22i	5d <sup>2</sup> 6s 6p		3	44730.3		1.19	N <sub>4</sub>			4	60834.0		1.29
23i			1	45398.4		1.52	R <sub>1,3</sub>				64129.1		
24i			2	46170.4		1.01	T <sub>1</sub>			6	64141.3		
25i			2	46419.4		0.87	O <sub>4</sub>			4	64505.9		
26i	5d <sup>2</sup> 6s 6p		0	46433.9									
27i	5d <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		3	46622.5		1.15							
								Pt II( <sup>2</sup> D <sub>3/2</sub> )	Limit		72300		

February 1955.

## Pt II

(Ir I sequence; 77 electrons)

Z=78

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^9 {}^2D_{3/2}$  $5d^9 {}^2D_{3/2}$  149723

I. P. 18.56 volts

The analysis is from Shenstone, who has observed the spectrum from 976.419 Å to 4514.17 Å and classified more than 550 lines. His spectrograms were made with three sources, namely, the arc and spark in air and the Schüller tube. He comments that "High even levels are calculated from arc or Schüller tube wave numbers which are approximately 2.5 greater than the spark wave numbers" given in his published table of classified lines.

Shenstone's notation is given in the left-hand column of the table, and his suggested term designations have been adopted. The limit terms in the "Config." column and the primes used in the "Desig." column to denote the different limits have been introduced by the writer. A single prime indicates that the Pt II term arises from the second highest limit term in Pt III; a double prime, the third limit term, etc. Since the Pt III spectrum has not yet been analyzed, the limit terms are assumed to lie in the same relative positions as indicated by the positions of the Pt II terms.

The doublet and quartet systems of terms are connected by observed intersystem combinations. Many levels have been assigned numbers only, "because the coupling is more nearly  $jj$  than  $LS$ ."

The ionization limit has been derived from the  $ns {}^4F_{3/2}$  series ( $n=6$  to 8) by means of a Ritz formula.

## REFERENCE

A. G. Shenstone, Phil. Trans. Roy. Soc. (London) [A] 237, No. 782, 453 (1938). (I P) (T) (C L)

Pt II

Pt II

Shenstone	Config.	Desig.	J	Level	Interval	Shenstone	Config.	Desig.	J	Level	Interval
5d 'D <sub>104</sub>	5P	5P 'D	2 1/2	0.0	-8419.9	44b <sub>4</sub>			4 1/2	78481.8	
5d 'D <sub>104</sub>			1 1/2	8419.9		45b <sub>4</sub>			3 1/2	78768.8	
6a 'F <sub>104</sub>	5P'(F)6a	6a 'F	4 1/2	4798.6	-4569.6	92b <sub>4</sub>			1 1/2	74000.1	
6a 'F <sub>104</sub>			3 1/2	9884.2		92b <sub>4</sub>			1 1/2	74841.8	
6a 'F <sub>104</sub>			2 1/2	13329.8		46b <sub>4</sub>			1 1/2	74819.6	
6a 'F <sub>104</sub>			1 1/2	15791.4		46b <sub>4</sub>			1 1/2	74746.8	
1 <sub>104</sub>	5P'(P)6a	6a' 'P	2 1/2	16821.0	-4348.0	47b <sub>4</sub>			2 1/2	74755.8	
2 <sub>104</sub>			1 1/2	21188.0		48b <sub>4</sub>			3 1/2	75185.8	
3 <sub>104</sub>			0 1/2	21717.5		48b <sub>4</sub>			0 1/2	75682.8	
6a 'F <sub>104</sub>	5P'(F)6a	6a 'F	3 1/2	18097.9	-5363.8	88a			3 1/2	76110.7	
6a 'F <sub>104</sub>			2 1/2	28461.7		49b <sub>4</sub>			1 1/2	76682.8	
4 <sub>104</sub>	5P'(D)6a	6a'' 'D	1 1/2	28878.9	9042.8	50b <sub>4</sub>			2 1/2	76821.1	
7 <sub>104</sub>			2 1/2	32919.2		51b <sub>4</sub>			1 1/2	76810.7	
6a' 'F <sub>104</sub>	5P' 6a'	6a' 'F	4 1/2	24879.6	-9767.8	52b <sub>4</sub>			4 1/2	77519.9	
6a' 'F <sub>104</sub>			3 1/2	24647.4		53b <sub>4</sub>			2 1/2	78044.0	
6a' 'F <sub>104</sub>			2 1/2	36484.2		54b <sub>4</sub>			4 1/2	78908.8	
6a' 'F <sub>104</sub>			1 1/2	37878.2		55b <sub>4</sub>			2 1/2	79608.1	
5 <sub>104</sub>	5P'(P)6a	6a' 'P	0 1/2	37256.1	4981.5	56b <sub>4</sub>			3 1/2	81897.9	
6 <sub>104</sub>			1 1/2	32237.6		57a			0 1/2	82636.5	
6a 'G <sub>104</sub>	5P'(G)6a	6a''' 'G	3 1/2	29030.7	231.6	80b <sub>4</sub>			1 1/2	82824.5	
6a 'G <sub>104</sub>			4 1/2	29262.8		58b <sub>4</sub>			2 1/2	82973.8	
6p 'D <sub>104</sub>	5P'(F)6p	6p 'D°	3 1/2	51408.8	-9782.0	91b <sub>4</sub>			2 1/2	80858.7	
24b <sub>4</sub>			2 1/2	61190.8		56b <sub>4</sub>			3 1/2	81897.9	
21b <sub>4</sub>			1 1/2	56588.8		57a			0 1/2	82636.5	
26b <sub>4</sub>			0 1/2	68788.8		58b <sub>4</sub>			2 1/2	82973.8	
6p 'G <sub>104</sub>	5P'(F)6p	6p 'G°	5 1/2	61058.6	7183.1	59b <sub>4</sub>			4 1/2	84183.8	
20b <sub>4</sub>			4 1/2	58875.5		60b <sub>4</sub>			3 1/2	85701.1	
22b <sub>4</sub>			3 1/2	57018.8		61b <sub>4</sub>			3 1/2	85701.1	
23b <sub>4</sub>	5P'(F)6p	6p 'F°	4 1/2	60907.8	-757.8	62b <sub>4</sub>			2 1/2	86608.8	
25b <sub>4</sub>			3 1/2	61665.6		63b <sub>4</sub>			4 1/2	86863.9	
27b <sub>4</sub>			4 1/2	62890.6				2 1/2	88482.0		
28b <sub>4</sub>			3 1/2	65739.8				3 1/2	88336.1		
29b <sub>4</sub>			1 1/2	64389.0				3 1/2	88482.0		
30b <sub>4</sub>			2 1/2	64757.6				4 1/2	89482.9		
31b <sub>4</sub>			2 1/2	65351.4				2 1/2	89754.9		
32b <sub>4</sub>			0 1/2	65587.4				3 1/2	9082.9		
33b <sub>4</sub>			1 1/2	66088.8				4 1/2	95803.5		
34b <sub>4</sub>			3 1/2	66484.6				2 1/2	96614.6		
35b <sub>4</sub>			2 1/2	68658.4				1 1/2			
36b <sub>4</sub>			0 1/2	69896.1				3 1/2	97630.9		
37b <sub>4</sub>			2 1/2	69953.7				2 1/2	98187.5		
38b <sub>4</sub>			4 1/2	70181.8				3 1/2	98817.9		
39b <sub>4</sub>			2 1/2	70879.8				2 1/2	99909.6		
40b <sub>4</sub>			2 1/2	71315.0				2 1/2	99797.7		
41b <sub>4</sub>			2 1/2	71948.9				2 1/2	100239.7		
42b <sub>4</sub>			2 1/2	72991.57				1 1/2	100612.0		
43b <sub>4</sub>			1 1/2	73086.7							
						7a 'F <sub>104</sub>	5P'(F)7a	7a 'F	4 1/2	95803.5	-811.1
						7a 'F <sub>104</sub>			3 1/2	96614.6	

Pt II—Continued

Pt II—Continued

Shenstone	Config.	Desig.	J	Level	Interval	Shenstone	Config.	Desig.	J	Level	Interval	
75hc	5d <sup>9</sup> 7c		1½	100798.9		6d <sup>4</sup> H <sub>3c</sub>	5d <sup>9</sup> (F)6d	6d <sup>4</sup> H	6½	104782.9	126.8	
76hc		3½	100803.8		9 <sub>3c</sub>				5½	104837.1		
8ac		2½	101199.3						4½			
77hc		3½	101342.2			11 <sub>3c</sub>	5d <sup>9</sup> (F)6d	6d <sup>4</sup> G	5½	105066.4		
78hc		2½	101398.7			12 <sub>3c</sub>			4½	105357.5		
79hc		3½	101515.9†						3½			
80hc		3½	101549.4				5d <sup>9</sup> (F)6d	6d <sup>4</sup> F	4½	106297.1	-126.3	
81hc		5½	101612.8			14 <sub>3c</sub>			3½	106433.4		
82hc		2½	101867.7						2½			
83hc		2½	101917.3				8c <sup>4</sup> F <sub>3c</sub>	5d <sup>9</sup> (F)8c	8c <sup>4</sup> F	4½	120626.6†	
84hc		4½	102086.6						3½			
85hc		2½	102415.9						2½			
86hc		2½	103493.9						1½			
87hc		3½	103517.4				15 <sub>3c</sub>	5d <sup>9</sup> 7d		4½	124047.0	
							16 <sub>3c</sub>	5d <sup>9</sup> 7d		4½	124487.4	
								Pt III(F <sub>1</sub> )	Limit	---	149723	

August 1954.

Pt II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup>	Observed Terms								
5d <sup>9</sup>	5d <sup>9</sup> <sup>3</sup> D								
5d <sup>7</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>4</sup> F								
	ns (n ≥ 6)			np (n ≥ 6)		nd (n ≥ 6)			
5d <sup>9</sup> (F)ns	{	6-8c <sup>4</sup> F 6c <sup>3</sup> F		6p <sup>4</sup> D <sup>o</sup>	6p <sup>4</sup> F <sup>o</sup>	6p <sup>4</sup> G <sup>o</sup>	6d <sup>4</sup> F	6d <sup>4</sup> G	6d <sup>4</sup> H
5d <sup>9</sup> (F)ns'		6s' <sup>4</sup> P 6s' <sup>3</sup> P							
5d <sup>9</sup> (D)ns''		6s'' <sup>3</sup> D							
5d <sup>9</sup> (G)ns'''				6s''' <sup>3</sup> G					

\*For predicted terms in the spectra of the Ir I isoelectronic sequence, see Vol. III, Introduction.

## GOLD

## Au I

79 electrons

Z=79

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 S_{1/2}$  $6s^2 S_{1/2}$  74416.0 K

I. P. 9.22 volts

The analysis is from Platt and Sawyer, who have revised and extended the earlier work by Thorsen, McLennan and McLay, and others. They list 315 classified lines in the interval 1494.57 Å and 10109.0 Å. Their limit has been derived by fitting an extended Ritz formula to the  $ns^2 S$  series ( $n=7$  to 10; 9 to 12), and the  $nd^2 D$  series ( $n=10$  to 13), the average deviation being  $\pm 3$  K.

They have assigned arbitrary numbers to the miscellaneous levels; these numbers have been retained in the left column of the table.

The three-place observed  $g$ -values in the last column are from a manuscript by Green and Maxwell, furnished in advance of publication. The writer has determined the 5 two-place  $g$ -values: 4 from the observations of Symons and Daley, and 1 from data in the 1955 manuscript.

Platt and Sawyer state that the levels from the  $5d^9 6s 6p$  configuration are divided into two widely separated groups "based, respectively, on  $5d^9 {}^2D_{3/2}$  and  $5d^9 {}^2D_{1/2}$  of Au III." This results from the fact that the Au I spectrum exhibits approximate  $jj$ -coupling, as is the case with Au II. The levels of Au II having as limits  ${}^2D_{3/2}$  and  ${}^2D_{1/2}$  in Au III are commonly designated  ${}^2D_{3,1}$  and  ${}^2D_{1,1}$  when the limit terms  ${}^2D$  in  $LS$ -notation have their levels grouped in pairs. In  $jj$ -coupling the pairs are described by the notation  $d_{3/2}^9 s_{1/2} | 3/2$  and  $d_{1/2}^9 s_{1/2} | 1/2$ , respectively. The complete configuration assignments have the form  $d_{3/2}^9$  and  $d_{1/2}^9$  combined with the  $j$ -values of the running  $p$ - and  $s$ -electrons, those for the running  $p$ -electron being  $1\frac{1}{2}$  and  $0\frac{1}{2}$ , and for an  $s$ -electron  $0\frac{1}{2}$ . The order of coupling of the vectors is not yet known for Au I.

In the table the subscripts for the electrons have been introduced by Trees and the writer to describe more fully the configurations of the observed levels in  $jj$ -coupling. Although the criteria for these assignments are not definitive, yet a good guide is available in Au I from a study of the observed  $g$ -values. For this purpose, the theoretical  $g$ -values for  $jj$ -coupling have been calculated from the formula by Mack, on the assumption that the  $d$ - and  $s$ -electrons together are coupled with the  $p$ -electron. These are given in the following table, which is complete for the  $5d^9 6s 6p$  array of levels. In the left column the  $jj$ -coupling notation is given for each of the four groups of levels. Beneath it is an arbitrary notation indicating the  $J$ -value of the limiting  $LS$ -level. This is included to facilitate comparison of the observed  $g$ -values with the theoretical Landé  $g$ -values in  $LS$ -coupling for the terms involved, which, in this case, are  ${}^4P^{\circ}D^{\circ}F^{\circ}$  and  ${}^2P^{\circ}D^{\circ}F^{\circ}$ . Observed  $g$ -values are entered where known. Leaders indicates that the level is known but no observed  $g$ -value is available. Three levels marked "absent" have not yet been found.

## Au I—Continued

g-values for j-coupling						
Configuration	J	4%	8%	2%	1%	0%
5d <sub>5/2</sub> 6s <sub>1/2</sub> 6p <sub>3/2</sub>	Theory Observed	1.333 -----	1.333 1.873	1.333 -----	1.333 1.422	
5d <sup>9</sup> 6s(1D <sub>2</sub> )6p <sub>3/2</sub>						
5d <sub>5/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>	Theory Observed		1.288 1.258	1.429 1.532		
5d <sup>9</sup> 6s(1D <sub>2</sub> )6p <sub>1/2</sub>						
5d <sub>5/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>	Theory Observed		1.200 Absent	1.000 -----	1.111 -----	1.889 Absent
5d <sup>9</sup> 6s(1D <sub>2</sub> )6p <sub>1/2</sub>						
5d <sub>5/2</sub> 6s <sub>1/2</sub> 6p <sub>3/2</sub>	Theory Observed			1.013 0.984	0.556 -----	0.444 -----
5d <sup>9</sup> 6s(1D <sub>2</sub> )6p <sub>3/2</sub>					1.187 1.16	

For the running 6d-electron no attempt has been made to indicate the *j*-values, as the data are too incomplete. The limit term, together with a suggested *J*-value denoting the limiting level involved, is, however, tentatively assigned.

Some *LS*-designations for miscellaneous levels, given in the literature, are quoted here, and others have been added provisionally by Trees and the writer, but no attempt has been made to group the levels into terms because of the departure from *LS*-coupling. For 5d<sup>9</sup> 6s 6p levels Racah has made a theoretical study of the arrangement of the terms in *LS*-coupling for Cu I, which, combined with the observed *g*-values has served as a general guide in making the present very tentative *LS*-term-assignments in Au I.

## REFERENCES

- J. E. Mack, *Phys. Rev.* **34**, 30 (1929).  
 A. S. M. Symons and J. Daley, *Proc. Phys. Soc. London* **41**, 431 (1929). (T) (C L) (Z E)  
 J. R. Platt and R. A. Sawyer, *Phys. Rev.* **60**, 866 (1941). (I P) (T) (C L)  
 G. Racah, *Phys. Rev.* **63**, 525 (1942).  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge National Laboratory ORNL-1469, Suppl. II, 31 (1955). (Summary hfs)  
 J. B. Green and H. N. Maxwell, *J. Opt. Soc. Am.* **45**, 98 (1955). (Z E)

As I

As I

PS	Config.	Design.	J	Level	Interval	Obs. <i>g</i>	PS	Config.	Design.	J	Level	Interval	Obs. <i>g</i>
	5d <sup>20</sup> (S)6e	6e *S	0%	0.0		1.997		5d <sup>20</sup> (S)9e	9e *S	0%	68880.5		
	5d <sup>20</sup> 6e <sup>2</sup>	6e <sup>2</sup> *D	2%	9161.2	-12274.0	1.193	2	5d <sup>20</sup> 6e <sup>2</sup> (D <sub>2</sub> )7e	*D	2%	68706.1		1.270
			1%	21435.2		0.804							
	5d <sup>20</sup> (S)6p	6p *P°	0%	37352.9	2815.4	0.661		5d <sup>20</sup> (S)8d	8d *D	1%	69971.2	24.2	
			1%	41174.2		1.234				2%	70007.6		
1°	5d <sup>20</sup> <sub>1</sub> 6e <sup>20</sup> <sub>1</sub> 6p <sup>20</sup> <sub>1</sub>	*P°	2%	48162.5		1.533	3	5d <sup>20</sup> (S)10e	10e *S	0%	70617.2		
2°	"	*P°	2%	45537.4		1.258	4	5d <sup>20</sup> 6e <sup>2</sup> (D <sub>2</sub> )7e	*D	1%	70652.2		0.905
3°	"	*P°	2%	46174.5		1.011		5d <sup>20</sup> 6e <sup>2</sup> (D <sub>2</sub> )7e	*D	2%	71222.0		
4°	5d <sup>20</sup> <sub>1</sub> 6e <sup>20</sup> <sub>1</sub> 6p <sup>20</sup> <sub>1</sub>	*D°	2%	46379.0				5d <sup>20</sup> (S)9d	9d *D	1%	71254.4	24.0	
5°	"	*P°	1%	47007.4		1.422				2%	71330.4		
6°	"	*P°	4%	48697.4				5d <sup>20</sup> (S)11e	11e *S	0%	71712.2		
7°	"	*D°	3%	51029.1		1.372		5d <sup>20</sup> (S)10d	10d *D	1%	72162.2	7.0	
8°	5d <sup>20</sup> <sub>1</sub> 6e <sup>20</sup> <sub>1</sub> 6p <sup>20</sup> <sub>1</sub>	*D°	1%	51231.4		1.064				2%	72170.2		
9°	5d <sup>20</sup> <sub>1</sub> 6e <sup>20</sup> <sub>1</sub> 6p <sup>20</sup> <sub>1</sub>	*P°	1%	51435.0				5d <sup>20</sup> (S)12e	12e *S	0%	72395.0		
10°	"	*D°	2%	51652.9		1.222		5d <sup>20</sup> (S)11d	11d *D	1%	72690.1	4.7	
11°	"	*P°	3%	52202.1						2%	72694.2		
12°	"	*P°	0%	53197.5				5d <sup>20</sup> (S)13e	13e *S	0%	72847.5		
	5d <sup>20</sup> (S)7e	7e *S	0%	54484.9		1.995		5d <sup>20</sup> (S)12d	12d *D	1%	73051.0	2.2	
13°	5d <sup>20</sup> <sub>1</sub> 6e <sup>20</sup> <sub>1</sub> 6p <sup>20</sup> <sub>1</sub>	*D°	0%	55732.5						2%	73054.2		
14°	"	*D°	1%	56106.0		1.16		5d <sup>20</sup> (S)14e	14e *S	0%	73162.1		
15°	"	*P°	2%	56916.5		0.964		5d <sup>20</sup> (S)13d	13d *D	1%	73208.1	4.5	
16°	"	*P°	1%	58245.1						2%	73212.0		
17°	5d <sup>20</sup> <sub>1</sub> 6e <sup>20</sup> <sub>1</sub> 6p <sup>20</sup> <sub>1</sub>	*P°	2%	59712.2			5	5d <sup>20</sup> (S)14d	14d *D	1%	73500.4	2.1	
18°	"	*D°	2%	61255.1		1.20				2%	73502.5		
19°	"	*D°	1%	61662.2				-----	-----				
20°	"	*P°	1%	63005.1				Au II(S <sub>2</sub> )	Limit		74419.0		
	5d <sup>20</sup> (S)7p	7p *P°	0%	60032.0			6	5d <sup>20</sup> 6e <sup>2</sup> (D <sub>2</sub> )6d		1%	76278.0		
			1%	60722.2	695.0	1.227	7	"		2%	76566.7		
	5d <sup>20</sup> (S)6d	6d *D	1%	61951.6	32.1	0.787	8	"		5%	76569.2		
			2%	62032.7		1.193	9	"		3%	76721.2		
	5d <sup>20</sup> (S)8e	8e *S	0%	64742.4		2.01	10	"		4%	76829.9		
	5d <sup>20</sup> (S)9p	9p *P°	0%	66905.2	205.0		11	"		1%	76906.4		
			1%	66910.2			12	"		2%	76955.0		
	5d <sup>20</sup> (S)7d	7d *D	1%	67469.4	41.2		13	5d <sup>20</sup> 6e <sup>2</sup> (D <sub>2</sub> )6d		2%	78667.2		
			2%	67510.7			14	"		1%	78819.9		
	5d <sup>20</sup> (S)5f	5f *P°	3%	67485.2	-4.7		15	"		1%	79010.2		1.10
			2%	67490.0			16	5d <sup>20</sup> 6e <sup>2</sup> (D <sub>2</sub> )6d		2%	80772.7		
1	5d <sup>20</sup> 6e <sup>2</sup> (D <sub>2</sub> )7e	*D	3%	67811.5		1.422	17	"		1%	81682.7		

February 1965.



## Au II

(Pt I sequence; 78 electrons)

Z=79

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} {}^1S_0$  $5d^{10} {}^1S_0$ , 165000 K

I. P. 20.5 volts

The analysis is from Platt and Sawyer, who have revised and extended the earlier work by McLennan and McLay, Rao, Mack and Fromer, and others. There are approximately 200 classified lines in the interval 820.42 Å to 8999.7 Å, but the work is still far from complete.

The spectra of the Pt I isoelectronic sequence are of special theoretical interest because they approach *jj*-coupling rather than *LS*- or *JL*-coupling. This appears more conspicuously in the spectra of higher ionization especially among the levels from the  $5d^9({}^2D)6p$  configuration. A detailed study of "The Four Vector Problem and Its Application to Energies and Intensities in Platinum-like Spectra" has been made by Goble. In *LS*-coupling this configuration gives rise to the terms  ${}^1,3(P^{\circ}D^{\circ}F^{\circ})$ . In *jj*-coupling the same number of energy levels and the same resultant *J*-values occur, but each level is defined by two *j*-values, one for the *d*-electron ( $1\frac{1}{2}$  or  $2\frac{1}{2}$ ), and one for the *p*-electron ( $0\frac{1}{2}$  or  $1\frac{1}{2}$ ).

In the table the writer has assigned *LS*-designations only for the terms from the  $5d^9({}^2D)ns$  configuration, although the *jj*-coupling applies here, also, as can be seen from the pairs of  $6s$ - and  $7s$ -levels. The levels from  $5d^9({}^2D)np$  have the *j*-values of the *d*- and *p*-electrons, as required for *jj*-coupling, indicated in the configuration column, the former being the same as the *J*-values of the limit term  ${}^2D$ . Shenstone has suggested that by analogy with Hg III the configuration assignments by Platt and Sawyer be interchanged for the levels labeled  $5^{\circ}$  and  $8^{\circ}$ . This has been done in the table. A table giving the transformation from *jj*- to *LS*-coupling may be found in the book by Condon and Shortley, p. 294.

The observed *g*-values are from a manuscript on "The Zeeman Effect of Gold", furnished by Green and Maxwell in advance of publication. They supersede the early work by Symons and Daley.

Platt and Sawyer state that their limit is estimated, with the help of Rydberg term tables, from the known terms of the  $5d^9 ns$  and  $5d^9 np$  series.

## REFERENCES

- A. S. M. Symons and J. Daley, Proc. Phys. Soc. London 41, 431 (1929). (Z E)  
 A. T. Goble, Phys. Rev. 48, 346 (1935).  
 J. R. Platt and R. A. Sawyer, Phys. Rev. 60, 866 (1941). (I P) (T) (C L)  
 E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra* p. 294 (Cambridge University Press, London, 1951).  
 A. G. Shenstone, Letter (October 1954).  
 J. B. Green and H. N. Maxwell, J. Opt. Soc. Am. 45, 96 (1955). (Z E)

Au II

Au II

PS	Config.	Desig.	J	Level	Interval	Obs. g	PS	Config.	Desig.	J	Level	Interval	Obs. g
1	5d°	5d° 18	0	0.0			12°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		2	86565.1		1.29
2	5d°(°D <sub>7</sub> )6e	6e °D	3	15089.0			14	5d° 6d°		0	91114.4		
3			2	17639.4	-2000.4		15	5d°(°D <sub>7</sub> )7e	7e °D	3	106172.2		1.20
4	5d°(°D <sub>13</sub> )6e		1	27764.5	-10125.1		16			2	106630.8	-458.6	0.95
5	5d°(°D <sub>13</sub> )6e	6e °D	2	29620.8		1.04	17	5d°(°D <sub>13</sub> )6d		1	116049.8		
6	5d° 6d°		4	40478.3		1.13	18	5d°(°D <sub>13</sub> )6d		4	116945.8		
7	5d° 6d°		2	42510.4		1.02	19	5d°(°D <sub>13</sub> )6d		2	117065.2		
8	5d° 6d°		3	52176.0		1.15	20	5d°(°D <sub>13</sub> )6d		1	117296.9		
9	5d° 6d°		0	55436.6			21	5d°(°D <sub>13</sub> )6d		3	117511.4		
10	5d° 6d°		2	58191.2		1.06	22	5d°(°D <sub>13</sub> )6d		3	117982.8		
11	5d° 6d°		1	59101.0			23	5d°(°D <sub>13</sub> )6d		2	118028.9		
12	5d° 6d°		4	61384.5			24	5d°(°D <sub>13</sub> )6d		4	118167.6		
1°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		2	63052.5		1.45	13°	5d°(°D <sub>13</sub> )7p <sub>13</sub>		2	119446.1		
2°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		3	65002.9		1.04	14°	5d°(°D <sub>13</sub> )7p <sub>13</sub>		3	120256.5		
13	5d° 6d°		2	68145.1		1.26	25	5d°(°D <sub>13</sub> )7e	7e °D	1	120822.1		
3°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		4	72494.8		1.15	15°	5d° 6e 6p		2	120952.0		
4°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		2	73177.8		1.12	26	5d°(°D <sub>13</sub> )7e	7e °D	2	121117.8		
5°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		1	73403.1		1.04	16°	5d°(°D <sub>13</sub> )7p <sub>13</sub>		2	121734.8		
6°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		3	74790.9		1.30	17°	5d°(°D <sub>13</sub> )7p <sub>13</sub>		1	121862.0		
7°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		2	76652.7		1.38	18°	5d°(°D <sub>13</sub> )7p <sub>13</sub>		4	123061.8		
8°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		1	81659.1			19°	5d°(°D <sub>13</sub> )7p <sub>13</sub>		3	123343.8		1.03
9°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		0	82613.0									
10°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		3	85699.8									
11°	5d°(°D <sub>13</sub> )6p <sub>13</sub>		1	85707.0		1.11		Au III(°D <sub>13</sub> )	Limit	...	165000		

February 1955.

## MERCURY

## Hg I

80 electrons

Z=80

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^4 4d^{10} 4f^{14} 5s^2 5p^4 5d^{10} 6s^2 \text{ } ^1S_0$  $6s^2 \text{ } ^1S_0$ , 84184.1 K

I. P. 10.43 volts

The first spectrum of mercury has attracted the attention of many investigators. Its lines exhibit isotope shifts, Paschen-Back Effect, hyperfine structure, Stark Effect, and other phenomena.

With the development of the electrodeless Hg<sup>180</sup> lamp, the spectrum has come into special prominence. Meggers has pointed out that with this source the observed wavelengths provide ideal and reproducible standards. Barrell has shown that these standards can be reproduced with an accuracy of  $\pm 1$  part in  $10^5$ . The line at 5460 Å is being considered by the International Committee on Weights and Measures as the primary standard of length.

More than 600 lines have been classified in Hg I. The observations extend from 745.95 Å to 39970 Å. Observed intersystem combinations connect the terms of different multiplicities. A number of "forbidden" transitions have been observed. Mrozowski discusses the structure of two such lines,  $\lambda 2967$  and  $\lambda 2269$ , as due to the "nuclear perturbational effect" in the odd isotopes. This has been discussed further by Kesler.

Shenstone and Russell comment that "The mercury spectrum contains long series but none of them is capable of giving an accurate limit" because of perturbations. The writer has obtained the value of the limit quoted in the table by adding the correction suggested by Walerstein, +2.8, to Paschen's value 84181.3. This limit gives fairly consistent values for converting the different lists of absolute term values later than Paschen's, into one complete array of terms starting from the ground state zero. The tabular values of the energy levels are not derived from a homogeneous line list, and the wavelengths are in general none too accurate, even though selected ones observed with the interferometer are measured with precision. Consequently, the narrow intervals given for some terms in the table are within the errors of observation and the terms may well be unresolved.

The three-place entries in the table are from Burns, Adams, and Longwell; and from Fowles, who has corrected the older values of  $8p \text{ } ^1P_1^o$  and confirmed the suggestion by Shenstone and Russell regarding  $6p' \text{ } ^1P^o$ . The two-place entries are quoted from Humphreys, who has observed Hg I in the infrared, and also from Fowles. Walerstein has extended the series of "even" terms, and his values have been used for higher series members. For the "odd" series having the  $^3S$  limit, most of the higher members are taken from Murakawa, but the terms  $np \text{ } ^3P^o$  ( $n=15$  to  $26$ ) and  $np \text{ } ^1P^o$  ( $n=17$  to  $19$ ) are from Kamiyama. Sawyer and Beese (1926) reported the  $6p^2 \text{ } ^3P_{1,2}$  levels. The level  $6p^2 \text{ } ^3P_0$  is from Garton and Rajaratnam, who designate the line reported by Selwyn at 1972.94 Å as  $6p \text{ } ^3P_1^o - 6p^2 \text{ } ^3P_0$ , in place of Sawyer's line at 1900.1 Å. The long series having as limit the  $^3D$  term in Hg II are from the lines observed in absorption by Beutler, in the extreme ultraviolet. These lines represent transitions between the ground term  $6s^2 \text{ } ^1S_0$  and levels above the ionization limit. Only the transition  $J=0$  to  $J=1$  is to be expected, so the missing components of these high triplet terms have not been indicated in the table. There are also three components of triplet terms from this limit below the ionization limit, which are listed separately, namely,  $6p' \text{ } ^3F_4^o$ ,  $6p' \text{ } ^3D_3^o$ , and  $6p' \text{ } ^3P_2^o$ . The  $J$ -values of the limit term and the values of the high  $^3D$  limits are quoted from Beutler. Shenstone also discusses auto-ionization in his paper entitled "Ultra-Ionization Potentials in Mercury Vapor."

The observed  $g$ -values are from Green and Loring. Both Paschen-Back Effect and hyperfine structures have been extensively observed for selected lines, but few  $g$ -values are reported. Only a small selection of the many references to this type of work is given below.

## Hg I—Continued

## REFERENCES

- A. Landé, *Phys. Zeit.* **23**, 417 (1921). (Z E)  
 A. Fowler, *Report on Series in Line Spectra*, p. 146 (Fleetway Press, Ltd., London, 1922). (I P) (T) (C L)  
 F. Paschen und R. Götz, *Seriengesetze der Linienspektren* p. 116 (Julius Springer, Berlin 1922). (I P) (T) (C L)  
 R. A. Sawyer, *J. Opt. Soc. Am. and Rev. Sci. Instr.* **13**, 440 (1926). (T) (C L)  
 H. Nagaoka and T. Mishima, *Proc. Imp. Acad. Japan* **2**, 383 (1926). (Z E)  
 W. C. van Geel, *Zeit. Phys.* **47**, 615 (1928). (Z E)  
 E. D. McAllister, *Phys. Rev.* **34**, 1142 (1929); *Phys. Rev.* **35**, 1585 (L) (1930). (T) (C L)  
 F. Paschen, *Ann. der Phys.* [5] **6**, 47 (1930). (T) (C L)  
 T. Takamini and T. Suga, *Sci. Papers Inst. Phys. Chem. Research (Tokyo)* **13**, 2, No. 232 (1930). (T) (C L)  
 A. G. Shenstone, *Phys. Rev.* **38**, 873 (1931). (I P)  
 A. G. Shenstone and H. N. Russell, *Phys. Rev.* **39**, 415 (1932). (C L)  
 H. Beutler, *Zeit. Phys.* **84**, 289 (1933); **86**, 710 (1933). (T) (C L)  
 J. B. Green and R. A. Loring, *Phys. Rev.* **46**, 888 (1934). (Z E)  
 I. Walerstein, *Phys. Rev.* **46**, 874 (1934). (T) (C L)  
 K. Murakawa, *Proc. Phys. Math. Soc. Japan* [3] **17**, 14 (1935). (T) (C L) (hfs)  
 K. Murakawa, *Proc. Phys. Math. Soc. Japan* [3] **18**, 345 (1936); *Zeit. Phys.* **106**, 168 (1937). (T) (C L)  
 T. Suga, M. Kamiyama and T. Sugiura, *Sci. Papers Inst. Phys. Chem. Research (Tokyo)* **34**, No. 741, 32 (1937). (C L) (G D)  
 E. H. Collins, *Phys. Rev.* **56**, 48 (1939). (Z E)  
 M. Kamiyama, *Bull. Inst. Phys. Chem. Research (Tokyo)* **23**, No. 10, 877 (1943). (T) (C L)  
 S. Mrosowski, *Phys. Rev.* **67**, 161 (1945). (C L)  
 K. Burns, K. B. Adams, and J. Longwell, *J. Opt. Soc. Am.* **40**, 339 (1950). (T) (C L)  
 W. F. Meggers and F. O. Westfall, *J. Research Nat. Bur. Std.* **44**, 447, RP 2091 (1950).  
 K. G. Kessler, *Phys. Rev.* **77**, 559 (L) (1950). (C L)  
 H. Barrell, *Proc. Roy. Soc. London [A]* **209**, No. 1096, 132 (1951).  
 K. Burns and K. B. Adams, *J. Opt. Soc. Am.* **43**, 56 (1952); **43**, 716 (1952). (T) (C L) (I S)  
 P. F. A. Klippenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 C. J. Humphreys, *J. Opt. Soc. Am.* **43**, 1027 (1953). (T) (C L)  
 G. R. Fowles, *J. Opt. Soc. Am.* **44**, 85, 760 (1954). (T) (C L) (I S)  
 W. R. S. Garton and A. Rajaratnam, *Proc. Phys. Soc. London [A]* **68**, 1107 (1955). (T) (C L)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, Suppl. II, 31 (1955). (Summary hfs)

Hg I

Hg I

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
$5d^{10} 6s^2$	$6s^2 \text{ } ^1S$	0	0.000			$5d^{10} 6s(2S)8p$	$8p \text{ } ^3P^o$	0	76447.243		
$5d^{10} 6s(2S)6p$	$6p \text{ } ^3P^o$	0	37645.080	1767.220	1.479		1	76487.067		19.824	
		1	39412.300	4630.677	1.503		2	76823.5		356.4	
		2	44042.977			$5d^{10} 6s(2S)8p$	$8p \text{ } ^1P^o$	1	76863.864		
$5d^{10} 6s(2S)6p$	$6p \text{ } ^1P^o$	1	54088.781		1.019	$5d^{10} 6s(2D_{3/2})6p$	$6p' \text{ } ^3F^o$	4	76945.41		
$5d^{10} 6s(2S)7s$	$7s \text{ } ^2S$	1	62350.456		2.003	$5d^{10} 6s(2S)7d$	$7d \text{ } ^1D$	2	77064.097		
$5d^{10} 6s(2S)7s$	$7s \text{ } ^1S$	0	63928.243			$5d^{10} 6s(2S)7d$	$7d \text{ } ^3D$	1	77084.632		
$5d^{10} 6s(2D_{3/2})6p$	$6p' \text{ } ^3P^o$	2	68836.60				2	77107.917			
							3	77129.535		23.285	
$5d^{10} 6s(2S)7p$	$7p \text{ } ^3P^o$	0	69516.66	145.23		$5d^{10} 6s(2S)5f$	$5f \text{ } ^3F^o$	2	77237.04		
		1	69661.89	1545.62			3	77239.20			
		2	71207.51				4	77286.89		2.16	
$5d^{10} 6s(2S)7p$	$7p \text{ } ^1P^o$	1	71296.15			$5d^{10} 6s(2S)5f$	$5f \text{ } ^1F^o$	3	77241.68		47.79
$5d^{10} 6s(2S)6d$	$6d \text{ } ^1D$	2	71333.182			$5d^{10} 6s(2S)9s$	$9s \text{ } ^2S$	1	78216.261		
$5d^{10} 6s(2S)6d$	$6d \text{ } ^3D$	1	71336.164	60.056		$5d^{10} 6s(2S)9s$	$9s \text{ } ^1S$	0	78404.387		
		2	71396.220	35.091	1.109	$5d^{10} 6s(2D_{3/2})6p$	$6p' \text{ } ^1D^o$	2	78676.7		
		3	71431.311		1.342	$5d^{10} 6s(2D_{3/2})6p$	$6p' \text{ } ^3P^o$	1	78813		
$5d^{10} 6s(2D_{3/2})6p$	$6p' \text{ } ^1D^o$	3	73119.2			$5d^{10} 6s(2S)9p$	$9p \text{ } ^3P^o$	0	79375.783		
$5d^{10} 6s(2S)8s$	$8s \text{ } ^2S$	1	73961.298				1	79412.745		36.962	
$5d^{10} 6s(2S)8s$	$8s \text{ } ^1S$	0	74404.590				2	79613.3		200.6	

Hg I—Continued

Hg I—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
5d <sup>10</sup> 6s(°S)8d	8d 1D	2	79660. 785			5d <sup>10</sup> 6s(°S)13s	13s °S	1	82591. 3		
5d <sup>10</sup> 6s(°S)8d	8d 1D	1	79678. 708			5d <sup>10</sup> 6s(°S)13s	13s °S	0	82616. 2		
		2	79690. 300	11. 592		5d <sup>10</sup> 6s(°S)13p	13p 1P°	0			
		3	79702. 634	12. 334				1	82765. 9	29. 1	
5d <sup>10</sup> 6s(°S)6f	6f 1F°	2	79743. 7					2	82785. 03		
		3	79745. 0	1. 3		5d <sup>10</sup> 6s(°S)12d	12d 1D	2	82807. 421		
		4	79748. 3	2. 3		5d <sup>10</sup> 6s(°S)12d	12d 1D	1	82811. 2	1. 3	
5d <sup>10</sup> 6s(°S)6f	6f 1F°	3	79745. 3					2	82812. 5	2. 8	
5d <sup>10</sup> 6s(°S)9p	9p 1P°	1	79964. 1					3	82815. 3		
5d <sup>10</sup> 6s(°S)10s	10s °S	1	80268. 056			5d <sup>10</sup> 6s(°S)10f	10f 1F°	2	82818. 7	1. 1	
5d <sup>10</sup> 6s(°S)10s	10s °S	0	80365. 653					3	82819. 8	0. 3	
5d <sup>10</sup> 6s(°S)10p	10p 1P°	0	80902. 87					4	82820. 1		
		1	80916. 686	14. 42		5d <sup>10</sup> 6s(°S)10f	10f 1F°	3	82820. 4		
		2	81022. 9	106. 2		5d <sup>10</sup> 6s(°S)13p	13p 1P°	1	82823. 94		
5d <sup>10</sup> 6s(°S)9d	9d 1D	2	81057. 749			5d <sup>10</sup> 6s(°S)14s	14s °S	1	82915. 6		
5d <sup>10</sup> 6s(°S)9d	9d 1D	1	81071. 027			5d <sup>10</sup> 6s(°S)14s	14s °S	0	82933. 2		
		2	81077. 8	6. 8		5d <sup>10</sup> 6s(°S)14p	14p 1P°	0			
		3	81085. 126	7. 8				1	83039. 3	21. 9	
5d <sup>10</sup> 6s(°S)7f	7f 1F°	2	81103. 9					2	83061. 2		
		3	81104. 6	0. 7		5d <sup>10</sup> 6s(°S)13d	13d 1D	2	83069. 2		
		4	81107. 1	2. 5		5d <sup>10</sup> 6s(°S)13d	13d 1D	1	83073. 4	0. 8	
5d <sup>10</sup> 6s(°S)7f	7f 1F°	3	81106. 5					2	83074. 2	2. 1	
5d <sup>10</sup> 6s(°S)10p	10p 1P°	1	81153. 614					3	83076. 3		
5d <sup>10</sup> 6s(°S)11s	11s °S	1	81416. 352			5d <sup>10</sup> 6s(°S)11f	11f 1F°	2	83079. 0	1. 1	
5d <sup>10</sup> 6s(°S)11s	11s °S	0	81473. 4					3	83080. 1	-0. 1	
5d <sup>10</sup> 6s(°S)11p	11p 1P°	0	81800. 0					4	83080. 0		
		1	81811. 876	11. 9		5d <sup>10</sup> 6s(°S)11f	11f 1F°	3	83080. 4		
		2	81873. 835	61. 959		5d <sup>10</sup> 6s(°S)14p	14p 1P°	1	83084. 0		
5d <sup>10</sup> 6s(°S)10d	10d 1D	2	81895. 0			5d <sup>10</sup> 6s(°S)15s	15s °S	1	83149. 9		
5d <sup>10</sup> 6s(°S)10d	10d 1D	1	81904. 5			5d <sup>10</sup> 6s(°S)15s	15s °S	0	83162. 9		
		2	81908. 7	4. 2		5d <sup>10</sup> 6s(°S)15p	15p 1P°	0			
		3	81913. 632	4. 9				1	83240. 1	18. 6	
5d <sup>10</sup> 6s(°S)8f	8f 1F°	2	81923. 5					2	83258. 7		
		3	81924. 3	0. 8		5d <sup>10</sup> 6s(°S)14d	14d 1D	2	83263. 9		
		4	81927. 8	3. 5		5d <sup>10</sup> 6s(°S)14d	14d 1D	1	83266. 9	1. 0	
5d <sup>10</sup> 6s(°S)8f	8f 1F°	3	81925. 3					2	83267. 9	1. 7	
5d <sup>10</sup> 6s(°S)11p	11p 1P°	1	81942. 444					3	83269. 6		
5d <sup>10</sup> 6s(°S)12s	12s °S	1	82124. 081			5d <sup>10</sup> 6s(°S)12f	12f 1F°	2	83271. 0	0. 9	
5d <sup>10</sup> 6s(°S)12s	12s °S	0	82160. 8					3	83271. 9	0. 6	
5d <sup>10</sup> 6s(°S)12p	12p 1P°	0						4	83272. 5		
		1	82379. 0			5d <sup>10</sup> 6s(°S)12f	12f 1F°	3	83273. 4		
		2	82422. 60	43. 6		5d <sup>10</sup> 6s(°S)15p	15p 1P°	1	83280. 5		
5d <sup>10</sup> 6s(°S)11d	11d 1D	2	82436. 2			5d <sup>10</sup> 6s(°S)16s	16s °S	1	83325. 0		
5d <sup>10</sup> 6s(°S)11d	11d 1D	1	82443. 0			5d <sup>10</sup> 6s(°S)16s	16s °S	0	83334. 8		
		2	82445. 9	2. 9		5d <sup>10</sup> 6s(°S)16p	16p 1P°	0			
		3	82449. 2	3. 3				1	83396. 7	10. 5	
5d <sup>10</sup> 6s(°S)9f	9f 1F°	2	82451. 9					2	83407. 2		
		3	82455. 2	0. 3		5d <sup>10</sup> 6s(°S)15d	15d 1D	2	83411. 9		
		4	82458. 7	1. 5		5d <sup>10</sup> 6s(°S)15d	15d 1D	1	83413. 9	0. 9	
5d <sup>10</sup> 6s(°S)9f	9f 1F°	3	82455. 3					2	83414. 8	0. 9	
5d <sup>10</sup> 6s(°S)12p	12p 1P°	1	82464. 05					3	83415. 7		

## Hg I—Continued

## Hg I—Continued

Config.	Desig.	J	Level	Interval	Obs. <i>g</i>	Config.	Desig.	J	Level	Interval	Obs. <i>g</i>
5d <sup>10</sup> 6s(°S)13f	13f °F°	2	83416. 1	1. 9		5d <sup>10</sup> 6s(°S)21s	21s °S	0	83772. 7		
		3	83418. 0					0			
		4	83420. 9								
5d <sup>10</sup> 6s(°S)13f	13f °F°	3	83418. 7			5d <sup>10</sup> 6s(°S)21p	21p °P°	0			
								1			
								2	83792. 8		
5d <sup>10</sup> 6s(°S)16p	16p °P°	1	83420. 3			5d <sup>10</sup> 6s(°S)20d	20d °D	2	83798. 3		
5d <sup>10</sup> 6s(°S)17s	17s °S	1	83459. 0			5d <sup>10</sup> 6s(°S)20d	20d °D	3	83802. 4	-0. 2	
5d <sup>10</sup> 6s(°S)17s	17s °S	0	83466. 4					2	83802. 6	-0. 3	
5d <sup>10</sup> 6s(°S)17p	17p °P°	0		8. 9		5d <sup>10</sup> 6s(°S)22s	22s °S	1	83817. 6		
		1	83512. 7					0			
		2	83521. 6								
5d <sup>10</sup> 6s(°S)16d	16d °D	2	83526. 7			5d <sup>10</sup> 6s(°S)22p	22p °P°	0			
								1			
								2	83534. 5		
5d <sup>10</sup> 6s(°S)16d	16d °D	1	83528. 5	0. 4		5d <sup>10</sup> 6s(°S)21d	21d °D	2	83842. 7		
								3	83843. 6	-0. 7	
								2	83844. 3	0. 3	
								1	83844. 0		
5d <sup>10</sup> 6s(°S)17p	17p °P°	1	83530. 7			5d <sup>10</sup> 6s(°S)23s	23s °S	1	83856. 6		
5d <sup>10</sup> 6s(°S)14f	14f °F°	2		7. 7		5d <sup>10</sup> 6s(°S)23p	23p °P°	0			
		3	83530. 9			0					
		4	83538. 6								
5d <sup>10</sup> 6s(°S)18s	18s °S	1	83564. 0					1			
								2	83868. 0		
5d <sup>10</sup> 6s(°S)18s	18s °S	0	83569. 6			5d <sup>10</sup> 6s(°S)22d	22d °D	3	83879. 0	-0. 3	
								2	83879. 3	0. 2	
								1	83879. 1		
5d <sup>10</sup> 6s(°S)18p	18p °P°	0		12. 0		5d <sup>10</sup> 6s(°S)24s	24s °S	1	83889. 7		
		1	83601. 9								
		2	83613. 9								
5d <sup>10</sup> 6s(°S)15f	15f °F°	2		6. 1		5d <sup>10</sup> 6s(°S)24p	24p °P°	0			
		3	83603. 9								
		4	83609. 3								
5d <sup>10</sup> 6s(°S)17d	17d °D	2	83617. 4			5d <sup>10</sup> 6s(°S)23d	23d °D	3	83909. 2	-0. 6	
								2	83909. 8		
								1			
5d <sup>10</sup> 6s(°S)17d	17d °D	1	83618. 5	1. 4		5d <sup>10</sup> 6s(°S)25s	25s °S	1	83918. 0		
				0. 4							
5d <sup>10</sup> 6s(°S)18p	18p °P°	1	83619. 9			5d <sup>10</sup> 6s(°S)25p	25p °P°	0			
								1			
								2	83920. 8		
5d <sup>10</sup> 6s(°S)19s	19s °S	1	83647. 9								
5d <sup>10</sup> 6s(°S)19s	19s °S	0	83651. 2			5d <sup>10</sup> 6s(°S)24d	24d °D	3	83933. 9	-1. 1	
								2	83935. 0		
								1			
5d <sup>10</sup> 6s(°S)19p	19p °P°	0		8. 1		5d <sup>10</sup> 6s(°S)26p	26p °P°	0			
		1	83677. 9								
		2	83686. 0								
5d <sup>10</sup> 6s(°S)18d	18d °D	2	83690. 7					1			
								2	83942. 0		
5d <sup>10</sup> 6s(°S)19p	19p °P°	1	83691. 3			5d <sup>10</sup> 6s(°S)25d	25d °D	3	83955. 7	-0. 5	
								2	83956. 2		
								1			
5d <sup>10</sup> 6s(°S)18d	18d °D	1	83692. 7	0. 1		5d <sup>10</sup> 6s(°S)26d	26d °D	3	83975. 6	-0. 4	
				0. 8				2	83976. 0		
								1			
5d <sup>10</sup> 6s(°S)20s	20s °S	1	83715. 4			5d <sup>10</sup> 6s(°S)27d	27d °D	3	83993. 4		
5d <sup>10</sup> 6s(°S)20s	20s °S	0	83718. 8					2			
								1			
5d <sup>10</sup> 6s(°S)20p	20p °P°	0									
		1									
		2	83745. 7								
5d <sup>10</sup> 6s(°S)19d	19d °D	2	83750. 3			5d <sup>10</sup> 6s(°S)28d	28d °D	3	84007. 4		
								2			
								1			
5d <sup>10</sup> 6s(°S)19d	19d °D	3	83753. 1	0. 5							
				-0. 9							
5d <sup>10</sup> 6s(°S)21s	21s °S	1	83771. 3			5d <sup>10</sup> 6s(°S)29d	29d °D	3	84021. 8		
								2			
								1			

Hg I—Continued

Hg I—Continued

Config.	Design.	J	Level	Interval	Obs. g	Config.	Design.	J	Level	Interval	Obs. g
-----						-----					
Hg II(8 <sub>1/2</sub> )	Limit	---	84184.1			Hg II(8 <sub>1/2</sub> )	Limit	---	119692		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6p	6p' <sup>1</sup> P°	1	88760	3884		5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 7p	7p' <sup>1</sup> P°	1	119653		
5d <sup>9</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> P	0	90096			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 7p	7p' <sup>3</sup> D°	1	120715		
		1	93980			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 8p	8p' <sup>1</sup> P°	1	122973		
		2				5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 8p	8p' <sup>3</sup> D°	1	127254		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6p	6p' <sup>3</sup> D°	1	93737			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 5f	5f' <sup>1</sup> P°?	1	127770		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 7p	7p' <sup>1</sup> P°	1	105881			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 9p	9p' <sup>1</sup> P°	1	129921		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 8p	8p' <sup>1</sup> P°	1	112272			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 9p	9p' <sup>3</sup> D°	1	130053		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 5f?	5f' <sup>3</sup> P°?	1	112726			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6f	6f' <sup>1</sup> P°?	1	130231		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 9p	9p' <sup>1</sup> P°	1	115035			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 10p	10p' <sup>1</sup> P°	1	131444		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 6f	6f' <sup>3</sup> P°?	1	115242			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 10p	10p' <sup>3</sup> D°	1	131535		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 10p	10p' <sup>1</sup> P°	1	116428			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 7f	7f' <sup>1</sup> P°?	1	131644		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 7f	7f' <sup>3</sup> P°?	1	116616			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 11p	11p' <sup>1</sup> P°	1	132345		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 11p	11p' <sup>1</sup> P°	1	117355			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 8f	8f' <sup>1</sup> P°?	1	132602		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 8f	8f' <sup>3</sup> P°?	1	117422			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 12p	12p' <sup>1</sup> P°	1	132921		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 12p	12p' <sup>1</sup> P°	1	117915			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 13p	13p' <sup>1</sup> P°	1	133312		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 9f	9f' <sup>3</sup> P°?	1	117964			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 14p	14p' <sup>1</sup> P°	1	133588		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 13p	13p' <sup>1</sup> P°	1	118307			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 15p	15p' <sup>1</sup> P°	1	133792		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 14p	14p' <sup>1</sup> P°	1	118565			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 16p	16p' <sup>1</sup> P°	1	133940		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 15p	15p' <sup>1</sup> P°	1	118722			5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 17p	17p' <sup>1</sup> P°	1	134057		
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 16p	16p' <sup>1</sup> P°	1	118912			-----					
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 17p	17p' <sup>1</sup> P°	1	119022		Hg II(8 <sub>1/2</sub> )	Limit	---	134732			
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D <sub>3/2</sub> ) 18p	18p' <sup>1</sup> P°	1	119119		-----						

February 1955.

Hg I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms	
	5d <sup>9</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>1</sup> S
5d <sup>9</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> P	
	n <sub>s</sub> (n ≥ 7)	n <sub>p</sub> (n ≥ 6)
5d <sup>9</sup> 6s( <sup>2</sup> S)n <sub>s</sub>	{ 7 to 25s <sup>2</sup> S 7 to 21s <sup>1</sup> S	6 to 26p <sup>1</sup> P° 6 to 19p <sup>1</sup> P°
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D)n <sub>s</sub> '	{	6 to 17p' <sup>1</sup> P° 6 to 10p' <sup>3</sup> D° 6p' <sup>1</sup> F° 6 to 18p' <sup>1</sup> P° 6p' <sup>3</sup> D°
	nd (n ≥ 6)	n <sub>f</sub> (n ≥ 5)
5d <sup>9</sup> 6s( <sup>2</sup> S)n <sub>s</sub>	{ 6 to 29d <sup>3</sup> D 6 to 21d <sup>1</sup> D	5 to 15f <sup>1</sup> F° 5 to 13f <sup>1</sup> F°
5d <sup>9</sup> 6s <sup>2</sup> ( <sup>2</sup> D)n <sub>s</sub> '	{	5 to 9f' <sup>1</sup> P° 5 to 8f' <sup>1</sup> P°

\*For predicted terms in the spectra of the Hg I isoelectronic sequence, see Vol. III, Introduction.

## Hg II

(Au I sequence; 79 electrons)

Z=80

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 S_{0,1}$  $6s^2 S_{0,1}$  151290 K

I. P. 18.751 volts

Although the doublet series having as a limit the term  $^1S_0$  in Hg III are well established in Hg II, the analysis needs to be revised and extended for terms from higher limits. In the table, nearly all the series terms are from Paschen, who revised and extended the earlier work by Carroll and others. In addition, he found the  $6s^2 ^3D$  term and the levels labeled Z, Y, U, X, C, B, and A in the left column of the table, headed "Author". McLennan, McLay, and Crawford extended the analysis in 1931 and added the levels that are designated by numerals in this column. Naudé (1929) suggested a slight improvement in Paschen's  $9 ^3D$  term value, and added the series terms labeled  $10 ^3D_{3,2}$ ,  $12 ^3D_3$ ,  $13 ^3D_3$ , and  $9 ^3F_3$  in Paschen's notation. He also published a number of quartet terms, most of which are included among the miscellaneous numbered levels. Three levels 76924, 77041, 91064, and one above the ionization limit have been omitted here.

As suggested by Edlén (letter, 1958), it has been assumed that the  $g$ - and  $h$ -orbits do not penetrate, and that the lowest  $^2G$  term is  $5g ^2G$ .

There are approximately 300 classified lines in the interval 893.10 Å to 10590 Å, including the infrared extension by Rasmussen, who added the  $5g ^2G$  term.

Paschen calls attention to the strong line observed in emission at 2814.93 Å, whose designation is  $6s^2 S_{0,1} - 6s^2 ^3D_{3,2}$ . This line is "forbidden" in two senses, in that it is produced by the transition between two "even" terms and also violates the  $J$ -rule.

As in Au I, the levels from the  $5d^9 6s 6p$  configuration in Hg II exhibit characteristics of  $jj$ -coupling. They have as limits the pairs  $^3D_{3,2}$  and  $^3D_1 ^3D_2$  in Hg III, which, in  $jj$ -coupling would be designated as  $d_{3/2}^9 s_{1/2} |_{32}$  and  $d_{3/2}^9 s_{1/2} |_{21}$ , respectively. The complete  $jj$ -coupling notation for the 23 levels to be expected from this configuration should indicate, also, the  $j$ -values of the running  $p$ -electron ( $0\frac{1}{2}$  or  $1\frac{1}{2}$ ). Further details regarding this notation may be found in the text for Au I. By comparison with isoelectronic spectra, Trees and the writer have extended the published configuration assignments in  $jj$ -coupling, very tentatively, to include 19 levels in this group. A number of "odd" levels still remain unassigned pending further study of the spectrum.

## REFERENCES

- F. Paschen, Akad. Wiss., Berlin (Phys.-Math. Kl.) Sitz. Parts 31-33, p. 536 (1928). (I P) (T) (C L) (E D)
- E. Rasmussen, Naturwiss. 13, 389 (1929). (T) (C L)
- S. M. Naudé, Ann. der Phys. [5] 3, 1 (1929). (T) (C L) (G D)
- J. C. McLennan, A. B. McLay and M. F. Crawford, Proc. Roy. Soc. London [A] 134, 41 (1931). (T) (C L)
- B. Venkatesachar and T. S. Subbaraya, Zeit. Phys. 73, 412 (1931). (T) (C L)
- T. S. Subbaraya, Zeit. Phys. 78, 541 (1932). (T) (C L)
- R. Ricard, Compt. Rend. 194, 781 (1932). (T) (C L)
- R. F. Baehar and S. Goudsmit, Atomic Energy States, p. 235 (McGraw-Hill Book Co., Inc., New York and London, 1932).
- S. Mrozowski, Phys. Rev. 61, 605 (1942). (T) (I S) (hfs)
- H. E. Walchli, A Table of Nuclear Moment Data, Oak Ridge National Laboratory ORNL-1469, 60 (1953). (Summary hfs).

## Hg II

## Hg II

Author	Config.	Desig.	J	Level	Interval	Author	Config.	Desig.	J	Level	Interval
1S	$5d^{10}(^1S)6s$	$6s^2 S$	0	0		$1h_{3/2}$	$5d_{3/2}^9 6s_{1/2} 6p_{3/2}$		$2\frac{1}{2}$	79704	
$^3D_3$	$5s^2 6s^2$	$6s^2 ^3D$	$2\frac{1}{2}$	35514	-15038	$2h_{3/2}$	"		$3\frac{1}{2}$	84909	
$^3D_2$			$1\frac{1}{2}$	50552		$3h_{3/2}$	"		$2\frac{1}{2}$	84934	
$2P_1$	$5d^{10}(^1S)6p$	$6p^2 ^1P^o$	$0\frac{1}{2}$	51485	9123	$4h_{3/2}$	"		$1\frac{1}{2}$	86177	
$2P_2$			$1\frac{1}{2}$	60608							



Hg II—Continued

Hg II—Continued

Author	Config.	Design.	J	Level	Interval	Author	Config.	Design.	J	Level	Interval
5a <sub>1</sub>	5d <sup>10</sup> (1S)6p <sub>1/2</sub>		4½	92569?		5D <sub>2</sub>	5d <sup>10</sup> (1S)8d	8d 1D	1½	124562	126
6a <sub>1</sub>	"		1½	94001		5D <sub>3</sub>			2½	124698	
7a <sub>1</sub>	"		3½	95186		3a <sub>1</sub>			3½	125299	
8a <sub>1</sub>	"		2½	95308		5F <sub>1</sub>	5d <sup>10</sup> (1S)9p	9p 1P°	0½	125467	362
2S	5d <sup>10</sup> (1S)7a	7a 1S	0½	95714		5F <sub>2</sub>			1½	125829	
						A			2½	126266	
						4a <sub>1</sub>			2½	126712	
9b <sub>1</sub>	5d <sup>10</sup> (1S)6p <sub>1/2</sub>		0½	96125		5S	5d <sup>10</sup> (1S)10a	10a 1S	0½	128424	
10b <sub>1</sub>	"		2½	97094		6F <sub>1</sub>	5d <sup>10</sup> (1S)7f	7f 1F°	3½	129793	-19
11°	5d <sup>10</sup> (1S)6p <sub>1/2</sub> ?		1½?	100859		6F <sub>2</sub>			2½	129812	
12b <sub>1</sub>	"		1½	103183		6G	5d <sup>10</sup> (1S)6g	6g 1G	3½, 4½	129042. 4	
Z	"	1P°	0½	103870		6D <sub>2</sub>	5d <sup>10</sup> (1S)9d	9d 1D	1½	129627	68
						6D <sub>3</sub>			2½	129695	
14°	"		2½?	104098		6P <sub>1</sub>	5d <sup>10</sup> (1S)10p	10p 1P°	0½	140128	201
3D <sub>2</sub>	5d <sup>10</sup> (1S)6d	6d 1D	1½	104983	560	6P <sub>2</sub>			1½	140329	
3D <sub>3</sub>			2½	105543			5a <sub>1</sub>			2½	140135
Y		1P°	1½	106086		6S	5d <sup>10</sup> (1S)11a	11a 1S	0½	141914. 5	
16°			3½	106213		7F <sub>1</sub>	5d <sup>10</sup> (1S)8f	8f 1F°	3½	142127. 3	-3. 4
U	5d <sup>10</sup> (1S)6p <sub>1/2</sub>	1P°	0½	106293		7F <sub>2</sub>			2½	142150. 7	
17b <sub>1</sub>	5d <sup>10</sup> (1S)6p <sub>1/2</sub>		0½	106612		7G	5d <sup>10</sup> (1S)7g	7g 1G	3½, 4½	142293. 4	
18b <sub>1</sub>			1½	106714		7D <sub>2</sub>	5d <sup>10</sup> (1S)10d	10d 1D	1½	142660	150
						7D <sub>3</sub>			2½	142709. 5	
3P <sub>1</sub>	5d <sup>10</sup> (1S)7p	7p 1P°	0½	108298	3672	7P <sub>1</sub>	5d <sup>10</sup> (1S)11p	11p 1P°	0½	142959	158
3P <sub>2</sub>			1½	111970			7P <sub>2</sub>			1½	
20b <sub>1</sub>			3½	108974		7S	5d <sup>10</sup> (1S)12a	12a 1S	0½	144150	
X	5d <sup>10</sup> (1S)6p <sub>1/2</sub>	1P°	1½	109189		8F <sub>1</sub>	5d <sup>10</sup> (1S)9f	9f 1F°	2½	144289	1
22b <sub>1</sub>	5d <sup>10</sup> (1S)6p <sub>1/2</sub>		1½	110603		8F <sub>2</sub>			3½	144290	
19d <sub>1</sub>			3½	110846		8G	5d <sup>10</sup> (1S)8g	8g 1G	3½, 4½	144401. 7	
23b <sub>1</sub>	5d <sup>10</sup> (1S)6p <sub>1/2</sub>		3½	111172		8D <sub>2</sub>	5d <sup>10</sup> (1S)11d	11d 1D	1½	144653. 2	26. 4
						8D <sub>3</sub>			2½	144679. 6	
19d <sub>2</sub>			2½	116200		4D <sub>1</sub>			0½	144910	
19d <sub>3</sub>			1½	117339		9F <sub>1</sub>	5d <sup>10</sup> (1S)10f	10f 1F°	3½	145768	-1
19d <sub>4</sub>			0½	119447		9F <sub>2</sub>			2½	145769	
3S	5d <sup>10</sup> (1S)8a	8a 1S	0½	121416		9G	5d <sup>10</sup> (1S)9g	9g 1G	3½, 4½	145847. 35	
C		1F°	2½	121960		9H	5d <sup>10</sup> (1S)9h	9h 1H°	4½, 5½	145849	
B			1½	122186		9D <sub>2</sub>	5d <sup>10</sup> (1S)12d	12d 1D	1½	146025. 6	27
						9D <sub>3</sub>			2½	146053	
4F <sub>1</sub>	5d <sup>10</sup> (1S)5f	5f 1F°	3½	123152	-257				2½	146647	
4F <sub>2</sub>			2½	123409			10G	5d <sup>10</sup> (1S)10g	10g 1G	3½, 4½	146880. 4
4D <sub>2</sub>	5d <sup>10</sup> (1S)7d	7d 1D	1½	125324	254	10H	5d <sup>10</sup> (1S)10h	10h 1H°	4½, 5½	146882	
4D <sub>3</sub>			2½	125578			10D <sub>2</sub>	5d <sup>10</sup> (1S)13d	13d 1D	1½	147014
4P <sub>1</sub>	5d <sup>10</sup> (1S)8p	8p 1P°	0½	126642	853	10D <sub>3</sub>			2½	147035	
4P <sub>2</sub>			1½	127795			11G	5d <sup>10</sup> (1S)11g	11g 1G	3½, 4½	147645. 0
4S	5d <sup>10</sup> (1S)9a	9a 1S	0½	132559		12D <sub>2</sub>	5d <sup>10</sup> (1S)15d	15d 1D	1½	148311	
4D <sub>1</sub>			3½	132714					2½		
5F <sub>1</sub>	5d <sup>10</sup> (1S)6f	6f 1F°	3½	133267. 5	-82	13D <sub>2</sub>	5d <sup>10</sup> (1S)16d	16d 1D	1½	148743	
5F <sub>2</sub>			2½	133350						2½	
5G	5d <sup>10</sup> (1S)5g	5g 1G	3½, 4½	133653			Hg III(1S <sub>2</sub> )	Limit		151280	

## Hg III

(Pt I sequence; 78 electrons)

Z=80

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 1S_0$  $5d^{10} 1S_0$ , 276000 ± K

I. P. 34.2 volts

Foster has revised the earlier work on this spectrum by Johns and others, added 13 new levels, and published a list of 99 classified lines between 610.46 Å and 8151.64 Å. The energy levels in the table are quoted from his paper. He has reobserved the spectrum from 2240 Å to 10000 Å, by means of an electrodeless discharge, and utilized measurements by others in the short-wave region for transitions between terms otherwise well established. His interpretation is supported by observations of isotope shifts.

Johns has published a list of nearly 500 classified lines between 570.91 Å and 8451.1 Å, attributed to Hg III, and 41 levels not reported by Foster. Most of the additional levels are assigned to the  $5d^9 6,7d$ ;  $5d^9 8s$ ;  $5d^9 7p$ ; and  $5d^9 5f$  configurations. This more extensive list needs further confirmation because some of the lines may not belong to the Hg III spectrum. Nevertheless, "a large part of the analysis as presented by Johns is confirmed . . ."

The spectra of the Pt I isoelectronic sequence are of special theoretical interest because they approach *jj*-coupling rather than *LS*-coupling. This appears more conspicuously in the spectra of higher ionization especially among the levels from the  $5d^9(^3D)6p$  configuration. A detailed study of "The Four Vector Problem and Its Application to Energies and Intensities in Platinum-like Spectra" has been made by Goble. In *LS*-coupling this configuration gives rise to the terms  $^1,3(P^{\circ}D^{\circ}F^{\circ})$ . In *jj*-coupling the same number of energy levels and the same resultant *J*-values occur, but each level is defined by two *j*-values, one for the *d*-electron ( $1\frac{1}{2}$  or  $2\frac{1}{2}$ ), and one for the *p*-electron ( $0\frac{1}{2}$  or  $1\frac{1}{2}$ ). The *j*-values for the *d*- and *p*-electrons, as required for *jj*-coupling, are indicated in the configuration column, the former being the same as the *J*-values of the limit term  $^3D$ .

In the table the writer has arranged the levels in numerical order and retained in the left column the arbitrary numbers used by the authors to designate the various levels. She has assigned *LS*-designations in the case of the  $^1,3D$  terms from the  $5d^9(^3D)ns$  configuration, although the *jj*-coupling applies here also.

Foster adopts *jj*-coupling notation for the levels having the configuration  $5d^8 6s^2$ , assigning the first group of three levels ( $J=4, 2, 0$ ) to " $5d_{3/2, 1/2}^8 6s^2$ ", the next group of four ( $J=3, 2, 1, 4$ ) to " $5d_{3/2, 1/2}^8 6s^2$ ", and the last group of two ( $J=2, 0$ ) to " $5d_{3/2, 1/2}^8 6s^2$ ". A table giving the transformation from *jj*- to *LS*-coupling may be found in the book by Condon and Shortley, p. 294.

Johns states that the limit is approximately 276000, which is quoted here. Mack and Fromer derive the value  $34.3 \pm 0.3$  from a Moseley diagram for the sequence. In making the diagram they used a Rydberg formula for the first two members of the  $5d^9(^3D_{3/2})ns$  series,  $J=3$ , and applied the same Ritz correction as for the Au I-like spectrum of the same stage of ionization.

## REFERENCES

- A. T. Goble, *Phys. Rev.* **48**, 346 (1935).  
 J. E. Mack and M. Fromer, *Phys. Rev.* **48**, 357 (1935). (I P) (T) (C L)  
 M. W. Johns, *Canadian J. Research [A]* **15**, 193 (1937). (I P) (T) (C L) (I S)  
 S. Mrosowski, *Phys. Rev.* **61**, 605 (1942). (I S)  
 E. W. Foster, *Proc. Roy. Soc. London [A]* **200**, 429 (1950). (T) (C L) (I S)  
 E. W. Foster, *Proc. Roy. Soc. London [A]* **208**, 367 (1951). (hfs)  
 E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra*, p. 294 (Cambridge University Press, London, 1951).  
 K. Murakawa, *Phys. Rev.* **93**, 1232 (1954). (hfs)

## Hg III

## Hg III

Author	Config.	Design.	J	Level	Author	Config.	Design.	J	Level
1 <sub>0</sub>	5d <sup>9</sup>	5d <sup>9</sup> 1S	0	0.0	51 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	158885.8
2 <sub>0</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6s	6s <sup>1</sup> D	3	42850.3	14 <sub>0</sub>	5d <sup>9</sup> 6s <sup>2</sup>	6s <sup>2</sup> 1S	0	158900.0
3 <sub>0</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6s	6s <sup>3</sup> D	2	40029.5	52 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	158861.4
4 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6s	6s <sup>1</sup> D	1	58405.8	53 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	160015.7
5 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6s	6s <sup>3</sup> D	2	61065.7	54 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	160170.0
6 <sub>0</sub>	5d <sup>9</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>3</sup> F	4	97893.8	23 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	160789.3
11 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		2	103549.4	55 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	160990.9
21 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		3	105697.8	24 <sub>1</sub>	5d <sup>9</sup> 6s 6p		4	161282.9?
7 <sub>1</sub>	5d <sup>9</sup> 6s <sup>2</sup>		2	106027.5	25 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	161451.4
9 <sub>1</sub>	5d <sup>9</sup> 6s <sup>2</sup>		3	112226.0	26 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	161461.8
51 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		4	117994.5	27 <sub>1</sub>	5d <sup>9</sup> 6s 6p		4	162155.0?
31 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		2	118542.0	28 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	162972.2
41 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		1	118607.4	29 <sub>1</sub>	5d <sup>9</sup> 6s 6p		4	163451.4
11 <sub>1</sub>	5d <sup>9</sup> 6s <sup>2</sup>		2	118926.5	56 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	163818.2
61 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		2	120927.8	30 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	164905.3
71 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		3	121602.0	31 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	165004.8?
8 <sub>0</sub>	5d <sup>9</sup> 6s <sup>2</sup>		0	122661.0	32 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	166179.4
10 <sub>1</sub>	5d <sup>9</sup> 6s <sup>2</sup>		1	122734.9	33 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	166311.1
12 <sub>0</sub>	5d <sup>9</sup> 6s <sup>2</sup>		4	126468.3	34 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	167580.3
81 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		1	128556.3	35 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	169063.0
91 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		0	130702.4	36 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	169666.8
13 <sub>2</sub>	5d <sup>9</sup> 6s <sup>2</sup>		2	133731.5	37 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	171366.7?
101 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		3	134588.0	57 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	171700.6
111 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		1	134998.7	38 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	173968.5
121 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )6p <sub>1/2</sub>		2	136479.0	39 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	175512.0
47 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	147142.4	40 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	177281.0
48 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	148426.0	15 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s		3	178428.2
131 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	149717.3	16 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s		2	179039.1
141 <sub>1</sub>	5d <sup>9</sup> 6s 6p		4	150276.0?	41 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	180246.8
151 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	150278.7	42 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	183544.2
161 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	150857.0	43 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	184698.4
171 <sub>1</sub>	5d <sup>9</sup> 6s 6p		1	151997.5	44 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	187028.5
181 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	153254.6?	27 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s		1	194078.6
191 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	154213.9	28 <sub>1</sub>	5d <sup>9</sup> ( <sup>2</sup> D <sub>3/2</sub> )7s		2	194476.0
491 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	154717.9	45 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	194904.0
201 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	155796.1	46 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	196377.9
211 <sub>1</sub>	5d <sup>9</sup> 6s 6p		3	156580.0					
501 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	157877.2		Hg IV( <sup>2</sup> D <sub>3/2</sub> )	Limit		276000
221 <sub>1</sub>	5d <sup>9</sup> 6s 6p		2	158204.9					

## Hg IV

(Ir I sequence; 77 electrons)

Z=80

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^9 \text{ } ^2D_{3/2}$  $5d^9 \text{ } ^2D_{3/2}$ 

K

I. P.

volts

The spectrum needs further study. By analogy with related spectra the ground level of Hg IV is predicted with certainty to be  $5d^9 \text{ } ^2D_{3/2}$ , as has been pointed out by Edlén. The analysis is from Subbaraya, who has classified about 200 lines between 938 Å and 7517 Å from wavelength measurements by Carroll, Bloch, Déjardin, and Ricard. A homogeneous line list containing accurate wavelengths is needed to confirm and extend the present work, since the limit of tolerance within the multiplets is inexcusably large.

No series are known. The doublet and quartet terms are connected by observed inter-system combinations.

## REFERENCES

- T. S. Subbaraya, Proc. Indian Acad. Sci. [A] 1, 39 (1935). (T) (C L)  
 B. Edlén, letter (February 1956).

## Hg IV

## Hg IV

Author	Config.	Design.	J	Level	Interval	Author	Config.	Design.	J	Level	Interval
1	5d <sup>9</sup> ( <sup>2</sup> F)6s	6s <sup>4</sup> F	4½	0		3°			3½	74708	
4			3½	7557	-7557	4°			3½	75338	
9			2½	12084	-4527	6°			2½	75779	
10			1½	15438	-3354	7°			3½	77045	
2	5d <sup>9</sup>	5d <sup>9</sup> <sup>4</sup> D	2½	2192		8°	5d <sup>9</sup> ( <sup>2</sup> F)6p	6p <sup>4</sup> D°	3½	85058	
7			1½	10876	-8184	18°			2½	88216	-3100
3	5d <sup>7</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>4</sup> F	4½	5653		19°			1½	89513	-1297
5			3½	7897	-2244	20°			0½	90387	-814
6			2½	9476	-1579	14°			2½	85091	
8			1½	10592	-1116	15°			3½	86380	
11	5d <sup>9</sup> ( <sup>2</sup> F)6s	6s <sup>2</sup> <sup>4</sup> P	2½	21011		16°			1½, 2½	87185	
12			1½	23270	-2259	21°			2½	91758	
14			0½	24564	-1294	23°			1½, 2½	101826	
13			5d <sup>9</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>4</sup> P	2½	24054				24°	2½
15	1½	25001			-947	25°	2½	102353			
16	0½	25902			-801	26°	1½, 2½	106210			
17	4½	42131				27°	2½	106385			
18			2½, 3½	44599		28°	1½	106748			
19			3½	53342		29°	1½, 2½	108798			
20			3½	55664		30°	2½	113648			
21			2½	57122		31°	2½	114508			
22			3½	57270							
23			3½	59490							
1°	5d <sup>9</sup> ( <sup>2</sup> F)6p	6p <sup>4</sup> F°	4½	70567							
5°			3½	76656	-5088						
9°			2½	79688	-4033						
12°			1½	83834	-3196						
2°	5d <sup>9</sup> ( <sup>2</sup> F)6p	6p <sup>4</sup> G°	5½	74419							
11°			4½	81039	-6620						
17°			3½	87826	-6786						
22°			2½	92237	-4412						

February 1958.

## Hg IV OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms		
5d <sup>9</sup> 5d <sup>7</sup> 6s <sup>2</sup>	5d <sup>9</sup> <sup>4</sup> D		
	6s <sup>2</sup> <sup>4</sup> P	6s <sup>2</sup> <sup>4</sup> F	
5d <sup>9</sup> ( <sup>2</sup> F)ns 5d <sup>9</sup> ( <sup>2</sup> F)ns'	ns (n ≥ 6)		np (n ≥ 6)
	6s <sup>2</sup> <sup>4</sup> P	6s <sup>2</sup> <sup>4</sup> F	6p <sup>4</sup> D°   6p <sup>4</sup> F°   6p <sup>4</sup> G°

\*For predicted terms in the spectra of the Ir I isoelectronic sequence, see Vol. III, Introduction.

## THALLIUM

Tl I

81 electrons

Z=81

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^2 P_{1/2}^0$  $6p^2 P_{3/2}^0$  49264.3 K

I. P. 6.106 volts

The analysis is chiefly from Fowler and Paschen-Götze, but some terms have been added from later papers. Meggers and Murphy have observed Tl I in the near infrared and added the terms  $7, 8^1 P^0$ . Beutler and Demeter have observed in absorption, ultraviolet combinations with the ground state giving the terms  $nS$  ( $n=8$  to 20) and  $nD$  ( $n=6$  to 23), thus extending the former series by one member, and the latter by five. For the  $^3D$  series they have observed only the level with  $J=1\frac{1}{2}$ , but for  $n>12$  these terms are listed in the table as unresolved. They give, also, the levels  $^4P_{3/2, 1/2}$ ,  $^3D_{3/2}$ , and  $^3P_{0/1}$  from the  $6p^2$  configuration. Clearman has measured 13 Tl I lines, adding the component  $6p^2 ^4P_{3/2}$  and the levels labeled  $2^0$  and  $3^0$  in the table.

The writer has utilized the later observations to improve as many values of the Tl I energy levels as possible. For levels above the ionization limit, rounded off values are tabulated.

Beutler and Demeter list, also, 9 lines observed between 651.42 Å and 891 Å (153510 K and 112250 K, respectively). They suggest that these lines are probably due to the transitions involving configurations  $5d^{10} 6s^2 6p_{3/2} - 5d^9 6s^2 6p_{3/2} np$  (or possibly  $nf$ ).

There are approximately 160 classified lines extending from 651 Å to 51057 Å, including a number of forbidden combinations. The observations by various authors do not yield precise term values even though the analysis is essentially complete. The spectrum exhibits hyperfine structure and isotope shift, and several papers deal with Paschen-Back effect and the Zeeman effect of hyperfine structure. Only a limited number of references to these topics is included below. Back and Wulff report the hyperfine structure separation of the ground level as 0.708K.

The series are long and well established. The limit quoted here is from Fowler. Beutler and Demeter state that the term  $10s^2 S$  is perturbed because its combination with the ground state is affected by a neighboring line classified as  $6p^2 P_{3/2}^0 - 6p^2 ^4P_{3/2}$ . This is based on their study of the series fitted to a Rydberg-Ritz formula.

## REFERENCES

- F. Paschen and R. Götze, *Seriengesetze der Linienspektren*, p. 132 (Julius Springer, Berlin, 1922). (I P) (T) (C L)
- A. Fowler, *Report on Series in Line Spectra*, p. 160 (Fleetway Press, London, 1922). (I P) (T) (C L)
- T. Takamine and M. Fukuda, *Jap. J. Phys.* **2**, 111 (1923). (C L)
- E. Back and J. Wulff, *Zeit. Phys.* **68**, 31 (1930). (Z E) (hfs)
- J. Wulff, *Zeit. Phys.* **68**, 70 (1931). (Z E) (hfs)
- H. Schöler and J. E. Keyston, *Naturwiss.* **19**, 320 (1931). (I S)
- H. Beutler and W. Demeter, *Zeit. Phys.* **91**, 143, 202, 218 (1934). (T) (C L)
- J. Okubo and S. Saitō, *Zeeman Verhandelingen* p. 253 (Martinus Nijhoff, The Hague, 1935); *Sci. Reports Tôhoku Imp. Univ.* [1] **26**, 141 (1936). (Z E) (hfs) (I S)
- H. E. Clearman, *J. Opt. Soc. Am.* **42**, 373 (1952). (T) (C L)
- W. F. Meggers and R. J. Murphy, *J. Research Nat. Bur. Std.* **48**, 334, RP 2320 (1952). (I P) (T) (C L)
- H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, 61 (1953). (Summary hfs)

Tl I					Tl I				
Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
6 <sup>o</sup> (S)6p	6p <sup>o</sup> P <sup>o</sup>	0½ 1½	0.0 7782.7	7792.7	6 <sup>o</sup> (S)12e	12e <sup>o</sup> S	0½	47654.7	
6 <sup>o</sup> (S)7a	7a <sup>o</sup> S	0½	26477.5		6 <sup>o</sup> (S)12p	12p <sup>o</sup> P <sup>o</sup>	0½, 1½	47847.7	
6 <sup>o</sup> (S)7p	7p <sup>o</sup> P <sup>o</sup>	0½ 1½	24169.9 25161.1	1001.2	6 <sup>o</sup> (S)12d	12d <sup>o</sup> D	1½, 2½	47878.0	
6 <sup>o</sup> (S)6d	6d <sup>o</sup> D	1½ 2½	26117.9 26199.9	82.0	6 <sup>o</sup> (S)14e	14e <sup>o</sup> S	0½	47982.2	
6 <sup>o</sup> (S)8e	8e <sup>o</sup> S	0½	28745.9		6 <sup>o</sup> (S)14p	14p <sup>o</sup> P <sup>o</sup>	0½, 1½	48189.6	
6 <sup>o</sup> (S)8p	8p <sup>o</sup> P <sup>o</sup>	0½ 1½	41998.1 41740.8	272.7	6 <sup>o</sup> (S)12d	12d <sup>o</sup> D	1½, 2½	48142.3	
6 <sup>o</sup> (S)7d	7d <sup>o</sup> D	1½ 2½	42011.4 42049.0	37.6	6 <sup>o</sup> (S)15e	15e <sup>o</sup> S	0½	48222.2	
6 <sup>o</sup> (S)5f	5f <sup>o</sup> F <sup>o</sup>	2½, 3½	42318.4		6 <sup>o</sup> (S)15p	15p <sup>o</sup> P <sup>o</sup>	0½, 1½	48331.2	
6 <sup>o</sup> (S)9e	9e <sup>o</sup> S	0½	43166.2		6 <sup>o</sup> (S)14d	14d <sup>o</sup> D	1½, 2½	48339.3	
6 <sup>o</sup> (S)9p	9p <sup>o</sup> P <sup>o</sup>	0½ 1½	44380.9 44662.5	181.6	6 <sup>o</sup> (S)16e	16e <sup>o</sup> S	0½	48399.5	
6 <sup>o</sup> (S)8d	8d <sup>o</sup> D	1½ 2½	44672.6 44692.7	20.1	6 <sup>o</sup> (S)16p	16p <sup>o</sup> P <sup>o</sup>	0½, 1½	48459.5	
6 <sup>o</sup> (S)6f	6f <sup>o</sup> F <sup>o</sup>	2½, 3½	44823.5		6 <sup>o</sup> (S)15d	15d <sup>o</sup> D	1½, 2½	48488.6	
6 <sup>o</sup> 6p <sup>a</sup>	6p <sup>a</sup> <sup>o</sup> P	0½ 1½ 2½	45220 49800 53050	4580 3250	6 <sup>o</sup> (S)17e	17e <sup>o</sup> S	0½	48534.8	
6 <sup>o</sup> (S)10e	10e <sup>o</sup> S	0½	45296.8		6 <sup>o</sup> (S)16d	16d <sup>o</sup> D	1½, 2½	48604.0	
6 <sup>o</sup> (S)10p	10p <sup>o</sup> P <sup>o</sup>	0½ 1½	45929.3 46045.6	104.3	6 <sup>o</sup> (S)18e	18e <sup>o</sup> S	0½	48639.0	
6 <sup>o</sup> (S)9d	9d <sup>o</sup> D	1½ 2½	46098.5 46110.3	11.8	6 <sup>o</sup> (S)17d	17d <sup>o</sup> D	1½, 2½	48696.5	
6 <sup>o</sup> (S)7f	7f <sup>o</sup> F <sup>o</sup>	2½, 3½	46185.3		6 <sup>o</sup> (S)19e	19e <sup>o</sup> S	0½	48726.2	
6 <sup>o</sup> (S)11e	11e <sup>o</sup> S	0½	46456.9		6 <sup>o</sup> (S)18d	18d <sup>o</sup> D	1½, 2½	48770.5	
6 <sup>o</sup> (S)11p	11p <sup>o</sup> P <sup>o</sup>	0½ 1½	46853.8 46917.1	63.3	6 <sup>o</sup> (S)20e	20e <sup>o</sup> S	0½	48796.2	
6 <sup>o</sup> (S)10d	10d <sup>o</sup> D	1½ 2½	46949.9 46958.0	8.1	6 <sup>o</sup> (S)19d	19d <sup>o</sup> D	1½, 2½	48828.6	
6 <sup>o</sup> (S)8f	8f <sup>o</sup> F <sup>o</sup>	2½, 3½	47004.6		6 <sup>o</sup> (S)20d	20d <sup>o</sup> D	1½, 2½	48878.2	
6 <sup>o</sup> (S)12e	12e <sup>o</sup> S	0½	47178.9		6 <sup>o</sup> (S)21d	21d <sup>o</sup> D	1½, 2½	48920.6	
6 <sup>o</sup> (S)12p	12p <sup>o</sup> P <sup>o</sup>	0½ 1½	47442.6 47477.4	34.8	6 <sup>o</sup> (S)22d	22d <sup>o</sup> D	1½, 2½	48957.7	
6 <sup>o</sup> (S)11d	11d <sup>o</sup> D	1½ 2½	47499.8 47504.1	4.37	6 <sup>o</sup> (S)23d	23d <sup>o</sup> D	1½, 2½	48988.8	
					Tl II (S <sub>0</sub> )	Limit		49264.2	
					6 <sup>o</sup> 6p <sup>a</sup>	6p <sup>a</sup> <sup>o</sup> D	1½ 2½	62000	
					6 <sup>o</sup> 6p <sup>a</sup>	6p <sup>a</sup> <sup>o</sup> P	0½ 1½	67150	
					6 <sup>o</sup> 6p(2P)7e	2°		75970	
						3°	1½	96740	

January 1955.

## Tl: OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>10</sup> +	Observed Terms			
6s <sup>2</sup> ( <sup>3</sup> S)6p	6p <sup>3</sup> P°			
6s 6p <sup>2</sup>	{ 6p <sup>2</sup> <sup>3</sup> P 6p <sup>2</sup> <sup>1</sup> P 6p <sup>2</sup> <sup>3</sup> D			
	<i>ns</i> (n ≥ 7)	<i>np</i> (n ≥ 7)	<i>nd</i> (n ≥ 6)	<i>nf</i> (n ≥ 5)
6s <sup>2</sup> ( <sup>3</sup> S) <i>ns</i>	7 to 30s <sup>3</sup> S	7 to 16p <sup>3</sup> P°	6 to 23d <sup>3</sup> D	5 to 8f <sup>3</sup> F°

\*For predicted terms in the spectra of the Tl I isoelectronic sequence, see Vol. III, Introduction.

## Tl II

(Hg I sequence; 80 electrons)

Z=81

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>4</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>10</sup> 6s<sup>2</sup> <sup>1</sup>S<sub>0</sub>

6s<sup>2</sup> <sup>1</sup>S<sub>0</sub> 164765 ± 5 K

I. P. 20.42 volts

Ellis and Sawyer have revised the early work on Tl II and almost doubled the number of classified lines by observing the spectrum with a hollow cathode source, from 639.08 Å to 9254 Å. The total number of classified lines is 247.

Hyperfine structure affects many of the lines and has been resolved by various investigators, including Smith, McLennan and Crawford, and others.

The Zeeman and Paschen-Back effects have also been observed for a few selected lines.

Perturbations in the series are discussed by Ellis and Sawyer. They derive the limit quoted above from the *ns* <sup>3</sup>S and *nd* <sup>3</sup>D series, n=7 to 12 and 6 to 11, respectively. A plot of the quantum defects for each of the series shows that they are unperturbed, according to these authors.

Observed intersystem combinations connect the singlet and triplet systems of terms.

## REFERENCES

- Stanley Smith, *Phys. Rev.* **25**, 235 (1930). (I P) (T) (C L) (hfs)  
 J. C. McLennan and A. M. I. A. W. Durnford, *Proc. Roy. Soc. London [A]* **129**, 48 (1930). (Z E)  
 J. C. McLennan and M. F. Crawford, *Proc. Roy. Soc. London [A]* **133**, 10 (1931). (T) (C L) (hfs)  
 A. M. Crooker, *Phil. Mag.* [7] **16**, 994 (1933). (Z E)

## Tl II

## Tl II

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
5d <sup>10</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>1</sup> S	0	0		5d <sup>10</sup> 6s( <sup>3</sup> S)6d	6d <sup>3</sup> D	1	116147	283
							2	116430	396
							3	116826	
5d <sup>10</sup> 6s( <sup>3</sup> S)6p	6p <sup>3</sup> P°	0	49451	2942	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>3</sup> P	0	117408	7930
		1	52393	9332			1	125338	3479
		2	61726				2	128817	
5d <sup>10</sup> 6s( <sup>3</sup> S)6p	6p <sup>1</sup> P°	1	75600		5d <sup>10</sup> 6s( <sup>3</sup> S)7p	7p <sup>3</sup> P°	0	119361	215
5d <sup>10</sup> 6s( <sup>3</sup> S)7s	7s <sup>3</sup> S	1	105225				1	119676	2453
							2	122080	
5d <sup>10</sup> 6s( <sup>3</sup> S)7s	7s <sup>1</sup> S	0	107996		5d <sup>10</sup> 6s( <sup>3</sup> S)7p	7p <sup>1</sup> P°	1	122570	
5d <sup>10</sup> 6s( <sup>1</sup> D)6p	1°	2	110387		5d <sup>10</sup> 6s( <sup>1</sup> D)6p	4°	2	125437	
5d <sup>10</sup> 6s( <sup>3</sup> S)6d	6d <sup>1</sup> D	2	115160		5d <sup>10</sup> 6s( <sup>1</sup> D)6p	5°	1	126204	



Tl II—Continued

Tl II—Continued

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>1</sup> D)6p	6°	3	139933		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)9p	9p <sup>1</sup> P°	0		
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>1</sup> D)6p	7°	2	139153				1	148319	476
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)8s	8s <sup>3</sup> S	1	133563				2	148395	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)8s	8s <sup>3</sup> S	0	134292		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)8d	8d <sup>1</sup> D	2	148465	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>1</sup> D)6p	8°	1	134392		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)9p	9p <sup>1</sup> P°	1	149063	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)5f	5f <sup>1</sup> F°	2	136216	-101	5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)10s	10s <sup>3</sup> S	1	151568	
		3	136115	115	5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)7f	7f <sup>1</sup> F°	2	152104	6
		4	136230				3, 4	152110	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)5f	5f <sup>1</sup> F°	3	136263		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)7f	7f <sup>1</sup> F°	3	152140	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)7d	7d <sup>1</sup> D	2	136891		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)6g	6g <sup>1,2</sup> G	3, 4, 5	152470	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)7d	7d <sup>1</sup> D	1	137927	126	5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)9d	9d <sup>1</sup> D	1	152819	28
		2	138052	150			2	152847	59
		3	138203				3	152906	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)8p	8p <sup>1</sup> P°	0			5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)9d	9d <sup>1</sup> D	2	153200	
		1	139995	939	5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)10p	10p <sup>1</sup> P°	0		
		2	140904				1		
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)8p	8p <sup>1</sup> P°	1	141000				2	153530?	
5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> D	2	141982		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)10p	10p <sup>1</sup> P°	1	153633?	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>1</sup> D)6p	10°	3	142781		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)11s	11s <sup>3</sup> S	1	155181	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>1</sup> D)6p	11°	1	143612		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)10d	10d <sup>1</sup> D	1	155963	19
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>1</sup> D)6p	12°	2	145092				2	155982	36
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)9s	9s <sup>3</sup> S	1	145415				3	156018	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)9s	9s <sup>3</sup> S	0	145591		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)10d	10d <sup>1</sup> D	2	156177	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)6f	6f <sup>1</sup> F°	4	146500	-34	5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)11p	11p <sup>1</sup> P°	1	156475?	
		3	146534	11	5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)12s	12s <sup>3</sup> S	1	157481	
		2	146523		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)11d	11d <sup>1</sup> D	1		
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)6f	6f <sup>1</sup> F°	3	146543				2	158027?	11
							3	158038?	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)5g	5g <sup>1,2</sup> G	3, 4, 5	147065		5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)11d	11d <sup>1</sup> D	2	158136?	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)8d	8d <sup>1</sup> D	1	147602	50					
		2	147652	95	Tl III( <sup>3</sup> S <sub>1/2</sub> )	Limit		164765 ± 5	
		3	147747						

December 1954.

Tl II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms		
5d <sup>10</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>3</sup> S		
5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> P 6p <sup>2</sup> <sup>1</sup> D		
	ns (n ≥ 7)	np (n ≥ 6)	nd (n ≥ 6)
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)ns	{ 7 to 12s <sup>3</sup> S 7 to 9s <sup>3</sup> S	6 to 10p <sup>1</sup> P° 6 to 11p <sup>1</sup> P°	6 to 11d <sup>1</sup> D 6 to 11d <sup>1</sup> D
	nf (n ≥ 5)	ng (n ≥ 5)	
5d <sup>10</sup> 6s <sup>2</sup> ( <sup>3</sup> S)ns	{ 5 to 7f <sup>1</sup> F° 5 to 7f <sup>1</sup> F°	5, 6g <sup>1,2</sup> G	

\*For predicted terms in the spectra of the Hg I isoelectronic sequence, see Vol. III, Introduction.

## TI III

(Au I sequence; 79 electrons)

Z=81

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 S_{6s}$  $6s^2 S_{6s}$ , 240000 K

I. P. 29.8 volts

The terms in the table are from McLennan, McLay, and Crawford (1929), who extended the work of Carroll. They list 22 classified lines between 1231.57 Å and 5927.8 Å and one at 8001 Å. The analysis is seriously incomplete. The inverted  $^3D$  term from  $5d^9 6s^2$ , whose lower component should lie between  $6p^2 P^o$  levels and whose interval is approximately 18000, is not reported in this paper.

P. Pattabhiramayya and A. S. Rao include this term in the list of 2 doublet and 6 quartet terms, which they report as an extension to the above analysis. Their work brings the total of classified lines to 89, but their data are omitted here pending further confirmation. The discrepancies within the multiplets between the observed and calculated wave numbers indicate that more precise wavelengths are needed to extend the analysis.

Crooker describes the Paschen-Back effect observed for  $\lambda 5362$ ,  $7s^2 S_{6s} - 7p^2 P_{3/2}$ . The two 1930 papers by McLennan and his associates discuss the observed hyperfine structure and the observed Zeeman effects of the hyperfine structure components. They include among their classified lines one designation involving the level  $5d^9 6s^2 ^3D_{13/2}$  and another involving the level  $5d^9 6s 6p^2 P_{3/2}$ . The respective new levels from these two lines are approximately 84595 and 155851. More combinations are required to confirm them.

The limit derived from the first three members of the  $^2S$  series by means of a Hicks formula is 240300. The authors have increased this value by 300 to give an effective total quantum number for  $5g^2 G$  that is nearly hydrogenic, i. e., 4.995.

## REFERENCES

- J. C. McLennan, A. B. McLay, and M. F. Crawford, Proc. Roy. Soc. London [A] **125**, 50 (1929). (I P) (T) (C L)  
 J. C. McLennan and E. J. Allin, Proc. Roy. Soc. London [A] **129**, 43 (1930). (hfs)  
 P. Pattabhiramayya and A. S. Rao, Indian J. Phys. **5**, 407 (1930). (T) (C L)  
 J. C. McLennan and A. M. I. A. W. Durnford, Proc. Roy. Soc. London [A] **129**, 48 (1930). (Z E) (hfs)  
 G. Arvidsson, Nature **126**, 565 (1930). (hfs)  
 A. M. Crooker, Phil. Mag [7] **16**, 994 (1933). (Z E)

## TI III

## TI III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$5d^{10}(^1S)6s$	$6s^2 S$	0	0		$5d^{10}(^1S)5f$	$5f^2 F^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	175593 178965	-1362
$5d^{10}(^1S)6p$	$6p^2 P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	64157 78970	14813	$5d^{10}(^1S)8s$	$8s^2 S$	0	183187	
$5d^{10}(^1S)7s$	$7s^2 S$	0	139209		$5d^{10}(^1S)7d$	$7d^2 D$	$1\frac{1}{2}$ $2\frac{1}{2}$	186356 186948	592
$5d^{10}(^1S)6d$	$6d^2 D$	$1\frac{1}{2}$ $2\frac{1}{2}$	145355 146669	1314	$5d^{10}(^1S)5g$	$5g^2 G$	$3\frac{1}{2}$ , $4\frac{1}{2}$	201011	
$5d^{10}(^1S)7p$	$7p^2 P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	157852 163654	5682	-----	-----	-----	-----	-----
					TI IV( $^1S_6$ )	Limit	-----	240600	

November 1954.

## Tl IV

(Pt I sequence; 78 electrons)

Z=81

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 1S_0$  $5d^{10} 1S_0$ , 400100 K

I. P. 50.7 volts

The analysis is from Mack and Fromer, who have revised and extended the earlier work of Rao and others. About 35 lines have been classified in the interval 531.260 Å to 1974.6 Å, and most of the results are based on observations by Arvidsson.

The spectra of the Pt I isoelectronic sequence are of special theoretical interest because they approach *jj*-coupling rather than *LS*- or *JI*-coupling. This appears more conspicuously in the spectra of higher ionization, especially among the levels from the  $5d^9(^2D)6p$  configuration. A detailed study of "The Four Vector Problem and Its Application to Energies and Intensities in Platinum-like Spectra" has been made by Goble. In *LS*-coupling this configuration gives rise to the terms  $^1S(^2D^2F^2)$ . In *jj*-coupling the same number of energy levels and the same resultant *J*-values occur, but each level is defined by two *j*-values, one for the *d*-electron (1½ or 2½), and one for the *p*-electron (0½ or 1½).

In the table the writer has assigned *LS*-designations only in the case of the  $^1D$  terms from the  $5d^9(^2D)ns$  configuration, although the *jj*-coupling applies here, also. The *j*-values for the *d*- and *p*-electrons, as required for *jj*-coupling, are indicated in the configuration column, the former being the same as the *J*-values of the limit term  $^2D$ . A table giving the transformation from *jj*- to *LS*-coupling may be found in the book by Condon and Shortley, p. 294.

The limit has been calculated by the writer from the ionization potential given by Mack and Fromer, which is obtained from a Moseley diagram for the sequence.

## REFERENCES

- G. Arvidsson, *Nature* **136**, 565 (1930). (hfs)  
 A. T. Goble, *Phys. Rev.* **48**, 346 (1935).  
 J. E. Mack and M. Fromer, *Phys. Rev.* **48**, 357 (1935). (I P) (T) (C L)  
 E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra* p. 294 (Cambridge University Press, London, 1951).

## Tl IV

## Tl IV

Config.	Desig.	<i>J</i>	Level	Interval	Config.	Desig.	<i>J</i>	Level	Interval
$5d^{10}$	$5d^{10} 1S$	0	0		$5d^9(^2D_{3/2})6p_{1/2}$		2	170354	
$5d^9(^2D_{3/2})6s$	$6s 3D$	3	75052	-3595 -15029	$5d^9(^2D_{3/2})6p_{1/2}$		3	172272	
$5d^9(^2D_{1/2})6s$		2	78647		$5d^9(^2D)6p$		1	175290	
$5d^9(^2D_{1/2})6s$	$6s 1D$	1	93676		$5d^9(^2D_{1/2})6p_{1/2}$		0	181083	
$5d^9(^2D_{3/2})6p_{3/2}$		2	147635		$5d^9(^2D_{1/2})6p_{1/2}$		3	187967	
$5d^9(^2D_{3/2})6p_{1/2}$		3	149841		$5d^9(^2D_{3/2})6p_{3/2}$		1	188253	
$5d^9(^2D_{1/2})6p_{3/2}$		2	166485		$5d^9(^2D_{1/2})6p_{1/2}$		2	190144	
$5d^9(^2D)6p$		1	167499						
$5d^9(^2D_{3/2})6p_{1/2}$		4	167672		Tl v( $^2D_{3/2}$ )	Limit		[400100 ±]	

August 1954.

## LEAD

## Pb I

82 electrons

Z=82

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^2 \text{ } ^3\text{P}_0$  $6p^2 \text{ } ^3\text{P}_0$ , 59821.0  $\pm$  0.5 K

I. P. 7.415 volts

Gieseler and Grotrian have published the most extensive analysis of Pb I. Although series have long been recognized, the spectrum has needed more thorough study. Consequently, Meggers has reobserved the Pb I spectrum from 2000 Å to 11360.0 Å, and from 1614.73 Å to 2000 Å utilized spectrograms furnished by Shenstone. With an electrodeless ultra-high-frequency source he has measured some 355 lines in the interval 6560.77 Å to longer waves. Of these, about 28 percent can be attributed to the known series.

The analysis has been revised and extended by Meggers for inclusion here, and some levels have been rearranged by the writer with the aid of the new observations in the long-wave region. There are approximately 240 classified lines to date, and very few observed lines between 1614 Å and 6560 Å remain unclassified. All level values have been revised on the basis of the new observations. The limit quoted by Bacher and Goudsmit has been verified by R. E. Trees by means of a Ritz formula applied to the  $nd \text{ } ^3\text{D}_1^o$  series,  $n=6$  to 20. Trees points out that this series is the one least likely to be perturbed.

The F-series are based chiefly on the evidence afforded by the Rydberg denominators. The LS-designations require further confirmation. They are included only to enable the user to identify the series for each J-value.

A number of tentative new levels that need further confirmation have been omitted from the table. They may belong to the configurations  $6p^3$  or  $6s 6p^2(^3\text{P})n\pi'$ . Four lines observed by Randall between 12564 Å and 15316 Å can be classified from the known terms. More observations in the infrared beyond the photographic limit, are needed.

Numerous papers deal with the Zeeman effect, hyperfine structure, and intensities of forbidden lines. Only a few references to these are listed here. The observed g-values in the table are quoted from Back.

The large interval of the ground term  $^3\text{P}^o$  in Pb II, 14081 K, results in a departure from LS-coupling in Pb I. Consequently, an arbitrary arrangement has been adopted in tabulating the levels. Pairs of related levels are listed together, with LS-designations indicating the series.

## REFERENCES

- H. Gieseler und W. Grotrian, *Zeit. Phys.* **24**, 374 (1925); **29**, 377 (1926). (I P) (T) (C L)  
 E. Back, *Zeit. Phys.* **37**, 193 (1926); **43**, 317 (1927). (Z E)  
 R. F. Bacher and S. Goudsmit, *Atomic Energy States*, p. 356 (McGraw-Hill Book Co., Inc., New York, N. Y. and London, 1932). (T)  
 S. Mrozowski, *Phys. Rev.* **68**, 1086 (1940). (hfs)  
 F. A. Jenkins and S. Mrozowski, *Phys. Rev.* **59**, 808 (1941); **60**, 225 (1941). (Z E) (hfs)  
 H. E. Clearman, *J. Opt. Soc. Am.* **42**, 375 (1952). (C L)  
 P. F. A. Klinkenberg, *Rev. Mod. Phys.* **24**, No. 2, 63 (1952). (Summary hfs)  
 K. Murakawa, *J. Phys. Soc. Japan* **8**, 382 (1953); **9**, 876 (1954). (I S) (hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab. ORNL-1469, Suppl. II 32 (1955). (Summary hfs)  
 R. E. Trees, unpublished material (May 1955). (I P)  
 W. F. Meggers, unpublished material (May 1955). (T) (C L)

Pb I

Pb I

Config.	Desig.	J	Level	Interval	Obs. $\rho$	Config.	Desig.	J	Level	Interval	Obs. $\rho$
6 <sup>s</sup> 6 <sup>p</sup>	6 <sup>p</sup> <sup>1</sup> P	0	0.00		0/0	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )8d	8d <sup>1</sup> D°	2	55084.90		
		1	7819.35	7819.35	1.501			1	55158.25	-74.05	
		2	10650.47	2831.12	1.269						
6 <sup>s</sup> 6 <sup>p</sup>	6 <sup>p</sup> <sup>1</sup> D	2	21457.90		1.230	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )6f	6f <sup>1</sup> F 6f <sup>2</sup> F	3	55352.34		16.24
		4						4	55368.58		
6 <sup>s</sup> 6 <sup>p</sup>	6 <sup>p</sup> <sup>1</sup> S	0	29466.81		0/0	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )9p	9p <sup>1</sup> D	1	55360.15		4.12
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7s	7s <sup>1</sup> P°	0	34959.90		0/0	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )6f	6f <sup>1</sup> F	2	55377.52		108.68
		1	35287.24	327.34	1.349			3	55486.20		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7p	7p <sup>1</sup> P	1	42918.68	-1482.24		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )10s	10s <sup>1</sup> P°	0	55706.24		14.28
		0	44400.92					1	55790.52		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7p	7p <sup>1</sup> D	1	44675.00	134.41		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )9d	9d <sup>1</sup> F°	2	55590.60		74.14
		2	44809.41					3	55600.74		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )6d	6d <sup>1</sup> F°	2	45443.28	885.55	0.796	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )9d	9d <sup>1</sup> D°	2	55663.29		-41.54
		3	46322.81		1.116			1	55604.83		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )6d	6d <sup>1</sup> D°	2	46060.90	-7.67	1.247	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7f	7f <sup>1</sup> F 7f <sup>2</sup> F	3	56730.11		3.11
		1	46082.57		0.864			4	56733.22		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7s	7s <sup>1</sup> P°	2	48182.67	-1250.90	1.496	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7f	7f <sup>1</sup> F	2	56743.47		38.96
		1	48439.67		1.131			3	56782.43		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )8s	8s <sup>1</sup> P°	1	48686.87	-39.29	1.304	6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )11s	11s <sup>1</sup> P°	0	56935.44		6.82
		0	48729.18		0/0			1	56942.28		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )8p	8p <sup>1</sup> P	1	51320.53	-465.37		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )10p	10p <sup>1</sup> P	1	57010.11		
		0	51785.90					0			
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )8p	8p <sup>1</sup> D	1	51916.80	27.14		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7p	7p <sup>1</sup> D	3	57371.75		
		2	51943.94								
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7d	7d <sup>1</sup> F°	2	52101.77	310.5		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )10d	10d <sup>1</sup> F°	2	57424.09		45.31
		3	52412.5					3	57489.40		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )7d	7d <sup>1</sup> D°	2	52311.37	-188.16		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )10d	10d <sup>1</sup> D°	2	57444.55		-26.69
		1	52499.53					1	57471.24		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )5f	5f <sup>1</sup> F	2	52850.4			6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )8f	8f <sup>1</sup> F 8f <sup>2</sup> F	3	57552.95		2.18
		3						4	57555.13		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )5f	5f <sup>1</sup> F 5f <sup>2</sup> F	3				6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )8f	8f <sup>1</sup> F	2			
		4	52857.81					3	57575.83		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )9s	9s <sup>1</sup> P°	0	53475.42	35.82		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )12s	12s <sup>1</sup> P°	0	57684.5		4.2
		1	53511.24					1	57688.73		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )9p	9p <sup>1</sup> P	1	54929.0	67.7		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )11d	11d <sup>1</sup> F°	2	57996.01?		32.59
		0	54861.29					3	58022.80		
6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )8d	8d <sup>1</sup> F°	2	55003.24	139.88		6 <sup>s</sup> 6 <sup>p</sup> (P <sub>6</sub> )11d	11d <sup>1</sup> D°	2	58012.14?		-17.88
		3	55143.22					1	58030.02		

Pb I—Continued

Pb I—Continued

Config.	Desig.	J	Level	Interval	Obs. g	Config.	Desig.	J	Level	Interval	Obs. g
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 9f	9f <sup>1</sup> F	3	58086. 22	1. 64		6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 17s	17s <sup>3</sup> P <sup>o</sup>	0	59080. 4		
	9f <sup>3</sup> F	4	58087. 86				1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 9f	9f <sup>3</sup> F	2	58101. 28			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 16d	16d <sup>3</sup> D <sup>o</sup>	2	59151. 3		
		3					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 13s	13s <sup>3</sup> P <sup>o</sup>	0	58179. 3			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 17d	17d <sup>3</sup> D <sup>o</sup>	2	59244. 9		
		1					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d <sup>3</sup> F <sup>o</sup>	2	58409. 447			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 18d	18d <sup>3</sup> D <sup>o</sup>	2	59320. 4		
		3					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d <sup>3</sup> D <sup>o</sup>	2	58411. 29			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 19d	19d <sup>3</sup> D <sup>o</sup>	2	59382. 1		
		1					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 10f	10f <sup>1</sup> F	3	58451. 437	1. 20		6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>3</sup> P <sup>o</sup>	1	59427. 0f		
	10f <sup>3</sup> F	4	58452. 63				1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 14s	14s <sup>3</sup> P <sup>o</sup>	0	58517. 67			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 20d	20d <sup>3</sup> D <sup>o</sup>	2	59432. 5		
		1					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 6d	6d <sup>3</sup> D <sup>o</sup>	3	58518. 3			Pb II( <sup>2</sup> P <sub>1/2</sub> )	Limit	--	59821. 0		
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 13d	13d <sup>3</sup> D <sup>o</sup>	2	58683. 4			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 8s	8s <sup>3</sup> P <sup>o</sup>	2	62620. 9		
		1					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 11f	11f <sup>1</sup> F	3	58712. 40			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 8p	8p <sup>3</sup> D	3	65751. 51		
	11f <sup>3</sup> F	4					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 15s	15s <sup>3</sup> P <sup>o</sup>	0	58761. 0			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 7d	7d <sup>3</sup> P <sup>o</sup>	1	65871. 3f		
		1					3				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 14d	14d <sup>3</sup> D <sup>o</sup>	2	58822. 4			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 7d	7d <sup>3</sup> D <sup>o</sup>	3	66035. 1		
		1					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 16s	16s <sup>3</sup> P <sup>o</sup>	0	58941. 8			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 9s	9s <sup>3</sup> P <sup>o</sup>	2	67481. 6		
		1					1				
6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 15d	15d <sup>3</sup> D <sup>o</sup>	2	59034. 3			6s <sup>2</sup> 6p( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d <sup>3</sup> P <sup>o</sup>	1	68925. 5f		
		1					3				
						Pb II( <sup>2</sup> P <sub>1/2</sub> )	Limit	---	73901		

May 1955.

Pb I OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>10</sup> +	Observed Terms	
6s <sup>2</sup> 6p <sup>2</sup>	{ 6p <sup>2</sup> <sup>3</sup> S	6p <sup>2</sup> <sup>3</sup> P
		6p <sup>2</sup> <sup>1</sup> D
6s <sup>2</sup> 6p( <sup>2</sup> P <sup>o</sup> ) n <sub>r</sub>	{ ns (n ≥ 7)	np (n ≥ 7)
	{ 7 to 17s <sup>3</sup> P <sup>o</sup> 7s <sup>1</sup> P <sup>o</sup>	7 to 10p <sup>3</sup> P      7 to 9p <sup>3</sup> D
6s <sup>2</sup> 6p( <sup>2</sup> P <sup>o</sup> ) n <sub>r</sub>	{ nd (n ≥ 6)	nf (n ≥ 5)
	{ 6 to 8d <sup>3</sup> P <sup>o</sup> 6 to 20d <sup>3</sup> D <sup>o</sup> 6 to 12d <sup>3</sup> F <sup>o</sup>	5 to 10f <sup>3</sup> F <sup>o</sup> 6 to 11f <sup>1</sup> F <sup>o</sup>

\*For predicted terms in the spectra of the Pb I isoelectronic sequence, see Vol. III, Introduction.

## Pb II

(II) sequence; 81 electrons)

Z=82

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^4 \text{ } ^3\text{P}_{0,1}^o$  $6p \text{ } ^3\text{P}_{0,1}^o$ , 121343 K

I. P. 15.028 volts

The analysis is from Earls and Sawyer, who have revised and extended the earlier work by Fr. Gieseler and others. "The spectrum was photographed from 800 Å to 10000 Å." Fourteen lines, observed also by Arvidsson in the vacuum region, were utilized in the analysis and in the determination of the limit. There are approximately 250 classified lines in the interval 840.25 Å to 9864.2 Å.

The limits of the  $^3\text{F}^o$  and  $^3\text{G}$  series have been used to derive the ionization limit, which has an estimated error of 2 or 3 K. The  $^3\text{S}$  and  $^3\text{D}$  series show irregularities due to perturbations caused by "intermingling levels from the  $6s \text{ } 6p^2$  configuration."

Green and Loring have observed the Zeeman effect of Pb II and published 15  $g$ -values based on the analysis by Fr. Gieseler. Earls and Sawyer have used these Zeeman observations to derive  $g$ -values for 10 "even" levels, by assuming the appropriate unperturbed theoretical  $g$ -values for the respective "odd" levels and applying the formulas of Shenstone and Blair for unresolved patterns. The writer has derived the observed  $g$ -values tabulated for both "even" and "odd" levels from the same observations and formulas, without assuming theoretical  $g$ -values, but by adopting the revisions in analysis suggested by Earls and Sawyer.

## REFERENCES

- J. B. Green and R. A. Loring, *Phys. Rev.* **42**, 459 (1933). (Z E)  
 J. L. Rose, *Phys. Rev.* **47**, 123 (1935). (hfs)  
 I. T. Earls and R. A. Sawyer, *Phys. Rev.* **47**, 115 (1935). (I P) (T) (C L) (Z E)  
 K. Murakawa, *J. Phys. Soc. Japan* **8**, 382 (1953). (I S) (hfs)

## Pb II

## Pb II

Config.	Desig.	$J$	Level	Interval	Obs. $g$	Config.	Desig.	$J$	Level	Interval	Obs. $g$
$6s \text{ } ^3\text{S} 6p$	$6p \text{ } ^3\text{P}^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	0 14081	14081	1.33	$6s \text{ } ^3\text{S} 6f$	$6f \text{ } ^3\text{F}^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	102868 102881	-13	1.14 0.83
$6s \text{ } 6p^2$	$6p^2 \text{ } ^3\text{P}$	$0\frac{1}{2}$ $1\frac{1}{2}$ $2\frac{1}{2}$	57911 66124 73905	8213 7781	1.64 1.52	$6s \text{ } ^3\text{S} 8d$	$8d \text{ } ^3\text{D}$	$1\frac{1}{2}$ $2\frac{1}{2}$	103394 104190	796	0.767 1.21
$6s \text{ } ^3\text{S} 7s$	$7s \text{ } ^3\text{S}$	$0\frac{1}{2}$	59448		2.01	$6s \text{ } ^3\text{S} 5g$	$5g \text{ } ^3\text{G}$	$3\frac{1}{2}, 4\frac{1}{2}$	103559		
$6s \text{ } ^3\text{S} 6d$	$6d \text{ } ^3\text{D}$	$2\frac{1}{2}$ $1\frac{1}{2}$	68964 69740	-776	1.29 0.81	$6s \text{ } 6p^2$	$6p^2 \text{ } ^3\text{S}$	$0\frac{1}{2}$	104297		
$6s \text{ } ^3\text{S} 7p$	$7p \text{ } ^3\text{P}^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	74459 77878	2813	0.60 1.33	$6s \text{ } ^3\text{S} 9p$	$9p \text{ } ^3\text{P}^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	104488 105080	598	
$6s \text{ } 6p^2$	$6p^2 \text{ } ^3\text{D}$	$1\frac{1}{2}$ $2\frac{1}{2}$	83083 88972	5889		$6s \text{ } ^3\text{S} 10s$	$10s \text{ } ^3\text{S}$	$0\frac{1}{2}$	107930		
$6s \text{ } 6p^2$	$6p^2 \text{ } ^3\text{P}$	$0\frac{1}{2}$ $1\frac{1}{2}$	88248 104669	16421		$6s \text{ } ^3\text{S} 7f$	$7f \text{ } ^3\text{F}^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	108589 108538	-9	
$6s \text{ } ^3\text{S} 8s$	$8s \text{ } ^3\text{S}$	$0\frac{1}{2}$	89180			$6s \text{ } ^3\text{S} 6g$	$6g \text{ } ^3\text{G}$	$3\frac{1}{2}, 4\frac{1}{2}$	108968		
$6s \text{ } ^3\text{S} 5f$	$5f \text{ } ^3\text{F}^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	92514 92589	-15	1.11 0.84	$6s \text{ } ^3\text{S} 9d$	$9d \text{ } ^3\text{D}$	$1\frac{1}{2}$ $2\frac{1}{2}$	109257 109336	79	
$6s \text{ } ^3\text{S} 7d$	$7d \text{ } ^3\text{D}$	$1\frac{1}{2}$ $2\frac{1}{2}$	94284 95304	1020	0.80 1.18	$6s \text{ } ^3\text{S} 10p$	$10p \text{ } ^3\text{P}^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	109500 109851	351	
$6s \text{ } ^3\text{S} 8p$	$8p \text{ } ^3\text{P}^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	96077 96238	1161		$6s \text{ } ^3\text{S} 11s$	$11s \text{ } ^3\text{S}$	$0\frac{1}{2}$	111574		
$6s \text{ } ^3\text{S} 9s$	$9s \text{ } ^3\text{S}$	$0\frac{1}{2}$	101346		1.96	$6s \text{ } ^3\text{S} 8f$	$8f \text{ } ^3\text{F}^o$	$3\frac{1}{2}$ $2\frac{1}{2}$	111940 111946	-6	
						$6s \text{ } ^3\text{S} 7g$	$7g \text{ } ^3\text{G}$	$3\frac{1}{2}, 4\frac{1}{2}$	112230		

Pb II—Continued

Pb II—Continued

Config.	Design.	<i>J</i>	Level	Interval	Obs. <i>g</i>	Config.	Design.	<i>J</i>	Level	Interval	Obs. <i>g</i>
6s(1S)10d	10d 1D	1½ 2½	112409 112467	58		6s(1S)12f	12f 1F°	3½ 2½	117590 117588		-2
6s(1S)11p	11p 1P°	0½ 1½	118578 118900	222		6s(1S)11g	11g 1G	3½, 4½	117600		
6s(1S)12s	12s 1S	0½	113912			6s(1S)14d	14d 1D	1½ 2½	117650 117666		16
6s(1S)9f	9f 1F°	3½ 2½	114145 114150	-5		6s(1S)13f	13f 1F°	3½ 2½	118181 118188		-1
6s(1S)8g	8g 1G	3½, 4½	114346			6s(1S)12g	12g 1G	3½, 4½	118183		
6s(1S)11d	11d 1D	1½ 2½	114466 114507	41		6s(1S)15d	15d 1D	1½ 2½	118223 118235		12
6s(1S)12p	12p 1P°	0½ 1½	114585 114735	150		6s(1S)14f	14f 1F°	3½ 2½	118588		
6s(1S)13s	13s 1S	0½	115496			6s(1S)13g	13g 1G	3½, 4½	118637		
6s(1S)10f	10f 1F°	3½ 2½	115653 115667	-4		6s(1S)16d	16d 1D	1½ 2½	118668 118679		11
6s(1S)9g	9g 1G	3½, 4½	115797			6s(1S)14g	14g 1G	3½, 4½	118996		
6s(1S)12d	12d 1D	1½ 2½	115885 115912	27		6s(1S)17d	17d 1D	1½ 2½	119022 119031		9
6s(1S)13p	13p 1P°	0½ 1½	115967 116078	105		6s(1S)15g	15g 1G	3½, 4½	119286		
6s(1S)14s	14s 1S	0½	116615			6s(1S)18d	18d 1D	1½ 2½	119315		
6s(1S)11f	11f 1F°	3½ 2½	116788 116790	-2		6s(1S)19d	19d 1D	1½ 2½	119548		
6s(1S)10g	10g 1G	3½, 4½	116833								
6s(1S)13d	13d 1D	1½ 2½	116899 116920	21		Pb III(1S <sub>0</sub> )	Limit		121343		
6s(1S)14p	14p 1P°	0½ 1½	116958								

January 1955.

Pb II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>2</sup> 3s <sup>2</sup> 3p <sup>2</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup> 4d <sup>10</sup> 5s <sup>2</sup> 5p <sup>2</sup> 5d <sup>10</sup> +	Observed Terms				
6s(1S)6p	6p 1P°				
6s 6p <sup>2</sup>	{	6p <sup>2</sup> 1S	6p <sup>2</sup> 1P	6p <sup>2</sup> 1D	
		6p <sup>2</sup> 3P			
		ns (n ≥ 7)	np (n ≥ 7)	nd (n ≥ 6)	nf (n ≥ 5)
6s(1S)ns		7 to 14s 1S	7 to 14p 1P°	6 to 19d 1D	5 to 14f 1F°
					5 to 15g 1G

\*For predicted terms in the spectra of the Tl I isoelectronic sequence, see Vol. III, Introduction.



## Pb III

(Hg I sequence; 80 electrons)

Z=82

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 \text{ } ^1S_0$  $6s^2 \text{ } ^1S_0$ ,  $257592 \pm 5 \text{ K}$ 

I. P. 31.93 volts

The configuration-interaction in the Pb III spectrum has been studied in great detail by Hume, Crawford, McLay, and Crooker. Wave functions of 52 levels have been determined in terms of the (*LS*) functions, and the configuration-perturbation parameters evaluated by fitting the calculated to the observed level values. Hume and Crawford give most of the details in their 1951 paper. The data in the table are, however, from an unpublished manuscript furnished by Crawford for inclusion here. The observed *g*-values are from Green and Loring.

The earlier papers by Stanley Smith, Crooker, and Rao and Narayan contain over 100 classified lines in the interval 995.75 Å to 5857.96 Å. About 400 lines are reported as classified, however.

The limit has been calculated from the  $6s \text{ } n\lambda$  sequence. The authors comment that the fine structure of the H-terms has not been resolved, but lines of G-H° multiplets are diffuse, indicating small intervals for the H-terms.

Both hyperfine structure and isotope shift affect the values of the observed energy levels. When an isotope shift is given for a level in the 1951 paper, the adopted value is the mean of the value for  $\text{Pb}^{208}$  and  $\text{Pb}^{209}$ .

In the table the levels are listed in order of increasing values. No attempt has been made to group them into their respective terms, because in many cases more than one designation is involved for a given level. On account of perturbations, *LS*-coupling notation does not apply throughout. For a number of levels the dominant contributor as given by Hume and Crawford is entered in the "Desig." column. The writer has made tentative designation assignments for  $6p'' \text{ } ^3F_2$ ,  $5f \text{ } ^1F_2$ ,  $6p'' \text{ } ^3D_2$ ,  $5,6,7,8g \text{ } ^1G_4$ ,  $6,7,8,9h \text{ } ^1H^{\circ}$ , and  $7s' \text{ } ^1P^{\circ}$ . The authors give configurations for these levels, but have not calculated the wave functions.

## REFERENCES

- Stanley Smith, Proc. Nat. Acad. Sci. 14, 878 (1928). (C L)  
 Stanley Smith, Phys. Rev. 30, 1 (1930). (T) (C L)  
 A. S. Rao and A. L. Narayan, Zeit. Phys. 59, 687 (1930). (T) (C L)  
 J. B. Green and R. A. Loring, Phys. Rev. 43, 459 (1933). (Z E) (C L)  
 A. M. Crooker, Canadian J. Research A 14, 115 (1936). (I P) (T) (C L) (hfs) (I S)  
 J. N. P. Hume and M. F. Crawford, Phys. Rev. 84, 496 (1951) and unpublished material (1951). (I P) (T) (hfs) (I S)  
 H. E. Walchli, A Table of Nuclear Moment Data, Oak Ridge Nat. Lab., ORNL-1469, 62 (1953). (Summary hfs)

## Pb III

## Pb III

Authors	Config.	Desig.	J	Level	Obs. <i>g</i>	Authors	Config.	Desig.	J	Level	Obs. <i>g</i>
1 <sub>0</sub>	$5d^{10} 6s^2$	$6s^2 \text{ } ^1S$	0	0		2 <sub>1</sub>	$5d^{10} 6s(6S)7s$	$7s \text{ } ^3S$	1	150063.7	1.98
13	$5d^{10} 6s(6S)6p$	$6p \text{ } ^3P^{\circ}$	0	60397		4 <sub>1</sub>	$5d^{10} 6s(6S)6d$	$6d \text{ } ^1D$	2	151894.5	1.06
21	$5d^{10} 6s(6S)6p$	$6p \text{ } ^3P^{\circ}$	1	64391		5 <sub>0</sub>	$5d^{10} 6s(6S)7s$	$7s \text{ } ^1S$	0	153783.4	
31	$5d^{10} 6s(6S)6p$	$6p \text{ } ^3P^{\circ}$	2	78984.6		5 <sub>2</sub>	$5d^{10} 6s(6D)6p$	$6p'' \text{ } ^3P^{\circ}$	2	154494.0	
41	$5d^{10} 6s(6S)6p$	$6p \text{ } ^1P^{\circ}$	1	95340.1		6 <sub>1</sub>	$5d^{10} 6p^3$	$6p^3 \text{ } ^3P$	1	155431.5	
2 <sub>0</sub>	$5d^{10} 6p^2$	$6p^2 \text{ } ^3P$	0	142551		7 <sub>1</sub>	$5d^{10} 6s(6S)6d$	$6d \text{ } ^3D$	1	157444.1	0.50

## Pb III—Continued

## Pb III—Continued

Authors	Config.	Desig.	J	Level	Obs. g	Authors	Config.	Desig.	J	Level	Obs. g
8 <sub>2</sub>	5d <sup>10</sup> 6s(°S)6d	6d °D	2	157925. 0	1. 15	20 <sub>2</sub>	5d <sup>10</sup> 6s(°S)5g	5g °G	3	217635. 3	0. 8
9 <sub>2</sub>	5d <sup>10</sup> 6s(°S)6d	6d °D	3	158956. 8	1. 33	21 <sub>2</sub>	5d <sup>10</sup> 6s(°S)5g	5g °G	5	217659. 1	
10 <sub>2</sub>	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> °P	2	164817. 9	1. 40	22 <sub>2</sub>	5d <sup>10</sup> 6s(°S)5g	5g °G	4	217660. 6	1. 06
7 <sub>1</sub>	5d <sup>10</sup> 6s(°S)7p	7p °P°	0	170917. 3		23 <sub>1</sub>	5d <sup>10</sup> 6s(°S)9s	9s °S	1	219344	
8 <sub>1</sub>	5d <sup>10</sup> 6s(°S)7p	7p °P°	1	171081. 4	1. 38	24 <sub>2</sub>	5d <sup>10</sup> 6s(°S)9s	9s °S	0	219910	
9 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °F°	4	173329		25 <sub>1</sub>	5d <sup>10</sup> 6s(°S)8d	8d °D	1	221205	
10 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °F°	2	173986. 3	0. 97	26 <sub>2</sub>	5d <sup>10</sup> 6s(°S)8d	8d °D	2	221307	
11 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °P°	1	174600. 9	1. 14	27 <sub>2</sub>	5d <sup>10</sup> 6s(°S)8d	8d °D	3	221600	
12 <sub>1</sub>	5d <sup>10</sup> 6s(°S)7p	7p °P°	2	176022. 9	1. 34	28 <sub>2</sub>	5d <sup>10</sup> 6s(°S)8d	8d °D	2	221938	
13 <sub>1</sub>	5d <sup>10</sup> 6s(°S)7p	7p °P°	1	177181. 4	1. 15	33 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )7s	7s' °P°	1	224740	0
14 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °D°	2	177906. 5	1. 08	34 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )7s	7s' °P°	0	226818	
11 <sub>2</sub>	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> °D	2	178432		35 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )6d		1	227584	
15 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °F°	3	178957. 0		36 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )6d		2	227807	
16 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °P°	1	184268. 1	0. 96	37 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )6d		3	228039	
12 <sub>2</sub>	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> °S	0	188615		29 <sub>2</sub>	5d <sup>10</sup> 6s(°S)6g	6g °G	4	229845	
17 <sub>1</sub>	5d <sup>10</sup> 6s(°S)5f	5f °F°	3	189785. 2	1. 09	30 <sub>2</sub>	5d <sup>10</sup> 6s(°S)6g	6g °G	3	229846	
18 <sub>1</sub>	5d <sup>10</sup> 6s(°S)5f	5f °F°	2	190287. 8	0. 67	31 <sub>2</sub>	5d <sup>10</sup> 6s(°S)6g	6g °G	5	229870. 5	
19 <sub>1</sub>	5d <sup>10</sup> 6s(°S)5f	5f °F°	4	190429. 0	1. 25	32 <sub>2</sub>	5d <sup>10</sup> 6s(°S)6g	6g °G	4	229870	
20 <sub>1</sub>	5d <sup>10</sup> 6s(°S)5f	5f °F°	3	190901. 2	1. 00	38°	5d <sup>10</sup> 6s(°S)6h	6h °H°		230085	
21 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °P°	0	192880		33 <sub>1</sub>	5d <sup>10</sup> 6s(°S)10s	10s °S	1	230954	
22 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °D°	3	196846. 7	1. 03	34 <sub>2</sub>	5d <sup>10</sup> 6s(°S)10s	10s °S	0	231244	
23 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °D°	1	197318. 6		35 <sub>1</sub>	5d <sup>10</sup> 6s(°S)9d	9d °D	1	232019	
13 <sub>2</sub>	5d <sup>10</sup> 6s(°S)8s	8s °S	1	197892. 8	2. 00	36 <sub>2</sub>	5d <sup>10</sup> 6s(°S)9d	9d °D	2	232097	
24 <sub>1</sub>	5d <sup>10</sup> 6s(°D)6p	6p'' °D°	2	199343. 9		37 <sub>2</sub>	5d <sup>10</sup> 6s(°S)9d	9d °D	3	232274	
14 <sub>2</sub>	5d <sup>10</sup> 6s(°S)8s	8s °S	0	199400. 6		38 <sub>2</sub>	5d <sup>10</sup> 6s(°S)9d	9d °D	2	232442	
15 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7d	7d °D	1	201398. 7	0. 50	39 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )6d		2	233680	
16 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7d	7d °D	2	201597. 3	1. 16	39 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7g	7g °G	4	237212	
17 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7d	7d °D	3	202046. 8	1. 33	40 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7g	7g °G	3	237219	
18 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7d	7d °D	2	203301. 6	0. 98	42 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7g	7g °G	4	237239	
25 <sub>1</sub>	5d <sup>10</sup> 6s(°S)8p	8p °P°	0	206809		41 <sub>2</sub>	5d <sup>10</sup> 6s(°S)7g	7g °G	5	237240	
26 <sub>1</sub>	5d <sup>10</sup> 6s(°S)8p	8p °P°	1	206979		40°	5d <sup>10</sup> 6s(°S)7h	7h °H°		237389	
27 <sub>1</sub>	5d <sup>10</sup> 6s(°S)8p	8p °P°	2	208922		43 <sub>2</sub>	5d <sup>10</sup> 6s(°S)8g	8g °G		242005	
28 <sub>1</sub>	5d <sup>10</sup> 6s(°S)8p	8p °P°	1	209318		44 <sub>2</sub>	5d <sup>10</sup> 6s(°S)8g	8g °G		242023	
29 <sub>1</sub>	5d <sup>10</sup> 6s(°S)6f	6f °F°	2	214434		41°	5d <sup>10</sup> 6s(°S)8h	8h °H°		242130	
30 <sub>1</sub>	5d <sup>10</sup> 6s(°S)6f	6f °F°	3	214477		42 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )7s	7s' °P°	1	242321	
31 <sub>1</sub>	5d <sup>10</sup> 6s(°S)6f	6f °F°	4	214486		43°	5d <sup>10</sup> 6s(°S)9h	9h °H°		245378	
32 <sub>1</sub>	5d <sup>10</sup> 6s(°S)6f	6f °F°	3	214846		44 <sub>1</sub>	5d <sup>10</sup> 6p(°P <sub>1/2</sub> )7s	7s' °P°	2	246144	
19 <sub>2</sub>	5d <sup>10</sup> 6s(°S)5g	5g °G	4	217633. 4	1. 01		Pb IV(°S <sub>01</sub> )	Limit	---	257592 ± 5	

## Pb III Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>6</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> +	Observed Terms		
	5d <sup>9</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>1</sup> S	
5d <sup>9</sup> 6p <sup>1</sup>	6p <sup>1</sup> <sup>1</sup> S      6p <sup>1</sup> <sup>3</sup> P      6p <sup>1</sup> <sup>1</sup> D		
	ns (n ≥ 6)	np (n ≥ 6)	nd (n ≥ 6)
5d <sup>9</sup> 6s( <sup>3</sup> S)ns	{ 7 to 10s <sup>3</sup> S 7 to 10s <sup>1</sup> S	6 to 8p <sup>3</sup> P° 6 to 8p <sup>1</sup> P°	6 to 8d <sup>3</sup> D 6 to 8d <sup>1</sup> D
5d <sup>9</sup> 6p( <sup>3</sup> P°)ns'	{ 7s' <sup>3</sup> P° 7s' <sup>1</sup> P°		
5d <sup>9</sup> 6s( <sup>1</sup> D)ns''	{	6p'' <sup>3</sup> P°      6p'' <sup>3</sup> D°      6p'' <sup>3</sup> F° 6p'' <sup>1</sup> P°      6p'' <sup>1</sup> D°	
	nf (n ≥ 5)	ng (n ≥ 5)	nħ (n ≥ 6)
5d <sup>9</sup> 6s( <sup>3</sup> S)ns	5, 6f <sup>3</sup> F° 5, 6f <sup>1</sup> F°	5 to 8g <sup>3</sup> G 5 to 8g <sup>1</sup> G	6 to 8ħ <sup>3</sup> H° 6 to 8ħ <sup>1</sup> H°

\*For predicted terms of the Hg I isoelectronic sequence, see Vol. III, Introduction.

## Pb IV

(Au I sequence; 79 electrons)

Z=82

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>10</sup> 6s <sup>3</sup>S<sub>1/2</sub>

6s <sup>3</sup>S<sub>1/2</sub> 341350 K

I. P. 42.81 volts

Further study of the spectrum is needed. The early work on this spectrum was extended by Schoepfle in 1935 to include 98 classified lines between 459.04 Å and 4496.21 Å. Later, Crawford, McLay, and Crooker revised and extended the analysis and published 134 classified lines in the interval 432.880 Å to 8749.7 Å. The data in the table are from this later paper, and the numbers used by these authors to designate miscellaneous levels are retained in column one. The wavelengths of Arvidsson have been used in both papers for the region 432 Å to 1438 Å. The terms 6, 7, 8s <sup>3</sup>S, 6p <sup>3</sup>P°, and 6,7d <sup>3</sup>D, and 6 miscellaneous levels are common to both papers, but Schoepfle lists 19 levels not retained by Crawford and his associates.

By applying a Hicks formula to the ns <sup>3</sup>S series (ns=6 to 8), Schoepfle derives the limit 340885, which gives an ionization potential of 42.25 volts. The limit quoted above "was obtained by assuming the sequence nħ <sup>3</sup>H° to be hydrogenic".

This spectrum exhibits evidence of *jj*-coupling in the levels from the 5d<sup>9</sup> 6s(<sup>1</sup>D)6p configuration. The notation for this coupling is discussed in the text for Au I. The authors assign the levels 1° through 4° to the configuration 5d<sup>9</sup><sub>2,4</sub> 6s<sub>0,1</sub> 6p<sub>0,1</sub>. By comparison with isoelectronic spectra, Trees and the writer have made the additional *jj*-coupling configuration assignments given in the table for this group; they are extremely tentative. Schoepfle suggests that the levels labeled in the table 16° and 22°, may be 7p <sup>3</sup>P°, and that 24° may be 5f <sup>3</sup>F°. The observed *g*-values are consistent with this 7p <sup>3</sup>P° assignment, but the overlapping of configurations makes a definitive choice difficult.

Pb IV—Continued

REFERENCES

J. B. Green and R. A. Loring, *Phys. Rev.* **43**, 440 (1933). (S E)  
 G. K. Schoopfe, *Phys. Rev.* **47**, 333 (1935). (I P) (T) (C L)  
 M. F. Crawford, A. B. McLay, and A. M. Crocker, *Proc. Roy. Soc. London [A]* **158**, No. 894, 455 (1937). (I P)  
 (T) (C L) (I S) (hfs)  
 A. L. Schawlow, J. N. P. Hume, and M. F. Crawford, *Phys. Rev.* **76**, 1877 (L) (1949). (hfs)

Pb IV

Pb IV

Authors	Config.	Desig.	J	Level	Interval	Obs. #	Authors	Config.	Desig.	J	Level	Interval	Obs. #
	5d <sup>10</sup> (S)6s	6s <sup>2</sup> S	0½	0			17°	5d <sup>10</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		0½	210369. 7		
	5d <sup>10</sup> (S)6p	6p <sup>2</sup> P°	0½ 1½	76158 97319	21061		18°	5d <sup>10</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>3/2</sub>		2½	213519. 3		
1	5d <sup>9</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>2</sup> D	2½ 1½	101252 122568	-21316		19°	5d <sup>10</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		1½	214842. 1		
2	5d <sup>9</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		2½	166369			20°			3½	214891. 8		
1°	"		3½	172667			21°	5d <sup>9</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		2½	217216		
2°	"		2½	173248			22°	" ?		1½	217851. 9		1. 29
3°	"		1½	175388			23°	"		2½	219461. 0		
4°	"		1½	184558. 8 186816. 8	2258. 0	0. 78 1. 17	24°	"		3½	221716. 1		
	5d <sup>10</sup> (S)6d	6d <sup>2</sup> D	1½ 2½	185103. 0		1. 92	25°			0½	231013		
	5d <sup>10</sup> (S)7s	7s <sup>2</sup> S	0½	188759			26°			2½	232633		
5°	5d <sup>9</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		1½	193776			27°			1½	235566		
7°	"		2½	193855				5d <sup>10</sup> (S)8s	8s <sup>2</sup> S	0½	249634. 5		
8°	"		3½	193954				5d <sup>10</sup> (S)7d	7d <sup>2</sup> D	1½ 2½	250401. 6 251419. 5	1017. 9	
9°	"		1½	194147				5d <sup>10</sup> (S)5g	5g <sup>2</sup> G	4½ 3½	270496 270498	-2	
10°	"		0½	197024				5d <sup>10</sup> (S)6h	6h <sup>2</sup> H°	4½, 5½	302543		
11°	"		2½	200081				5d <sup>10</sup> (S)7h	7h <sup>2</sup> H°	4½, 5½	305516		
12°	5d <sup>9</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		1½	201480				5d <sup>10</sup> (S)8h	8h <sup>2</sup> H°	4½, 5½	313939		
13°	"		0½	208584. 0									
14°	5d <sup>9</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		3½	209051. 1									
15°	5d <sup>9</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub>		1½	209722. 4		0. 68							
16°	5d <sup>9</sup> <sub>1/2</sub> 6s <sub>1/2</sub> 6p <sub>1/2</sub> ?		0½					Pb v(S <sub>2</sub> )	Limit		341350		

September 1954.

## Pb v

(Pt I sequence; 78 electrons)

Z=82

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} {}^1S_0$  $5d^{10} {}^1S_0$ , 555000 K

I. P. 69.8 volts

The analysis is from Schoepfle, who has extended the early work of Mack and the subsequent investigation by Mack and Fromer. The levels have been based on the observations by Arvidsson, who separated the lines belonging to the spectra of different stages of ionization. About 200 lines have been classified in the interval 284.329 Å to 1401.56 Å.

The spectra of the Pt I isoelectronic sequence are of special theoretical interest because they approach *jj*-coupling rather than *LS*- or *JL*-coupling. This appears more conspicuously in the spectra of higher ionization, especially among the levels from the  $5d^9({}^2D)6p$  configuration. A detailed study of "The Four Vector Problem and Its Application to Energies and Intensities in Platinum-like Spectra" has been made by Goble. In *LS*-coupling this configuration gives rise to the terms  ${}^1,3(P^{\circ}D^{\circ}F^{\circ})$ . In *jj*-coupling the same number of energy levels and the same resultant *J*-values occur, but each level is defined by two *j*-values, one for the *d*-electron ( $1\frac{1}{2}$  or  $2\frac{1}{2}$ ), and one for the *p*-electron, ( $0\frac{1}{2}$  or  $1\frac{1}{2}$ ).

In the table the writer has assigned *LS*-designations only in the case of the  ${}^1,3D$  terms from the  $5d^9({}^2D)ns$  configuration, although the *jj*-coupling applies here, also. "Odd" levels are in italics throughout. The *j*-values for the *d*- and *p*-electrons as required for *jj*-coupling, are indicated in the configuration column, the former being the same as the *J*-values of the limit term  ${}^2D$ . A table giving the transformation from *jj*- to *LS*-coupling may be found in the book by Condon and Shortley, p. 294.

Schoepfle uses as the ionization limit the value 562600 derived from a Ritz formula, from which he apparently derived the absolute value of the level  $5d^9({}^2D_{3/2})6p_{1/2}$ , *J*=3. The corresponding ionization potential is 69.7 volts. Mack and Fromer quote an ionization potential derived from a Moseley diagram for the sequence. This estimate is based on a two-member series,  $5d^9({}^2D_{3/2})ns$  (*n*=6,7), *J*=3, represented by a Rydberg formula to which the same Ritz correction is applied as for the spectrum of the same stage of ionization in the Au I sequence. Their limit is quoted here.

## REFERENCES

- G. Arvidsson, *Ann. der Phys.* [5] **12**, 787 (1932).  
 A. T. Goble, *Phys. Rev.* **48**, 346 (1935).  
 J. E. Mack and M. Fromer, *Phys. Rev.* **48**, 357 (1935). (I P) (T) (C L)  
 G. K. Schoepfle, *Phys. Rev.* **50**, 538 (1936). (I P) (T) (C L)  
 E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra*, p. 294 (Cambridge University Press, London, 1951).

Pb v

Pb v

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
5d <sup>10</sup>	5d <sup>10</sup> 1S	0	0				3	326888	
5d <sup>9</sup> (D <sub>3/2</sub> )6s	6s 1D	3	110768	-3937 -18006			4	327438	
		2	114705				3	328284	
5d <sup>9</sup> (D <sub>5/2</sub> )6s		1	132711				3	329672	
5d <sup>9</sup> (D <sub>3/2</sub> )6s	6s 1D	2	135997			3	329935		
		2	141988			3	331487		
5d <sup>9</sup> (D <sub>5/2</sub> )6p <sub>3/2</sub>		2	194808			2	331749		
5d <sup>9</sup> (D <sub>3/2</sub> )6p <sub>3/2</sub>		3	197131			2	332358		
		3	199048			2	332501		
		3	210514			2	335013		
		2	211487			1	335013		
		3	212878		5d <sup>9</sup> (D <sub>3/2</sub> )7s	7s 1D	3	338440	-794
		2	213131			2	339234		
		2	215804			2	341595		
5d <sup>9</sup> (D <sub>5/2</sub> )6p <sub>3/2</sub>		2	217069			2	350805		
		3	218043			3	358771		
5d <sup>9</sup> (D <sub>3/2</sub> )6p <sub>3/2</sub>		1	219488			2	359988		
5d <sup>9</sup> (D <sub>5/2</sub> )6p <sub>1/2</sub>		4	221069		5d <sup>9</sup> (D <sub>5/2</sub> )7s	7s 1D	1	360496	
5d <sup>9</sup> (D <sub>3/2</sub> )6p <sub>1/2</sub>		2	223909		5d <sup>9</sup> (D <sub>3/2</sub> )7s	7s 1D	2	361073	
5d <sup>9</sup> (D <sub>5/2</sub> )6p <sub>1/2</sub>		3	226511		5d <sup>9</sup> (D)7p?		2	397410	
5d <sup>9</sup> (D <sub>3/2</sub> )6p <sub>1/2</sub>		2	227748		5d <sup>9</sup> (D)7p?		3	427086	
5d <sup>9</sup> (D <sub>5/2</sub> )6p <sub>1/2</sub>		1	227835		5d <sup>9</sup> (D)7p?		2	427580	
		2	232040			2	450173		
		2	233830		5d <sup>9</sup> (D <sub>3/2</sub> )8s	8s 1D	3	452624	-343
		0	237680			2	452967		
		1	239418			3	473642		
5d <sup>9</sup> (D <sub>3/2</sub> )6p <sub>1/2</sub>		3	244681		5d <sup>9</sup> (D <sub>5/2</sub> )8s	8s 1D	1	476582	
5d <sup>9</sup> (D <sub>5/2</sub> )6p <sub>1/2</sub>		1	245877		5d <sup>9</sup> (D <sub>3/2</sub> )8s	8s 1D	2	476894	
5d <sup>9</sup> (D <sub>3/2</sub> )6p <sub>1/2</sub>		2	247595		5d <sup>9</sup> (D <sub>5/2</sub> )9s	9s 1D	3	546508	
5d <sup>9</sup> (D <sub>5/2</sub> )6p <sub>1/2</sub>		2	253588			2	554440		
		2	253588		5d <sup>9</sup> (D)8p?		3	554440	
		1	260674		Pb vi(D <sub>3/2</sub> )	Limit		555000	
		3	283483		5d <sup>9</sup> (D)8p?		3	571267	
		3	297112		5d <sup>9</sup> (D)8p?		2	575919	
		2	299708		5d <sup>9</sup> (D)8p?		3	577526	
		3	317397		5d <sup>9</sup> (D)8p?		2	604580	
		2	326216						

August 1954.

## BISMUTH

Bi I

83 electrons

Z=83

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^3 \text{ } ^3\text{S}_{1/2}$  $6p^3 \text{ } ^3\text{S}_{1/2}$  58790 K

I. P. 7.287 volts

Although the Bi I spectrum has been studied by a number of investigators, yet no complete homogeneous set of observations exists. The spectrum needs to be reobserved and accurately measured over the entire photographic range.

Series were recognized by Thorsen in 1926. They were extended by Clearman in 1952 from observations in the short-wave region; his revised values of the limits are quoted here. Mrozowski has studied the hyperfine structure in detail and also extended the analysis.

The present list of energy levels is based mostly upon the terms given by Mrozowski and Clearman with revisions by the writer. She has prepared a line list from unpublished measurements by Meggers and Murphy, by Humphreys and Paul (infrared), and from Clearman's paper. She has recalculated the level values from these wavelengths and made additional tentative assignments of configurations for many miscellaneous levels. A number of published levels are omitted here, pending further confirmation. Some new levels have been added on evidence furnished by the series assignments. In general, however, further confirmation of many  $J$ -values, configurations, and levels is needed.

There are approximately 180 classified lines between 1362.84 Å and 22551.6 Å.

Most of the levels are listed in increasing numerical order in the table, because of the departure from  $LS$ -coupling. Some  $LS$ -designations are, however, indicated on the evidence afforded by the observed  $g$ -values.

## REFERENCES

- P. A. van der Harst, Proc. K. Akad. Wetensch. Amsterdam **22**, Part 1, 311 (1920). (Z E)  
 V. Thorsen, Zeit. Phys. **40**, 642 (1926). (T) (C L)  
 P. Zeeman, E. Beck, und S. Goudsmit, Zeit. Phys. **66**, 1 (1930). (Z E) (hfs)  
 S. Mrozowski, Phys. Rev. **63**, 526 (1942); **66**, 169 (1946). (T) (C L) (hfs)  
 H. E. Clearman, J. Opt. Soc. Am. **42**, 376 (1952). (I P) (T) (C L)  
 W. F. Meggers and R. J. Murphy, unpublished material (1953). (C L)  
 H. E. Walchli, A Table of Nuclear Moment Data, Oak Ridge Nat. Lab., ORNL-1469, 63 (1953). (Summary hfs)  
 C. J. Humphreys and E. Paul, Jr., U. S. Naval Ord. Lab., NAVORD Report 4589, 25 (1956). (C L)

Bi I

Bi I

Config.	Design.	J	Level	Interval	Obs. g	Config.	Design.	J	Level	Interval	Obs. g
6p <sup>2</sup>	6p <sup>2</sup> 6°	1½	0.00			6p <sup>2</sup> (P <sub>1</sub> )8s	8s °P?	0½	61133?		
6p <sup>2</sup>	6p <sup>2</sup> 1D°	1½ 2½	11418.09 15457.66	4018.63	1.225 1.20	6p <sup>2</sup> (P <sub>1</sub> )6d		0½?	61495?		
6p <sup>2</sup>	6p <sup>2</sup> 1P°	0½ 1½	21661.0 33164.84	11503.8	0.667	6p <sup>2</sup> (P <sub>1</sub> )6d		1½	61812		
6p <sup>2</sup> (P <sub>1</sub> )7s	7s °P	0½	32588.17		2.088	6p <sup>2</sup> (P <sub>1</sub> )6d		1½	62744		
6p <sup>2</sup> (P <sub>1</sub> )7p		0½?	41185.18			6p <sup>2</sup> (P <sub>1</sub> )8p		0½?	62870.87		
6p <sup>2</sup> (P <sub>1</sub> )7p		1½	48940.6			6p <sup>2</sup> (P <sub>1</sub> )8p		1½?	63200.6		
6p <sup>2</sup> (P <sub>1</sub> )6d	6d 1D	1½	43912.67		0.79	6p <sup>2</sup> (P <sub>1</sub> )8p		0½?	63733.87		
6p <sup>2</sup> (P <sub>1</sub> )6d	6d 1D	2½	44816.86		1.20	6p <sup>2</sup> (P <sub>1</sub> )8s	8s °P	2½	63858		
6p <sup>2</sup> (P <sub>1</sub> )7s	7s °P	1½	44865.06		1.676	6p <sup>2</sup> (P <sub>1</sub> )8s	8s °P?	1½	64023		
6p <sup>2</sup> (P <sub>1</sub> )7s	7s °P?	0½	45915.57		1.30	6p <sup>2</sup> (P <sub>1</sub> )7d?		2½	64122		
6p <sup>2</sup> (P <sub>1</sub> )8s	8s °P	0½	47373.23			6p <sup>2</sup> (P <sub>1</sub> )7d		0½	64238		
6p <sup>2</sup> (P <sub>1</sub> )7s	7s °P	2½	48489.88		1.41	6p <sup>2</sup> (P <sub>1</sub> )7d		1½	64275		
6p <sup>2</sup> (P <sub>1</sub> )7s	7s °P?	1½	49456.6		0.98	6p <sup>2</sup> (P <sub>1</sub> )7d		2½	64575?		
6p <sup>2</sup> (P <sub>1</sub> )8p		0½?	49997.48?			6p <sup>2</sup> (P <sub>1</sub> )9s	9s °P	1½	65483		
6p <sup>2</sup> (P <sub>1</sub> )7d	7d 1D	1½	51019.11			6p <sup>2</sup> (P <sub>1</sub> )9s	9s °P?	0½	65612		
6p <sup>2</sup> (P <sub>1</sub> )7d	7d 1D	2½	51158.65			6p <sup>2</sup> (P <sub>1</sub> )9p		0½?	66584.9?		
6p <sup>2</sup> (P <sub>1</sub> )9s	9s °P	0½	52255.36			6p <sup>2</sup> (P <sub>1</sub> )9p		0½?	66992.1?		
6p <sup>2</sup> (P <sub>1</sub> )9p		0½?	53471.?			6p <sup>2</sup> (D <sub>1</sub> )7s		1½?	67117		
6p <sup>2</sup> (P <sub>1</sub> )8d	8d 1D	2½	53878.6			6p <sup>2</sup> (P <sub>1</sub> )8d		0½	67216?		
6p <sup>2</sup> (P <sub>1</sub> )7p		0½	55893.74			6p <sup>2</sup> (P <sub>1</sub> )8d?		1½?	67328		
6p <sup>2</sup> (P <sub>1</sub> )8d	8d 1D	1½	53976.84			6p <sup>2</sup> (P <sub>1</sub> )10s?	10s °P?	1½	67832		
6p <sup>2</sup> (P <sub>1</sub> )10s	10s °P	0½	54559.7			6p <sup>2</sup> (P <sub>1</sub> )7d		0½?	68049?		
6p <sup>2</sup> (P <sub>1</sub> )7p		1½	54570.53			6p <sup>2</sup> (D <sub>1</sub> )7s		2½?	68086		
6p <sup>2</sup> (P <sub>1</sub> )9d	9d 1D	1½	55424.3			6p <sup>2</sup> (P <sub>1</sub> )10p		0½?	68365.5?		
6p <sup>2</sup> (P <sub>1</sub> )9d	9d 1D	2½	55478.0			6p <sup>2</sup> (P <sub>1</sub> )9d		0½	68764		
6p <sup>2</sup> (P <sub>1</sub> )11s	11s °P	0½	55822.2			6p <sup>2</sup> (P <sub>1</sub> )9s	9s °P	2½	68967?		
6p <sup>2</sup> (P <sub>1</sub> )7p		1½?	56088.3			6p <sup>2</sup> (P <sub>1</sub> )9s	9s °P	1½	69120?		
6p <sup>2</sup> (P <sub>1</sub> )7p		0½?	56275.4			6p <sup>2</sup> (P <sub>1</sub> )8d		0½?	70931?		
6p <sup>2</sup> (P <sub>1</sub> )10d	10d 1D	1½	56320			6p <sup>2</sup> (P <sub>1</sub> )10s	10s °P	1½	71499?		
6p <sup>2</sup> (P <sub>1</sub> )6d		1½	56570.50			-----					
6p <sup>2</sup> (P <sub>1</sub> )12s	12s °P	0½	56591?			Bi II(P <sub>1</sub> )	Limit		72112		
6p <sup>2</sup> (P <sub>1</sub> )6d		2½	57076			-----					
6p <sup>2</sup> (P <sub>1</sub> )6d		0½	57607.0			6p <sup>2</sup> (P <sub>1</sub> )9d		0½?	72463?		
6p <sup>2</sup> (P <sub>1</sub> )6d		1½	57767.8			6p <sup>2</sup> (P <sub>1</sub> )10d		0½?	73376?		
6p <sup>2</sup> (P <sub>1</sub> )6d		2½	58273.64			-----					
6p <sup>2</sup> (P <sub>1</sub> )6d		3½	58377.6			Bi II(P <sub>2</sub> )	Limit		75815		
Bi II(P <sub>1</sub> )	Limit		58790			-----					
6p <sup>2</sup> (P <sub>1</sub> )8s	8s °P	1½	60817			Bi II(D <sub>1</sub> )	Limit		92720		



## Bi II

(Pb I sequence; 82 electrons)

Z=83

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^2 \ ^3P_0$ . $6p^2 \ ^3P_0$ , 134600 K

I. P. 16.68 volts

The energy levels and limit are from Crawford and McLay, the authors who assigned the arbitrary numbers listed in the first column of the table. In 1947, Murakawa and Suwa revised and extended the *jj*-coupling assignments of these authors and also suggested *LS*-designations for nearly all of the levels. Their results are quoted in the second and third columns of the table.

The level at 112645 K is from this paper, and the levels labeled 25<sub>2</sub> and 21<sub>1</sub> are found only in the earlier one. The level at 117004 K is listed as both "even" ( $6p_{3/2} 6f_{5/2} \ ^3F_2$ ) (1947) and "odd" ( $6p_{3/2} 5g$ ) (1934). On the basis of the combinations in the 1934 paper, the latter configuration has been adopted in the table. This level has been "assumed hydrogenic" for the calculation of the limit.

There are approximately 190 lines in the interval 937.95 Å to 8863 Å, including some "forbidden" transitions. The hyperfine structure exhibited by a number of the lines is discussed in detail by Fisher and Goudsmit and others. The over-all separation is wide for a number of the energy levels.

## REFERENCES

- R. A. Fisher and S. Goudsmit, *Phys. Rev.* **37**, 1057 (1931). (hfs)  
 M. F. Crawford and A. B. McLay, *Proc. Roy. Soc. [A]* **143**, 540 (1934). (I P) (T) (C L) (hfs)  
 K. Murakawa and S. Suwa, *Reports Inst. Sci. Tech. Tokyo Univ.* **1**, 121 (1947). (I P) (T) (hfs)  
 J. E. Mack, *Rev. Mod. Phys.* **32**, 64 (1950). (Summary hfs)  
 C. D. Cole and S. Mrozowski, *Bull. Am. Phys. Soc.* **38**, 29 (1953). (C L)

## Bi II

## Bi II

Authors	Config.	Desig.	J	Level	Interval	Authors	Config.	Desig.	J	Level	Interval	
1 <sub>0</sub>	$6p_{3/2} 6p_{3/2}$	$6p^2 \ ^3P$	0	0	13324 3706	10f	$6p_{1/2} 7s$	$^1P^o$	1	89883		
2 <sub>1</sub>	$6p_{3/2} 6p_{1/2}$		1	13324			11f	6s	$^3D^o$	1	94440	
3 <sub>2</sub>	$6p_{3/2} 6p_{1/2}$		2	17030			12f	6s $6p^o$	$^3D^o$	2	94928	
4 <sub>2</sub>	$6p_{1/2} 6p_{1/2}$	$6p^2 \ ^1D$	2	33936		13f	6s $6p^3$	$^3D^o$	3	96057		
5 <sub>0</sub>	$6p_{1/2} 6p_{1/2}$	$6p^2 \ ^3S$	0	44173		14f	$6p_{1/2} 6d$	$^3P^o$	2	99011		
14	$6p_{3/2} 7s$	$^1P^o$	0	89133		15f	$6p_{1/2} 6d$	$^3P^o$	1	100492		
21	$6p_{3/2} 7s$	$^3P^o$	1	69598		16f	$6p_{3/2} 8s$	$^3P^o$	1	101341		
43	6s $6p^3$	$6p^2 \ ^3S^o$	2	76147		17 <sub>0</sub>	$6p_{1/2} 6d$	$^3P^o$	0	101464		
53	$6p_{3/2} 6d_{3/2}$	$^3D^o$	2	79089		18f	$6p_{1/2} 6d$	$^1D^o$	2	102330		
6f	$6p_{3/2} 6d_{3/2}$	$^3D^o$	1	80575		19f	$6p_{1/2} 6d$	$^3D^o$	3	103003		
7f	$6p_{3/2} 6d_{3/2}$	$^3F^o$	2	82047		10 <sub>2</sub>	$6p_{3/2} 5f_{3/2}$	$^3F$	2	105083		
8f	$6p_{3/2} 6d_{3/2}$	$^3F^o$	3	82255		11 <sub>2</sub>	$6p_{1/2} 7p_{3/2}$	$^3P$	2	105269		
6 <sub>1</sub>	$6p_{3/2} 7p_{3/2}$	$^3D$	1	84281		12 <sub>2</sub>	$6p_{3/2} 5f$	$^3G$	3	105287		
7 <sub>0</sub>	$6p_{3/2} 7p_{3/2}$	$^3P$	0	87078		13 <sub>2</sub>	$6p_{3/2} 5f$	$^3F$	3	105449		
8 <sub>1</sub>	$6p_{3/2} 7p_{1/2}$	$^3P$	1	83566		20f	$6p_{3/2} 7d$	$^3D^o$	2	105526		
9f	$6p_{1/2} 7s$	$^3P^o$	2	83769		14 <sub>1</sub>	$6p_{3/2} 5f_{3/2}$	$^3F$	4	105726		
9 <sub>2</sub>	$6p_{3/2} 7p_{1/2}$	$^3D$	2	83789								

## Bi II—Continued

## Bi II—Continued

Authors	Config.	Desig.	<i>J</i>	Level	Interval	Authors	Config.	Desig.	<i>J</i>	Level	Interval
15 <sub>1</sub>	6p <sub>3/2</sub> 8p <sub>3/2</sub>	<sup>1</sup> D	1	106447		19 <sub>1</sub>	6p <sub>1/2</sub> 7p <sub>1/2</sub>	<sup>1</sup> D	2	109904	
21 <sub>1</sub>	6p <sub>3/2</sub> 7d	<sup>1</sup> D°	1	106611			6p <sub>1/2</sub> 7p <sub>1/2</sub>	<sup>1</sup> S	0	112645	
21 <sub>0</sub>	6p <sub>3/2</sub> 8p <sub>3/2</sub>	<sup>1</sup> P	0	107976		23 <sub>1</sub>	6p <sub>3/2</sub> 6f	<sup>1</sup> G	3	115933	
22 <sub>1</sub>	6p <sub>3/2</sub> 7d <sub>3/2</sub>	<sup>1</sup> F°	3	108007		24 <sub>1</sub>	6p <sub>3/2</sub> 6f <sub>3/2</sub>	<sup>1</sup> F	4	115990	
23 <sub>1</sub>	6p <sub>1/2</sub> 6d	<sup>1</sup> P°	1	108186		25 <sub>1</sub>	6p <sub>3/2</sub> 6f	<sup>1</sup> F	3	116089	
16 <sub>1</sub>	6p <sub>1/2</sub> 7p <sub>1/2</sub>	<sup>1</sup> D	3	108278		26 <sub>1</sub>	6p <sub>3/2</sub> 6f <sub>3/2</sub>	<sup>1</sup> F	2	116212	
17 <sub>1</sub>	6p <sub>3/2</sub> 8p <sub>1/2</sub>	<sup>1</sup> P	1	108404		25°	6p <sub>3/2</sub> 5p <sub>1/2</sub>		3	117004	
22 <sub>1</sub>	6p <sub>3/2</sub> 8p <sub>1/2</sub>	<sup>1</sup> D	2	108886							
18 <sub>1</sub>	6p <sub>1/2</sub> 7p <sub>1/2</sub>	<sup>1</sup> S	1	109104			Bi III( <sup>1</sup> P <sub>1/2</sub> )	Limit		134600	
24 <sub>1</sub>	6p <sub>3/2</sub> 7d	<sup>1</sup> F°	2	109157							

January 1955.

## Bi III

(Tl I sequence; 81 electrons)

Z=83

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^2 P^0_{3/2}$  $6p^2 P^0_{3/2}$  206180 K

I. P. 25.56 volts

The analysis is from Crawford and McLay, who have revised and extended their earlier work and that of Lang. There are 106 classified lines between 590.73 Å and 8934 Å. Arvidsson's measurements have been used in the short-wave region, 590 Å to 1486 Å, to supplement those of the authors.

The limit is based on the assumption that the term  $7h^2 H^0$  is hydrogenic.

By analogy with Pb II, the writer has tentatively assigned *LS*-designations to the terms of the  $6s 6p^2$  configuration. The authors adopt the following numerical labels and *jj*-coupling nomenclature for these levels:

Desig.	Config.	Desig.	Config.
4 <sub>1/2</sub>	6s 6p <sub>3/2</sub> 6p <sub>3/2</sub>	6 <sub>1/2</sub>	6s 6p <sub>3/2</sub> 6p <sub>1/2</sub>
1 <sub>1/2</sub>	6s 6p <sub>3/2</sub> 6p <sub>1/2</sub>	3 <sub>3/2</sub>	6s 6p <sub>1/2</sub> 6p <sub>1/2</sub>
2 <sub>3/2</sub>	6s 6p <sub>3/2</sub> 6p <sub>1/2</sub>	7 <sub>1/2</sub>	6s 6p <sub>1/2</sub> 6p <sub>3/2</sub>
5 <sub>1/2</sub>	6s 6p <sub>3/2</sub> 6p <sub>1/2</sub>	8 <sub>1/2</sub>	6s 6p <sub>1/2</sub> 6p <sub>3/2</sub>

Fisher and Goudsmit report wide hyperfine structure intervals for these levels.

## REFERENCES

- R. A. Fisher and S. Goudsmit, *Phys. Rev.* **37**, 1057 (1931). (hfs)  
 G. Arvidsson, *Ann. der Phys.* **13**, 802 (1932).  
 M. F. Crawford and A. B. McLay, *Proc. Roy. Soc. [A]* **143**, 540 (1934). (I P) (T) (C L) (hfs)  
 F. M. Kelly, R. Richmond, and M. F. Crawford, *Phys. Rev.* **80**, 295 (L) (1950). (hfs)  
 H. E. Walchli, *A Table of Nuclear Moment Data*, Oak Ridge Nat. Lab., ORNL-1469, 63 (1953). (Summary hfs)

## Bi III

## Bi III

Authors	Config.	Desig.	<i>J</i>	Level	Interval	Authors	Config.	Desig.	<i>J</i>	Level	Interval
4 <sub>1/2</sub> 1 <sub>1/2</sub> 2 <sub>1/2</sub>	6s <sup>2</sup> ( <sup>1</sup> S)6p	6p <sup>1</sup> P°	0 $\frac{1}{2}$ 1 $\frac{1}{2}$	0 20768	20788		6s <sup>2</sup> ( <sup>1</sup> S)8p	8p <sup>1</sup> P°	0 $\frac{1}{2}$ 1 $\frac{1}{2}$	154198 156481	2223
	6s 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> P	0 $\frac{1}{2}$ 1 $\frac{1}{2}$ 2 $\frac{1}{2}$	70254 83038 89236	12784 6198		6s <sup>2</sup> ( <sup>1</sup> S)6f	6f <sup>1</sup> F°	3 $\frac{1}{2}$ 2 $\frac{1}{2}$	168186 168848	-56
	6s <sup>2</sup> ( <sup>1</sup> S)7s	7s <sup>1</sup> S	0 $\frac{1}{2}$	95075			6s <sup>2</sup> ( <sup>1</sup> S)5g	5g <sup>1</sup> G	4 $\frac{1}{2}$ 3 $\frac{1}{2}$	166234 166237	-3
5 <sub>1/2</sub> 7 <sub>1/2</sub>	6s <sup>2</sup> ( <sup>1</sup> S)6d	6d <sup>1</sup> D	1 $\frac{1}{2}$ 2 $\frac{1}{2}$	96154 102446	6292		6s <sup>2</sup> ( <sup>1</sup> S)9s	9s <sup>1</sup> S	0 $\frac{1}{2}$	167289	
	6s 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> P	0 $\frac{1}{2}$ 1 $\frac{1}{2}$	108052 130966	22914		6s <sup>2</sup> ( <sup>1</sup> S)8d	8d <sup>1</sup> D	1 $\frac{1}{2}$ 2 $\frac{1}{2}$	169292 169657	365
6 <sub>1/2</sub> 3 <sub>1/2</sub>	6s 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> D	1 $\frac{1}{2}$ 2 $\frac{1}{2}$	108586 116414	7828		6s <sup>2</sup> ( <sup>1</sup> S)6g	6g <sup>1</sup> G	4 $\frac{1}{2}$ 3 $\frac{1}{2}$	178467 178470	-3
	6s <sup>2</sup> ( <sup>1</sup> S)7p	7p <sup>1</sup> P°	0 $\frac{1}{2}$ 1 $\frac{1}{2}$	116993 122128	5135		6s <sup>2</sup> ( <sup>1</sup> S)6h	6h <sup>1</sup> H°	4 $\frac{1}{2}$ , 5 $\frac{1}{2}$	178719	
8 <sub>1/2</sub>	6s 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> S	0 $\frac{1}{2}$	130986?			6s <sup>2</sup> ( <sup>1</sup> S)7g	7g <sup>1</sup> G	4 $\frac{1}{2}$ 3 $\frac{1}{2}$	185850 185853	-3
	6s <sup>2</sup> ( <sup>1</sup> S)5f	5f <sup>1</sup> F°	3 $\frac{1}{2}$ 2 $\frac{1}{2}$	137455 137555	-100		6s <sup>2</sup> ( <sup>1</sup> S)7h	7h <sup>1</sup> H°	4 $\frac{1}{2}$ , 5 $\frac{1}{2}$	186084	
	6s <sup>2</sup> ( <sup>1</sup> S)8s	8s <sup>1</sup> S	0 $\frac{1}{2}$	145227			6s <sup>2</sup> ( <sup>1</sup> S)8g	8g <sup>1</sup> G	4 $\frac{1}{2}$ 3 $\frac{1}{2}$	190642 190644	-2
	6s <sup>2</sup> ( <sup>1</sup> S)7d	7d <sup>1</sup> D	1 $\frac{1}{2}$ 2 $\frac{1}{2}$	149086 149796	710		Bi IV ( <sup>1</sup> S <sub>0</sub> )	Limit	-----	206180	

January 1955.

## Bi IV

(Hg I sequence; 80 electrons)

Z=83

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>10</sup> 6s<sup>2</sup> <sup>1</sup>S<sub>0</sub>6s<sup>2</sup> <sup>1</sup>S<sub>0</sub>, 365500 K

I. P. 45.3 volts

The analysis is incomplete, and is complicated because the spectrum exhibits both hyperfine structure and configuration-interaction. The early work by Arvidsson and by Stanley Smith has been extended by McLay and Crawford to include 139 classified lines in the range from 420.7 Å to 5347 Å. The energy levels in the table are quoted from the paper by these last two authors.

By analogy with Pb III the writer has added tentative *LS*-designations except for the terms <sup>1</sup>P°D°F° having the configuration 5d<sup>9</sup> 6s<sup>2</sup>(<sup>1</sup>D)6p, where the *jj*-coupling notation as given by the authors, is tabulated. For further use of this notation the 1933 reference should be consulted.

The limit has been "obtained by applying a Rydberg formula to 6s 6d 8<sub>2</sub> and 6s 7d 17<sub>2</sub>".

## REFERENCES

- S. Smith, *Nature* **127**, 855 (1931). (T) (C L)  
 G. Arvidsson, *Ann. der Phys.* [5] **12**, 787 (1932). (T) (C L) (hfs)  
 A. B. McLay and M. F. Crawford, *Phys. Rev.* **44**, 986 (1933). (I P) (T) (C L) (hfs)  
 F. M. Kelly, R. Richmond, and M. F. Crawford, *Phys. Rev.* **80**, 295 (L) (1950). (hfs)

Bi IV

Bi IV

Authors	Config.	Desig.	J	Level	Authors	Config.	Desig.	J	Level
1 <sub>a</sub>	5d <sup>10</sup> 6s <sup>2</sup>	6s <sup>2</sup> <sup>1</sup> S	0	0	12 <sub>i</sub>	5d <sup>9</sup> <sub>5/2</sub> 6s <sup>2</sup> 6p <sub>1/2</sub>		4	225617
1 <sub>b</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6p	6p <sup>3</sup> P <sup>o</sup>	0	70963	13 <sub>i</sub>	5d <sup>9</sup> <sub>3/2</sub> 6s <sup>2</sup> 6p <sub>3/2</sub>		1	228187
2 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6p	6p <sup>3</sup> P <sup>o</sup>	1	75926	14 <sub>i</sub>	5d <sup>9</sup> <sub>5/2</sub> 6s <sup>2</sup> 6p <sub>1/2</sub>		2	230637
3 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6p	6p <sup>3</sup> P <sup>o</sup>	2	86423	15 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7p	7p <sup>1</sup> P <sup>o</sup>	1	231913
4 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6p	6p <sup>1</sup> P <sup>o</sup>	1	114608	16 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7p	7p <sup>3</sup> P <sup>o</sup>	2	232627
2 <sub>a</sub>	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>3</sup> P	0	166552	17 <sub>i</sub>	5d <sup>9</sup> 6s <sup>2</sup> 6p <sub>1/2</sub> ?		3	233268
3 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6d	6d <sup>1</sup> D	2	184160	18 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 5f	5f <sup>3</sup> F <sup>o</sup> ?	3	233955
4 <sub>a</sub>	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> P	1	185014	19 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 5f	5f <sup>1</sup> F <sup>o</sup>	2	235172
5 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6d	6d <sup>3</sup> D	1	197151	20 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 5f	5f <sup>3</sup> F <sup>o</sup>	4	235657
6 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7s	7s <sup>1</sup> S	1	197446	21 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 5f	5f <sup>1</sup> F <sup>o</sup> ?	3	236518
7 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6d	6d <sup>3</sup> D	2	197829	22 <sub>i</sub>	5d <sup>9</sup> <sub>5/2</sub> 6s <sup>2</sup> 6p <sub>1/2</sub>		1	237701
8 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 6d	6d <sup>1</sup> D	3	199769	13 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 8s	8s <sup>1</sup> S	1	266674
9 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7s	7s <sup>1</sup> S	0	201581	14 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7d	7d <sup>1</sup> D	1	267576
10 <sub>a</sub>	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>3</sup> P	2	202434	15 <sub>i</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7d	7d <sup>3</sup> D	2	267856
11 <sub>a</sub>	5d <sup>10</sup> 6p <sup>2</sup>	6p <sup>2</sup> <sup>1</sup> D	2	217823	16 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 8s	8s <sup>3</sup> S	0	268191
7 <sub>i</sub>	5d <sup>9</sup> <sub>3/2</sub> 6s <sup>2</sup> 6p <sub>3/2</sub>		2	224292	17 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7d	7d <sup>3</sup> D	3	268690
8 <sup>o</sup> 9 <sup>o</sup> 10 <sup>o</sup> 11 <sup>o</sup>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7p	7p <sup>3</sup> P <sup>o</sup>	0, 1	224524. 6	18 <sub>a</sub>	5d <sup>10</sup> 6s( <sup>2</sup> S) 7d	7d <sup>1</sup> D	2	269758
224600. 2									
224607. 0									
224611. 9									
						Bi v( <sup>2</sup> S <sub>1/2</sub> )	Limit	-----	365506

December 1954.

Bi v

(Au I sequence; 79 electrons)

Z=83

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>10</sup> 6s <sup>2</sup>S<sub>1/2</sub>

6s <sup>2</sup>S<sub>1/2</sub> 451700 K

I. P. 56.0 volts

In 1932 Arvidsson observed the spectrum in the extreme ultraviolet and classified three pairs of doublets. Later, Schoepfle extended the analysis with the aid of Arvidsson's measurements; his data are used in the table. There are 20 classified lines between 355.769 A and 1139.548 A.

Schoepfle states that the hyperfine separation of the ground term is 13 K. The combination of this term with 6p <sup>3</sup>P<sub>0/2</sub> gives rise to two lines having respective wave numbers 87754.1 and 87767.8. Similarly, its combination with 6p <sup>3</sup>P<sub>1/2</sub> gives the two components 115673.9 and 115686.8. Schoepfle adopts an average value for each of the <sup>3</sup>P<sup>o</sup> levels, namely, 87760 and 115681. In order to list the energy levels from the ground state zero, the writer has adopted the lower of the hyperfine-structure levels instead of the average of the two, and adjusted the values of other levels in the table based on combinations with 6p <sup>3</sup>P<sup>o</sup>, accordingly. The small correction (-6 or -7 K) is perhaps negligible when it is considered that the observed Bi v lines lie in the far ultraviolet region. Arvidsson estimates that the measured wavelengths are probably accurate to about 0.03 A.

REFERENCES

G. Arvidsson, Ann. der Phys. [5] 12, 787, (1932). (C L) (hfs)  
 G. K. Schoepfle, Phys. Rev. 47, 232 (1935). (I P) (T) (C L) (hfs)  
 F. M. Kelly, R. Richmond, and M. F. Crawford, Phys. Rev. 80, 295 (L) (1950). (hfs)

## Bi v

Config.	Desig.	<i>J</i>	Level	Interval
$5d^{10}(^1S_0)6s$	$6s\ ^1S$	$0\frac{1}{2}$	0	
$5d^{10}(^1S_0)6p$	$6p\ ^3P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	87754 116674	27920
$5d^9\ 6s^2$	$6s^2\ ^3D$	$2\frac{1}{2}$ $1\frac{1}{2}$	103656 133772	-30116
	1	$1\frac{1}{2}$	223224	
$5d^{10}(^1S_0)6d$	$6d\ ^3D$	$1\frac{1}{2}$ $2\frac{1}{2}$	229289 232540	3251
$5d^{10}(^1S_0)7s$	$7s\ ^1S$	$0\frac{1}{2}$	233340	
$5d^{10}(^1S_0)7p$	$7p\ ^3P^o$	$0\frac{1}{2}$ $1\frac{1}{2}$	270625 281081	10456
$5d^{10}(^1S_0)7d$	$7d\ ^3D$	$1\frac{1}{2}$ $2\frac{1}{2}$	317807 318629	1322
$5d^{10}(^1S_0)8s$	$8s\ ^1S$	$0\frac{1}{2}$	320430	
-----	-----	-----	-----	-----
Bi vi( $^1S_0$ )	<i>Limit</i>	-----	451700	

September 1954.

## Bi vi

(Pt i sequence; 78 electrons)

*Z*=83Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} ^1S_0$  $5d^{10} ^1S_0$ , 712000 K

I. P. 88.3 volts

The analysis is from Schoepfle, who has extended the work of Mack and Fromer. Both investigations are based on the observations of Arvidsson, who reported the first regularities. There are in all, 91 classified lines in the interval 214.318 Å to 1394.46 Å, rejecting one dubious level above the ionization limit; and a level at 176585 which cannot be explained theoretically, as pointed out by Mack.

The spectra of the Pt i isoelectronic sequence are of special theoretical interest because they approach *jj*-coupling rather than *LS*- or *JL*-coupling. This appears more conspicuously in the spectra of higher ionization, especially among the levels from the  $5d^9(^3D)6p$  configuration. A detailed study of "The Four Vector Problem and Its Application to Energies and Intensities in Platinum-like Spectra" has been made by Goble. In *LS*-coupling this configuration gives rise to the terms  $^1,^3(P^oD^oF^o)$ . In *jj*-coupling the same number of energy levels and the same resultant *J*-values occur, but each level is defined by two *j*-values, one for the *d*-electron ( $1\frac{1}{2}$  or  $2\frac{1}{2}$ ), and one for the *p*-electron ( $0\frac{1}{2}$  or  $1\frac{1}{2}$ ).

In the table, the writer has assigned *LS*-designations only in the case of the  $^3D$  terms from the  $5d^9(^3D)ns$  configuration, although the *jj*-coupling applies here also. The *j*-values for the *d*- and *p*-electrons as required for *jj*-coupling are indicated in the configuration column, the former being the same as the *J*-values of the limit term  $^3D$ . A table giving the transformation from *jj*- to *LS*-coupling may be found in the book by Condon and Shortley, p. 294.

Schoepfle uses as the ionization limit the value 761800 derived by a Ritz formula from which he apparently determined the absolute value of the  $5d^9(^3D_{3/2})6p_{1/2}$ , *J*=3, level. The corresponding ionization potential is 94.4 volts.

Mack and Fromer quote an ionization potential derived from a Moseley diagram for the sequence. This estimate is based on a two-member series,  $5d^9(^3D_{3/2})ns$  (*n*=6,7), *J*=3 represented by a Rydberg formula to which the same Ritz correction is applied as for the spectrum of the same stage of ionization in the Au i sequence. Their limit is quoted here.

## Bi VI—Continued

## REFERENCES

- G. Arridsson, *Ann. der Phys.* [5] **12**, 787 (1933). (T) (C L) (hfs)  
 A. T. Goble, *Phys. Rev.* **48**, 346 (1935).  
 J. E. Mack and M. Fromer, *Phys. Rev.* **48**, 357 (1935). (I P) (T) (C L) (hfs)  
 G. K. Schoepfle, *Phys. Rev.* **50**, 538 (1936). (I P) (T) (C L)  
 E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra*, p. 204 (Cambridge University Press, London, 1951).  
 J. E. Mack, letter (June 1956).

Bi VI					Bi VI				
Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
$5d^{10}$	$5d^{10}^1S$	0	0				2	373679	
$5d^9(^3D_{3/2})6s$	$6s^3D$	3	149495	-4242			2	401592	
$5d^9(^1D_{1/2})6s$		2	153737	-21280	$5d^9(^3D_{3/2})7s$	$7s^3D$	3	426587	-896
$5d^9(^3D_{1/2})6s$	$6s^1D$	1	175017		$5d^9(^1D_{1/2})7s$		2	427483	-24797
		2	178479		$5d^9(^3D_{3/2})7s$	$7s^1D$	1	452280	
$5d^9(^3D_{3/2})6p_{3/2}$		2	237881		$5d^9(^1D_{1/2})7s$	$7s^3D$	2	452914	
$5d^9(^3D_{3/2})6p_{1/2}$		2	244781		$5d^9(^3D_{3/2})7s$	$7s^1D$	1	539649	
$5d^9(^1D_{1/2})6p_{3/2}$		3	247165		$5d^9(^3D)7p?$		3	558731	
$5d^9(^1D_{1/2})6p_{1/2}$		2	250191		$5d^9(^3D)7p?$		3	562057	
$5d^9(^3D_{3/2})6p_{3/2}$		2	270803				2	580252	
$5d^9(^3D_{3/2})6p_{1/2}$		1	274158		$5d^9(^3D_{3/2})8s$	$8s^3D$	3	606519	-235
$5d^9(^1D_{1/2})6p_{3/2}$		4	277866				2	606754	
$5d^9(^1D_{1/2})6p_{1/2}$		2	280930		$5d^9(^1D_{1/2})8s$	$8s^1D$	1	631304	
$5d^9(^3D_{3/2})6p_{3/2}$		1	284036				2	644278	
$5d^9(^3D_{3/2})6p_{1/2}$		3	284058		$5d^9(^3D_{3/2})9s$	$9s^3D$	3	711313	
$5d^9(^1D_{1/2})6p_{3/2}$		0	298862				2		
$5d^9(^1D_{1/2})6p_{1/2}$		3	305272				1		
$5d^9(^3D_{3/2})6p_{3/2}$		3	305272						
$5d^9(^3D_{3/2})6p_{1/2}$		1	305903		Bi VII( $^3D_{3/2}$ )	Limit		712000	
$5d^9(^1D_{1/2})6p_{3/2}$		2	308613		$5d^9(^3D)8p?$		3	744960	
$5d^9(^1D_{1/2})6p_{1/2}$		1	315254		$5d^9(^3D)8p?$		3	749193	
		2	319259						

August 1956.

## POLONIUM

## Po I

84 electrons

Z=84

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^4 \text{ } ^3P_1$  $6p^4 \text{ } ^3P_1$ , 67960 K

I. P. 8.43 volts

The analysis is from the paper by Charles, Hunt, Fish, and Timma, who have classified some 50 out of approximately 150 lines observed between 1919.4 Å and 9374.80 Å. They report that a study of the Zeeman effect is in progress. Their limit is from the  $ns \text{ } ^4S^{\circ}$  series ( $n=7, 8$ ), corrected by extrapolating the differences of the effective quantum numbers for Tl I, Pb I, and Bi I.

Mrozowski has classified 7 additional lines in the above list, and added the term  $6p^4 \text{ } ^1S_0$  and the 2 levels 53762 and 54471. He attributes 3 of these lines to forbidden transitions of the multipole type, observed in emission, stating that they are "just the three strongest multipole lines to be expected."

Most of the levels and designations listed below have been reported independently by Vernyi, Zaidel, and Shvebel'blit, whose paper has just become available.

## REFERENCES

- G. W. Charles, D. J. Hunt, G. Fish, and D. L. Timma, J. Opt. Soc. Am. **45**, 869 (1955). (I P) (T) (C L)  
 S. Mrozowski, J. Opt. Soc. Am. **46**, 663 (1956). (T) (C L)  
 E. A. Vernyi, A. N. Zaidel, and K. G. Shvebel'blit, Doklady Akad. Nauk. SSSR **104**, 710 (1955); A. E. R. E. (Harwell, England) Lib/Trans. 681 (1956). (T) (C L)

Po I					Po I						
Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval		
$6p^4$	$6p^4 \text{ } ^3P$	2	0	-16831 9317	$6p^4 \text{ } (^4S^{\circ}) 8s$	$8s \text{ } ^4S^{\circ}?$	2	55923			
		1	16831				$6p^4 \text{ } (^4S^{\circ}) 8s$	$8s \text{ } ^3S^{\circ}?$		1	56268
		0	7514							9 <sup>o</sup>	1
$6p^4$	$6p^4 \text{ } ^1D$	2	21679		$6p^4 \text{ } (^4S^{\circ}) 8p?$	10	1 or 2	59354			
$6p^4 \text{ } (^4S^{\circ}) 7s$	$7s \text{ } ^4S^{\circ}$	2	39081			11 <sup>o</sup>	2	59469			
$6p^4 \text{ } (^4S^{\circ}) 7s$	$7s \text{ } ^3S^{\circ}$	1	40802		$6p^4 \text{ } (^4S^{\circ}) 8p?$	12	1 or 2	59583			
$6p^4$	$6p^4 \text{ } ^1S$	C	42718			13 <sup>o</sup>	1 or 2	61818			
		1 <sup>o</sup>	51712		$6p^4 \text{ } (^4S^{\circ}) 9p?$	14	1 or 2	62704			
		2 <sup>o</sup>	52099			$6p^4 \text{ } (^4S^{\circ}) 9p?$	15	1 or 2	62806		
		3 <sup>o</sup>	52532		16 <sup>o</sup>		1 or 2	62884			
		4 <sup>o</sup>	53027		17 <sup>o</sup>	1 or 2	62959				
		5 <sup>o</sup>	53762		$6p^4 \text{ } (^4S^{\circ}) 10p?$	18	1 or 2	64451			
6 <sup>o</sup>	54471										
		2	55465		$Po \text{ II } (^8P_4)$	Limit		67960			

September 1956.

## RADON

## Rn I

86 electrons

Z=86

Ground State  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^6 {}^1S_0$  $6p^6 {}^1S_0$ , 86692.5 K

I. P. 10.746 volts

The analysis is by Rasmussen, who has classified 172 lines in the interval 3316.14 Å to 10161.45 Å, in addition to the two resonance lines 1451.56 Å and 1786.07 Å. The series are well determined from a Ritz formula.

Edlén has noted that both the run of the quantum defects and the observed combinations indicate that the levels given by Rasmussen in the Paschen notation as  $1s_2$  and  $1s_1$ , are, respectively,  $3d_6$  and  $3d_5$ . This change has been made in the table, and Edlén's predicted positions of the  $1s_2$  and  $1s_1$  levels entered in brackets.

By analogy with Xe I, the writer has introduced *JL*-coupling notation for Rn I in conformity with the designations adopted in these volumes for other spectra of the inert-gas type.

Rasmussen gives the following *LS*-designations for the different types of series:

Paschen	<i>LS</i>	Paschen	<i>LS</i>	Paschen	<i>LS</i>	Paschen	<i>LS</i>
$s_2$	${}^3P_2$	$p_6$	${}^3D_3$	$d'_1$	${}^3F_4$	X	${}^3D_1$
$s_1$	${}^3P_1$	$p_7$	${}^3D_1$	$d_1$	${}^3P_2$	Y	${}^3D_2$
$s_2$	${}^3P_0$	$p_6$	${}^3D_3$	$d_1$	${}^3F_4$	Z	${}^3F_4$
$s_1$	${}^3P_1$	$p_7$	${}^3P_0$	$d'_1$	${}^3D_2$	U	${}^3D_2$
$p_{10}$	${}^3S_1$	$d_6$	${}^3P_0$	$d'_1$	${}^3D_2$	V	${}^3F_3$
$p_9$	${}^1D_2$	$d_5$	${}^3P_1$	$d_1$	${}^3D_1$	W	${}^3F_2$

## REFERENCES

- E. Rasmussen, Zeit. Phys. **80**, 726 (1933). (I P) (T) (C L)  
 B. Edlén, Ark. Mat. Astr. Fys. (Stockholm) **29A**, No. 21, 4 (1943).

## Rn I

## Rn I

Author	Config.	Desig.	<i>J</i>	Level	Author	Config.	Desig.	<i>J</i>	Level
$p_9$	$6p^6$	$6p^6 {}^1S$	0	0.0	$3d_1$	$6p^6({}^3P_{1,2})6d$	$6d [1\frac{1}{2}]^{\circ}$	2 1	70222.82
$1s_2$ $1s_1$	$6p^6({}^3P_{1,2})7s$	$7s [1\frac{1}{2}]^{\circ}$	2 1	54820.35 55989.03	$3p_{10}$	$6p^6({}^3P_{1,2})8p$	$8p [0\frac{1}{2}]$	1	77039.38
$1s_2$ $1s_1$	$6p^6({}^3P_{1,2})7s$	$7s' [0\frac{1}{2}]^{\circ}$	0 1	[85876] [87053]	$3p_9$ $3p_8$	"	$8p [2\frac{1}{2}]$	2 3	77161.69 7604.53
$2p_{10}$	$6p^6({}^3P_{1,2})7p$		1	66244.97	$3p_7$ $3p_6$	"	$8p [1\frac{1}{2}]$	1 2	77677.83 77827.82
$2p_9$ $2p_8$	"		2 3	66707.53 68039.48	$3p_5$	"	$8p [0\frac{1}{2}]$	0	78163.27
$2p_7$ $2p_6$	"	$7p [1\frac{1}{2}]$	1 2	68332.10 68789.93	$4d_9$ $4d_1$	$6p^6({}^3P_{1,2})7d$	$7d [0\frac{1}{2}]^{\circ}$	0 1	77594.82 77816.18
$2p_5$	"	$7p [0\frac{1}{2}]$	0	69743.98	$4d'_1$ $4d_1$	"	$7d [3\frac{1}{2}]^{\circ}$	4 3	78088.48 78230.67
$3d_6$ $3d_5$	$6p^6({}^3P_{1,2})6d$	$6d [0\frac{1}{2}]^{\circ}$	0 1	67906.52 68891.34	$4d_3$ $4d_2$	"	$7d [1\frac{1}{2}]^{\circ}$	2 1	78173.30
$3d'_1$ $3d_1$	"	$6d [3\frac{1}{2}]^{\circ}$	4 3	69798.00 70440.48	$4d'_1$ $4d_1$	"	$7d [2\frac{1}{2}]^{\circ}$	2 3	78475.97 78688.88



Rn I—Continued

Rn I—Continued

Author	Config.	Desig.	J	Level	Author	Config.	Desig.	J	Level
3e <sub>0</sub> 3e <sub>1</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9e	9e [1½] <sup>o</sup>	2 1	79696. 50 79762. 87	6X 6Y	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 7f	7f [1½]	1 2	83575. 07 83575. 55
4X 4Y	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 5f	5f [1½]	1 2	79639. 03 79644. 93	6Z	"	7f [4½]	5 4	83592. 53
4Z	"	5f [4½]	5 4	79690. 3	6U 6V	"	7f [2½]	3 2	83607. 96 83610. 71
4U 4V	"	5f [2½]	3 2	79745. 40 79750. 02	6W	"	7f [3½]	3 4	83622. 40
4W	"	5f [3½]	3 4	79785. 18	5e <sub>1</sub> 5e <sub>1</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 11e	11e [1½] <sup>o</sup>	2 1	83595. 18 83630. 49
4p <sub>10</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9p	9p [0½]	1	80998. 06	7d <sub>0</sub> 7d <sub>0</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10d	10d [0½] <sup>o</sup>	0 1	84075. 08 84111. 60
4p <sub>0</sub> 4p <sub>0</sub>	"	9p [2½]	2 3	81049. 29 81253. 54	7d <sub>1</sub> 7d <sub>1</sub>	"	10d [3½] <sup>o</sup>	4 3	84162. 11 84177. 33
4p <sub>1</sub> 4p <sub>1</sub>	"	9p [1½]	1 2	81283. 24 81351. 52	7d <sub>2</sub> 7d <sub>2</sub>	"	10d [1½]	2 1	84169. 80
4p <sub>2</sub>	"	9p [0½]	0	81510. 27	7d <sub>3</sub> 7d <sub>3</sub>	"	10d [2½] <sup>o</sup>	2 3	84210. 00 84236. 50
5d <sub>0</sub> 5d <sub>0</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 8d	8d [0½] <sup>o</sup>	0 1	81234. 15 81329. 96	6p <sub>0</sub> 6p <sub>0</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 11p	11p [2½]	2 3	84099. 35
5d <sub>1</sub> 5d <sub>1</sub>	"	8d [3½] <sup>o</sup>	4 3	81514. 40	6p <sub>1</sub> 6p <sub>1</sub>	"	11p [1½]	1 2	84108. 5 84131. 6
5d <sub>2</sub> 5d <sub>2</sub>	"	8d [1½] <sup>o</sup>	2 1	81490. 50 81966. 30	6p <sub>2</sub>	"	11p [0½]	0	84183. 06
5d <sub>3</sub> 5d <sub>3</sub>	"	8d [2½] <sup>o</sup>	2 3	81619. 30 81695. 16	7X 7Y	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 8f	8f [1½]	1 2	84407. 70 84410. 41
5X 5Y	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 6f	6f [1½]	1 2	82190. 55 82195. 97	7Z	"	8f [4½]	5 4	84419. 04
5Z	"	6f [4½]	5 4	82218. 92	6s <sub>1</sub> 6s <sub>1</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12s	12s [1½] <sup>o</sup>	2 1	84425. 40
5U 5V	"	6f [2½]	3 2	82246. 73 82251. 06	7p <sub>0</sub> 7p <sub>0</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12p	12p [2½]	2 3	84746. 35
5W	"	6f [3½]	3 4	82270. 68	7p <sub>1</sub> 7p <sub>1</sub>	"	12p [1½]	1 2	84767. 3
4s <sub>1</sub> 4s <sub>1</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10s	10s [1½] <sup>o</sup>	2 1	82211. 18 82275. 97	8d <sub>1</sub> 8d <sub>1</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 11d	11d [3½] <sup>o</sup>	4 3	84785. 85 84795. 53
5p <sub>10</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10p	10p [0½]	1	82927. 18	8d <sub>2</sub> 8d <sub>2</sub>	"	11d [1½] <sup>o</sup>	2 1	84791. 24
5p <sub>0</sub> 5p <sub>0</sub>	"	10p [2½]	2 3	82953. 90 83064. 98	8d <sub>3</sub> 8d <sub>3</sub>	"	11d [2½] <sup>o</sup>	2 3	84816. 50 84833. 91
5p <sub>1</sub> 5p <sub>1</sub>	"	10p [1½]	1 2	83079. 34 83116. 71	8Z	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9f	9f [4½]	5 4	84954. 68
5p <sub>2</sub>	"	10p [0½]	0	83203. 97	8U	"	9f [2½]	3 2	84960. 71
6d <sub>0</sub> 6d <sub>0</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 9d	9d [0½] <sup>o</sup>	0 1	83042. 53 83096. 24	9d <sub>1</sub> 9d <sub>1</sub>	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 12d	12d [3½] <sup>o</sup>	4 3	85204. 47 85210. 91
6d <sub>1</sub> 6d <sub>1</sub>	"	9d [3½] <sup>o</sup>	4 3	83172. 08 83199. 40	9Z	6p <sup>2</sup> ( <sup>2</sup> P <sub>1/2</sub> ) 10f	10f [4½]	5 4	85322. 08
6d <sub>2</sub> 6d <sub>2</sub>	"	9d [1½] <sup>o</sup>	2 1	83186. 66 83449. 38					
6d <sub>3</sub> 6d <sub>3</sub>	"	9d [2½] <sup>o</sup>	2 3	83254. 04 83297. 22					
						Rn II( <sup>2</sup> P <sub>1/2</sub> )	Limit		86692. 5

Rn I OBSERVED LEVELS\*

Configuration	Observed Terms			
	$ns$ ( $n \geq 7$ )	$np$ ( $n \geq 7$ )	$nd$ ( $n \geq 6$ )	$nf$ ( $n \geq 5$ )
$6p^4$	$7, 9-12s^1p^3$	$7-10p^1s$	$6-11d^1p^2$	$5-9f^1D \quad 5-10f^1F$
$6p^4(3P^0)ns$	$7, 9-12s^1p^3$	$7-11p^1p^2$ $7-10p^1D$	$7-11d^1D^0$	
<i>Jl</i> Coupling Notation				
$6p^4(3P^1_{1/2})ns$	Observed Pairs			
	$ns$ ( $n \geq 7$ )	$np$ ( $n \geq 7$ )	$nd$ ( $n \geq 6$ )	$nf$ ( $n \geq 5$ )
	$7, 9-12s^1(1/2)^0$	$7-11p^1(0/2)$ $7-12p^1(2/2)$ $7-12p^1(1/2)$	$6-10d^1(0/2)$ $6-12d^1(3/2)$ $6-11d^1(1/2)$ $7-11d^1(2/2)$	$5-9f^1(1/2)$ $5-10f^1(4/2)$ $5-7, 9f^1(2/2)$ $5-7f^1(3/2)$

\*For predicted levels in the spectra of the Rn I isoelectronic sequence, see Vol. III, Introduction.

Rn II

(At I sequence; 85 electrons)

Z=86

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^6 7p^{1/2}$

$6p^4 \ ^3P^1_{1/2}$  K

I. P. volts

The spectrum is still unknown. Edlén has suggested that the one unclassified line at 3235.83 Å, listed by Rasmussen as Rn I, may be Rn II. In discussing magnetic-dipole transitions in the configurations  $5p^4$  and  $6p^4$  of xenon and radon, Edlén states that although the interval of the ground term in Rn II is not known, "an approximate value around 31,000 cm<sup>-1</sup> was indirectly deduced from a certain perturbation in the observed  $nd_4$  series of Rn I." The wave number of the above line is 30895.1, and the transition probability is given by Edlén as 531 sec<sup>-1</sup>. "The clue to the analysis of Rn II" may be, therefore, to start with the term:

Design.	J	Level	Interval
$6p^4 \ ^3P^0$	$1\frac{1}{2}$ $0\frac{1}{2}$	$0.0$ $30895.1$	$-30895.1$

REFERENCE

B. Edlén, Phys. Rev. **65**, 248, 1944. (C L)

February 1955.

## RADIUM

## Ra I

88 electrons

Z=88

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^6 7s^2 {}^1S_0$  $7s^2 {}^1S_0$ , 42577.35 K

I. P. 5.277 volts

The analysis is from Rasmussen, with the revisions suggested by Russell on the basis of the comparison of Ra I with the analogous spectra Be I, Mg I, Ca I, Sr I, and Ba I. In the first column of the table Rasmussen's notation is entered. The changes in analysis can be detected by comparing the designations in this column with those in column three.

There are 69 classified lines between 2955.65 Å and 9932.21 Å. Observed intersystem combinations connect the singlet and triplet systems of terms.

The limit is from the  ${}^3F^{\circ}$  series. Rasmussen's value has been increased by 627.66 K to agree with the revisions introduced by Russell in the connection between certain groups of terms.

## REFERENCES

- E. Rasmussen, *Zeit. Phys.* **87**, 607 (1934). (I P) (T) (C L)  
 H. N. Russell, *Phys. Rev.* **46**, 989 (1934). (I P) (T) (C L)

## Ra I

## Ra I

Author	Config.	Desig.	J	Level	Interval	Author	Config.	Desig.	J	Level	Interval
1 ${}^1S_0$	$7s^2$	$7s^2 {}^1S$	0	0.00		3 ${}^1D_2$	$7s({}^3S)6d$	$6d {}^1D$	2	17081.45	
2 ${}^1P_0$	$7s({}^3S)7p$	$7p {}^3P^{\circ}$	0	15078.44	920.94 2689.16	2 ${}^1P_1$	$7s({}^3S)7p$	$7p {}^1P^{\circ}$	1	20715.71	
2 ${}^1P_1$			1	15999.38		2 ${}^3S_1$	$7s({}^3S)8s$	$8s {}^3S$	1	26754.05	
2 ${}^1P_2$			2	16688.54		$d_p {}^3F_2$	$6d({}^3D)7p$	$7p' {}^3F^{\circ}$	2	28038.06	2079.73 2250.00
3 ${}^3D_1$	$7s({}^3S)6d$	$6d {}^3D$	1	13715.85	278.12	$d_p {}^3F_3$		3	30117.78		
3 ${}^3D_2$			2	13993.97	713.38	$d_p {}^3F_4$		4	32367.78		
3 ${}^3D_3$			3	14707.35							

Ra I—Continued

Ra I—Continued

Author	Config.	Design.	J	Level	Interval	Author	Config.	Design.	J	Level	Interval
$dp^1D_2$	$6d(^3D)7p$	$7p^1D^0$	2	30918.14		$3^3S_1$	$7s(^3S)9s$	$9s^3S$	1	34475.78	
$dp^3P_1$	$7s(^3S)8p$	$8p^3P^0$	0	31085.88	477.41	$4^3F_2$ $4^3F_3$ $4^3F_4$	$7s(^3S)8f$	$8f^3F^0$	2	35255.65	12.37
			1	31563.89	311.15				3	35968.08	
			2	31874.44					4	35994.40	
$4^1D_1$ $dd^3P_1$	$7p^3$	$7p^3^3P$	0			$5^3F_2$ $5^3F_4$	$7s(^3S)6f$	$6f^3F^0$	2		7.38
			1	31248.61	1692.53				3	37928.80	
			2	32941.13					4	37989.64	
$dd^3P_1$ $4^1D_1$ $4^1D_2$	$7s(^3S)7d$	$7d^3D$	1	32000.82	-7.41	$6^3F_2$ $6^3F_4$	$7s(^3S)7f$	$7f^3F^0$	2		6.02
			2	31993.41	203.87				3	39360.96	
			3	32197.28					4	39366.98	
$dd^1D_1$	$7p^3$	$7p^3^1D$	2	32214.84		$7^3F_1$ $7^3F_2$ $7^3F_4$	$7s(^3S)8f$	$8f^3F^0$	2	40819.5	0.0 12.1
$dp^3D_1$ $dp^3D_2$ $dp^3D_3$	$6d(^3D)7p$	$7p^3^3D^0$	1	32229.87	276.62				3	40819.5	
			2	32506.59	690.87				4	40831.58	
			3	33197.46							
$3^1P_1$	$7s(^3S)8p$	$8p^1P^0$	1	32857.68		Ra II ( $^3S_{40}$ )	Limit	---		42577.35	
	$6d(^3D)7p$	$7p^3^3P^0$	0	33782.41	41.29						
$dp^3P_1$			1	33853.70	559.21						
			2	34388.91							

March 1955.

Ra I OBSERVED TERMS\*

Configuration $1s^2 2s^2 2p^6 3s^2 3p^4$ $3d^{10} 4s^2 4p^6 4d^{10} 4f^{14}$ $5s^2 5p^6 5d^{10} 6s^2 6p^4 +$	Observed Terms			
$7s^2$	$7s^2^1S$			
$7p^3$	{	$7p^3^3P$		$7p^3^1D$
		$ns (n \geq 7)$		$np (n \geq 7)$
$7s(^3S)ns$	{	$8s, 9s^3S$	$7, 8p^3^3P^0$ $7, 8p^3^1P^0$	$6, 7d^3D$ $6d^1D$
$6d(^3D)ns'$	{		$7p^3^3P^0$ $7p^3^3D^0$ $7p^3^3F^0$ $7p^3^1D^0$	$5-8f^3F^0$

\*For predicted terms in the spectra of the Ra I isoelectronic sequence, see Vol. III, Introduction.

## Ra II

(Fr I sequence; 87 electrons)

Z=88

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^6 7s^2 \text{ } ^1\text{S}_{00}$  $7s^2 \text{ } ^1\text{S}_{00}$  81842.31 K

I. P. 10.144 volts

The analysis is by Rasmussen, who has classified 64 lines in the range 1888.7 Å to 9453.57 Å, from observations made with a hollow-cathode source. The spectrum gives well-established series of doublet terms. Rasmussen stresses the similarity of this spectrum and Ba II.

Revised values of the  $6f \text{ } ^3\text{F}^{\circ}$  term are from the infrared combination with  $7d \text{ } ^3\text{D}$  given in the later reference.

## REFERENCES

E. Rasmussen, Zeit. Phys. 66, 24 (1933). (I F) (T) (C L)

E. Rasmussen, Zeit. Phys. 87, 615 (1934). (C L)

## Ra II

## Ra II

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
$6p^4(^3\text{S})7s$	$7s \text{ } ^3\text{S}$	0½	0.00		$6p^4(^3\text{S})10s$	$10s \text{ } ^3\text{S}$	0½	66837.88	
$6p^4(^3\text{S})6d$	$6d \text{ } ^3\text{D}$	1½ 2½	12084.38 18743.11	1658.73	$6p^4(^3\text{S})9d$	$9d \text{ } ^3\text{D}$	1½ 2½	68264.07 68394.86	130.79
$6p^4(^3\text{S})7p$	$7p \text{ } ^3\text{P}^{\circ}$	0½ 1½	21351.20 26208.86	4857.66	$6p^4(^3\text{S})6g$	$6g \text{ } ^3\text{G}$	3½, 4½	69560.85	
$6p^4(^3\text{S})8s$	$8s \text{ } ^3\text{S}$	0½	48405.01		$6p^4(^3\text{S})11s$	$11s \text{ } ^3\text{S}$	0½	71172.92	
$6p^4(^3\text{S})7d$	$7d \text{ } ^3\text{D}$	1½ 2½	48744.04 49240.48	496.44	$6p^4(^3\text{S})10d$	$10d \text{ } ^3\text{D}$	1½ 2½	72043.06 72123.78	80.72
$6p^4(^3\text{S})5f$	$5f \text{ } ^3\text{F}^{\circ}$	2½ 3½	48937.38 49872.31	284.33	$6p^4(^3\text{S})7g$	$7g \text{ } ^3\text{G}$	3½, 4½	72824.37	
$6p^4(^3\text{S})8p$	$8p \text{ } ^3\text{P}^{\circ}$	0½ 1½	50608.01 52392.05	1784.04	$6p^4(^3\text{S})11d$	$11d \text{ } ^3\text{D}$	1½ 2½	74434.04 74488.51	54.47
$6p^4(^3\text{S})9s$	$9s \text{ } ^3\text{S}$	0½	59165.23		$6p^4(^3\text{S})8g$	$8g \text{ } ^3\text{G}$	3½, 4½	74941.91	
$6p^4(^3\text{S})6f$	$6f \text{ } ^3\text{F}^{\circ}$	2½ 3½	59515.48 59815.59	300.11	$6p^4(^3\text{S})12d$	$12d \text{ } ^3\text{D}$	1½ 2½	76080.74	
$6p^4(^3\text{S})8d$	$8d \text{ } ^3\text{D}$	1½ 2½	61734.88 61973.82	238.94	$6p^4(^3\text{S})9g$	$9g \text{ } ^3\text{G}$	3½, 4½	76393.12	
$6p^4(^3\text{S})9p$	$9p \text{ } ^3\text{P}^{\circ}$	0½ 1½	63410.41		$6p^4(^3\text{S})10g$	$10g \text{ } ^3\text{G}$	3½, 4½	77431.31	
$6p^4(^3\text{S})5g$	$5g \text{ } ^3\text{G}$	3½, 4½	64150.65		$6p^4(^3\text{S})11g$	$11g \text{ } ^3\text{G}$	3½, 4½	78195.21	
$6p^4(^3\text{S})7f$	$7f \text{ } ^3\text{F}^{\circ}$	2½ 3½	66521.86 66891.82	169.26	Ra III ( $^3\text{S}_0$ )	Limit		81842.31	

March 1955.

## ACTINIUM

## Ac I

89 electrons

Z=89

Ground state  $1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^6 7s^2 6d^1 D_{3/2}$  $s^1 D_{3/2}$ 

K

I. P.

volts

The observation, description, and analyses of the emission spectra of actinium have been undertaken especially for inclusion here. The analysis is by Meggers, who has measured and interpreted arc and hollow-cathode spectrograms made at the Argonne National Laboratory by Fred and Tomkins. Thus far, 87 Ac I lines, with wavelengths ranging from 2968.82 Å to 7866.10 Å, have been classified as combinations among 40 energy levels. An estimated value of  $s^1 P_{3/2}^o$  is entered in brackets in the table. This term has not yet been found because of the lack of infrared observations.

No series have been found. From a study of the screening constants, Finkelnburg and Humbach have extrapolated an ionization potential of  $6.9 \pm 0.6$  volts.

## REFERENCES

W. F. Meggers, M. Fred, and F. S. Tomkins, *J. Opt. Soc. Am.* **41**, 867 (A) (1951). (T)

M. Fred, F. S. Tomkins, and W. F. Meggers, *Phys. Rev.* **98**, 1514 (L) (1955). (hfs)

W. Finkelnburg und W. Humbach, *Naturwiss.* **43**, 35 (1955). (I P)

W. F. Meggers, M. Fred, and F. S. Tomkins, *J. Research Nat. Bur. Std.* **58**, 297, RP2763 (1957). (T) (C L)

Ac I

Ac I

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval																																																																																																																																						
6d 7s <sup>2</sup>	a <sup>1</sup> D	1½	0.00	2231.43	6d 7s(a <sup>1</sup> D)7p	y <sup>1</sup> D <sup>o</sup>	1½	26066.04	467.12																																																																																																																																						
		2½	2231.43				2½	26553.16		6d <sup>2</sup> (a <sup>1</sup> F)7s	a <sup>1</sup> F	1½	9217.28	646.31	6d 7s(a <sup>1</sup> D)7p	y <sup>1</sup> F <sup>o</sup>	2½	26836.20	1732.20	2½	9863.59	3½	26568.40	3½	10906.02			4½	12078.07			7s <sup>2</sup> (a <sup>1</sup> S)7p	z <sup>1</sup> P <sup>o</sup>	0½			6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> P <sup>o</sup>	0½			1½	[10000±]	1½	30396.61	6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> F <sup>o</sup>	1½	15712.90	1227.82	6d <sup>2</sup> (a <sup>1</sup> F)7p	z <sup>1</sup> G <sup>o</sup>	2½	31494.68	724.94	2½	14940.72	3½	32219.62	3½	17683.87	4½	32867.39	4½		5½	33489.76	6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> D <sup>o</sup>	0½	17199.71	1812.75	6d <sup>2</sup> (a <sup>1</sup> F)7p		1½	31800.35	562.37	1½	19012.48	2½	32495.67	2½	21195.87	1½	32918.40	3½	23476.94	2½	33673.66	6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> D <sup>o</sup>	1½	17736.28	214.45	6d <sup>2</sup> (a <sup>1</sup> F)7p		1½	33756.43		2½	17950.71	3½	34380.25	6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> P <sup>o</sup>	0½	22401.52	399.58	6d <sup>2</sup> (a <sup>1</sup> F)7p		3½	34380.25		1½	22801.10	2½	34658.47	2½	23898.88	1097.76		6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> F <sup>o</sup>	2½	23916.54	1052.46	6d <sup>2</sup> (a <sup>1</sup> F)7p		4½	34788.18		3½	24969.30	4½	35370.00	6d 7s(a <sup>1</sup> D)7p	y <sup>1</sup> P <sup>o</sup>	0½	25729.03	1280.81	6d <sup>2</sup> (a <sup>1</sup> F)7p		4½
6d <sup>2</sup> (a <sup>1</sup> F)7s	a <sup>1</sup> F	1½	9217.28	646.31	6d 7s(a <sup>1</sup> D)7p	y <sup>1</sup> F <sup>o</sup>	2½	26836.20	1732.20																																																																																																																																						
		2½	9863.59				3½	26568.40																																																																																																																																							
		3½	10906.02																																																																																																																																												
		4½	12078.07																																																																																																																																												
7s <sup>2</sup> (a <sup>1</sup> S)7p	z <sup>1</sup> P <sup>o</sup>	0½			6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> P <sup>o</sup>	0½																																																																																																																																								
		1½	[10000±]				1½	30396.61		6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> F <sup>o</sup>	1½	15712.90	1227.82	6d <sup>2</sup> (a <sup>1</sup> F)7p	z <sup>1</sup> G <sup>o</sup>	2½	31494.68	724.94	2½	14940.72	3½	32219.62	3½	17683.87	4½	32867.39	4½		5½	33489.76	6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> D <sup>o</sup>	0½	17199.71	1812.75	6d <sup>2</sup> (a <sup>1</sup> F)7p		1½	31800.35	562.37	1½	19012.48	2½	32495.67	2½	21195.87	1½	32918.40	3½	23476.94	2½	33673.66	6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> D <sup>o</sup>	1½	17736.28	214.45	6d <sup>2</sup> (a <sup>1</sup> F)7p		1½	33756.43		2½	17950.71	3½	34380.25	6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> P <sup>o</sup>	0½	22401.52	399.58	6d <sup>2</sup> (a <sup>1</sup> F)7p		3½	34380.25		1½	22801.10	2½	34658.47	2½	23898.88	1097.76		6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> F <sup>o</sup>	2½	23916.54	1052.46	6d <sup>2</sup> (a <sup>1</sup> F)7p		4½	34788.18		3½	24969.30	4½	35370.00	6d 7s(a <sup>1</sup> D)7p	y <sup>1</sup> P <sup>o</sup>	0½	25729.03	1280.81	6d <sup>2</sup> (a <sup>1</sup> F)7p		4½	35370.00		1½	27009.84																																
6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> F <sup>o</sup>	1½	15712.90	1227.82	6d <sup>2</sup> (a <sup>1</sup> F)7p	z <sup>1</sup> G <sup>o</sup>	2½	31494.68	724.94																																																																																																																																						
		2½	14940.72				3½	32219.62																																																																																																																																							
		3½	17683.87				4½	32867.39																																																																																																																																							
		4½					5½	33489.76																																																																																																																																							
6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> D <sup>o</sup>	0½	17199.71	1812.75	6d <sup>2</sup> (a <sup>1</sup> F)7p		1½	31800.35	562.37																																																																																																																																						
		1½	19012.48				2½	32495.67																																																																																																																																							
		2½	21195.87				1½	32918.40																																																																																																																																							
		3½	23476.94				2½	33673.66																																																																																																																																							
6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> D <sup>o</sup>	1½	17736.28	214.45	6d <sup>2</sup> (a <sup>1</sup> F)7p		1½	33756.43																																																																																																																																							
		2½	17950.71				3½	34380.25																																																																																																																																							
6d 7s(a <sup>1</sup> D)7p	s <sup>1</sup> P <sup>o</sup>	0½	22401.52	399.58	6d <sup>2</sup> (a <sup>1</sup> F)7p		3½	34380.25																																																																																																																																							
		1½	22801.10				2½	34658.47																																																																																																																																							
		2½	23898.88				1097.76																																																																																																																																								
6d 7s(a <sup>1</sup> D)7p	z <sup>1</sup> F <sup>o</sup>	2½	23916.54	1052.46	6d <sup>2</sup> (a <sup>1</sup> F)7p		4½	34788.18																																																																																																																																							
		3½	24969.30				4½	35370.00																																																																																																																																							
6d 7s(a <sup>1</sup> D)7p	y <sup>1</sup> P <sup>o</sup>	0½	25729.03	1280.81	6d <sup>2</sup> (a <sup>1</sup> F)7p		4½	35370.00																																																																																																																																							
		1½	27009.84																																																																																																																																												

June 1967.

## Ac I Observed Terms\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>10</sup> 6s <sup>2</sup> 6p <sup>6</sup> +	Observed Terms			
6d 7s <sup>2</sup>	a <sup>3</sup> D			
	ns (n ≥ 7)	np (n ≥ 7)		
7s <sup>2</sup> (a <sup>1</sup> S)ns	{			
6d 7s(a <sup>1</sup> D)ns		s <sup>1</sup> P <sup>o</sup> y <sup>1</sup> P <sup>o</sup>	s <sup>1</sup> D <sup>o</sup> s <sup>1</sup> D <sup>o</sup>	
6d 7s(a <sup>1</sup> D)ns		s <sup>1</sup> P <sup>o</sup>	y <sup>1</sup> D <sup>o</sup>	y <sup>1</sup> F <sup>o</sup>
6d <sup>2</sup> (a <sup>1</sup> F)ns		a <sup>1</sup> F		s <sup>1</sup> G <sup>o</sup>

\*For predicted terms in the spectra of the Ac I isoelectronic sequence, see Vol. III, Introduction.

## Ac II

(Ra I sequence; 88 electrons)

Z=89

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>10</sup> 6s<sup>2</sup> 6p<sup>4</sup> 7s<sup>2</sup> <sup>1</sup>S<sub>0</sub>a<sup>1</sup>S<sub>0</sub> 97300 K

I. P. 12.1 volts

The analysis is by Meggers, who has measured and interpreted spark and hollow-cathode spectrograms made at the Argonne National Laboratory by Fred and Tomkins. This work has been carried out especially for inclusion here. Meggers has furnished manuscript in advance of publication. He states that "The Ac II spectrum is more complex than Ac I because the 5f-electron contributes many excited odd levels. Thus far, 226 Ac II lines with wavelengths from 2261.75 Å to 7886.82 Å have been classified as combinations of 27 even and 38 odd levels."

Racah has suggested the *jj*-coupling arrangement for the levels from the 5f(<sup>2</sup>F<sup>o</sup>)7p configuration. The tentative *LS*-designations ascribed to these levels agree reasonably well both with the intensities of the observed combinations and with the theory for 2-electron spectra by Racah.

The limit is from the 7s(<sup>2</sup>S)ns series (n=7, 8), and has been derived from a Rydberg formula, corrected by analogy with Ra II. From a study of screening constants Finkelnburg and Humbach interpolate an ionization potential of 11.5 ± 0.4 volts.

## REFERENCES

- G. Racah, *Physica* 16, Nos. 7-8, 651 (1950).  
 W. F. Meggers, M. Fred, and F. S. Tomkins, *J. Opt. Soc. Am.* 41, 867 (A) (1951). (T)  
 M. Fred, F. S. Tomkins, and W. F. Meggers, *Phys. Rev.* 96, 1514 (L) (1955). (hfs)  
 W. Finkelnburg und W. Humbach, *Naturwiss.* 42, 35 (1955). (I P)  
 W. F. Meggers, M. Fred, and F. S. Tomkins, *J. Research Nat. Bur. Std.* 58, 297, RP2763 (1957). (I P) (T)  
 (C L)

## Ac II

## Ac II

Config.	Design.	J	Level	Interval	Config.	Design.	J	Level	Interval
7s	e <sup>1</sup> S	0	0.00		6d(°D)7p	y <sup>1</sup> F°	3	41857.05	
6d(°D)7s	e <sup>1</sup> D	1	4739.63	527.53	6d(°D)7p	y <sup>1</sup> P°	1	4,199.94	
		2	5267.16	2159.36	6d(°D)5f	y <sup>1</sup> D°	2	44705.33	
		3	7426.52		6d(°D)5f	s <sup>1</sup> G°	3	45807.08	1620.61
6d(°D)7s	e <sup>1</sup> D	2	9087.54				4	47487.67	1723.87
6s	e <sup>1</sup> F	2	13236.46	1712.75			5	49151.54	
		3	14949.21	1807.69	6d(°D)5f	y <sup>1</sup> D°	1	48518.41	960.64
		4	16756.90				2	49479.05	580.65
6s	e <sup>1</sup> P	0	17737.10	1278.22			3	50059.70	
		1	19015.32	3184.13	6d(°D)5f	z <sup>1</sup> P°	0	49780.60	455.00
		2	22199.45				1	50335.60	1021.91
6s	b <sup>1</sup> D	2	19203.02				2	51857.51	
6s	e <sup>1</sup> G	4	20848.23		7s(°S)8s	e <sup>1</sup> S	1	51680.55	
7s(°S)7p	z <sup>1</sup> P°	0	30956.40	1224.12	6d(°D)5f	z <sup>1</sup> F°	3	53355.30	
		1	33180.53	9698.35	7s(°S)8s	e <sup>1</sup> S	0	53374.01	
		2	31878.87		5f(°F <sub>3/2</sub> )7p <sub>1/2</sub>	e <sup>1</sup> F	2	54633.05	
6d(°D)7p	z <sup>1</sup> F°	2	36448.96	4727.64	"	e <sup>1</sup> G	3	54644.11	
		3	31174.60	7944.42	6d(°D)5f	z <sup>1</sup> P°	1	56152.3	
		4	39119.03		5f(°F <sub>3/2</sub> )7p <sub>1/2</sub>	e <sup>1</sup> G	4	57683.78	
7s(°S)5f	y <sup>1</sup> F°	2	33201.11	1679.98	"	e <sup>1</sup> D	3	57944.57	
		3	33831.09	3084.46	5f(°F <sub>3/2</sub> )7p <sub>1/2</sub>	e <sup>1</sup> F	3	59071.9	
		4	33965.55		"	e <sup>1</sup> D	2	59807.5	
6d(°D)7p	z <sup>1</sup> D°	1	39250.40	4054.56	"	e <sup>1</sup> G	4	59889.05	
		2	33304.96	3667.98	"	e <sup>1</sup> D	1	60063.0	
		3	36973.84		5f(°F <sub>3/2</sub> )7p <sub>1/2</sub>	e <sup>1</sup> F	3	62350.68	
7s(°S)7p	z <sup>1</sup> P°	1	33388.61		"	e <sup>1</sup> F	4	62869.81	
6d(°D)7p	z <sup>1</sup> D°	2	35397.13		"	e <sup>1</sup> D	2	63581.56	
7s(°S)5f	z <sup>1</sup> F°	3	36144.35		"	e <sup>1</sup> G	5	65392.46	
6d(°D)7p	y <sup>1</sup> P°	0	36780.01	75.49	5f(°F <sub>3/2</sub> )7p <sub>1/2</sub>	e <sup>1</sup> F	3	62350.68	
		1	36855.50	1516.14	"	e <sup>1</sup> F	4	62869.81	
		2	38371.64		"	e <sup>1</sup> D	2	63581.56	
6d(°D)5f	z <sup>1</sup> H°	4	39807.14	1109.27	"	e <sup>1</sup> G	5	65392.46	
		5	40916.41						
		6							
6d(°D)5f	z <sup>1</sup> F°	2	41578.57	1697.28					
		3	43275.85	1764.45					
		4	45040.30						
6d(°D)5f	s <sup>1</sup> G°	4	41637.79						
					Ac III(°S <sub>3/2</sub> )	Limit		97300	

June 1957.



## As II OBSERVED TERMS\*

Configuration 1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>2</sup> 3p <sup>4</sup> 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>6</sup> 4d <sup>10</sup> 4f <sup>14</sup> 5s <sup>2</sup> 5p <sup>6</sup> 5d <sup>10</sup> 6s <sup>2</sup> 6p <sup>4</sup> +	Observed Terms		
7s	a <sup>1</sup> S		
6d	a <sup>3</sup> P    b <sup>1</sup> D    c <sup>3</sup> F    a <sup>1</sup> G		
	ns (n ≥ 7)	np (n ≥ 7)	nf (n ≥ 5)
7s(8)ns	{ a <sup>3</sup> S a <sup>1</sup> S	{ s <sup>3</sup> P° s <sup>1</sup> P°	{ y <sup>3</sup> F° s <sup>1</sup> F°
6d(1D)ns	{ a <sup>1</sup> D a <sup>3</sup> D	{ y <sup>3</sup> P°    s <sup>3</sup> D°    s <sup>3</sup> F° y <sup>1</sup> P°    s <sup>1</sup> D°    y <sup>1</sup> F°	{ z <sup>3</sup> P°    y <sup>3</sup> D°    z <sup>3</sup> F°    s <sup>3</sup> G°    s <sup>3</sup> H° z <sup>1</sup> P°    y <sup>1</sup> D°    z <sup>1</sup> F°    s <sup>1</sup> G°
5f(2F°)ns	{	{ e <sup>3</sup> D    e <sup>3</sup> F    e <sup>3</sup> G e <sup>1</sup> D    e <sup>1</sup> F    e <sup>1</sup> G	

\*For predicted terms in the spectra of the Ra I isoelectronic sequence, see Vol. III, Introduction.

## Ac III

(Fr I sequence; 87 electrons)

Z=89

Ground state 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>10</sup> 4s<sup>2</sup> 4p<sup>6</sup> 4d<sup>10</sup> 4f<sup>14</sup> 5s<sup>2</sup> 5p<sup>6</sup> 5d<sup>10</sup> 6s<sup>2</sup> 6p<sup>6</sup> 7s<sup>2</sup> S<sub>0,1</sub>7s<sup>2</sup> S<sub>0,1</sub>

K

I. P. volts

The analysis is by Meggers, who first detected strong Ac III lines on spark spectrograms made at the Argonne laboratory by Fred and Tomkins. This work has been done especially for inclusion here, and the results have been furnished in advance of publication.

Meggers has ascribed eight lines to Ac III, in the range 2626.44 Å to 5193.21 Å, because they appeared in the arcs only near the cathode, and were greatly enhanced and hazy in the spark. He reports that two lines exhibit hyperfine components having a separation of 1.73 K, which led him to conclude that they represent the combination from the ground state 7s<sup>2</sup> S to 7p<sup>2</sup> P°. Further observations over a longer spectral range are needed to extend the analysis.

By analogy with Ra II and Th IV Lang estimates an ionization potential near 20 volts.

## REFERENCES

R. J. Lang, Canadian J. Research [A] 14, 46 (1936). (I P).

W. F. Meggers, M. Fred, and F. S. Tomkins, J. Opt. Soc. Am. 41, 867 (A) (1951).

W. F. Meggers, M. Fred, and F. S. Tomkins, J. Research Nat. Bur. Std. 58, 297, RP2763 (1957). (T) (C L) (his).

## Ac III

## Ac III

Config.	Desig.	J	Level	Interval	Config.	Desig.	J	Level	Interval
6p <sup>4</sup> (1S)7s	7s <sup>2</sup> S	0½	0.0		6p <sup>4</sup> (1S)5f	5f <sup>2</sup> F°	2½ 3½	23454.5 23080.2	2625.7
6p <sup>4</sup> (1S)6d	6d <sup>2</sup> D	1½ 2½	801.0 4203.9	3402.9	6p <sup>4</sup> (1S)7p	7p <sup>2</sup> P°	0½ 1½	29465.9 38063.0	8597.1

June 1957.

Page	Spectrum	Remarks																																																																																																																														
1	H I	Add references: C. J. Humphreys, <i>J. Research Nat. Bur. Std.</i> <b>50</b> , 1, RP 2380 (1953) (T) (CL); Sixth series and other series members in far infrared. W. E. Lamb, Jr. and T. M. Sanders, Jr., <i>Phys. Rev.</i> [2] <b>103</b> , 313 (1956); Fine structure of $n=3$ .																																																																																																																														
3	D	Add reference: G. Herzberg, <i>Proc. Roy. Soc. London [A]</i> <b>234</b> , 516 (1956); Fine structure.																																																																																																																														
4	He I	Add references: W. V. Houston, <i>Proc. Nat. Acad. Sci.</i> <b>13</b> , 91 (1927); Fine structure. C. J. Humphreys and H. J. Kostkowski, <i>J. Research Nat. Bur. Std.</i> <b>49</b> , 73, RP 2345 (1952) (T) (CL); Revised term values from infrared observations. G. Herzberg, <i>Am. Assn. Adv. Sci. Symposium (December 1957) (I P)</i> ; Limit $198310.4 \pm 0.3$ ; I. P. 24.581. T. H. Maiman and W. E. Lamb, Jr., <i>Phys. Rev.</i> <b>98</b> , 1194 (A) (1955); Fine structure. C. J. Humphreys and E. Paul, Jr., U. S. Naval Ord Lab., NAVORD Report 4589, 25 (1956) (T) (CL); Infrared observations. A. G. Shenstone, unpublished material (1957); Revised level values based on recent observations:																																																																																																																														
		<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="3">He I</th> <th colspan="3">He I</th> </tr> <tr> <th>Desig.</th> <th><i>J</i></th> <th>Level</th> <th>Desig.</th> <th><i>J</i></th> <th>Level</th> </tr> </thead> <tbody> <tr><td>8s <sup>2</sup>S</td><td>1</td><td>196455.68</td><td>13s <sup>2</sup>S</td><td>1</td><td>197625.00</td></tr> <tr><td>8s <sup>1</sup>S</td><td>0</td><td>196529.03</td><td>13s <sup>1</sup>S</td><td>0</td><td>197641.53</td></tr> <tr><td>8d <sup>3</sup>D</td><td>3, 2, 1</td><td>196589.26</td><td>13d <sup>1</sup>D</td><td>2</td><td>197655.34</td></tr> <tr><td>8d <sup>1</sup>D</td><td>2</td><td>196589.73</td><td>13d <sup>3</sup>D</td><td>3, 2, 1</td><td>197655.38</td></tr> <tr><td>9s <sup>2</sup>S</td><td>1</td><td>196856.29</td><td>14s <sup>2</sup>S</td><td>1</td><td>197720.62</td></tr> <tr><td>9s <sup>1</sup>S</td><td>0</td><td>196907.13</td><td>14s <sup>1</sup>S</td><td>0</td><td>197733.82</td></tr> <tr><td>9d <sup>3</sup>D</td><td>3, 2, 1</td><td>196949.41</td><td>14d <sup>1</sup>D</td><td>2</td><td>197744.84</td></tr> <tr><td>9d <sup>1</sup>D</td><td>2</td><td>196949.63</td><td>14d <sup>3</sup>D</td><td>3, 2, 1</td><td>197744.86</td></tr> <tr><td>10s <sup>2</sup>S</td><td>1</td><td>197139.53</td><td>15s <sup>2</sup>S</td><td>1</td><td>197797.37</td></tr> <tr><td>10s <sup>1</sup>S</td><td>0</td><td>197176.32</td><td>15s <sup>1</sup>S</td><td>0</td><td>197808.10</td></tr> <tr><td>10d <sup>3</sup>D</td><td>3, 2, 1</td><td>197207.05</td><td>15d <sup>1</sup>D</td><td>2</td><td>197817.01</td></tr> <tr><td>10d <sup>1</sup>D</td><td>2</td><td>197207.22</td><td>15d <sup>3</sup>D</td><td>3, 2, 1</td><td>197817.08</td></tr> <tr><td>11s <sup>2</sup>S</td><td>1</td><td>197347.13</td><td>16s <sup>2</sup>S</td><td>1</td><td>197860.11</td></tr> <tr><td>11d <sup>3</sup>D</td><td>3, 2, 1</td><td>197397.63</td><td>16d <sup>1</sup>D</td><td>2</td><td>197875.87</td></tr> <tr><td>11d <sup>1</sup>D</td><td>2</td><td>197397.84</td><td>16d <sup>3</sup>D</td><td>3, 2, 1</td><td>197876.17</td></tr> <tr><td>12s <sup>2</sup>S</td><td>1</td><td>197503.77</td><td>17s <sup>2</sup>S</td><td>1</td><td>197911.78</td></tr> <tr><td>12s <sup>1</sup>S</td><td>0</td><td>197524.83</td><td>17d <sup>1</sup>D</td><td>3, 2, 1</td><td>197925.16</td></tr> <tr><td>12d <sup>3</sup>D</td><td>3, 2, 1</td><td>197542.55</td><td>18d <sup>1</sup>D</td><td>3, 2, 1</td><td>197966.47</td></tr> <tr><td>12d <sup>1</sup>D</td><td>2</td><td>197542.69</td><td>19d <sup>1</sup>D</td><td>3, 2, 1</td><td>198001.27</td></tr> </tbody> </table>	He I			He I			Desig.	<i>J</i>	Level	Desig.	<i>J</i>	Level	8s <sup>2</sup> S	1	196455.68	13s <sup>2</sup> S	1	197625.00	8s <sup>1</sup> S	0	196529.03	13s <sup>1</sup> S	0	197641.53	8d <sup>3</sup> D	3, 2, 1	196589.26	13d <sup>1</sup> D	2	197655.34	8d <sup>1</sup> D	2	196589.73	13d <sup>3</sup> D	3, 2, 1	197655.38	9s <sup>2</sup> S	1	196856.29	14s <sup>2</sup> S	1	197720.62	9s <sup>1</sup> S	0	196907.13	14s <sup>1</sup> S	0	197733.82	9d <sup>3</sup> D	3, 2, 1	196949.41	14d <sup>1</sup> D	2	197744.84	9d <sup>1</sup> D	2	196949.63	14d <sup>3</sup> D	3, 2, 1	197744.86	10s <sup>2</sup> S	1	197139.53	15s <sup>2</sup> S	1	197797.37	10s <sup>1</sup> S	0	197176.32	15s <sup>1</sup> S	0	197808.10	10d <sup>3</sup> D	3, 2, 1	197207.05	15d <sup>1</sup> D	2	197817.01	10d <sup>1</sup> D	2	197207.22	15d <sup>3</sup> D	3, 2, 1	197817.08	11s <sup>2</sup> S	1	197347.13	16s <sup>2</sup> S	1	197860.11	11d <sup>3</sup> D	3, 2, 1	197397.63	16d <sup>1</sup> D	2	197875.87	11d <sup>1</sup> D	2	197397.84	16d <sup>3</sup> D	3, 2, 1	197876.17	12s <sup>2</sup> S	1	197503.77	17s <sup>2</sup> S	1	197911.78	12s <sup>1</sup> S	0	197524.83	17d <sup>1</sup> D	3, 2, 1	197925.16	12d <sup>3</sup> D	3, 2, 1	197542.55	18d <sup>1</sup> D	3, 2, 1	197966.47	12d <sup>1</sup> D	2	197542.69	19d <sup>1</sup> D	3, 2, 1	198001.27
He I			He I																																																																																																																													
Desig.	<i>J</i>	Level	Desig.	<i>J</i>	Level																																																																																																																											
8s <sup>2</sup> S	1	196455.68	13s <sup>2</sup> S	1	197625.00																																																																																																																											
8s <sup>1</sup> S	0	196529.03	13s <sup>1</sup> S	0	197641.53																																																																																																																											
8d <sup>3</sup> D	3, 2, 1	196589.26	13d <sup>1</sup> D	2	197655.34																																																																																																																											
8d <sup>1</sup> D	2	196589.73	13d <sup>3</sup> D	3, 2, 1	197655.38																																																																																																																											
9s <sup>2</sup> S	1	196856.29	14s <sup>2</sup> S	1	197720.62																																																																																																																											
9s <sup>1</sup> S	0	196907.13	14s <sup>1</sup> S	0	197733.82																																																																																																																											
9d <sup>3</sup> D	3, 2, 1	196949.41	14d <sup>1</sup> D	2	197744.84																																																																																																																											
9d <sup>1</sup> D	2	196949.63	14d <sup>3</sup> D	3, 2, 1	197744.86																																																																																																																											
10s <sup>2</sup> S	1	197139.53	15s <sup>2</sup> S	1	197797.37																																																																																																																											
10s <sup>1</sup> S	0	197176.32	15s <sup>1</sup> S	0	197808.10																																																																																																																											
10d <sup>3</sup> D	3, 2, 1	197207.05	15d <sup>1</sup> D	2	197817.01																																																																																																																											
10d <sup>1</sup> D	2	197207.22	15d <sup>3</sup> D	3, 2, 1	197817.08																																																																																																																											
11s <sup>2</sup> S	1	197347.13	16s <sup>2</sup> S	1	197860.11																																																																																																																											
11d <sup>3</sup> D	3, 2, 1	197397.63	16d <sup>1</sup> D	2	197875.87																																																																																																																											
11d <sup>1</sup> D	2	197397.84	16d <sup>3</sup> D	3, 2, 1	197876.17																																																																																																																											
12s <sup>2</sup> S	1	197503.77	17s <sup>2</sup> S	1	197911.78																																																																																																																											
12s <sup>1</sup> S	0	197524.83	17d <sup>1</sup> D	3, 2, 1	197925.16																																																																																																																											
12d <sup>3</sup> D	3, 2, 1	197542.55	18d <sup>1</sup> D	3, 2, 1	197966.47																																																																																																																											
12d <sup>1</sup> D	2	197542.69	19d <sup>1</sup> D	3, 2, 1	198001.27																																																																																																																											
		July 1947.																																																																																																																														
6	He II	J. E. Mack (letter, December 1953) states that the term 15s <sup>2</sup> S, etc. should read 436958.026, etc., instead of 436957.026, etc. Add references: R. Wilson, <i>Mon. Not. Roy. Astron. Soc.</i> <b>113</b> , 557 (1953); Series members 5g <sup>2</sup> G— $n$ h <sup>2</sup> H <sup>o</sup> ( $n=13$ to 34). G. W. Series, <i>Proc. Roy. Soc. London [A]</i> <b>226</b> , 377 (1954); Fine structure.																																																																																																																														
8	Li I	Add reference: K. Lidén and N. Starfelt, <i>Ark. Fys.</i> <b>5</b> , 127 (1952) (I P).																																																																																																																														
10	Li II	2s <sup>1</sup> S should read 491361 instead of [490079]. Add reference: H. Schüler, <i>Zeit. Phys.</i> <b>66</b> , 431 (1930); Fine structure.																																																																																																																														

\*At the end of Volume II a list of corrections to Volume I was appended. These are not repeated, but additional revisions are given.

Page	Spectrum	Remarks
12	Be I	Add reference: W. R. Bozman, C. H. Corliss, W. F. Meggers and R. E. Trees, J. Research Nat. Bur. Std. <b>50</b> , 131, RP 2399 (1953) (T) (C L); Observed intersystem combination gives $x = -1.18$ .
14	Be III	Add reference: B. Edlén, Ark. Fys. <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
16	B I	Add references: H. E. Clearman, J. Opt. Soc. Am. <b>42</b> , 373 (1952) (T) (C L). E. W. Burke, Jr. and J. E. Mack, J. Opt. Soc. Am. <b>46</b> , 100 (1956) (T).
19	B IV	Add reference: B. Edlén, Ark Fys. <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
21	C I	Limit should read <b>90814</b> , I. P. 11.256. Add references: B. Edlén, Accad. Naz. Lincei Volta Conv. (Rome) <b>11</b> , 58 (1952) and unpublished material (September 1952); Wavelengths and revised term values. L. Minnhagen, Ark. Fys. <b>7</b> , No. 33, 413 (1954) (T) (C L).
24	C II	Add reference: S. Glad, Ark. Fys. <b>7</b> , No. 2, 7 (1954) (I P) (T) (C L); Revised analysis.
26	C III	Add reference: K. Bockasten, Ark. Fys. <b>9</b> , No. 30, 457 (1955) (I P) (T) (C L); Revised analysis: Revised limit is <b>386213.9</b> , I. P. 47.871; $x = +51.7$ .
29	C IV	Add reference: K. Bockasten, Ark Fys. <b>10</b> , No. 40, 567 (1956) (I P) (T) (C L); Revised analysis.
30	C V	Add reference: B. Edlén, Ark. Fys. <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
32	N I	The term $3s' ^2D$ is unresolved and should read 99663. The lack of resolution has been pointed out by J. C. Boyce in private conversation. Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): $2p^3 ^2D_{2\frac{3}{2}}$ should read <b>19223.9</b> $^2D_{1\frac{3}{2}}$ should read <b>19233.1</b> Limit should read <b>117214</b> , I. P. 14.53.
35	N II	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955). (C L) K. B. Eriksson, unpublished material reported by B. Edlén (March 1957); Revised analysis: Revised limit is <b>238751.1 ± 1</b> , I. P. 29.593; $x = -382.6$ . $4p ^1S_0$ should read 206911.0.
43	N VI	Add reference: B. Edlén, Ark Fys. <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
45	O I	Add references: D. O. Davis and K. W. Meissner, J. Opt. Soc. Am. <b>43</b> , 510 (1953) Fine structure. G. Kvitte, Ark. Mat. o Naturvidenskab (Oslo) [B] <b>52</b> , 65 (1954) (T) (C L); Series extended. I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L).
47	O II	Limit should read <b>233244</b> , I. P. 35.108. Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L).
50	O III	Limit should read <b>442807</b> , I. P. 54.886. Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L).
59	O VII	Add reference: B. Edlén, Ark. Fys. <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
66	F IV	Limit should read <b>703020</b> , I. P. 87.14. Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L).
75	F VIII	Add reference: B. Edlén, Ark. Fys. <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
83	Ne III	Add references: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): $2p^4 ^3P_1$ should read <b>642.9</b> $2p^4 ^1D_2$ should read <b>25840.8</b> W. Finkelnburg und W. Humbach, Naturwiss. <b>42</b> , 35 (1955) (I P): Interpolated ionization potential $63.5 ± 0.1$ .
84	Ne IV	Limit should read <b>782768</b> , I. P. 97.02. Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): $2p^3 ^2D_{2\frac{3}{2}}$ reads <b>40950 + x</b> -44.8 $^2D_{1\frac{3}{2}}$ should read <b>40996 + x</b> $2p^3 ^3P_{0\frac{3}{2}}$ should read <b>62150 + x</b> 6.4 $^3P_{1\frac{3}{2}}$ should read <b>62157 + x</b> $2p^4 ^2D_{2\frac{3}{2}}$ should read <b>253799 + x</b> -23 $^2D_{1\frac{3}{2}}$ should read <b>253822 + x</b> Other doublet terms should be revised accordingly.

\*At the end of Volume II a list of corrections to Volume I was appended. These are not repeated, but additional revisions are given.

Page	Spectrum	Remarks
86	Ne v	Limit should read 1018634, I. P. 126.3.
		Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L).
89	Na I	Add reference: P. Risberg, <i>Ark. Fys.</i> <b>10</b> , No. 41, 583 (1956) (I P) (T) (C L) (G D); Revised analysis.
96	Na v	Limit should read 116312, I. P. 138.37.
98	Na VI	Limit should read 1388419, I. P. 172.09.
106	Mg I	Add reference: R. A. Fisher and F. E. Eshbach, <i>J. Opt. Soc. Am.</i> <b>43</b> , 1030 (1953) (T) (C L); Revised term values from near infrared observations.
108	Mg II	Add reference: P. Risberg, <i>Ark. Fys.</i> <b>9</b> , No. 31, 483 (1955) (I P) (T) (C L) (G D); Revised analysis.
114	Mg VI	Limit should read 1504581, I. P. 186.49.
117	Mg VII	Limit should read 1814430, I. P. 224.90.
123	Mg XI	Add reference: B. Edlén, <i>Ark. Fys.</i> <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
135	Al VII	Limit should read 1947390, I. P. 241.38.
136	Al VIII	Limit should read 2295500, I. P. 284.53.
143	Al XII	Add reference: B. Edlén, <i>Ark. Fys.</i> <b>4</b> , No. 28, 441 (1952) (I P) (T) (C L); Revised analysis.
144	Si I	A. G. Shenstone, private communication (October 1957) reports following new terms: $3s 3p^2$ $^4S_2^o$ 33326.16  $3s 3p^2(^4P)4s$ $^4P_1$ 81724.8                      99.5 $^4P_2$ 81824.3                      152.2 $^4P_3$ 81976.5  $3s 3p^2(^4P)3d$ $^4P_3$ 94292.3                      -73.8 $^4P_2$ 94366.1                      -47.7 $^4P_1$ 94413.8  He suggests also, that the term designated $3p^2 ^3D^o$ (48399.15 etc.) be deleted, and that the series terms $nd ^3D^o$ ( $n=4$ to 8) have $n$ decreased by 1, ( $n=3$ to 7), since the lowest $^3D^o$ term has the configuration $3s 3p^2$ .  Add references: M. A. El'yashevich and O. N. Nikitina, <i>Dokl. Akad. Nauk SSSR</i> , <b>111</b> , No. 2, 325 (1956); <i>Phys. Abstr.</i> <b>60</b> , No. 713, 383 (1957) (T) (C L); Report $3s 3p^2 ^4S_2^o$ 33326.28. C. C. Kiess (July 1957), confirms the term $3s 3p^2 ^4S_2^o$ from two Si I lines he has measured at 3020.020A (20) and 3006.738A (10). The term designated $3d ^3D^o$ (45276.20 etc.) should have the configuration $3s 3p^2$ .  A. G. Shenstone and S. L. Blatt, letter (Dec. 1957) report the following terms: $3p ^2P_{0\frac{1}{2}}$ 0.00                      287.45 $4p ^4P_{0\frac{1}{2}}$ 134017.4                      62.3 $^2P_{1\frac{1}{2}}$ 287.45 $^4P_{1\frac{1}{2}}$ 134079.7                      134.3 $^4P_{2\frac{1}{2}}$ 134214.0  $3p^2 ^4P_{0\frac{1}{2}}$ 42824.39                      108.52 $^4P_{1\frac{1}{2}}$ 42932.91                      175.17 $^4P_{2\frac{1}{2}}$ 43108.08 $4p ^4S_{1\frac{1}{2}}$ 134906.0  $x$ is, therefore, approximately -1257 and the values of the quartet terms should be revised to fit the above data.
148	Si III	B. Edlén, letter (July 1956) notes that the $3p^2 ^1D$ and $3d ^1D$ terms are dubious.
156	Si VIII	Limit should read 2445110, I. P. 303.07.
157	Si IX	Limit should read 2831470, I. P. 350.96.
163	P I	Add reference: W. C. Martin, Dissertation, Princeton University (June 1956) (I P) (T) (C L); Revised analysis: Limit 84580, I. P. 10.484.
164	P II	Add reference: W. C. Martin, Dissertation, Princeton University (June 1956); Revised analysis: Limit 159100, I. P. 19.72; $x = -6753.2$ .

\*At the end of Volume II a list of corrections to Volume I was appended. These are not repeated, but additional revisions are given.

Page	Spectrum	Remarks
166	P III	Add reference: W. C. Martin, unpublished material (February 1957); Revised wavelengths and term values; $4d^2D_{3/2}$ should read 172428.63 $^2D_{3/2}$ should read 172429.40      -0.77  Add $4p^4D_{3/2}$ 207036.2                      325.4 $^4D_{3/2}$ 207361.6
168	P IV	W. C. Martin, letter (July 1957); Revised analysis: delete $3d^1D^0$ and $5p^1P^0$ .
174	P IX	Limit should read <del>3997600</del> , I. P. 371.6.
176	P X	Limit should read <del>3423000</del> , I. P. 424.3.
181	S I	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L).
183	S II	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L): $3p^3^2D_{3/2}^1$ should read 14853.0                      31.8 $^2D_{3/2}^1$ should read 14884.8  $3p^3^2P_{3/2}^0$ should read <del>24524.9</del> 46.7 $^2P_{3/2}^0$ should read 24571.6
185	S III	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L).
194	S X	Limit should read <del>3666000?</del> , I. P. 447.07.
195	Cl I	Add reference: S. Avellán, <i>Ark. Fys.</i> <b>8</b> , No. 3, 211 (1954) (I P) (T) (C L): $3p^3^1P_{3/2}^0$ should read <del>882.50</del> ; Limit should read <del>104995.46</del> .
199	Cl III	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L): $3p^3^2D_{3/2}^0$ should read 18118.6.
201	Cl IV	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L).
210	Cl XI	Add reference: W. Finkelnburg und W. Humbach, <i>Naturwiss.</i> <b>42</b> , 35 (1955) (I P): From screening constants, interpolated I. P. is $530.9 \pm 0.2$ .
211	Ar I	Add references: C. J. Humphreys and H. J. Kostkowski, <i>J. Research Nat. Bur. Std.</i> <b>49</b> , 73, RP 2345 (1952) (C L); Infrared observations. T. A. Littlefield and D. T. Turnbull, <i>Proc. Roy. Soc. London [A]</i> <b>218</b> , 577 (1953) (T) (C L). K. Burns and K. B. Adams, <i>J. Opt. Soc. Am.</i> <b>43</b> , 1020 (1953) (T) (C L); Revised level values. G. H. Dieke and H. M. Crosswhite, <i>Ordnance Project No. TB 2-0001 (488)</i> 38 pp. (1954) (C L); Bibliography, wavelength list, intensities. C. J. Humphreys and E. Paul, Jr., <i>U. S. Naval Ord. Lab., NAVORD Report 4589</i> , 40 (1956) (T) (C L); Infrared observations.
218	Ar III	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L): $3p^4^1P_1$ should read 1112.1 $3p^4^1D_2$ should read 14010.0 $3p^4^1S_0$ should read 33265.7
220	Ar IV	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L): $3p^3^2D_{3/2}^0$ should read <del>21090.3</del> 129.2 $^2D_{3/2}^0$ should read <del>21219.5</del>  $3p^3^1P_{3/2}^0$ should read <del>34855.4</del> 177.3 $^1P_{3/2}^0$ should read <del>35032.7</del>
222	Ar V	Add reference: I. S. Bowen, <i>Astroph. J.</i> <b>121</b> , 306 (1955) (C L): new term $3p^3^1S_0$ 37914.
225	Ar IX	Add reference: W. Finkelnburg und W. Humbach, <i>Naturwiss.</i> <b>42</b> , 35 (1955) (I P): From screening constants interpolated I. P. is $422.6 \pm 0.2$ .
226	Ar XIV	Add reference: B. Edlén, <i>Mon. Not. Roy. Astron. Soc.</i> <b>114</b> , 700 (1954) (T); Interval of ground term of A XIV should read 22657, derived from the coronal line observed at $\lambda 4412.4A$ , which replaces $\lambda 4359$ previously identified as [A XIV].

\*At the end of Volume II a list of corrections to Volume I was appended. These are not repeated, but additional revisions are given.

Page	Spectrum	Remarks												
227	K I	Add reference: P. Risberg, Ark. Fys. <b>10</b> , No. 41, 594 (1956) (I P) (T) (C L) (G D); Revised analysis.												
233	K IV	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): 3p <sup>4</sup> <sup>1</sup> D <sub>2</sub> should read 16384.0 3p <sup>4</sup> <sup>1</sup> S <sub>2</sub> should read 38546												
234	K V	Add references: W. Finkelnburg und W. Humbach, Naturwiss. <b>42</b> , 35 (1955) (I P): From screening constants, interpolated I. P. is 82.6±0.4. I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): 3p <sup>3</sup> <sup>2</sup> D <sub>1,2</sub> <sup>o</sup> should read 24012.7      236.8 <sup>2</sup> D <sub>3,4</sub> <sup>o</sup> should read 24249.5												
245	Ca II	Add reference: B. Edlén and P. Risberg, Ark. Fys. <b>10</b> , No. 39, 553 (1956) (I P) (T) (C L) (G D); Revised analysis.												
249	Ca V	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): 3p <sup>4</sup> <sup>3</sup> P <sub>1</sub> should read 2406.0 3p <sup>4</sup> <sup>1</sup> D <sub>2</sub> should read 18830.1												
251	Ca VI	Add reference: W. Finkelnburg und W. Humbach, Naturwiss. <b>42</b> , 35 (1955) (I P): From screening constants, interpolated I. P. is 109±1.												
253	Ca VIII	Add reference: W. Finkelnburg und W. Humbach, Naturwiss. <b>42</b> , 35 (1955) (I P): From screening constants, interpolated I. P. is 143.3±0.4.												
258	Ca XV	Add reference: B. Edlén, Mon. Not. Roy. Astron. Soc. <b>114</b> , 700 (1954) (T): From coronal observations, ground term of Ca xv is: <table style="margin-left: 40px;"> <tr> <td>2p<sup>3</sup> <sup>3</sup>P</td> <td>0</td> <td>0</td> <td>17556</td> </tr> <tr> <td></td> <td>1</td> <td>17556</td> <td>18355</td> </tr> <tr> <td></td> <td>2</td> <td>35911</td> <td></td> </tr> </table>	2p <sup>3</sup> <sup>3</sup> P	0	0	17556		1	17556	18355		2	35911	
2p <sup>3</sup> <sup>3</sup> P	0	0	17556											
	1	17556	18355											
	2	35911												
259	Sc I	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] <b>48</b> , 247 (1952) (I P): Revised limit 52750, I. P. 6.54.												
263	Sc III	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] <b>48</b> , 247 (1952) (I P).												
267	Sc VII	Add reference: W. Finkelnburg und W. Humbach, Naturwiss. <b>42</b> , 35 (1955) (I P): From screening constants, interpolated I. P. is 139±1.												
268	Sc VIII	I. S. Bowen in conversation (July 1953) has noted that the analysis is incorrect.												
273	Ti I	Add references: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] <b>48</b> , 247 (1952) (I P): Revised limit 55010, I. P. 6.82. A. K. Wardakee, J. Opt. Soc. Am. <b>45</b> , 354 (1955) (C L).												
281	Ti III	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] <b>48</b> , 247 (1952) (I P). The <i>J</i> -value for <sup>2</sup> S should read "1".												
287	Ti VIII	I. S. Bowen in conversation (July 1953) has noted that the analysis is incorrect. Add reference: W. Finkelnburg u W. Humbach, Naturwiss. <b>42</b> , 35 (1955) (I P): From screening constants, interpolated I. P. is 172±1.												
288	Ti X	Add reference: W. Finkelnburg u W. Humbach, Naturwiss. <b>42</b> , 35 (1955) (I P): From screening constants, interpolated I. P. is 216.9±0.6.												
301	V III	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp [A] <b>48</b> , 247 (1952) (I P).												
XXXVI	Sc I sequence	For configurations having F-limit terms, under " <i>n</i> " add S <sup>o</sup> and P <sup>o</sup> as predicted terms; for those having G-limit terms, add P <sup>o</sup> and D <sup>o</sup> as predicted terms.												

\*At the end of Volume II a list of corrections to Volume I was appended. These are not repeated, but additional revisions are given.

## NOTE ADDED IN PROOF

- |     |       |   |
|-----|-------|---|
| 124 | A1/i  | B. Edlén (letter, February, 1958) states that the term 38929.42 etc. designated 4d <sup>1</sup> D should be assigned to 3s 3p <sup>2</sup> , and, consequently, the <i>n</i> -values for all higher <i>nd</i> <sup>1</sup> D terms should be decreased by one unit. |
| 266 | Sc VI | Add reference: B. Edlén, Phys. Rev. <b>63</b> , 434 (1942) (T): 3s 3p <sup>4</sup> <sup>1</sup> P <sub>1</sub> 224474.  |

Page	Spectrum	Remarks																																				
1	Cr I	Add references: C. C. Kiess, J. Research Nat. Bur. Std. <b>51</b> , 247, RP 2457 (1953) (I P) (T) (C L) (Z E). P. Brix, J. T. Eisinger, H. Lew, and G. Wessel, Phys. Rev. [2] <b>92</b> , 647 (1953) (Z E).																																				
10	Cr II	In the array of observed terms the second entry under "Config." should read $3d^2 4s^2$ instead of $3d^3 4s$ . G. Racah (1953) has suggested that the configuration of the term $c^4P$ should read $3d^4(b^4P)4s$ instead of $3d^3 4s^2$ .																																				
14	Cr III	Add references: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] <b>48</b> , 247 (1952) (I P). F. L. Moore, Jr. (Thesis, Princeton 1949), Univ. Microfilm Publ. No. 10, 972, 172 pp.; Dissertation Abstract (Ann Arbor, Mich.) <b>15</b> , 432 (1955) (I P) (T) (C L).																																				
27	Mn I	Add references: M. A. Catalán, J. Research Nat. Bur. Std. <b>47</b> , 502, RP 2278 (1951) (Z E). M. L. Espinosa, An. Real. Soc. Esp. [A] <b>48</b> , 223 (1952) (Z E). M. A. Catalán (1953) in private conversation has suggested that the term $c^4F$ is somewhat dubious.																																				
32	Mn II	Add references: R. E. Trees, Phys. Rev. [2] <b>83</b> , 756 (1951): Notes that the configurations of $a^4G$ and $b^4G$ as published in "AEL" should be interchanged in accordance with Curtis' 1952 paper. O. García-Riquelme, L. Iglesias, y R. Velasco, An. Real. Soc. Esp. [A] <b>53</b> , 77 (1957) (I P) (T) (C L). L. Iglesias, J. Opt. Soc. Am. <b>46</b> , 449 (1956); <b>47</b> , 852 (1957) (T) (C L) (E D).																																				
40	Mn v	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L).																																				
41	Mn vi	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L).																																				
49	Fe I	Add references: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] <b>48</b> , 247 (1952) (I P). B. Edlén, Trans. Intern. Astron. Union <b>9</b> , 219 (1955) (T); Revised term values. J. Blackie and T. A. Littlefield, Proc. Roy. Soc. London [A] <b>234</b> , 398 (1956) (C L).																																				
55	Fe II	Add reference: M. Sales, An. Real. Soc. Esp. [A] <b>49</b> , 15 (1953) (T) (C L) (E D).																																				
60	Fe III	Add references: R. E. Trees, Phys. Rev. [2] <b>84</b> , 1089 (1951) (T). I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L). S. Glad, Ark. Fys. <b>10</b> , No. 22, 291 (1956) (I P) (T) (C L) (E D); Extension of analysis.																																				
66	Fe v	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L).																																				
67	Fe vi	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): The following terms should read:																																				
		<table border="0"> <tr> <td><math>a^4F</math></td> <td>1½</td> <td>0.0</td> <td></td> </tr> <tr> <td></td> <td>2½</td> <td>510.9</td> <td>510.9</td> </tr> <tr> <td></td> <td>3½</td> <td>1189</td> <td>678</td> </tr> <tr> <td></td> <td>4½</td> <td>2002</td> <td>813</td> </tr> <tr> <td><math>a^4P</math></td> <td>0½</td> <td>18738.1</td> <td></td> </tr> <tr> <td></td> <td>1½</td> <td>18943</td> <td>205</td> </tr> <tr> <td></td> <td>2½</td> <td>19612</td> <td>669</td> </tr> <tr> <td><math>a^4G</math></td> <td>3½</td> <td>20617</td> <td></td> </tr> <tr> <td></td> <td>4½</td> <td>21315</td> <td>698</td> </tr> </table>	$a^4F$	1½	0.0			2½	510.9	510.9		3½	1189	678		4½	2002	813	$a^4P$	0½	18738.1			1½	18943	205		2½	19612	669	$a^4G$	3½	20617			4½	21315	698
$a^4F$	1½	0.0																																				
	2½	510.9	510.9																																			
	3½	1189	678																																			
	4½	2002	813																																			
$a^4P$	0½	18738.1																																				
	1½	18943	205																																			
	2½	19612	669																																			
$a^4G$	3½	20617																																				
	4½	21315	698																																			
69	Fe vii	Add reference: I. S. Bowen, Astroph. J. <b>121</b> , 306 (1955) (C L): The following terms should read:																																				
		<table border="0"> <tr> <td><math>a^4F</math></td> <td>2</td> <td>0.0</td> <td></td> </tr> <tr> <td></td> <td>3</td> <td>1050.2</td> <td>1050.2</td> </tr> <tr> <td></td> <td>4</td> <td>2326</td> <td>1276</td> </tr> <tr> <td><math>a^4D</math></td> <td>2</td> <td>17474.3</td> <td></td> </tr> <tr> <td><math>a^4P</math></td> <td>1</td> <td>20428.5</td> <td></td> </tr> <tr> <td></td> <td>2</td> <td>21270</td> <td>842</td> </tr> </table>	$a^4F$	2	0.0			3	1050.2	1050.2		4	2326	1276	$a^4D$	2	17474.3		$a^4P$	1	20428.5			2	21270	842												
$a^4F$	2	0.0																																				
	3	1050.2	1050.2																																			
	4	2326	1276																																			
$a^4D$	2	17474.3																																				
$a^4P$	1	20428.5																																				
	2	21270	842																																			

Page	Spectrum	Remarks
78	Co I	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] 48, 247 (1952) (I P): Limit should read 63430. The configuration assignments of $b^3D$ and $b^3G$ should be interchanged.
85	Co III	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] 48, 247 (1952) (I P).
103	Ni III	Add references: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] 48, 247 (1952) (I P). A. G. Shenstone, J. Opt. Soc. Am. 44, 749 (1954) (I P) (T) (C L); Revised analysis. O. García-Riquelme, An. Real. Soc. Esp. [A] 51, 59 (1955) (C L) (E D).
111	Cu I	Add reference: K. Murakawa, J. Phys. Soc. Japan 11, No. 7, 774 (1956) (hfs).
121	Cu III	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] 48, 247 (1952) (I P).
124	Zn I	Add reference: W. R. S. Garton and A. Rajaratnam, Proc. Phys. Soc. London [A] 68, 1107 (1955) (T) (C L): Suggests that the level at 80795 should have the designation $4p^3^3P_1$ .
128	Zn III	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] 48, 247 (1952) (I P).
130	Ga I	Add reference: W. R. S. Garton, Proc. Phys. Soc. London [A] 65, 268 (1952) (T) (C L).
133	Ga III	Add reference: M. A. Catalán y R. Velasco, An. Real. Soc. Esp. [A] 48, 247 (1952) (I P).
135	Ge I	Add references: G. V. Deverall, K. W. Meissner, and G. J. Zissis, Phys. Rev. [2] 96, 1463 (1954) (T) (C L); Precision measurements. R. D. Van Veld and K. W. Meissner, J. Opt. Soc. Am. 46, 598 (1956) (T) (C L). K. L. Andrew and K. W. Meissner, J. Opt. Soc. Am. 47, 850 (1957) (T) (C L): Report new term: $4s 4p^3^3S_2^o$ , 41926.726.
144	As II	A. M. Crooker and R. E. Bedford, unpublished material (December 1955) (I P) (T) (C L); Revised analysis: Limit should read 150290, I. P. 18.63.
146	As III	Add references: R. E. Bedford and A. M. Crooker, Phys. Rev. [2] 96, 845 (A) (1954) (T); unpublished material (1955) (I P) (T): Limit should read 228670, I. P. 28.34.
147	As IV	Add reference: R. E. Bedford and A. M. Crooker, Phys. Rev. [2] 96, 845 (A) (1954) (T).
156	Se V	Add reference: W. Finkelnburg and W. Humbach, Naturwiss. 42, 35 (1955) (I P): From screening constants, interpolated I. P. is $68.3 \pm 0.1$ .
159	Br I	J. L. Tech, unpublished material (July 1957) (T) (C L) (Z E): Reports new description and analysis of the spectrum being made at the National Bureau of Standards by J. L. Tech and C. H. Corliss.
163	Br III	Add reference: Y. B. Rao, Indian J. Phys. 30, 371 (1956) (T).
164	Br IV	Add reference: W. Finkelnburg und W. Humbach, Naturwiss. 42, 35 (1955) (I P): From screening constants, interpolated I. P. is 47.3.
165	Br V	Add reference: W. Finkelnburg und W. Humbach, Naturwiss. 42, 35 (1955) (I P): From screening constants, interpolated I. P. is $59.7 \pm 0.1$ .
166	Br VI	Add reference: W. Finkelnburg und W. Humbach, Naturwiss. 42, 35 (1955) (I P): From screening constants, interpolated I. P. is $88.6 \pm 0.2$ .
167	Br VII	Add reference: W. Finkelnburg und W. Humbach, Naturwiss. 42, 35 (1955) (I P): From screening constants, interpolated I. P. is $103.0 \pm 0.4$ .



Page	Spectrum	Remarks
169	Kr I	Add reference: C. J. Humphreys and E. Paul, Jr., U. S. Naval Ord. Lab., NAVORD Report 4600, 29 (1957) (T) (C L); Infrared observations. In 1952 ref., page no. 73 should be inserted. The level at 105989.60 should have the $J$ -values 4,5.
184	Rb II	Add reference: H. Kopfermann, A. Steudel, und J. O. Trier, Zeit. Phys. 144, 9 (1956) (hfs); Fine structure, corrections to analysis.
191	Sr II	The designations for the $^2G$ series should have $n$ increased by 1, i. e. $n=5$ to 11.
196	Y I	Add references: M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. [A] 48, 328 (1952) (I P); Revised limit 51447, I. P. 6.38. L. F. H. Bovey, Proc. Phys. Soc. London [A] 68, 79 (1955) (C L); Infrared observations.
199	Y II	Add references: M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. [A] 48, 328 (1952) (I P); Revised limit 98690, I. P. 12.23. L. F. H. Bovey, Proc. Phys. Soc. London [A] 68, 79 (1955) (C L); Infrared observations.
201	Y III	Add reference: M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. [A] 53, 85 (1957) (I P); Revised limit 165500, I. P. 20.51.
205	Zr I	Add reference: M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. [A] 48, 328 (1952) (I P); Revised limit 55145, I. P. 6.84. The level $^2G_3^o$ 36941.65 has Obs. $g=0.87$ .
209	Zr II	Add references: C. C. Kiess, J. Opt. Soc. Am. 43, 1024 (1953) (T) (C L); Extension of analysis from infrared observations; revises $4d^2 d^2D$ to read: $d^2D$ $2\frac{1}{2}$ 27640.60                      -59.36 $1\frac{1}{2}$ 27699.96 M. A. Catalán y F. R. Rico, letter (December 1956): revised limit should read 105900, I. P. 13.13.
212	Zr III	Add references: C. C. Kiess, J. Research Nat. Bur. Std. 56, 167, RP 2663 (1956) (I P) (T) (C L) (Z E); Revised analysis. M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. [A] 53, 85 (1957) (I P); Revised limit 185400, I. P. 22.98.
213	Zr IV	Add reference: C. C. Kiess, J. Research Nat. Bur. Std. 56, 167, RP 2663 (1956) (I P) (T) (C L) (G D); Revised analysis: Limit should read 276970, I. P. 34.33.
214	Zr VI	C. C. Kiess in conversation (September 1955) has reported the analysis incorrect.
216	Nb I	Add reference: M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. [A] 48, 328 (1952) (I P); Revised limit 55511, I. P. 6.88.
221	Nb II	Add reference: M. A. Catalán y F. R. Rico, letter (December 1956): Revised limit should read 115500, I. P. 14.32.
223	Nb III	Add references: L. Iglesias, An. Real. Soc. Esp. [A] 50, 135 (1954) (T) (C L); Analysis extended from ultraviolet observations. L. Iglesias, J. Opt. Soc. Am. 45, 856 (1955) (I P) (T) (C L); Revised analysis. M. A. Catalán y F. R. Rico, An. Real. Soc. Esp. [A] 53, 85 (1957) (I P); Revised limit 202000, I. P. 25.04.
xviii	Ni I sequence	For configurations having F-limit terms, under " $n_f$ " add $S^o$ and $P^o$ as predicted terms; for those having G-limit terms, add $P^o$ and $D^o$ as predicted terms. Similarly, under " $n_g$ ", for F-limit terms, add P and D as predicted terms; and for those having G-limit terms, add S, P, D, F as predicted terms.

NOTE ADDED IN PROOF

130	Ga I	B. Eldén (letter, February 1958) states that the $n$ -values for $n^1F^o$ terms should be increased as follows: 4 to 5, 5 to 7; 6 to 8; 7 to 9. The terms $4^1F^o$ and $6^1F^o$ are still missing.
202	Y v	B. Edlén (letter, March 1958) reports the analysis incorrect.