	ATION	PAGE	· · · · · · · · · · · · · · · · · · ·		Form Approved y
	4. 120		r Vong the time ton Nong timents ter Plunes timetorate		Simplify presiming wishing and structures stretching wishing and burgen estimate or any stret asta or Operations and Property 1/15
I ARMAN WAT ANDE MUNI KURI MINI KURI (III (III))	DATE	.ne huoget Par	PORT TYPE A	ND DATE	18E) Washington, IC 225C3 S COVERED
A. TITLE AND SUBT. TLEA RETROSPEC SUBLUHORF DE AIR FORMU TO HAZAR ON ENDISE DU	TIVE ST	NILE	TEXPOSE	SERTAT	IDING NUMBERS
6. AUTHOR(S)					\bigvee
Jim ALAN DAVIS,	B.S., D.	.V.M	1		
7. PERFORMING ORGANIZATION NAME(S) AND ADI	DRESS(ES)			B. PERI REP	FORMING ORGANIZATION ORT NUMBER
AFIT Student Attending:	20			AF1	T/C1/C1A
University of Ieki	AS			140	1-066
9. SPONSORING/MONITORING AGENCY NAME(S) A DEPARTMENT OF THE AIR FORCE AFIT/CI	ND ADDRESS(I	E S)		10. SPO AGE	NSORING/MONITORING NCY REPORT NUMBER
2950 P STREET WRIGHT-PATTERSON AFB OH 45433-	7765				
11. SUPPLEMENTARY NOTES			'depy	94-	22511
12a. DISTRIBUTION / AVAILABILITY STATEMENT			-	1	STRIBUTION COUL
Approved for Public Release I/ Distribution Unlimited MICHAEL M. BRICKER, SMSgt, US/	AW 190-1 AF				
LITET ADMINISTRATION 13. ABSTRACT (Maximum 200 words)				<u> </u>	·····
DTIC					
C ELECTE					
					·
and a strength			DTIC QUAL	ITY INS	
ř			•		
ج ۸ - (11016/7 TED-AC		<u></u>			
N.O.	ヤ	1 Q		1	16. PRICE CODE
7. SECURITY CLASSIFICATION 18. SECURITY CLA	SSIFICATION	19. SECU	RITY CLASSIFIC		20. LIMITATION OF ABSTRA
OF REPORT OF THIS PAGE		OF A	BSTRACT		
N 7540-01-280-5500		*		512	andard form 298 (Rev. 2-89

Prestriend by 1751 the plane

Best Available Copy

A RETROSPECTIVE STUDY OF A 1992 SUBCOHORT OF AIR FORCE PERSONNEL EXPOSED TO HAZARDOUS NOISE BY AIR FORCE SKILL CODE

? - > ... :

By

JIM ALAN DAVIS, B.S., D.V.M.

Accesi	on For	• •
NTIS	CRASI V	ہ۔۔۔ !
Und .	.0	
Justifi	cation	
By Distrib	uticit /	-1
4	vulability	· - 4
Dist	Avail and the Special	-1
A-1		

APPROVED:

Sharon P. Cooper, Ph.D.

Michael Decker, Ph.D., A.M.

Marcus M. Key, M.D., M.I.H.

DEDICATION

To my wife, Brenda, who has been my dearest and best friend for many years.

"A wife of noble character who can find? She is worth far more than rubies." Proverbs 31:10

8. 16. de 1

· ·

, , , ,

"Many women do noble things, but you surpass them all." Proverbs 31: 29

A RETROSPECTIVE STUDY OF A 1992 SUBCOHORT OF AIR FORCE PERSONNEL EXPOSED TO HAZARDOUS NOISE BY AIR FORCE SKILL CODE

, r , r

By

JIM ALAN DAVIS, B.S., D.V.M.

THESIS

Presented to the Faculty of The University of Texas

Health Science Center at Houston

School of Public Health

in Partial Fulfillment

of the Requirements

for the Degree of

MASTER OF PUBLIC HEALTH

THE UNIVERSITY OF TEXAS HEALTH SCIENCE CENTER AT HOUSTON SCHOOL OF PUBLIC HEALTH Houston, Texas May 1994

ACKNOWLEDGMENTS

1

-

In a work like this there are always many participants. The brevity of the document is not indicative of the enormous input from so many different individuals. I was fortunate to be surrounded by such exemplary individuals to encourage, brainstorm, and solve seemingly insurmountable problems.

Special thanks to the staff at the School of Public Health. In particular Dr. Sharon Cooper and Dr. Marcus Key for providing guidance and acting as my mentors so unselfishly. Thanks also to Dr. Michael Decker for his thorough review of the project. Other professors that I had in occupational, environmental, and epidemiology courses greatly expanded my knowledge base which allowed completion of study. Those special individuals are: Dr. George Delclos, Dr. Larry Whitehead, Dr. Kim Waller, Dr. Fred Annegers, Dr. Mary Smith, and Dr. Keith Burau.

To Ms. Julie Shadoan of Fort Detrick, MD for her many hours of work, I express a special thanks. For her tireless efforts to overcome program errors we seemed to encounter at every turn, I am extremely grateful. This project could not have been completed without her insight, skills, persistence, optimism.

The Air Force has been most gracious in their support. Thank so much for each one that contributed so unselfishly. Lt.Col Ben Sierra-Irizarry and SSgt. Todd Goins have provided data input and ideas on several occasions. Maj. Kevin Grayson and Maj. John Allen were extremely helpful in the initial generation of this study. Other individuals that were also extremely helpful were: Mr. Phil Polish, Maj. Cari Sherris, Mr. Joe McInturf, and Mr. Tom Moore.

A RETROSPECTIVE STUDY OF A 1992 SUBCOHORT OF AIR FORCE PERSONNEL EXPOSED TO HAZARDOUS NOISE BY AIR FORCE SKILL CODE

Jim Alan Davis, D.V.M. The University of Texas Health Science Center at Houston School of Public Health, 1994

Supervising Professor: Sharon Cooper, Ph.D.

This study measured the magnitude of risk of the 1992 cohort of employees (enlisted, officer, and civilian) on the United States Air Force Hearing Conservation Program by occupational codes (Air Force Skill Codes, AFSC). Some 160,062 individuals that received annual audiograms during this period were eligible for analysis. To enhance precision, females were excluded from the study since they were less than 6% of the subcohort resulting in 151,512 potential male employees to analyze. The study further restricted this group to view only the most recent audiogram for each individual that had a valid service and rank code. The final number of this subcohort was 107,421. Only AFSCs with 100 or more employees were analyzed. This represented 117 AFSCs with 93,854 audiograms (87.4% of 107,421). The age and race distribution of the 107,421 audiograms were used as the standard for direct adjustment of each AFSC. Risk of Permanent Threshold Shift (PTS) ranged from several AFSCs with 0% to civilian aircraft mechanics with 11.4%. The overall crude risk for the standard was 2.3%. Six of the top ten risk categories were civilian even though only 28% of the AFSCs analyzed were civilian. Due to wide confidence intervals, the Air Force should use caution in implementing new procedures from these findings until they are replicated in a subsequent year group.

TABLE OF CONTENTS

Page

DEDICATIONii
ACKNOWLEDGMENTSiv
ABSTRACTv
LIST OF TABLES
LIST OF FIGURES
SECTION I INTRODUCTION 1
A. Specific Aims1
B. Significance
C. Hearing Loss History and Susceptibility
D. Risk Factors as Confounders4
SECTION II METHODS AND PROCEDURES
A. Selection and Recruitment of Study Population8
B. Selection of Comparison or Standard Group13
C. Data Collection and Data Management Strategies
D. Protocols or Instruments Used
E. Independent, Dependent, and Confounding Variables
F. Method of Data Analysis15
SECTION III RESULTS
SECTION IV DISCUSSION
SECTION V RECOMMENDATIONS
APPENDICES
A. Appendix A25
B. Appendix B

の一年二月、日本の日本の一日の一日

TABLE OF CONTENTS

Page

	C. Appendix C	
	D. Appendix D	30
	E. Appendix E	
	F. Appendix F	
	G. Appendix G	43
BIBLIO	OGRAPHY	
VITA		

LIST OF TABLES

• .

..

1 af - 1

5

-

1.	Distribution of initial 1992 cohort for all races and AFSCs by sex and military status	10
2.	Frequency distribution of male cohort by age and racel (Derivation of standard weight)	6
3.	Comparison of crude rates of the 117 AFSCs(N=93,854) with the crude rate of the 1992 Air Force Annual Hearing Conservation Report (AFHCR) and adjusted rates of the 117 AFSCs	7
4.	Summary of ten AFSCs with highest and lowest risk of developing PTS 1	9

Page

LIST OF FIGURES

1. DD Forms 2216(Hearing Conservation Data form) for 1991 and 1992 in the HCDR by total submitted for data entry and number of those being annual audiograms	9
2. Number of employees in each AFSC by Enlisted, Officer, and Civilian	
3. Number of AFSCs in different adjusted rate categories	18
4. Distributition of risk of developing PTS among AFSCs with over 100 employees in each AFSC	22

7

114

Page

SECTION I INTRODUCTION

A RETROSPECTIVE STUDY OF A 1992 SUBCOHORT OF AIR FORCE PERSONNEL EXPOSED TO HAZARDOUS NOISE BY AIR FORCE SKILL CODE

A. SPECIFIC AIMS

This research is designed to provide information on the risk of developing permanent hearing loss across occupations in the United States Air Force. All Air Force individuals exposed to hazardous noise (over 85 dB-8 hour weighted, over 115 dB intermittent noise, or over 140 dB impact or impulse noise) and free of significant hearing loss are entered into the Air Force Hearing Conservation Program (AFHCP).¹

The AFHCP has been conducted by hundreds of highly trained technicians, physicians, and allied support personnel for many years. These efforts have resulted in a database (Air Force Hearing Conservation Data Registry - HCDR) with over three million audiometric records. ² Each year the Air Force Annual Hearing Conservation Report is compiled to summarize data for the previous year. ³ The report contains a myriad of useful analytical and epidemiologic information. Incidence of Permanent Threshold Shifts (PTS) is reported annually in this report as required by Air Force regulations. ⁴ It also contains a frequency analysis of PTS for various career fields for officers, enlisted, and civilian employees, but the career fields cover broad areas. The career fields are not specific for occupations identified by Air Force Skill Codes (AFSC).

With scores of different Air Force Skill Codes on the AFHCP, it is extremely

beneficial, in terms of targeting appropriate interventions, to identify those AFSCs showing the highest risk of hearing loss compared to other AFSCs.

Therefore, the specific aims of this study are :

(1) to identify the ten AFSCs showing the highest risk for hearing loss, and

(2) to identify the ten AFSCs on the AFHCP that have the lowest risk of developing permanent hearing loss.

B. SIGNIFICANCE

The Department of Labor (DOL) and Veterans Administration (VA) pay millions of dollars annually as a result of claims of hearing loss by Department of Defense (DOD) eivilian employees and former military personnel. The DOL reported the average cost in civilian Air Force employee claims was \$4,969,417 annually from 1989 to 1992. ⁴ On average, 993 civilian employees made claims annually to the DOL during this period. The VA only pays claims to prior military personnel with service connected hearing loss. In 1992 the VA had 7,531 Air Force cases on the active roll where hearing loss was considered the primary disability. There were 47,699 Air Force cases on the active roll during the same period where hearing loss was considered a secondary disability. Primary disability recipients received a total of \$25,431,576 in 1992 alone. Financial payments received by individuals in the secondary disability category are not kept by the VA. ³ When Air Force costs are combined with other DOD agencies, hearing loss claims paid by the United States government are substantial.

Any effort to reduce hearing loss will ultimately have an impact by reducing claims paid and alleviating the long term physical disabilities so many service connected personnel experience. By identifying the ten AFSCs with the greatest risk for hearing loss,

Air Force management and Military Public Health officers can search for factors that may put these individuals at increased risk and focus preventive medicine efforts on these high risk occupations. Remediation measures can then be designed to circumvent further preventable hearing losses. Amelioration measures may appear to be beneficial enough that they may be instituted Air Force wide.

By identifying the ten AFSCs on the AFHCP with the lowest risk of hearing loss, AF management will be able to further analyze the AFHCP. Factors such as lower levels of noise exposure, better education programs and different types of hearing protection could be identified as protective for these occupations. The discovery of these protective factors could of value to the Air Force and other DOD agencies in further curtailing occupational hearing loss.

C. HEARING LOSS HISTORY and SUSCEPTIBILITY

1. S. S.

Loss of hearing from occupational noise has been known for over 300 years. ⁶ Development of technology during this century has brought louder and louder machines and industrial processes into the workplace. ¹⁴ In 1830 "osbroke" spoke of certain occupations predisposed to hearing loss.

> The blacksmiths' deafness is a consequence of their employment; it creeps on them gradually, 'n general at about forty or fifty years of age. At first the patient is insensible of weak impressions of sound; the deafness increases with a ringing and noise in the ears, slight vertigo, and pain in the cranial bones, periodical or otherwise, and often violent.

As the Industrial Revolution advanced, it also brought hearing loss to millions of

Americans in industry ⁷. Noise is now the most common hazard in the workplace. ¹⁰ Over 40 million Americans suffer hearing loss from various sources. ⁷ Over 7,900,000 U.S. workers are exposed to noise levels at or above 80 dB. ¹⁰ It is estimated that more than 20 billion dollars have been paid out in compensation for hearing loss to workers. Noiseinduced hearing loss is listed by the Center for Disease Control as one of the ten leading occupational diseases in the U.S. ¹¹

It has long been recognized that some individuals are more susceptible to hearing loss than others. ^{2,12,14} There has been much speculation about potential factors causing this inter-individual variation including: heredity, diet, blood and nutrition supply, stress, drugs, metabolism, race, age, sex, eye color, non-occupational exposure to noise, smoking, previous hearing loss, systemic diseases (cardiovascular, hypothyroidism, diabetes, hyperlipoproteinemia), use of hearing protection, consumption of salicylates, plasma LDLcholesterol concentration, gunfire, and tinnitus. ^{27,12,13,15-19} If we can learn more about factors that predispose people to hearing loss, we may be able to reduce hearing loss.

D. RISK FACTORS as CONFOUNDERS

Of all the potential confounders listed above as potentially affecting susceptibility to hearing loss, one is likely to affect the study. Presbycusis, a slow progressive deterioration of hearing not attributable to any cause except aging, has been found in many occupational hearing loss studies as a major factor in hearing loss. ^{10,19} Lutman and Spencer²⁰ studied 2,162 subjects free of material conductive impairment. Study participants ranged from 17 to 80 years. Their findings showed increased hearing threshold levels with age. Bauer *et al.* " studied 47,388 noise exposed workers in Austria and found that age was a dominating factor in determining hearing threshold loss. Ribak *et al.* ²¹ studied 777

aircrew members of the Israeli Air Force for permanent hearing threshold shifts. It was discovered that age was highly correlated with hearing loss (p<0.001) while aircraft type and accumulated flying time played a minor role. Fitzpatrick, in a subsequent study of 211 U.S. Army aviators, found noise exposure as measured by total flight hours (1,000 to 4,000 hours) increased threshold shifts four-fold (p<0.001), but he also found hearing loss with clear progression in each age group (p<0.001).²

Gender also seems to play an important role in susceptibility to hearing loss. One auditory brainstem response study has shown women have different lengths of time for brainstem response than men. Explanation for this was hypothesized: lighter ossieles in women, shorter cochleas, and variations in the skuil (men have thicker skull bones than women and different shapes of the skull bones). " Royster et al. 23 studied audiometric test data on 14,000 industrial employees from several different types of industries in North Carolina. The study found that black females exhibited the lowest hearing threshold levels. White males had the highest hearing threshold levels. Black males and white females had similar hearing threshold levels. Royster et al.²³ stated "that meaningful evaluations of industrial audiometric data bases are not possible unless race and sex compositions of the population are considered." Johnson's 24 review of field studies on industrial exposure also showed an increase in noise induced permanent threshold shift (NIPTS) in men compared with women. Johnson postulates that teenage boys are involved in far more noise activities than teenage girls. He states this trend is expected to continue through adulthood. Male non-occupational exposures (chain saws, guns, etc.) predispose them to NIPTS. In the study by Bauer "previously mentioned, of the 47,388 noise exposed workers, sex and age were found to be the dominating factors in determining hearing threshold.

The effect of race on hearing sensitivity has been studied and may affect data interpretation. As mentioned earlier, Royster *et al.* 2 noted a difference in hearing threshold

level 3 by race. This study was performed in 1979. An earlier study of patients at Johns Hopkins Hospital in 1930 by Bunch and Raiford ²³ indicated better hearing in blacks than whites. Karsai *et al.* ²⁶ studied a population of 836 longshoreman to analyze hearing differences in men from several different subcultures. Black Americans had superior hearing compared to four other groups, all of which were white. The four other groups (Italian, Irish, Yugoslav, and White American) were not significantly different.

The amount of melanin in the skin and eye appear to be correlated with susceptibility to hearing loss. This hypothesis is supported by studies like those in the previous paragraph discussing differences in darker and lighter races. LaFerriere et al. 27 discovered the density of melanocytes throughout the labyrinth varies according to skin pigmentation. The pale person will have hardly any melanin in the stria vascularis. Barrenas and Lindgren¹² cited the important discovery of the correlation of eye color and inner ear melanin by Bonaccorsi in 1965. The results of studies correlating eye color with hearing loss are mixed. Hood et al. 2 tested 38 subjects for temporary threshold shifts (TTS). They were divided into four groups by eye-color: dark brown, light brown, greengrey, and blue. The results confirmed earlier studies that TTS is related to melanin content of the iris. Carter et al. "studied 118 otologically normal Australian soldiers. In this study left ears from soldiers with eye color indicating no melanin pigmentation of the iris had poorer hearing at 3 kHz (p<0.05) than soldiers with melanin pigment. Although the majority of studies do indicate eye color is correlated to hearing loss, a few do not. For instance, Karlovich (1975) tested 45 young adults for auditory fatigue and found no significant differences in brown, green-gray, and blue iris categories. ³⁰ Hood et al.²⁰ attempted to explain why Karlovich produced results different from others. He stated the methodology Karlovich used of pulsed tones rather than continuous tones may have produced these misleading results. The protective effects of melanin in the inner ear may be

due to binding of noise induced free radicals. This may protect the hair cells against a hazardous overload produced by excessive noise. ¹²³¹ Carlin and MCroskey 22 (1980) studied 100 industrial employees and concluded that eye color may be related to the effects of noise on hearing. They suggested any future study of PTS or TTS should consider eye color.

Nonoccupational noise may be a confounder. Occasional exposure to loud rockand-roll and music will not produce significant hearing damage. Habitual exposure though may cause damage between 2,000 and 8,000 kHz. ⁷ Also gunfire can show a detectable effect on hearing threshold, in particular on the left car. ^{7,39} Very little can be done to control for this variable since its impact cannot be clearly defined.

Smoking has also been implicated as affecting hearing susceptibility. The results of studies have varied. Barone *et al.* ¹⁰ studied 2,348 noise-exposed white males and found smokers had a statistically significant increased risk of noise-induced hearing loss. Pyykko *et al.* ¹⁰ carried out a detailed analysis of risk factors for hearing loss on 199 Finnish forestry workers. They concluded the effect smoking has on hearing loss is still uncertain. Their results showed no significant difference in smokers versus nonsmokers.

SECTION II METHODS AND PROCEDURES

A. SELECTION and RECRUITMENT of STUDY POPULATION

Since all individuals in the HCDR are exposed to hazardous noise (as defined by the Air Force) and are otologically normal, all individuals in the registry could be potential candidates for this study. The study population was selected from the HCDR with three primary criteria in mind.

Accuracy of data is of paramount importance so the results would be valid and information bias would be minimized. Though the HCDR has been active for several years, confidence in the quality of data has improved significantly in recent years. During the last few years a thorough quality control has been implemented to insure data collected at base level is entered properly and completely into the HCDR. For this reason, selection of a more recent subcohort would provide the highest quality data.

A large enough sample size is also important so that adjusted risks would be reliable. As with any study, sample size may mean the difference between statistically significant results and just interesting information. The larger the sample size the better, but it still must be kept small enough to be manageable for analysis. In 1991 alone the HCDR had 198,890 DD Forms 2216 entered into the database. The DD Form 2216 (Hearing Conservation Data form) is a multipurpose form used by the Air Force to record audiograms for several different purposes such as 90-day follow-ups, termination, annual, etc. After identifying those audiograms marked with the code for annual audiograms, 167,144 were analyzed for hearing loss. ³ The 1992 number of DD Forms 2216 entered into the HCDR was 198,484. Once again when only those coded for annual audiograms were

8

isolated 160,062 were analyzed for hearing loss (Figure 1).³⁴



One last concern is that the data being analyzed should be current enough to reflect current working conditions in the United States Air Force. Currency of data will help Air Force management use the results of this study immediately. The more current the data, the more confidence can be placed that inferences from the results in the study are applicable to the present work force.

To support the three objectives stated above, part of the cohort of 1992 individuals entered on the HCDR was selected for the study population. New accessions are not allowed to start working in an AFSC unless they are cleared for medically normal hearing. New accessions into the military workforce for 1992 were not included since it could not be documented they were disease free the 12 months prior to their audiogram. The 1992 cohort that were on the AFHCP in 1991 received their annual audiogram sometime in 1992. This group should adequately meet all three requirements making the study data accurate, reliable, and immediately usable by the Air Force. Since sex was implicated in previous literature as being a possible risk factor for disease, it was considered in this study. Incorporation of sex (a dichotomous variable) would double the number of strata in the final analysis. This effect of doubling cells to be analyzed could cause many unstable cells (cells < 5) if not enough women were in this subcohort. Of the 160,062 only 8,550 were female (Table 1). Since this group was less than 6% of the subcohort and would probably decrease the precision of the study, it was decided to further restrict the study to just males. Since the goal of the study is to identify high risk AFSCs, removing sex as a confounding variable in the design phase was parsimonious.

Mi	litary	Civil	ian	A	All	
N	%	N	%	N	%	
6,873	5.18	1,677	6.13	8,550	5,34	
125,822	94.82	25,690	93.87	151,512	94.66	
132,695	100.00	27,367	100.00	160,062	100.00	
	Mi N 6,873 125,822 132,695	Military N % 6,873 5.18 125,822 94.82 132,695 100.00	Military Civil N % N 6,873 5.18 1,677 125,822 94.82 25,690 132,695 100.00 27,367	Military Civilian N % N % 6,873 5.18 1,677 6.13 125,822 94.82 25,690 93.87 132,695 100.00 27,367 100.00	Military Civilian Al N % N % N 6,873 5.18 1,677 6.13 8,550 125,822 94.82 25,690 93.87 151,512 132,695 100.00 27,367 100.00 160,062	

() 1

Franking Street

1 - 1 E

This study further restricted the 151,512 annual audiograms from males to view only the most recent audiogram for each individual that had a valid service (Regular Air Force, Reserve, National Guard, or other) and rank (officer, enlisted, civilian grade) code. The final number of individuals/audiograms in this subcohort was 107,421. The decrease in the 1992 male subcohort from 151,512 to 107,421 was probably related to the individual receiving more than one annual audiogram, double testing (i.e. Jan 92 then again Dec 92) where windows of annual audiogram might overlap, improper coding on the form, or improper data entry.

Further restrictions were necessary. The individual unit of analysis in this study is the AFSC. If an AFSC had too few individuals in it, its adjusted risk would have such wide confidence intervals the point estimate would lose meaningful precision. Also the study is geared to have Air Force-wide application, therefore it was also important to select AFSCs with large enough numbers of individuals to significantly impact the Air Force. Therefore, some cut-off point would need to be determined for the number of employees per AFSC to be analyzed.

The subcohort of 107,421 was analyzed to see how many AFSCs it represented and the number of employees in each. When AFSCs where broken out initially it resulted in 4,648 different codes (Appendix A). There were several possible reasons for such a large number. The main reason is many AFSCs have suffix identifiers that the database treats separate AFSCs. For instance, a 908XX, 90830, 90850, and 90870 all represent the 908 series for an enlisted public health technician. Military Personal Center and the Civilian Personnel Center at Randolph AFB were consulted on all the different coding variations of AFSCs for civilian, officer, and enlisted. To combine individual codes to make meaningful groupings, a strategy was developed. Any AFSC code with over seven employees was combined with its parent code. The number seven was chosen because even if three codes with seven each where combined, the parent code would still only be 21; a number too small to analyze. Rarely did any AFSC code with seven individuals have another code with more than two individuals that would have contributed to their parent code therefore the rationale for the method seemed reasonable, although arbitrary. AFSCs were combined in the manner described above resulting in 2,718 total AFSC codes still

listed.

The distribution of AFSCs after this combining made the decision for cutoffs simpler (Figure 2). The preponderance of employees were in AFSCs with more than 100 persons. One hundred seventeen AFSCs represented 93,854 individuals with a cut-off of 100 persons per AFSC (Appendix B). Thus 87. 4% of the subcohort of 107,421 could now be analyzed with a manageable number of AFSCs.



B. SELECTION of COMPARISON or STANDARD GROUP

1. 1. N

One of the formidable hurdles of any retrospective cohort study is to identify an adequate comparison group. The comparison group in a cohort study is utilized to compare the disease experience in an unexposed group to that of the exposed study group. Usually a comparison is chosen in one of these 2 ...: internal comparison group, external comparison group, or comparisons with the general population. ³⁵ Since the goal of this study is to rank the magnitude of effect in each AFSC, a comparison group was not necessary.

A standard group though was needed for adjustment in the analysis. Since different AFSCs would have different proportions of race and age groups, adjustment using a standard distribution of these potential confounders would remove any bias they might produce in the crude point estimate for that AFSC. For direct standardization the standard group can be one of the groups under study, a combination of groups under study, or some external group. The actual values of the adjusted rates would depend on which standard is chosen. Therefore the choice of the standard would alter the magnitude of effect in the adjusted rate, however, the magnitude in differences of adjusted rates would not vary. ³⁴³⁷ This means the irregardless of the standard chosen the ranking of rates would not be altered. For convenience the subcohort of 107,421 males were selected to be the standard.

C. DATA COLLECTION and DATA MANAGEMENT STRATEGIES

Data and preliminary analysis were obtained from the HCDR at Ft. Detrick, MD. The researcher had extensive communications with Armstrong Lab (Brooks AFB, Texas) Air Force Military Personnel Center (Randolph AFB, Texas) and Civilian Personnel Center (Randolph AFB, Texas) to resolve apparent inconsistencies in occupational coding. Collaboration between the researcher and HCDR was necessary to adjust occupational codes and perform the raw data analysis. Further analysis were accomplished by the researcher on a 386-IBM compatible personal computer using a software spreadsheet.

D. PROTOCOLS or INSTRUMENTS USED

All data was furnished courtesy of the U.S. Air Force. No data contained any personal information such as social security numbers.

No instruments were used in this study.

E. INDEPENDENT, DEPENDENT & CONFOUNDING VARIABLES

INDEPENDENT VARIABLE:

The independent variable for this study was each AFSC under study.

DEPENDENT VARIABLE:

The dependent variable for this study was the presence or absence of permanent hearing threshold shift (PTS). PTS was defined as an increase (an average of 10 dB at 2,000, 3,000, and 4,000 Hz, either ear) in the hearing threshold relative to the baseline audiogram after a 40 hour noise free period. ¹

CONFOUNDING VARIABLES:

Sex is regarded in the literature as a possible confounding variable. By restriction it was removed from the study during the design phase.

Adjustment was made for age and race during the analysis of this study. Age was readily available on the current HCDR. Race was used as a surrogate for melanin in the inner ear. The data on race was obtained from another DOD database and was merged with the HCDR database.

F. METHOD of DATA ANALYSIS

As with any retrospective cohort study, all participants should be disease free at the beginning of the review period. At the beginning of 1992 all individuals that had been on the AFHCP in 1991 should have had normal hearing. If one of these individuals had a PTS in 1991 they would normally be removed from a hazardous noise area thereby removing them from hazardous noise. Occasionally, by direction of a physician or audiologist, an individual with a PTS in 1991 may have been reassigned a new baseline for his hearing level and considered to have normal hearing. These individuals would stay on the AFHCP and continue in their work centers. Each one of the individuals on the AFHCP in 1991 should have received their annual audiogram sometime during 1992 unless they changed jobs, left the military, or were removed from a hazardous noise area for medical reasons. During the twelve month period of 1992, annual audiograms entered into the HCDR were categorized as either normal or a permanent threshold shift (PTS).

To develop the standard, the subcohort of 107,421 was stratified by age and race. The percentages generated in this analysis would be used later as the multiplicand with the stratum rates of each AFSC (Table 2).

	Wh	White		k	Other		
AGE	N	%	N	%	N	%	
18-27	20,865	19.42	2,225	2.07	6,402	5.96	
28-37	32,159	29.94	3,707	3.45	11,378	10.59	
38-47	13,464	12.53	1,536	1.43	7,425	6.91	
> 47	4,563	4.25	5 39	0.50	3,158	2.94	

. . .

۲,

The measure of disease frequency studied was cumulative incidence (CI). This measure is also often referred to simply as risk. Since CI is the proportion of people that have become diseased in a given period, it fits this study well. \leq

CI = <u>mumber of new cases of a disease (PTS) during a siven period of time</u> total population at risk (all AFSCs)

The crude cumulative incidence for each AFSC was adjusted for age and race by direct adjustment using 12 age/race strata. The adjusted cumulative incidence yielded a more accurate point estimate. A 95% confidence interval around the adjusted point estimate was also calculated (Appendix C & D).

SECTION III RESULTS

The overall crude rate of PTS for the study group of 107,421 was 2.52%. This figure matches the results of the 1992 Air Force Annual Hearing Conservation Report which calculated 2.5% for a crude rate. 3

Analyzing the group of 117 AFSCs under study and comparing them to the 1992 Air Force Annual Hearing Conservation Report revealed similar crude rates in officers, enlisted, and civilians (Table 3). If active sotable in the adjusted rates in these categories is the increase in the enlisted rates and the figure as n the civilians. By removing the effect of age and race the enlisted and civilian overall adjusted rates are almost identical yet both are distinctly higher than the officer category.

Table 3. Comparison of crude rates of the 117 AFSCs(N=93,854) with the crude rate of the 1992 Air Force Annual Hearing Conservation Report (AFHCR) and adjusted rates of the 117 AFSCs.							
	Officer	Enlisted	Civilian	All			
AFHCR Crude	1.7	2.0	4.8	2.5			
117 AFSCs Crude	1.41	1.98	5.10	2.28			
117 AFSCs Adjusted	1.45	2.72	2.83	2.49			

The range of adjusted rates (risks) for all 117 AFSCs ranged from 0% to 11.38% (Figure 3.). The majority of the AFSCs feil between adjusted rates of 0.5% and 4.5%. The

distribution of the risk is skewed to the right with several AFSCs as outlyers.



The adjusted risk for each one of the 117 AFSCs were summarized in a single table. The table was then sorted from highest risk to lowest risk according to enlisted, officer, and civilian (Appendix E). Lastly, the ten AFSCs with the highest and lowest risks of hearing losses were summarized (Table 4). The actual calculations of the highest (Appendix F) and lowest (Appendix G) were reported also.

Only four AFSCs were above an adjusted rate of 7.0%. Of those, three were civilian: Aircraft mechanic, Structural civil engineer, and vehicle maintenance. The one enlisted category above an adjusted rate of seven was radar technician. The highest risk calculated was on the civilian Aircraft mechanic at 11.38%.

It is interesting to note that six of the AFSCs with the greatest hearing loss were

Table 4. Summary of Ten AFSCs with the highest and lowest risk of developing PTS								
Ten AFSC (Ordered	at highes highest to	t risk of l b least)	PTS	Ten AFSC at lowest risk of F (Ordered lowest to highest)				
Title	AFSC	Risk	Confid. Interval	Title	AFSC	Risk	Confid. Intervai	
Aircraft Mechanic	C-8840	11.38%	17.92% 4.85%	A.L. Missile Systems	E-466XX	0.00%	0.00% 0.00%	
Structural civ.eng.	C-552XX	8.61%	15.86% 1.37%	Nuclear Weapons	E-463XX	0 .00%	0.00% 0.00%	
Vehicle Mainten.	C-472XX	8.37%	15.34% 1.39%	Instru- mentalist	E-872XX	0.00%	0.00% 0.00%	
Radar Technician	E-303XX	8.30%	10.87% 5.73%	SpecialOps Pilot	0-1315	0.00%	0.00% 0.00%	
Aeromedic Specialist	E-901XX	6.72%	12.00% 1.43%	PilotTrainee	0-0006	0.00%	0.00% 0.00%	
Avionics Technician	E-451XX	6.26%	10.08% 2.43%	Aircraft Pneudraui	C-8268	0.26%	0.75% -0.24%	
Jet Eng. Mechanic	E-426XX	6.00%	9.29% 2.72%	Medical Service	E-902XX	0.27%	0.47% 0.06%	
Machine Tool Oper	C-3431	5.75%	8.67% 2.83%	Combat Control	E-273XX	0.30%	0.88% -0.28%	
Aircraft Ordinance	C-6652	5.64%	10.23% 1.04%	SpecialOps Nav	O-1585	0.42%	1.23% -0.39%	
Aircraft Mainten.	C-452XX	5.44%	9.10% 1.79%	Electronic Com.	E-304XX	0.51%	1.22% -0.19%	

12. 61

1

÷17

14 1 11

civilian categories. This is particularly important in light of the fact that out of the 117 AFSCs analyzed only 33 (28%) were civilian. It appears that several civilian categories have higher risk of PTS than other categories even after adjusting for age, race, and sex.

The spread of the confidence intervals was greatly affected by the number within an

AFSC. For instance, the civilian AFSC 3431 (Machine tool operator) had a much narrower range with 289 employees versus the civilian AFSC 8840 (Aircraft mechanic) with 108 employees. There is significant overlap of many confidence intervals. Certainly the lower limits of the civilian Aircraft mechanic (4.85%) and the enlisted Radar technician (5.73%) are well above the group of AFSCs noted in Figure 3.

艰险

SECTION IV DISCUSSION

Millions of dollars are paid annually to claimants of civilian DOD employees and past military personnel. All three services contribute substantially to this cost. In addition to cost, these injuries inflict undue disabilities on many of these individuals.

The data in this study are extremely valuable and should be replicated in the future with each subsequent year group. By analyzing data directly by the AFSC, managers at base level will be able to review programs for specific high and low risk groups. High risk groups can be monitored more closely, trained better, or other preventive measures can be implemented to take them off the high risk group. Conversely, low risk groups can be analyzed to see what is being done in that occupation to keep them in such a low risk. Perhaps some of their mechanisms can be duplicated in other areas to decrease the risk of hearing loss. This investigative work will surely generate paradigm shifts in the way the Air Force manages its AFHCP.

In particular the Air Force should pay particular attention to certain high risk civilian AFSCs. Since the civilians overall adjusted rate is very similar to the enlisted overall adjusted rate, it is particularly alarming to find a disproportionate number of civilian AFSCs in the top ten risk rankings.

Finding the enlisted AFSC 901XX (Aeromedical Specialist) seems out of place as one of the top ten risks for hearing loss. Many individuals in this occupation are on the hearing conservation program since they are required to fly periodically. Yet to the researcher their exposure seems minimal when compared to most other AFSCs on the AFHCP. The question of why this group appears in the top ten list may have two plausible answers. With wide 95% confidence intervals, it may be in its ranking by chance from the multiple comparisons issue. Another explanation may be that since this is the AFSC that administers the audiograms in the Air Force, their ability to take audiograms may be different from that of the rest of the subcohort.

The Air Force should not take any major action on this study alone. Until a AFSC has reoccurred in the highest risk or lowest risk category 2 years in a row it may not be valid. Validity of this study may be affected by two factors.

Multiple comparison issues has been cited by authors a potential problem when analyzing large amounts of data. In essence, it states that because of the laws of probability, given a large enough sample, some variables will be statistically significant when they really are due to chance of occurrence. ³⁷ The application here is that an AFSC that was in the left end of the curve (high risk) in 1992 may end up in the center next year. It may have just randomly appeared as a high risk this year when in fact it is not a historically high risk group (Figure 4).



As second area validity may have been sacrificed was when the 4,648 AFSC codes were combined to the 2,718 AFSC codes. Although experts at Randolph AFB were consulted, misclassification of exposure may have occurred. Some decisions whether to combine codes into a parent code where complicated and time consuming. The researcher did his best to double check all decisions and entries. Even though these efforts were made validity may have been altered.

Precision of the results are also a concern. Since the goal was to analyze AFSCs, removing females early in the study was probably wise. In the future though, as women increase in percentages of the workforce, they should also be examined. Stratifying by age was prudent and would be a well recognized method to control for such a strong risk factor.

Race and/or melanin content of the inner ear has been implicated as a risk factor and was therefore adjusted for in the analysis. This adjustment may have decreased precision for no reason at all if race is not a strong risk factor in the military setting. Future studies should be performed to evaluate race as a risk factor for occupationally related hearing loss in the military.

Certain potential confounders were not controlled for in the design or analysis. Smoking, eye color, nonoccupational noise exposures have not been recorded on any easily accessible database in the Air Force; therefore, they were not adjusted for. It is hoped that smoking and nonoccupational noise exposures are evenly distributed through the AFSCs so they will not skew the results.

The greatest concern for precision surrounds the issue of what number of employees is a reasonable number to use as a cut-off for analysis. One hundred was used in this study with very rational reasoning. Unfortunately, from looking at the results closely one can see that even with this number of employees the adjusted risk can be easily altered by one or two PTSs in the AFSC. This factor is reflected in the wide confidence intervals. In future studies this cut-off may need to be altered.

SECTION V RECOMMENDATIONS

The validity and precision issues that have been raised should not minimize the value of this study. If the study is replicated with 1993 data and similar results are found, Air Force management can confidently approach managers and Military Public Health Officers at base level to follow through with their investigative work to improve the AFHCP.

The Air Force is currently combining several databases at a central location. Within one to two years the access and quality of the data should be better. Race information will be easier to obtain. Generation of AFSCs that will not require manual and computer combining to parent codes will be likely. An Air Force computer programmer should build a program that will automatically generate this study. Even though this study was labor intensive, by using the avenues soon available much of the work intensity should be alleviated. Appendix A Example of printout with AFSCs broken out into 4648 different AFSC codes ANALYSIS FOR EACH OCCUPATIONAL GROUP

	MILITA	RY	CIVILI	CIVILIAN		ALL	
	N	*	N	x	N	*	
OCCUPATIONAL CODE		÷					
452XB	1	0.00	0	٥	1	0.00	
452XQ	167	0.15	9	0.01	176	0.16	
452X1	147	0.13	4	0.00	151	0.13	
452X2	233	0.21	2	0.00	235	0.21	
452X3	77	0.07	٥	٥	77	0.07	
452X4	1,569	1.38	18	0.02	1,587	1,40	
452X5	237	0.21	1	0.00	238	0.21	
4 52X7	19	0.02	٥	٥	19	0.02	
452X9	5	0.00	2	0.00	7	0.01	
45272	1	0.00	0	0	1	0.00	
452Y4	ł	0.00	0	e	:	0.00	
4520	1	0.00	٥	٥	1	0.00	
45200	26	0.02	2	0.00	28	0.02	
4521	4	0.00	٥	0	4	0.00	
45210	1	0.00	٥	o	1	0.00	
45211	а	0.01	û	C	8	0.01	
45212	1	0.00	٥	٥	1	0.00	
45214	6	0.01	٥	0	6	0.01	
45215	1	0.00	٥	G	1	0.00	
45216	1	0.00	٥	0	ì	0.00	
45222	۱	0.00	٥	٥	1	0.00	
45223	1	0.00	Û	0	ĩ	0.00	
45224	\$	0.00	0	٥	1	0.00	
4523	ı	0.00	٥	٥	1	0.00	
45230	2	0.00	1	0.00	3	0.00	
45231	37	0.03	٥	٥	37	0.03	
45232	56	0.05	1	0.00	57	0.05	

(CONTINUED)

1 4 5 4 E

to the second second

.

the set of the set of the set

Appendix B

	1.05 4			Officer*	•	1	Civilia	n* **	
		Arex uper	totals	AFSC	Number	totals	AFSC	Number	totais
	Codes	AFSC		Codes	in AFSC		Codes	In AFSC	
1	454	8682	8682	1115	3756	3756	3806	2284	2284
2	452	7886	16568	1045	2577	6333	8852	2225	4509
3	457	5993	22561	1065	1803	8136	571	568	5077
4	462	3507	26068	1355	1639	9775	8602	566	5643
5	113	2951	29019	1055	1559	11334	3414	565	6208
6	114	2645	31664	1545	858	12192	81	525	6733
7	455	2205	33869	1535	846	13038	2892	482	7215
8	458	2083	35952	1575	658	13696	4102	445	7660
9	571	1960	37912	1745	618	14314	454	410	8070
10	631	1840	39752	1235	609	14923	5378	342	8412
11	811	1778	41530	1525	596	15519	8255	309	8721
12	461	1322	42852	1555	563	16082	3431	289	9010
13	272	1319	44171	9356	480	16562	551	277	9287
i4	551	1265	45436	2225	383	16945	5402	250	9537
15	552	1042	46478	1825	341	17286	3703	245	9782
16	902	1007	47485	1325	335	17621	8268	223	10005
17	472	1004	48489	9016	277	17898	6652	216	10221
18	112	960	49449	1455	267	18165	2604	215	10436
19	605	935	50384	1025	240	18405	6968	205	10641
20	542	907	51291	1565	239	18644	3500	200	10841
21	545	886	52177	1145	219	18863	552	196	11037
22	431	822	52999	1435	204	19067	457	191	11228
23	456	645	53644	1425	201	19268	545	187	11415
24	411	600	54244	6	193	19461	458	164	11579
25	208.	513	54757	1495	192	19653	472	157	11736
26	116.	390	55147	9766	161	19914	431	138	11874
27	423	382	55529	2255	157	19971	452	121	11995
28	753	321	55850	1716	152	20123	4604	118	12113
29	901	315	56165	2245	149	20272	2805	114	12227

The state of the second second

* These codes sometimes end in a 2 digit suffix. Two periods stand for the suffix which could be any 2 numerals. In other references this designation is often XX rather than two periods.

** If these codes have less than four digits, it can be assumed it is preceded by enough zeros to make it a four digit code.

Appendix B

	Enliste	d*		Officer**			Civilia	n* **	
; 1	AFSC	Number	totals	AFSC	Number	totais	AFSC	Number	totals
<u> </u>	Codes	in AFSC		Codes	In AFSC		Codes	in AFSC	
30	427	281	56446	1585	148	20420	542	109	12336
31	118	280	56726	1315	146	20566	8840	108	12444
32	117	271	56997	1595	143	20709	5413	104	12548
33	361	271	57268	2865	120	20829	8800	101	12649
34	207	264	57532	4024	119	20948	3705	99	12748
35	304	264	57796	2295	115	21063	8862	98	12846
36	426	230	58026	36	107	21170	5705	92	12938
37	871	229	58255	66	88	21258	6907	88	13026
38	273	218	58473	51115	86	21344	5309	87	13113
39	566	210	58683	1445	81	21425	2854	87	13200
40	115	208	58891	1365	80	21505	5823	86	13286
41	122	169	59060	2265	73	21578	6912	84	13370
42	602	163	59223	1035	73	21651	5350	84	13454
43	303	156	59379	1406	68	21719	3416	83	13537
44	872	147	59526	9025	64	21783	90850	81	13618
45	911	137	59663	1634	63	21846	4352	81	13699
46	451	130	59793	1335	55	21901	605	79	13778
47	463	125	59918	4016	54	21955	4204	79	13857
48	466	117	60035	7	54	22009	427	74	13931
49	453	97	60132	2	48	22057	4206	74	14005
50	645	86	60218	26	47	22104	3809	70	14075
51	275	76	60294	940	47	22151	7408	68	14143
52	996	64	60358	2724	42	22193	3858	66	14209
53	901	63	60421	2815	39	22232	5352	66	14275
54	997	62	60483	5525	38	22270	2606	60	14335
55	111.	60	60543	5516	38	22308	455	59	14394
56	231	59	60602	2825	37	22345	566	59	14453
57	328	57	60659	1465	37	22382	7407	58	14511
58	425	55	60714	8	33	22415	2810	58	14569

Air Force Skill Codes(AFSC) containing 100 or more employees

* These codes sometimes end in a 2 digit suffix. Two periods stand for the suffix which could be any 2 numerals. In other references this designation is often XX rather than two periods.

** If these codes have less than four digits, it can be assumed it is preceded by enough zeros to make it a four digit code.

ž

Appendix B

Air Force Skill Codes(AFSC) containing 100 or more employees

1	Enliste	kd*	1	Officer*			Civilia	n* **	
	AFSC	Number	totals	AFSC	Number	totals	AFSC	Number	totals
!	Codes	in AFSC		Codes	In AFSC		Codes	in AFSC	
59	316	54	60768	2716	33	22448	3707	58	14627
60	392	53	60821	6054	30	22478	462	55	14682
61	908	47	60868	46	30	22508	4605	55	14737
62	464	47	60706	76	29	22411	5716	54	14565
63	251	44	60750	2875	27	22438	5439	51	14616
64	995	40	60790	8075	27	22465	3359	50	14666
!]								
	1								
	1								
				N/107,42	21	No. of	AFSC:	% abov	e 117
!	total for	≥ 100	93854	87.37%			117		1
[total for	≥ 80	95422	88.83%			135		115%
1	total for	r≥70	96020	89.39%			143		122%
	total for	≥ 50	97722	90.97%			172		147%

* These codes sometimes end in a 2 digit suffix. Two periods stand for the suffix which could be any 2 numerals. In other references this designation is often XX rather than two periods.

** If these codes have less than four digits, it can be assumed it is preceded by enough zeros to make it a four digit code.

Appendix C

Race. Ave	Annual Audio	No. of PTS (diseased) Risk	Stratum Crude	Standard weight(w) Risk	Stratum Adjusted	Conf. addends
1.white, 18-27	n	d	n/d	w 0.1942	(n/d)w	*(da) 1.da)
2.white,28-37				0.2994		
3.white,38-47				0.1253		
4.white, >47				0.0425		
5.black, 18-27	1			0.0207		
6.black,28-37				0.0345		
7.black_38-47				0.0143		
8.black, >47				0.0050		
9.0ther,18-27				0.0596		
10.0ther.28-37				0.1059		
11.0ther,38-47				0.0691		
12.0ther, >47				0.0294		
Crude Ris	k calculation:	·····		Sums =	Adjusted Rate	
				Confidence_ Interval		

ANALYSIS FOR EACH AFSC CODE

Confidence interval for direct adjustment:

Point Estimate +or- 1.96

ŝ

Ч. Т

 $\sqrt{\leq W_i \left(\frac{(d/n_{.})(l-d/n_{.})}{n_i}\right)}$

Appendix D

Race. Are	Annual Audio	No. of PTS (diseased) Risk	Stratuta Crude	Standard weight(w) Risk	Stratum Adjusted	Conf. addends
i.white, 18-27	4	0	0	0.1942	0	0
2.white, 28-37	41	1	0.024390	0.2994	0.007302	0.000052
3.white,38-47	31	3	0.096774	0.1253	0.012125	0.000044
4.white, >47	18	2	0.111111	0.0425	0.004722	0.000009
5.black,18-27	3	0	0	0.0207	0	0
6.black,28-37	3	1	0.333333	0.0345	0.011500	0.000088
7.black,38-47	5	0	0	0.0143	0	0
8.black, >47	3	0	0	0.0050	0	0
9.other, 18-27	31	1	0.322580	0.0596	0.001922	0.000003
10.other,28-37	71	4	0.056338	0.1059	0. 00-0 66	0.000008
11.other, 38-47	61	8	0.131147	0.0691	0.009062	0.000008
12.other, >47	18	3	0,166666	0.0294	0.004900	0.000006
Crude Risl	colculation:			Sums-	0.057501	0.000221
Totals	289	23	0.079584	Confidence_ Interval	0.086700 0.028302	

ANALYSIS FOR Civilian AFSC 3431

J.

Confidence interval for direct adjustment:

Point Estimate +or- 1.96
$$\bigvee W_i^2 \left(\frac{(d/n_i)(1-d/n_i)}{n_i} \right)$$

Appendix E

1		ENLIS	TED *			OFFIC	ER			CIVILI	AN *
	N	AFSC	AdjRisk		N	AFSC	AcjPisk		N	AFSC	AdjRisk
1	156	303	0.08302		219	1145	0.05026		108	8840	0.11388
2	315	901	0.06719		115	2295	0.04588		196	552	0.08613
3	130	451	0.06257		201	1425	0.04169		157	472	0.08369
4	230	426	0.06001		143	1595	0.03068		289	3431	0.0575
5	281	427	0.05061		277	9016	0.02643		216	6652	0.05637
6	210	566	0.05012		563	1555	0.0242		121	452	0.05444
7	1322	461	0.04785		267	1455	0.02406		445	4102	0.04003
8	163	602	0.04176		240	1025	0.02261		104	5413	0.03936
<u> </u>	600	411	0.04113		239	1565	0.02215		277	551	0.0376
10	822	431	0.03872		120	2865	0.02098		245	3703	0.03602
11	2083	458	0.03859	1	618	1745	0.01937		309	8255	0.0304
12	169	122	0.03783		335	1325	0.01877		114	2805	0.02971
13	1840	631	0.03642	1	152	1716	0.01858		2284	3806	0.02962
14	7886	452	0.03509	Ţ	3756	1115	0.01605		342	5378	0.0277
15	5993	457	0.03508	1	157	2255	0.01506		2225	8852	0.02516
16	907	542	0.03473	1	204	1435	0.01464		164	458	0.02506
17	390	116	0.03441	1	858	1545	0.01459		565	3414	0.02282
18	1042	552	0.03075	T	383	2225	0.01349		525	0081	0.02282
19	1265	551	0.03073	1	107	0036	0.01317		109	542	0.02259
20	2205	455	0.02999	T	596	1525	0.01261		215	2604	0.02253
21	382	423	0.0299	1	846	1535	0.01213	1	566	8602	0.02071
22	645	455	0.02939	T	1559	1055	0.01195	1	410	454	0.02063
23	208	115	0.02786	T	1639	1355	0.01005	1	482	2892	0.02031
24	8682	454	0.0272	T	1803	1065	0.00849	7	138	431.	0.01961
25	960	112	0.02606	T	192	1495	0.00839	1	205	6968	0.01946
26	3507	462	0.0249	T	658	1575	0.00781	Ì	187	545	0.01929
27	1004	472	0.02387	T	480	9356	0.0078	1	568	571	0.01865
28	935	605	0.02078	Ţ	2577	1045	0.00696	1	250	5402	0.01657
29	885	545	0.02007	t	119	4024	0.00665	1	118	4604	0.01446
30	321	753	0.01893	T	341	1825	0.0063	1	101	8800	0.01317
31	2645	114	0.01825	T	609	1235	0.0058	1	200	3500	0.00708

Adjusted Risk for all AFSCs with 100 or more individuals

* These codes sometimes end in a 2 digit suffix. Two periods stand for the suffix which could be any 2 numerals. In other references this designation is often XX rather than two periods. Appendix E

Adjusted Risk for all AFSCs with 100 or more individuals

۲.- A

		ENLIS	TED *			OFFIC	ER			CIVILI	AN *
	N	AFSC	AdjRisk		N	AFSC	AdjRisk		N	AFSC	AdjRisk
32	1319	272	0.01743		149	2245	0.00543	Γ	191	457	0.00548
33	2:51	113	0.01677		161	9766	0.00533		223	8268	0.00255
34	1778	811	0.01637		148	1585	0.00415				
35	1960	571	0.01604		193	0006	0				
36	271	361	0.01391		146	1315	0				
37	271	117	0.01053								
38	264	207	0.00966					Γ			
39	280	118	0.00944								
40	513	208	0.00745								
41	137	911	0.0068								
42	229	871	0.0055								
43	264	304	0.00512								
44	218	273	0.00299								
45	1007	902	0.00267								
46	147	872	0								
47	125	463	0								
48	:17	466	Û	ļ							

* These codes sometimes end in a 2 digit suffix. Two periods stand for the suffix which could be any 2 numerals. In other references this designation is often XX rather than two periods. •

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ACDENDS
WHITE	18-27	e	0	0.000024000	0.1942	000000000 0.	0.60600000
	28-37	30	ŝ	0.16666667	0.2994	0.049900000	0.000415002
	36-47	29	6	0.137931034	0.1253	0.017282759	0.000064374
	> 47	61	2	0.105263158	0.0425	0.004473684	0.000008954
BLACK	18-27	o	o	0.00000000	0.0207	0.00000000	0.00000000
	28-37	Q	0	0.00000000.0	0.0345	0.000000000	0.000000000
	38-47	n	0	0.00000000	0.0143	0.000000000	0.00000000
	~ 47	9	0	0.00000000	0.0050	0.00000000	000000000000000
OTHER	18-27	o	0	0.00000000	0.0596	0.00000000	0,00000000
	28-37	4	-	U.25000000	0.1059	0.026475000	0.000525694
	38-47	7	-	0.142857143	0.0691	0.009871429	0.000083524
	× 47	10	7	0,20000000	0.0294	0.005880000	0.000013830
TOTALS		108	15	0.138888889		0.113882871	0.001111377
			CONFIDENCE	INTERVAL = 0.	179224028(+)		
				0.0	048541714(-		
				 0	048541714(-		

ANALVSIS FOR CIVILIAN AFSC: 8840

. .

. . .

これをいう かったい しんしょう 教育学校教授 アイ・マイン 一次に 日本の

9 24 J. C. W. 4. 4.

4-21

· 역 인정이 아직 아이 이상이 아이

33

Appendix F

Ten AFSCs(Air Force Skill Code) with the Highest Risk of developing PTS

ANALVSIS FOR CIVILIAN AFSC: 552XX

F

•

.

4.

1.1

an state and the

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	1	0	0.00000000	0.1942	0.000000000	0.00000000
	28-37	11 -	2	0.181818182	0.2994	0.054436364	0.001212266
	38-47	5.4	е	0.055555556	0.1253	0.00696111	0.000015255
	> 47	53	8	0.150943396	0.0425	0.006415094	0.000004368
BLACK	18-27	-	0	0.00000000.0	0.4207	0.00000000	0.000000000
	28-37	2	0	0.00000000	0.0345	0.00000000	0.000000000
	38-47	4	٥	0.000000000	0.0143	0.00000000	0.000000000
	> 47	n	0	0.00000000	0.0050	0.00000000.0	0.00000000
OTHER	18-27	2	0	0.00000000	0.0596	0.00000000	0.00000000
	28-37	6	-	0.111111110	0.1059	0.011766667	0.000123071
	38-47	25	-	0.04000000	0.0691	0.002764000	0.000007334
	~ 47	31	4	0.129032258	0.0294	0.003793548	0.000003134
TOTALS		196	61	0.096938776		0.086136784	0.001365427

0.013711475(-)

CONFIDENCE INTERVAL = 0.158562094(+)

ANALYSIS FOR CIVILIAN AFSC: 472XX

• • •

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	2	٥	0.00000000	0.1942	00000000000000	0.00000000
	28-37		-	1 80606060 . 0	0.2994	0.027218182	0.000673481
	38-47	39	4	0.102564103	0.1253	0.012851282	0.000037054
	> 47	55	12	0.218181818	0.0425	0.009272727	0.000005602
BLACK	18-27	0	0	0.000000000	0.0207	0.00000000	0.00000000
	28-37	-	0	0.000000000	0.0345	0.00000000	0.000000000
	38-47	8	0	0.00000000	0.0143	0.00000000	0.000000000
	~ 47	-	0	0,00000000	0.0050	0.00000000	0.00000000
OTHER	18-27	0	0	0.00000000	0.0596	0.00000000	0,00000000
	28-37	4	-	0.25000000	0.1059	0.026475000	0.000525694
	38-47	15	-	0.066666667	0.0691	0.004606667	C.000019807
	47	18	2	0.11111111.0	0.0294	0.003266667	0.000004743
TOTALS		157	12	0.133757962		0.083690524	0.001266381

0.013941482(-)

CONFIDENCE INTERVAL = 0.153439567(+)

ANALVSIS FOR ENLISTED AFSC: 303XX

A STATE OF STATE

1. 5 5.

1

2

2 k 1

4

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFICZNCE ADDENDS
WHITE	18-27	53	o	0.00000000	0.1942	0,00000000 O	0,00000000,0
	28-37	56	0	0.00000000	0.2994	0.00000000	0.00000000.0
	38-47	8	-	0.11111111.0	0.1253	0.01392222	0.000172292
	> 47	-	0	0.00000000.0	0.0425	0.00000000	0.000000000.0
BLACK	18-27	6	0	0.00000000	0.0207	0.00000000	0.00000000
	28-37	٩	0	0.00000000	0.0345	0.00000000	0.00000000.0
	38-47	0	0	0.00000000	0.0143	0.00000000.0	0.00000000.0
	74 <	0	o	D.00000000	0.0050	0.00000000	0.00000000.0
отнек	18-27	ß	0	0.000000000	0.0596	0.000000000	0.00000000
	28-37	13	0	0.000000000	0.1059	0.00000000	0.00000000
	36-47	-	-	1.000000000	0.0691	0.069100000	0,00000000
	~ 47	0	0	0.00000000	0.0294	0.00000000	0.000000000
TOTALS		156	2	0.012820513		0.08302222	0.000172292

0.057295268(-)

CONFIDENCE INTERVAL = 0.108749176(+)

ANALYSIS FOR ENLISTED AFSC: 901XX

.

.

4

.

지수

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	79	0	0.00000000.0	0.1942	0.00000000	0.00000000
	28-37	5 DB	-	0.009259259	0.2994	0.002772722	0.000007614
	38-47	25	ß	0.20000000	0.1253	0.025060000	0.000100481
	> 47	7	0	0.00000000.0	0.0425	0.00000000	0.00000000
BLACK	18-27	14	0	0.00000000	0.0207	0.00000000	0.00000000
	28-37	24	٥	0.00060000	0.0345	0.00000000	0.000000000
	38-47	۵	0	0,00000000	0.0143	0.00000000	0.00000000
	> 47	0	0	0.00000000	0.2050	0.00000000	0.00000000
OTHER	18-27	34	0	0.00000000.0	0,0596	0.00000000	0.00000000
	28-37	22	-	0.045454545	0.1059	0.004813636	0.000022118
	38-47	2	-	0.500000000	0.0691	0.034550000	0.000596851
	> 47	o	٥	0.00000000	0.0294	0.00000000	0.00000000
TOTALS		315	æ	0.025396825		0.067195859	0.000727064

0.014346184(-)

CONFIDENCE INTERVAL = 0.120045533(+)

0

ANALVSIS FOR ENLISTED AFSC: 451XX

4

• •

AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
18-27	25	0	0.00000000	0.1942	0.00000000.0	0.0000000000000
28-37	53	2	0.037735849	0.2994	0.011298113	0.000061415
38-47	25	ß	0.20000000	0.1253	0.025060000	0.000100481
> 47	2	a	0.00000000000	0,0425	0.00000000	0.00000000
18-27	e	o	0.00000000	0.0207	0,00000000	0.00000000
28-37	2	0	0,00000000.0	0.0345	0,00000000	0.000000000
38-47	-	C	0.000000000	0.0143	0.00000000	0.000000000
1 47	0	0	0.00000000	0.0050	0.00000000	0,00000000
18-27	2	0	0.00000000	0.0596	0.00000000	00000000000000
28-37	ŋ	0	0.00000000.0	0.1059	0.00000000	0.00000000
38-47	g	-	0.16666667	0.0691	0.011516667	0.000110528
< 47	3	-	0.50000000	0,0294	0.014700000	0.000108045
	130	G	0.069230769		0.062574780	0.000380469

0.024343784(-)

CONFIDENCE INTERVAL = 0.100805776(+)

E.

Progr N

-

....

ANALVSIS FOR ENLISTED AFSC: 426XX

. . .

4

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM Adjusted risk	CONFIDENCE ADDENDS
WHITE	18-27	24	-	0.041666667	0.1942	0.008091667	0.000062747
	28-37	62		0.036585366	0.2994	0.010953659	0.000038531
	38-47	39	3	0.076923077	0.1253	0.009638462	0.000028585
	~ 47	11	2	0.181818182	0.0425	0.007727273	0.000024427
BLACK	18-27	-	0	0.000000000	0.0207	0.000000000000	0.000000000
	28-37	11	0	0.00000000	0.0345	0.000000000	0.000000000
	38-47	ų	٥	0.00000000	0.0143	0.000000000	0.00000000
	~ 47	2	0	0.00000000	0.0050	0.00000000	0.000000000
OTHER	18-27	2	-	0.142857143	0.0596	0.008514286	0.000062137
	28-37	29	0	0.000000000	0.1659	0.00000000	0.000000000
	38-47	15	2	0.133333333	0.0691	0.009213333	0.000036784
	~ 47	α	-	0.20000000	0.0294	0.005880000	0.000027660
TOTALS		230	61	0.056521739		0.060018679	0.000280870

0.021170703(--)

CONFIDENCE INTERVAL = 0.092866654(+)

ANALYSIS FOR CIVILIAN AFSC: 3431

and differences

• • •

¢

CONFIDENCE	AUDENDS	u . uuuuuuuu	0.000052025	0.000044269	11660000000000000000000000000000000000	0.000088167	0,000000000	0.000000000	0.000003577	0,000008398	0.0000080	C.000006669	0.000221935
STRATUM AD HISTED DI SU	D DODDODOD		0.010101000	0.00472225	0.00000000	0.011500000	0.00000000	0.00000000	0.001922581	0.005966197	0.009052295	0.004900000	0.057501541
STANDARD WEIGHT	0.1942	0 2094	0 1253	0.0425	0.0207	0.0345	0.0143	0,0050	0.0596	0.1059	0.0691	0.0294	
STRATUM Crude Risk	0.000000000	6.024390244	0.096774194	0.1111111	0.000000000	0.333333333	0,00000000	0,00000000	0.032258065	0,056338028	0.131147541	0.166666667	0.07958477\$
NUM WITH PTS	٥	-	ũ	2	0	-	0	٥	-	4	G)	ũ	23
NUM ANNUAL AUDIOGRAMS	4	141	31	18	ო	e	5	e,	31	12	61	81	289
AGE	18-27	28-37	38-47	> 47	18-27	28-37	38-47	~ 47	18-27	28-37	38-47	> 47	
RACE	WHITE				BLACK			,	0 THER				TOTALS

0.028302504(-)

alan yang baran salah salah salah 🦗 alam dalam yang baran kata baran yang bara sala yang baran salah yang

CONFIDENCE INTERVAL = 0.086700577(+)

ANALYSIS FOR CIVILIAN AFSC: 6652

÷,

S . 16. 5

Ę

1 3 1

¢

ONF I DENCE ADDENDS	.000413867	.000018033	.000013352	.000012640	.000000000	.000000000	.000000000	.000000000	.000000000	166170000.	. 000000000	.00020008	.000549291
STRATUM CUUSTED RISK	0.02157778 0	0.004277143 0	0.010081609 0	0.006710526 0	0.000000000	0,00000000	0.000000000	0.000000000	0.00000000 0	0.008825000 0	0.00000000 0	0.004900000	0.056372056 0
STANDARD WEIGHT	0.1942	0.2994	0.1253	0.0425	0.0207	0.0345	0.0143	0.0050	0.0596	0.1059	0.0691	0.0294	
STRA.UM CRUDE RISK	0.11111111	0.014285714	0.080459770	0.157894737	0.00000000	0.000000000	0.00000000	0.00000000	0,000000000	0.083333333	0.00000000	0.166666667	0.064814815
NUM WITH PTS	~	-	7	9	٥	0	O	0	0	-	0	-	14
NUM ANNUAL AUDIOGRAMS	đ	. 70	87	19	0	-	S	0	0	12	٢	ŋ	216
AGE	18-27	16-82	38-47	> 47	18-27	26-37	38-47	> 47	18-27	28-37	38-47	× 47	
RACE	WHITE				BLACK				OTHER				TOTALS

0.010435639(-)

CONFIDENCE INTERVAL = 0.102308474(+)

ANALYSIS FOR CIVILIAN AFSC: 452XX

......

: :-::

to a to a to

. . .

.

RACE	46 E	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	J.	o	0.000000000	0.1942	0.00000000	0.00000000
	28-37	27	5	0.074074074	0.2994	0.022177778	0.000227710
	38-47	24	e	0.125000000	0.1253	0.015662500	0.000071550
	> 47	23	4	0.173913043	0.0425	0.007391304	0.000011283
BLACK	18-27	Q	0	0.00000000	0.0207	0.00000000	0.000000000
	28-37	2	0	0.00000000.0	0.0345	0.00000000	0.000000000
	38-47	0	o	0,00000000	0.0143	0.00000000	0.0000000000000
	> 47	2	a	0.000000000	G.0050	0.00000000	0.00000000
OTHER	18-27	2	0	0.00000000	0.0596	0.00000000	0.000000000.0
	28-37	1 5	0	0.00000000.0	0.1059	0.00000000	0.00000000
	38-47	15	2	0.133333333	0.0691	0.009213333	0.000036784
	5 47	01	0	0.00000000	0,0294	0.00000000	0.000000000
TOTALS		121	Ξ	1 60606060 . 0		0.054444915	0.000347326

(-)66691610.0

CONFIDENCE INTERVAL = 0.090972832(+)

τ....

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PIS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WH1TE	18-27	44	0	0.000000000	0.1942	0.00000000	0.00000000
	28-37	1	0	0.000000000	0.2994	0.00000000	0.00000000
	38-47	ß	0	0000000000000	0.1253	0.00000000	0.00000000
	× 47	٥	0	0.00000000.0	0.0425	0.00000000	0.00000000
BLACK	18-27	9	0	0.00000000.0	0.0207	0.00000000	0.60000000
	28-37	-	0	0,00000000	0.0345	0.00000000	0.000000000
	38-47	-	a	0.00000000	0.0143	0.00000000	0.00000000
	2 47	0	o	0.00000000.0	0.0050	0.00000000	0.000000000
OTHER	18-27	6	0	0.00000000	0.0596	0.00000000	0.00000000
	28-37	Q	0	0.000000000	0.1059	0.00000000	0.00000000
	38-47	-	a	0,00000000	0.0691	0.00000000.0	0.00000000
	> 47	0	0	0000000000000	0.0294	0.00000000	0.00000000
TOTALS		117	٥	0.00000000		0.00000000	0.00000000
			CONFIDENCE	INTERVAL = 0.0	(+)00000000		
) 0000000000		

43

ANALVSIS FOR ENLISTED AFSC: 466XX

18. **12** -

G. 0000000000 (-)

Appendix G

Ten AFSCs(Air Force Skill Code) with the Lowest Risk of developing PTS

ANALVSIS FOR ENLISTED AFSC: 463XX

121 - 2121 BEER AT 81 - 121

ā

C. Succession

 ٠

.

1

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PIS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM Adjusted risk	CONFIDENCE ADDENDS
WHITE	18-27	62	٥	0.00000000	0.1942	0.00000000	0.000000000
	28-31	30	٥	0.000000000	0.2994	0.00000000.0	0.00000000
	38-47	1	0	0.00000000	0.1253	0.00000000.0	0 . 000000000
	> 47	0	0	0.00000000	0.0425	0.00000000.0	0.000000000
BLACK	18-21	-	0	0.000000000	0.0207	0.00000000.0	0.00000000
	28-37	ę	0	0.000000000	0.0345	0,00000000	0.00000000
	38-47	-	0	0,00000000	0.0143	0.00000000.0	0.00000000
	> 47	0	0	0.00000000	0:00:0	0.000000000	0.00000000
OTHER	18-27	13	C	0.00000000	0.0596	0.000000000	0.000000000
	28-37	7	0	0.00000000	0.1059	0.00000000	0.00000000
	38-47	-	0	0.00000000	0.0691	0.00000000	0.000000000
	× 47	0	0	0.000000000	0.0294	0.00000000	0.00000000
TOTALS		125	G	0.00000000		0.00000000	0.00000000

(-)000000000).0

CONFIDENCE INTERVAL = 0.40000000(+)

Į

1 15

ANALVSIS FOR ENLISTED AFSC: 872XX

and a state of the second state

· ·

NF I DENCE DOENDS	000000000	000000000	000000000	000000000	000000000	000000000	000000000	000000000	000000000	000000000	000009600	000000000	000000000
03 <	0.	<u>о</u>	0	<u>.</u>	0.	0.	0	ō	0	0	0	0	0
STRATUM ADJUSTED RISK	000000000000000000000000000000000000000	0.00000000	0.00000000	0.00000000000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000	0.00000000
STANDARD WEIGHT	0.1942	0.2994	0.1253	0.0425	0.0207	0.0345	0.0143	0.0050	0.0596	0.1050	0.0691	0.0294	
STRATUN CRUDE RISK	0.00000000	0.00000000	0.000000000	D.00000000	0.000000000	0.00000000.0	0.00000000.0	0.00000000	0.000000000	0.000000000	0.00000000.0	0.00000000.6	0.00000000.0
NUM WITH PTS	0	0	C	٥	0	0	٥	0	0	O	0	C	٥
NUM ANNUAL AUDIOGRAMS	Q	69	47	**	o	N	4	0	-	ej	8	-	147
AGC	18-27	28-37	38-47	× 47	18-27	28- <i>.</i> 17	38-47	> 47	18-27	28-37	38-47	~ 47	
RACE	WHITE				BLACK				OTHER				TOTALS

CONFIDENCE INTERVAL = 0.00000000(+)

ANALYSIS FOR OFFICER AFSC: 1315

ı

÷ .

.

ž. .

and the state of the

The last provide the second

: a > *

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PIS	STRATUM Crude RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	11	a	0.00000000	0.1942	0.00000000	0.00000000.0
	28-37	76	0	0.00000000	0.2994	0.00000000	0.00000000.0
	38-47	25	0	0,000000000	0.1253	0.00000000	0.00000000
	> 47	-	0	0.00000000	0.0425	0.00000000	0.00000000.0
BLACK	18-27	0	0	0,00000000	0.0207	0.00000000	000000000.0
	28-37	E	0	0.00000000	0.0345	0.00000000	0.00000000
	38-47	-	-	0.00000000	0.0143	0.00000000	0.00000000
	> 47	0	0	0.00000000.0	0.0050	0.00000000	0.00000000
ОТНЕЯ	18-27	ũ	0	0.000000000	0.0596	0.00000000	0.00000000
	28-37	17	0	0.00000000.0	0.1059	0.00000000	0.00000000000
	38-47	8	O	0.000000000	0.0691	0.000000000	0.00000000
	7 4 7	-	Ð	0,00000000	0.0294	0.00000000	0.00000000
TOTALS		146	٥	0.000000000		0.00000000.0	000000000000000000000000000000000000000

0.000000000.0

CONFIDENCE INTERVAL = 0.000000000(+)

_

A &

ANALVSIS FOR OFFICER AFSU: 0006

• • •

-

-

•

.

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WH1 TE	18-27	102	J	0.000000000	0.1942	0.00000000	0.00000000
	28-37	15	۵	0,000000000	U.2994	0,00000000	0.0000000000
	38-47	8	o	0.00000000	0.1253	0,00000000	0.000000000
	47	4	٥	0.00000000	0.0425	0.00000000.0	0.00000000
BLACK	18-27	-	0	0.00000000	0.0207	0,00000000	0.000000000
	28-37	0	0	0.00000000	0.0345	0 00000000	0.00000000
	38-47	0	0	0.00000000	0.0143	0.000000000	0.00000000.0
	> 47	0	0	0.00000000	0.0050	0.00000000.0	0.00000000
OTHER	18-27	34	0	0.000000000.0	0.0596	0.00000000.0	0.000000000
	28-37	26	0	0,00000000	0.1059	0.00000000.0	0.000000000
	38-47	2	0	0.00000000.0	0,0691	0.00000000	0.00000000
	> 47	-	0	0.00000000	0.0294	0,00000000	0.000000000
TUTALS		661	٥	0.00000000		0,00000000	0.00000000

CONFIDENCE INTERVAL = 0.000009000(+) 0.000000000(-) -

ANALVSIS FOR CIVILIAN AFSC: 8268

.....

ī

. . .

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	S	a	0.00000000	0.1942	0.00000000	0.00000000
	28-37	. 65	0	0.00000000	0.2994	0.00000000	0.00000000
	38-47	48	0	0.00000000	0.1253	0.00000000000000	0.00000000
	> 47	42	o	0.00000000	0.6425	0.000000000	0.00000000
BLACK	16-27	0	Ċ	0.0000000000000	0.0207	0.00000000	0,00000000
	28-37	2	0	0,00000000	0.0345	0.00000000	0.00000000
	38-47	5	0	0.00000000	0.0143	0.00000000	0.00000000
	> 47	2	o	0.00000000	0.0050	0.00000000	0.00000000
OTHER	18-27	2	C	0.00000000	0.0596	0.00000000	0.000000000
	28-37	14	٥	0.00000000	0.1059	0.00000000	0.000000000
	38-47	27	-	0.037037037	0.0691	0.002559259	0.000006307
	~ 47	17	o	0.00000000	6.0294	0.00000000	0,00000000
TOTALS		223	-	0,004464305		0.002559259	0.00006307

-.002363121(-)

CONFIDENCE INTERVAL = 0.007481639(+)

AWALVSIS FOR ENLISTED AFSC: 902XX

ないがい。

• •

,

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM WITH PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	31	0	0.00000000	0.1942	0.00000000	0.000000000
	28-37	- 67	0	0.00000000	0.2994	0.00000000	0.000000000
	38-47	11	0	0.00000000	0.1253	0.00000000	0.00000000
	> 47	ব	0	0.00000000	0.0425	0.00000000.0	0.00000000
BLACK	18-27	9	0	0.00000000	0.0207	0.00000000	0.00000000
	28-37	13	0	0.000000000	0,0345	0.00000000	0.000000000
	36-47	Ģ	0	0.00000000	0.0143	0.00000000	0.000000000
	> 47	0	O	0.00000000	0.0050	0.00000000	0.000000000
OTHER	18-27	296	-	0.003378378	0.0596	0.000201351	0.00000040
	28-37	321	0	0.00000000	0.1059	0.00000000	0.000000000
	38-47	197	4	0.020304569	0.0691	0.001403046	0.000000482
	> 47	55	7	0.036363636	0.0294	0.001069091	0.000000551
TOTALS		1,007	٢	0.006951341		0.002673488	0.000001073

0.000642977(-)

CONFIDENCE INTERVAL = 0.004703999(+)

ANALVSIS FOR ENLISTED AFSC: 273XX

nut i

1 2 1

.

RACE	AGE	NUM ANNUAL AUDIOGRAMS	NUM W.TH	STRATUM CRUDE 11SK	STANDARD WEIGHT	STRATUM Adjusted risk	CONFIDENCE ADDENDS
WHITE	18-27	61	a	0,00000000	0.1942	0.00000000	0.00000000
	28-37	·100	-	0.01000000	0.2994	0.002994000	0.000008874
	38-47	61	0	0.00000000	0.1253	0.00000000	0.00000000.0
	~ 47	-	o	0.000000000	0.0425	0.00000000	00000000000000
BLACK	13-27	-	0	0.0000000000000000000000000000000000000	0.0207	0.000000000	0.000000000000
	28-37	3	C	0.000000000	0.0345	0.00000000	0.000000000
	38-47	-	0	0.00000000	0.0143	0.00000000	0.00000000
	> 47	٥	O	0.00000000	0.0050	0,00000000	0.00000000.0
OTHER	18-27	11	0	0.000000000	0.0596	0.000000000	0.000000000
	28-37	15	0	0,00000000	0.1059	0.00000000	0.00000000
	38-47	e	0	0.000000000	0.0691	0.00000000	0.000000000
	- 47	0	0	0.00000000	0.0294	0.000000000	0.00000000
TOTALS		218	-	0.004587156		0.002994000	0.000038874

-.002844825(-)

CONFIDENCE INTERVAL = 0.000832825(+)

ANALVSIS FOR OFFICER AFSC: 1585

l.

ないないのないないないないで、こう

2 a 3

.

RACE	AGE	NUM ANIVUAL AUDIOGRAMS	NUM WITH PIS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM Adjusted risk	CONFIDENCE ADDENDS
WHITE	18-27		0	0.00000000	0.1942	0.00000000	0.00000000
	28-37	72	-	0.013888889	0.2994	0.004158333	0.000017052
	38-47	15	0	0.00000000	0.1253	0.00000000	0.00000000
	> 47	र	0	0.00000000	0.0425	0.00000000	0,00000000
BLACK	18-27	0	٥	0.00000000	0.0207	0.00000000	0.00000000
	28-37	e	0	0.00000000	0.0345	0.00000000	0.00000000
	38-47	0	٥	0,00000000	0.0143	0.00000000	0.00000000
	> 47	0	o	0.00000000	0.0050	0.00000000	0.00000000
OTHER	18-27	0	0	0.00000000	0.0596	0.00000000	0.00000000
	28-37	50	o	0.000000000	0.1059	0.00000000	0.000000000
	38-47	12	0	0.00000000	0.0691	0.00000000	0.000000000
	× 47	-	O	0.00000000.0	0.0294	0.00000000	0.00000000
TOTALS		1 48	-	0.006756757		0.004158333	0.000017052

-.003935203(-)

CONFIDENCE INTERVAL = 0.012251869(+)

51

ANALYSIS FOR ENLISTED AFSC: 304XX

e • • •

A STATE OF A

States and the states and the

•

-

2000

RACE	AGE	NUM ANNUAL AUDIOGRAMS	HTIW MUN PTS	STRATUM CRUDE RISK	STANDARD WEIGHT	STRATUM ADJUSTED RISK	CONFIDENCE ADDENDS
WHITE	18-27	67	0	0.00000000.0	0.1942	0.0000000000000000	0.00000000
	28-37	117	2	0.017094017	0.2994	0,005117949	0.000012873
	38-47	12	a	0.00000000	0,1253	0.00000000	0 . 000000000
	> 47	a	٥	0,00000000	0.0425	0.00000000	0.00000000
BLACK	18-27	13	0	0.00000000.0	0.0207	0.00000000	0.00000000
	28-37	15	0	0,00000000	0.0345	0.00000000	0,00000000
	38-47	-	0	0.00000000.0	0.0143	0.00000000	0.00000000.0
	× 47	0	٥	0.00000000	0.0050	0.000000000	0.00000000
OTHER	18-27	61	0	0.000000000	0.0596	0.00000000	0.00000000
	28-37	15	0	0.00000000	0.1059	0.00000000	0000000000000
	36-47	5	Q	0.00000000	0.0691	0.00000000	0.00000000.0
	> 47	0	0	0.00000000	0.0284	0.00000000	0.00000000
TOTALS		264	8	0.007575758		0.005117949	0.000012873

-.001914280(-)

CONFIDENCE INTERVAL = 0.012150178(+)

••••

....

Bibliography

۰.

- 1. United States Air Force. Air Force Regulation 161-20. Hearing Conservation Program 1991.
- 2. Grayson JK. Previous hearing loss and susceptibility to future permanent threshold shifts. *Military Medicine* 1992; 5: 248,249.
- 3. United States Air Force. Air Force Annual Hearing Conservation Report 1991.
- 4. Department of Labor Statistics. Washington, D.C.; personal communication with Ralph Slighter, OWCP, on 29Mar93.
- 5. Department of Veterans Affairs. Washington, D.C.; personal communication with Mike Wells(VA statistician) on 24May93.
- 6. Franks JR, Davis RR, Kreig EF. Analysis of a hearing conservation program data base: factors other than workplace noise. *Ear and Hearing* 1989; 10(5): 273-280.
- 7. Sataloff RT, Sataloff J. Occupational Hearing Loss. New York, New York: Marcel Dekker, Inc., 1993.
- 8. Harrison RK. Hearing conservation implementing and evaluating a program. AAOHN Journal 1989; 37: No.4, 107-111.
- 9. Fosbroke J. Practical observations on the pathology and treatment of deafness. Lancet 1830 & 1831; 1: 645.
- Barone JA, Peters JM, Garabrant DH, Bernstein L, Krebsbach R. Smoking as a risk factor in noise-induced hearing loss. *Journal of Occupational Medicine* 1987; 29(9): 741-745.
- 11. CDC. Leading Work-Related Diseases and Injuries United States. MMWR 1983; 32(2):24-32.
- 12. Barrenas ML, Lindgren F. The influence of eye colour on susceptibility to TTS in humans. British Journal of Audiology 1991; 25: 303-307.
- 13. Harrison RK. Hearing conservation implementing and evaluating a program. AAOHN Journal 1989; 37: No.4, 107-111.
- 14. Attias J, Pratt H. Auditory-evoked potential correlates of susceptibility to noiseinduced hearing loss. Audiology 1985; 24: 149-156.
- 15. Pyykko I, Koskimies K, Starck J, Pekkarinen J, Farkkila M, Inaba R. Risk factors in the genesis of sensorineural hearing loss in Finnish forestry workers. British Journal of Industrial Medicine 1989; 46: 439-446.
- 16. Barrenas ML, Lindgren F. The influence of inner ear melanin on susceptibility to TTS in humans. Scandinavian Audiolgy 1990; 19: 97-102.

- 17. Trune DR, Mitchell C, Phillips DS. The relative importance of head size, gender and age on the auditory brainstem response. Hearing Research 1988; 32: 165-174.
- 18. Bauer P, Korpert K, Neuberger M, Raber A, Schwetz F. Risk factors for hearing loss at different frequencies in a population of 47,388 noise-exposed workers. Journal of the Acoustical Society of America 1991; 90(6): 3086-3098.
- 19. LaDou J. Occupational Medicine. Norwalk, Connecticut: Appleton and Lange, 1990.
- 20. Lutman ME, Spencer HS. Occupational noise and demographic factors in hearing. ACTA Oto-laryngologica 1991; Suppl. 476: 74-84.
- 21. Ribak J. The association of age, flying time, and aircraft type with hearing loss of aircrew in the Israeli Air Force. Aviation, Space, and Environmental Medicine 1985; 56: 322-327.
- 22. Fitzpatrick DT. An analysis of noise-induced hearing loss in Army helicopter pilots. Aviation, Space, and Environmental Medicine 1988; 59(10): 937-941.
- 23. Royster LH, Royster JD, Thomas WG. Representative hearing levels by race and sex in North Carolina industry. Journal of the Acoustical Society of America 1980; 68: 551-566.
- 24. Johnson DL. Field studies: industrial exposures. Journal of the Acoustical Society of America 1991; 90(1): 170-174.

1. K 1. 1

- 25. Bunch CC, Raiford TS. Race and sex variations in auditory acuity. Archives of Otolaryngology 1931; 13: 423-434.
- 26. Karsai L, Bergman M Choo YB. Hearing in ethnically different longshoremen. Archives of Otolaryngology 1972; 96: 499-504.
- 27. LaFerriere KA, Kaufman AI, Hawkins JE, Johnson L-G. Melanocytes of the vestibular labyrinth and their relationship to microvasculature. Annals of Otology, Rhinology, and Larngology 1974; 83: 685-694.
- 28. Hood JD, Poole JP, Freedman L. The influence of eye colour upon temporary threshold shift. Audiology 1976; 15: 449-464.
- Carter NL, Keen K, Waugh RL, Murray N, Bulteau VG. The relations of eye colour and smoking to noise-induced permanent threshold shift. *Audiology* 1981; 20: 335-346.
- 30. Karlovich RS. Comments on the relations between auditory fatigue and iris pigmentation. Audiology 1975; 14: 238-243.
- 31. Hedin CA. Smoker's Melanosis may explain the lower hearing loss and lower frequency of Parkinson's Disease found among tobacco smokers a new hypothesis. *Medical Hypothesis* 1991; 35: 247-249.

- 32. Carlin MF, McCroskey RL. Is eye color a predictor of noise-induced hearing loss? Ear and Hearing 1980; 1(4): 191-196.
- 33. Pirila T. Sorri M, Jounio-Ervasti K, Sipilå P, Karjalainen H. Hearing asymmetry among occupationally noise-exposed men and women under 60 years of age. Scandinavian Audiolgy 1991; 20: 217-222.
- 34. United States Air Force. Air Force Annual Hearing Conservation Report 1992.

1

. મંં, ≭. .

- 35. Ahlbom A, Norell S. Introduction to Modern Epidemiology. Chestnut Hill, MA: Epidemiology Resources Inc., 1984.
- 37. Hennekens CH, Buring JE. Epidemiology in medicine. Boston: Little, Brown and Company, 1987.
- 38. Rothman KJ. Modern Epidemiology. Boston: Little, Brown and Company, 1986.

Vita

Jim Alan Davis was born in Crystal City, Texas, on August 21, 1955, the son of Bert and Billie Margaret Davis. After graduating from Boling High School in 1973, he attended Wharton County Junior College, Wharton, Texas, until December 1974. He then transferred to Texas A&M, College Station, Texas, and graduated with a Bachelor of Science in Veterinary Medicine in 1977. His education continued at Texas A&M until he graduated with a Doctorate of Veterinary Medicine in August 1978. From 1978 to 1981 he was in the Army Veterinary Corps and was stationed at Ft. McClellan, Alabama, and then Fort Sam Houston, Texas. He left the military in 1981 and went into private veterinary practice in Houston, Texas until 1987. In August 1987 he joined the Air Force and has served as a Military Public Health officer to the present. He was statione: 'at Loring AFB, Maine from 1987 to 1991; Ellsworth AFB, South Dakota, from 1991 to 1992; and University of Texas at Houston as a student from 1992 to 1994. He is married to Brenda Darlene Davis and has three children : J. Andrew, age 19; Kristin, age 15; and Joshua, age 13.

> Permanent address : Rr. 1 Box 84A Boling, Texas 77420

This thesis was typed by the author.