GL-TR-90-0551

Scintillation and In-situ Irregularity Parameters in the Auroral Oval and Polar Cap

AD-A282 459

Eileen MacKenzie Sunanda Basu Paul Pruneau

Boston College Institute for Space Research Newton MA 02159



1. J. M.

28 December 1990

Scientific Report No. 1

Approved for public release; distribution unlimited

15402 8 94

GEOPHYSICS LABORATORY AIR FORCE SYSTEMS COMMAND UNITED STATES AIR FORCE HANSCOM AIR FORCE BASE, MASSACHUSETTS 01731-5000

DTIC QUALITY INSPECTED 1

Addension For MTL: GRADI

DTIC TAB

By.

Dist

A-1

Unsmiowing

Justification

Distribution/

Availability Codes Avail and/or

Special



5

П

AUG 0 3 1994

1

REPRODUCED BY U.S. DEPARTMENT OF COMMERCE NATIONAL TECHNICAL INFORMATION SERVICE SPRINGFIELD, VA 22161 "This technical report has been reviewed and is approved for publication"

Contract Manager

endan

Acting Branch Chief

FOR THE COMMANDER

ROBERT A. SKRIVANEK Division Director

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. all others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify GL/IMA, Hanscom AFB, MA 01731. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

UNCL	ASS	IF	IED
------	-----	----	-----

SECURITY	CLASSIFICAT									
				REPORT DOCUM	ENTATION PAG	E				
1. REPORT SECURITY CLASSIFICATION			IN RESTRICTIVE	Ib. RESTRICTIVE MARKINGS						
UNCLASSIFIED			None							
7. SECURITY CLASSIFICATION AUTHORITY			D. DISTRIBUTION/AVAILABILITY OF REPORT							
26. DECLASSIFICATION/DOWNGRADING SCHEDULE			Approved for public release; distribution unlimited.							
A PERFORMING ORGANIZATION REPORT NUMBER(S)				5. MONITORING ORGANIZATION REPORT NUMBER(5) GL-TR-90-0351						
6. NAME OF PERFORMING ORGANIZATION St. OFFICE SYMBOL			Te. NAME OF MORITORING ORGANIZATION							
Institute for Space Research				Geophysics Laboratory						
6c. ADDIRESS (City, State and ZIP Code)			7h, ADDRESS (City, State and ZIP Code)							
Newton MA 02159				Hanscom AFB MA 01731-5000						
. NAME OF FUNDING/SPONSORING Bb. OFFICE SYMBOL			B. OFFICE SYMBOL	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER						
ORGANIZATION · (If applicable)		(If applicable)	F19628-90-K-0007							
Bc. ADDRESS (City, Stele and ZIP Code)			10. SOURCE OF FUI							
			-		PROGRAM ELEMENT NO.	PHOJECT NO.	TASK NO.	WORK UNIT		
1. TITLE (Include Security Chamilication) Scintillation & In-Sity Irregula			arity Para-	61102F	2310	G9	AM			
12. PERSON	AL AUTHOR	(5)	<u>orar</u> and		J		L			
Eileen	MacKenzi	e, Suna	nda Basu,	Paul Pruneau						
Soloni	SF HEFORT		TISE TIME CO	$\frac{1}{2} \frac{1}{2} \frac{1}$	1990 December 28 36					
16. SUPPLE	MENTARY NO	OTATION								
7. COSATI CODES 18. SUBJECT TEIMIS			18. SUBJECT TEINIS (C	ontinue on reverse if ne	cessary and identi	ly by block number	,			
FIELD	GROUP	SUB. GR.		Amplitude & Pha	Phase Scintillation Auroral Oval					
				In-situ Density	y Polar Cap					
J		L		Electron Densi	ty Deviation	Sui	nspot Cycle	Variation		
IS. ABSTRA Th	CT <i>(Continue</i> e orbitit	<b>on reverse li</b> ng HiLai	(necessary and t_sate11i	identify by block number te launched by i	) the Defense Nu	clear Agend	v in 1983 o	ffered		
a uniqu	e opportu	unity fo	or studyi	ng ionospheric a	scintillation	parameters	in relation	to		
in-situ	irregula	arity pa	arameters	. An earlier re	eport (AFGL-TR	-87-0245) s	summarized of	ur		
finding	s during	the sur	nspot min	imum period of 3	1983-1985 for	data obtain	ned at Troms	D,		
Norway.	In this	s report	t, we ext	end our study to	the sunspot	maximum per	iod of 1988	-1989.		
For com	pleteness	s, we au	igment the	e HiLat scintil	lation data by	phase and	amplitude			
scintii	tor at T	orpnolog	y obtain ring the	ed with the nell	1990 To com	-geostation	ary polar b	eacon		
Jatitud	e region		rovide ec	period 1966 Co	hology using	nolar beach	e two disting	e from		
Thule.	Greenland	d for th	he period	1987 to 1990.	Some new find	ings emerge	e on the UT	and		
seasona	1 variat:	ion of 1	high lati	tude scintillat:	ion as a resul	t of this	study.			
			<b>y-</b>				-			
-										
U DIST HIS UTURIAVALLABILITY OF ABSTRACT			UNCLASSIFIC							
2. NAME C	F RESPONSI		IDUAL		276 TELEPHONE NL	IMBER	ZZE. OFFICE SYMI			
James Whalen				(Include Area Code) (617) 377-4766		GL/LIS				
	1473 82	APR		EDITION OF 1 JAN 23 P		IIN	CLASSIFIED			
21011				Control of Tarrer 191.						

## ACKNOWLEDGEMENTS

3

The authors thank Santimay Basu of the Ionospheric Physics Division, Phillips Laboratory, who made the report possible by making the scintillation data available and for providing helpful advice. We also thank F.J. Rich of the Space Physics Division, Phillips Laboratory, for providing the ion density data from HiLat. We appreciate the efforts of R.C. Livingston of SRI International for his help with the radio beacon data analysis, and Elizabeth Galligan of Boston College for assistance with the polar beacon satellite data analysis.

The work was supported by GL Contract F19628-90-K-0007 with Boston College.

iii

## **1. INTRODUCTION**

t.

The morphological features of scintillation regions have been examined in a wide variety of studies (cf. reviews by Aarons, 1982; Basu et al., 1988 and references therein). However, even such a plethora of earlier publications has failed to clearly delineate the different variables on which scintillation depends, particularly at high latitudes. This is because recent studies have shown that in the polar and auroral environment, large-scale ionization transport in the form of "patches" has an important role to play in determining scintillation behavior (Buchau et al., 1985; Weber et al., 1984; 1986; Basu et al., 1989; 1990). Thus, Tsunoda (1988), in his comprehensive review, points out that attention has been diverted from "local" type of mechanisms, such as particle precipitation, to processes such as global electrodynamics and convective control of patch formation by the interplanetary magnetic field (IMF).

It is the intent of this report to extend the measurements of scintillations and in-situ parameters presented in an earlier study using HiLat (MacKenzie et al., 1987, to be referred to as Report I) by more recent measurements using the same satellite during the current sunspot maximum. We also augment these measurements with those of long-term scintillations from polar beacon satellites at Tromso and Thule which allow us to better study the effects of transport.

## 2. THE HILAT DATA BASE

As detailed in Report I, the HiLat satellite was designed to provide quantitative information on high latitude scintillation through radio beacon experiments, accompanied by insitu data to define the background ionospheric processes (Fremouw et al., 1985). We concentrate on the HiLat satellite observations performed by the Air Force Geophysics Laboratory at Tromso, Norway ( $66^{\circ} \Lambda$ ). Under quiet magnetic conditions, this station is within the nightside auroral oval while the dayside oval is poleward of the station. This report will focus on observations made in 1988-89, a period of increasing solar sunspot number. Figure 1 illustrates the drastic variation of both the monthly mean and smoothed sunspot numbers from December 1983 to March 1990. During the time frame of the observations shown in Report I, namely 1984-85, smoothed sunspot number declined from approximately 60 to 15. Within the time frame of observations to be examined in this report, namely 1988-89, smoothed sunspot number number increased from approximately 60 to 160.

3

The phase and amplitude scintillation magnitudes are examined in the spatial and temporal frames of invariant latitude (A) and magnetic local time (MLT). As in the previous analysis, scintillation measurements have been made on a sliding 30-sec window with values computed every 15 sec at the midpoint of the window for the length of each pass. These have then been merged into seasonal data bases, deleting data acquired below a satellite elevation angle of 20°. The parameters to be discussed in this report are the 137-MHz phase and intensity scintillation observations and the in-situ ion density at the satellite altitude. For definitions of phase and amplitude scintillation indices, the reader is referred to Report I. The phase and intensity scintillations are assumed to pertain to the 350 km sub-ionospheric intersection point of the satellite (830 km). Each 15-sec scintillation value, both phase and intensity, has been corrected for ionospheric zenith angle (i) by means of a (sec i)<sup>1/2</sup> correction. Median values of the corrected rms phase deviation,  $\sigma_{\phi}$ , and intensity scintillation, S<sub>4</sub>, and the ion density have been calculated in 2.5°A - 1-hr MLT bins for each season. Bins with fewer than 10 points have been

omitted. These seasonal data bases have been further sorted into two groups: those observed under magnetically quiet conditions (Kp < 3.5) and those under magnetically disturbed conditions (Kp > 3.5). Results of such statistical analyses are given below.

## 2.1. RMS Phase Deviation

Figures 2a - 2c illustrate, in polar plots of invariant latitude-MLT, the median rms phase deviation,  $\sigma_{\phi}$ , of 137 MHz scintillation for 3 equinoxes: Feb-Apr 1984, Feb-Apr 1988, and Feb-Apr 1989. The 1984 season is included to facilitate comparison with data in the earlier report. The 1984 and 1988 seasons have similar features and scintillation levels: low  $\sigma_{0}$  values are present on the dayside while maximum  $\sigma_{\phi}$  is seen in the region of L-shell enhancement on the nightside. The Feb-Apr 1989 data shows a similar pattern but with much higher  $\sigma_0$  in each bin. A common scale for all panels results in a deceptive broadening of the L-shell enhancement region. Figure 3 presents the rms phase deviation,  $\sigma_{\varphi}$ , in a different format. This figure utilizes a midnight-noon cut through the dial plots of Figures 2b and 2c to obtain in a more quantitative manner the variation of  $\sigma_{\phi}$  with invariant latitude in the 2-hr MLT time period centered on midnight and noon. We have broadened this data set to include all seasons between Feb-Apr 1988 and Feb-Apr 1989. In these cuts, both the gradual increase of rms phase deviation,  $\sigma_{\varphi}$ , with each season and the pronounced enhancement on the nightside over the latitude range  $65^{\circ}$  -  $67.5^{\circ}\Lambda$  can be seen. This enhancement, occurring where the satellite ray path achieves alignment with the magnetic L-shell oriented irregularities, was the most stable morphological feature in Report I. This has also been noted by MacDougall (1990).

## 2.2. Intensity Scintillation, S<sub>4</sub>

Figures 4a - 4c illustrate, in polar plots of invariant latitude-MLT, the median 137 MHz intensity scintillation magnitudes, S4, for 3 equinoxes: Feb-Apr 1984, Feb-Apr 1988, and Feb-Apr 1989. The 1984 and 1988 seasons are similar in features and scintillation level, while the 1989 data show increased intensity scintillation magnitudes particularly in the midnight and noontime sectors. The nighttime enhancement of intensity scintillation index, S4, in the region

of alignment of the ray path with the magnetic L-shell is less pronounced than was seen for the phase scintillation. The L-shell alignment of the km-scale irregularities causing intensity scintillation appears weaker than the tens of kilometers scale that controls the phase scintillations. In Figure 5, the intensity scintillation, S<sub>4</sub>, is presented in the format of a midnight-noon cut through the dial plots of Figures 4b and 4c. Additional seasons have been included. In these cuts, the intensity scintillation, S<sub>4</sub>, is seen to increase with each season.

# 2.3. Ion Density

The in-situ ion density has been observed to vary with both solar activity and season. Report I has shown the invariant latitude-MLT variation of this parameter during 1984-85, a low sunspot period. The median ion density showed a seasonal maximum in Feb-Jul with a daytime diurnal maximum. Figures 6a - 6c illustrate the invariant latitude-MLT ion density variation during Feb-Apr 1984, Feb-Apr 1988, and Feb-Apr 1989. The 1984 and 1988 data are comparable while the Feb-Apr 1989 data show an enormous increase; the minimum ion density in 1989 is an order of magnitude greater than those of 1984 or 1988. Figure 6c has been redone on a different scale in Figure 6d to exhibit some of the variation lost when the scale, appropriate for 1984 and 1988, is used for 1989. Figure 7 illustrates the midnight-noon cuts through the polar plots of Figures 6b and 6d. Additional seasons between Feb-Apr 1988 and Feb-Apr 1989 are included. Although sunspot number was increasing throughout 1988, the large increase in ion density does not show in the statistics until Feb-Apr 1989, probably because the seasonal variations noted in Report I get masked by the ramp-like increase in smoothed sunspot number.

# 3. SCINTILLATION MEASUREMENTS FROM QUASI-GEOSTATIONARY SATELLITES

Additional quasi-geostationary scintillation data bases are available to augment the HiLat orbiting satellite statistics. The major difference between HiLat scintillation data and those obtained from quasi-geostationary satellites is that in the latter, irregularity anisotropy does not play a major role so that long-term trends are more readily discernible. Measurements using several polar beacon satellites at 250 MHz are available for similar time periods. Auroral oval data are obtained in the Scandinavian sector from Tromso, while polar cap data are available from Thule. These polar beacon satellites do not provide uniform local time coverage. They also tend to explore different latitude ranges at different times of the year. Results from each station are given below.

#### 3.1. Measurements at Tromso

At Tromso, the polar beacons provide 24-hour coverage in a combination of data from within the auroral oval with that from slightly below it. Basu et al. (1988) have shown very low intensity scintillation statistics during the sunspot minimum period of 1984-86, with fades of 5 dB approximately 5% of the time. From the current sunspot maximum 250 MHz data, both intensity and phase scintillation data are utilized. This has been subdivided to focus on only auroral location data for nighttime hours of 21-03 MLT. Figure 8 illustrates the median S<sub>4</sub> and rms phase deviation,  $\sigma_{\phi}$ , statistics seasonally from November 1985 to January 1990. The S<sub>4</sub> median shows low values through the sunspot minimum years (1985-1987) increasing late in 1988. The  $\sigma_{\phi}$  median exhibits less pronounced variation. However, both S<sub>4</sub> and  $\sigma_{\phi}$  show a pronounced seasonal variation with minimum scintillation occurrence seen in the Summer months. A similar minimum was observed in the Goose Bay auroral scintillation data (Basu et al. 1988). We shall discuss this point further in Section 4.

### 3.2. Measurements at Thule

The polar cap data, available from Thule, likewise explores different latitude ranges between 85° - 88° invariant at different times of the year. However, this range is within the

central polar cap so that no subdivision of the data has been performed. Figures 9a and 9b illustrate the percent occurrence of varying intensity scintillation fade levels over a whole solar cycle, 1979-1990, while Figure 10 illustrates the median S<sub>4</sub> and rms phase deviation,  $\sigma_{\varphi}$ , statistics seasonally over a shorter time frame (1987-1989). As was seen with the Tromso data, a larger variation with increased sunspot number is seen in S<sub>4</sub> than in  $\sigma_{\varphi}$ . Both Figures 9 and 10 show a minimum in scintillation occurrence and magnitude during the local Summer months.

ż

To study this behavior in greater detail, we present the scintillation fade occurrence  $> 4 \, dB$ during sunspot minimum (July 1985 - June 1986) and sunspot maximum (July 1989 - June 1990) periods in Figures 11 and 12. It should be noted that a fade of 4 dB is equivalent to peak-to-peak fluctuations of 6 dB (Whitney, 1974). Figure 12 clearly shows minimal scintillation occurrence during Summer, and a diffuse maximum between October to February. Largest percentage occurrence is observed between 15 - 06 UT. To study the occurrence of intense scintillation events, we show in Figure 13, contours of a fade level >10 dB (peak-topeak fluctuations of 15 dB) during the high sunspot year of 1989-90. The same seasonal pattern is emphasized with the highest activity being confined to 18 - 24 UT. There is a general tendency for somewhat greater scintillation occurrence in the Fall as compared with the Spring, as Figures 9a and 9b also depict. It is noteworthy that the time of maximum occurrence is not related to total darkness, as the sunset and sunrise times at 110-km altitude indicate for the equinoxes. At the solstices, of course, there is either no sunrise in the winter months or no sunset in the summer months. Thus, the occurrence of scintillation at Thule does not seem to be controlled by local conditions. We shall discuss this point further in the next Section.

# 4. **DISCUSSION**

The major morphological finding of the HiLat scintillation study, namely, the enhancement of scintillation where the satellite ray path was aligned with the magnetic L-shell oriented irregularities, has been discussed at length in Report I. Longer term variations, such as seasonal dependence of scintillations, are harder to identify in HiLat data (Rino and Matthews, 1980) because of the great sensitivity of the magnitude of scintillations on the irregularity anisotropy. Sunspot cycle effects are clearly evident both on scintillations with the sunspot number can be thought of as being caused by a modulation in the background density as both the scintillation and ion density magnitude increase by the same factor. Of course, here we make the tacit assumption that the in-situ data measured at 830 km altitude mimics the density variation at the maximum of the F-layer because it is that density which controls the scintillations.

The additional geo-stationary scintillation data from Tromso and Thule presented in Section 3 provide much new information when considered in tandem with earlier published work from Thule (Aarons et al., 1981), Goose Bay (Basu et al., 1988), and Narssarsuaq (Basu, 1975; Basu and Aarons, 1980). If we start with the Thule data which refers to the central polar cap, then we find that the UT seasonal and solar cycle dependence of scintillations are generally consistent with transport of plasma from sub-cusp latitudes taken in conjunction with the shorting effect of a sunlit E-region in the summer on F-region irregularities. Some unexplained time shifts do occur, however. For instance, the highest scintillation occurrence is observed at Thule between 18-24 UT (cf. Figure 13), whereas Figure 4 of Buchau et al. (1985) shows the UT variation of maximum F-region density at a sub-cusp location (73° invariant at noon magnetic local time) to peak about three hours earlier. It is generally believed that sub-cusp plasma is convected across the polar cap and structures in the anti-sunward convection field giving rise to scintillations (Weber et al., 1984; Basu et al., 1990). On this basis, one expects maximum scintillation occurrence to follow maximum density behavior with a phase lag of 30 mins to an hour, depending on the velocity range of these patches varying between 300 and 700 ms<sup>-1</sup>. However, a phase lag of three hours is difficult to understand and factors such as transit delays and other irregularity source mechanisms have to be investigated.

The seasonal behavior of the Tromso polar beacon scintillations is very similar to that observed at Thule during sunspot maximum, with the predominant variation being an annual one with minimum in the Summer. As mentioned earlier, Goose Bay polar beacon data had yielded similar results. Thus all three sets taken together seem to indicate that, on the average, transport and E-region shorting both are important considerations. Earlier annual scintillation morphology studies based on Narssarsuaq ATS-3 data, which explored the equatorward edge of the auroral oval, showed dramatically different occurrence statistics with a summer maximum and a winter minimum (Basu, 1975; Basu and Aarons, 1980). These earlier measurements were also at a frequency of 137 MHz so could respond to much weaker irregularities. However, this great difference in the occurrence characteristics between the poleward and equatorward regions of the auroral oval merit much further study. It is hoped that the routine polar beacon measurements from Sondrestrom started in 1990 will be helpful in providing further clues regarding this disagreement.

### REFERENCES

Aarons, J. (1982) Global morphology of ionospheric scintillation, Proc. IEEE, 70: 360.

- Aarons, J., J.P. Mullen, H.E. Whitney, A.L. Johnson, and E.J. Weber (1981) UHF scintillation activity over polar latitudes, *Geophys. Res. Lett.*, 8: 277.
- Basu, S., E. MacKenzie and Su. Basu (1988) Ionospheric constraints on VHF/UHF communications links during solar maximum and minimum periods, *Radio Sci.*, 23: 363.
- Basu, S., Su. Basu, C.E. Valladares, E.J. Weber, J. Buchau, G.J. Bishop, and B.W. Reinisch (1989) Coordinated observations of high latitude ionospheric turbulence, SPI Conference Proceedings and Reprint Series, 8: 137.
- Basu, Su. (1975) Universal time seasonal variations of auroral zone magnetic activity and VHF scintillations, J. Geophys. Res., 80: 4725.
- Basu, Su. and J. Aarons (1980) The morphology of high-latitude VHF scintillation near 70°W, Radio Sci., 15: 59.
- Basu, Su., S. Basu, E.J. Weber, and G.J. Bishop (1990) Plasma structuring in the polar cap, J. Geomag. Geoelectr., 42: 763.
- Buchau, J., E.J. Weber, D.N. Anderson, H.C. Carlson, Jr., J.G. Moore, B.W. Reinisch, and R.C. Livingston (1985) Ionospheric structures in the polar cap: their origin and relation to 250-MHz scintillation, *Radio Sci.*, 20: 325.
- Fremouw, E.J., H.C. Carlson, T.A. Potemra, P.F. Bythrow, C.L. Rino, J.F. Vickrey, R.L. Livingston, R.E. Huffman, C.-I. Meng, D.A. Hardy, F.J. Rich, R.A. Heelis, W.B. Hanson, and L.A. Wittwer (1985) HiLat satellite mission, *Radio Sci.*, 20: 416.
- MacDougall, J. (1990) Distribution of irregularities in the northern polar region determined from HiLat observations, *Radio Sci.*, 25: 115.
- MacKenzie, E., S. Basu, and Su. Basu (1987) Ionospheric scintillations/TEC and in-situ density measurements at an auroral location in the European sector, AFGL-TR-87-0245, 14 August, ADA205543.
- Rino, C.L. and S.J. Matthews (1980) On the morphology of auroral zone radio wave scintillation, J. Geophys. Res., 85: 4139.
- Tsunoda, R.T. (1988) High-latitude F-region isregularities: a review and synthesis, Rev. Geophys., 26: 719.
- Weber, E.J., J. Buchau, J.G. Moore, J.R. Sharber, R.C. Livingston, J.D. Winningham, and B.W. Reinisch (1984) F-layer ionization patches in the polar cap, J. Geophys. Res., 89: 1683.
- Weber, E.J., J.A. Klobuchar, J. Buchau, H.C. Carlson, Jr., R.C. Livingston, O. de la Beaujardiere, M. McCready, J.G. Moore, and G.J. Bishop (1986), Polar cap F layer patches: structure and dynamics, J. Geophys. Res., 91: 12121.
- Whitney, H.E. (1974) Notes on the relationship of scintillation index to probability distributions and their uses for system design, AFCRL-TR-74-0004, AD778092.

## FIGURE CAPTIONS

- Figure 1. Variation of monthly mean and smoothed sunspot numbers between December 1983 and March 1990.
- Figure 2a. Variation of Tromso HiLat median rms phase deviation,  $\sigma_{\phi}$  (137 MHz), with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1984.

i

- Figure 2b. Variation of Tromso HiLat median rms phase deviation,  $\sigma_{\varphi}$  (137 MHz), with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1988.
- Figure 2c. Variation of Tromso HiLat median rms phase deviation,  $\sigma_{\varphi}$  (137 MHz), with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1989.
- Figure 3. Seasonal variations of 50th and 90th percentiles of Tromso HiLat rms phase deviation,  $\sigma_{\varphi}$  (137 MHz), with Invariant Latitude in 2-hr MLT bins along the midnight-noon meridian under quiet magnetic conditions in 1988-89.
- Figure 4a. Variation of Tromso HiLat median S<sub>4</sub> (137 MHz), with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1984.
- Figure 4b. Variation of Tromso HiLat median S<sub>4</sub> (137 MHz), with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1988.
- Figure 4c. Variation of Tromso HiLat median S<sub>4</sub> (137 MHz), with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1989.
- Figure 5. Seasonal variation of 50th and 90th percentiles of Tromso HiLat intensity scintillation, S<sub>4</sub> (137 MHz) with invariant latitude in 2-hr MLT bins along the midnight-noon meridian under quiet magnetic conditions in 1988-89.
- Figure 6a. Variation of Tromso HiLat median ion density with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1984.
- Figure 6b. Variation of Tromso HiLat median ion density with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1988.
- Figure 6c. Variation of Tromso HiLat median ion density with Invariant Latitude and magnetic local time under quiet magnetic conditions during Feb-Apr 1989.
- Figure 6d. Replotted version of Fig. 6c with a revised density scale in order to highlight increased ion density.
- Figure 7. Seasonal variation of 50th and 90th percentiles of Tromso HiLat median ion density with invariant latitude in 2-hr MLT bins along the midnight-noon meridian under quiet magnetic conditions in 1988-89.

- Figure 8. Seasonal variations of 50th and 90th percentiles of Tromso intensity scintillation, S4 (250 MHz) and rms phase deviation,  $\sigma_{\phi}$  (250 MHz) from quasi-geostationary satellites under quiet magnetic conditions within the auroral oval (~ 69°A) around magnetic local midnight (21-03 MLT) during 1985-1989.
- Figure 9a. Occurrence of 5-, 10-, and 15- dB fades at 250 MHz observed using quasigeostationary satellites at Thule during the period 1979-1986. All local times and all magnetic activity data are included.
- Figure 9b. Occurrence of 5-, 10-, and 15- dB fades at 250 MHz observed using quasigeostationary satellites at Thule during the period 1987-1990. All local times and all magnetic activity data are included.
- Figure 10. Seasonal variations of 50th and 90th percentiles of Thule intensity scintillation, S4 (250 MHz) and rms phase deviation,  $\sigma_{\phi}$  (250 MHz) from quasi-geostationary satellites under all magnetic conditions around midnight (21-03 UT) during 1987-1989.
- Figure 11. Contours of percent occurrence of fades > 4 dB at 250 MHz observed using quasigeostationary satellites at Thule during the solar minimum period of July 1985 through June 1986 for all magnetic activity levels.
- Figure 12. Contours of percent occurrence of fades > 4 dB at 250 MHz observed using quasigeostationary satellites at Thule during the solar maximum period of July 1989 through June 1990 for all magnetic activity levels.
- Figure 13. Contours of percent occurrence of fades > 10 dB at 250 MHz observed using quasi-geostationary satellites at Thule during the solar maximum period of July 1989 through June 1990 for all magnetic activity levels.



J









Figure 2c.



Figure 3.



Figure 4a.



Figure 4b.





S<sub>4</sub>

. . . . .



Figure 5.



!





Figure 6c.



Figure 6d.

TROMSO

Kp < 3.5







Figure 8.



Figure 9a.







;

OCCURRENCE OF IS MINUTE FADE DEPTH INDICES (dB RELATIVE TO MEDIAN SIGNAL)

Figure 9b.



Figure IO.





•

ł



31

٠.





•

<u>م</u>

•

\_