

REPORT DOCUMENTATION PAGE *Dist: A*

Form Approved  
OMB No 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 30 Sept 1994 3. REPORT TYPE AND DATES COVERED Annual Technical 9/1/93-8/31/94

4. TITLE AND SUBTITLE AUGMENTATION AWARD FOR MONOCLONAL ANTIBODY DETECTION OF CHLORINATED BENZENES ON CONTAMINATED SEDIMENTS 5. FUNDING NUMBERS AFOSR F49620-93-1-0523

6. AUTHOR(S) Deborah J. Mossman *61103 F 3484/45*


7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Missouri-Columbia Dept. of Civil Engineering 5100 Rockhill Road Kansas City, MO 64110-2499 8. PERFORMING ORGANIZATION REPORT NUMBER AFOSR-TR-94 0667

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Office of Scientific Research/NL Building 410 Bolling AFB, DC 20332-6448 10. SPONSORING/MONITORING AGENCY REPORT NUMBER *Dr Kozumbo* *388* 94-33891

11. SUPPLEMENTARY NOTES *388* 

12a. DISTRIBUTION / AVAILABILITY STATEMENT *A*

13. ABSTRACT (Maximum 200 words) The nonextractive immunoassay techniques developed with enzyme immunoassay procedures are being applied to fluorescent immunoassay visualization. Sorbed contaminants can be viewed using this modified fluorescent immunoassay techniques and and epifluorescent microscope to observe the microdistribution of sorbed contaminants.



14. SUBJECT TERMS fluorescent immunoassay, sorption 15. NUMBER OF PAGES 2 16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED 18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED 19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED 20. LIMITATION OF ABSTRACT

AD-A285 896

(3)

**AUGMENTATION AWARD FOR MONOCLONAL ANTIBODY DETECTION OF  
CHLORINATED BENZENES ON CONTAMINATED SEDIMENTS**

Deborah J. Mossman  
Dept. of Civil Engineering  
University of Missouri-Columbia  
5100 Rockhill Road  
Kansas City, MO 64110-2499

30 September 1994

AASERT Technical Report

Prepared for  
AIR FORCE OFFICE OF SCIENTIFIC RESEARCH  
Bolling AFB, DC 20332-6448

Engineering and Services Laboratory  
HQ AFESC/RDVW  
Tyndall AFB, FL 32403

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

DTIC QUALITY INSPECTED 6

94 11 1 06 1

## **Progress of AASERT Student**

The AASERT student has been using a Zeiss Axiophot epifluorescent microscope to examine the feasibility of fluorescent immunoassay (FIA) techniques for visualization of contaminants sorbed to soil particles. The first task completed was to select a fluorochrome which would be compatible with the soil FIA. In the excitation/emission wavelength ranges used in common fluorochromes, the mineral particles exhibit significant yellow-green autofluorescence. The fluorochromes commercially available conjugated to secondary antibodies are fluorescein, rhodamine, phycoerythrin, and Texas Red™. At present, the excitation/emission filter set which is needed for Texas Red is not available in the laboratory. Fluorescein emits in the yellow-green range and rhodamine was too weak an emitter. Work completed during the first grant year used R-phycoerythrin (RPE), which emits in the orange range, for fluorescent immunoassay examination of soil contamination.

A variety of soil sample types were examined with the FIA method: brick crushed to be retained on sieve #45, brick #80, builders sand #45, kaolinite, crushed obsidian #170, St. Peter's sandstone #45, St. Peter's sandstone #80, S-2 soil (obtained from Soil Testing Laboratory at Kansas State University) #400, and S-2 soil #200. The contaminant sorbed on the soils was 2,4-dinitrobenzene sulfonate (DNB), and the primary antibody was anti-DNB developed in rabbit (rb  $\bar{\alpha}$ DNB) (Sigma). The samples were contaminated by soaking the soil samples in a pH 10 DNB solution. The labelled secondary antibody was commercially-obtained antibodies to rabbit developed in goat labelled with R-phycoerythrin (gt  $\bar{\alpha}$ rb-RPE). All results are photodocumented with daylight ASA400 Ektachrome slide film.

Clean and contaminated samples are first blocked using a protein solution to prevent nonspecific attachment of antibodies during the procedure. The samples are then incubated and agitated in the primary antibody solution (rb  $\bar{\alpha}$ DNB). The samples are then washed to remove unbound antibodies, and the fluorochrome-labelled secondary antibody (gt  $\bar{\alpha}$ rb-RPE) is added, and the sample and solution are incubated and agitated. The sample is washed again, and can then be viewed under the epifluorescence microscope.

There is a distinction between contaminated samples and uncontaminated samples when treated with FIA. The distinction was clearly observed in the brick samples (which exhibit low autofluorescence) and the St. Peter's sandstone. There was less distinct difference in the obsidian and S-2 soil, and no discernible difference between contaminated and uncontaminated samples using the kaolinite and builders sand. Experiments were conducted varying the blocking concentration (used to reduce nonspecific binding of the antibodies which will yield false positives), and two different blocking solutions were tried. Chicken egg albumin block reduced autofluorescence, but did not appear to prevent non-specific binding. Skim milk block autofluoresced somewhat, but fulfilled the primary goal of preventing nonspecific binding. Therefore skim milk has been used as the protein component of the block solution in the experiments.

An oral presentation titled "Fluorescent Immunoassay Visualization of Sorbed Pollutants" was given by W.K. Moore (AASERT student) at the Ninth Annual Hazardous Waste Research Conference, 8-10 June 1994, Montana State University, Bozeman, Montana.

## **Academic Performance of AASERT Student**

The graduate student supported under the AASERT grant has maintained a 4.0/4.0 grade point average during the first grant year.