



Final Report, N00014-90-J-1834
"Southern Norwegian Sea Surveys"
Principal Investigator:
Alexander Shor
Hawaii Institute of Geophysics
University of Hawaii at Manoa

The research grant N00014-90-J-1834 supported a 75-day geophysical field program on Reykjanes Ridge in the North Atlantic Ocean and Aegir Ridge in the Norway Basin of the Norwegian Sea. The field program took place in summer 1990 on R/V Maurice Ewing of Lamont-Doherty Geological Observatory; the program was carried out jointly by researchers at the Naval Research Laboratory Code 5110, Lamont-Doherty Geological Observatory and University of Hawaii. University of Hawaii responsibility included providing and operating the SeaMARC II seafloor mapping system during the course of the field program, processing the SeaMARC II side-scan sonar and bathymetric data to produce seafloor charts and sonar mosaics, and participating in the geophysical interpretation of the data.

The research cruise was successfully completed in September 1990 in Bergen, Norway. Most SeaMARC II data processing was completed during the cruise, and copies of mosaics and charts were delivered to NRL and Hawaii for analysis and interpretation. Some necessary post-processing was completed over the six months following the field effort, and those results also were distributed among the collaborators for analysis. A copy of the ONR "Geology and Geophysics Program Summary for FY90" report written after field program completion is attached, providing information on objectives and project status at the end of 1990. Publications resulting from this field program include:

- Shor, A.N., C.E. Nishimura, M. Czarnecki and P.R. Vogt, 1990. Lava extrusion from the 1989 Reykjanes Ridge seismic swarm? Probably yes (SeaMARC II). Eos, v. 71, p. 1602 (abstract).
- Vogt, P.R., D.A. Chayes, H.S. Fleming and A.N. Shor, 1990. SeaMARC II/Hydrosweep/marine geophysical investigation of the extinct Aegir Ridge, Norway Basin. Eos, v. 71, p. 1408 (abstract).
- Appelgate, B., and A. Shor, 1991. The Reykjanes Ridge: Along-axis variation in tectonic and volcanic morphology from SeaMARC II and Hydrosweep. Eos, 1991 Fall Meeting Supplement, p. 467 (abstract).
- Jung, W-Y., P.R. Vogt, D. Chayes and H.S. Fleming, 1991. Detailed gravity/magnetic mapping of the extinct Aegir Ridge. Eos, 1991 Fall Meeting Supplement, p. 486 (abstract).
- Vogt, P.R., K. Crane, S. Pfirman, E. Sundvor, D. Chayes, N. Cherkis, H. Fleming, C. Nishimura and A. Shor, 1991. SeaMARC II sidescan sonar imagery and swath bathymetry in the Nordic Basin: Sedimentary processes elucidated by correlation with 3.5 kHz and other data. Eos, 1991 Fall Meeting Supplement, p. 486 (abstract).
- Appelgate, B., and A.N. Shor, 1992. Reykjanes Ridges and the northernmost Mid-Atlantic Ridge: Segmentation and tectonic signature of a slow-spreading ridge near the Iceland hot spot, Eos, 1992 Fall Meeting Supplement, p. 538 (abstract).
- Appelgate, B., and A.N. Shor (submitted 6/93), The northern Mid-Atlantic and Reykjanes Ridges: Spreading center morphology between 55°50'N and 63°00'N. Journal of Geophysical Research.

This document has been approved for public release and sale; its distribution is unlimited.

**GEOLOGY AND GEOPHYSICS
PROGRAM SUMMARY FOR
FY90**

MARCH 1991

ONR CODE 1125GG



Approved for Public Release: Distribution Unlimited

**Ocean Sciences Directorate
Office of Naval Research
800 N. Quincy St.
Arlington, Virginia 22217-5000**

Project Objectives (continued)

1. Define the morphologic expression of the extinct Aegir Ridge system. In particular, we plan to examine relief, orientation and spacing of "abyssal hill" fabric as a function of spreading rate, and the relative contribution of tectonics (faulting) and volcanism (seamounts) to seafloor relief. Comparison of Aegir Ridge survey data with other slow spreading rate ridge surveys, including the Knipovich Ridge SeaMARC II surveys carried out in 1989 and 1990, will provide a data set for identifying characteristic morphologies associated with a dying ridge system. Simultaneous gravity data acquisition with detailed bathymetry and sediment thickness will allow modelling of the flexural properties of the ridge and estimates of isostatic readjustment of the ridge to the new thermal regime caused by cessation of spreading. The Hydrosweep bathymetry and SeaMARC II side-scan sonar data will be the principal information used for interpretation of the central basin, augmented by closely spaced seismic reflection lines for exploring buried crust on the Aegir Ridge flanks.
2. Identify and define the tectonic and morphologic character of the northern and southern ends of the Aegir Ridge system. The tectonic signature at both ends of Aegir Ridge is complex (Old Jan Mayen Fracture Zone to the north; Iceland-Faeroes Ridge to the south), and the character of the intersection between spreading segments with both offsets is poorly defined.
3. Define the sediment sources and depositional regime of the Norway Basin. The combined use of full coverage bathymetry and side-scan sonar with closely spaced seismic reflection profiling will provide a powerful data set for interpreting spatial and temporal variations in depositional processes.

Present Status and Progress During the Current Year

The field program was successfully completed in September 1990. On-board processing of Hydrosweep and SeaMARC II data and the preparation of final charts and sonar mosaics at sea allowed preliminary interpretation during the cruise. However, detailed study was just beginning as FY90 ended. Preliminary results will be presented at the Fall 1990 AGU Meeting in San Francisco.

Publications for FY90

Vogt, P.R., Chayes, D.A., Fleming, H.S., Shor, A.N., 1990. SeaMARC II/Hydrosweep/Marine Geophysical Investigation of the Extinct Aegir Ridge, Norway Basin. *Eos*, 71, 1408 (abstract).