




**EDGEWOOD**

RESEARCH DEVELOPMENT & ENGINEERING CENTER

U.S. ARMY CHEMICAL AND BIOLOGICAL DEFENSE COMMAND

ERDEC-TR-212

**COMPUTATION OF ELECTROMAGNETIC SCATTERING PARAMETERS  
FOR LOGNORMAL DISTRIBUTIONS OF MAGNETIC SPHERES:  
THEORY AND ALGORITHMS**



Merrill E. Milham

RESEARCH AND TECHNOLOGY DIRECTORATE

October 1994

Approved for public release; distribution is unlimited.

19950112 084



Aberdeen Proving Ground, MD 21010-5423

#### **Disclaimer**

**The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.**

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)			2. REPORT DATE 1994 October	3. REPORT TYPE AND DATES COVERED Final, 91 Jan - 94 Jan	4. TITLE AND SUBTITLE <b>Computation of Electromagnetic Scattering Parameters for Lognormal Distributions of Magnetic Spheres: Theory and Algorithms</b>	5. FUNDING NUMBERS <b>PR-1O162622A552</b>
6. AUTHOR(S) <b>Milham, Merrill E.</b>						
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>DIR, ERDEC,* ATTN: SCBRD-RTE, APG, MD 21010-5423</b>			8. PERFORMING ORGANIZATION REPORT NUMBER <b>ERDEC-TR-212</b>			
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING / MONITORING AGENCY REPORT NUMBER			
11. SUPPLEMENTARY NOTES <b>*When this study was conducted, ERDEC was known as the U.S. Army Chemical Research, Development and Engineering Center, and the author was assigned to the Research Directorate.</b>						
12a. DISTRIBUTION / AVAILABILITY STATEMENT <b>Approved for public release; distribution is unlimited.</b>			12b. DISTRIBUTION CODE			
13. ABSTRACT (Maximum 200 words)  <b>In this report, relevant parts of the scattering theory for magnetic spheres are presented. Mass extinction coefficients, and the lognormal size distribution are defined. The theory and algorithms for integrating scattering parameters over size distributions are developed. The integrations are carried out in terms of dimensionless scattering, and size distribution parameters, which are simply related to the usual mass scattering coefficients. Fortran codes, which implement the algorithmic design, are presented, and examples of code use are given. Code listings are included.</b>						
14. SUBJECT TERMS <b>Magnetic sphere Complex permeability Complex permittivity</b>			15. NUMBER OF PAGES <b>32</b>			
16. PRICE CODE						
17. SECURITY CLASSIFICATION OF REPORT <b>UNCLASSIFIED</b>	18. SECURITY CLASSIFICATION OF THIS PAGE <b>UNCLASSIFIED</b>	19. SECURITY CLASSIFICATION OF ABSTRACT <b>UNCLASSIFIED</b>	20. LIMITATION OF ABSTRACT <b>UL</b>			

**14. SUBJECT TERMS (Continued)**

**Lognormal distributions**

**Dimensionless extinction coefficients**

## PREFACE

The work described in this report was authorized under Project No. 10162622A552, Smoke Technology. This work was started in January 1991 and completed in January 1994.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

This report has been approved for release to the public. Registered users should request additional copies from the Defense Technical Information Center; unregistered users should direct such requests to the National Technical Information Service.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution	
Availability Codes	
Dist	Aval and/or Special
R'	

**Blank**

## **CONTENTS**

1.	INTRODUCTION .....	7
2.	LIGHT SCATTERING AND CLOUD MACROPHYSICS .....	7
3.	NUMERICAL METHODS - ALGORITHM DESIGN .....	8
4.	RESULTS AND DISCUSSION .....	9
	LITERATURE CITED .....	13
	APPENDIXES	
	A. PROGRAM LISTINGS .....	15
	B. SAMPLE OUTPUT .....	31

**Blank**

**COMPUTATION OF ELECTROMAGNETIC SCATTERING PARAMETERS  
FOR LOGNORMAL DISTRIBUTIONS OF MAGNETIC SPHERES:  
THEORY AND ALGORITHMS**

**1. INTRODUCTION**

A previous report<sup>1</sup> described the theory, algorithmic design, and testing of a subroutine for computing the electromagnetic scattering from a magnetic sphere. In this report the methodology for using this subroutine to compute the mass extinction, scattering, absorption, and backscatter coefficients for lognormally distributed particulate ensembles is developed.

In what follows the relevant parts of Mie scattering theory are presented. Mass extinction coefficients, and the lognormal size distribution are defined, and the theory and algorithms for integrating Mie scattering parameters over size distributions are developed. The integrations are carried out in terms of dimensionless scattering, and size distribution parameters, which are simply related to the usual mass scattering coefficients.

**2. LIGHT SCATTERING AND CLOUD MACROPHYSICS**

For spheres, the efficiencies, the ratio of the optical cross section to the geometric cross section, for extinction ( $Q_e$ ), scattering ( $Q_s$ ), absorption ( $Q_a$ ), and backscatter ( $Q_b$ ) are:<sup>2</sup>

$$Q_e = \frac{2}{X^2} \sum_{n=1}^{\infty} (2n+1) \operatorname{Re}(a_n + b_n) \quad (1)$$

$$Q_s = \frac{2}{X^2} \sum_{n=1}^{\infty} (2n+1) [ |a_n|^2 + |b_n|^2 ] \quad (2)$$

$$Q_a = Q_e - Q_s \quad (3)$$

$$Q_b = \frac{1}{X^2} \left| \sum_{n=1}^{\infty} (-1)^n (2n+1) (a_n - b_n) \right| \quad (4)$$

where

$$a_n = \frac{\sqrt{\tilde{\mu}} J_{n+\frac{1}{2}}(X) J'_{n+\frac{1}{2}}(mX) - \sqrt{\tilde{\epsilon}} J_{n+\frac{1}{2}}(mX) J'_{n+\frac{1}{2}}(X)}{\sqrt{\tilde{\mu}} H_{n+\frac{1}{2}}^{(2)}(X) J'_{n+\frac{1}{2}}(mX) - \sqrt{\tilde{\epsilon}} J_{n+\frac{1}{2}}(mX) H_{n+\frac{1}{2}}^{(2)}(X)} \quad (5)$$

$$b_n = \frac{\sqrt{\epsilon} J_{n+1/2}(X) J'_{n+1/2}(mX) - \sqrt{\mu} J_{n+1/2}(mX) J'_{n+1/2}(X)}{\sqrt{\epsilon} H_{n+1/2}^{(2)}(X) J'_{n+1/2}(mX) - \sqrt{\mu} J_{n+1/2}(mX) H_{n+1/2}^{(2)}(X)} \quad (6)$$

$J_{n+1/2}(z)$ ,  $H_{n+1/2}(z)$  are the Bessel functions and Hankel functions (second kind) of half-integer order.<sup>3</sup>  $X = \pi D/\lambda$  is the size parameter where,  $D$ , is the diameter in  $\mu\text{m}$ ;  $\lambda$ , the wavelength in  $\mu\text{m}$ ; and  $m = \sqrt{\mu\epsilon}$ , the refractive index for which  $\mu$ , the permeability, and  $\epsilon$ , the permittivity, are complex quantities. Primes denote derivatives with respect to the argument.

For ensembles of spherical particles, the mass extinction, scattering, absorption, or backscatter coefficients (in square meters per gram) at wavelength  $\lambda$  are defined as

$$\alpha = \sum_i A_i / \sum_i m_i \quad (7)$$

where  $A_i$  is the optical cross section for extinction, scattering, absorption, or backscatter of the  $i$ th particle (in square meters),  $m_i$  is the mass of the  $i$ th particle (in grams), and the sum extends over the ensemble of particles contained in the optical path. For a continuous distribution of particles, the expression for  $\alpha$  becomes

$$\alpha = \int \alpha(D) dm \quad (8)$$

$\alpha(D)$  is the coefficient of extinction, scattering, absorption, or backscatter for a particle of size  $D$ ,  $= 3Q/2\rho D$  where  $Q$  is given by the expressions for  $Q_e$ ,  $Q_s$ ,  $Q_a$ , and  $Q_b$  (Equations 1-4); by convention an additional factor of  $4\pi$  is included in the denominator of this expression when  $Q = Q_b$ .<sup>4</sup>  $dm$  is the mass-distribution function which for the purposes of this work is chosen to be the lognormal distribution function:

$$dm = (2\pi \ln^2 \sigma_g)^{-1/2} \exp \left\{ -\frac{1}{2} [\ln(D/D_m) / \ln \sigma_g]^2 \right\} d(\ln D) \quad (9)$$

### 3. NUMERICAL METHODS - ALGORITHM DESIGN

Evaluation of the indicated quadratures over lognormal particle-size distributions can be a difficult numerical problem. The determination of the  $Q$ 's is a computationally intense procedure that must be carried out over a fine integration mesh to ensure accurate results; this is especially true for  $Q_e$  with no absorption ( $\text{Im}(\tilde{m}) = 0$ ) or  $Q_s$ , which have substantial resonant structure.

Because the  $Q$ 's are evaluated in terms of the dimensionless size parameter,  $X$ , it is convenient to formulate the quadrature problem in terms of dimensionless quantities. The  $\alpha$ 's can be put in a dimensionless form as follows:

$$\alpha_d = \alpha \cdot \lambda \cdot \rho \quad (10)$$

Where  $\rho$  is the density in g cm<sup>-3</sup>. Also, converting  $D$ ,  $D_m$  in the particle size distribution to  $X$ ,  $X_m$  leaves the functional form of the distribution unchanged.

After converting these quantities to dimensionless form the integrations are carried out by establishing minimum ( $X_o$ ) and maximum ( $X$ ) size parameters that are the end points for  $N$  evenly spaced abscissae on a logarithmic grid and then applying an appropriate numerical integration scheme such as Simpson's rule. The  $X$  grid is given by

$$X_0, RX_0, \dots, R^{N-1} X_0 \quad (11)$$

where  $X_o = X_m/\sigma_g^n$ ,  $X_i = X_m \sigma_g^n = R^{i-1} X_o$ , and  $R = \Delta \ln X = \sigma_g^{2n/(N-1)}$ .  $n$ , the number of geometric standard deviations, is chosen to be three or greater so that regions outside the grid contribute a numerically negligible amount to the integral. The dimensionless coefficients are then given by the sum

$$\alpha_d \approx (9\pi/8 \ln^2 \sigma_g)^{1/2} \sum_{j=1}^N X_j Q \exp\left\{-\frac{1}{2} \left[ (\ln X_j - \ln X_m) / \ln \sigma_g \right]^2\right\} \Delta(\ln X) \quad (12)$$


When  $Q = Q_b$ , then the expression for  $\alpha_d$  must be multiplied by an additional factor of  $1/4\pi$ . The Figure shows the effect of varying the number of terms in the sum in Equation 12 for the dimensionless extinction coefficient for  $m = (3.0, 0.)$ ;  $\mu = (1.0, 0.)$ .

#### 4. RESULTS AND DISCUSSION

The listings of a main (driver) program (disint), which computes either dimensioned or dimensionless extinction, scattering, absorption, and backscatter coefficients is displayed in Appendix A along with the listings for the auxiliary routines: dxtcs, magsph, zbjy, and simp. The Bessel function subroutine package by Amos<sup>5</sup> is required to implement these routines.

Appendix B lists the input to disint and then displays the resulting output for three sample calculations. The first two sample problems compute the dimensionless extinction coefficients for a mass median size parameter of 10, a refractive index of (1.5, -0.1). The first example uses a very narrow distribution ( $\sigma_g = 1.01$ ); the results of this calculation can be inverted by the relationship,

$$Q \approx \frac{2X_m}{3\pi} \alpha \quad (13)$$



and the results compared with the calculation for monodisperse particle calculation of Wiscombe<sup>6</sup> (Table). The next example gives the results for a particle with same refractive index and a geometric standard deviation of 1.4. The final example is a sample problem that exercises the code for the case of magnetic spheres with  $\mu = \epsilon = (2.24, -3)$ , which are lognormally distributed with an MMD of 10  $\mu\text{m}$  and a  $\sigma_g$  of 1.5. It is known that the backscatter vanishes for spheres with these material properties;<sup>1,7</sup> this sample problem illustrates this point.

Table. Efficiency Factor Comparisons

Distribution	$Q_c$	$Q_s$	$Q_t$
Monodisperse	2.4598	1.2351	1.2246
Lognormal <sup>a</sup>	2.4593	1.2346	1.2246

<sup>a</sup> $\sigma_g = 1.01$

**Blank**

## LITERATURE CITED

1. Milham, Merrill E., Electromagnetic Scattering by Magnetic Spheres: Theory and Algorithms, ERDEC-TR-207, U.S. Army Edgewood Research, Development and Engineering Center, MD, October 1994, UNCLASSIFIED Report.
2. Van de Hulst, H.C., Light Scattering by Small Particles, Dover, NY, 1981.
3. Abramowitz, M., and Stegun, I.A., Handbook of Mathematical Functions, GPO, Washington, DC, 1964.
4. Bohren, C.F., and Huffman, D.R., Absorption and Scattering of Light by Small Particles, Wiley, New York, NY, 1983.
5. Amos, D.E., "Algorithm 644: A Portable Package for Bessel Functions of a Complex Argument and Nonnegative Order," ACM Trans Math Software Vol 12, pp 265-273, 1986.
6. Wiscombe, W.J., Mie Scattering Calculations: Advances in Technique and Fast, Vector-Speed Computer Codes, NCAR/TN-140+STR, National Center for Atmospheric Research, Boulder, CO, 1979.
7. Kerker, M., Wang, D.-S., and Giles, C.L., "Electromagnetic Scattering by Magnetic Spheres," J. Opt. Soc. Am. Vol. 73, pp 765-767 (1983).

**Blank**

## APPENDIX A: PROGRAM LISTINGS

```

program disint
c***** ****
c
c Program disint computes extinction, scattering, absorption, and
c backscatter coefficients for lognormal distributions of homogeneous
c magnetic or nonmagnetic spheres.
c
c Merrill Milham      >>> version 1.0 <<<      JANUARY 1994
c
c inputs:
c           ALL INPUT IS IN LIST DIRECTED FORMAT
c
c line #1
c           mmd = mass median diameter (micron)      (real*8)
c           sigmag = geometric standard deviation    (real*8)
c           rho = density of the particulate material
c                   if rho .le. zero dimensionless
c                   extinction, scattering, absorption
c                   and backscatter coefficients are
c                   computed. (g cm-3)                  (real*8)
c           ndfp = number of logarithmically equally spaced
c                   abscissa values over which the various
c                   scattering coefficients are to be
c                   integrated                         (integer)
c           nsig = number of sigmas on each side of the mmd over
c                   which the scattering coefficients are to be
c                   integrated. No correction is made for excluded
c                   distribution tails.                 (real*8)
c
c line #2
c           nwl = number of wavelengths and material
c                   properties to be read             (integer)
c           flag = 'r' for refractive index data or
c                   'p' for permittivity data        (character*1)
c           wl = wavelength, up to 10 values (micron) (real*8)
c           mp = complex index or permittivity values
c                   to be read, up to 10 values     (complex*16)
c
c line #3
c           muu = complex permeability values to be read
c                   nwl values are required       (complex*16)
c
c output-
c line 1: mmx or mmd (micron)  sigmag  density (g cm-3) ndfp  nsig
c line 2: wavelength (micron)  ext. coef. (m sq / g)  scat. coef. (m sq / g)
c line 3:                      abs. coef. (m sq / g)  bsc. coef. (m sq/g/sr)
c line 4: permittivity permeability
c line 5: refractive index
c
c           If dimensionless quantities are computed, the output is so
c           labeled.
c
c subroutines used:

```

```

c
c          dxtcs - returns integrated extinction coefficients for a
c          homogeneous magnetic sphere or homogeneous
c          nonmagnetic sphere if the permeability = (1,0)
c
c*****=====
c
implicit none
real*8 pi
parameter (pi=3.14159265358979d0)
real*8 mmx,sigmag,nsig,dxe,dxs,dxa,dxb
real*8 mmd,xc,rho,wl(10),xe,xs,xa,xb
integer ndfp,i,nwl
complex*16 muu(10),mu,mp(10),eps
real*8 one,zero
parameter (one=1.d0,zero=0.d0)
character*1 flag
logical dflag
c
write(*,*) 'read(*,*) mmd,sigmag,rho,ndfp,nsig'
read(*,*) mmd,sigmag,rho,ndfp,nsig
write(*,*) 'read(*,*) nwl,flag,(wl(i),mp(i),i=1,nwl)'
read(*,*) nwl,flag,(wl(i),mp(i),i=1,nwl)
write(*,*) 'read(*,*) (muu(i),i=1,nwl)'
read(*,*) (muu(i),i=1,nwl)
write(*,*)
*
dflag=rho.le.zero
if(dflag) then
rho=one
write(*,1050)
write(*,1100) mmd,sigmag,rho,ndfp,nsig
else
write(*,1000) mmd,sigmag,rho,ndfp,nsig
end if
write(*,*)
c
do 100 i=1,nwl
*
if(dflag) then
mmx=mmd
wl(i)=one
else
mmx=pi*mmd/wl(i)
end if
*
if(flag.eq.'p') then
eps=mp(i)
else if(flag.eq.'r') then
eps=(mp(i))**2
else
stop 'material properties input error'

```

```

*      end if
*
*      mu=muu(i)
*
*      call dxtcs(mmx,sigmag,ndfp,nsig,eps,mu,dxe,dxs,dxa,dxb)
*
*      xc=1.d0/wl(i)/rho
*      xe=xc*dxe
*      xs=xc*dxs
*      xa=xc*dxa
*      xb=xc*dxb
*
*      write(*,1200) wl(i),xe,xs,xa,xb
*      write (*,1300) eps,mu
*      if(flag.eq.'r') write(*,1400) mp(i)
*      write(*,*) 
*
100 continue
c
200 stop 'done disint'
c
1000 format('mmd = ',f5.2,' sigmag = ',f5.2,' rho = ',f5.2,
&           ' ndfp = ',i5,' nsig = ',f5.2)
1050 format('dimensionless extinction coefficients will be computed')
1100 format('mmx = ',f5.2,' sigmag = ',f5.2,' rho = ',f5.2,
&           ' ndfp = ',i5,' nsig = ',f5.2)
1200 format('wl = ',f8.4,' ext = ',e10.5,' scat = ',e10.5,/13x,
&           ' abs = ',e10.5,' bsc = ',e10.5)
1300 format('eps = ',2f6.3,' mu = ',2f6.3)
1400 format('refractive index = ',2f6.3)
c
      end
c

```

```

subroutine dxtcs(mmx,sigmag,ndfp,nsig,eps,mu
& ,dxe,dxs,dxa,dxb)
c-----
c>>> subroutine dxtcs computes dimensionless extinction, scattering, <<
c>>> absorption, and backscatter coefficients for particulate ensembles <<
c>>> of spheres with lognormal particle size distributions <<
c
c      Merrill Milham      >>> version 1.0 <<<          JANUARY 1994
c
c      Inputs:
c          mmx = mass median size parameter                      (real*8)
c          sigmag = geometric standard deviation                  (real*8)
c          ndfp = number of equally spaced (logarithmically)    points at which integrand is to be
c                   evaluated                                         (integer)
c          nsig = number of geometric standard deviations on both sides of the mmx over which the integration extends (real*8)
c          eps = complex permittivity of particle material (complex*16)
c          mu = complex permeability of particle material (complex*16)
c
c      Outputs:
c          dxe = dimensionless extinction coefficient           (real*8)
c          dxs = dimensionless scattering coefficient          (real*8)
c          dxa = dimensionless absorption coefficient         (real*8)
c          dxb = dimensionless backscatter coefficient        (real*8)
c
c      subroutines used:
c          simp(real*8 function ) - does Simpson's rule
c          integration
c
c-----
c      implicit none
*
real*8 mmx,sigmag,nsig
integer ndfp
complex*16 eps,mu
real*8 dxe,dxs,dxa,dxb
*
integer mpsd
parameter (mpsd=3001)
real*8 tsume(mpsd),tsums(mpsd),tsumb(mpsd),dx,dc
real*8 lnsg,ez2x,r,xo,z0,z
real*8 theta,qext,qsca,qbac,g
real*8 simp
complex*16 s1,s2
integer i
*
real*8 pi
real*8 dco,sqrt2
parameter (pi=3.14159265358979d0)

```

```

parameter (dco=1.5d0*1.25331413731550d0,sqrt2=1.41421356237309d0)
real*8 zero,one,two,fourpi
parameter (zero=0.d0,one=1.d0,two=2.d0,fourpi=4.d0*pi)

c
lnsg=dlog(sigmag)
dc=dco/lnsg
r=sigmag**two*nsig/dble(ndfp-1)
xo=mmx/(sigmag**nsig)/r
zo=one/sqrt2/lnsg
c
90 if(ndfp.gt.mpsd) stop 'dxtcs: arrays too small $90'
*
theta=zero
do 100 i=1,ndfp
xo=xo*r
z=zo*dlog(xo/mmx)
ez2x=dexp(-z*z)/xo
*
call magsph(xo,eps,mu,0,theta,qext,qsca,qbac,g,s1,s2)
*
tsume(i)=qext*ez2x
tsums(i)=qsca*ez2x
tsumb(i)=qbac*ez2x
*
100 continue
c
dx=dlog(r)
dxe=dc*simp(tsume,ndfp,dx)
dxs=dc*simp(tsups,ndfp,dx)
dxa=ddim(dxe,dxs)
dxb=dc/fourpi*simp(tsumb,ndfp,dx)
c
return
c
end

```

```

subroutine magsph(x,eps,mu,numang,theta,
&                               qext,qsca,qbac,g,s1,s2)
c ****
c Subroutine magsph computes the scattering cross sections and angular
c scattering from a magnetic sphere. If the number of scattering angles
c is set to zero, only the cross sections (efficiencies) are returned.
c
c Merrill Milham           >>> version 2.0 <<<          SEPT 1993
c
c Inputs:
c      x = size parameter of the sphere                      (real*8)
c      eps = complex permittivity: epsr -i*epsi            (complex*16)
c      mu = complex permeability: mur - i*mui            (complex*16)
c      numang = number of scattering angles                 (integer)
c                  between 0 & 90 deg.
c      theta = scattering angles in degrees                  (real*8)
c                  theta(i) are entered between 0 & 90 deg.
c                  theta must increase monotonically. Results for
c                  supplementary angles (180 deg. - theta(i)) are
c                  also returned.
c
c Outputs:
c      qext = extinction efficiency                         (real*8)
c      qsca = scattering efficiency                        (real*8)
c      qbac = backscatter efficiency                     (real*8)
c      g = asymmetry factor                            (real*8)
c      s1 = scattered amplitude                          (complex*16)
c      s2 = scattered amplitude                          (complex*16)
c
c Subroutines used:
c
c      zbjy - returns one-half integer order J & Y Bessel functions
c
c References:
c
c      M. Kerker, D.-S. Wang, and C. L. Giles, "Electromagnetic scattering
c      by magnetic spheres," J. Opt. Soc. Am., 73, 765-767 (1983).
c
c      D. E. Amos, "Algorithm 644:A portable package for Bessel functions of
c      a complex argument and nonnegative order," ACM Trans. on Math.
c      Software, 12, 265-273 (1986).
c
c      M. Abramowitz and I. A. Stegun, "Handbook of Mathematical Functions,"
c      NBS Applied Math. Series 55, US Dept. of Commerce, Washington, DC
c      (1955).
c
c      W. J. Wiscombe, "Mie Scattering Calculations: Advances in Technique and
c      Fast, Vector-Speed Computer Codes," NCAR Tech. Note, NCAR/TN-140+STR
c      (1979)

```

```

c ****
c
c      implicit none
c
c      real*8 x
c      complex*16 eps,mu
c      integer numang
c      real*8 theta(1)
*
c      real*8 qext,qsca,qbac,g
c      complex*16 s1(1),s2(1)
c
c      integer al,nangl,nangl2
c      real*8 third
c      parameter (third=1.d0/3.d0,al=5100,nangl=255,nangl2=(nangl+1)/2)
c      complex*16 sp(nangl2),sm(nangl2),sps(nangl2),sms(nangl2)
*
c      complex*16 m,mcl,mxi,s,t,u,v,an,bn,xp
c      real*8 xi,dn,dnn,rn,tnp1,thetan
c      real*8 xmu(nangl),pi(nangl),pil(nangl),tau(nangl)
*
c      real*8 bjr(al),byr(al)
c      real*8 cjr(al),cji(al),cyr(al),cyi(al)
c      real*8 bjn,byn,bjl
c      real*8 sc,ca,t1,t2,t3,t4
c      complex*16 cjn,cyn,cjl,cyl,h2n,h2l
c      complex*16 anl,bnl,bs,anp,bnp,abp,abm,anpm,bnpm
c      complex*16 zt1,zt2
*
c      integer kstop,k,mm,n,j,j2,j3
*
c      real*8 cpi,zero,one,two,rad,half,fnu
c      complex*16 cdblei,cdble1,cdble0
c      parameter (cpi=3.1415926535897932384d0,zero=0.d0,one=1.d0)
c      parameter (two=2.d0,rad=cpi/180.d0,half=0.5d0,fnu=half)
c      parameter (cdblei=(0.d0,1.d0),cdble1=(1.d0,0.d0))
c      parameter (cdble0=(0.d0,0.d0))
c
c      kstop=idint(x+4.d0*x**third+4.d0)
*
c      if(kstop.le.al) then
c          continue
c          else
c              print*, 'magsph arrays too small: kstop =',kstop,' al =',al
c              stop
c      end if
c
c      if(numang.eq.0) then
c          s1(1)=cdble0
c          s2(1)=cdble0
c          else
c

```

```

    if(numang.le.nangl2) then
        continue
        else
    print*,numang,'scattering angles input: only',nangl2,' allowed'
        stop
    endif
*
    do 100 n=1,numang
    thetan=dabs(theta(n))
    theta(n)=thetan
*
        if(thetan.le.90.d0) then
            continue
            else
    print*, 'theta('',n,'')=',thetan,'scattering angles must be < 90 deg'
            stop
        end if
*
        thetan=rad*thetan
        xmu(n)=dcos(thetan)
*
        sp(n)=cdble0
        sm(n)=cdble0
        sps(n)=cdble0
        sms(n)=cdble0
        pi(n)=half
        pil(n)=zero
*
    100 continue
*
    end if
c
    m=zsqrt(mu*eps)
    mc1=m/mu
*
    xi=one/x
    mxi=xi/m
c
    call zbjy(x,m,kstop,fnu,bjr,byr,cjr,cji)
c
    bj1=bjr(1)
    cjl=dcmplx(cjr(1),cji(1))
    cyl=dcmplx(cyr(1),cyi(1))
    h2l=dcmplx(bjr(1),-byr(1))
*
    qext=zero
    qsca=zero
    bss=cdbl0
    g=zero
*
    anl=cdbl0
    bn1=cdbl0

```

```

dn=one
rn=one
tnp1=one
mm=1
c
do 300 k=2,kstop
*
tnp1=tnp1+two
t1=dn-rn
ca=one+rn
sc=rn
*
dnn=dn+one
rn=one/dnn
*
sc=sc+rn
*
bjn=bjr(k)
byn=byr(k)
cjn=dcmplx(cjr(k),cji(k))
cyn=dcmplx(cyr(k),cyi(k))
h2n=dcmplx(bjn,-byn)
*
xp=dn*mxi
s=cjl-xp*cjn
u=s*h2n
s=s*bjn
*
xp=dn*dcmplx(xi,zero)
t=cjn*(bjl-xp*bjn)
v=cjn*(h2l-xp*h2n)
*
an=(s-mc1*t)/(u-mc1*v)
bn=(mc1*s-t)/(mc1*u-v)
abp=an+bn
abm=an-bn
*
zt1=dconjg(an)
zt2=dconjg(bn)
qext=qext+tnp1*dble(abp)
qsca=qsca+tnp1*(an*zt1+bn*zt2)
bs=bs-(dn+half)*mm*abm
g=g+t1*dble(an*zt1+bn*zt2)+sc*dble(an*zt2)
*
if(numang.eq.0) then
  continue
  else
    anp=sc*abp
    bnp=sc*abm
    anpm=mm*anp
    bnpm=mm*bnp
*

```

```

do 375 j=1,numang
t1=xmu(j)*pi(j)
t4=t1-pil(j)
tau(j)=dn*t4-pil(j)
t2=pi(j)+tau(j)
t3=pi(j)-tau(j)
*
sp(j)=sp(j)+anp*t2
sms(j)=sms(j)+bnpm*t2
sm(j)=sm(j)+bnp*t3
sps(j)=sps(j)+anpm*t3
*
pil(j)=pi(j)
pi(j)=t1+ca*t4
*
375 continue
end if
*
dn=dnn
mm=-mm
anl=an
bnl=bn
*
bj1=bjn
cj1=cjn
cyl=cyn
h2l=h2n
*
300 continue
c
if(numang.eq.0) then
  continue
  else
    j2=2*numang
    do 500 j=1,numang
      j3=j2-j
      s1(j)=sp(j)+sm(j)
      s2(j)=sp(j)-sm(j)
      s1(j3)=sps(j)+sms(j)
      s2(j3)=sps(j)-sms(j)
    500 continue
*
  end if
c
xi=two*xi*xi
qext=xi*qext
qsca=xi*qsca
xi=two*xi
qbac=xi*bs*dconjg(bs)
g=xi/qsca*g
c
return

```

**c**

**end**

```

subroutine zbjy(x,m,nstop,fnu,
&                                bjr,byr,cjr,cji)
c ****
c
c Subroutine zbjy gets J & Y Bessel functions for use in
c sphere (fnu=0.50) or cylinder (fnu=0.0d0) scattering calculations.
c Scaled or nonscaled functions for argument z = m*x are returned depending
c on the magnitude of the product of the size parameter and the complex
c refractive index (zabs(z)). Nonscaled functions are returned if the
c imaginary part of the refractive index is zero. Nonscaled functions are
c returned for argument x.
c
c      Merrill Milham      >>> version: 2.0 <<<           JANUARY 1994
c
c inputs:
c      x = the size parameter of the cylinder.(real*8)
c      m = the complex refractive index, n - ik.(complex*16)
c      nstop = the highest order of the Bessel functions.(integer)
c      fnu = 0.5d0 for sphere calculations or
c            0.0d0 for cylinder calculations. (real*8)
c
c outputs:
c
c      bjr = real part of J(x) Bessel functions.(array: real*8)
c      byr = real part of Y(x) Bessel functions.(array: real*8)
c      cjr = real part of J(m*x) Bessel functions.(array: real*8)
c      cji = imag. part of J(m*x) Bessel functions.(array: real*8)
c
c subroutines used:
c
c      zbesj - returns J Bessel functions
c      zbesy - returns Y Bessel functions
c
c Reference: D. E. Amos, "Algorthim 644: A Portable Package for Bessel
c Functions of a Complex Argument and Nonnegative Order,"
c ACM Transcations on Mathematical Software, 12,265-273(1986).
c ****
c
c      implicit none
c      automatic cwrkr,cwrki,bji,byi
*
c      integer al
c      real*8 zero,xll,two
c      parameter (al=5100,zero=0.0d0,xll=1.d-1,two=2.d0)
*
c      real*8 bjr(1),byr(1)
c      real*8 x,fnu,cjr(1),cji(1)
c      real*8 zr,zi,dlmach
c      real*8 cwrkr(al),cwrki(al)
c      real*8 r1m5,elim,aa,alim

```

```

complex*16 m,z
integer kode,ierr,nz,nstop,n
integer k1,ilmach,k2,k
logical zflag,eflg1,eflg2

c
kode=1

c
call zbesj(x,zero,fnu,kode,nstop,bjr,cji,nz,ierr)

*
eflg1=ierr.eq.0
eflg2=nz.eq.0
if (eflg1.and.eflg2) then
    continue
else if (.not.eflg2.and.eflg1) then
    nstop=nstop-nz
    print*, 'zbesj error: ierr =',ierr,'nz =',nz
    else
    print*, 'inputs =',x,zero,fnu,kode,nstop
    print*, 'zbesj called from subroutine zbjy'
end if

c
call zbesy(x,zero,fnu,kode,nstop,byr,cji,nz,cwrkr,cwrki,ierr)

*
eflg1=ierr.eq.0
eflg2=nz.eq.0
if (eflg1.and.eflg2) then
    continue
else if (.not.eflg2.and.eflg1) then
    nstop=nstop-nz
    print*, 'zbesy error: ierr =',ierr,'nz =',nz
continue
    else
    print*, 'zbesy error: ierr =',ierr,'nz =',nz
    print*, 'inputs =',x,zero,fnu,kode,nstop
    print*, 'zbesy called from subroutine zbjy'
end if

c
z=m*x
zr=dble(z)
zi=dimag(z)
if(zi.ne.zero) then
*
k1 = ilmach(15)
k2 = ilmach(16)
rlm5 = dimach(5)
k = min0(iabs(k1),iabs(k2))
elim = 2.303d0*(dble(float(k))*rlm5-3.0d0)
k1 = ilmach(14) - 1
aa = rlm5*dble(float(k1))
aa = aa*2.303d0
alim = elim + dmax1(-aa,-41.45d0)
*

```

```

if(dabs(zi).gt.alim) kode=2
                           else
      continue
end if
c
call zbesj(zr,zi,fnu,kode,nstop,cjr,cji,nz,ierr)
*
eflg1=ierr.eq.0
eflg2=nz.eq.0
if (eflg1.and.eflg2) then
  continue
else if (.not.eflg2.and.eflg1) then
  nstop=nstop-nz
  print*, 'zbesj error: ierr =',ierr,'nz =',nz
continue
else
  print*, 'zbesj error: ierr =',ierr,'nz =',nz
  print*, 'inputs =',zr,zi,fnu,kode,nstop
  print*, 'zbesj called from subroutine zbjy'
end if
c
zflag=dabs(zi).lt.two*d1mach(1).and.x.lt.xll
if (.not.zflag) then
  continue
else
  do 100 n=1,nstop
    cji(n)=zero
100  continue
end if
*
zflag=dabs(zr).lt.two*d1mach(1).and.x.lt.xll
if (.not.zflag) then
  continue
else
  cji(1)=zero
  do 200 n=2,nstop
    if(mod(n,2).eq.0) then
      cjr(n)=zero
    else
      cji(n)=zero
    end if
200  continue
end if
c
return
c
end

```

```

real*8 function simp (y,n,dx)
c*****
c      function simp does simpson's rule integration
c      inputs:
c          y = array containing ordinate values      (real*8)
c          n = number of ordinate values          (integer)
c          dx = abscissa increment(constant) between
c                ordinate values                  (real*8)
c      output:
c          simp = estimated value of the integral   (real*8)
c*****
c
c      implicit none
c      real*8 s1,s2,s4,third
c      real*8 dx,y(1)
c      integer n,ne,i
c      parameter (third=1.d0/3.d0)
c
c      if(n.lt.5) then
c         write(6,1000)
1000 format(' error function : simp')
         write(6,'('' number of integration points must be .ge. 5 n = ''
+           ,i10)') n
         stop
c
c      else if(mod(n,2) .ne.1) then
c         write(6,1000)
c         write(6,'('' number of integration points must be odd. n = '',i10)
+           ') n
         stop
c
c      else
c         continue
c      end if
c
c      ne=n-3
c      s1=y(1)+y(n)
c      s2=0.d0
c      s4=y(n-1)
c
c      do 100 i=2,ne,2
c         s4=s4+y(i)
c         s2=s2+y(i+1)
100   continue
c
c      s4=4.d0*s4
c      s2=2.d0*s2
c
c      simp=third*dx*(s1+s2+s4)
c
c      return

```

**Blank**

## APPENDIX B: SAMPLE OUTPUT

### INPUT

```
10,1.01,-1,151,5  
1,'r',1,(1.5,-.1)  
(1,0)
```

### OUTPUT

```
read(*,*) mmd,sigmag,rho,ndfp,nsig  
read(*,*) nwl,flag,(wl(i),mp(i),i=1,nwl)  
read(*,*) (muu(i),i=1,nwl)  
  
dimensionless extinction coefficients will be computed  
  
mmx = 10.00 sigmag = 1.01 rho = 1.00 ndfp = 151 nsig = 5.00  
  
wl = 1.0000 ext = .11589E+01 scat = .58179E+00  
abs = .57709E+00 bsc = .33692E-02  
eps = 2.240-0.300 mu = 1.000 0.000  
refractive index = 1.500-0.100
```

### INPUT

```
10,1.4,-1,151,5  
1,'r',1,(1.5,-.1)  
(1,0)
```

### OUTPUT

```
read(*,*) mmd,sigmag,rho,ndfp,nsig  
read(*,*) nwl,flag,(wl(i),mp(i),i=1,nwl)  
read(*,*) (muu(i),i=1,nwl)  
  
dimensionless extinction coefficients will be computed  
  
mmx = 10.00 sigmag = 1.40 rho = 1.00 ndfp = 151 nsig = 5.00  
  
wl = 1.0000 ext = .12361E+01 scat = .63179E+00  
abs = .60428E+00 bsc = .35034E-02  
eps = 2.240-0.300 mu = 1.000 0.000  
refractive index = 1.500-0.100
```

### INPUT

```
10,1.4,-1,151,5  
1,'p',1,(2.24,-.3)  
(2.24,-.3)
```

### OUTPUT

```
read(*,*) mmd,sigmag,rho,ndfp,nsig
read(*,*) nwl,flag,(wl(i),mp(i),i=1,nwl)
read(*,*) (muu(i),i=1,nwl)

dimensionless extinction coefficients will be computed

mmx = 10.00 sigmag = 1.40 rho = 1.00 ndfp = 151 nsig = 5.00
wl = 1.0000 ext = .12292E+01 scat = .57928E+00
      abs = .64991E+00 bsc = .42778E-32
eps = 2.240-0.300 mu = 2.240-0.300
```