


# NAVAL POSTGRADUATE SCHOOL

## Monterey, California



**MODELING AND STATISTICAL ANALYSIS OF  
BIOASSAY DATA: MEDAKA CELL PROLIFERATION  
UNDER DEN AND TCE**

Donald P. Gaver  
Patricia A. Jacobs

DTIC QUALITY INSPECTED 2

October 1994

Approved for public release; distribution is unlimited.

Prepared for:  
U.S. Army Biomedical Research & Development Laboratory  
Ft. Detrick, MD 21702-5010

19941227 060

NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943-5000


Rear Admiral T. A. Mercer  
Superintendent


Harrison Shull  
Provost

This report was prepared for and funded by the U.S. Army Biomedical Research and Development Laboratory, Ft. Detrick, MD.

Reproduction of all or part of this report is authorized.


This report was prepared by:

  
DONALD P. GAVER, JR.  
Professor of Operations Research

  
PATRICIA A. JACOBS  
Professor of Operations Research

Reviewed by:

Released by:

  
PETER PURDUE  
Professor and Chairman  
Department of Operations Research

  
PAUL J. MARTO  
Dean of Research

| REPORT DOCUMENTATION PAGE  |  |   | Form Approved<br>OMB No. 0704-0188                        |  |
|--|--|---|---|--|
| Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503. |  |   |   |  |
| 1. AGENCY USE ONLY (Leave blank)   | 2. REPORT DATE<br>October 1994                           | 3. REPORT TYPE AND DATES COVERED<br>Technical           |   |  |
| 4. TITLE AND SUBTITLE<br>Modeling and Statistical Analysis of Bioassay Data; Medaka Cell Proliferation under DEN and TCE   |  |   | 5. FUNDING NUMBERS<br>RLLDG002JA                          |  |
| 6. AUTHOR(S)<br>Donald P. Gaver and Patricia A. Jacobs   |  |   |   |  |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br>Naval Postgraduate School<br>Monterey, CA 93943  |  |   | 8. PERFORMING ORGANIZATION REPORT NUMBER<br>NPS-OR-94-014 |  |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)<br>U.S. Army Biomedical Research & Development Laboratory<br>Ft. Detrick, MD 21702-5010  |  |   | 10. SPONSORING / MONITORING AGENCY REPORT NUMBER          |  |
| 11. SUPPLEMENTARY NOTES<br>The views expressed in this report are those of the authors and do not reflect the official policy or position of the Department of Defense or the U.S. Government.   |  |   |   |  |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT<br>Approved for public release; distribution is unlimited.  |  |   | 12b. DISTRIBUTION CODE                                    |  |
| 13. ABSTRACT (Maximum 200 words)<br>The response of <i>medaka</i> liver to the chemicals DEN and TCE is analyzed statistically. The analysis illustrates the application of methods useful in <i>environmetrics</i> , i.e. environmental statistics. It suggests an overall dose-response <i>effect</i> but not an easily-interpreted dose-response functional <i>relationship</i> .   |  |   |   |  |
| 14. SUBJECT TERMS<br>box plots, variance-stabilizing transformation, multiple comparisons, analysis of variance (ANOVA)  |  |   | 15. NUMBER OF PAGES<br>48                                 |  |
|  |  |   | 16. PRICE CODE  |  |
| 17. SECURITY CLASSIFICATION OF REPORT<br>Unclassified  | 18. SECURITY CLASSIFICATION OF THIS PAGE<br>Unclassified | 19. SECURITY CLASSIFICATION OF ABSTRACT<br>Unclassified | 20. LIMITATION OF ABSTRACT<br>UL                          |  |

# Modeling and Statistical Analysis of Bioassay Data: Medaka Cell Proliferation under DEN and TCE

by  
**Donald P. Gaver**  
and  
**Patricia A. Jacobs**

**Department of Operations Research  
Naval Postgraduate School  
Monterey, CA 93943**

|                                      |   |
|--------------------------------------|---|
| Accession For                        |   |
| NTIS                                 | CRA&I <input checked="" type="checkbox"/> |
| DTIC                                 | TAB <input type="checkbox"/>              |
| Unannounced <input type="checkbox"/> |   |
| Justification _____                  |   |
| By _____                             |   |
| Distribution / _____                 |   |
| Availability Codes                   |   |
| Dist                                 | Avail and/or Special                      |
| A-1                                  |   |

## 1. INTRODUCTION

This report describes an analysis of cell proliferation data, by liver slice, from an experiment using Japanese Medaka. Previous work has used summary data from the same experiment (Gaver and Jacobs, 1994a,b). Another relevant reference is to Morris (1993). A brief description of the experiment follows.

The medaka are exposed to differing levels of DEN and TCE in tanks of water. The treatment groups are: control, 10 mg/ℓ DEN, 100 mg/ℓ DEN, 0.1 mg/ℓ TCE, (10 mg/ℓ DEN with 0.1 mg/ℓ TCE), 1 mg/ℓ TCE, and (10 mg/ℓ DEN with 1 mg/ℓ TCE). Each treatment group has two replicate tanks. Eight animals in each tank were sacrificed on 4 August, 1993; this is sacrifice B. Eight additional animals in each tank were sacrificed on 20 August 1993; this is sacrifice D.

Each sacrificed fish was exposed to BrdU for 72 hours prior to sacrifice; any cell that is in S-phase during this time has a BrdU marker. Each sacrificed fish is frozen and sliced longitudinally into 7-micron sections. A third of the slices are stained with another agent. This agent stains nuclei with the BrdU marker black; these nuclei are called *positive*. It is 5 of the latter stained slices that are analyzed subsequently.

Five slices containing a portion of the liver are considered for each fish. A region of interest (ROI) is marked on the slice; the ROI is chosen to attempt to maximize the number of hepatocytes and minimize the number of nonhepatocytes present. The area of all of the hepatocytes within the region of interest is measured, and the area of positive nuclei within the region of interest is measured. The number of hepatocytes in the ROI, and the number of positive hepatocytes in the ROI were also counted for half the fish in sacrifice B.

A count measure of cell proliferation, the *count index* (CI), for a slice is the number of positive hepatocytes in the ROI divided by the number of hepatocytes in the ROI, multiplied by 100. Evaluation of this measure is very labor intensive. As an alternative, the following *area index* (AI) is used

$$AI = \text{Area Index} = \frac{\text{Area of positive nuclei in the ROI}}{\text{Area of hepatocytes in the ROI}} \times 100.$$

The area index is easier to obtain; however, it does not quantify cells in S-phase exactly as does the CI since cells are of different size, as are the areas resulting from the slicing process, which are the result of a random intersection with the cell.

Figure 1 displays a plot of the count index (divided by 100) versus area index (divided by 100), computed by slice, for those fish of sacrifice B for which both measures are available, along with a simple unweighted least-squares-fitted

straight line; also displayed is the least-squares line equation with the standard errors of the coefficients displayed in parentheses below the coefficients. There appears to be a satisfactory linear relationship between the count index and area index, indicating that AI and CI are generally measuring the same response. However the variability of the area index increases as the count index increases; this increase is generally associated with high DEN and TCE dose levels; its biological interpretation is not yet available. It suggests that cell sizes become more variable under dosage.

Figure 2 is a display of the slice area indices (divided by 100), by fish, for the two control tanks. Note the variability between fish and the somewhat greater variability between fish in tank 2 as compared to those in tank 1.

Several slice area datum appear to be missing or are of doubtful validity. These have been deleted from analytical consideration. They are listed in Table 1. An alternative might have been to use robust statistical procedures throughout; such procedures automatically down-weight highly discrepant observations. Furthermore, examination of the weights indicates discrepancy so that explanations can be sought. It seems likely that robust methods should be more widely used in environmental toxicology. Robust statistical methods are discussed seriously in Cox and Hinkley (1974). A less advanced treatment appears in Koopmann (1987).

A summary of the findings of the data analysis is as follows.

- a. Available data from sacrifice B suggests that there is a reasonably strong linear association between the count index and the area index. For ease of analysis the area index has been used throughout.

**TABLE 1**  
Slices Not Considered in the Data Analysis

| <b>MISSING SLICES: SACRIFICE B</b>        |               |      |                    |   |
|---|---------------|------|--------------------|---|
| Missing Measurements are left blank.      |               |      |                    |   |
| ID  | No. of Slices | Tank | Treatment          | Reason  |
| 9273802                                   | All           | 8    | 0.1 TCE            | Blank field   |
| 9273811                                   | All           | 9    | 10 DEN,<br>0.1 TCE | Blank field   |
| <b>MISSING SLICES: SACRIFICE D</b>        |               |      |                    |   |
| Missing Values appear to be coded with 0. |               |      |                    |   |
| ID  | No. of Slices | Tank | Treatment          | Reason  |
| 9273986                                   | All           | 2    | Control            | Area ROI = Positive Area = 0                            |
| 9285049                                   | All           | 10   | 1 TCE              | Area ROI = Positive Area = 0                            |
| 9274001                                   | All           | 4    | 10 DEN             | All positive areas = 0 with 2 slices having 1 mit. hep. |
| 9285079                                   | All           | 14   | 10 DEN,<br>1 TCE   | All positive areas = 0 with 3 slices having 1 mit. hep. |
| 9285022                                   | 1             | 7    | 0.1 TCE            | Area of ROI = 247.16, others of order 7500              |
| 9285043                                   | 1             | 9    | 10 DEN,<br>0.1 TCE | Positive area of slice = 0;<br>0 mit. cells             |
| 9285073                                   | 1             | 13   | 10 DEN,<br>1 TCE   | Positive area of slice = 0;<br>0 mit. cells             |

- b. There is evidence that the variances of the slice area indices over fish exposed to various treatments are approximately equal to the corresponding means. This relationship can lead to misleading conclusions if standard statistical procedures are used uncritically; furthermore, the results are less efficient than necessary. However, the variances of the *square roots* of the slice area

indices for fish subjected to the various treatment (DEN and TCE) levels appear approximately constant, i.e. far less dependent on the corresponding mean values. Consequently, the square roots of the area indices are used in subsequent data analyses. The underlying reason for the above data behavior is that positive counts are rare random events, hence tend to be approximately Poisson distributed. The square root transformation is known to stabilize the variance of such counts; see Miller (1986), p. 59. The same transformation should, and here does, stabilize the variance of count-associated areas.

- c. The means of the fish mean square root of area indices are considered for both sacrifices. There is generally more treatment effect for sacrifice B than for sacrifice D: examine  $p$ -values in lines 3 of Tables 2 and 3 for the overall analysis of variance indications to see that sacrifice D  $p$ -values are always larger than those for sacrifice B. Four out of five of the treatment means are significantly larger (at the 95% level) than that for the control for sacrifice B. There is no significant difference (95% level) between the treatment means and the control mean for sacrifice D. For sacrifice B, the treatment means for two out of the three levels of TCE with 10 mg/ℓ DEN are significantly larger than those without DEN for sacrifice B; there is no significant difference for sacrifice D.
- d. The control mean for sacrifice D is significantly larger than the control mean for sacrifice B. The other treatment means for sacrifice D are not significantly different than those for sacrifice B.



- e. There is some suggestion that in the later sacrifice D, the presence of TCE *lowers* the mean of the area index. The biological mechanisms likely to explain this behavior are not yet available.

The above results are from one-way analysis of variance (ANOVA) of the square roots of the area indices, augmented by *multiple comparison* methods; the latter allow all possible pairwise comparisons to be made with a specified experiment-wise error rate, here 5%.

#### **Brief Overall Summary of Findings to Date**

The above can be briefly, and simplistically, summarized as follows. While it can be said that there is a statistically significant difference between mean responses ( $\sqrt{\text{AI}}$ ) to the various treatments with DEN and TCE, no simple and interpretable dose-response patterns have been found. In particular, response does not appear to increase (or decrease) systematically with dose increase, where "dose" includes time of exposure as well as increases in chemical concentration dose levels. It remains to be seen whether the latter inconclusivity is lessened by the analysis of more data (later sacrifices), by finding that experimental problems or biases occurred, *or*, more exciting, that the dose-responses observed can be explained by biological mechanism, and that the findings essentially reappear when further experiments and data analyses are conducted.

Section 2 presents results of graphical displays of the data. Section 3 presents results of exploratory analyses of variance. Results of exploratory linear regression models are presented in Section 4. Multiple comparison results appear in Section 5.

## 2. GRAPHICAL SUMMARIES OF THE SLICE AREA INDICES

Figure 3a (respectively 3b) displays boxplots of the slice area indices by tank for sacrifice B, (respectively sacrifice D). The boxplots may be viewed as a graphical one-way analysis of variance. There is some tank effect within a treatment group. The greatest dose-response effect is clearly for the 100 DEN treatment.

Figures 4a and 4b display plots of the mean of the area indices for each tank versus the variance of the area indices for each tank. Also displayed is a 45° line. There appears to be a linear relationship between the mean and variance; in fact, the variances appear to be approximately equal to the means. This relationship between the means and variances may lead to misleading results if analysis of variance techniques are applied directly to the slice area indices; Miller (1986) and Box (1954) discuss the effects of inequality of variance on one-way analysis of variance. It is believed that such effects may well appear often in environmental toxicology, particularly where *counts*, or count-like phenomena, are found.

As noted earlier, one standard transformation that can be applied to data with variances approximately equal to means to attempt to make the variance of the transformed data more nearly constant is the *square root transformation*; see Miller (1986). Figures 5a and 5b display plots of the mean of the square root of the area indices for each tank versus the variance of the square root of the area indices for each tank. The variances now appear unrelated to the corresponding means. Figures 6a and 6b display boxplots of the square roots of the slice area indices by tank. Note that the lengths (heights between quartiles) of the boxes are less variable than are those for the boxplots of the raw area indices themselves. In the remainder of this paper the square root of area indices will be used.

### 3. EXPLORATORY ANALYSES OF VARIANCE

Results of exploratory analyses of variances appear in Tables 2 – 4. The basic data are summaries of area indices and the square roots of area indices for each fish. Since the boxplots indicate that the 100 DEN treatment is associated with much larger area indices, the analyses of variances were done with and without 100 DEN. Table 2 shows the results by tank. Recall that small  $p$ -values will indicate tank, hence treatment, effect. The first row of the table indicates that the tank means of the fish mean area index are significantly different. However the second row of the table indicates that the tank means of the logarithms of the variance (log variance) of area indices for each fish are also significantly different. The graphical analysis has already indicated this difference. Now recall that one of the assumptions of analysis of variance is that the data come from populations with equal variance, so apply the square-root transformation. Rows 3 and 4 report results using the square roots of the area indices. Note that the results of row 4 indicate that there is no significant difference between the tank means of the log variance of the slice square root of area index for each fish. However, the results of row 3 indicate that there *is* significant difference between the tank *means* of the mean of the slice square root area indices for each fish; note that the  $p$ -values for the tank means are smaller using the fish mean square root of area indices rather than the raw (untransformed) area indices for the analysis of variance (without the 100 DEN treatment). Thus, the difference in variances of the raw (untransformed) area indices appears to have masked some of the difference in means, presumably resulting from treatment effects.

Table 3 displays results for analyses of variance, with the two tanks in each treatment combined. Once again, the analysis of variance using the mean slice

square root of the area indices for each fish indicates that there is a significant difference between treatment means of the means, even without 100 DEN.

**TABLE 2**  
***p*-Values for Exploratory ANOVA of Area Indices (AI) by Tank**  
**(AI = [(Positive Area)/ROI Area] × 100)**  
**(Small *p*-Values Indicate Chemical-Tank Effect)**

| Data  | Sacrifice B           |                      | Sacrifice D           |                        |
|---|-----------------------|----------------------|-----------------------|------------------------|
|   | with 100 DEN          | without 100 DEN      | with 100 DEN          | without 100 DEN        |
| Mean slice AI for each fish in a tank                     | < 10 <sup>-16</sup>   | 3 × 10 <sup>-5</sup> | 6 × 10 <sup>-16</sup> | 2.9 × 10 <sup>-2</sup> |
| Log variance of slice AI for each fish in a tank          | 4 × 10 <sup>-11</sup> | 2 × 10 <sup>-5</sup> | 3 × 10 <sup>-2</sup>  | 0.54                   |
| Mean slice $\sqrt{AI}$ for each fish in a tank            | 4 × 10 <sup>-16</sup> | 8 × 10 <sup>-7</sup> | 1 × 10 <sup>-14</sup> | 1.6 × 10 <sup>-2</sup> |
| Log variance of slice $\sqrt{AI}$ for each fish in a tank | 0.16                  | 0.25                 | 0.89                  | 0.97                   |

**TABLE 3**  
***p*-Values for Exploratory ANOVA of Area Indices (AI) by Treatment**  
**(AI = [(Positive Area)/ROI Area] × 100)**  
**Treatment: 2 Tanks Combined**  
**(Small *p*-Values Indicate Treatment Effect)**

| Data   | Sacrifice B           |                      | Sacrifice D           |                      |
|--|-----------------------|----------------------|-----------------------|----------------------|
|  | with 100 DEN          | without 100 DEN      | with 100 DEN          | without 100 DEN      |
| Mean slice AI for each fish in a treatment                     | 1 × 10 <sup>-16</sup> | 1 × 10 <sup>-5</sup> | 1 × 10 <sup>-16</sup> | 2 × 10 <sup>-2</sup> |
| Log variance of slice AI for each fish in a treatment          | 4 × 10 <sup>-13</sup> | 3 × 10 <sup>-6</sup> | 1 × 10 <sup>-4</sup>  | 0.33                 |
| Mean slice $\sqrt{AI}$ for each fish in a treatment            | 2 × 10 <sup>-16</sup> | 9 × 10 <sup>-7</sup> | 4 × 10 <sup>-16</sup> | 9 × 10 <sup>-3</sup> |
| Log variance of slice $\sqrt{AI}$ for each fish in a treatment | 0.19                  | 0.46                 | 0.56                  | 0.89                 |

Table 4 displays results of an analysis of variance of the mean square root of area indices for each fish in a treatment but without the control. Once again there is evidence of significant differences between the treatment means.

We conclude that there is a definite treatment effect, i.e. response to different treatment levels, even if no treatment = control and the “strong” 100 DEN treatment responses are removed.

**TABLE 4**  
*p*-Values for Exploratory ANOVA of Area Indices (AI) by Treatment  
 Without Control  
 (AI = [(Positive Area)/ROI Area] × 100)  
 Treatment: 2 Tanks Combined  
 (Small *p*-Values Indicate Treatment Effect)

| Data                             | Sacrifice B     |                      | Sacrifice D           |                      |
|----------------------------------|-----------------|----------------------|-----------------------|----------------------|
|                                  | with<br>100 DEN | without<br>100 DEN   | with<br>100 DEN       | without<br>100 DEN   |
| Mean $\sqrt{\text{AI}}$ for fish | 0               | $1.9 \times 10^{-3}$ | $5.9 \times 10^{-15}$ | $7.4 \times 10^{-3}$ |

#### 4. EXPLORATORY LINEAR REGRESSION

Tables 5 – 8 report results of fitting exploratory linear regression models to the square root of the slice area indices for each fish. For Tables 5 and 6, the covariates are the level of DEN minus its mean; the level of TCE minus its mean; and an interaction term: {(level of DEN minus its mean) times (level of TCE minus its mean)}. The means were subtracted to give the interaction term a value other than 0 if the level of DEN or the level of TCE is 0. The linear regressions were fit with and without the 100 DEN treatment. The results of Table 5 are for sacrifice B and those of Table 6 are for sacrifice D.

**TABLE 5**  
**Sacrifice B:  $\sqrt{AI}$**   
**Linear Regression Coefficient Estimates with Standard Error**  
**and 95% Normal Confidence Intervals**

| <i>WITHOUT 100 DEN</i>                                |                       |                       |   |                |        |
|---|-----------------------|-----------------------|---|----------------|--------|
| CONSTANT  | DEN- $\overline{DEN}$ | TCE- $\overline{TCE}$ | (DEN- $\overline{DEN}$ ) $\times$<br>(TCE- $\overline{TCE}$ ) | R <sup>2</sup> | s.e.   |
| (SE)<br>[CI]  | (SE)<br>[CI]          | (SE)<br>[CI]          | (SE)<br>[CI]  | —<br>—         | —<br>— |
| 1.88  | 0.037                 | 0.526                 | 0.025   | 0.15           | 0.50   |
| (0.067)   | (0.005)               | (0.148)               | (0.010)   |                |        |
| [1.75,2.01]   | [0.028,0.046]         | [0.237,0.816]         | [0.005,0.045]   |                |        |
| $\overline{DEN} = 5.0 \quad \overline{TCE} = 0.372$   |                       |                       |   |                |        |
| <i>WITH 100 DEN</i>                                   |                       |                       |   |                |        |
| 1.77  | 0.029                 | 0.710                 | 0.039   | 0.56           | 0.50   |
| (0.040)   | (0.003)               | (0.117)               | (0.008)   |                |        |
| [1.69,1.85]   | [0.024,0.033]         | [0.48,0.94]           | [0.02,0.05]   |                |        |
| $\overline{DEN} = 18.81 \quad \overline{TCE} = 0.318$ |                       |                       |   |                |        |

**TABLE 6**  
**Sacrifice D:  $\sqrt{AI}$**   
**Linear Regression Coefficient Estimates with Standard Error**  
**and 95% Normal Confidence Intervals**

| <i>WITHOUT 100 DEN</i>                                |                       |                       |   |                |        |
|---|-----------------------|-----------------------|---|----------------|--------|
| CONSTANT  | DEN- $\overline{DEN}$ | TCE- $\overline{TCE}$ | (DEN- $\overline{DEN}$ ) $\times$<br>(TCE- $\overline{TCE}$ ) | R <sup>2</sup> | s.e.   |
| (SE)<br>[CI]  | (SE)<br>[CI]          | (SE)<br>[CI]          | (SE)<br>[CI]  | —<br>—         | —<br>— |
| 1.89  | 0.028                 | -0.402                | -0.025  | 0.08           | 0.50   |
| (0.07)  | (0.005)               | (0.151)               | (0.01)  |                |        |
| [1.76,2.03]   | [0.02,0.04]           | [-0.70,-0.11]         | [-0.05,-0.005]  |                |        |
| $\overline{DEN} = 4.88 \quad \overline{TCE} = 0.37$   |                       |                       |   |                |        |
| <i>WITH 100 DEN</i>                                   |                       |                       |   |                |        |
| 1.68  | 0.013                 | -0.050                | 0.0005  | 0.45           | 0.50   |
| (0.04)  | (0.003)               | (0.12)                | (0.008)   |                |        |
| [1.60,1.76]   | [0.008,0.018]         | [-0.29,0.19]          | [-0.02,0.02]  |                |        |
| $\overline{DEN} = 19.05 \quad \overline{TCE} = 0.315$ |                       |                       |   |                |        |

In both tables the values of  $R^2$  are small when the 100 DEN data are excluded. This implies that a linear function of the above explanatory variables does not explain the data well. However, the standard errors of the estimates are also small. This behavior suggests that, although there may be an association between levels of DEN and TCE and the square root of the area index, that association is not linear. Note that for sacrifice B all of the estimates of the coefficients are significantly positive, suggesting that increasing levels of DEN and TCE are associated with higher area indices. However, for sacrifice D, the estimate of the coefficient of the level of TCE and the estimate of the coefficient of the interaction term coefficient are significantly negative for the regression, even without the 100 DEN treatment. This suggests that for the later sacrifice, exposure to TCE may have an inhibitory effect. The only estimate of covariate that is significantly different from 0 for sacrifice D when 100 DEN is included is exposure to DEN.

Tables 7 and 8 report results of fitting linear regressions with the covariate being the level of TCE exposure to data from fish not exposed to DEN and to data from fish exposed to 10 mg/ $\ell$  DEN; the dependent variable is the square root of the slice area index. The  $R^2$  values are very small, indicating a lack of linear fit, but the estimate coefficients for TCE in the regressions using data from fish exposed to 10 mg/ $\ell$  DEN are formally significant. Once again, this behavior suggests that there may be an association but the association is not linear.

For sacrifice B, there was no significant effect for the level of TCE for the fish not exposed to DEN; for those fish exposed to 10 mg/ $\ell$  DEN the coefficient for level of TCE is significantly positive indicating that increasing levels of TCE are associated with increasing (square roots of) area indices.

**TABLE 7**  
**Sacrifice B:  $\sqrt{AI}$**   
**Linear Regression Coefficient Estimates with Standard Error**  
**and 95% Normal Confidence Intervals**  
**(Replicate Tanks Pooled)**

| <i>NO DEN</i>            |                     |                     |                          |                |
|--------------------------|---------------------|---------------------|--------------------------|----------------|
| CONSTANT<br>(SE)<br>[CI] | TCE<br>(SE)<br>[CI] | DEN<br>(SE)<br>[CI] | R <sup>2</sup><br>—<br>— | s.e.<br>—<br>— |
| 1.18                     | 0.056               | -                   | 0.002                    | 0.55           |
| (0.047)                  | (0.080)             | -                   |                          |                |
| [1.09, 1.27]             | [-0.10, 0.21]       | -                   |                          |                |
| <i>10 DEN</i>            |                     |                     |                          |                |
| 1.47                     | 0.31                | -                   | 0.09                     | 0.44           |
| (0.037)                  | (0.06)              | -                   |                          |                |
| [1.40, 1.54]             | [0.18, 0.43]        | -                   |                          |                |
| <i>0 DEN and 10 DEN</i>  |                     |                     |                          |                |
| 1.14                     | 0.18                | 0.038               | 0.14                     | 0.50           |
| (0.038)                  | (0.051)             | (0.005)             |                          |                |
| [1.06, 1.21]             | [0.08, 0.28]        | [0.29, 0.047]       |                          |                |

In sacrifice D, for those fish not exposed to DEN, the estimate of the coefficient of TCE is not significantly different than 0. However, for those fish exposed to 10 mg/l DEN the estimate of the coefficient of TCE is significantly *negative*, suggesting that for the fish of the later sacrifice that were exposed to DEN, the *greater* the level of TCE exposure, the *smaller* the (square root of) the area index. This effect calls for biological explanation.



**TABLE 8**

Sacrifice D:  $\sqrt{AI}$   
 Linear Regression Coefficient Estimates with Standard Error  
 and 95% Normal Confidence Intervals  
 (Replicate Tanks Pooled)

| <i>NO DEN</i>            |                     |                     |                          |                |
|--------------------------|---------------------|---------------------|--------------------------|----------------|
| CONSTANT<br>(SE)<br>[CI] | TCE<br>(SE)<br>[CI] | DEN<br>(SE)<br>[CI] | R <sup>2</sup><br>—<br>— | s.e.<br>—<br>— |
| 1.35                     | 0.059               | —                   | 0.004                    | 0.43           |
| (0.037)                  | (0.062)             | —                   |                          |                |
| [1.27, 1.42]             | [-0.063, 0.181]     | —                   |                          |                |
| <i>10 DEN</i>            |                     |                     |                          |                |
| 1.71                     | -0.189              | —                   | 0.02                     | 0.56           |
| (0.048)                  | (0.083)             | —                   |                          |                |
| [1.61, 1.80]             | [-0.352, -0.026]    | —                   |                          |                |
| <i>0 DEN and 10 DEN</i>  |                     |                     |                          |                |
| 1.39                     | -0.061              | 0.027               | 0.07                     | 0.50           |
| (0.038)                  | (0.052)             | (0.005)             |                          |                |
| [1.32, 1.47]             | [-0.16, 0.04]       | [0.017, 0.036]      |                          |                |

## 5. MULTIPLE COMPARISONS

The exploratory analyses of variances strongly rejected the null hypothesis that all the treatment means (even without the 100 DEN treatment) of the fish mean square root of the area indices are equal. Rejection of the null hypothesis does not indicate specifically which means are not equal. A method for discovering which means differ is called a multiple comparisons procedure. There are a number of different *multiple comparisons* procedures in the literature; see Miller (1981). We will use two of them.

## 5.1 Simultaneous Confidence Intervals using Studentized Range Distribution

The first procedure uses the studentized range distribution to construct simultaneous confidence statements about the true values of all differences of the treatment means. Table 9 describes the procedure to obtain simultaneous 95% confidence intervals for all differences of treatment means for one sacrifice without the 100 DEN treatment. The original procedure requires that there be an equal number of fish in each treatment. However, Ott *et al.* suggest step 4 in Table 9 if the number of fish in each treatment do not differ by much.

### 5.1a Treatment Means Minus Control Mean

Figure 7 presents some of the 95% simultaneous confidence intervals of the differences of treatment means for sacrifice B. It shows the confidence intervals for the treatment means minus the control mean for the fish mean square root of the area indices. Note that 4 out of the 5 intervals are significantly above 0 indicating that the treatments are associated with a larger mean square root area indices than those for the control. The greatest difference is that for the treatment of 10 mg/ℓ DEN with 1 mg/ℓ TCE.

Figure 8 presents some of the 95% simultaneous confidence intervals for the later sacrifice D. It shows the confidence intervals for the treatment mean minus the control mean. None of the treatment means is significantly different from the control mean.

### 5.1b Treatment Means for Treatments with Exposure to 10 mg/ℓ DEN Minus Those without Exposure to 10 mg/ℓ DEN

Figure 9 displays the sacrifice B 95% simultaneous confidence intervals for the difference between the treatment means for mean fish square root of the area indices for those treatments *with* 10 mg/ℓ DEN minus the treatment means for those treatments *without* 10 mg/ℓ DEN, by level of TCE exposure. Note that the

treatment means with 10 mg/ℓ DEN is significantly larger than that without for 0 mg/ℓ TCE and 1 mg/ℓ TCE. There is no significant difference for 0.1 mg/ℓ TCE.

**TABLE 9**

**To Obtain Tukey (Studentized Range Distribution) Simultaneous 95% Confidence Intervals for Treatment Mean Differences, One Sacrifice**

|    |   |
|----|---|
| 1. | There are 6 treatments: Control; 10 DEN; 0.1 TCE; 10 DEN with 0.1 TCE; 1 TCE; and 10 DEN with 1 TCE.  |
| 2. | There are ~88 within degrees of freedom $\left( \sum_{i=1}^6 (n_i - 1) \right)$ where $n_i$ is the number of fish in treatment $i$ .  |
| 3. | The 0.05 percentage point for the studentized range for 60 within-degrees-of-freedom and 6 treatment means is 4.16, from published tables, <i>BIOMETRIKA Tables for Statisticians</i> , Vol. 1. This is larger than the percentage point for 88 within-degrees-of-freedom. Thus, the constructed confidence intervals will be conservative: one can truly say that <i>all</i> pairwise difference comparisons are made with (95%) confidence. |
| 4. | Since the number of fish per treatment differs somewhat (due to missing fish) the harmonic mean of the number of fish per treatment is used $n = \frac{6}{\frac{1}{n_1} + \dots + \frac{1}{n_6}}$ where $n_i$ is the number of fish in treatment $i$ .  |
| 5. | The mean square within is $\frac{\sum_{i=1}^6 \sum_j (y_{ij} - \bar{y}_i)^2}{\sum_{i=1}^6 (n_i - 1)} = \text{MS}(\text{within})$ where $y_{ij}$ is the mean $\sqrt{\text{AI}}$ for fish $j$ in treatment $i$ and $\bar{y}_i$ is the mean of the mean $\sqrt{\text{AI}}$ for the fish in treatment $i$ .   |
| 6. | 95% confidence intervals for all pairs of means $\mu_i$ and $\mu'_i$ $(\bar{y}_i - \bar{y}_{i'}) \pm (4.16) \sqrt{\text{MS}(\text{within})/n}$  |

Figure 10 displays a similar plot for the later sacrifice D. There is no significant difference between the treatment means with 10 mg/ℓ DEN and those without.

### 5.1c Treatment Means for Sacrifice D Minus Treatment Means for Sacrifice B

Simultaneous confidence intervals are computed for all differences of the treatment means of the fish mean square root of the area indices for sacrifices B and D combined. Figure 11 displays six of the 95% simultaneous confidence intervals. It displays the 95% confidence intervals for the difference in treatment means between sacrifice D and sacrifice B. The only significant difference is for the control where the mean of the fish mean square root of area indices for sacrifice D is significantly larger than that for sacrifice B.

### 5.2 Studentized Maximum Modulus Confidence Intervals

Simultaneous confidence intervals for the treatment means themselves can be constructed using the studentized maximum modulus procedure; cf. Miller (1981). The procedure is as follows for the mean fish square root area index for the three treatment groups (10 mg/ℓ DEN, 0 mg/ℓ TCE), (10 mg/ℓ DEN, 0.1 mg/ℓ TCE), and (10 mg/ℓ DEN, 1 mg/ℓ TCE) for 1 sacrifice.

*To Obtain Simultaneous Confidence Intervals for 3 Treatment Means using the Studentized Maximum Modulus Distribution*

1. Compute the within degrees of freedom for the three treatments.

$$d = \sum_{i=1}^3 (n_i - 1)$$

where  $n_i$  is the number of fish in treatment  $i$ .

2. Compute the mean square within

$$MS(\text{within}) = \frac{\sum_{i=1}^3 \sum_j (y_{ij} - \bar{y}_i)^2}{\sum_{i=1}^3 (n_i - 1)}$$

where  $y_{ij}$  is the mean square root of the area indices for fish  $j$  in treatment  $i$  and  $\bar{y}_i$  is the mean of the fish means in treatment  $i$ .

3. Find the upper 0.05 point of the studentized maximum modulus distribution with parameters 3 (treatments) and  $d$  degrees of freedom,  $m(3, d)$ . Tables can be found in Miller (1981).
4. The three simultaneous 95% confidence intervals are

$$\bar{y}_i \pm m(3, d) \sqrt{MS(\text{within})/n_i}.$$

Figure 12 displays the 95% simultaneous confidence intervals for sacrifice B for the means of the fish mean of the square root of the slice area indices for those treatments having fish exposed to 10 mg/ℓ DEN. The means appear about the same for 0 mg/ℓ TCE and 0.1 mg/ℓ TCE. The mean for 1 mg/ℓ TCE appears to be somewhat larger.

Figure 13 displays the 95% simultaneous confidence intervals for sacrifice D for the means of those treatments with 10 mg/ℓ DEN by level of TCE. There is some suggestion that the presence of TCE is associated with a *lower* mean of the fish mean square root of the slice area indices.

## CONCLUSION

The above analyses illustrate the use of statistical methods appropriate for the kinds of data obtained by the medaka experiments. The methods of

transformation, analysis of variance, and multiple comparisons are useful and powerful for the initial data analyses, suggesting some surprising dose-response relations that are worthy of careful further biological investigation and explanation. Alternative methods can also be applied, and should yield the same general insights.

## REFERENCES

- Bickel, P. J. and Doksum, K. A. *Mathematical Statistics: Basic Ideas and Selected Topics*, Holden-Day, Inc., San Francisco, 1977.
- Box, G. E. P. "Some theorems on quadratic forms applied in the study of analysis of variance problems, I. Effect of inequality of variance in the one-way classification." *Annals of Mathematical Statistics*, **25**, pp. 290-302.
- Cox, D. R. and Hinkley, D. V. *Theoretical Statistics*, Chapman and Hall, New York, 1974.
- Gaver, D. P. and Jacobs, P. A. "Assessment of liver modification and cell proliferation in medaka under DEN and TCE dosage, using data available 4/26/94, suggestions for statistical analysis and results from exploratory data analysis." Working Paper, April 26, 1994.
- Gaver, D. P. and Jacobs, P. A. "Comparison of area indices in medaka livers for sacrifices at different dose time combinations using data available 5/19/94." Working Paper, May 19, 1994.
- Huff, J. "Absence of morphologic correlation between chemical toxicity and chemical carcinogenesis." *Environmental Health Perspectives*, **101** (Suppl. 5) pp. 45-54, 1993.
- Koopmann, L. H. *Introduction to Contemporary Statistical Methods*, Duxbury Press, Boston, 1987.
- Miller, R. G. Jr. *Beyond ANOVA, Basics of Applied Statistics*, John Wiley & Sons, New York, 1986.
- Miller, R. G. Jr. *Simultaneous Statistical Inference*, Second Edition, Springer-Verlag, New York, 1981.
- Morris, R. W. "Analysis of cell proliferation data." *Environmental Health Perspectives*, **101** (Suppl. 5) pp. 73-78, 1993.
- Ott, L. and Hildebrand, D. K. *Statistical Thinking for Managers*, PWS Publishers, Boston, 1983.

# AREA INDEX VERSUS COUNT INDEX BY SLICE

SCATTER PLOT, SSZ=275





Figure 1



# SACRIFICE B, CONTROL

## RATIO POSITIVE AREA TO ROI AREA—TANK 1



## RATIO POSITIVE AREA TO ROI AREA—TANK 2




Figure 2  
22

SACRIFICE B: AREA INDICES

$((\text{POSITIVE AREA}) \div (\text{AREA ROI})) \times 100$  BY SLICE FROM ALL FISH




Figure 3a  
23

SACRIFICE D: AREA INDICES

$((\text{POSITIVE AREA}) \div (\text{AREA ROI})) \times 100$  BY SLICE FROM ALL FISH





Figure 3b  
24

# SACRIFICE B: PLOT MEAN VS VARIANCE FOR EACH TANK

VARIANCE OF RATIO OF (POSITIVE AREA ÷ AREA ROI) × 100  
USING SLICES WITHIN A TANK



# SACRIFICE D: PLOT MEAN VS VARIANCE FOR EACH TANK

VARIANCE OF RATIO OF (POSITIVE AREA ÷ AREA ROI) × 100  
USING SLICES WITHIN A TANK




Figure 4b  
26

# SACRIFICE B: SQUARE ROOT AREA INDEX BY TANK




Figure 5a  
27

# SACRIFICE D: SQUARE ROOT AREA INDEX BY TANK




Figure 5b  
28

SACRIFICE B

SQUARE ROOT AREA INDICES  $((\text{POSITIVE AREA}) \div (\text{AREA ROI})) \times 100$

BY SLICE FROM ALL FISH




Figure 6a  
29



SACRIFICE D

SQUARE ROOT AREA INDICES ((POSITIVE AREA) ÷ (AREA ROI)) × 100  
 BY SLICE FROM ALL FISH




Figure 6b  
30

SACRIFICE B: SQUARE ROOT OF AREA INDICES

MEAN OF TREATMENT MEANS - MEAN OF CONTROL MEANS  
 +/- 95 PERC TUKEY SIMUL CONF LIMITS FOR ALL DIFF




Figure 7  
31

# SACRIFICE D: SQUARE ROOT OF AREA INDICES

MEAN OF TREATMENT MEANS - MEAN OF CONTROL MEANS  
+/- 95 PERC TUKEY SIMUL CONF LIMITS FOR ALL DIFF




Figure 8<sup>32</sup>

SACRIFICE B:MEAN(10 DEN)-MEAN(ODEN)

95 PERCENT STUDENTIZED RANGE SIMULTANEOUS CONFIDENCE INTERVALS  
FOR ALL TREATMENT MEAN DIFFERENCES




Figure 9  
33

SACRIFICE D:MEAN(10 DEN) - MEAN(ODEN)

95 PERCENT STUDENTIZED RANGE SIMULTANEOUS CONFIDENCE INTERVALS  
FOR ALL TREATMENT MEAN DIFFERENCES




Figure 10  
34

# SQUARE ROOT OF AREA INDICES

MEAN OF MEANS BY TREATMENT FOR SACR D-SACR B  
 95 PERCENT CONF LIMITS OVER ALL DIFF




Figure 11

SACRIFICE B:10 DEN; SQUARE ROOT OF AREA INDICES  
 STUDENTIZED MAXIMUM MODULUS SIMULTANEOUS CONFIDENCE INTERVALS  
 95 PERCENT




Figure 12  
 36

SACRIFICE D:10 DEN; SQUARE ROOT OF AREA INDICES  
STUDENTIZED MAXIMUM MODULUS SIMULTANEOUS CONFIDENCE INTERVALS  
95 PERCENT




Figure 13  
37



## INITIAL DISTRIBUTION LIST

1. Research Office (Code 08) .....1  
Naval Postgraduate School  
Monterey, CA 93943-5000
2. Dudley Knox Library (Code 52) .....2  
Naval Postgraduate School  
Monterey, CA 93943-5002
3. Defense Technical Information Center.....2  
Cameron Station  
Alexandria, VA 22314
4. Department of Operations Research (Code OR).....1  
Naval Postgraduate School  
Monterey, CA 93943-5000
5. Prof. Donald P. Gaver (Code OR/Gv).....5  
Naval Postgraduate School  
Monterey, CA 93943-5000
6. Prof. Patricia A. Jacobs (Code OR/Jc).....5  
Naval Postgraduate School  
Monterey, CA 93943-5000
7. Dr. J. Abrahams, Code 1111, Room 607 .....1  
Mathematical Sciences Division, Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000
8. Dr. John Bailar .....1  
468 N St. NW  
Washington, DC 20024
9. Dr. David Brillinger.....1  
Statistics Department  
University of California  
Berkeley, CA 94720

10. Dr. David Burman.....1  
 AT&T Bell Telephone Laboratories  
 600 Mountain Avenue  
 Murray Hill, NJ 07974
  
11. Prof. Brad Carlin.....1  
 School of Public Health  
 University of Minnesota  
 Mayo Bldg. A460  
 Minneapolis, MN 55455
  
12. Dr. Robert Carpenter .....1  
 NAMRI/Navy Toxicology Detachment  
 Bldg. 433, Area B  
 Wright-Patterson AFB, OH 45433-6503
  
13. Center for Naval Analyses.....1  
 4401 Ford Avenue  
 Alexandria, VA 22302-0268
  
14. Prof. H. Chernoff.....1  
 Department of Statistics  
 Harvard University  
 1 Oxford Street  
 Cambridge, MA 02138
  
15. Dr. Edward G. Coffman, Jr.....1  
 AT&T Bell Telephone Laboratories  
 600 Mountain Avenue  
 Murray Hill, NJ 07974
  
16. Dr. John Copas.....1  
 Dept. of Mathematics,  
 University of Birmingham  
 P. O. Box 363  
 Birmingham B15 2TT  
 ENGLAND
  
17. Professor Sir David Cox.....1  
 Nuffield College  
 Oxford, OXI INF  
 ENGLAND

|     |                                   |   |
|-----|-----------------------------------|---|
| 18. | Professor H. G. Daellenbach.....  | 1 |
|     | Department of Operations Research |   |
|     | University of Canterbury          |   |
|     | Christchurch, NEW ZEALAND         |   |
| 19. | Dr. S. R. Dalal.....              | 1 |
|     | Bellcore                          |   |
|     | 445 South Street                  |   |
|     | Morristown, NJ 07962-1910         |   |
| 20. | Dr. D. F. Daley.....              | 1 |
|     | Statistic Dept. (I.A.S.)          |   |
|     | Australian National University    |   |
|     | Canberra, A.C.T. 2606             |   |
|     | AUSTRALIA                         |   |
| 21. | Dr. B. Doshi.....                 | 1 |
|     | AT&T Bell Laboratories            |   |
|     | HO 3M-335                         |   |
|     | Holmdel, NJ 07733                 |   |
| 22. | Naihua Duan.....                  | 1 |
|     | RAND Corporation                  |   |
|     | Santa Monica, CA 90406            |   |
| 23. | Prof. Bradley Efron.....          | 1 |
|     | Statistics Dept.                  |   |
|     | Sequoia Hall                      |   |
|     | Stanford University               |   |
|     | Stanford, CA 94305                |   |
| 24. | Dr. Guy Fayolle.....              | 1 |
|     | I.N.R.I.A.                        |   |
|     | Dom de Voluceau-Rocquencourt      |   |
|     | 78150 Le Chesnay Cedex            |   |
|     | FRANCE                            |   |
| 25. | Dr. M. J. Fischer.....            | 5 |
|     | Defense Communications Agency     |   |
|     | 1860 Wiehle Avenue                |   |
|     | Reston, VA 22070                  |   |

26. Prof. George S. Fishman.....1  
 Curr. in OR & Systems Analysis  
 University of North Carolina  
 Chapel Hill, NC 20742
27. Henry S. Gardner .....1  
 U.S. Army Biological Research & Development Laboratory  
 Ft. Detrick, MD 21702-5010
28. Dr. Neil Gerr .....1  
 Office of Naval Research  
 Arlington, VA 22217
29. Prof. Linda V. Green.....1  
 Graduate School of Business  
 Columbia University  
 New York, NY 10027
30. Dr. D. C. Hoaglin .....1  
 Department of Statistics  
 Harvard University  
 1 Oxford Street  
 Cambridge, MA 02138
31. Prof. D. L. Iglehart.....1  
 Dept. of Operations Research  
 Stanford University  
 Stanford, CA 94350
32. Institute for Defense Analysis.....1  
 1800 North Beauregard  
 Alexandria, VA 22311
33. Prof. J. B. Kadane.....1  
 Dept. of Statistics  
 Carnegie-Mellon University  
 Pittsburgh, PA 15213
34. Dr. Jon Kettenring.....1  
 Bellcore  
 445 South Street  
 Morris Township, NJ 07962-1910

35. Koh Peng Kong.....1  
 OA Branch, DSO  
 Ministry of Defense  
 Blk 29 Middlesex Road  
 SINGAPORE 1024
36. Dr. Prabha Kumar.....1  
 Defense Communications Agency.....1  
 1860 Wiehle Avenue  
 Reston, VA 22070
37. Dr. A. J. Lawrence.....1  
 Dept. of Mathematics,  
 University of Birmingham  
 P. O. Box 363  
 Birmingham B15 2TT  
 ENGLAND
38. Prof. M. Leadbetter.....1  
 Department of Statistics  
 University of North Carolina  
 Chapel Hill, NC 27514
39. Prof. J. Lehoczky .....1  
 Department of Statistics  
 Carnegie-Mellon University  
 Pittsburgh, PA 15213
40. Dr. Colin Mallows.....1  
 AT&T Bell Telephone Laboratories  
 600 Mountain Avenue  
 Murray Hill, NJ 07974
41. Prof. R. Douglas Martin.....1  
 Department of Statistics, GN-22  
 University of Washington  
 Seattle, WA 98195
42. Prof. M. Mazumdar .....1  
 Dept. of Industrial Engineering  
 University of Pittsburgh  
 Pittsburgh, PA 15235

43. Dr. Sati Mazumdar .....1  
 Biostatistics Dept.  
 University of Pittsburgh  
 Graduate School of Public Health  
 Pittsburgh, PA 15261
44. Prof. Carl N. Morris.....1  
 Statistics Department  
 Harvard University  
 1 Oxford St.  
 Cambridge, MA 02138
45. Prof. F. W. Mosteller .....1  
 Department of Statistics  
 Harvard University  
 1 Oxford St.  
 Cambridge, MA 02138
46. Operations Research Center, Rm E40-164.....1  
 Massachusetts Institute of Technology  
 Attn: R. C. Larson and J. F. Shapiro  
 Cambridge, MA 02139
47. Dr. John Orav.....1  
 Biostatistics Department  
 Harvard School of Public Health  
 677 Huntington Ave.  
 Boston, MA 02115
48. Dr. Jim Petty .....1  
 National Biological Survey  
 4200 New Haven Road  
 Columbia, MO 65201
49. Prof. Maria Rieders.....1  
 Dept. of Industrial Eng.  
 Northwestern Univ.  
 Evanston, IL 60208
50. Dr. John E. Rolph.....1  
 RAND Corporation  
 1700 Main St.  
 Santa Monica, CA 90406

51. Prof. M. Rosenblatt .....1  
 Department of Mathematics  
 University of California, San Diego  
 La Jolla, CA 92093
52. Prof. Frank Samaniego .....1  
 Statistics Department  
 University of California  
 Davis, CA 95616
53. Prof. W. R. Schucany .....1  
 Dept. of Statistics  
 Southern Methodist University  
 Dallas, TX 75222
54. Prof. D. C. Siegmund .....1  
 Dept. of Statistics  
 Sequoia Hall  
 Stanford University  
 Stanford, CA 94305
55. Prof. H. Solomon .....1  
 Department of Statistics  
 Sequoia Hall  
 Stanford University  
 Stanford, CA 94305
56. Andrew Solow .....1  
 Woods Hole Oceanographic Institute  
 Woods Hole, MA 02543
57. Prof. W. Stuetzle .....1  
 Department of Statistics  
 University of Washington  
 Seattle, WA 98195
58. Dr. D. J. Svendsgaard .....1  
 Health Effects Research Lab  
 U. S. Environmental Protection Agency  
 Biostatistics Branch MD55  
 Research Triangle Park, NC 27711

59. Prof. J. W. Tukey .....1  
 Statistics Dept., Fine Hall  
 Princeton University  
 Princeton, NJ 08540
60. Prof. David L. Wallace .....1  
 Statistics Dept., University of Chicago  
 5734 S. University Ave.  
 Chicago, IL 60637
61. Dr. Ed Wegman .....1  
 George Mason University  
 Fairfax, VA 22030
62. Dr. Alan Weiss .....1  
 Rm. 2C-118  
 AT&T Bell Laboratories  
 600 Mountain Avenue  
 Murray Hill, NJ 07974-2040
63. Dr. P. Welch .....1  
 IBM Research Laboratory  
 Yorktown Heights, NY 10598
64. Prof. Roy Welsch .....1  
 Sloan School  
 M.I.T.  
 Cambridge, MA 02139