

AD _____


CONTRACT NO: DAMD17-91-c-1048

TITLE: Further Development and Validation of the Frog Embryo Teratogenesis Assay-Xenopus (FETAX).

PRINCIPAL INVESTIGATOR: John A. Bantle

CONTRACTING ORGANIZATION: Oklahoma State University
430 LSW
Zoology Dept.
Stillwater, OK 74078

REPORT DATE: 12/30/1994



TYPE OF REPORT: Midterm

19950809 015

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 8

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND DATES COVERED	
	12/30/1994	Midterm: March 8, 1991-Nov. 30, 1994	
4. TITLE AND SUBTITLE		5. FUNDING NUMBERS	
Further Development and Validation of the Frog Embryo Teratogenesis Assay- Xenopus (FETAX).		DAMD17-91-C-1048	
6. AUTHOR(S) John A. Bantle,			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER	
Oklahoma State University Stillwater, OK 74078			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			
11. SUPPLEMENTARY NOTES Phase III testing of FETAX using 6 different chemicals			
12a. DISTRIBUTION/AVAILABILITY STATEMENT		12b. DISTRIBUTION CODE	
Approved for public release; distribution unlimited			
13. ABSTRACT (Maximum 200 words) FETAX is a 96-h whole embryo developmental toxicity screening assay that can be used in ecotoxicology and in detecting mammalian developmental toxicants when an <i>in vitro</i> metabolic activation system is employed. A standardized ASTM guide for the conduct of FETAX has been published. As part of the ASTM process, a three phase interlaboratory validation study was undertaken to evaluate the repeatability and reliability of FETAX. Seven different laboratories participated in the study. In Phase I, FETAX proved to be more repeatable and reliable than many other bioassays. However, some excessive variation was observed in a few laboratories. Some of this variation may have been due to an initial lack of experience with the assay by some technicians. Phase II showed far less variability than Phase I. Nonteratogens showed the most consistent results while more variability was observed for the two teratogens tested. Interlaboratory coefficient of variation values for all FETAX endpoints ranged from 7.3 to 54.7. Phase III, which is reported here, used coded samples and test concentration ranges selected by each laboratory. Phase III yielded results similar to Phase I. Analysis of the causes of variation suggested that some technicians judged malformations more severely than others and that the ranges of concentrations tested by some of the laboratories varied greatly. A new protocol for selecting concentrations was written to reduce variation from this source. Testing to date suggests that FETAX is a repeatable and reliable bioassay.			
14. SUBJECT TERMS		15. NUMBER OF PAGES 185	
FETAX, <u>Xenopus</u> , teratogen, interlaboratory validation, developmental toxicity		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	
19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified		20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Avail and/or Date	Specie
P-1	


Dr. J. A. Bantle 1/9/94
PI - Signature Date

TABLE OF CONTENTS

FRONT COVER.....	i
SF298 REPORT DOCUMENTATION PAGE.....	ii
FOREWORD.....	iii
TABLE OF CONTENTS.....	iv
ABSTRACT.....	1
INTRODUCTION.....	2
MATERIALS AND METHODS.....	4
BODY.....	7
CONCLUSIONS.....	11
REFERENCES.....	17
TABLES.....	26
FIGURES.....	35
APPENDIX (supplementary data for each test compound)	
MONOSODIUM GLUTAMATE.....	42
ASCORBIC ACID.....	65
β -AMINOPROPIONITRILE.....	88
SODIUM ACETATE.....	113
SODIUM ARSENATE.....	138
COPPER SULFATE.....	160

Accession For	
NTIS GRA&I	<input type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Distr	Avail and/or Specia
A-1	

Abstract - The Frog Embryo Teratogenesis Assay- *Xenopus* (FETAX) is a 96-h whole embryo developmental toxicity screening assay that can be used in ecotoxicology and in detecting mammalian developmental toxicants when an *in vitro* metabolic activation system is employed. A standardized American Society for Testing and Materials (ASTM) guide for the conduct of FETAX has been published along with a companion atlas that helps in embryo staging and identifying malformations. As part of the ASTM process, a three phase interlaboratory validation study was undertaken to evaluate the repeatability and reliability of FETAX. Seven different laboratories participated in the study. In Phase I, FETAX proved to be more repeatable and reliable than many other bioassays. However, some excessive variation was observed in a few laboratories. Some of this variation may have been due to an initial lack of experience with the assay by some technicians. Phase II showed far less intra- and interlaboratory variability than Phase I. Nonteratogens showed the most consistent results while more variability was observed for the two teratogens tested. Interlaboratory coefficient of variation values for all FETAX endpoints ranged from 7.3 to 54.7. Phase III, using coded samples and test concentration ranges selected by each laboratory showed results similar to Phase I. Analysis of the causes of variation suggested that some technicians judged malformations more severely than others and that the ranges of concentrations tested by some of the laboratories varied greatly. A new protocol for selecting concentrations for initial testing were written to reduce variation from this source. Testing to date suggests that FETAX is as repeatable and reliable as other standard bioassays.

INTRODUCTION

FETAX (Frog Embryo Teratogenesis Assay- *Xenopus*) is a 96-h whole embryo developmental toxicity test that utilizes the embryos of the South African clawed frog, *Xenopus laevis*. FETAX was initially designed as an indicator of potential human developmental health hazards and this has been enhanced by the development of an *in vitro* metabolic activation system using Aroclor 1254- and isoniazid-induced rat liver microsomes.¹⁻⁹ FETAX has undergone extensive validation using single chemicals of known mammalian developmental toxicity and additional work is currently being done.¹⁰

FETAX is also applicable to aquatic toxicity assessments and is well suited for testing complex mixtures such as industrial effluents.¹¹⁻¹⁴ This amphibian developmental toxicity test may help in studies designed to discover the reasons for the reported world-wide disappearance of amphibians even in pristine locations.^{15,16} This decline may be due in part to normal population fluctuations caused by climatological factors or by anthropogenic factors.¹⁷ However, in at least one case, frog eggs failed to develop in pond water but developed normally when moved to the laboratory.¹⁸ It is, therefore, possible that some decline may be due to chemical pollution and FETAX may be used to investigate the extent and causes of the decline.

An American Society for Testing and Materials *New Standard Guide for the conduct of FETAX* was published along with a companion *Atlas of Abnormalities* that helps in embryo staging and identifying malformations.¹⁹⁻²⁰ As part of the ASTM process, an interlaboratory validation study (ILS) was undertaken to determine the repeatability and reliability of FETAX. A three phase experimental plan with seven participants was designed. Phase I was a training and protocol evaluation phase in which the identity of the three test materials was known. Because they had been previously tested in FETAX, the same concentrations needed to establish the 96-h LC50 and EC50 (malformation) were used by all laboratories. Phase I²¹ showed that proper technician training was important in providing repeatable and reliable data. Based on Phase I results the following protocol modifications were necessary: 1) a 10% malformation rate in FETAX solution controls is acceptable, instead of 7% as specified in the ASTM Guide, 2) a change from separately pipetting toxicant and diluent directly into dishes to a procedure where toxicant and diluent are mixed in 25 ml volumes in 50 ml flasks prior to aliquoting into dishes and, 3) the elimination of 6-aminonicotinamide as the positive control. Phase I results in terms of variability and correspondence to historical data were very good with only occasional high variation observed in some laboratories. Phase II was designed to be similar to Phase I except that the identity of the test materials was not

known²². All technicians had greater experience in Phase II than in Phase I and the nature of the test compounds may have played a role in the results obtained. Phase II showed far less intra- and interlaboratory variability than Phase I. Nonteratogens showed the most consistent results while more variability was observed for the two teratogens tested. Interlaboratory coefficient of variation values for all FETAX endpoints ranged from 7.3 to 54.7%. The most variable endpoint was the MCIG and the least variable was the LC50. Phase III, reported here, was designed to test FETAX using six test compounds not previously tested in FETAX in a blind testing format with each laboratory responsible for determining the concentrations to be tested. The *in vitro* metabolic activation system for FETAX was not employed in either Phases I, II or III.

MATERIALS AND METHODS

Interlaboratory Study Organization

Each laboratory participating in this interlaboratory study included a principal investigator and, with the exception of Dr. Doug Dawson, one or two primary technicians. At least six different laboratories tested each compound in the study. Each laboratory utilized one technician with the exception of one laboratory which utilized two independent technicians. For ease of discussion, we will refer to seven laboratories being used in total. The technicians performed FETAX while the principal investigators compiled and reported the data to a central coordinator. The experimental results were screened by the coordinator to see that they complied with the standard protocol established in the ASTM Standard Guide ¹⁹ as well as the modifications made to the standard protocol following Phase I.

Interlaboratory Study Phase III Procedure

Tests were performed as specified in the ASTM Standard Guide ¹⁹. All technicians generated their own concentration ranges based on range-finding studies as outlined in the Standard Guide¹⁹. Technicians did not know which chemical they were testing as all samples were coded. Phase I modifications were used as previously mentioned ²¹.

Samples of test solutions were retained from selected concentrations and were frozen in specially supplied borosilicate glass tubes with Teflon-lined lids (Fisher®, Dallas, TX). At the end of the experiment, the samples from each laboratory were shipped on dry ice to Dr. Bantle's laboratory for concentration analysis in the event that the cause of excess variability had to be identified. Embryos preserved in 3% v/v formalin were stored in glass scintillation vials (RPI®, Elkhart, IN) and were available to be sent to Dr. Bantle's laboratory for verification of results when necessary.

Assay Procedure

Xenopus culture, breeding procedures, and egg sorting were described previously in the ASTM Standard Guide ¹⁹ and the Atlas of Abnormalities ²⁰. Adult *Xenopus laevis* frogs were obtained from Xenopus I (Ann Arbor, Michigan).

Test Compounds

Chemicals tested in Phase III were monosodium glutamate (CAS # 6106-04-3), ascorbic acid (CAS # 50-81-7), β -aminopropionitrile (CAS # 2079-89-2), sodium acetate (CAS # 6131-90-4), sodium arsenate (CAS # 10048-95-0), and copper sulfate (CAS # 7758-99-8). The chemicals were purchased from Sigma Chemical Co. (St. Louis, MO) in bulk quantities from the same lot. One technician individually weighed the compounds using a calibrated analytical balance. The amount was then placed in 100 ml serum vials (Fisher) and capped with rubber septa and sealed with aluminum lids. Each vial contained enough material to prepare test solutions for each 24-h interval of the experiment; four vials were needed for each test. The chemicals were sent to a coordinator who coded the samples and then shipped enough chemical to each laboratory for three definitive concentration-response experiments. Instructions shipped with each chemical included: 1) the stock concentration, 2) concentrations from which to take samples for chemical analysis, and 3) a Material Safety Data Sheet, available for use in an emergency, sealed in an envelope to prevent disclosure of the identity of the substance.

Data Analysis

Probit analysis, using the method of Litchfield-Wilcoxon ²³ was used to determine the 96-h LC50 (median lethal concentration), 96-h EC50 (concentration inducing malformations in 50% of the surviving embryos), and 95% percent confidence intervals. When the homogeneity test failed, either the trimmed Spearman-Karber ²⁴ or EPA probit method was used instead of the Litchfield-Wilcoxon probit analysis. Teratogenic hazard was determined using a Teratogenic Index [TI=LC50/EC50 (malformation)]. Head-tail length (growth) was measured using an IBM-compatible computer equipped with digitizing software (eg. Jandel Scientific®, Corte Madera, CA). For each test, the Minimum Concentration to Inhibit Growth (MCIG) was calculated using the t-Test for grouped observations ($p<0.05$).

The coefficient of variation (CV) values for the 96-h LC50, EC50, TI and MCIG was calculated according to Steel and Torrie²⁵. Analysis of the data was also carried out using ASTM E691-87 (Interlaboratory Study) software and methods²⁶.

BODY

RESULTS

Monosodium Glutamate

Three complete concentration-response tests (replicates) were performed by each laboratory. The 96-h LC50, 96-h EC50, and MCIG were determined for each replicate. All laboratories reported data except for Laboratory 3 which had difficulty in obtaining data for this compound. Monosodium glutamate often permitted the growth of microorganisms which increased variability. Intralaboratory CV values for the 96-h LC50 ranged from 2.9 to 54% and individual LC50 values ranged from a low of 0.48 to a high of 16.44 mg/ml (Table 1). The 96-h EC50 intralaboratory CV values ranged from 7.4 to 84.3%. Individual EC50 values ranged from 0.08 to 13.96 mg/ml and mean TI values ranged from 1.2 to 15.4, but the intralaboratory CV values for the mean TI were from 8.2 to 449.2%. The intralaboratory CVs for the MCIG ranged from 0 to 117.9% and the mean MCIG values ranged from 0.38 to 7.67 mg/ml.

Fig. 1 shows a graphic analysis using ASTM methodology²⁶. Intralaboratory variability (repeatability) was given by the k values (Fig. 1A). The k value is the intralaboratory consistency statistic found by dividing the cell standard deviation for one laboratory by the repeatability standard deviation of the material²⁶. The repeatability standard is calculated by dividing the sum of the squared cell standard deviations by the number of laboratories. The ASTM software²⁶ generates a value for the limits of variability which is represented by a solid horizontal line(s) in each figure. Only values above the solid horizontal line on the figure were considered to have deviated excessively and only Laboratory 1 showed excessively high variation for the MCIG. The h values presented in Fig. 1B shows the interlaboratory variation (reliability). The h value is the interlaboratory consistency statistic found by dividing the cell deviation (i.e., the deviation of the cell average from the average of the cell averages) divided by the standard deviation of the cell averages²⁶. All interlaboratory measurements were well within acceptable limits (solid horizontal lines on Fig. 1B).

Ascorbic Acid

Laboratory 6 experienced a pH problem with this compound and elected to discontinue testing since its participation was voluntary and uncompensated at this time. Intralaboratory CV values for the 96-h LC50 ranged from 4.6 to 39.5% and individual LC50 values ranged from a low of 1.06 to a high of 16.88 mg/ml (Table 2). The 96-h

EC50 CV values ranged from 2.4 to 14.1%. Individual EC50 values ranged from 0.70 to 13.68 mg/ml and mean TI values ranged from 1.0 to 2.5, while the intralaboratory CV values for the mean TI were from 1.5 to 31.5%. The intralaboratory CVs for the MCIG ranged from 5.7 to 35.4% and the mean MCIG values ranged from 0.37 to 14.67 mg/ml.

Figure 2 shows the ASTM graphic analysis method for ascorbic acid²⁶. Intralaboratory variability was given by the k values (Fig. 2A). None of the values were above the solid horizontal line and were not considered to have deviated excessively. The h values presented in Fig. 2B show the interlaboratory variation. In this case, none of the laboratories exceeded the limits of allowable variation (solid horizontal line) ²⁶.

β-Aminopropionitrile

Intralaboratory CV values for the 96-h LC50 ranged from 7.0 to 112.8% and individual LC50 values ranged from a low of 2.9 to a high of 105.0 mg/L (Table 3). The 96-h EC50 CV values ranged from 5.2 to 66.2%. Individual EC50 values ranged from 0.04 to 2.22 mg/L and mean TI values ranged from 4.5 to 1221.0, while the intralaboratory CV values for the mean TI were from 12.2 to 213.6%. The intralaboratory CVs for the MCIG ranged from 14.1 to 86.9% and the mean MCIG values ranged from 0.08 to 11.73 mg/L.

Fig. 3 shows the ASTM graphic analysis method for β-Aminopropionitrile. Intralaboratory variability indicated by the k values in Fig 3A showed excess variation for the TI (Laboratory 1), the MCIG (Laboratory 2) and the EC50 (malformation) (Laboratory 4). The h values presented in Fig. 3B shows the interlaboratory variation. Laboratory 1 exceeded the ASTM variation limit ²⁶ for the TI while Laboratory 2 exceeded the limit for the MCIG.

Sodium Acetate

Intralaboratory CV values for the 96-h LC50 ranged from 2.2 to 19.5% and individual LC50 values ranged from a low of 5.36 to a high of 12.32 mg/ml (Table 4). The 96-h EC50 CV values ranged from 4.8 to 33.8%. Individual EC50 values ranged from 1.08 to 9.23 mg/ml and mean TI values ranged from 0.9 to 7.5, while the intralaboratory CV values for the mean TI were from 3.3 to 44.8%. The intralaboratory CVs for the MCIG ranged from 0 to 84.1% and the mean MCIG values ranged from 0.5 to 9.33 mg/ml.

Fig. 4 shows the ASTM graphic analysis method for sodium acetate ²⁶. Intralaboratory variability indicated by the k values showed that variation was acceptable

(Fig 4A). The h values presented in Fig 4B show that none of the laboratories had variation exceeding the ASTM variation limits (solid horizontal lines) ²⁶.

Sodium Arsenate

Laboratory 4 withdrew from the project after losing critical staff. Laboratory 6 provided the data for the six laboratory minimum required by the ASTM software. Intralaboratory CV values for the 96-h LC50 ranged from 6.6 to only 22.0 % and individual LC50 values ranged from a low of 1.53 to a high of 3.65 mg/ml (Table 5). The 96-h EC50 CV values ranged from 9.9 to 38.1%. Individual EC50 values ranged from 0.21 to 2.01 mg/ml and mean TI values ranged from 1.4 to 7.0, while the intralaboratory CV values for the mean TI were from 9.8 to 47.3%. The intralaboratory CVs for the MCIG ranged from 0 to 82.7% and the mean MCIG values ranged from 0.47 to 1.5 mg/ml.

Fig. 5 shows the ASTM graphic analysis method for sodium arsenate ²⁶. Intralaboratory variability indicated by the k values showed that variation was acceptable (Fig 5A). The h values presented in Fig 5B show that none of the laboratories had variation exceeding the ASTM variation limits (solid horizontal lines) although the MCIG for laboratory 7 was on the limit line.

Copper Sulfate

Intralaboratory CV values for the 96-h LC50 ranged from 5.1 to 32.8% and individual LC50 values ranged from a low of 0.29 to a high of 2.74 mg/L (Table 6). The 96-h EC50 CV values ranged from 2.7 to 74.5%. Individual EC50 values ranged from 0.08 to 3.95 mg/L and mean TI values ranged from 0.8 to 5.6, while the intralaboratory CV values for the mean TI were from 3.4 to 57.5%. The intralaboratory CVs for the MCIG ranged from 12.9 to 84.1% and the mean MCIG values ranged from 0.04 to 0.92 mg/L.

The ASTM graphic analysis method for copper sulfate is shown in Fig. 6 ²⁶. Intralaboratory variability indicated by the k values showed that variation was generally acceptable except for the EC50 (malformation) for laboratory 5 which exceeded the limit considerably (Fig 6A). The h values presented in Fig 6B show that none of the laboratories had variation exceeding the ASTM variation limits (solid horizontal lines) ²⁶.

Interlaboratory Variability as Determined by CV Values

Table 7. shows interlaboratory summary results for all six test compounds. CV values above 75% were considered somewhat high while values above 150% were considered excessive based on bioassay results reported by Parkhurst et al. 27. Interlaboratory CV values were very good except for monosodium glutamate (84.5% and β -aminopropionitrile (108.2%). For the EC50 (malformation), three of the six test compounds produced somewhat high results (monosodium glutamate- 134.9%, β -aminopropionitrile-85.1% and copper sulfate- 111.4%). Except for ascorbic acid, most of the values for the TI were either high or excessive. CV values for the MCIG were better with values for ascorbic acid, sodium acetate and sodium arsenate better within reasonable limits (Table 7).

CONCLUSIONS

Monosodium Glutamate

Monosodium glutamate has not been previously tested in FETAX. Previous reports suggest little developmental toxicity attributable to monosodium glutamate administered as a brief treatment in mice²⁸. Monosodium glutamate was typical of compounds with low developmental toxicity tested in the FETAX ILS in that overall variation was fairly low for this study. This was similar to Phase II findings²². The chief cause of the observed interlaboratory variation was Laboratory 5 which reported very low LC50 (Laboratory mean 0.56 mg/ml) and EC50 (malformation) (Laboratory mean 0.24 mg/ml) values while Laboratory 6 reported LC50 and EC50 (malformation) values of 15.96 and 12.96 mg/ml, respectively. The interlaboratory means were 6.52 and 3.43, respectively. A greater problem was that the mean laboratory TI ranged from a low of 1.19 which indicated little teratogenic hazard to a high of 15.4 which suggested strong teratogenic hazard. The intralaboratory variation was quite low suggesting other factors were affecting interlaboratory variability. There were some reports of bacterial contamination when this compound was tested. Some test compounds permit or even support the growth of microbial organisms which can then affect test results. Antibiotic treatments were avoided in this study but may be used as long as suitable controls are used to account for possible interactions¹⁹. Laboratory 3 was unable to successfully test this compound, perhaps because of contamination.

Ascorbic Acid

Previous FETAX tests with ascorbic acid indicated LC50, EC50 (malformation) and MCIG values of 19.7, 12.13 and 10 mg/ml. The TI was 1.62. Table 7 shows that the six laboratories that tested ascorbic acid in the ILS study had interlaboratory mean values of 10.70, 7.61 and 6.28 mg/ml, respectively, for the LC50, EC50 (malformation) and MCIG. The TI was 1.4. The lot numbers of ascorbic acid were different between the trials and the ILS numbers were reduced by Laboratory 4 who reported very low values for the LC50, EC50 (malformation) and MCIG endpoints (Table 2). The reason for these low values is unknown. Interestingly, the resultant TI was not greatly different from the other laboratories. On the whole, results were judged to be very good. Ascorbic acid was considered to pose little developmental hazard and this was confirmed in this study. The relationship of these tests results to the mammalian literature has already been discussed²⁹.

β -Aminopropionitrile

This chemical has not been previously tested in FETAX although the mammalian literature indicates that it is a lathrygen in rats (30,31). Results from this study seem to confirm this as a TI value of 55.3 was reported (Table 7). The LC50, EC50 (malformation) and MCIG values were 34.76, 0.63 and 2.11 mg/L, respectively. Once again, there was a fairly significant range observed between laboratories, especially for

the LC50 (Table 3). For example, Laboratories 1 and 7 reported values near 80 mg/L, while Laboratories 3, 5 and 6 reported values at least 10 fold lower. Interestingly, the EC50 (malformation) endpoint was less variable and these differences caused great variability in the TI as seen in the high interlaboratory CV value of 991.6% (Table 7).

Sodium Acetate

Sodium acetate has been previously tested in FETAX but the exposure period was for 5 days instead of the standard four and, therefore, the data was not directly comparable to this study. The five day exposure data was 4.2, 3.3 and 2.5 mg/ml for the LC50, EC50 (malformation) and MCIG, respectively (32). The TI was 1.3. Predictably, the results of the four day exposure study was higher at 9.26, 5.00 and 4.48 mg/ml for the LC50, EC50 (malformation) and MCIG, respectively (Table 7). The TI was 1.9 which indicated a modest degree of teratogenic hazard. The interlaboratory variability for the LC50 and EC50 (malformation) was within acceptable ranges but the CV for the TI was high (Table 7). The CV values for TIs tended to be higher because the LC50 is divided by the EC50 (malformation) thereby causing a multiplication effect on the resultant CV values for the TI.

Sodium Arsenate

Sodium arsenate has not been previously tested in FETAX. This study found that the LC50, EC50 (malformation) and MCIG were 2.30, 0.97 and 1.11 mg/ml, respectively (Table 7). The TI was 2.4 and indicated modest teratogenic activity. Malformations were moderate to severe in nature. Interlaboratory variability was generally low with this compound except for the TI again (Table 7). Between laboratories the EC50 (malformation) proved considerably more variable than the LC50.

Mammalian studies indicated some teratogenicity in rodents such as mice ³³, hamsters ³⁴ and rats ³⁵. The FETAX studies reported here indicate less developmental hazard due to arsenate than the mammalian studies. Limited human studies of accidental exposures did not detect human developmental toxicity.

Copper Sulfate

Fort et al., ³⁶ tested copper sulfate in FETAX. They found that the mean LC50, EC50 (malformation) and MCIG were 0.94, 0.81 and 0.75 mg/L, respectively. The TI was only 1.15 which indicated little developmental hazard. Data from this study showed excellent correspondence to the historical data as the values were 1.23, 0.83 and 0.35 mg/L and the TI was 1.5 (Table 7). Most of the laboratories reported consistent results but the mean laboratory TI values ranged from 0.8 to 5.58.

The effects of copper on mammalian development has been reported by Everson et al. ³⁷ and Kalter ³⁸. Some effects on neural tissue are reported but copper is not

considered a strong teratogen. The main effects are embryotoxicity and FETAX confirms this.

Endpoint Variability

Table 8 shows that the LC50 was the least variable endpoint based on comparison of CV values in five of six tests. The MCIG was the most variable in three of six tests. These results were not surprising because the concentration ranges were designed to define the LC50 and EC50 values and the LC50 is the more objective endpoint compared to the EC50.

Sources of Variation

In analyzing the results of this study and Phases I and II, it became apparent that at least one technician typically judged malformations more harshly than any other throughout the four years of testing. This pattern was not repeated for the LC50 estimation. This occurred even though there was extensive technician training and practical experience. Another technician, reported high EC50 (malformation) and LC50 values in two of the three phases and was second highest in the third Phase. This may have been due to problems in sample dilution because the LC50 tracked the EC50 in this Laboratory.

The chief difference between Phase II which yielded outstanding results and Phase III which provide good but far more variable results was the ability of participants to choose their own test concentrations. Analysis of the concentrations selected by participants revealed that some selected very narrow ranges of close steps between concentrations while other selected far broader ranges. Different toxicants yielded concentration-response curves with different slopes and this seemed to have an effect on results depending on the concentrations selected. In an effort to standardize the selection of test concentration yet still allow each laboratory to select their own concentration the following protocol was written. While complicating the procedure, it seems to be yielding better results at this stage of the study. Further tests are now being conducted to determine whether variability is reduced.

Recommended Protocol for Testing New Compounds Using FETAX

The following sequence is now being followed when testing a new chemical for a three way round robin test program currently in progress. This procedure will help to standardize initial-range finding experiments and help reduce some of the variability found in the present study. The metabolic activation system (MAS) is composed of Aroclor 1254-induced rat liver microsomes and NADPH generator system⁵.

Initial Range-Finding Test (No MAS used)

The range-finding tests consists of a series of at least seven concentrations that differ by a factor of ten. This is adequate to delineate the concentration range to establish the 96-h LC50 and EC50 (malformation)¹⁹. Growth inhibition data are not collected from range-finding tests. Investigators are encouraged to add concentrations whenever possible to find the best approximation of the LC50 and EC50. Once the data are collected from the range-finding tests, the LC50 and EC50 should be estimated using probit analysis, trimmed Spearman-Karber or the two point method. The two point method is used when regular statistical programs fail to generate any useful data. It involves drawing a line from the low response to the high response and the 50% response concentration is estimated from where the line intercepts the 50% response point. If the data allows the Litchfield-Wilcoxon test or trimmed Spearman-Karber methods to be used, then Litchfield-Wilcoxon is used when the data passes homogeneity (Shapiro-Wilks) and normality (Chi Square) tests and trimmed Spearman-Karber is used when the data fails these tests. Range-finding tests may bypass the homogeneity requirement here but not in replicate tests discussed below. The Toxstat version 3.0 program from the Univ. of Wyoming is used to make homogeneity and normality assessments (Fish Physiology and Toxicology Laboratory, Dept. of Zoology and Physiology, P.O. Box 3166, University of Wyoming, Laramie, WY, 82071).

Estimation of the unactivated LC50 and EC50 (No MAS used).

This is a one or two step process depending on the nature of the test compound and the results of the first test in this series. The test is performed using the sliding scale of concentrations presented in Table 9. The table presents values from 0.001 to 100. From 0.001 to 0.1 the steps are 0.005. From 0.1 to 1, the steps are 0.05 and from 1 to 10, the steps (differences between concentrations) are 0.5. Lastly, the steps are 5.0 from concentration 10 to 100. The value on the table closest to the LC50 is found. Then three values are chosen immediately below and three values immediately above this point. The same is performed for the EC50. A test is performed and the data collected. The LC50 and EC50 values with confidence limits are then generated. After the successful conclusion of this test, a determination is made as to whether a metabolic activation system should be used in subsequent testing. From the data obtained in above, calculate the LC5, LC16, LC50, LC84 and LC95 and the EC5, EC16, EC50, EC84 and EC95. By determining the above values, the concentrations to be tested in the definitive tests below are established and the slope of the concentration-response curves are accounted for. Additional concentrations may be added to ensure obtaining an LC50 and EC50, but the same concentrations must be used for each replicate (definitive) test.

Replicate (Definitive) Tests (MAS used).

Three tests are then performed with and without MAS each with a separate clutch of embryos. The ten concentrations determined above are used, with and without MAS. The unactivated experiments should yield good LC50 and EC50 values. If they do not,

the tests must be repeated. If the MAS test yields LC50 and EC50 values on the first test by an acceptable probit analysis method, the results should be reported. If the addition of MAS alters toxicity such that LC50 and EC50 values can not be determined, then more range finding is done similar to that for the unactivated tests above until the proper concentration range for MAS is defined. Tests with and without MAS are then performed on the same clutch of embryos until three replicate tests are obtained.

Controls to be performed for the initial range finding step and the estimation of the unactivated LC50 and EC50 (No MAS used).

Note: dishes are plastic unless otherwise noted.

- 1.) FETAX solution. Use four dishes of 20 embryos each.
- 2.) FETAX + cosolvent (if required). Use four dishes of 20 embryos each.

Controls to be performed with each replicate (definitive) test:

- 1.) FETAX/Antibiotics solution. Use four dishes of 20 embryos each.
FETAX/Antibiotics solution is the diluent for all test concentrations.
- 2.) FETAX/Antibiotics solution + cosolvents if one is used. Use four dishes of 20 embryos each. Refer to ASTM guide¹⁹ on when to use for data comparison.
If a cosolvent is required, then the following controls (except the cyclophosphamide controls) must contain the cosolvent.
- 3.) Four mg/ml cyclophosphamide positive controls with MAS. Two dishes of 20 embryos each.
- 4.) Microsomes alone (no generator) + four mg/ml cyclophosphamide. Two dishes of 20 embryos each.
- 5.) Microsomes alone (no generator) + test compound at the LC50 concentration.
2 dishes of 20 embryos each.
- 6.) Generator system only plus test compound at the LC50 concentration. Two dishes of 20 embryos each.
- 7.) MAS alone. Two dishes of 20 embryos each.

Final Conclusions

The results of this study indicated that FETAX is still a repeatable and reliable assay but that variation due to technician bias in terms of judging malformations and in selecting test concentrations must be controlled. Bacterial contamination still causes variability but this can be detected by noting sample turbidity and variation in replicates of the same concentration. Antibiotics can be used to control contamination but care must be exercised to account for possible interaction. Variability is generally lower for nonteratogens than strong teratogens.

ACKNOWLEDGMENT

This work was supported by U.S. Army Medical Research and Development Command Contract #DAMD17-91-C-1048. The views, opinions, and/or findings contained in this article are those of the authors and should not be construed as official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations. Research was conducted in compliance with the Animal Welfare Act, and other Federal Statutes and regulations relating to animals and experiments involving animals and adheres to principles stated in the *Guide for Care and Use of Laboratory Animals*, NIH Publication No. 86-23, 1985 edition).

REFERENCES

1. J.N. Dumont, T.W. Schultz, M.V. Buchanan and G.L. Kao, Frog embryo teratogenesis assay: *Xenopus* (FETAX) - A short-term assay applicable to complex environmental mixtures. In *Symposium on the Application of Short-Term Bioassays in the Analysis of Complex Environmental Mixtures III*, ed. by M. D. Waters, S.S. Sandhu, J. Lewtas, L. Claxton, N. Chernoff, and S. Nesnow, pp. 393-405. Plenum Press, New York (1983).
2. J. A. Bantle and D. A. Dawson, Uninduced rat liver microsomes as an *in vitro* metabolic activation system for the Frog Embryo Teratogenesis Assay-*Xenopus* (FETAX). In *Aquatic Toxicology and Hazard Assessment*, Vol. 10, ed. by W.J. Adams, G. A. Chapman and W. G. Landis, ASTM STP 971, pp. 316-326. American Society for Testing and Materials, Philadelphia (1988).
3. J. A. Bantle, D. J. Fort and B. L. James, Identification of developmental toxicants using the Frog Embryo Teratogenesis Assay-*Xenopus* (FETAX). *Hydrobiol* **188/189**, 577-585 (1989).

4. D. A. Dawson, D. J. Fort, G. L. Smith, D. L. Newell and J. A. Bantle, Evaluation of the developmental toxicity of nicotine and cotinine with Frog Embryo Teratogenesis Assay: *Xenopus* (FETAX). *Teratogen. Carcinogen. Mutagen.* **8**, 329-338 (1988).
5. D. J. Fort, D. A. Dawson and J. A. Bantle, Development of a metabolic activation system for the Frog Embryo Teratogenesis Assay: *Xenopus* (FETAX). *Teratogen. Carcinogen. Mutagen.* **8**, 251-263 (1988).
6. D. J. Fort and J. A. Bantle, Use of Frog Embryo Teratogenesis Assay - *Xenopus* and an exogenous metabolic activation system to evaluate the developmental toxicity of diphenylhydantoin. *Fund. Appl. Toxicol.* **14**, 720-733 (1990a).
7. D. J. Fort and J. A. Bantle, Analysis of the mechanism of isoniazid-induced developmental toxicity with the Frog Embryo Teratogenesis Assay - *Xenopus* (FETAX), *Teratogen. Carcinog. Mutagen.* **10**, 463-476 (1990b).

8. D. J. Fort, J. R. Rayburn, D. J. DeYoung and J. A. Bantle, Assessing the efficacy of an Aroclor 1254-induced exogenous metabolic activation system for FETAX. *Drug Chem. Toxicol.* **14**(1&2), 143-160 (1991).
9. D. J. Fort, J. R. Rayburn and J. A. Bantle, Evaluation of acetaminophen-induced developmental toxicity using FETAX, *Drug Chem. Toxicol.* **15**, 329-350 (1992).
10. J. A. Bantle, FETAX - A developmental toxicity assay using frog embryos. In: *Fundamentals of Aquatic Toxicology*, 2nd ed.. ed. by G. M. Rand and S. R. Petrocelli, Hemisphere Publishing, New York (1993), In Press.
11. J. N. Dumont and T. W. Schultz, Effects of coal-gasification sour water on *Xenopus laevis* embryos. *J. Environ. Sci. Health A***15**, 127-138 (1980).
12. D. A. Dawson, C. A. McCormick and J. A. Bantle, Detection of teratogenic substances in acidic mine water samples using the Frog Embryo Teratogenesis Assay-*Xenopus*(FETAX). *J. Appl. Toxicol.* **5**(4), 234-244 (1985).
13. D. A. Dawson, E. Stebler, S. A. Burks and J. A. Bantle, Teratogenicity of metal contaminated sediment extracts to frog (*Xenopus laevis*) and fathead

- minnow (*Pimephales promelas*) embryos. *Environ. Toxicol. Chem.* **7**(1), 27-34 (1988).
14. J. A. Bantle, D. J. Fort and D. A. Dawson, Bridging the gap from short-term teratogenesis assays to human health hazard assessment by understanding common modes of teratogenic action. In *Aquatic Toxicology and Hazard Assessment*, Vol. 12, ed. by W. G. Landis, W. H. Van der Schalie, ASTM STP 1096, pp. 46-58. American Society for Testing and Materials, Philadelphia, (1989).
15. D. B. Wake and H. J. Morowitz, Declining amphibian populations-a global phenomenon, report of a workshop sponsored by the Board on Biology, National Research Council, held at Irvine, CA, February 19-20, (1990).
16. D. B. Wake, Declining amphibian populations. *Science* **253**, 860 (1991).
17. J. H. K. Pechman, D. E. Scott, R. D. Semlitsch, J. P. Caldwell, L. J. Vitt and J. W. Gibbons, Declining amphibian populations: The problem of separating human impacts from natural fluctuations. *Science* **253**, 892-895 (1991).

18. Science Briefings, New task force on declining amphibians. *Science* **253**, 509 (1991).
19. American Society for Testing and Materials, *New standard guide for conducting the Frog Embryo Teratogenesis Assay-Xenopus (FETAX)*, ASTM E1439-91, Special Publications Philadelphia, (1991).
20. J. A. Bantle, J. N. Dumont, R. Finch and G. Linder, *Atlas of Abnormalities: A Guide for the Performance of FETAX*. Oklahoma State Publications Department (1991).
21. J.A. Bantle, D.T. Burton, D.A. Dawson, J.N. Dumont, R.A. Finch, D.J. Fort, G. Linder, R.A. Rayburn, D. Buchwalter, M.A. Maurice, and S.D. Turley. Initial interlaboratory validation study of FETAX: Phase I testing. *Journal of Applied Toxicology*, 14(3),213-223 (1994).
22. J.A. Bantle, D.T. Burton, D.A. Dawson, J.N. Dumont, R.A. Finch, D.J. Fort, G. Linder, R.A. Rayburn, D. Buchwalter, A.M. Gaudet-Hull, M.A. Maurice and S.D. Turley. Initial interlaboratory validation study of FETAX: Phase II testing. *Environmental Toxicology and Chemistry*, 13(10), 1629-1637 (1994).

23. R. J. Tallarida and R. B. Murray, *Manual of Pharmacologic Calculations with Computer Programs*. 2nd Edn, Springer-Verlag, New York (1987).
24. M. A. Hamilton, R. C. Russo and R. V. Thurston, Trimmed Spearman-Karber method for estimating median lethal concentrations in toxicity bioassays. *Environ. Sci. Tech.* **11**(7), 714-718 (1977).
25. G. D. Steel and J. H. Torrie, *Principles and Procedures of Statistics: A Biometrical Approach*. 2nd Edn, ed. by C. Napier and J.W. Maisel. McGraw-Hall Book Company, New York (1980).
26. American Society for Testing and Materials, *Standard Practice for Conducting an Interlaboratory Study to Determine the Precision of a Test Method*, ASTM E691-87, Annual book of ASTM standards, **14.02**, 419-438, Philadelphia (1988).

27. B. R. Parkhurst, W. Warren-Hicks and L. E. Noel, Performance characteristics of effluent toxicity tests: Summarization and evaluation of data. *Envir. Toxicol. Chem.* **2**, 771-791 (1992).
28. U. Murakami and M. Inouye, Brain lesions in the mouse fetus caused by maternal administration of monosodium glutamate. *Congenital Anomalies* **11**:171-177, 1971.
29. D.J. DeYoung, J.A. Bantle and D.J. Fort, Developmental toxicity of ascorbic acid, sodium selenate, coumarin, serotonin, and 13-cis retinoic acid tested in FETAX. *Drug and Chemical Toxicology*, **14**,127-141 (1991).
30. F.W. Stamler, Reproduction in rats fed lathrus peas or aminonitriles. *Proc. Soc. Exp. Biol. Med.* **90**: 294-298 (1955).

31. A.J. Steffak, A.C. Verrusio and C.A. Watkins, Cleft palate in rodents after maternal treatment with various lathyrogenic agents. *Teratology* 5: 33-40 (1972).
32. D.J Deyoung, M.A. Hull and J.A. Bantle, Differences in sensitivity to developmental toxicants as seen in *Xenopus* and *Pimephales* embryos. *Bulletin of Environmental Contamination and Toxicology, Submitted (1995)*.
33. M.N. Baxley, R.D. Hood, G.C. Vedel, W.P. Harrison, and G.M. Szezech, Prenatal toxicity of orally administered sodium arsenate in Mice. *Bulletin of Environmental Contamination and Toxicology*, 26:749-756, (1981).
34. A.R. Beaudoin, Teratogenicity of sodium arsenate in rats. *Teratology* 10: 153-158, (1974).
35. V.H. Ferm, A. Saxon and B.M. Smith, The teratogenic profile of sodium arsenate in the golden hamster. *Arch. Environ. Health* 22: 557-560, (1971).

36. D.J. Fort, E.L. Stover and J.A. Bantle, Integrated ecological hazard assessment of waste site soil extracts using FETAX and short-term fathead minnow teratogenesis assay, *Environmental Toxicology and Risk Assessment: Transboundary Issues in Pollution- Air, Surface, and Groundwater*: Fourth Volume, ASTM STP 1262, T. W. LaPoint, E.E. Little, F.T. Price, Eds., American society for Testing and Materials, Philadelphia, (1995)- In Press.
37. R.L. Brinster and P.C. Cross, Effect of copper on the preimplantation mouse embryo. *Nature* 238, 398-399, (1972).
38. F.J. DiCarlo, Syndromes of cardiovascular malformations induced by copper citrate in hamsters. *Teratology*. 21, 89-101, (1980).

Table 1. Results of the Interlaboratory Study for Monosodium Glutamate.

Laboratory	Replicate (n)	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV1 (%)	EC50 (mg/ml)	Mean EC50 (mg/ml)	EC50 CV (%)	Mean TI TI ² (%)	TI CV (%)	MCIG ³ (mg/ml)	Mean MCIG (mg/ml)	MCIG CV (%)
1	1	4.07	4.07	0.46	0.51	26.6	15.4	29.3	1.3	10.0	3.92	110.1
	2	5.64	7.80	54.0	0.37	0.69	—	—	—	—	—	—
	3	13.68	—	—	—	—	—	—	—	—	—	—
2	1	8.02	—	—	1.32	—	—	—	—	—	—	—
	2	4.15	7.56	34.6	0.86	1.02	20.5	7.4	41.1	0.3	4.0	117.9
	3	10.50	—	—	0.89	—	—	—	—	—	—	—
3	1	—	—	—	—	—	—	—	—	—	—	—
	2	—	—	—	—	—	—	—	—	—	—	—
	3	—	—	—	—	—	—	—	—	—	—	—
4	1	1.35	—	—	0.08	—	—	—	—	—	—	—
	2	1.07	1.07	21.4	1.39	0.65	84.3	1.7	449.2	—	—	—
	3	0.79	—	—	0.48	—	—	—	—	—	—	—
5	1	0.66	—	—	0.26	—	—	—	—	—	—	—
	2	0.48	0.56	13.4	0.21	0.24	9.7	2.3	8.2	0.4	0.42	5.7
	3	0.54	—	—	0.26	—	—	—	—	—	—	—
6	1	16.10	—	—	11.66	—	—	—	—	—	—	—
	2	16.44	15.96	2.9	13.96	12.96	7.4	1.2	8.2	7.5	7.5	7.50
	3	15.34	—	—	13.27	—	—	—	—	—	—	—
7	1	5.47	—	—	3.90	—	—	—	—	—	—	—
	2	6.52	6.17	8.0	5.67	5.19	17.7	1.2	11.3	7.0	10.0	22.2
	3	6.51	—	—	5.99	—	—	—	—	—	—	—

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

Table 2. Results of the Interlaboratory Study for Ascorbic Acid.

Laboratory	Replicate (n)	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV1 (%)	EC50 (mg/ml)	Mean EC50 (mg/ml)	EC50 CV (%)	Mean TI2 (%)	TI CV (%)	MCIG ³ (mg/ml)	Mean MCIG (mg/ml)	MCIG CV (%)
1	1	11.04	7.01							5.0		
	2	6.98	10.98	29.5	5.31	6.60	14.1	1.7	16.8	10.0	8.33	28.3
	3	14.91			7.48					10.0		
2	1	15.21	7.00							2.0		
	2	8.85	13.06	22.8	6.69	6.29	12.7	2.1	31.5	2.0	2.67	35.4
	3	15.11			5.17					4.0		
3	1	15.61	11.33							2.0		
	2	16.88	14.83	13.9	11.99	11.60	2.4	1.3	12.9	4.0	3.33	28.3
	3	12.00			11.49					4.0		
4	1	2.78	0.82							0.3		
	2	1.61	1.82	39.5	0.70	0.74	7.6	2.5	31.3	0.3	0.37	25.7
	3	1.06			0.70					0.5		
5	1	9.26	7.24							9.0		
	2	10.21	9.91	4.6	7.69	7.43	2.6	1.3	3.6	8.0	8.33	5.7
	3	10.25			7.35					8.0		
6	1											
	2											
	3											
7	1	12.20	11.94							15.0		
	2	14.26	13.61	7.3	13.47	13.03	6.0	1.0	1.5	16.0	14.67	8.5
	3	14.36				13.68				13.0		

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

Table 3. Results of the Interlaboratory Study for β -aminopropionitrile.

Laboratory	Replicate (n)	Mean LC50 (mg/L)	LC50 CV ¹ (%)	EC50 (mg/L)	Mean EC50 (mg/L)	EC50 CV (%)	Mean TI ² (%)	TI CV (%)	Mean MCIG ³ (mg/L)	MCIG CV (%)
1	1	93.70		0.07					0.08	
	2	45.40	79.37	30.4	0.08	0.07	25.5	1221.0	58.7	0.08
	3	99.00		0.04					0.10	
2	1	9.50		0.021					0.20	
	2	65.40	29.43	86.6	0.55	0.30	57.9	97.9	30.9	25.00
	3	13.40		0.15					10.00	
3	1	5.30		0.28					0.25	
	2	2.90	4.17	23.6	1.28	0.93	49.2	4.5	167.2	0.25
	3	4.30		1.24					0.50	
4	1	12.60		2.22					0.50	
	2	105.00	40.53	112.8	0.63	1.15	66.2	35.4	213.6	0.75
	3	4.00		0.59					0.10	
5	1	5.50		0.67					1.00	
	2	4.70	4.90	8.8	0.66	0.69	5.2	7.1	12.2	2.00
	3	4.50		0.74					1.00	
6	1	7.70		0.21					0.10	
	2	7.40	7.93	7.0	0.22	0.20	11.8	40.1	20.6	0.15
	3	8.70		0.17					0.35	
7	1	60.00		1.10					1.00	
	2	76.00	77.00	18.6	1.20	1.07	11.7	72.2	30.8	0.50
	3	95.00		0.90					0.50	

¹LCV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean EC50 by the Mean LC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

Table 4. Results of the Interlaboratory Study for Sodium Acetate.

Laboratory	Replicate (n)	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV1 (%)	EC50 (mg/ml)	Mean EC50 (mg/ml)	EC50 CV (%)	Mean TI2 (%)	TI CV (%)	MCIG ³ (mg/ml)	Mean MCIG (mg/ml)	MCIG CV (%)
1	1	6.96	6.21	10.6	1.41	2.32	33.8	2.6	44.8	6.0	3.00	72.0
	2	6.30	5.36	5.36	3.37	2.37	33.8	2.6	44.8	6.0	3.00	2.0
	3	5.36										
2	1	11.20	10.96	2.3	1.18	2.01	1.45	27.1	7.5	22.9	0.5	0.50
	2	11.07	10.62	10.62	1.17							
	3	10.62										
3	1	12.00	11.69	4.2	5.12	7.64	7.21	21.7	1.6	27.6	10.0	9.33
	2	11.00	12.06	12.06	8.88							
	3	12.06										
4	1	11.69	9.72	17.4	5.23	6.62	6.12	10.3	1.6	28.3	1.0	1.67
	2	9.92	7.56	7.56	6.52							
	3	7.56										
5	1	7.76	7.05	13.3	7.92	8.26	7.50	11.3	0.9	3.3	8.0	7.67
	2	7.67	5.72	5.72	6.31							
	3	5.72										
6	1	12.32	11.68	11.99	2.2	8.77	9.23	8.74	4.8	1.4	5.9	5.0
	2	11.68	11.96	11.96	8.21							
	3	11.96										
7	1	8.91	7.31	7.23	19.5	2.30	1.64	30.6	4.4	24.7	0.50	5.67
	2	7.31	5.46	5.46	1.08							
	3	5.46										

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

Table 5. Results of the Interlaboratory Study for Sodium Arsenite.

Laboratory	Replicate (n)	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV (%)	EC50 (mg/ml)	Mean EC50 (mg/ml)	EC50 CV (%)	Mean TI CV (%)	TI CV (%)	MCIG ³ (mg/ml)	Mean MCIG (mg/ml)	MCIG CV (%)
1	1	2.53	2.09	14.9	0.58	0.40	38.1	5.3	39.3	0.30	0.10	0.47
	2	1.87	1.87	0.21	0.21	0.40	0.43	32.4	7.0	38.7	1.00	0.0
	3	1.87		0.40			0.25			1.00	1.00	0.0
2	1	2.65			0.59					1.00		
	2	3.65	3.03	14.6	0.45	0.43						
	3	2.79		0.25								
30	1	1.53			1.27					1.50		
	2	2.03	2.07	22.0	1.91	1.40				1.00		
	3	2.64		1.01						1.00		
4	1									1.17		
	2										20.2	
	3											
5	1	1.77			1.23					1.50		
	2	2.47	2.07	14.1	1.54	1.28				1.50		
	3	1.98		1.08						1.50		
6	1	2.89			2.01					1.00		
	2	2.50	2.45	15.4	1.64	1.76				1.00		
	3	1.97		1.64						1.00		
7	1	1.89			0.39					1.00		
	2	2.16	2.08	6.6	0.56	0.52				2.50		
	3	2.20		0.62						1.00		

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

Table 6. Results of the Interlaboratory Study for Copper Sulfate.

Laboratory	Replicate (n)	LC50 (mg/L)	Mean LC50 (mg/L)	LC50 CV1 (%)	EC50 (mg/L)	Mean EC50 (mg/L)	EC50 CV (%)	Mean TI ² CV (%)	TI CV (%)	MCIG ³ (mg/L)	Mean MCIG (mg/L)	MCIG CV (%)
1	1	1.00	1.56	32.8	0.32	0.21	0.28	17.7	5.6	33.4	0.70	22.0
	2	1.45	2.24		0.31						0.40	0.57
	3										0.60	
2	1	0.63	0.79	0.77	13.9	0.15	0.24	0.20	19.0	3.8	10.5	0.20
	2										0.40	0.27
	3	0.89				0.22					0.20	35.4
3	1	0.93	0.66	0.85	15.6	0.33	0.45	25.8	1.9	8.0	0.05	51.4
	2										0.05	
	3	0.95				0.56					0.01	
4	1	0.29	0.50	0.46	26.8	0.08	0.11	0.20	74.5	2.3	57.5	0.01
	2										0.03	0.04
	3	0.58				0.41					0.08	
5	1	1.70	2.07	2.17	19.8	2.47	1.61	2.68	36.1	0.8	34.6	0.75
	2										1.00	
	3	2.74				3.95					1.00	0.92
6	1	0.88	0.82	0.88	5.1	0.81	0.76	0.79	2.7	1.1	3.4	0.25
	2										0.20	35.4
	3	0.93				0.80					0.10	
7	1	1.49	2.09	1.93	16.2	1.48	0.89	1.10	24.7	1.8	36.6	0.50
	2										0.30	0.43
	3	2.20				0.92					0.50	21.8

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

Table 7. Results of the Interlaboratory Study.

Chemical	Mean LC50 ¹	LC50 ²	Mean EC50 ¹	EC50 ²	Mean TI ³	TI CV (%)	Mean MCIG ⁴	MCIG CV (%)
Monosodium Glutamate ⁵	6.52	84.5	3.43	134.9	1.9	318.3	3.75	98.8
Ascorbic Acid ⁵	10.70	44.5	7.61	53.0	1.4	46.3	6.28	78.1
β -Aminopropionitrile	34.76	108.2	0.63	85.1	55.3	991.6	2.11	261.1
Sodium Acetate	9.26	26.2	5.00	59.4	1.9	135.8	4.48	74.4
Sodium Arsenate	2.30	21.7	0.97	59.7	2.4	120.8	1.11	44.5
Copper Sulfate	1.23	54.6	0.83	111.4	1.5	127.3	0.35	87.2

¹All values are mg/ml except β -aminopropionitrile and copper sulfate which are mg/L.

²CV=Coefficient of Variation.

³TI=Mean Teratogenic Index calculated by dividing the mean LC50 by the mean EC50 for all laboratories.

⁴MCIG=Minimum Concentration to Inhibit Growth.

⁵Only six laboratories reported.

Table 8. Comparison of the Phase III Interlaboratory Variability of FETAX Endpoints Based on CV Values.

Compound	Least Variable	Mid-Range	Most Variable
Monosodium Glutamate	LC50	MCIG	EC50
Ascorbic Acid	LC50	EC50	MCIG
β -Aminopropionitrile	EC50	LC50	MCIG
Sodium Acetate	LC50	EC50	MCIG
Sodium Arsenate	LC50	MCIG	EC50
Copper Sulfate	LC50	MCIG	EC50

Table 9. Concentration Series to be Used in the First Round of Testing.

Low Series	Medium Series	High Series
0.001	0.10	20
0.005	0.15	25
0.010	0.20	30
0.015	0.25	40
0.020	0.30	45
0.025	0.35	50
0.030	0.40	55
0.035	0.45	60
0.040	0.50	65
0.045	0.55	70
0.050	0.60	75
0.055	0.65	80
0.060	0.70	85
0.065	0.75	90
0.070	0.80	95
0.075	0.85	100
0.080	0.90	
0.085	0.95	
0.090	1.0	
0.095	1.5	
0.10	2.0	
0.15	2.5	
0.20	3.0	
0.25	3.5	
0.30	4.0	
0.35	4.5	
0.40	5.0	
0.45	5.5	
0.50	6.0	
0.55	6.5	
0.60	7.0	
0.65	7.5	
0.70	8.0	
0.75	8.5	
0.80	9.0	
0.85	9.5	
0.90	10	
0.95	15	

FIGURE LEGENDS

Figure 1. Intralaboratory k and interlaboratory h values for monosodium glutamate. (—) acceptable limit of variation. **A.** k values for intralaboratory variation. **B.** h values for interlaboratory variation.

Figure 2. Intralaboratory k and interlaboratory h values for ascorbic acid. (—) acceptable limit of variation. **A.** k values for intralaboratory variation. **B.** h values for interlaboratory variation.



Figure 3. Intralaboratory k and interlaboratory h values for β -aminopropionitrile. (—) acceptable limit of variation. **A.** k values for intralaboratory variation. **B.** h values for interlaboratory variation.

Figure 4. Intralaboratory k and interlaboratory h values for sodium acetate. (—) acceptable limit of variation. **A.** k values for intralaboratory variation. **B.** h values for interlaboratory variation.

Figure 5. Intralaboratory k and interlaboratory h values for sodium arsenate. (—) acceptable limit of variation. **A.** k values for intralaboratory variation. **B.** h values for interlaboratory variation.

Figure 6. Intralaboratory k and interlaboratory h values for copper sulfate. (—) acceptable limit of variation. **A.** k values for intralaboratory variation. **B.** h values for interlaboratory variation.


Figure 1

**Laboratory****Laboratory**


■ LC50 ■ EC50 ■ MCIG ■ TI

p3b

Figure 2




Laboratory



Laboratory


■ LC50 ■ EC50 ■ MCIG ■ TI

Figure 3




p3d

Figure 4




Laboratory



Laboratory


■ LC50 ■ EC50 ■ MCIG ■ TI

Figure 5



■ LC50 ■ EC50 ■ MCIG ■ TI

Figure 6



■ LC50 ■ EC50 ■ MCIG ■ TI

APPENDIX 1

INTERMEDIARY DATA TABLES AND

FIRST PAGES FOR EACH COMPOUND

MONOSODIUM GLUTAMATE

PHASE III COMPOUND A DATA (P3A) MSG

FETAX ILS STUDY; 7 LABS, 8 LEVELS, 3 REPLICATES

PAGE 1

ORIGINAL DATA TABLE WITH MATERIALS

		ORIGINAL DATA TABLE WITH MATERIALS								SUBTOTALS																				
		LC50				EC50				CILO				CIHI				LOCMORT		LOEC/MALF		SUBTOTALS		STD:		CV:				
		LAB	A	B	C	D	E	F	G	H	I	J	K	L	M	N	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10	T11	T12	T13	T14
BANTLE/ RAYBURN	1	4.07	2.88	5.76	0.46	0.30	0.70	10.00	8.8	5.00	0.25	LC50	7.80	4.21	54.0															
	2	5.64	5.06	6.29	0.37	0.24	0.58	1.25	15.2	7.50	0.10	EC50	0.51	0.13	26.6															
	3	13.68	11.49	16.28	0.69	0.54	0.89	0.50	19.8	5.00	0.50	MCIG	3.92	4.31	110.1															
BANTLE/ HULL	1	2	8.02	7.33	8.77	1.32	1.01	1.72	4.00	6.1	8.00	0.25	LC50	7.56	2.61	34.6														
	2	4.15	3.83	4.49	0.86	0.63	1.18	0.25	4.8	6.00	0.75	EC50	1.02	0.21	20.5															
	3	10.50	7.99	13.78	0.89	0.24	3.32	0.25	11.8	12.00	14.00	MCIG	1.50	1.77	117.9															
FINCH	1	3																												
	2																													
	3																													
LINDER/ BUCHWALTER	1	4	1.35	0.00	1.30E+07	0.08	0.00	0.00	16.9	2.00	5.00	LC50	1.07	0.23	21.4															
	2	1.07	0.51	2.25	1.39	0.53	3.66	0.25	0.8	2.00	2.00	EC50	0.65	0.55	84.3															
	3	0.79	0.59	1.06	0.48	0.41	0.58	0.50	1.6	1.00	0.50	MCIG	0.38	0.13	33.3															
BURTON/ TURLEY	1	5	0.66	0.60	0.72	0.26	0.20	0.32	0.40	2.5	0.15	LC50	0.56	0.07	13.4															
	2	0.48	0.42	0.56	0.21	0.16	0.27	0.45	2.3	0.15	EC50	0.24	0.02	9.7																
	3	0.54	0.50	0.59	0.26	0.20	0.32	0.40	2.1	0.15	MCIG	0.42	0.02	5.7																
DAWSON	1	6	16.10	15.80	16.50	11.66	9.00	15.00	7.50	1.4	17.50	7.50	LC50	15.96	0.46	2.9														
	2	16.44	16.20	16.60	13.96	12.60	15.50	7.50	1.2	17.50	12.50	EC50	12.96	0.56	7.4															
	3	15.34	15.00	15.70	13.27	12.90	13.60	7.50	1.2	15.00	6.00	MCIG	7.50	0.00	0.0															
STOVER/ FORT	1	7	5.47	4.84	6.46	3.90	3.08	4.93	10.00	1.4	1.00	LC50	6.17	0.49	8.0															
	2	6.52	6.25	6.80	5.67	5.37	5.99	7.00	1.2	5.00	3.00	EC50	5.19	0.92	17.7															
	3	6.51	6.32	6.72	5.99	5.69	6.30	6.00	1.1	6.00	5.00	MCIG	7.67	1.70	22.2															
GRAND TOTAL																														
LC50		6.52	5.51	84.5																										
EC50		3.43	4.6247	134.9																										
MCIG		3.75	3.74	98.8																										
T1		1.90	6.05	318.3																										
LOEC (mort)		7.37	5.45	73.9																										
LOEC (malf)		3.73	4.46137	119.5																										

Table 1. Results of the Interlaboratory Study for Monosodium Glutamate.



Laboratory	Replicate (n)	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV ¹ (%)	Mean EC50 (mg/ml)	EC50 CV (%)	Mean TI ² (%)	TI CV (%)	Mean MCIG ³ (mg/ml)	MCIG CV (%)
1	1	4.07	54.0	0.46	26.6	15.4	29.3	1.3	3.92	110.1
	2	5.64	7.80	0.37	0.51	0.69				
	3	13.68								
2	1	8.02	34.6	1.32	0.86	1.02	20.5	7.4	41.1	4.0
	2	4.15	7.56							
	3	10.50								
3	1									
	2									
	3									
4	1	1.35	21.4	0.08	0.65	84.3	1.7	449.2	0.3	0.38
	2	1.07								
	3	0.79								
5	1	0.66								
	2	0.48	0.56	13.4	0.21	0.24	9.7	2.3	8.2	0.4
	3	0.54								
6	1	16.10	15.96	2.9	11.66	12.96	7.4	1.2	8.2	7.5
	2	16.44								
	3	15.34								
7	1	5.47	6.17	8.0	3.90	5.67	5.19	17.7	1.2	11.3
	2	6.52								
	3	6.51								

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

ASTM Analysis of Monosodium Glutamate



FETAX Summary Sheet

Test No. DEF 1 *[Signature]*

Test Material	P3A	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	11-30-92
Composition/Purity		Test End Date	12-4-92
Solvent	NONE	Conc.	Test Units (i.e., mg/ml) MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.1	7.1	7	7.1	
	Control		7.5	7.7	7.4	7.2
	Highest Concentration		7.12	7.15	7	6.9

No. Dead or Malformed		X 100 = %	
Total Number		Mortality Record	Malformation Record
FETAX Control	1 : 100	X 100 = 1%	5 : 99 X 100 = 5.1%
Solvent Control	C19 : E19	X 100 = G19	H19 : J19 X 100 = L19
Control Length	9.87	Solvent Control Length	
Minimum Concentration to Inhibit Growth (MCIG)		10.00	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.5	0.1	T-test
LOEL	5	0.25	T-test
LC50	4.07	EC50 0.460	
95% CL	2.88	5.76	95% Confidence Limits 0.303 ---- 0.698
Test Teratogenic Index (TI = LC50/EC50):			8.85
95% Confidence limits			5.14 -- 15.24

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

JRB
Test No. DEF 2

Test Material	P3A	Investigator	JAMES RAYBURN	
Source	OAK RIDGE	Laboratory	BANTLE	
CAS No.	Lot No.	Test Start Date:	12-7-92	
Composition/Purity		Test End Date	12-11-92	
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML	

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.2	7.3	7.2	7.2	
	Control		7.6	7.4	7.5	7.3
	Highest Concentration		7.5	7.7	7.8	7.8

No. Dead or Malformed					
----- X 100 = %					
Total Number	Mortality Record		Malformation Record		
FETAX Control	7 : 100	X 100 = 7%	4 : 93	X 100 = 4.3%	
Solvent Control	:	X 100 =	:	X 100 =	
Control Length	C20	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)	J21 / .25				

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	5	N.A.	T-test
LOEL	7.5	0.1	T-test
LC50	5.638	EC50 0.372	
95% CL	5.055 -- 6.289	95% Confidence limits	0.241 ---- 0.576

Test Teratogenic Index (TI = LC50/EC50): 15.15

95% Confidence limits 9.66 -- 23.75

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. DEF 3

Test Material	P3A	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	12-13-92
Composition/Purity		Test End Date	12-17-92
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.2	7.1	7	7.1	
	Control		7.6	7.4	7.6	7.3
	Highest Concentration		7.3	7.4	7.3	7.2

No. Dead or Malformed		X 100 = %			
Total Number		Mortality Record		Malformation Record	
FETAX Control	5 : 100	X 100 =	5%	4 : 95	X 100 = 4.2%
Solvent Control	C19 : E19	X 100 =	G19	H19 : J19	X 100 = L19
Control Length	C20	Solvent Control Length		J20	
Minimum Concentration to Inhibit Growth (MCIG)		J21 0.5			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.75	0.25	T-test
LOEL	5	0.5	T-test
LC50	13.676	EC50 0.694	
95% CL	11.488 -- 16.279	95% Confidence Limits	0.538 ---- 0.894
Test Teratogenic Index (TI = LC50/EC50):			19.71
95% Confidence limits			14.49 -- 26.82

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No.

1

Test Material	Unknown P3A	Investigator	Mendi A. Hull
Source	B4	Laboratory	Bantle / OSU
CAS No.	B5	Lot No. E5	Test Start Date: Feb 1 1993
Composition/Purity	C6	Test End Date	Feb 5 1993
Solvent	B7	Conc. E7	Test Units (i.e., mg/ml) mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.0	7.2	7.0	7.0	X
	Control	X	7.8	7.9	7.9	7.6
	Highest Concentration	X	7.9	8.0	7.7	7.7

No. Dead or Malformed			
----- X 100 = %			
Total Number	Mortality Record		Malformation Record
FETAX Control	5 : 100	X 100 = 5%	6 : 95 X 100 = 6.3%
Solvent Control	C19 : E19	X 100 = G19	H19 : J19 X 100 = L19
Control Length	0.9	Solvent Control Length	J20
Minimum Concentration to Inhibit Growth (MCIG)	4 mg/ml		

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	6	N.A.	T-test
LOEL	8	0.25	T-test
LC50	8.018	EC50 1.3164	
95% CL	7.332 -- 8.767	95% Confidence limits	1.0067 ---- 1.7214
Test Teratogenic Index (TI = LC50/EC50):			6.09
95% Confidence limits			4.59 -- 8.08

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No.

2

Test Material	Unknown P3A	Investigator	Mendi A. Hull
Source	B4	Laboratory	Bantle / OSU
CAS No.	B5	Lot No.	E5
Composition/Purity	C6	Test Start Date:	Feb 8 1993
Solvent	B7	Conc.	E7
		Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.1	7.0	7.1	7.2	X
	Control	X	7.1	7.7	7.4	7.4
	Highest Concentration	X	7.7	7.8	8.2	7.7

No. Dead or Malformed	X 100 = %				
MALFORMATION EXCEED ASTM LIMIT					
Total Number	Mortality Record		Malformation Record		
FETAX Control	10 : 100	X 100 = 10%	7 : 90	X 100 = 7.8%	
Solvent Control	C19 : E19	X 100 = G19	H19 : J19	X 100 = L19	
Control Length	0.9 cm	Solvent Control Length		J20	
Minimum Concentration to Inhibit Growth (MCIG)					0.25 mg/ml

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	4	0.5	T-test
LOEL	6	0.75	T-test
LC50	4.148	EC50 0.861	
95% CL	3.830 -- 4.494	95% Confidence limits 0.626 ---- 1.183	

Test Teratogenic Index (TI = LC50/EC50): 4.82

95% Confidence limits 3.47 -- 6.69

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No.

3

Test Material	Unknown P3A	Investigator	Mendi A. Hull
Source	B4	Laboratory	Bantle / OSU
CAS No.	B5	Test Start Date:	Mar 1 1993
Composition/Purity	C6	Test End Date	Mar 5 1993
Solvent	B7	Conc.	E7
		Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.0	7.0	7.0	6.9	X
	Control	X	7.0	7.2	7.4	7.6
	Highest Concentration	X	7.5	7.6	7.7	7.8

No. Dead or Malformed	X 100 = %				
MALFORMATION EXCEED ASTM LIMIT					
Total Number	Mortality Record		Malformation Record		
FETAX Control	7 : 100	X 100 = 7%	9 : 93	X 100 = 9.7%	
Solvent Control	C19 : E19	X 100 = G19	H19 : J19	X 100 = L19	
Control Length	C20	Solvent Control Length		J20	
Minimum Concentration to Inhibit Growth (MCIG)	0.25 0.25				

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	10	1	T-test
LOEL	12	14	T-test
LC50	10.495	EC50 0.886	
95% CL	7.994 -- 13.779	95% Confidence limits	0.237 ---- 3.321

Test Teratogenic Index (TI = LC50/EC50): 11.84

95% Confidence limits 3.07 -- 45.61

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : F33 X 100 = G33	H33 : J33 X 100 = I33
2500 mg/L	C34 : L34 X 100 = G34	H33 : J34 X 100 = I34

CL = Confidence limits

FETAX Summary Sheet

Test No. 1

Test Material	P3A	Investigator	J3
Source		Laboratory	J4
CAS No.	Lot No.	Test Start Date:	5/17/93
Composition/Purity		Test End Date	5/21/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	71	7.2	7.1	7	
	Control		7.3	7.2	7.1	7.3
	Highest Concentration		7.5	7.7	7.9	8.1

No. Dead or Malformed		MALFORMATION EXCEED ASTM LIMITS			
Total Number	X 100 = %	Mortality Record		Malformation Record	
FETAX Control	2 : 100	X 100 =	2%	10 : 98	X 100 = 10.2%
Solvent Control	:	X 100 =		:	X 100 =
Control Length (mm)		Solvent Control Length (mm)			
Minimum Concentration to Inhibit Growth (MCIG)				mg/ml	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.25	0.5	T-test
LOEL	2	5	T-test
LC50	1.354	EC50 0.076	
95% CL	1.46E-07	-- 1.3E+07	95% Confidence limits 1E-10 ---- 6E+07
	Test Teratogenic Index (TI = LC50/EC50):		
	95% Confidence limits 9.35E-11 -- 3.36E+12		

Percent effect	LC	EC
5	3.1484E-05	2E-07
16	0.0021389	3E-05
50	1.35387185	0.076
84	856.966253	171.3
95	58218.6083	26615

FETAX Summary Sheet

Test No. 2

Test Material	P3A	Investigator	Gillett
Source		Laboratory	ERL-C
CAS No.	Lot No.	Test Start Date:	5/17/93
Composition/Purity		Test End Date	5/21/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7	7.1	7.1	7.2	
	Control		7.5	7.2	7.3	7.5
	Highest Concentration		7.4	7.7	7.8	7.9

No. Dead or Malformed	X 100 = %	Mortality Record		Malformation Record			
Total Number							
FETAX Control	4 : 100	X 100 = 4%		6 : 96	X 100 = 6.3%		
Solvent Control	:	X 100 =		:	X 100 =		
Control Length (mm)	Solvent Control Length (mm)						
Minimum Concentration to Inhibit Growth (MCIG)	25				mg/ml		

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	1	1	T-test
LOEL	2	2	T-test
LC50	1.069	EC50 1.386	
95% CL	0.509	-- 2.246	95% Confidence limits 0.525 ---- 3.661
Test Teratogenic Index (TI = LC50/EC50):			0.771
95% Confidence limits			0.227 -- 2.618

Percent	LC	EC
effect		
5	0.1473788	0.138
16	0.3226661	0.343
50	1.0693508	1.386
84	3.543946	5.604
95	7.7589951	13.97

FETAX Summary Sheet

Test No.	3
Test Material	P3A
Investigator	Gillett
Source	Laboratory
CAS No.	Lot No.
Test Start Date:	5/16/93
Composition/Purity	Test End Date
Solvent	Conc.
	Test Units (i.e., mg/ml)
	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
		Stock	7.1	7.2	7.2	7.1
	Control			7.4	7.3	7.3
	Highest Concentration			7.2	7.4	8

No. Dead or Malformed X 100 = %	MALFORMATION EXCEED ASTM LIMITS			
Total Number	Mortality Record			
FETAX Control	2 : 100	X 100 =	2%	10 : 98 X 100 = 10.2%
Solvent Control	:	X 100 =	:	X 100 =
Control Length (mm)	Solvent Control Length (mm)			
Minimum Concentration to Inhibit Growth (MCIG)	0.5 mg/ml			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.5	0.25	T-test
LOEL	1	0.5	T-test
LC50	0.790	EC50 0.482	
95% CL	0.588 -- 1.063	95% Confidence limits	0.405 ---- 0.575
	Test Teratogenic Index (TI = LC50/EC50): 1.638		
	95% Confidence limits 1.161 -- 2.312		

Percent	LC	EC
effect		
5	0.4188964	0.279
16	0.5383593	0.346
50	0.7901094	0.482
84	1.1595839	0.672
95	1.4902799	0.835

FETAX SUMMARY SHEET

Test No. 2

Test Material	UNKNOWN P3A	Investigator	TURLEY
Source	OSU	Lab	U.MD-WREC
CAS No.	-	Test Start Date	NOVEMBER 17, 1992
Composition/Purity	-	Test End Date	NOVEMBER 21, 1992
Solvent	-	Test Units (i.e., µg/ml)	mg/ml

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH					
Stock	7.20	7.22	7.17	7.24	7.26
Control	7.80	7.72	7.84	7.76	7.85
Highest Conc.	7.35	7.31	7.37	7.35	7.39

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	1 : 100 X 100 = 1 %	8 : 99 X 100 = 8 %
Solvent Control	: X 100 = %	: X 100 = %
Control Length 10.09 mm	Solvent Control Length — mm	
Minimum Concentration to Inhibit Growth (MCIG) 0.45 mg/ml		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	0.10	Bonferroni T-test
LOEL	FAIL HOMOGENEITY	0.15	Bonferroni T-test
LC ₅₀	0.48 mg/ml (Trimmed Spearman-Karber)	EC ₅₀ 0.21 mg/ml (Trimmed Spearman-Karber)	
95% Confidence Limits	0.42 - 0.56	95% Confidence Limits 0.16 - 0.27	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		2.29	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

Test No. 3

Test Material	UNKNOWN P3A	Investigator	TURLEY
Source	O.S.U	Lab	UMD-WREC
CAS No.		Test Start Date	NOV. 30
Composition/Purity		Test End Date	DEC. 3, 1992
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.19	7.16	7.21	7.22	7.19
Control	7.79	7.84	7.90	7.82	7.86
Highest Conc.	7.30	7.29	7.36	7.31	7.36

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	2 : 100 X 100 = 2 %	8 : 98 X 100 = 8 %
Solvent Control	—	— : — X 100 = — %
Control Length 10.13 mm	Solvent Control Length	— mm
Minimum Concentration to Inhibit Growth (MCIG) 0.40 mg/ml		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LC ₅₀ 0.54 mg/ml Trimmed Spearman-Kubber	EC ₅₀ 0.26 mg/ml Trimmed Spearman-Kubber		
95% Confidence limits 0.50 - 0.59 mg/ml	95% Confidence Limits 0.20 - 0.32 mg/ml		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) 2.08			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	— : — X 100 = — %	— : — X 100 = — %
2500 mg/L	— : — X 100 = — %	— : — X 100 = — %

170

FETAX SUMMARY SHEET

Test No. 4

Test Material	UNKNOWN P3A	Investigator	TURLEY
Source	O.S.U	Lab	UMD-WREC
CAS No.	Lot No.	Test Start Date	12/4/92
Composition/Purity		Test End Date	12/8/92
Solvent	Cone.	Test Units (i.e., mg/ml)	mg/ml

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH	7.13	7.16	7.19	7.14	7.18
Stock	7.13	7.16	7.19	7.14	7.18
Control	7.65	7.63	7.61	7.64	7.69
Highest Conc.	7.21	7.18	7.26	7.24	7.28
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %				
Total Number	2 : 100 X 100 = 2 %	7 : 98 X 100 = 7 %			
Solvent Control	— : — X 100 = — %	— : — X 100 = — %			
Control Length 10.03 mm	Solvent Control Length	— mm			
Minimum Concentration to Inhibit Growth (MICIG) 0.40					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Pi Ferroni T-test
LOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LC ₅₀ 0.60 mg/ml (Trimmed Spearman-Karber)	EC ₅₀ 0.26 mg/ml (Trimmed Spearman-Karber)		
95% Confidence limits (0.60 - 0.72)	95% Confidence Limits 0.30 - 0.32		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		2.54	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	— : — X 100 = — %	— : — X 100 = — %
2500 mg/L	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

		Test No. <u>1</u>
Test Material	P3A	
Source	Investigator <u>J.A. DAWSON</u>	
CAS No.	Lot No.	Lab <u>WT/CMH</u>
Composition/Purity	Test Start Date <u>11/30/92</u>	
Solvent	Cone.	Test End Date <u>12/4/92</u> Test Units (l.s., mg/ml) <u>mg/ml</u>

<u>pH</u>	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.10	7.10	7.13	7.13	—
Control	7.83	7.69	7.58	7.43	7.37
Highest Cone.	—	7.62	7.74	7.96 ^{de}	8.19 ^{de}

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		0 : 100 X 100 = 0 %	2 : 100 X 100 = 2 %
Solvent Control		_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
Control Length mm		Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG)		7.5	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	15 both	< 7.5 u.u.	
LOEL	175 both	7.5 u.u.	
LC ₅₀	16.10	EPA	EC ₅₀ 11.66
95% Confidence Limits (15.8-16.5)	Prob.t		95% Confidence Limits (9.0-15.0) >L1W
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		138 - L1W/EC ₅₀	

EPA
 F-test
 12.35
 (u.u. - 7.5)

ISK
 12.1
 (11.8-13.0)

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L Not	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L dose	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

J.A.J.

12/14/92

FETAX SUMMARY SHEET

Test No. 2

Test Material	P3A	Investigator	J.A. DAWSON
Source		Lab	AT/CCVY
CAS No.	Lot No.	Test Start Date	12/8/92
Composition/Purity		Test End Date	12/12/92
Solvent	Conc.	Test Units (i.e., mg/ml)	1mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.11	7.11	7.10	7.10	-
Control	7.87	7.80	7.62	7.62	7.62
Highest Conc.	-	7.64	7.77	7.95 ^{ac}	8.02 ^{ad}

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		0 : 100 X 100 = 0 %	3 : 100 X 100 = 3 %
Solvent Control		_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
Control Length mm	-	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG) 7.5			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	15.0 both	100 both	
LOEL	17.5 both	125 both	
LC ₅₀	16.44 TSK	EC ₅₀ 13.96 CFW	
95% Confidence Limits 16.2-16.6			95% Confidence Limits (12.6 15.5)
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.18

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

Test No. 3

Test Material	P3A	Investigator	D. A. DAWSON
Source		Lab	WRI/CVM
CAS No.	Lot No.	Test Start Date	1/4/93
Composition/Purity		Test End Date	1/8/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.12	7.10	7.12	7.07	—
Control	7.77	7.68	7.54	7.56	7.51
Highest Conc.	—	7.63	7.74	7.78	7.81
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	$\times 100 = \%$		Total Number	<u>0</u> : <u>100</u> $\times 100 = \underline{\quad}$ %	<u>0</u> : <u>100</u> $\times 100 = \underline{\quad}$ %
Solvent Control	<u> </u> : <u> </u> $\times 100 = \underline{\quad}$ %	<u> </u> : <u> </u> $\times 100 = \underline{\quad}$ %	Control Length mm	Solvent Control Length mm	mm
Minimum Concentration to Inhibit Growth (MCIG)			7.5		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	12.5 both	—	hazardarity failed for EC ₅₀ - NOEL-LOEL
LOEL	15.0	—	
LC ₅₀	15.34	EC ₅₀ 13.27	
95% Confidence limits (15.0 - 15.7)	>TSK	95% Confidence Limits (12.9 - 13.6) >TSK	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.16

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L 1/6	<u> </u> : <u> </u> $\times 100 = \underline{\quad}$ %	<u> </u> : <u> </u> $\times 100 = \underline{\quad}$ %
2500 mg/L 1/4	<u> </u> : <u> </u> $\times 100 = \underline{\quad}$ %	<u> </u> : <u> </u> $\times 100 = \underline{\quad}$ %

FETAX SUMMARY SHEET

Test No. 1

Test Material	P3A	Investigator	Fort
Source	ILS, Phase 3	Lab	SBL
CAS No.	Lot No.	Test Start Date	12/14/92
Composition/Purity		Test End Date	12/18/92
Solvent	Conc.	Test Units (i.e., mg/ml)	

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.3	7.2	7.3	7.4	
Control	7.9	7.9	7.8	7.9	
Highest Conc.	7.6	7.5	7.5	7.6	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	1 : 100 X 100 = 1 %	6 : 99 X 100 = 0 %
Solvent Control	— : — X 100 = — %	— : — X 100 = — %
Control Length 97.3 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG)	10.0	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.1	0.1	Dilution
LOEL	1.0	1.0	Williams
LC ₅₀	5.47	EC ₅₀ 3.90	
95% Confidence Limits	4.64 - 6.46	95% Confidence Limits	3.08 - 4.93
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.40

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	— : — X 100 = — %	— : — X 100 = — %
2500 mg/L	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

		Test No. 2				
Test Material	P3A					
Source	ILS					
CAS No.	Lot No.					
Composition/Purity		Test Start Date 2/5/93				
Solvent	Conc. Test Units (i.e., mg/ml) mg/ml					
<u>pH</u>		DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock		7.3	7.4	7.4	7.4	
Control		9.0	7.9	7.9	7.9	
Highest Conc.		7.5	7.5	7.6	7.5	
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD		
No. Dead or Malformed		$\times 100 = \%$				
Total Number		$0 : 100 \times 100 = 0 \%$		$0 : 100 \times 100 = 0 \%$		
Solvent Control		$: \quad \times 100 = \quad \%$		$: \quad \times 100 = \quad \%$		
Control Length 97.0 mm		Solvent Control Length mm				
Minimum Concentration to Inhibit Growth (MCIG) 7.0						

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	4.0	4.0	Williams Test
LOEL	5.0	3.0	
LC ₅₀	6.52	EC ₅₀	5.67
95% Confidence limits	6.25 - 6.80	95% Confidence Limits	5.37 - 5.99
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.15

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	$: \quad \times 100 = \quad \%$	$: \quad \times 100 = \quad \%$
2500 mg/L	$: \quad \times 100 = \quad \%$	$: \quad \times 100 = \quad \%$

FETAX SUMMARY SHEET

DF
Test No.

3

Test Material	P3A	Investigator	Fort
Source	TLS	Lab	SBL
CAS No.	Lot No.	Test Start Date	2/6/93
Composition/Purity		Test End Date	2/10/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.4	7.5	7.5	7.5	
Control	8.0	7.9	7.9	7.8	
Highest Conc.	7.6	7.6	7.6	7.6	
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %				
Total Number					
Solvent Control					
Control Length 96.0 mm	Solvent Control Length	mm			
Minimum Concentration to Inhibit Growth (MCIG)	6.0				

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	5.0	4.0	Williams Test
LOEL	6.0	5.0	
LC ₅₀	6.51	EC ₅₀	5.99
95% Confidence limits	6.32 - 6.72	95% Confidence Limits	5.69 - 6.30
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		1.09	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

ASCORBIC ACID

PHASE III COMPOUND B DATA (P3B) ASCORBIC ACID

FETAX ILS STUDY; 7 LABS, 8 LEVELS, 3 REPLICATES
PAGE 1

FEIAX

ORIGINAL DATA TABLE WITH MATERIALS

Table 2. Results of the Interlaboratory Study for Ascorbic Acid.



Laboratory	Replicate (n)	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV1 (%)	EC50 (mg/ml)	Mean EC50 (mg/ml)	EC50 CV (%)	Mean TI2 (%)	TI CV (%)	MCIG ³ (mg/ml)	Mean MCIG (mg/ml)	MCIG CV (%)
1	1	11.04	7.01							5.0		
	2	6.98	10.98	29.5	5.31	6.60	14.1	1.7	16.8	10.0	8.33	28.3
	3	14.91		7.48						10.0		
2	1	15.21	7.00							2.0		
	2	8.85	13.06	22.8	6.69	6.29	12.7	2.1	31.5	2.0	2.67	35.4
	3	15.11		5.17						4.0		
3	1	15.61	11.33							2.0		
	2	16.88	14.83	13.9	11.99	11.60	2.4	1.3	12.9	4.0	3.33	28.3
	3	12.00		11.49						4.0		
4	1	2.78	0.82							0.3		
	2	1.61	1.82	39.5	0.70	0.74	7.6	2.5	31.3	0.3	0.37	25.7
	3	1.06		0.70						0.5		
5	1	9.26	7.24							9.0		
	2	10.21	9.91	4.6	7.69	7.43	2.6	1.3	3.6	8.0	8.33	5.7
	3	10.25		7.35						8.0		
6	1											
	2											
	3											
7	1	12.20				11.94				15.0		
	2	14.26	13.61	7.3	13.47	13.03	6.0	1.0	1.5	16.0	14.67	8.5
	3	14.36				13.68				13.0		

¹CV=Coefficient of Variation.



²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

ASTM Analysis of Ascorbic Acid



p3b



■ LC50 ■ EC50 ■ MCIG ■ TI

FETAX Summary Sheet

Test No. DEF 1 *JHAB*

Test Material	P3B	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	11-30-92
Composition/Purity		Test End Date	12-4-92
Solvent	NONE	Conc.	Test Units (i.e., mg/ml) MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.1	8	8.2	7.5	
	Control		7.2	7.5	7.2	7
	Highest Concentration		7.57	7.51	7.3	7.6

No. Dead or Malformed		X 100 = %			
Total Number		Mortality Record		Malformation Record	
FETAX Control	1 : 100	X 100 =	1%	5 : 99	X 100 = 5.1%
Solvent Control		X 100 =		:	X 100 =
Control Length	C20	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)		5.0 J21			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	10	5	T-test
LOEL	12	7.5	T-test
LC50	11.038	EC50 7.014	
95% CL	9.967 -- 12.225	95% Confidence limits	6.179 ---- 7.961
Test Teratogenic Index (TI = LC50/EC50):			1.57
95% Confidence limits			1.34 -- 1.85

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No. DEF 2

Test Material	P3B	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	12-7-92
Composition/Purity		Test End Date	12-11-92
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

		Day 0	Day 1	Day 2	Day 3	Day 4
pH	Stock	7.5	7.2	7.6	7.4	
	Control		7.4	7.3	7.6	7.3
	Highest Concentration		7.5	7.3	7.3	7.3

No. Dead or Malformed	X 100 = %				
Total Number	Mortality Record			Malformation Record	
FETAX Control	10 : 100	X 100 =	10%	6 : 90	X 100 = 6.7%
Solvent Control	:	X 100 =	:	X 100 =	
Control Length	C20	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)			10	J21	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	6	4	T-test
LOEL	8	6	T-test
LC50	6.975	EC50 5.314	
95% CL	6.052 -- 8.039	95% Confidence limits	4.483 ---- 6.300
	Test Teratogenic Index (TI = LC50/EC50):		
	95% Confidence limits		

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. DEF 3

Test Material	P3B	Investigator	JAMES RAYBURN <i>JRB</i>
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	12-13-92
Composition/Purity		Test End Date	12-17-92
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH	Stock	Day 0	Day 1	Day 2	Day 3	Day 4
	Control		7.3	7.5	7.3	7
	Highest Concentration		7.2	7.4	7.1	7

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		4 : 100 X 100 = 4%		4 : 96 X 100 = 4.2%			
Solvent Control		C19 : E19 X 100 = G19		H19 : J19 X 100 = L19			
Control Length		C20		Solvent Control Length		J20	
Minimum Concentration to Inhibit Growth (MCIG)				10		J21	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used				
NOEL	6	6	T-test				
LOEL	18	8	T-test				
LC50	14.907		EC50	7.475			
95% CL	14.078	--	15.785	95% Confidence limits	6.786	----	8.235
Test Teratogenic Index (TI = LC50/EC50):				1.99			
95% Confidence limits				1.78	--	2.23	

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No.	1
Test Material	Unknown B6A
Source	B4
CAS No.	B5
Composition/Purity	C6
Solvent	B7
Conc.	E7
Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.0	8.0	7.7	7.9	X
	Control	X	7.7	7.4	7.5	7.6
	Highest Concentration	X	7.9	8.0	7.7	7.4

No. Dead or Malformed		X 100 = %			
Total Number		Mortality Record		Malformation Record	
FETAX Control	9 : 100	X 100 =	9%	6 : 91	X 100 = 6.6%
Solvent Control	C19 : E19	X 100 =	G19	H19 : J19	X 100 = L19
Control Length	1 cm	Solvent Control Length		J20	
Minimum Concentration to Inhibit Growth (MCIG)				2 mg/ml	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	14	4	T-test
LOEL	16	6	T-test
LC50	15.209	EC50 7.004	
95% CL	14.264 -- 16.217	95% Confidence Limits	6.252 ---- 7.846
	Test Teratogenic Index (TI = LC50/EC50):		
	95% Confidence Limits		

POSTIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. 2

Test Material	Unknown P3B	Investigator	Mendi A. Hull
Source	B4	Laboratory	Bantle / OSU <i>JMB</i>
CAS No.	B5	Test Start Date:	Feb 8 1993
Composition/Purity	C6	Test End Date	Feb 12 1993
Solvent	B7	Conc.	E7
		Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.9	7.9	8.0	7.9	X
	Control	X	7.7	7.4	7.5	7.4
	Highest Concentration	X	7.2	7.0	6.8	6.9

No. Dead or Malformed		MALFORMATION EXCEED ASTM LIM			
X 100 = %		Mortality Record		Malformation Record	
Total Number		10 : 100 X 100 = 10%		9 : 90 X 100 = 10.0%	
FETAX Control		10 : 100 X 100 = 10%		9 : 90 X 100 = 10.0%	
Solvent Control		C19 : E19 X 100 = G19		H19 : J19 X 100 = L19	
Control Length		0.9 cm		Solvent Control Length J20	
Minimum Concentration to Inhibit Growth (MCIG)				2 mg/ml	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	7	7	T-test
LOEL	8	8	T-test
LC50	8.854	EC50 6.692	
95% CL	7.487 -- 10.472	95% Confidence limits 5.982 ---- 7.487	

Test Teratogenic Index (TI = LC50/EC50): 1.32

95% Confidence limits 1.08 -- 1.62

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No. 3

Test Material	Unknown P3B	Investigator	Mendi A. Hull
Source		Laboratory	Bantle / OSU <i>JMH</i>
CAS No.	Lot No.	Test Start Date:	Mar 1 1993
Composition/Purity		Test End Date	Mar 5 1993
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.8	7.9	8.0	8.1	X
	Control	X	7.4	7.2	7.5	7.8
	Highest Concentration	X	6.8	6.8	6.5	6.7

No. Dead or Malformed		MALFORMATION EXCEED ASTM LIM			
Total Number	X 100 = %	Mortality Record		Malformation Record	
FETAX Control	3 : 100	X 100 =	3%	10 : 97	X 100 = 10
Solvent Control	:	X 100 =	:	X 100 =	
Control Length	1 cm	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)		4 mg/ml			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	10	N.A.	T-test
LOEL	12	2	T-test
LC50	15.114	EC50 5.172	
95% CL	13.234 -- 17.260	95% Confidence limits 3.949 ---- 6.772	
Test Teratogenic Index (TI = LC50/EC50):			2.92
95% Confidence limits			2.16 -- 3.95

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	: X 100 =	: X 100 =
2500 mg/L	: X 100 =	: X 100 =

CL= Confidence limits

FETAX Summary Sheet

Test No. 3-P3B

Test Material	P3B	Investigator	FINCH
Source	OSU	Laboratory	USABRDL
CAS No.	-	Lot No.	-
Composition/Purity	-	Test Start Date:	10 MAY 93
Solvent	-	Test End Date	14 MAY 93
Conc.	-	Test Units (i.e., mg/ml)	MG/ML

pH	Stock	Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	6.9	7.0	7.5	7.0	-
	Control	-	7.9	7.9	7.9	7.9
	Highest Concentration	-	7.5	7.4	7.7	7.2

No. Dead or Malformed	X 100 = %		
Total Number	Mortality Record	Malformation Record	
FETAX Control	0 : 100 X 100 = ✓ 0%	7 : 100 X 100 =	7.0% ✓
Solvent Control	-	-	-
Control Length	9.641	Solvent Control Length	-
Minimum Concentration to Inhibit Growth (MCIG)	-	-	2.0 ✓

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	14	10	T-test
LOEL	16	12	T-test
LC50	15.608 ✓	EC50 11.331 ✓	
95% CL	15.019 --	16.220	95% Confidence limits 10.673 ---- *****
Test Teratogenic Index (TI = LC50/EC50):			1.38 ✓
95% Confidence limits			1.28 -- 1.48

CL = Confidence limits		

CL = Confidence limits

FETAX Summary Sheet

Test No. 5-P3B

Test Material	P3B	Investigator	FINCH
Source	OSU	Laboratory	USABRDE
CAS No.	Lot No.	Test Start Date	17 MAY 93
Composition/Purity	-	Test End Date	21 MAY 93
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.5	7.0	7.2	7.2	7.2
	Control		7.9	7.9	7.9	7.9
	Highest Concentration		7.8	7.3	7.7	7.3

No. Dead or Malformed		
X 100 = %		
Total Number	Mortality Record	Malformation Record
FETAX Control	1 : 100 X 100 = ✓ 1%	5 : 99 X 100 = 5.1%
Solvent Control	--	--
Control Length	10.295	Solvent Control Length
Minimum Concentration to Inhibit Growth (MCIG)		4.0 ✓

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	16	8	T-test
LOEL	18	14	T-test
LC50	16.876 ✓	EC50 11.992	12.4
95% CL	15.497 --	18.377	95% Confidence limits 11.054 --- 13.009

Test Teratogenic Index (TI = LC50/EC50): 1.41

95% Confidence limits 1.25 -- 1.58

CL = Confidence limits

FETAX Summary Sheet

Test No.: 4-P3B

Test Material	Item #:	12345	Investigator	FINCH
Source	OSU		Laboratory	USABRDL
CAS No.	E5	Lat. No. E5	Test Start Date	
Composition/Purity	(C)		Test End Date	
Solvent	Corn oil		Test Units (Kg/Embryos/ml)	MG/Ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	E11	E11	G11	J11	
	Control		E12	G12	J11	K12
	Highest Concentration		E13	B13	J11	K13

No. Dead or Malformed	X 100 = %		
Total Number			
FETAX Control	4 : 100	X 100 = 4%	6 : 96 X 100 = 6.3%
Solvent Control	E19 : E19	X 100 = G19	H19 : J19 X 100 = L19
Control Length	C20	Solvent Control Length	J20
Minimum Concentration to Inhibit Growth (MCIG)			J21

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	16	18	T-test
LOEL	18	20	T-test
LC50	11.99629	EC50 11.49150	
95% CL	7.69582	18.69988	95% Confidence limits 9.67021 --- 13.65583
Test Teratogenic Index (TI = LC50/EC50):			1.04
95% Confidence limits			0.65 --- 1.68

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	E33 : E33	X 100 = G33
2500 mg/L	E34 : E34	X 100 = G34

CL= Confidence limits

FETAX SUMMARY SHEET

Test Material: P3B	Test No. Def #1
Source: Bantle	Investigator: Gillett
CAS No. — Lot No. —	Laboratory: CERL
Composition / Purity: —	Test Start Date: 06/28/93
Solvent: N/A Conc. —	Test End Date: 07/02/93
	Test Units:(i.e., mg/ml) mg/ml

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.2	7.1	7.1	7.2	X
Control	X	7.5	7.4	7.4	7.6
Highest Conc.	X	6.9	7.1	7.5	7.2

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	6 : 100 X 100 = 6 %	4 : 94 X 100 = 4.3 %
Solvent Control: N.A.	— : — X 100 = — %	— : — X 100 = — %
Control Length: 0.907 ± 0.045 mm	Solvent Control Length: — mm	
Minimum Concentration to Inhibit Growth (MCIG): 0.3 mg/ml		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.0 mg/ml	0.7 mg/ml	Williams
LOEL	5.0 mg/ml	1.0 mg/ml	Williams
LC ₅₀	2.78 mg/ml	EC ₅₀ 0.82 mg/ml	
95% Confidence Limits 2.24 - 3.45		95% Confidence Limits 0.69 - 0.97 mg/ml	
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) - 3.39			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	— : — X 100 = — %	— : — X 100 = — %
2.5 mg/ml	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

Test Material: P3B	Test No. Def. #2
Source: Bantle	Investigator: Gillett
CAS No. — Lot No. —	Laboratory: Corvallis Env. Res. Lab
Composition / Purity: —	Test Start Date: 8/2/93
Solvent: — Conc. —	Test End Date:
	Test Units:(i.e., mg/ml)

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.0	7.1	7.3	7.4	X
Control	X	7.0	7.4	7.1	7.2
Highest Conc.	X	6.9	7.1	7.2	7.1

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed x 100 = % Total Number	8 : 100 X 100 = 8 %	10 : 92 X 100 = 10.8 %
Solvent Control: NA 0.83 ± 0.74	— : — X 100 = — %	— : — X 100 = — %
Control Length: 0.616 ± 0.074 mm	Solvent Control Length:	mm
Minimum Concentration to Inhibit Growth (MCIG): 0.3 .		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.7 mg/ml	0.3 mg/ml	Williams
LOEL	1.0 mg/ml	0.5 mg/ml	Williams
LC_{50} 1.61		EC_{50} 0.70	
95% Confidence Limits 1.26 - 2.04 mg/ml		95% Confidence Limits 0.48 - 1.03 mg/ml	
TEST TERATOGENIC INDEX (TI = LC_{50}/EC_{50}) - 2.3			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	— : — X 100 = — %	— : — X 100 = — %
2.5 mg/ml	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

		Test No. <u>3</u> DEF.
Test Material: <u>P3B</u>		Investigator: <u>Gillett</u>
Source: <u>Banfile</u>		Laboratory: <u>CERL</u>
CAS No. _____	Lot No. _____	Test Start Date: <u>8-10-93</u>
Composition / Purity: _____		Test End Date: <u>8-15-93</u>
Solvent: <u>N/A</u> Conc. _____		Test Units:(i.e., mg/ml) <u>mg/ml</u>

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.3	7.1	7.1	7.1	X
Control	X	7.4	7.5	7.4	7.4
Highest Conc.	X	6.9	7.2	6.8	8.0

FETAX CONTROL	MORTALITY	MALFORMATION
<u>No. Dead or Malformed</u> X 100 = <u> </u> <u>Total Number</u>	<u> </u> : <u>100</u> X 100 = <u> </u> %	<u> </u> : <u>99</u> X 100 = <u>9.1</u> %
Solvent Control: <u>N/A</u>	<u> </u> : <u> </u> X 100 = <u> </u> %	<u> </u> : <u> </u> X 100 = <u> </u> %
Control Length: <u>0.98</u> ± <u>.105</u> mm	Solvent Control Length: <u> </u> mm	
Minimum Concentration to Inhibit Growth (MCIG): <u>.5</u> mg/ml		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	<u>0.7</u> mg/ml	<u>0.3</u> mg/ml	Williams
LOEL	<u>1.0</u> mg/ml	<u>0.5</u> mg/ml	Williams
LC ₅₀	<u>1.06</u> mg/ml	EC ₅₀ <u>0.70</u> mg/ml	
95% Confidence Limits	<u>0.93 - 1.22</u> mg/ml	95% Confidence Limits <u>0.51-0.96</u> mg/ml	
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) - <u>1.51</u>			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
<u>0.0055</u> mg/ml	<u> </u> : <u> </u> X 100 = <u> </u> %	<u> </u> : <u> </u> X 100 = <u> </u> %
<u>2.5</u> mg/ml	<u> </u> : <u> </u> X 100 = <u> </u> %	<u> </u> : <u> </u> X 100 = <u> </u> %

FETAX SUMMARY SHEET

YV

Test Material UNKNOWN P3B		Test No. 1			
Source OSU		Investigator TURLEY			
CAS No.		Lab U.MD-WREC			
Composition/Purity		Test Start Date 1/21/93			
Solvent -		Test End Date 1/25/93			
Conc.		Test Units (i.e., µg/ml) mg/ml			
pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.33	7.40	7.40	7.38	7.39
Control	7.61	7.63	7.66	7.62	7.69
Highest Conc.	7.40	7.42	7.44	7.48	7.48
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed Total Number		$X 100 = \%$		$2 : 100 X 100 = 2 \% \quad 6 : 98 X 100 = 6 \%$	
Solvent Control		$- : - X 100 = - \% \quad - : - X 100 = - \%$			
Control Length 10.21 mm		Solvent Control Length mm			
Minimum Concentration to Inhibit Growth (MCIG) 9.0 mg/ml					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LC ₅₀	9.26 Trimmed Spearman-Karber	EC ₅₀ 7.24 Trimmed Spearman-Karber	
95% Confidence limits (8.93-9.61)		95% Confidence Limits (6.66-7.86)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.28

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	$\quad : \quad X 100 = \quad \% \quad$	$\quad : \quad X 100 = \quad \% \quad$
2500 mg/L	$\quad : \quad X 100 = \quad \% \quad$	$\quad : \quad X 100 = \quad \% \quad$

FETAX SUMMARY SHEET

YD
77

Test No. 2

Test Material	UNKNOWN P3B	Investigator	TURLEY
Source	OSU	Lab	U MD - WREC
CAS No.	Lot No.	Test Start Date	1/21/93
Composition/Purity		Test End Date	1/25/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.36	7.40	7.40	7.40	7.36
Control	7.71	7.65	7.60	7.71	7.63
Highest Conc.	7.59	7.57	7.55	7.56	7.57

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	2 : 100 X 100 = 2 %	6 : 98 X 100 = 6 %
Solvent Control	: X 100 = %	: X 100 = %
Control Length 10.02 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG)	8.0 mg/ml	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LC ₅₀ 10.21 mg/ml Truncated Spearman-Karber	EC ₅₀ 7.69 mg/ml Truncated Spearman-Karber		
95% Confidence limits (9.79 - 10.66)	95% Confidence Limits (7.10 - 8.33)		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)	1.33		

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

81"

Test No. 3

Test Material UNKNOWN P3B		Investigator TURLEY
Source O.S.U.		Lab UMD-WREC
CAS No.	Lot No.	Test Start Date JAN 27, 1993
Composition/Purity		Test End Date JAN 31, 1993
Solvent	Conc.	Test Units (i.e., mg/ml) mg/ml

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH	7.42	7.40	7.41	7.44	7.46
Stock	7.42	7.40	7.41	7.44	7.46
Control	7.69	7.71	7.73	7.61	7.67
Highest Conc.	7.58	7.59	7.63	7.58	7.57
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %	3 : 100 X 100 = 3 %	7 : 97 X 100 = 7 %		
Total Number					
Solvent Control					
Control Length 10.16 mm	Solvent Control Length mm				
Minimum Concentration to Inhibit Growth (MCIG) 8.0 mg/ml					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LC ₅₀ 10.25 mg/ml	Trimmed Spearman-Karber	EC ₅₀ 7.35 mg/ml Trimmed Spearman-Karber	
95% Confidence limits 9.65 - 10.90		95% Confidence Limits 6.69 - 8.07	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) 1.39			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

Test No. 1

Test Material	P3B	Investigator	FORT
Source	ILS, Floor E	Lab	SBL
CAS No.	Lot No.	Test Start Date	12/14/92
Composition/Purity		Test End Date	12/18/92
Solvent	Conc.	Test Units (i.e., mg/ml)	

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	3.5/7.0	3.5/7.0	3.5/7.0	3.5/7.0	
Control	8.0	7.9	7.9	7.9	
Highest Conc.	7.3	7.4	7.5	7.5	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	() : 100 X 100 = 0 %	: 100 X 100 = %
Solvent Control	— : — X 100 = — %	— : — X 100 = — %
Control Length 95.0 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG) 15.0		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	5.0	5.0	Williams
LOEL	10.0	10.0	Williams
LC ₅₀	12.20	EC ₅₀ 11.94	
95% Confidence Limits	11.33 - 13.13	95% Confidence Limits	10.59 - 13.46
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.07

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	— : — X 100 = — %	— : — X 100 = — %
2500 mg/L	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

DK

Test No. 2

Test Material	P3-B	Investigator	Fort
Source	FETAX ILS	Lab	SBL
CAS No.	Lot No.	Test Start Date	2/18/93
Composition/Purity		Test End Date	3/4/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock					
Control	7.8	7.8	7.7	7.8	
Highest Conc.	5.0 / 7.0	5.2 / 1.0	4.9 / 7.0	5.1 / 7.0	
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %				
Total Number		0 : 100 X 100 = 0 %		1 : 100 X 100 = 1 %	
Solvent Control					
Control Length 94.3 mm		Solvent Control Length mm			
Minimum Concentration to Inhibit Growth (MCIG) 16.0					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	11.0	11.0	Williams
LOEL	12.0	12.0	"
LC ₅₀	14.26	EC ₅₀	13.47
95% Confidence limits	14.00 - 14.54	95% Confidence Limits	13.14 - 13.80
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		1.06	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

Test No. 3
DF

Test Material	P3-B	Investigator	Fort
Source	FETAX ILS	Lab	SBL
CAS No.		Test Start Date	3/2/93
Composition/Purity		Test End Date	3/6/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/lQ

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH					
Stock					
Control	8.0	7.9	7.9	7.9	
Highest Conc.	5.0 / 7.0	4.9 / 7.0	4.8 / 7.0	5.1 / 7.0	

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number			
Solvent Control			
Control Length 90.0 mm	Solvent Control Length mm		
Minimum Concentration to Inhibit Growth (MCIG)		13.0	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	12.0	12.0	Williams
LOEL	13.0	13.0	"
LC ₅₀	14.36	EC ₅₀ 13.68	
95% Confidence limits	14.08 - 14.65	95% Confidence Limits	13.43 - 13.92
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		1.05	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

β -AMINOPROPIONITRILE

PHASE III COMPOUND C DATA (P3C) B-AMINOPROPIONITRILE

FETAX ILS STUDY; 7 LABS, 8 LEVELS, 3 REPLICATES
PAGE 1
units are in mg/L

Table 3. Results of the Interlaboratory Study for β -aminopropionitrile.



Laboratory	Replicate (n)	LC50 (mg/L)	Mean LC50 (mg/L)	CV1 (%)	EC50 (mg/L)	Mean EC50 (mg/L)	CV (%)	EC50 (mg/L)	Mean TI ² (%)	TI (%)	MCIG ³ (mg/L)	Mean MCIG (mg/L)	MCIG CV (%)
1	1	93.70	93.70	0.07	25.5	1221.0	58.7	0.08	0.08	0.08	14.1		
	2	45.40	79.37	30.4	0.08	0.07							
	3	99.00		0.04									
2	1	9.50	29.43	86.6	0.021	0.30	57.9	97.9	30.9	0.20	25.00	11.73	86.9
	2	65.40		0.15									
	3	13.40											
3	1	5.30	5.30	0.28	0.93	49.2	4.5	167.2	0.25	0.25	0.33	35.4	
	2	2.90	4.17	23.6	1.28								
	3	4.30		1.24									
4	1	12.60	12.60	2.22	0.63	1.15	66.2	35.4	213.6	0.50	0.45	59.5	
	2	105.00	40.53	112.8	0.63								
	3	4.00		0.59									
5	1	5.50	4.90	8.8	0.67	0.69	5.2	7.1	12.2	1.00	0.00133	35.4	
	2	4.70		0.66									
	3	4.50		0.74									
6	1	7.70	7.93	7.0	0.21	0.20	11.8	40.1	20.6	0.10	0.15	0.20	54.0
	2	7.40		0.22									
	3	8.70		0.17									
7	1	60.00	77.00	18.6	1.10	1.20	1.07	11.7	72.2	1.00	0.50	0.67	35.4
	2	76.00		0.90									
	3	95.00											

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

ASTM Analysis of β -aminopropionitrile



FETAX Summary Sheet

Test No. DEF 1

Test Material	P3C	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	11-30-92
Composition/Purity		Test End Date	12-4-92
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	6.5	6.6	6.6	6.5	
	Control		7.6	7.2	7.2	7.3
	Highest Concentration		6.6	6.7	7.3	7.4

No. Dead or Malformed	X 100 = %		
Total Number	Mortality Record		
FETAX Control	9 : 100	X 100 = 9%	5 : 91 X 100 = 5.5%
Solvent Control		X 100 =	: X 100 =
Control Length	C20	Solvent Control Length	
Minimum Concentration to Inhibit Growth (MCIG) J21 0.00008			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.075	2.5E-05	T-test
LOEL	0.1	5E-05	T-test
LC50	0.094	EC50 6.93E-05	
95% CL	0.065 -- 0.135	95% Confidence limits	4.59E-05 --- 1.0E-04
Test Teratogenic Index (TI = LC50/EC50):			1351.44
95% Confidence limits			780.52 -- 2339.95

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality		Malformation	
5.5 mg/L	C33 : E33	X 100 = G33	H33 : J33	X 100 = L33
2500 mg/L	C34 : E34	X 100 = G34	H33 : J34	X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. DEF 2

Test Material	P3C	Investigator	JAMES RAYBURN <i>JRB</i>
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	12-7-92
Composition/Purity		Test End Date	12-11-92
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	6.6	6.5	6.7	6.5	
	Control		7.3	7.2	7.5	7.3
	Highest Concentration		7.2	7.4	7.1	6.7

No. Dead or Malformed		X 100 = %			
Total Number		Mortality Record		Malformation Record	
FETAX Control	5 : 100	X 100 =	5%	5 : 95	X 100 = 5.3%
Solvent Control	:	X 100 =		X 100 =	
Control Length	C20	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)				0.000088	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.00075	2.5E-05	T-test
LOEL	0.05	5E-05	T-test
LC50	0.045	EC50 8.26E-05	
95% CL	0.033 -- 0.063	95% Confidence limits	6E-05 ---- 1.1E-04

Test Teratogenic Index (TI = LC50/EC50): 550.09

95% Confidence limits 354.20 -- 854.33

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. DEF 3

Test Material	P3C	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	12-13-92
Composition/Purity		Test End Date	12-17-92
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	6.7	6.7	6.6	6.6	
	Control		7.7	7.4	7.2	7.4
	Highest Concentration		7.3	7.4	7.3	7

No. Dead or Malformed	X 100 = %		Mortality Record		Malformation Record	
Total Number						
FETAX Control	7 : 100 X 100 =		7% 4 : 93 X 100 =		4.3%	
Solvent Control	C19 : E19 X 100 = G19		H19 : J19 X 100 = L19			
Control Length	C20		Solvent Control Length		J20	
Minimum Concentration to Inhibit Growth (MCIG)	0.00010					

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.075	N.A.	T-test
LOEL	0.1	1E-05	T-test
LC50	0.099	EC50 4.34E-05	
95% CL	0.081	0.121	95% Confidence limits 3.24E-05 --- 5.8E-05
Test Teratogenic Index (TI = LC50/EC50):			2281.47
95% Confidence limits			1602.02 -- 3249.09

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX SUMMARY SHEET

Test Material:	Unknown PBC	Test No.	3 - Def 1 JMB
Source:		Investigator:	Hull
CAS No.	Lot No.	Laboratory:	Banville OSU/GB
Composition / Purity:		Test Start Date:	12-17-92
Solvent:	Conc.	Test End Date:	12-21-92
		Test Units:(i.e., mg/ml)	mg/ml

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.6, 7.5	7.5, 7.7	8.1, 7.7	7.5, 7.6	X
Control	X	7.5	7.8	8.0	8.0
Highest Conc.	X	7.6	8.3	8.3	8.1

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	<u>10</u> : <u>100</u> X 100 = <u>10</u> %	<u>7</u> : <u>90</u> X 100 = <u>7.7</u> %
Solvent Control:	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
Control Length: mm	Solvent Control Length: mm	
Minimum Concentration to Inhibit Growth (MCIG):		0.0002

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.001	0.000075	Williams
LOEL	0.01	0.001	Williams
LC ₅₀	0.00948	EC ₅₀ 0.000206	
95% Confidence Limits	0.00167 - 0.0535	95% Confidence Limits	0.00017 - 0.00024
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀)		4.7777 16.02 A	

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2.5 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX Summary Sheet

Test No. 2 *JAHB*

Test Material	Unknown P3C	Investigator	Mendi A. Hull
Source	B4	Laboratory	Bantle / OSU
CAS No.	B5	Lot No. E5	Test Start Date: Feb 1 1993
Composition/Purity	C6	Test End Date	Feb 5 1993
Solvent	B7	Conc. E7	Test Units (i.e., mg/ml) mg/ml

		Day 0	Day 1	Day 2	Day 3	Day 4
pH	Stock	7.6, 7.6	7.7, 7.4	7.6, 7.2	7.6, 7.4	X
	Control	X	7.4	7.7	7.7	7.6
	Highest Concentration	X	7.2	7.7	7.8	7.5

No. Dead or Malformed	MALFORMATION EXCEED ASTM LIMITS			
Total Number	Mortality Record		Malformation Record	
FETAX Control	8 : 100	X 100 = 8%	9 : 92	X 100 = 9.8%
Solvent Control	C19 : E19	X 100 = G19	H19 : J19	X 100 = L19
Control Length	0.9	Solvent Control Length J20		
Minimum Concentration to Inhibit Growth (MCIG)	0.025			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.025	0.0003	T-test
LOEL	0.1	0.0005	T-test
LC50	6.54E-02	EC50 5.45E-04	
95% CL	1.97E-02 -- 2E-01	95% Confidence limits 0.0340E-04 ---- 0.001	
Test Teratogenic Index (TI = LC50/EC50):			1.20E+02
95% Confidence limits			3.31E+01 -- 4.4E+02

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. 3

Test Material	Unknown P3C	Investigator	Mendi A. Hull
Source	B4	Laboratory	Bantle / OSU <i>JMB</i>
CAS No.	B5	Lot No.	E5
Composition/Purity	C6	Test Start Date:	Mar 8 1993
Solvent	B7	Test End Date	Mar 12 1993
Conc.	E7	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.6	7.1	7.4	7.6	X
	Control	X	7.1	7.1	6.8	6.6
	Highest Concentration	X	7.5	7.4	7.0	7.0

No. Dead or Malformed	X 100 = %		
Total Number	Mortality Record		Malformation Record
FETAX Control	3 : 100	X 100 = 3%	4 : 97 X 100 = 4.1%
Solvent Control	C19 : E19	X 100 = G19	H19 : J19 X 100 = L19
Control Length	320 cm	Solvent Control Length	J20
Minimum Concentration to Inhibit Growth (MCIG)			0.01 mg/me

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.01	7.5E-05	T-test
LOEL	0.025	0.0001	T-test
LC50	0.013	EC50 1.51E-04	
95% CL	0.010 -- 0.018	95% Confidence limits	1E-04 ---- 1.8E-04
Test Teratogenic Index (TI = LC50/EC50):			88.57
95% Confidence limits			64.09 -- 122.40

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No. 1-P3C

Test Material	PAC	Investigator	FINCH
Source	OSU	Laboratory	USABERD
CAS No.	85	Lot No. E5	Test Start Date:
Composition/Purity	CB		26 JUN 93
Solvent	None	Test End Date:	2 JULY 93
		Test Units (i.e., mm/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	E11	F11	G11	J11	K11
	Control		F12	G12	J11	K12
	Highest Concentration		F13	G13	J11	K13

No. Dead or Malformed	X 100 = %		Mortality Record		Malformation Record			
	Total Number		0 : 100	X 100 =	0%	4 : 100	X 100 =	4.0%
FETAX Control								
Solvent Control	G19	E19	X 100 = G19		H19	J19	X 100 = L19	
Control Length	O20		Solvent Control Length		J20			
Minimum Concentration to Inhibit Growth (MCIG)					J21			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.0025	0.0001	T-test
LOEL	0.005	0.00025	T-test
LC50	0.00530	EC50 0.00028	
95% CL	0.00453	- 0.00620	95% Confidence limits
			0.00023 --- 0.00035
Test Teratogenic Index (TI = LC50/EC50):			18.65
95% Confidence limits			14.29 - 24.34

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
1.5 mg/L	E33 E33	X 100 = G33 H33 J33 X 100 = L33
2500 mg/L	E34 E34	X 100 = G34 H33 J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. 2-PSC

Test Material	P30	Investigator	FINCH
Source	OSU	Laboratory	USABRI
CAS No.	85	Lat. No.	E5
Composition/Purity	CD	Test Start Date	
Solvent	None	Test End Date	
		Test Units (L or mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	E11	E11	G11	J11	
	Control		E12	G12	J11	K12
	Highest Concentration		E13	B13	J11	K13

No. Dead or Malformed	X 100 = %		
Total Number			
FETAX Control	4 : 100	X 100 = 4%	6 : 96 X 100 = 6.3%
Solvent Control	C19 E19	X 100 = G19	H19 J19 X 100 = L19
Control Length	C20	Solvent Control Length	J20
Minimum Concentration to Inhibit Growth (MCIG)			J21

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.0025	0.0005	T-test
LOEL	0.005	0.0025	T-test
LC50	0.00288	EC50 0.00128	
95% CL	0.00237	-- 0.00350	95% Confidence limits 0.00106 --- 0.00154
Test Teratogenic Index (TI = LC50/EC50):			2.25
95% Confidence limits			1.72 -- 2.94

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	E33 E33 X 100 = G33	H33 J33 X 100 = L33
2500 mg/L	E34 E34 X 100 = G34	H33 J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No. 3-PJC

Test Material:	P2C	Investigator:	FINCH
Source:	CSU	Laboratory:	USABRD/L
CAS No.:	65-15-7	Test Start Date:	
Composition/Purity:	CB	Test End Date:	
Solvent:	Cover	Test Units (i.e., mg/ml):	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	E11	F11	G11	J11	
	Control		F12	G12	J11	K12
	Highest Concentration		F13	B13	J11	K13

No. Dead or Malformed	X 100 = %					
	Mortality Record			Malformation Record		
Total Number						
FETAX Control	3 : 100	X 100 =	3%	0 : 97	X 100 =	0.0%
Solvent Control	C19 E19	X 100 =	G19	H19 J19	X 100 =	L19
Control Length	C20	Solvent Control Length:		J20		
Minimum Concentration to Inhibit Growth (MCIG)				J21		

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.0025	0.0005	T-test
LOEL	0.005	0.00075	T-test
LC50	0.00427	EC50 0.00124	
95% CL	0.00378	- 0.00482	95% Confidence limits 0.00103 - 0.00149

Test Teratogenic Index (TI = LC50/EC50): 3.45

95% Confidence limits 2.76 - 4.30

POSTIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 E33 X 100 = G33	H33 J33 X 100 = L33
2500 mg/L	C34 E34 X 100 = G34	H33 J34 X 100 = L34

CL= Confidence limits

FETAX SUMMARY SHEET

Test Material: P3C-1		Test No. Definitive #1
Source: Bantle		Investigator: Gillett
CAS No. Lot No.		Laboratory: CERL
Composition / Purity:		Test Start Date: 09/27/93
Solvent:	Conc.	Test End Date: 10/01/93
		Test Units:(i.e., mg/ml) 1 mg/ml

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.0	7.0	7.1	7.0	
Control	X	7.5	7.6	7.3	
Highest Conc.	X	7.6	7.5	not taken	

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = X Total Number	: X 100 = <u>6</u> %	: X 100 = <u>2</u> %
Solvent Control:	: X 100 = <u> </u> %	: X 100 = <u> </u> %
Control Length: 9.5645 mm	Solvent Control Length: <u> </u> mm	
Minimum Concentration to Inhibit Growth (MCIG): 0.0005 mg/ml		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.0006		Williams
LOEL	0.0008		Williams
LC ₅₀	.0126	EC ₅₀ .00222	
95% Confidence Limits	.0111 - .0144	95% Confidence Limits	.00200 - .00247

$$\text{TEST TERATOGENIC INDEX (TI)} = \frac{\text{LC}_{50}}{\text{EC}_{50}} = \frac{.0126}{.00222} = 5.6757$$

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	: X 100 = <u> </u> %	: X 100 = <u> </u> %
2.5 mg/ml	: X 100 = <u> </u> %	: X 100 = <u> </u> %

FETAX SUMMARY SHEET

Test Material:	P3C-2	Test No.
Source:	Bantle	Investigator: G. Neff
CAS No.	Lot No.	Laboratory: CERL
Composition / Purity:		Test Start Date: 10/11/93
Solvent:	Conc.	Test End Date: 10/15/93
		Test Units:(i.e., mg/ml)

--pH-- Stock <i>7.04</i>	Day 1	Day 2	Day 3	Day 4	Day 5
	7.0	7.0	7.0	7.1	X
Control	X	7.6	7.4	7.2	7.5
Highest Conc.	X	7.5	7.7	7.7	7.6

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	____ : ____ X 100 = <u>3</u> %	____ : ____ X 100 = <u>4</u> %
Solvent Control:	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
Control Length: <u>0.9448</u> mm	Solvent Control Length:	mm
Minimum Concentration to Inhibit Growth (MCIG): <u>0.00075</u> mg/ml		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	.025	.005 Kruskal-Wallis	Williams
LOEL	.05	.01 Kruskal-Wallis	Williams
LC ₅₀	.105	EC ₅₀ .000631	
95% Confidence Limits	.090 - .123	95% Confidence Limits	.000412 - .000964
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) = <u>.1664</u> / <u>.000412</u> = <u>.400484</u>			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2.5 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

		Test No.	Acf. #3	
Test Material:	P3C		Investigator:	Gillatt/wilbom
Source:	Bantle		Laboratory:	ERL-C
CAS No.	Lot No.	Test Start Date: 10/22/93		
Composition / Purity:		Test End Date: 10/26/93		
Solvent:	Conc.	Test Units:(i.e., mg/ml)		

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.2	7.1	7.1	7.1	7.6
Control	—	Not taken	7.2	7.2	7.6
Highest Conc.	—	Not taken	7.8	7.7	7.6

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
Solvent Control:	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
Control Length: 10.0359 mm	Solvent Control Length: mm	
Minimum Concentration to Inhibit Growth (MCIG): 0.0001 mg/ml		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	.001	.0001	Williams
LOEL	.005	.0005	Williams
LC ₅₀	.004	EC ₅₀ .000589	
95% Confidence Limits	.00358 - .00638	95% Confidence Limits	.000467 - .000744
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) = 6.79			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2.5 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

Test No. 1

Test Material	Unknown P3C	Investigator	TURLEY
Source	OSU	Lab	U.M.D.-WREC
CAS No.	Lot No.	Test Start Date	2/2/93
Composition/Purity		Test End Date	2/6/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/L

— pH —	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	6.99	6.92	6.87	6.88	6.91
Control	7.67	7.69	7.74	7.70	7.67
Highest Conc.	7.56	7.51	7.49	7.52	7.55

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		4 : 100 X 100 = 4 %	5 : 100 X 100 = 5 %
Solvent Control		_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
Control Length 10.01 mm	Solvent Control Length mm		
Minimum Concentration to Inhibit Growth (MCIG) 1.0 mg/L			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	4.0	0.50	Bonferroni T-test
LOEL	5.0	0.60	Bonferroni T-test
LC ₅₀ 0.549 mg/L Trimmed Spearman-Karber	EC ₅₀ 0.67 mg/L Trimmed Spearman-Karber		
95% Confidence limits 5.04 - 5.98	95% Confidence Limits 0.59 - 0.75		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) 8.19			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

Test No. 2

Test Material	Unknown P3C	Investigator	TURLEY
Source	O.S.U	Lab	U.MD-WREC
CAS No.	Lot No.	Test Start Date	2/19/93
Composition/Purity		Test End Date	2/13/93
Solvent	None	Test Units (i.e., mg/L)	mg/L

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	6.99	6.90	6.89	6.94	6.98
Control	7.71	7.70	7.68	7.72	7.68
Highest Conc.	7.60	7.58	7.56	7.48	7.56
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %				
Total Number	2 : 100 X 100 = 2 %	I : 100 X 100 = 7 %			
Solvent Control					
Control Length 10.07 mm	Solvent Control Length	mm			
Minimum Concentration to Inhibit Growth (MCIG) 2.00 mg/L					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	4.00 mg/L	0.25 mg/L	Bonferroni T-test
LOEL	5.00 mg/L	0.50 mg/L	Bonferroni T-test
LC ₅₀	4.70 (Trimmed Spearman-Karber)	EC ₅₀ 0.66 mg/L (Trimmed Spearman-Karber)	
95% Confidence limits	4.49 - 4.91	95% Confidence Limits (0.57 - 0.77)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			7.12

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2500 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

Test No. 3

Test Material	Unknown P3C	Investigator	TURLEY
Source	O.S.U	Lab	U.MD-WREC
CAS No.	Lot No.	Test Start Date	2/20/93
Composition/Purity		Test End Date	2/24/93
Solvent	Conc.	Test Units (l.e., mg/ml)	mg/L

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	6.94	6.87	6.92	6.90	6.98
Control	7.68	7.69	7.65	7.63	7.72
Highest Conc.	7.51	7.48	7.49	7.51	7.53

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	3 : 100 X 100 = 3 %	7 : 100 X 100 = 7 %
Solvent Control	: X 100 = %	: X 100 = %
Control Length 10.09 mm	Solvent Control Length	mm
Minimum Concentration to Inhibit Growth (MCIG) 1.00 mg/L		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LOEL	FAIL HOMOGENEITY	FAIL HOMOGENEITY	Bonferroni T-test
LC ₅₀ 4.46 mg/L Truncated Spearman-Karber	EC ₅₀ 0.74 mg/L Truncated Spearman-Karber		
95% Confidence limits 4.08 - 4.89	95% Confidence Limits 0.63 - 0.88		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) 6.03			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

		Test No. /
Test Material P3C		Investigator J.G. DAWSON
Source	Lab AT/CV4	
CAS No.	Lot No.	Test Start Date 12/18/92
Composition/Purity		Test End Date 12/12/92
Solvent	Conc.	Test Units (l.e., mg/ml) 1/10,000

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.09	7.11	7.09	7.09	-
Control	7.91	7.67	7.64	7.62	7.63
Highest Conc.	-	7.80	7.84	7.84	7.83

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		0 : 100 X 100 = 0 %	3 : 100 X 100 = 3 %
Solvent Control		: X 100 = %	: X 100 = %
Control Length mm		Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG) 0.0001			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	< 0.006 btl	0.0001 btl	
LOEL	0.006 btl	0.0002 btl	
LC ₅₀	0.0077 TSK	EC ₅₀ 0.00021 LHW	0.00018 TSK
95% Confidence limits (0.0073 - 0.0082) TSK	95% Confidence Limits (0.00008 - 0.0005)	(0.00008 - 0.0005)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		42.8 - TSK - EC ₅₀	36.7 - LHW - EC ₅₀

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

J.A.I.
12/14/92

FETAX SUMMARY SHEET

Test No. 2

Test Material	PSC	Investigator	<i>DR. ANSARI</i>
Source		Lab	UTICVM
CAS No.	Lot No.	Test Start Date	1/4/93
Composition/Purity		Test End Date	1/8/93
Solvent	Conc.	Test Units (i.e., mg/mL)	<i>xx/mi</i>

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.07	7.05	7.12	7.04	—
Control	7.79	7.75	7.51	7.55	7.52
Highest Conc.	—	7.86	7.76	7.69	7.61
FETAX CONTROL					
No. Dead or Malformed	MORTALITY RECORD		MALFORMATION RECORD		
Total Number	X 100 = %	0 : 100 X 100 = 0 %	2 : 100 X 100 = 2 %		
Solvent Control		: X 100 = %	: X 100 = %		
Control Length mm	Solvent Control Length mm				
Minimum Concentration to Inhibit Growth (MCIG) <i>0.00015</i>					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.04 <i>b2d</i>	0.0001 <i>b2d</i>	
LOEL	0.06	0.00015 <i>b2d</i>	
LC ₅₀	0.0074	<i>76%</i>	EC ₅₀ 0.000218 <i>LEW</i>
95% Confidence limits (0.0070 - 0.0077)	<i>76%</i>		95% Confidence Limits (0.00019 - 0.00025)
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			33.9

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L <i>Not</i>	: X 100 = %	: X 100 = %
2500 mg/L <i>100%</i>	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

Test No. 3

Test Material	P3C	Investigator	D. A. DAWSON
Source		Lab	UT/CVM
CAS No.	Lot No.	Test Start Date	11/11/93
Composition/Purity		Test End Date	11/15/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.08	7.09	7.05	7.07	—
Control	7.82	7.72	7.61	7.59	7.51
Highest Conc.	—	7.81	7.75	7.67	7.27

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed		
Total Number	X 100 = %	
Solvent Control	1 : 100 X 100 = 1 %	1 : 99 X 100 = 1.01 %
Control Length mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (NCIG)	0.02% ²⁵	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.006 6-AN	<0.0001 NIL	
LOEL	0.018 6-AN	0.0001 NIL	
LC ₅₀	0.011-15K 0.017-EP4	EC ₅₀ 0.000162-LFW 0.00017-TSK	0.000165-EP7
95% Confidence Limits	(0.0091-0.012)-TSK (0.011-0.018)-EP4	95% Confidence Limits (0.00014-0.00018)-LFW (0.00015-0.00018)-TSK	(0.00015-0.00018)
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		64.7-TSK 52.7-EP4	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L Not	— : — X 100 = — %	— : — X 100 = — %
2500 mg/L clear	— : — X 100 = — %	— : — X 100 = — %

D.A.D.
11/15/93

FETAX SUMMARY SHEET

Test No. 1

DF

Test Material	P3C	Investigator	FORT
Source	ILS, Phase 3	Lab	SBL
CAS No.	Lot No.	Test Start Date	12/14/92
Composition/Purity		Test End Date	12/18/92
Solvent	Cone.	Test Units (i.e., mg/ml)	

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.5	7.5	7.4	7.5	
Control	8.0	8.0	7.9	7.9	
Highest Cone.	7.6	7.7	7.7	7.7	

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		0 : 0 X 100 = 0 %	0 : 0 X 100 = 0 %
Solvent Control	—	— : — X 100 = — %	— : — X 100 = — %
Control Length 95.0 mm	Solvent Control Length — mm		
Minimum Concentration to Inhibit Growth (MCIG) 0.001			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.005	0	Williams
LOEL	0.01	0.0005	Williams
LC ₅₀	0.06	EC ₅₀	0.0011
95% Confidence limits	0.05 - 0.07	95% Confidence Limits 0.0009 - 0.0015	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) 54.5			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	— : — X 100 = — %	— : — X 100 = — %
2500 mg/L	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

Test No. 2

Test Material	P3C	Investigator	Fort
Source	FETAX ILS	Lab	SBL
CAS No.	Lot No.	Test Start Date	1/21/93
Composition/Purity		Test End Date	1/25/93
Solvent	Conc.	Test Units (i.e., mg/ml)	Mg/L

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.2	7.3	7.4	7.4	
Control	7.8	7.8	7.9	7.9	
Highest Conc.	7.5	7.5	7.5	7.6	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	0 : 100 X 100 = 0 %	0 : 100 X 100 = 0 %
Solvent Control	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
Control Length 90.3 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG) 0.0005		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.005	0.0005	Williams
LOEL	0.05	0.00075	Williams
LC ₅₀	0.076	EC ₅₀ 0.0012	
95% Confidence limits	0.063 - 0.092	95% Confidence Limits 0.0010 - 0.0014	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) 63.33			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

Test No. 3

Test Material	P3C	Investigator	Fort
Source	FETAX ILS	Lab	SBL
CAS No.		Test Start Date	1/23/93
Composition/Purity		Test End Date	1/27/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.3	7.2	7.3	7.4	
Control	7.9	8.0	7.9	7.9	
Highest Conc.	7.7	7.8	7.9	7.7	

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		0 : 100 X 100 = 0 %	0 : 100 X 100 = 0 %
Solvent Control		0 : X 100 = 0 %	0 : X 100 = 0 %
Control Length 91.6 mm		Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG)		0.0005	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.005	0.0005	Williams Test
LOEL	0.05	0.00075	Williams Test
LC ₅₀	0.095	EC ₅₀ 0.0009	
95% Confidence Limits	0.083 - 0.107	95% Confidence Limits	0.0008 - 0.0011
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			105.56

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	0 : X 100 = 0 %	0 : X 100 = 0 %
2500 mg/L	0 : X 100 = 0 %	0 : X 100 = 0 %

SODIUM ACETATE

PHASE III COMPOUND D DATA (P3D) SODIUM ACETATE

FETAX ILS STUDY; 7 LABS, 8 LEVELS, 3 REPLICATES
PAGE 1
ORIGINAL DATA TABLE WITH MATERIALS

units in mg/ml

ORIGINAL DATA TABLE WITH MATERIALS										SUBTOTALS:									
		LC50	C1LO	C1HI	EC50	C1LO	C1HI	MCG	Tl	LOEC/MORT	LOEC/MALF	J						STD:	CV:
BANTLE/ RAYBURN	LAB	A	5.34	9.07	2.32	1.32	4.10	1.00	3.0	12.00	1.00		LC50	6.21	0.657	10.6			
	1	6.96	5.34	9.07	2.32	0.85	1.63	0.50	9.4	12.00	0.20		EC50	2.37	0.801	33.8			
	2	6.30	5.29	7.49	1.41	0.65	3.05	6.00	4.5	0.25			MCG	3.00	2.160	72.0			
BANTLE/ HULL	3	5.36	4.62	6.21	3.37	2.57	4.43	2.00	1.6	4.00	4.00		Tl	2.62	1.184	45.2			
	1	2	11.20	9.52	13.17	2.01	1.08	3.73	0.50	5.6	8.00	0.50	LC50	10.96	0.249	2.3			
	2	11.07	10.16	12.06	1.18	0.85	1.17	0.84	1.62	0.50	9.1	12.00	0.25	EC50	1.45	0.394	27.1		
FINCH	3	10.62	10.14	11.13	1.17	0.84							MCG	0.50	0.000	0.0			
	1	3	12.00	10.84	13.29	5.12	4.14	6.33	10.00	2.3	14.00	4.00	LC50	11.69	0.486	4.2			
	2	11.00	9.27	13.06	7.64	7.03	8.30	9.00	1.4	12.00	9.00	0.00	EC50	7.21	1.564	21.7			
LINDER/ BUCHWALTER	3	12.06	11.42	12.74	8.88	8.26	9.55	9.00	1.4	11.00	8.00	0.25	MCG	9.33	0.471	5.1			
	1	4	11.69	10.98	12.44	5.23	4.37	6.25	1.00	2.2	13.00	1.00	LC50	9.72	1.692	17.4			
	2	9.92	9.49	10.38	6.62	6.03	7.26	1.00	1.5	9.00	5.00	0.25	EC50	6.12	0.633	10.3			
BURTON/ TURLEY	3	7.56	7.18	7.97	6.52	5.90	7.21	3.00	1.5	9.00	0.25		MCG	1.67	0.943	56.6			
	1	5	7.76	7.49	8.04	7.92	7.28	8.61	8.00	1.0	7.00		Tl	1.59	0.330	20.8			
	2	7.67	7.23	8.13	8.26	7.84	8.71	8.00	0.9	7.00	6.00	0.25	LC50	7.05	0.941	13.3			
DAVISON	3	5.72	5.42	6.02	6.31	5.83	6.83	7.00	0.9	4.00	6.00	0.25	EC50	7.50	0.851	11.3			
	1	6	12.32	11.90	12.80	8.77	8.20	9.40	5.00	1.4	7.00	7.00	0.05	MCG	7.67	0.471	6.1		
	2	11.68	10.30	13.20	9.23	7.10	12.10	5.00	1.3	10.00	5.00	0.00	LC50	7.23	0.262	2.2			
STOVER/ FORT	3	11.96	11.70	12.30	8.21	7.80	8.60	7.00	1.5	10.00	7.00	0.00	EC50	8.74	0.417	4.8			
	1	7	8.91	7.68	10.35	1.55	1.20	1.99	0.50	5.8	5.00	0.05	MCG	5.67	0.943	16.6			
	2	7.31	6.39	8.36	2.30	1.96	2.69	7.50	3.2	2.50	1.00	0.00	LC50	1.64	0.502	30.6			
GRAND TOTAL	3	5.46	4.63	6.43	1.08	0.99	1.19	2.50	5.1	2.50	1.00	0.00	EC50	3.50	2.944	84.1			
													Tl	4.40	1.098	25.0			
													CV:						

Table 4. Results of the Interlaboratory Study for Sodium Acetate.



Laboratory	Replicate (n)	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV1 (%)	EC50 (mg/ml)	Mean EC50 (mg/ml)	EC50 CV (%)	Mean TI ² (%)	TI CV (%)	MCIG ³ (mg/ml)	Mean MCIG (mg/ml)	MCIG CV (%)
1	1	6.96	6.21	10.6	2.32	1.41	2.37	33.8	2.6	44.8	6.0	3.00
	2	6.30	5.36	3.37							2.0	72.0
	3											
2	1	11.20	11.07	10.96	2.3	1.18	2.01	1.45	27.1	7.5	22.9	0.5
	2										0.5	0.0
	3	10.62			1.17							
3	1	12.00	11.00	11.69	4.2	5.12	7.64	7.21	21.7	1.6	27.6	10.0
	2										9.0	9.33
	3	12.06			8.88							5.1
4	1	11.69	9.92	9.72	17.4	5.23	6.62	6.12	10.3	1.6	28.3	1.0
	2										1.0	1.67
	3	7.56	7.56			6.52					3.0	56.6
5	1	7.76	7.67	7.05	13.3	7.92	8.26	7.50	11.3	0.9	3.3	8.0
	2										8.0	7.67
	3	5.72				6.31					7.0	6.1
6	1	12.32	11.68	11.99	2.2	8.77	9.23	8.74	4.8	1.4	5.9	5.0
	2										5.0	5.67
	3	11.96						8.21			7.0	
7	1	8.91	7.31	7.23	19.5	1.55	2.30	1.64	30.6	4.4	24.7	0.50
	2										7.50	3.50
	3	5.46							1.08		2.50	84.1

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

ASTM Analysis of Sodium Acetate



FETAX Summary Sheet

Test No. DEF 1 *JRK*

Test Material	P3D	Investigator	JAMES RAYBURN	
Source	OAK RIDGE	Laboratory	BANTLE	
CAS No.	Lot No.	Test Start Date:	3-25-93	
Composition/Purity		Test End Date	3-29-93	
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML	

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.9	7.9	7.8	7.8	
	Control		7.8	7.5	7.3	7.1
	Highest Concentration		7.6	7.3	7.2	7.1

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		9 : 100	X 100 =	9%	6 : 91	X 100 =	6.6%
Solvent Control		:	X 100 =	:	X 100 =		
Control Length	20 9.18		Solvent Control Length				
Minimum Concentration to Inhibit Growth (MCIG)				1 mg/ml			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	6	0.75	T-test
LOEL	(12)	1	T-test
LC50	6.961	EC50 2.323	
95% CL	5.340	9.074	95% Confidence limits 1.316 ---- 4.101

Test Teratogenic Index (TI = LC50/EC50):	3.00
95% Confidence limits	1.60 -- 5.61

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No. DEF 2 *MHD*

Test Material	P3D	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	3-29-93
Composition/Purity		Test End Date	4-2-93
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.8	7.8	7.7	7.7	
	Control		7.7	7.5	7.4	7.2
	Highest Concentration		7.7	7.5	7.3	7.1

No. Dead or Malformed	MALFORMATION EXCEED ASTM LIMIT				
X 100 = %	Mortality Record		Malformation Record		
Total Number					
FETAX Control	10 : 100	X 100 = 10%	7 : 90	X 100 = 7.8%	
Solvent Control	:	X 100 =	:	X 100 =	
Control Length	9.688 mg/ml	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)			6 mg/ml		

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	N.A.	1	T-test
LOEL	0.25	2	T-test
LC50	6.296	EC50 1.409	
95% CL	5.293 -- 7.489	95% Confidence limits	0.649 --- 3.059

Test Teratogenic Index (TI = LC50/EC50): 4.47

95% Confidence limits 2.02 -- 9.89

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. DEF 3

Test Material	P3D	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	4-19-93
Composition/Purity		Test End Date	4-23-93
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.8	7.8	7.7	7.7	
	Control		7.7	7.5	7.4	7.2
	Highest Concentration		7.7	7.5	7.3	7.1

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		8 : 100 X 100 = 8%		9 : 92 X 100 = 9.8%			
Solvent Control		: X 100 =		: X 100 =			
Control Length		9.23		Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)				2 / mg/ml			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used				
NOEL	2	2	T-test				
LOEL	4	4	T-test				
LC50	5.356		EC50	3.374			
95% CL	4.617	--	6.213	95% Confidence Limits	2.570	----	4.429
	Test Teratogenic Index (TI = LC50/EC50):				1.59		
	95% Confidence Limits				1.16	--	2.16

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No. 1

Test Material	P3D	Investigator	Mendi A. Hull
Source	Sigma	Laboratory	OSU Bantle
CAS No.	Lot No.	Test Start Date:	8 Mar 93
Composition/Purity		Test End Date	12 Mar 93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.1	8	8.2	8	
	Control		7.6	7.4	7	7.1
	Highest Concentration		7.8	7.5	7	8

No. Dead or Malformed	X 100 = %	Mortality Record		Malformation Record	
Total Number					
FETAX Control	3 : 100	X 100 = 3%	7 : 97	X 100 = 7.2%	
Solvent Control	:	X 100 =	:	X 100 =	
Control Length (mm)	1 cm	Solvent Control Length (min)			
Minimum Concentration to Inhibit Growth (MCI)	0.5				mg/ml

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	6	0.25	T-test
LOEL	8	0.5	T-test
LC50	11.197	EC50 2.006	
95% CL	9.517 -- 13.173	95% Confidence limits 1.078 ---- 3.730	
Test Teratogenic Index (TI = LC50/EC50):			5.58
95% Confidence limits			2.94 -- 10.60

FETAX Summary Sheet

Test No. 2

Test Material	P3D	Investigator	Mendi A. Hull
Source		Laboratory	Bantle / OSU
CAS No.	Lot No.	Test Start Date:	Mar 15 1993
Composition/Purity		Test End Date	Mar 19 1993
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.1	8.0	8.2	8.0	X
	Control	X	7.6	7.4	7.0	7.1
	Highest Concentration	X	7.8	7.5	7.0	8.0

No. Dead or Malformed	X 100 = %	
Total Number	Mortality Record	Malformation Record
FETAX Control	10 : 100 X 100 = 10%	6 : 90 X 100 = 6.7%
Solvent Control	: X 100 =	: X 100 =
Control Length	1 cm	Solvent Control Length
Minimum Concentration to Inhibit Growth (MCIG)	0.5 mg/ml	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	10	N.A.	T-test
LOEL	12	0.25	T-test
LC50	11.071	EC50 1.176	
95% CL	10.161 -- 12.063	95% Confidence limits	0.849 ---- 1.630
Test Teratogenic Index (TI = LC50/EC50):			9.41
95% Confidence limits			6.72 -- 13.19

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

FETAX Summary Sheet

Test No.	3
Test Material	P3D
Source	Laboratory
CAS No.	Lot No.
Composition/Purity	Test Start Date:
Solvent	Conc.
	Test Units (i.e., mg/ml)
	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.1	8.0	8.2	8.0	X
	Control	X	7.6	7.4	7.0	7.1
	Highest Concentration	X	7.8	7.5	7.0	8.0

No. Dead or Malformed		Mortality Record		Malformation Record	
Total Number	X 100 = %				
FETAX Control	2 : 100	X 100 =	2%	7 : 98	X 100 = 7.1%
Solvent Control	:	X 100 =		:	X 100 =
Control Length	1 cm	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)		0.5 mg/ml			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	10	0.5	T-test
LOEL	12	0.75	T-test
LC50	10.623	EC50 1.169	
95% CL	10.139 -- 11.130	95% Confidence limits 0.842 --- 1.623	
	Test Teratogenic Index (TI = LC50/EC50):		9.09
	95% Confidence limits 6.53 -- 12.66		

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	: X 100 =	: X 100 =
2500 mg/L	: X 100 =	: X 100 =

CL = Confidence limits

FETAX Summary Sheet

Test No.: 1(200-01)

Test Material	P3D	Investigator	FINCH
Source	OSU	Laboratory	USABRDE
CAS No.	Lot No.	Test Start Date	21 JUN 93
Composition/Purity		Test End Date	25 JUN 93
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.6	8.6	8.6	8.6	8.6
	Control		7.7	7.7	7.8	7.7
	Highest Concentration		8.2	8.1	8.2	8.1

No. Dead or Malformed		X 100 = %	
Total Number		Mortality Record	Malformation Record
FETAX Control	1 : 100	X 100 = ✓ 1%	5 : 99 X 100 = 5.1%
Solvent Control		X 100 = .	X 100 = .
Control Length	9.567	Solvent Control Length	.
Minimum Concentration to Inhibit Growth (MCIG)			10.0 ✓

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	12	2	T-test
LOEL	14	4	T-test
LC50	12.000 ✓ (?)	EC50 5.116 (?)	
95% CL	10.835 -- 13.290	95% Confidence Limits 4.139 --- 6.325	
	Test Teratogenic Index (TI = LC50/EC50): 1.17 (?) 2.35		
	95% Confidence Limits 1.85 -- 2.97		

CL= Confidence limits

FETAX Summary Sheet

Test No: 2(200-02)

Test Material:	P3D	Investigator:	FINCH
Source:	OSU	Laboratory:	USABRDL
CAS No.:	-	Test Start Date:	28 JUN 93
Composition/Purity:	-	Test End Date:	2 JULY 93
Solvent:	-	Test Units (i.e., mg/ml):	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
		Stock	8.6	8.6	8.7	8.7
	Control			8.0	7.9	7.7
	Highest Concentration		8.2	8.2	8.4	8.3

No. Dead or Malformed	X 100 = %	
Total Number	Mortality Record	Malformation Record
FETAX Control	1 : 100 X 100 = ✓ 1%	5 : 99 X 100 = 5.1%
Solvent Control	X 100 = -	X 100 = -
Control Length	9.361	Solvent Control Length -
Minimum Concentration to Inhibit Growth (MCIG)		9.0 ✓

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	11	6	T-test
LOEL	12	9	T-test
LC50	11.000 ✓	EC50 7.640 ~8.3 (?)	~1.3 (?) 1.44
95% CL	9.269 - 13.055	95% Confidence Limits 7.033 --- 8.300	
	Test Teratogenic Index (TI = LC50/EC50):	~1.3 (?)	1.44
	95% Confidence Limits	1.19 -	1.74

CL= Confidence limits

FETAX Summary Sheet

Test No. 3(200-03)

Test Material	P3D	Investigator	FINCH
Source	CSU	Laboratory	USABRDL
CAS No.	65	Lot No.	E5
Composition/Purity	CP	Test Start Date:	28 JUN 93
Solvent	COP	Test End Date:	2 JULY 93
		Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	E11	F11	G11	J11	
	Control		F12	G12	J11	K12
	Highest Concentration		I13	B13	J11	K13

No. Dead or Malformed	X 100 = %		
Total Number			
FETAX Control	1 : 100	X 100 = 1%	2 : 99 X 100 = 2.0%
Solvent Control	C19 : E19	X 100 = G19	H19 : J19 X 100 = I.10
Control Length	C20	Solvent Control Length	J20
Minimum Concentration to Inhibit Growth (MCIG)			J21 q.D

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	10	5	T-test
LOEL	11	8	T-test
LC50	12.061	EC50 8.882	
95% CL	11.418 -- 12.740	95% Confidence limits	8.260 ---- 9.551
Test Teratogenic Index (TI = LC50/EC50):			1.36
95% Confidence limits			1.24 -- 1.49

POSTIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL = Confidence limits

CS8 CT8
CS9 CT9

FETAX SUMMARY SHEET

giant against

Test Material:	P3D	Test No.	Definitive #1
Source:	Bantle	Investigator:	Gillett
CAS No.	Lot No.	Laboratory:	CERL
Composition / Purity:		Test Start Date:	11/01/93
Solvent:	Conc.	Test End Date:	11/05/93
		Test Units:(i.e., mg/ml)	mg/ml

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.0	7.0	7.0	7.1	X
Control	X	7.5	7.7	8.0	7.8
Highest Conc.	X	7.6	7.6	7.8	7.7

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	X 100 = %	X 100 = %
Solvent Control:	X 100 = %	X 100 = %
Control Length: 9.0739 mm	Solvent Control Length:	mm
Minimum Concentration to Inhibit Growth (MCIG):	1.0 mg/ml	

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	11.0	0.5	Williams
LOEL	13.0	1.0	Williams
LC ₅₀	11.69	EC ₅₀ 5.23	
95% Confidence Limits	10.18-12.44	95% Confidence Limits	4.37-6.25
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) - 2.23			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	X 100 = %	X 100 = %
2.5 mg/ml	X 100 = %	X 100 = %

FETAX SUMMARY SHEET

Test Material:	P3D	Test No.	Def. #4
Source:	Bantle	Investigator:	Gillet
CAS No.	Lot No.	Laboratory:	CERL
Composition / Purity:		Test Start Date:	12/09/93
Solvent:	Conc.	Test End Date:	12/13/93
		Test Units:(i.e., mg/ml)	mg/ml

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.0	7.2	7.1	7.1	—
Control	7.4	7.4	7.3	7.5	7.5
Highest Conc.	7.5	7.4	7.7	7.4	—

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
Solvent Control:	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
Control Length: 9.7844 mm	Solvent Control Length:	mm
Minimum Concentration to Inhibit Growth (MCIG):	1 mg/ml	

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	7.0	3.0	Williams
LOEL	9.0	5.0	Williams
LC ₅₀	9.92	EC ₅₀ 6.62	
95% Confidence Limits	9.49 - 10.38	95% Confidence Limits	6.03 - 7.26
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) - 1.498			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2.5 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

		Test No. Definitive # 3
Test Material:	P3D	Investigator: Gillett/W./born
Source:	Bantle	Laboratory: C-ERL
CAS No.	Lot No.	Test Start Date: 11/19/93
Composition / Purity:		Test End Date: 11/23/93
Solvent:	Conc.	Test Units:(i.e., mg/ml) mg/ml

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.0	6.94	7.0	7.0	7.0
Control	X	7.42	7.5	7.4	7.6
Highest Conc.	X	7.87	7.8	7.7	7.7

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	____ : ____ X 100 = 0 %	____ : ____ X 100 = 0 %
Solvent Control:	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
Control Length: ^{10.09} _{10.09} mm	Solvent Control Length:	mm
Minimum Concentration to Inhibit Growth (MCIG):		3.0 mg/ml

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	7.0	Less than 0.25	Williams
LOEL	9.0	0.25	Williams
LC ₅₀	7.56	EC ₅₀ 6.52	
95% Confidence Limits	7.18 - 7.97	95% Confidence Limits	5.9 - 7.21
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) - 1.159			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2.5 mg/ml	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

7/10
Test No. 3

Test Material UNKNOWN P3D		Investigator TURLEY
Source OSU		Lab UMD/WREC
CAS No.	Lot No.	Test Start Date 4/17/93
Composition/Purity		Test End Date 4/21/93
Solvent	Conc.	Test Units (i.e., mg/ml) mg/ml

		DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH						
Stock	40 mg/ml	8.53	8.41	8.55	8.60	8.51
Control		7.77	7.69	7.70	7.67	7.63
Highest Conc.		8.00	7.97	8.11	8.05	7.99
FETAX CONTROL						
No. Dead or Malformed	X 100 = %	MORTALITY RECORD		MALFORMATION RECORD		
Total Number		1 : 100 X 100 = 1 %		6 : 99 X 100 = 6 %		
Solvent Control		_____ : _____ X 100 = _____ %		_____ : _____ X 100 = _____ %		
Control Length 9.98 mm		Solvent Control Length		mm		
Minimum Concentration to Inhibit Growth (MCIG) 8.0 mg/ml						

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	6.0	FAIL HOMOGENEITY	Bonferroni T-test
LOEL	7.0	FAIL HOMOGENEITY	Bonferroni T-test
LC ₅₀	7.76 (TTC Truncated Spearman-Karber)	EC ₅₀ 7.92 mg/ml Truncated Spearman-Karber	
95% Confidence limits	7.49 - 8.04	95% Confidence Limits 7.28 - 8.61	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			0.98

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

PB

Test Material UNKNOWN BY		Test No. 4			
Source OSU	Investigator TURLEY				
CAS No.	Lab UMD/WREC				
Composition/Purity		Test Start Date 4/17/93			
Solvent	Conc.	Test End Date 4/21/93			
		Test Units (l.p., mg/ml) mg/ml			
— pH —	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	8.53	8.41	8.55	8.60	8.51
Control	7.77	7.69	7.70	7.67	7.63
Highest Conc.	8.00	7.97	8.11	8.05	7.99
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %	1 : <u>49</u> X 100 = <u>1</u> %	6 : <u>59</u> X 100 = <u>6</u> %		
Total Number					
Solvent Control	: X 100 = %	: X 100 = %	: X 100 = %		
Control Length 10.08 mm	Solvent Control Length			mm	
Minimum Concentration to Inhibit Growth (MCIG) 8.0 mg/ml					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	5.0	6.0	Bonferroni T-test
LOEL	6.0	7.0	Bonferroni T-test
LC ₅₀ 7.67 mg/ml Trimmed Spearman-Kaber	EC ₅₀ 8.26 mg/ml Trimmed Spearman-Kafer		
95% Confidence Limits 7.23 - 8.23	95% Confidence Limits 7.84 - 8.71		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		0.93	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

DTS

Test No. 2

Test Material	UNKNOWN P3D	Investigator	TURLEY
Source	OSU	Lab	UMD/WREC
CAS No.	Lot No.	Test Start Date	3/13/93
Composition/Purity		Test End Date	4/3/93
Solvent	Cone.	Test Units (i.e., mg/ml)	mg/ml

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH					
Stock 40 mg/l	8.48	8.41	8.51	8.43	8.55
Control	7.65	7.71	7.60	7.59	7.63
Highest Cone.	8.01	8.09	7.97	7.99	8.04

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		1 : 100 X 100 = 1 %	68 : 99 X 100 = 68 %
Solvent Control		_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
Control Length 9.79 mm	Solvent Control Length mm		
Minimum Concentration to Inhibit Growth (MCIG) 7.00			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	3.0	5.0	BonFerroni T-test
LOEL	4.0	6.0	BonFerroni T-test
LC ₅₀ 5.72 mg/ml Truncated Spearman-Kaber	EC ₅₀ 6.31 mg/ml Truncated Spearman-Kaber		
95% Confidence limits 5.42-6.02	95% Confidence Limits 5.83-6.83		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			0.91

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

		Test No.
Test Material	P3D	Investigator D.A. DAWSON
Source		Lab UT/CVM
CAS No.	Lot No.	Test Start Date 5/12/93
Composition/Purity		Test End Date 5/16/93
Solvent	Conc.	Test Units (i.e., mg/ml) 1mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.44	7.41	7.44	7.43	—
Control		7.69	7.60	7.60	7.52
Highest Conc.	—	7.92 (9)	7.89 (7)	7.94 (7)	7.91 (7)

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	<u>0</u> : <u>100</u> X 100 = <u>0</u> %	<u>3</u> : <u>100</u> X 100 = <u>3</u> %
Solvent Control	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
Control Length mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG) 5.		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	5.0 mill.	5.0 B/F	
LOEL	7.0 mill.	7.0 B/F	
LC ₅₀	12.32	EC ₅₀ 8.77	
95% Confidence Limits (11.7-12.8)		95% Confidence Limits (8.2-9.4)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			(12.32) / 8.77 = 1.40

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

D.A.D.

FETAX SUMMARY SHEET

Test No. 2

Test Material	P3D	Investigator	J.A. DAWSON
Source		Lab	UT/CMV
CAS No.		Test Start Date	5/17/93
Composition/Purity		Test End Date	5/21/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.41 ad.	7.44 ad.	7.42 ad.	7.45 ad.	—
Control	7.59	7.56	7.62	7.54	7.47
Highest Conc.	—	7.93	7.88	7.99	7.97

FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %				
Total Number		0 : 100 X 100 = 0 %		0 : 100 X 100 = 0 %	
Solvent Control		0 : 100 X 100 = 0 %		0 : 100 X 100 = 0 %	
Control Length mm		Solvent Control Length mm			
Minimum Concentration to Inhibit Growth (MCIG) 5.					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	8.5 birth	<5 week.	
LOEL	10 birth	5 week.	
LC ₅₀	11.68	EC ₅₀ 9.23	
95% Confidence Limits (10.3 - 13.2)		95% Confidence Limits (7.1 - 12.1)	

$$\text{TEST TERATOGENIC INDEX (TI)} = \text{LC}_{50} / \text{EC}_{50} = 1.27$$

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2500 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

Test No. 3

Test Material	P3D	Investigator	D.A. Dawson
Source		Lab	UT/CDM
CAS No.		Test Start Date	5/18/93
Composition/Purity		Test End Date	5/22/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/lst

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.41 H ₂ O	7.41 rd.	7.46 adj.	7.44 adj.	-
Control	7.56	259	7.47	7.46	7.42
Highest Conc.	-	1.18	7.74	8.04	7.93
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %	0 : 100 X 100 = 0 %	1 : 100 X 100 = 1 %		
Total Number					
Solvent Control					
Control Length mm		Solvent Control Length mm			
Minimum Concentration to Inhibit Growth (MCIG) 7.					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	8.5 will.	5 Both	
LOEL	10 will.	7 Both	
LC ₅₀ 11.96		EC ₅₀ 8.21 Q	
95% Confidence Limits (11.7 - 12.3)		95% Confidence Limits (7.8 - 8.6)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.46

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

J.A.D.

5/22/93

FETAX SUMMARY SHEET

		Test No.				
Test Material	P3 D		Investigator	Fort		
Source	QA/QC Corrections		Lab	SBL		
CAS No.			Test Start Date	3/29/93		
Composition/Purity				Test End Date	4/2/93	
Solvent	Conc.	Test Units (i.e., mg/ml)				
		DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH	Stock	7.5	7.4	7.5	7.5	
	Control	7.9	7.9	8.0	7.9	
	Highest Conc.	7.6	7.6	7.7	7.7	
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD		
No. Dead or Malformed	X 100 = %	Total Number	<u>0</u> : <u>0</u> X 100 = <u>0</u> %	<u>1</u> : <u>100</u> X 100 = <u>1</u> %		
Solvent Control			<u> </u> : <u> </u> X 100 = <u> </u> %	<u> </u> : <u> </u> X 100 = <u> </u> %		
Control Length 96.1 mm	Solvent Control Length		mm			
Minimum Concentration to Inhibit Growth (MCIG)		0.5				

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.0	0.01	Williams
LOEL	5.0	0.05	
LC ₅₀	8.91	EC ₅₀ 1.55	
95% Confidence Limits 7.68 - 10.35		95% Confidence Limits 1.20 - 1.99	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) <u> </u> 5.75			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	<u> </u> : <u> </u> X 100 = <u> </u> %	<u> </u> : <u> </u> X 100 = <u> </u> %
2500 mg/L	<u> </u> : <u> </u> X 100 = <u> </u> %	<u> </u> : <u> </u> X 100 = <u> </u> %

FETAX SUMMARY SHEET

Test No. 2

Test Material	PSD	Investigator	Fort
Source	QA/QC Corrections	Lab	SBL
CAS No.		Test Start Date	4/26/93
Composition/Purity		Test End Date	4/30/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.2	7.2	7.3	7.2	
Control	7.9	7.9	7.8	7.9	
Highest Conc.	7.5	7.5	7.4	7.4	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	1 : 100 X 100 = 1 %	3 : 99 X 100 = 3 %
Solvent Control	: X 100 = %	: X 100 = %
Control Length 98.9 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG)	7.5	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.75	1.0	Williams Test
LOEL	1.0	2.5	
LC ₅₀	7.31	EC ₅₀ 2.30	
95% Confidence limits	6.39 - 8.36	95% Confidence Limits	1.96 - 2.69
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			3.18

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

Test No. 3

Test Material	P3D	Investigator	Fort
Source	QA/QC Corrections	Lab	SBL
CAS No.	Resorcinol	Test Start Date	4/26/93
Composition/Purity		Test End Date	4/30/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.5	7.4	7.5	7.5	
Control	7.9	7.9	7.9	8.0	
Highest Conc.	7.7	7.6	7.6	7.6	

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		2 : 100 X 100 = 2%	0 : 98 X 100 = 0%
Solvent Control			
Control Length 98.0 mm	Solvent Control Length mm		
Minimum Concentration to Inhibit Growth (MICIG) 2.5			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.0	0.75	Williams Test
LOEL	2.5	1.0	
LC ₅₀	5.46	EC ₅₀ 1.08	
95% Confidence limits	4.63 - 6.43	95% Confidence Limits	0.99 - 1.19
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			5.06

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

SODIUM ARSENATE

PHASE III COMPOUND E DATA (P3E) SODIUM ARSENATE

EETAX II STUDY: 7 LABS, 8 LEVELS, 3 REPLICATES

PAGE 1
FELAX LEG SIEBEN, 1 BUCH, SEITE 11

PAGE 1 DATA TAKEN WITH MATERIALS

units are in mg/ml

III/118

ORIGINAL DATA TABLE WITH MATERIALS

LOEC/MALF										SUBTOTALS:				STD:				CV:				
LAB		LC50	C1LO	C1HI	EC50	C1LO	C1HI	MCIG	G	H	I	J	LC50	EC50	MCIG	Tl	LC50	EC50	MCIG	Tl		
BANTLE/ RAYBURN	1	2.53	B	1.79	C	3.59	D	0.58	E	0.43	F	0.80	0.30	0.10	0.30	0.40	0.47	0.47	0.40	0.311	14.9	
	2	1.87		1.31		2.69		0.21		0.14		0.33		0.10		0.30	0.30	0.30	0.151	38.1		
	3	1.87		1.33		2.63		0.40		0.26		0.61		1.00		4.7	2.00	0.30	0.386	82.7		
BANTLE/ HULL	1	2	2.65	2.06	3.41	0.59	0.47	0.74	1.00	4.5	1.25	0.50	0.50	0.50	0.50	0.43	0.43	0.43	0.140	32.4		
	2	3.65		3.12		4.28		0.45		0.35		0.58		1.00		8.1	4.00	0.08	0.00	0.00	0.0	
	3	2.79		2.23		3.48		0.25		0.16		0.38		1.00		11.2	1.50	0.08	0.00	0.00	0.0	
FINCH	1	3	1.53	1.31	1.78	1.27	0.98	1.64	1.50	1.2	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.454	22.0		
	2	2.03		1.75		2.35		1.91		1.64		2.22		1.00		1.1	1.00	2.00	1.40	0.378	27.1	
	3	2.64		2.36		2.95		1.01		0.82		1.25		1.00		2.6	1.50	0.50	1.17	0.236	20.2	
LINDER/ BUCHWALTER	1	4																	LC50	#DIV/0!		
	2																		EC50	#DIV/0!		
	3																		MCIG	#DIV/0!		
BURTON/ TURLEY	1	5	1.77	1.61	1.94	1.23	1.03	1.46	1.50	1.4	1.50	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.293	14.1	
	2	2.47		2.31		2.65		1.54		1.35		1.74		1.50		1.6	1.50	1.50	1.50	0.192	14.9	
	3	1.98		1.82		2.15		1.08		0.90		1.30		1.50		1.8	1.50	1.00	0.000	0.000	0.0	
DAWSON	1	6	2.89	2.69	3.09	2.01	1.86	2.17	1.00	1.4	2.00	1.50	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.377	15.4	
	2	2.50		2.34		2.67		1.64		1.49		1.80		1.00		1.5	2.00	1.50	1.76	0.174	9.9	
	3	1.97		1.83		2.13		1.64		1.50		1.81		1.00		1.2	1.00	1.00	1.00	0.000	0.000	
STOVER/ FORT	1	7	1.89	1.62	2.20	0.39	0.31	0.49	1.00	4.8	1.00	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.138	6.6		
	2	2.16		1.83		2.56		0.56		0.49		2.50		0.64		3.9	0.75	0.50	0.52	0.097	18.6	
	3	2.20		1.83		2.65		0.62		0.54		0.70		1.00		3.5	0.75	0.25	1.50	0.707	47.1	
GRAND TOTAL																STD:	CV:				5.54E-01	
																LC50	0.97	0.5761	59.7			
																EC50	1.11	0.49	44.5			
																MCIG	2.38	2.88	120.8			
																Tl	1.51	0.73	48.5			
																LOEC (mo)	0.80	0.55560	69.8			
																LOEC (ma)						

Table 5. Results of the Interlaboratory Study for Sodium Arsenite.



Laboratory	Replicate	LC50 (mg/ml)	Mean LC50 (mg/ml)	LC50 CV1 (%)	EC50 (mg/ml)	Mean EC50 (mg/ml)	CV (%)	Mean TI ² (%)	CV (%)	MCIG ³ (mg/ml)	Mean MCIG (mg/ml)	MCIG CV (%)
1	1	2.53	2.09	14.9	0.58	0.21	0.40	38.1	5.3	39.3	0.30	82.7
	2	1.87			0.40						0.10	0.47
	3	1.87									1.00	
2	1	2.65			0.59						1.00	0.0
	2	3.65	3.03	14.6	0.45	0.43	0.25	32.4	7.0	38.7	1.00	1.00
	3	2.79									1.00	
3	1	1.53			1.27						1.50	0.0
	2	2.03	2.07	22.0	1.91	1.40	1.01	27.1	1.5	47.3	1.00	20.2
	3	2.64									1.00	
4	1											
	2											
	3											
5	1	1.77			1.23						1.50	0.0
	2	2.47	2.07	14.1	1.54	1.28	1.08	14.9	1.6	10.0	1.50	1.50
	3	1.98									1.50	
6	1	2.89			2.01						1.00	0.0
	2	2.50	2.45	15.4	1.64	1.76	1.64	9.9	1.4	9.8	1.00	1.00
	3	1.97									1.00	
7	1	1.89			0.39						1.00	0.0
	2	2.16	2.08	6.6	0.56	0.52	0.62	18.6	4.0	13.9	2.50	47.1
	3	2.20									1.00	

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

ASTM Analysis of Sodium Arsenate



FETAX Summary Sheet

Test No. DEF 1

Test Material	P3E	Investigator	JAMES RAYBURN
Source	OAK RIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	3-25-93
Composition/Purity		Test End Date	3-29-93
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.6	7.6	7.8	7.8	
	Control		7.8	7.5	7.3	7.1
	Highest Concentration		7.6	7.3	7.2	7.1

No. Dead or Malformed		X 100 = %			
Total Number		Mortality Record		Malformation Record	
FETAX Control	10 : 100	X 100 = 10%	6 : 90	X 100 = 6.7%	
Solvent Control	:	X 100 =	:	X 100 =	
Control Length	020 9.27	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)		0.3 mg/ml	d21		

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	1	0.1	T-test
LOEL	1.5	0.3	T-test
LC50	2.531	EC50 0.584	
95% CL	1.786 -- 3.586	95% Confidence limits	0.427 --- 0.798
Test Teratogenic Index (TI = LC50/EC50):			
95% Confidence limits			

POSTIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No.	2
Investigator	JAMES RAYBURN
Source	OAK RIDGE
Laboratory	BANTLE
CAS No.	Lot No.
Test Start Date:	APRIL 26 1993
Composition/Purity	Test End Date
Solvent	Test Units (i.e., mg/ml)
Conc	MG/ML

pH

	Day 0	Day 1	Day 2	Day 3	Day 4
Stock	7.8	8	7.9	7.7	
Control		7.3	7.7	7.7	7.8
Highest Concentration		7.5	7.2	7.4	6.9

No. Dead or Malformed

X 100 = %

Total Number

FETAX Control	Mortality Record			Malformation Record				
	10	: 100	X 100 =	10%	7	: 90	X 100 =	7.8%
Solvent Control	G19	: E19	X 100 =	G19	H19	: J19	X 100 =	L19
Control Length	G20	: 93	Solvent Control Length					
Minimum Concentration to Inhibit Growth (MCIG)				0.1				

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.1	0.1	T-test
LOEL	1.5	0.3	T-test
LC50	1.874	EC50	0.212
95% CL	1.307	--	95% Confidence limits
	2.687		0.135

			0.331
Test Teratogenic Index (TI = LC50/EC50):			8.86
95% Confidence limits			4.99 -- 15.74

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.6 mg/L	X 100 =	X 100 =
2600 mg/L	X 100 =	X 100 =

CL = Confidence limits

FETAX Summary Sheet

Test No. 4113

Test Material	P3E	Investigator	JAMES RAYBURN
Source	OAKRIDGE	Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	APRIL 26 1993
Composition/Purity		Test End Date:	APRIL 30 1993
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	E11	F11	G11	J11	
	Control		F12	G12	J11	K12
	Highest Concentration		F13	B13	J11	K13

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		9 : 100	X 100 =	9%	9 : 91	X 100 =	9.9%
Solvent Control		C19 : E19	X 100 =	G19	H19 : J19	X 100 =	L19
Control Length	220	8.5	Solvent Control Length		J20		
Minimum Concentration to Inhibit Growth (MCIG)		1 mg/ml	dt				

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.01	0.1	T-test
LOEL	2	0.3	T-test
LC50	1.869	EC50 0.402	
95% CL	1.330 -- 2.628	95% Confidence limits	0.263 ---- 0.613

Test Teratogenic Index (TI = LC50/EC50):	4.65
95% Confidence limits	2.70 -- 8.00

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	X 100 =	X 100 =
2500 mg/L	X 100 =	X 100 =

CL = Confidence limits

FETAX Summary Sheet

Test No. 1

Test Material	P3E	Investigator	Mendi A. Hull <i>[Signature]</i>
Source		Laboratory	Bantle / OSU
CAS No.	Lot No.	Test Start Date:	Mar 15 1993
Composition/Purity		Test End Date	Mar 19 1993
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.1	8.0	8.2	8.0	X
	Control	X	7.7	7.6	7.6	7.4
	Highest Concentration	X	8.0	7.8	7.5	7.2

No. Dead or Malformed		MALFORMATION EXCEED ASTM LIMIT			
X 100 = %		Mortality Record		Malformation Record	
<u>Total Number</u>					
<u>FETAX Control</u>		4 : 100 X 100 = 4%		10 : 96 X 100 = 10.4%	
<u>Solvent Control</u>		: X 100 =		: X 100 =	
<u>Control Length</u>		0.9 cm	Solvent Control Length		
<u>Minimum Concentration to Inhibit Growth (MCIG)</u>				1.0 mg/ml	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.25	0.25	T-test
LOEL	1.25	0.5	T-test
LC50	2.648	EC50 0.590	
95% CL	2.055 -- 3.412	95% Confidence limits	0.472 ---- 0.736
Test Teratogenic Index (TI = LC50/EC50):			4.49
95% Confidence limits			3.21 -- 6.29

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No.	2
Test Material	P3E
Investigator	Mendi A. Hull
Source	Bantle / OSU
CAS No.	Lot No.
Test Start Date:	Jun 14 1993
Composition/Purity	Test End Date
Solvent	Conc.
Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.0	8.0	8.1	8.0	X
	Control	X	7.8	7.6	7.7	7.6
	Highest Concentration	X	8.0	7.7	7.6	7.3

No. Dead or Malformed	MALFORMATION EXCEED ASTM LIMIT				
X 100 = %		Mortality Record		Malformation Record	
Total Number					
FETAX Control	7 : 100 X 100 = 7%		10 : 93 X 100 = 10.8%		
Solvent Control	: X 100 =		: X 100 =		
Control Length	1 cm	Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)		1.0 mg/ml			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	2	0.25	T-test
LOEL	4	0.5	T-test
LC50	3.654	EC50 0.448	
95% CL	3.124 -- 4.275	95% Confidence limits	0.345 ---- 0.581
Test Teratogenic Index (TI = LC50/EC50):			8.16
95% Confidence limits			6.02 -- 11.06

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. 3

Test Material	P3E	Investigator	Mendi A. Hull
Source		Laboratory	Bantle / OSU
CAS No.	Lot No.	Test Start Date:	Mar 15 1993
Composition/Purity		Test End Date	Mar 19 1993
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	8.0	8.1	8.0	8.0	X
	Control	X	7.5	7.5	7.8	7.4
	Highest Concentration	X	8.0	7.7	7.9	7.6

No. Dead or Malformed	X 100 = %	
Total Number	Mortality Record	Malformation Record
FETAX Control	0 : 100 X 100 = 0%	10 : 100 X 100 = 10.0%
Solvent Control	: X 100 =	: X 100 =
Control Length	1 cm	Solvent Control Length
Minimum Concentration to Inhibit Growth (MCIG)	1.0 mg/ml	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.1	N.A.	T-test
LOEL	1.5	0.075	T-test
LC50	2.785	EC50 0.247	
95% CL	2.230 -- 3.479	95% Confidence Limits 0.162 ---- 0.376	
Test Teratogenic Index (TI = LC50/EC50):			11.29
95% Confidence limits			7.00 -- 18.21

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

Concentration	Mortality	Malformation
5.5 mg/L	C33 : E33 X 100 = G33	H33 : J33 X 100 = L33
2500 mg/L	C34 : E34 X 100 = G34	H33 : J34 X 100 = L34

CL= Confidence limits

FETAX Summary Sheet

Test No. 1(201-01)

Test Material	P3E	Investigator	FINCH
Source	OSU	Laboratory	USABRDL
CAS No.	Lat. No.	Test Start Date:	12 JULY 93
Composition/Purity		Test End Date:	16 JULY 93
Solvent	Copic	Test Units (i.e., mg/ml)	MG/ML

pH	Stock	Day 0	Day 1	Day 2	Day 3	Day 4
	Control		7.8	7.7	7.7	7.7
	Highest Concentration		7.6	7.3	7.4	7.4

No. Dead or Malformed	X 100 = %	Mortality Record		Malformation Record	
		Total Number			X 100 =
FETAX Control	1 : 100	X 100 =	✓ 1%	2 : 99	X 100 = 2.0%
Solvent Control	--	X 100 = --	--	--	X 100 = --
Control Length	9.718	Solvent Control Length	--	--	--
Minimum Concentration to Inhibit Growth (MCIG)				1.5	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.5	0.5	T-test
LOEL	1	1	T-test
LC50	1.526 ✓	EC50 1.271 ✓	()
95% CL	1.310 --	1.778	95% Confidence limits 0.983 ---- 1.642
Test Teratogenic Index (TI = LC50/EC50):			1.20 ✓
95% Confidence limits			0.89 -- 1.62

CL = Confidence limits

FETAX Summary Sheet

Test No. 2(201-02)

Test Material	P3E	Investigator	FINCH
Source	CSU	Laboratory	USABRDL
CAS No.	Lot No.	Test Start Date:	12 JULY 93
Composition/Purity	-	Test End Date	16 JULY 93
Solvent	Conc.	Test Units (i.e., mg/ml)	MG/ML

pH		Day 1	Day 2	Day 3	Day 4
	Stock	8.9	7.6	7.6	7.6
	Control		7.8	7.7	7.7
	Highest Concentration		7.6	7.3	7.4

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		3 : 100 X 100 = ✓ 3%		7 : 97 X 100 = 7.2%			
Solvent Control		X 100 = -		X 100 = -		X 100 = -	
Control Length		9.754		Solvent Control Length			
Minimum Concentration to Inhibit Growth (MCIG)				1.0 ✓			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.5	1.5	T-test
LOEL	1	2	T-test
LC50	2.030	~1.8?	EC50 1.907 ~2.2?
95% CL	1.752	- 2.352	95% Confidence limits 1.636 --- 2.221
Test Teratogenic Index (TI = LC50/EC50):		~0.8? 1.06	
95% Confidence limits		0.86 -- 1.32	

CL = Confidence limits

FETAX Summary Sheet

Test No. 3(201-03)

Test Material	P3E	Investigator	FINCH
Source	OSU	Laboratory	USABRDL
CAS No.		Test Start Date:	19 JUL 93
Composition/Purity		Test End Date	23 JUL 93
Solvent		Test Units (i.e., mg/ml)	MG/ML

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.9	7.8	7.9	7.8	
	Control		7.7	7.6	7.6	7.7
	Highest Concentration		7.7	7.6	7.6	7.6

No. Dead or Malformed	X 100 = %	
Total Number		
FETAX Control	1 : 100 X 100 = ✓ 1%	6 : 99 X 100 = 6.1%
Solvent Control	X 100 = -	X 100 = -
Control Length	9.839	Solvent Control Length
Minimum Concentration to Inhibit Growth (MCIG)		1.0

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	1	N.A.	T-test
LOEL	1.5	0.5	T-test
LC50	2.640	2.5	EC50 1.014 ~ 1.3
95% CL	2.361	- 2.952	95% Confidence limits 0.824 ---- 1.250
Test Teratogenic Index (TI = LC50/EC50): ~ 1.4 ? 2.60 - ?			
95% Confidence limits 2.05 -- 3.30			

CL = Confidence limits

FETAC SUMMARY SHEET

Test No. 1

Test Material	UNKNOWN P3E	Investigator	TURLEY
Source	OSU	Lab	UMD/WREC
CAS No.	Lot No.	Test Start Date	5/20/93
Composition/Purity		Test End Date	5/24/93
Solvent	Conc.	Test Units (1.0, mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	9.16	9.10	9.07	9.13	9.15
Control	7.61	7.57	7.60	7.55	7.60
Highest Conc.	8.48	8.40	8.38	8.44	8.46

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		2 : 100 X 100 = 2 %	4 : 98 X 100 = 4 %
Solvent Control		_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
Control Length 10.16 mm	Solvent Control Length mm		
Minimum Concentration to Inhibit Growth (MCIG) 1.5 mg/ml			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.0 mg/ml	0.5 mg/ml	Bonferroni T-test
LOEL	1.5 mg/ml	1.0 mg/ml	Bonferroni T-test
LC ₅₀ 1.77 mg/ml (Trimmed Spearman-Karber)	EC ₅₀ 1.23 mg/ml (Trimmed Spearman-Karber)		
95% Confidence limits (1.61 - 1.94)	95% Confidence Limits (1.03 - 1.46)		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.44

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETA SUMMARY SHEET

Test No. 2

Test Material	UNKNOWN P3E	Investigator	TURLEY
Source	O.S.U	Lab	JMD/WREC
CAS No.	Lot No.	Test Start Date	June 15, 1993
Composition/Purity		Test End Date	June 19, 1993
Solvent	Conc.	Test Units (i.e., mg/ml)	

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH Stock	9.19	9.14	9.15	9.09	9.06
Control	7.59	7.55	7.61	7.55	7.49
Highest Conc. (5.0)	8.31	8.44	8.39	8.33	8.27

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number		
Solvent Control		
Control Length 9.89 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG) 1.5 mg/ml		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	Fail Normality/Homogeneity	1.0 mg/ml	BonFerroni T-test
LOEL	Fail Normality/Homogeneity	1.5 mg/ml	BonFerroni T-test
LC ₅₀	2.47 mg/ml (Truncated Spearman-Katner)	EC ₅₀ 1.54 mg/ml (Truncated Spearman-Katner)	
95% Confidence limits	2.31 - 2.65	95% Confidence Limits 1.35 - 1.74	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		1.60	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

Test No. 3

Test Material UNKNOWN P3E	Investigator TURLEY
Source O.S.U.	Lab UMD/WREC
CAS No.	Lot No.
Composition/Purity	Test Start Date: June 24, 1993
Solvent	Test End Date: June 28, 1993
	Test Units (i.e., mg/ml) mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	9.11	9.19	9.09	9.13	9.14
Control	7.55	7.59	7.61	7.57	7.53
Highest Conc. S.D./mg/l	8.40	8.43	8.40	8.42	8.38
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %	2 : 100 X 100 = 2 %	8 : 98 X 100 = 8 %		
Total Number					
Solvent Control					
Control Length 10.02 mm	Solvent Control Length mm				
Minimum Concentration to Inhibit Growth (MCIG) 1.5 mg/ml					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.0 mg/ml	0.5 mg/ml	BonFerroni T-test
LOEL	1.5 mg/ml	1.0 mg/ml	Bonferroni - T-test
LC ₅₀	1.98 (Trimmed Spearman-Kaber)	EC ₅₀ 1.08 (Trimmed Spearman-Kaber)	
95% Confidence limits 1.82 - 2.15		95% Confidence Limits (0.90 - 1.30)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.83

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

		Test No. <u>1</u>
Test Material	P3E	Investigator <u>D.A. DAWSON</u>
Source		Lab UT /CVM
CAS No.	Lot No.	Test Start Date <u>5/3/93</u>
Composition/Purity		Test End Date <u>5/7/93</u>
Solvent	Conc.	Test Units (1.0., mg/ml) <u>mg/ml</u>

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.39 adj	7.40 adj	7.41 adj	7.41 adj	—
Control	7.74	7.63	7.52	7.57	7.61
Highest Conc.	—	7.59	7.61	7.67	7.71

FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed		$\times 100 = \%$			
Total Number		<u>0</u> : <u>100</u> $\times 100 =$ <u>0</u> %		<u>2</u> : <u>100</u> $\times 100 =$ <u>2</u> %	
Solvent Control		<u> </u> : <u> </u> $\times 100 =$ <u> </u> %		<u> </u> : <u> </u> $\times 100 =$ <u> </u> %	
Control Length mm		Solvent Control Length mm			
Minimum Concentration to Inhibit Growth (MCIG)		1.0			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.5 both	1.0 both	
LOEL	2.0 both	1.5 both	
LC ₅₀	2.81 L&W	EC ₅₀ 2.01 L&W	
95% Confidence Limits	2.09 - 3.09	95% Confidence Limits (1.86 - 2.17)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		1.44	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L not	<u> </u> : <u> </u> $\times 100 =$ <u> </u> %	<u> </u> : <u> </u> $\times 100 =$ <u> </u> %
2500 mg/L d/e	<u> </u> : <u> </u> $\times 100 =$ <u> </u> %	<u> </u> : <u> </u> $\times 100 =$ <u> </u> %

D.A.D.

FETAX SUMMARY SHEET

Test No. 2

Test Material	P3E	Investigator	J.A. Dawson
Source		Lab	AT/CVM
CAS No.	Lot No.	Test Start Date	5/10/83
Composition/Purity		Test End Date	5/14/83
Solvent	Cone.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.37 adj	7.38 adj	7.39 adj	7.40 adj	—
Control	7.72	7.61	7.66	7.46	7.45
Highest Conc.	—	7.60	7.56	7.56	7.52

FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %				
Total Number		0 : 100 X 100 = 0 %		3 : 100 X 100 = 3 %	
Solvent Control		_____ : _____ X 100 = _____ %		_____ : _____ X 100 = _____ %	
Control Length mm		Solvent Control Length mm			
Minimum Concentration to Inhibit Growth (MCIG)		1.0			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.5 both	1.0 both	
LOEL	2.0 both	1.5 both	
LC ₅₀	2.50	L & W	EC ₅₀ 1.64 L & W
95% Confidence Limits (2.34 - 2.67)		95% Confidence Limits (1.49 - 1.80)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		1.52	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L lot	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L dme	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX SUMMARY SHEET

Test No. 3

Test Material	P3E	Investigator	D.A. DAWSON
Source		Lab	WT/CVM
CAS No.	Lot No.	Test Start Date	5/11/93
Composition/Purity		Test End Date	5/15/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.38 adj.	7.39 adj.	7.41 adj.	7.39 adj.	--
Control	7.70	7.60	7.52	7.43	7.39
Highest Conc.	—	7.61	7.55	7.59	7.54

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		0 : 100 X 100 = 0 %	3 : 100 X 100 = 3 %
Solvent Control		0 : 100 X 100 = 0 %	0 : 100 X 100 = 0 %
Control Length mm	Solvent Control Length mm		
Minimum Concentration to Inhibit Growth (MCIG) 1.0			

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	< 1.0 M/L	< 1.0 BOTH	
LOEL	1.0 M/L	1.0 BOTH	
LC ₅₀	1.97	EC ₅₀ 1.64	
95% Confidence Limits (1.83 - 2.13)		95% Confidence Limits (1.58 - 1.81)	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀) 1.20			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L Not	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2500 mg/L 0/48	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

		Test No.																										
Test Material	P3E	Investigator Farr																										
Source	QA/QC Corrections	Lab SBL																										
CAS No.	Lot No.	Test Start Date 3/29/93																										
Composition/Purity		Test End Date 4/2/93																										
Solvent	Conc.	Test Units (i.e., mg/ml) mg/ml																										
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%;">pH</th> <th style="width: 20%;">DAY 1</th> <th style="width: 20%;">DAY 2</th> <th style="width: 20%;">DAY 3</th> <th style="width: 20%;">DAY 4</th> <th style="width: 20%;">DAY 5</th> </tr> </thead> <tbody> <tr> <td>Stock</td> <td>7.5</td> <td>7.6</td> <td>7.4</td> <td>7.3</td> <td></td> </tr> <tr> <td>Control</td> <td>8.0</td> <td>8.1</td> <td>8.0</td> <td>8.0</td> <td></td> </tr> <tr> <td>Highest Conc.</td> <td>7.5</td> <td>7.7</td> <td>7.7</td> <td>7.5</td> <td></td> </tr> </tbody> </table>					pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	Stock	7.5	7.6	7.4	7.3		Control	8.0	8.1	8.0	8.0		Highest Conc.	7.5	7.7	7.7	7.5	
pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5																							
Stock	7.5	7.6	7.4	7.3																								
Control	8.0	8.1	8.0	8.0																								
Highest Conc.	7.5	7.7	7.7	7.5																								
FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD																									
No. Dead or Malformed Total Number		$\times 100 = \%$ 0 : 0 $\times 100 = 0\%$	$\times 100 = \%$ 0 : 0 $\times 100 = 0\%$																									
Solvent Control		$\times 100 = \%$: : $\times 100 = \%$	$\times 100 = \%$: : $\times 100 = \%$																									
Control Length 96.3 mm		Solvent Control Length mm																										
Minimum Concentration to Inhibit Growth (MCIG) 1.0																												

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.5	0.05	Williams Test
LOEL	1.0	0.1	
LC ₅₀	(PF) 0.76	1.89	EC ₅₀ 0.39
95% Confidence limits	0.72 - 0.81		95% Confidence Limits 0.31 - 0.49
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			4.85
(DF) 1.62 - 2.20			

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

Test No.

2

Test Material	PBE	Investigator	Fort
Source	QA/QC Corrections	Lab	SBL
CAS No.	Resorcinol	Test Start Date	4/26/93
Composition/Purity		Test End Date	4/30/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.4	7.3	7.2	7.2	
Control	7.9	7.9	7.8	7.9	
Highest Conc.	7.5	7.4	7.5	7.5	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number		
Solvent Control	: X 100 = %	: X 100 = %
Control Length 97.8 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG)	2.5	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.5	0.25	Williams test
LOEL	0.75	0.5	
LC ₅₀	2.1b	EC ₅₀	0.56
95% Confidence limits	1.83 - 2.56	95% Confidence Limits	0.49 - 0.64
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			3.86

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

FETAX SUMMARY SHEET

Test No. 3

Test Material	P3E	Investigator	Fort
Source	QA/QC Correlation	Lab	SBL
CAS No.	Resources	Lot No.	
Composition/Purity		Test Start Date	4/26/93
Solvent	Cone.	Test End Date	4/30/93
		Test Units (i.e., mg/ml)	mg/ml

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.4	7.3	7.2	7.2	
Control	7.8	7.9	7.8	7.9	
Highest Conc.	7.4	7.5	7.5	7.5	

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		0 : 100 X 100 = 0 %	0 : 100 X 100 = 0 %
Solvent Control		0 : _____ X 100 = _____ %	0 : _____ X 100 = _____ %
Control Length 97.1 mm	Solvent Control Length mm		
Minimum Concentration to Inhibit Growth (MCIG)	1.0		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.5	0.1	Williams Test
LOEL	0.75	0.25	
LC ₅₀	2.20	EC ₅₀	0.62
95% Confidence limits	1.83 - 2.65	95% Confidence Limits	0.54 - 0.70
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		3.55	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

COPPER SULFATE

PHASE III COMPOUND F DATA (P3F) CuSO(4)
FETAX ILS STUDY; 7 LABS, 8 LEVELS, 3 REPLICATES
PAGE 1
ORIGINAL DATA TABLE WITH MATERIALS

FETAX ILS STUDY; 7 LABS, 8 LEVELS, 3 REPLICATES Units are in mg/l PAGE 1

Table 6. Results of the Interlaboratory Study for Copper Sulfate.


Laboratory	Replicate (n)	Mean LC50 (mg/L)	LC50 CV ¹ (%)	Mean EC50 (mg/L)	EC50 CV (%)	Mean TI ² (%)	TI CV (%)	Mean MCIG ³ (mg/L)	MCIG CV (%)
1	1	1.00		0.32				0.70	
	2	1.45	1.56	32.8	0.21	0.28	17.7	5.6	33.4
	3	2.24		0.31				0.40	0.57
2	1	0.63		0.15				0.20	
	2	0.79	0.77	13.9	0.24	0.20	19.0	3.8	10.5
	3	0.89		0.22				0.40	0.27
3	1	0.93		—				0.05	
	2	0.66	0.85	15.6	0.33	0.45	25.8	1.9	8.0
	3	0.95		0.56				0.05	0.04
4	1	0.29		0.08				0.01	
	2	0.50	0.46	26.8	0.11	0.20	74.5	2.3	57.5
	3	0.58		0.41				0.03	0.04
5	1	1.70		2.47				0.75	
	2	2.07	2.17	19.8	1.61	2.68	36.1	0.8	34.6
	3	2.74		3.95				1.00	0.92
6	1	0.88		0.81				0.25	
	2	0.82	0.88	5.1	0.76	0.79	2.7	1.1	3.4
	3	0.93		0.80				0.10	0.20
7	1	1.49		1.48				0.50	
	2	2.09	1.93	16.2	0.89	1.10	24.7	1.8	36.6
	3	2.20		0.92				0.30	0.43
								0.50	21.8

¹CV=Coefficient of Variation.

²TI=Mean Teratogenic Index calculated by dividing the Mean LC50 by the Mean EC50 for each laboratory.

³MCIG=Minimum Concentration to Inhibit Growth.

ASTM Analysis of Copper Sulfate



FETAX Summary Sheet

Test No. DEF. 1

Test Material	P3F	Investigator	JAMES RAYBURN <i>[Signature]</i>
Source		Laboratory	BANTLE
CAS No.	Lot No.	Test Start Date:	APRIL 25 1994
Composition/Purity		Test End Date	APRIL 29 1994
Solvent	NONE	Conc.	Test Units (i.e., mg/ml) MG/L

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	6.8	6.7	6.8	6.8	
	Control		7.7	7.7	7.7	7.3
	Highest Concentration		6.8	7	7	7.1

No. Dead or Malformed	X 100 = %	Mortality Record		Malformation Record	
		Total Number			X 100 =
FETAX Control	3 : 100	X 100 =	3%	8 : 97	X 100 = 8.2%
Solvent Control	:	X 100 =		:	X 100 =
Control Length (mm)	8.7	Solvent Control Length (mm)			
Minimum Concentration to Inhibit Growth (MCIG)		0.7		MG/L	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.3	0.1	T-test
LOEL	16 0.9	0.2	T-test
LC50	0.995	EC50 0.317	
95% CL	0.894 -- 1.107	95% Confidence limits	0.275 ---- 0.365
Test Teratogenic Index (TI = LC50/EC50):			3.14
95% Confidence limits			2.63 -- 3.75

FETAX Summary Sheet

Test No.	2
Investigator	James Rayburn <i>JRB</i>
Source	Laboratory Bantle
CAS No.	Lot No. 09-JUN-94
Composition/Purity	Test End Date 12-JUN-94
Solvent	Conc. Test Units (I.e., mg/ml) mg/l

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7	7	7	7.1	
	Control		7.5	7.6	7.1	7.4
	Highest Concentration		7.5	7.3	7.2	7.3

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control	4 : 100	X 100 =	4%	10 : 96	X 100 =	10.4%	
Solvent Control	:	X 100 =		:	X 100 =		
Control Length (mm)		9.7	Solvent Control Length (mm)				
Minimum Concentration to Inhibit Growth (MCIG)				0.4	mg/l		

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	1	0.1	T-test
LOEL	2	0.2	T-test
LC50	1.448	EC50 0.206	
95% CL	1.160 -- 1.808	95% Confidence Limits	0.151 ---- 0.282
Test Teratogenic Index (TI = LC50/EC50):			7.03
95% Confidence Limits			4.79 -- 10.32

FETAX Summary Sheet

Test No. 3

Test Material	P3F	Investigator	James Rayburn
Source		Laboratory	Bantle
CAS No.	Lot No.	Test Start Date:	July 7, 1994
Composition/Purity		Test End Date	July 11, 1994
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/l

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.2	7	7.1	7	
	Control		7.6	7.7	7.2	7.1
	Highest Concentration		7	7.3	7.2	7.1

No. Dead or Malformed	X 100 = %	Mortality Record		Malformation Record	
Total Number					
FETAX Control	3 : 100	X 100 =	3%	9 : 97	X 100 = 9.3%
Solvent Control	:	X 100 =		:	X 100 =
Control Length (mm)	9.49	Solvent Control Length (mm)			
Minimum Concentration to Inhibit Growth (MCIG)		0.6 mg/l			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.05	0.1	T-test
LOEL	1	0.2	T-test
LC50	2.239	EC50 0.312	
95% CL	1.723 -- 2.911	95% Confidence limits	0.276 ---- 0.353
Test Teratogenic Index (TI = LC50/EC50):			7.18
95% Confidence limits			5.37 -- 9.59

Percent effect	LC	EC
5	0.8367732	0.162
16	1.234964	0.21
50	2.2394243	0.312
84	4.0608643	0.464
95	5.9932858	0.602

FETAX Summary Sheet

Test No.	1
Investigator	Mendi A. Hull <i>MB</i>
Source	B5
CAS No.	B4
Composition/Purity	Lot No. E5 C6
Solvent	B7 Conc. E7
Test Units (i.e., mg/ml)	mg/L

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.5	7.5	7.5	7.5	
	Control		7.6	7.5	7.4	7.6
	Highest Concentration		7.7	8	8	7.9

No. Dead or Malformed		X 100 = %			
Total Number		Mortality Record		Malformation Record	
<u>FETAX Control</u>	<u>10 : 100</u>	X 100 =	<u>10%</u>	<u>6 : 90</u>	X 100 = <u>6.7%</u>
<u>Solvent Control</u>	:	X 100 =		:	X 100 =
<u>Control Length (mm)</u>	0.9	Solvent Control Length (mm)	J20		
<u>Minimum Concentration to Inhibit Growth (MCIG</u>		0.2		mg/L	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.4	0.1	T-test
LOEL	0.6	0.2	T-test
LC50	0.634	EC50 0.147	
95% CL	0.479 -- 0.838	95% Confidence limits	0.115 ---- 0.187
Test Teratogenic Index (TI = LC50/EC50):			4.32
95% Confidence limits			2.98 -- 6.26

FETAX Summary Sheet

Test No.	2
Investigator	Mendi A. Hull <i>JAHB</i>
Laboratory	OSU/Bantle
Test Start Date:	11/19/93
Test End Date	11/23/93
Test Units (i.e., mg/ml)	mg/L

Test Material	P3F	Investigator	Mendi A. Hull <i>JAHB</i>
Source	B5	Laboratory	OSU/Bantle
CAS No.	B4	Lot No.	E5
Composition/Purity	C6	Test Start Date:	11/19/93
Solvent	B7	Conc.	E7
		Test End Date	11/23/93
		Test Units (i.e., mg/ml)	mg/L

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.5	7.5	7.5	7.5	
	Control		7.5	7.4	7.4	7.8
	Highest Concentration		7.8	7.6	7.8	7.7

No. Dead or Malformed	X 100 = %	Mortality Record		Malformation Record	
Total Number					
FETAX Control	9 : 100	X 100 =	9%	5 : 91	X 100 = 5.5%
Solvent Control	:	X 100 =		:	X 100 =
Control Length (mm)	0.9	Solvent Control Length (mm)		J20	
Minimum Concentration to Inhibit Growth (MCIG)		0.4			mg/L

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.4	0.1	T-test
LOEL	0.6	0.2	T-test
LC50	0.788	EC50 0.243	
95% CL	0.605 -- 1.027	95% Confidence limits	0.180 ---- 0.326
Test Teratogenic Index (TI = LC50/EC50):			3.25
95% Confidence limits			2.18 -- 4.83

FETAX Summary Sheet

Test No.	3
Test Material	P3F
Source	B5
CAS No.	B4
Composition/Purity	C6
Solvent	B7
Conc.	E7
Investigator	Mendi A. Hull <i>JMHS</i>
Laboratory	OSU/Bantle
Lot No.	E5
Test Start Date:	12/2/93
Test End Date	12/6/93
Test Units (i.e., mg/ml)	mg/L

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.5	7.5	7.5	7.5	
	Control		7.6	7.5	7.5	7.6
	Highest Concentration		7.5	7.4	7.4	7.5

No. Dead or Malformed		X 100 = %			
Total Number		Mortality Record		Malformation Record	
FETAX Control	1 : 100	X 100 =	1%	7 : 99	X 100 = 7.1%
Solvent Control	:	X 100 =		:	X 100 =
Control Length (mm)	0.9	Solvent Control Length (mm)		J20	
Minimum Concentration to Inhibit Growth (MCIG)		0.2		mg/L	

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.6	0.01	T-test
LOEL	4	0.1	T-test
LC50	0.887	EC50 0.218	
95% CL	0.681	-- 1.154	95% Confidence Limits 0.156 ---- 0.303

Test Teratogenic Index (TI = LC50/EC50): 4.08

95% Confidence Limits 2.67 -- 6.22

FETAX SUMMARY SHEET

		Test No. Definitive #
Test Material: P3F		Investigator: Gillett / Wilborn
Source: Bantle		Laboratory: CERL
CAS No. — Lot No. —		Test Start Date: 01/06/94
Composition / Purity: —		Test End Date: 01/10/94
Solvent: — Conc. —		Test Units:(i.e., mg/ml) mg/l

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.4	7.0	7.6	7.5	7.5 X
Control	X	7.5	6.9 ^{co} 7.1	7.4	7.4
Highest Conc.	X	7.6	6.9	7.4	7.2

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed x 100 = % Total Number	12 : 100 X 100 = 12%	6 : 48 X 100 = 6.8 %
Solvent Control: N.A.	— : — X 100 = — %	— : — X 100 = — %
Control Length: 9.325 mm	Solvent Control Length: N.A. mm	
Minimum Concentration to Inhibit Growth (MCIG): .005		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.1 mg/L	0.01 mg/L	Williams
LOEL	0.5 mg/L	0.025 mg/L	Williams
LC ₅₀ 0.29 mg/L (Spearman-Karber)		EC ₅₀ 0.08 mg/L (Spearman-Karber)	
95% Confidence Limits 0.22 - 0.38 mg/L		95% Confidence Limits 0.05 - 0.12 mg/L	
TEST TERATOGENIC INDEX (TI = LC ₅₀ /EC ₅₀) - 3.625			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	— : — X 100 = — %	— : — X 100 = — %
2.5 mg/ml	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

Test Material: P3F	Test No. Definitive #2
Source: Battelle	Investigator: G. Hall
CAS No. — Lot No. —	Laboratory: CERL
Composition / Purity: —	Test Start Date: 01/14/94
Solvent: — Conc. —	Test End Date: 01/18/94 Test Units:(i.e., mg/ml) mg/l

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.0	7.2	7.5	7.1	X
Control	X	7.5	7.3	7.4	7.5
Highest Conc.	X	7.6	7.2	6.6	7.0

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed X 100 = % Total Number	13 : 100 X 100 = 13%	16 : 87 X 100 = 18%
Solvent Control: N.A.	— : — X 100 = — %	— : — X 100 = — %
Control Length: 8.828 mm	Solvent Control Length: N.A. mm	
Minimum Concentration to Inhibit Growth (MCIG): 0.025		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.1 mg/L	0.05 mg/L	Williams
LOEL	0.5 mg/L	0.075 mg/L	Williams
LC_{50} 0.50 mg/L (Spearman-Karber)		EC_{50} 0.11 mg/L (Spearman-Karber)	
95% Confidence Limits 0.38-0.65 mg/L		95% Confidence Limits 0.06-0.19 mg/L	
TEST TERATOGENIC INDEX (TI = LC_{50}/EC_{50}) - 4.545			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	— : — X 100 = — %	— : — X 100 = — %
2.5 mg/ml	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

		Test No. Definitive #3
Test Material: P3F		Investigator: Gillett
Source: Bantle		Laboratory: CERL
CAS No. — Lot No. —		Test Start Date: 01/28/94
Composition / Purity: —		Test End Date: 02/01/94
Solvent: — Conc. —		Test Units:(i.e., mg/ml) mg/L

--pH--	Day 1	Day 2	Day 3	Day 4	Day 5
Stock	7.2	7.3	7.2	7.4	X
Control	X	7.4	7.3	7.3	7.4
Highest Conc.	X	6.8	6.7	6.9	7.3

FETAX CONTROL	MORTALITY	MALFORMATION
No. Dead or Malformed x 100 = % Total Number	2 : 100 X 100 = 2 %	1 : 98 X 100 = 1 %
Solvent Control: N.A.	— : — X 100 = — %	— : — X 100 = — %
Control Length: 9.475 mm	Solvent Control Length: N.A.	mm
Minimum Concentration to Inhibit Growth (MCIG): 0.075 mg/L		

STATISTICAL RESULTS:

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.1 mg/L	0.075 mg/L	Williams
LOEL	0.5 mg/L	0.1 mg/L	Williams
LC_{50} 0.58 mg/L (Spearman-Karber)		EC_{50} 0.41 mg/L (Spearman-Karber)	
95% Confidence Limits 0.51-0.66 mg/L		95% Confidence Limits 0.19-0.86 mg/L	
TEST TERATOGENIC INDEX (TI = LC_{50}/EC_{50}) - 1.415			

POSITIVE CONTROL: 6-AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
0.0055 mg/ml	— : — X 100 = — %	— : — X 100 = — %
2.5 mg/ml	— : — X 100 = — %	— : — X 100 = — %

FETAX SUMMARY SHEET

178

Test No. 1

Test Material	UNKNOWN P3F	Investigator	TURLEY
Source	OSU	Lab	UMD - WREC
CAS No.	Lot No.	Test Start Date	11/24/93
Composition/Purity		Test End Date	11/30/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/L

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.13	7.11	7.15	7.13	
Control	7.65	7.60	7.67	7.70	
Highest Conc.	7.29	7.33	7.30	7.31	
FETAX CONTROL		MORTALITY RECORD		MALFORMATION RECORD	
No. Dead or Malformed	X 100 = %				
Total Number		2 : 100 X 100 = 2 %		6 : 98 X 100 = 6.1 %	
Solvent Control					
Control Length 1002 mm		Solvent Control Length mm			
Minimum Concentration to Inhibit Growth (MCIG) 0.75 mg/L					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	0.50	0.75	Bonferroni T-test
LOEL	0.75	1.00	Bonferroni T-test
LC ₅₀ 1.70 Trimmed Spearman-Karber	EC ₅₀ 2.47 Trimmed Spearman-Karber		
95% Confidence Limits 1.34 - 2.15	95% Confidence Limits 0.32 - 19.30		
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		0.69	

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2500 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

47B

Test No. 2

Test Material	UNKNOWN P3F	Investigator	TURLEY
Source	1511	Lab	UMD-WREC
CAS No.	Lot No.	Test Start Date	12/20/93
Composition/Purity		Test End Date	12/24/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/L

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH					
Stock	7.16	7.12	7.14	7.20	
Control	7.71	7.69	7.69	7.66	
Highest Conc.	7.20	7.23	7.26	7.23	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed		
Total Number	X 100 = %	X 100 = %
Solvent Control	1 : 80 X 100 = 1.25 %	4 : 79 X 100 = 5.1 %
Control Length 9.29 mm	Solvent Control Length mm	
Minimum Concentration to Inhibit Growth (MCIG)	1.0	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	FAIL HOMOGENEITY	1.0	Bonferroni T-test
LOEL	FAIL HOMOGENEITY	3.0	Bonferroni T-test
LC ₅₀ 2.07	Trimmed - Spearman-Karber	EC ₅₀ 1.61	Trimmed Spearman-Karber
95% Confidence Limits 1.77 - 2.41		95% Confidence Limits 1.21 - 2.14	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)		1.33	1.29

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %
2500 mg/L	____ : ____ X 100 = ____ %	____ : ____ X 100 = ____ %

FETAX SUMMARY SHEET

413

Test No. 3

Test Material	UNANONN P3F	Investigator	TURLEY
Source	OSU.	Lab	UMD-WREC
CAS No.	Lot No.	Test Start Date	12/21/93
Composition/Purity		Test End Date	12/27/93
Solvent	Conc.	Test Units (i.e., mg/ml)	ml/L

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
pH					
Stock	7.09	7.11	7.13	7.16	
Control	7.77	7.69	7.69	7.73	
Highest Conc.	7.30	7.28	7.30	7.31	
FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD			
No. Dead or Malformed	X 100 = %				
Total Number					
Solvent Control					
Control Length 9.69 mm	Solvent Control Length	mm			
Minimum Concentration to Inhibit Growth (MCIG) 1.0 mg/L					

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	1.0	2.0	Bonferroni T-test
LOEL	2.0	3.0	Bonferroni T-test
LC ₅₀ 2.74 Trimmed Spiegelman-Kabba		EC ₅₀ 3.95	
95% Confidence limits 1.97-3.65		95% Confidence Limits 3.13-4.99	
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			0.69 -

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %
2500 mg/L	_____ : _____ X 100 = _____ %	_____ : _____ X 100 = _____ %

FETAX Summary Sheet

Test No. 1

Test Material	P3F	Investigator	Doug Dawson
Source		Laboratory	Ashland Univ
CAS No.	Lot No.	Test Start Date:	7/12/94
Composition/Purity		Test End Date	7/16/94
Solvent	Stock Co 2.5 mg/	Test Units (i.e., mg/ml)	mg/L

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	6.94	9.63	6.93	6.9	
	Control		7.33	7.24	7.4	7.27
	Highest Concentration		7.48	7.42	7.4	--

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		0 : 100 X 100 = 0%		2 : 100 X 100 = 2.0%			
Solvent Control		: X 100 =		:		X 100 =	
Control Length (mm) C20		Solvent Control Length (mm) J20					
Minimum Concentration to Inhibit Growth (MCIG) G21		0.25		mg/L			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.25	0.625	T-test
LOEL	0.375	0.75	T-test
LC50	0.883	EC50 0.808	
95% CL	0.791 --	0.986	95% Confidence limits 0.722 ---- 0.905
Test Teratogenic Index (TI = LC50/EC50):			1.09
95% Confidence limits			0.93 -- 1.28

Percent effect	LC	EC
5	0.4607641	0.448
16	0.5958838	0.565
50	0.8829404	0.808
84	1.3082815	1.156
95	1.6919368	1.46

FETAX Summary Sheet

Test No. 2

Test Material	P3F	Investigator	Doug Dawson
Source		Laboratory	Ashland Univ
CAS No.	Lot No.	Test Start Date:	7/18/94
Composition/Purity		Test End Date	7/22/94
Solvent	Stock Co 2.5 mg/	Test Units (i.e., mg/ml)	mg/L

pH		Day 0	Day 1	Day 2	Day 3	Day 4
	Stock	7.07	7.07	7.07	7.1	
	Control		7.44	7.41	7.3	7.19
	Highest Concentration		7.5	7.43	7.4	7.27

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		0 : 100 X 100 = 0%		3 : 100 X 100 = 3.0%			
Solvent Control		: X 100 =		: X 100 =			
Control Length (mm)		Solvent Control Length (mm)					
Minimum Concentration to Inhibit Growth (MCIG)		0.25 mg/L					

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used				
NOEL	0.5	0.375	T-test				
LOEL	0.625	0.75	T-test				
LC50	0.815		EC50	0.759			
95% CL	0.736	-- 0.902	95% Confidence Limits	0.675	----	0.855	
			Test Teratogenic Index (TI = LC50/EC50):	1.07			
			95% Confidence Limits	0.92	--	1.25	

Percent effect	LC	EC
5	0.4879727	0.466
16	0.5976449	0.565
50	0.8148417	0.759
84	1.1109725	1.02
95	1.3606643	1.238

FETAX Summary Sheet

Test No. 3

Test Material	P3F	Investigator	Doug Dawson
Source		Laboratory	Ashland Univ
CAS No.	Lot No.	Test Start Date:	7/21/94
Composition/Purity		Test End Date	7/25/94
Solvent	Stock Co 2.5 mg/	Test Units (i.e., mg/ml)	mg/L

pH	Stock	Day 0	Day 1	Day 2	Day 3	Day 4
		7.09	7.07	7.11	7.1	
	Control		7.51	7.42	7.3	7.16
	Highest Concentration		7.42	7.39	7.3	7.29

No. Dead or Malformed		X 100 = %		Mortality Record		Malformation Record	
Total Number							
FETAX Control		0	: 100	X 100 =	0%	4	: 100 X 100 = 4.0%
Solvent Control		:		X 100 =	:	X 100 =	
Control Length (mm)				Solvent Control Length (mm)			
Minimum Concentration to Inhibit Growth (MCIG)				0.1 mg/L			

Test Material/Compound Results

Test	Mortality	Malformation	Statistical Test Used
NOEL	0.5	0.375	T-test
LOEL	0.625	0.5	T-test
LC50	0.933	EC50 0.802	
95% CL	0.831	-- 1.049	95% Confidence limits 0.709 ---- 0.908
	Test Teratogenic Index (TI = LC50/EC50):		
	95% Confidence limits 0.98 -- 1.38		

Percent effect	LC	EC
5	0.5032428	0.418
16	0.6425083	0.541
50	0.9334985	0.802
84	1.3562773	1.189
95	1.7316082	1.538

FETAX SUMMARY SHEET

Test Material	CuSO ₄ P3F	Investigator	M. JACKSON
Source		Lab	STOEVER BIO LAB
CAS No.	Lot No.	Test Start Date	8/24/93
Composition/Purity		Test End Date	8/28/93
Solvent	Conc.	Test Units (i.e., mg/ml)	mg/L

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.8	7.8	7.8	7.8	
Control	7.8	7.8	7.8	7.8	
Highest Conc.	7.8	7.8	7.8	7.8	

FETAX CONTROL		MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %		
Total Number		7 : 100 X 100 = 7 %	4 : 93 X 100 = 4 %
Solvent Control	NA	: X 100 = %	: X 100 = %
Control Length 90.9 mm		Solvent Control Length NA mm	
Minimum Concentration to Inhibit Growth (MCIG)		0.5	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL	NA	NA	Failed variance assumption
LOEL	NA	NA	"
LC ₅₀	1.49	EC ₅₀	1.48
95% Confidence limits	1.31 - 1.70	95% Confidence Limits	1.18 - 1.85
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			1.00

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	: X 100 = %	: X 100 = %
2500 mg/L	: X 100 = %	: X 100 = %

LOC
DR

FETAX SUMMARY SHEET

Test No. 4 K

Test Material	P3F	Investigator
Source	OSU /BANTLE	Lab
CAS No.	Lot No.	Test Start Date 12/10/93
Composition/Purity		Test End Date 12/14/93
Solvent	Cone.	Test Units (i.e., mg/mL)

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.4	7.4	7.4	7.4	
Control	8.0	8.0	7.9	8.1	
Highest Conc.	7.6	7.4	7.5	7.5	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number	0 : 100 X 100 = 0 %	4 : 100 X 100 = 4 %
Solvent Control	0 : X 100 = %	0 : X 100 = %
Control Length 93.4 mm	Solvent Control Length 93.4 mm	
Minimum Concentration to Inhibit Growth (MCIG) 0.3		

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL			
LOEL			
LC ₅₀	2.09	EC ₅₀	0.89
95% Confidence limits	2.00 - 2.18	95% Confidence Limits	0.85 - 0.94
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			2.35

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	0 : X 100 = %	0 : X 100 = %
2500 mg/L	0 : X 100 = %	0 : X 100 = %

* Ca measured = 3.9 mg/L Corrected for weight
 Gravimetric = 10.0 mg/L error

FETAX SUMMARY SHEET

Test No. VI 15

Test Material	P3F	Investigator
Source	OSU / BANTLE	Lab
CAS No.	Lot No.	Test Start Date
Composition/Purity		12/10/93
Solvent	Conc.	Test End Date
		12/14/93
		Test Units (i.e., mg/ml)

pH	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
Stock	7.4	7.4	7.4	7.4	
Control	8.0	8.0	8.0	8.0	
Highest Conc.	7.6	7.6	7.5	7.5	

FETAX CONTROL	MORTALITY RECORD	MALFORMATION RECORD
No. Dead or Malformed	X 100 = %	
Total Number		
Solvent Control	0 : 100 X 100 = 0 %	0 : 100 X 100 = 0 %
Control Length 94.4 mm	: X 100 = %	: X 100 = %
Minimum Concentration to Inhibit Growth (MCIG) 0.5	Solvent Control Length mm	

TEST MATERIAL/COMPOUND : RESULTS

TEST	MORTALITY	MALFORMATION	STATISTICAL TEST USED
NOEL			
LOEL			
LC ₅₀	2.20	EC ₅₀	0.92
95% Confidence Limits	2.10 - 2.51	95% Confidence Limits	0.84 - 1.03
TEST TERATOGENIC INDEX (TI = LC ₅₀ / EC ₅₀)			2.39

POSITIVE CONTROL: 6 AMINONICOTINAMIDE (6-AN) RESULTS

CONCENTRATION	MORTALITY	MALFORMATION
5.5 mg/L	0 : 100 = %	0 : 100 = %
2500 mg/L	0 : 100 = %	0 : 100 = %

* Cu measured = 3.9 mg/L
 Grav. metric = 10.0 mg/L

Corrected
for error
(D.F.)