

AECD - 2056 (LADC - 76)

UNITED STATES ATOMIC ENERGY COMMISSION

THE MATHEMATICAL DEVELOPMENT OF THE END-POINT METHOD **MARKEN**

by

S. Frankel S. Goldberg

Los Alamos Scientific Laboratory

This document is reproduced as a project report and is without editorial preparation. The manuscript has been submitted to The Physical Review for possible ience and Technology Project publication.

FILE COPY Library of Congress TO BE RETURNED

DTIC QUALITY INSPECTED 5

04803

TRANSINE

1

MUN

Accesio	n For	
NTIS DTIC Unanno Justifici	CRA&I]
By Distribu	ition /	
A	vailability Code	S
Dist	Avail and/or Special	
A-1		

prof 1°1

Date of Manuscript: Date Declassified:

April 10, 1945 June 17, 1948

Issuance of this document does not constitute authority for declassification of classified copies of the same or similar content and title and by the same authors.

Technical Information Branch, Oak Ridge, Tennessee AEC, Oak Ridge, Tenn., 5-2-49--850-A1487

DESTRIBUTION

Approved for public releases Distaguant Universities

Printed in U.S.A. VPRICE/20 CENTS

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

THE MATHEMATICAL DEVELOPMENT OF THE END-POINT METHOD

By S. Frankel and S. Goldberg

ABSTRACT

The end-point method is mathematically developed and its application to the Milne kernel studied in detail. The general solution of the Wiener-Hopf integral equation is first obtained. The Milne kernel appears in applying this method to the integral equation describing the diffusion and multiplication of neutrons in multiplying and scattering media. The neutrons are treated as monochromatic, isotropically scattered and of the same total mean free path in all materials involved. Only problems with spherical symmetry are treated, these being reducible to equivalent infinite slab problems. Solutions are obtained for tamped and untamped spheres; in the former case both growing and decaying exponential asymptotic solutions in the tamper are treated in detail. Appendix I treats the effects of the approximations inherent in the end-point method (cf. LADC - 79). Appendix II gives the solution of the inhomogeneous Wiener-Hopf equation.

INTRODUCTION

The general development of the end-point method and some of its applications are described in LADC - 79. It is the purpose of this report to supplement this general description with an explicit mathematical development of the end-point method and a detailed study of its application to the Milne kernel. This is the kernel entering in the integral equation describing the diffusion and multiplication of neutrons in multiplying and scattering materials where the neutrons are treated as monochromatic, isotropically scattered, and of the same total mean free path in all materials involved. The end-point method to the treatment of problems in which the materials involved and the neutron distribution are both spherically symmetric, these problems being reducible to equivalent infinite-slab problems. In LADC - 79 it was shown that the end-point results may be applied loosely to problems of somewhat more complicated geometry and give more or less accurate approximations to the truth. These applications depend primarily on loose analogies rather than mathematical argument and will not be treated here.

Much of this report will be, in part, repetition of material treated in LADC 79. Here the emphasis will be primarily on the clear mathematical development of the methods of application presented there.

Chapter I

THE WIENER-HOPF METHOD

The integral equation,

$$n(x) = \int_{0}^{\infty} dx' n(x') K(x - x')$$
(1.0)
AECD - 2056

is known as the equation of Wiener and Hopf. With certain reasonable restrictions on the character of K and n this equation can be solved exactly. Before examining the method of solving this equation developed by Wiener and Hopf, it is useful to examine the simpler equation,

$$n(x) = \int_{-\infty}^{\infty} dx' \ n(x') \ K(x - x')$$
(1.1)

Since this equation is homogeneous, if $n_0(x)$ is a solution then $an_0(x)$ also satisfies the equation for any constant, a. Because of the infinite limits of integration and the "displacement" character of the kernel (K depends only on the difference, x - x'), $n_0(x - b)$ must also be a solution. If the solution, $n_0(x)$, is unique (except for a multiplicative factor) then $n_0(x - b) = an_0(x)$ for some a. Hence $n_0(x) = e^{kx}$. This suggests looking for exponential solutions of (1.1).

$$n(x) = e^{kx} = \int_{-\infty}^{\infty} dx' e^{kx'} K(x - x')$$
$$= e^{kx} \int_{-\infty}^{\infty} dy e^{-ky} K(y)$$
$$(1.2)$$
$$\int_{-\infty}^{\infty} dy e^{-y} K(y) = 1$$

Any solution of this "characteristic equation" gives a value of k for which e^{kx} satisfies equation 1.1. If there is more than one solution to the characteristic equation, then any linear combination of the exponentials determined by them will satisfy equation 1.1.

These considerations will be relevant to the study of the equation 1.0 if K decays rapidly for large |y|. If this is the case, for large x, equation 1.0 approximates equation 1.1, and it may be expected that with increasing x the solutions of equation 1.0 will approach asymptotically the exponential solutions of equation 1.1. If this is the case, the asymptotic exponential part of the solution of equation 1.0 may be separated from the remainder of the solution by Laplace or Fourier transformation. The use of the Laplace transform is further suggested by the fact that the left hand term of equation 1.2 is the Laplace transform of the kernel.

Taking the Laplace transform of equation 1.1 gives:

$$\int_{-\infty}^{\infty} dx \ e^{-kx} \ n(x) = \int_{-\infty}^{\infty} dx \ e^{-kx} \int_{-\infty}^{\infty} dx' \ n(x') \ K(x - x')$$
$$= \int_{-\infty}^{\infty} dx' \ n(x') \ e^{-kx'} \int_{-\infty}^{\infty} dy' \ e^{-ky'} \ K(y)$$
$$\int_{-\infty}^{\infty} dx' \ e^{-kx} \ n(x) \left(\int_{-\infty}^{\infty} dy' \ e^{-ky'} \ K(y) - 1 \right) = 0$$

This last equation shows that the Laplace transform of n(x) must vanish for all values of k which do not satisfy the characteristic equation 1.2.

An application of the same technique to equation 1.0 does not lead immediately to a factored equation because of the finite lower limit. To get around this difficulty Wiener and Hopf introduced the following trick.

Define n(x) = f(x) + g(x)

where $f(x) \equiv 0$ for x < 0

$$g(x) \equiv 0 \text{ for } x \ge 0$$

This permits writing equation 1.0 in the form

$$f(\mathbf{x}) + g(\mathbf{x}) = \int_{-\infty}^{\infty} d\mathbf{x}' f(\mathbf{x}') \mathbf{K}(\mathbf{x} - \mathbf{x}')$$

Now, taking the Laplace transform gives

$$\int_{-\infty}^{\infty} dx f(x) e^{-kx} + \int_{-\infty}^{\infty} dx g(x)e^{-kx} = \int_{-\infty}^{\infty} dx e^{-kx} \int_{-\infty}^{\infty} dx' f(x') K(x - x')$$
$$= \int_{-\infty}^{\infty} dx' e^{-kx'} f(x') \int_{-\infty}^{\infty} dy e^{-ky} K(y)$$
$$F(k) \equiv \int_{-\infty}^{\infty} dx f(x) e^{-kx}$$
$$G(K) \equiv \int_{-\infty}^{\infty} dx g(x) e^{-kx}$$
$$\underline{K}(k) \equiv \int_{-\infty}^{\infty} dx K(x)e^{-kx}$$
$$G(k) = F(k) \left(\underline{K}(k) - 1\right) \equiv F(k) P(k)$$

we have

Defining

This equation will hold for any value of k for which all three integrals exist. We therefore impose conditions on the kernel and solution of equation 1.0, which ensure the existence of a suitable region in the complex plane in which all three integrals exist. We require that K(y) decay at least as rapidly as an exponential for large (positive or negative) y.

$$K(y) = c(e^{-C}|y|), c > 0.$$
(1.4)

Then K(k) will exist for -c < R(k) < c. We further assume that

$$f(x) = c(e^{dx}), d < c$$
 (1.5)

The kernels of primary interest are symmetric. For these, if the "largest" value of c satisfying equation 1.4 is chosen, equation 1.5 is not a restrictive condition, since f(x) must approach asymptotically an exponential, e^{kx} , for some k satisfying $\underline{K}(k) = 1$ and therefore within the range of convergence of $\underline{K}(k)$. The form of equation 1.3 clearly requires that g(x) decay (for large negative x) at least as fast as e^{Cx} . Thus G(k) exists for all k having R(k) < c. The three integrals will therefore all exist throughout a vertical strip in the complex k-plane defined by d < R(k) < c.



Figure 1.

(1.3)

Within this "common strip" all three integrals are convergent and equation 1.3 must be satisfied. Outside this strip the nonconvergent integrals will be defined by analytic extension (and need not be analytic) in such a way that the equation is still satisfied.

Within and to the right of the common strip, F(k) exists and is analytic. [It is clear from its definition that in this range any derivative of F(k) exists.] Similarly within and to the left of the strip, G(k) exists and is analytic. K(k), hence also P(k), exists and is analytic within the strip but may have singularities on either side of it. We make the further assumption that F(k) and G(k) have no roots in their respective regions of analyticity. (Cf. Paley and Wiener, Fourier Transforms, p. 51). We further require that there exist a sub-strip within the common strip within which P(k) has no roots. [This must be true if P(k) has only a finite number of zeros in the common strip. This will actually be the case, Cf. Titchmarsh, Fourier Integrals, p. 339.]

We have now a sub-strip within which $\log P(k)$ is analytic; within which, and to the right, $\log F(k)$ is analytic; within which, and to the left, $\log G(k)$ is analytic, and within which the three satisfy

$$\log P(k) = \log G(k) - \log F(k)$$

This equation will be satisfied throughout the plane by the analytic extensions.

It is now easy to find functions, F and G, satisfying this equation and the analyticity conditions. For values of k within the sub-strip we express $\log P(k)$ by means of a Cauchy integral:

$$\log P(k) = (1/2\pi i) \int_{C} \frac{dk'}{k' - k} \log P(k')$$
$$= (1/2\pi i) \int_{R} \frac{dk'}{k' - k} \log P(k')$$
$$+ (1/2\pi i) \int_{L} \frac{dk'}{k' - k} \log P(k')$$

where the contour of integration consists of two vertical lines in the sub-strip, one running up to the right of k, the other down to its left.



Figure 2.

We have now decomposed log P(k) into two parts, one certainly analytic within the strip and to the left, the other within and to the right. These may be identified with log G(k) and $-\log F(k)$, and give a solution to the equation 1.0.

$$\log F(k) = -\frac{1}{2\pi i} \int_{L} \frac{dk'}{k' - k} \log P(k) + \text{constant}$$

$$\log G(k) = \frac{1}{2\pi i} \int_{R} \frac{dk'}{k' - k} \log P(k') + \text{constant}$$
(1.6)

This contour integral representation of log F(k) determines F(k), hence also f(x).

$$f(x) = \frac{1}{2\pi i} \int_{\delta - i\infty}^{\delta + i\infty} e^{kx} F(k) dk \qquad (1.7)$$

where δ is chosen to make F(k) regular along the contour. In particular, δ may be taken in the substrip. Since F(k) is analytic to the right of the sub-strip, the contour may be translated to the right as far as desired. For negative values of x this may be used to show that f(x) vanishes.

If f(x) contains a term Ae^{kox} (e.g., as its asymptotic solution), then its Laplace transform, F(k) will contain a corresponding term.

$$\int_{0}^{\infty} dx e^{-kx} A e^{k_0 x} = A/(k - k_0)$$

Thus a pure exponential term in f(x) manifests itself in F(k) as a simple pole, and the coefficients of the two may be identified. The coefficient of the singularity is most easily determined by expanding log F(k) about the singularity.

$$\log F(k) = -\log(k - k_0) + \log A + 0(k - k_0)$$

The asymptotic solution will be determined by all of the singularities of F(k) on the imaginary axis and in the right half-plane. If there are no singularities on or to the right of the imaginary axis the solution, f(x), will approach zero asymptotically. A more useful asymptotic solution however, will be that determined by the first singularities to the left of the imaginary axis:

An important special case of this general treatment is that for which the kernel, K(y), is symmetric and for which the characteristic equation has only a single pair of conjugate roots on the imaginary axis. If these two roots are at $\pm ik_0$, then the solution will be of the form

$$F(k) = B\left[\sin k_0 (x + x_0) + h(x)\right], h(x) \rightarrow 0 \text{ as } x \rightarrow +\infty$$
(1.8)

Since the equation is homogeneous, B is undetermined; x_0 , however, can be evaluated.

$$F(k) = \int_{0}^{\infty} dx \ e^{-kx} \ B\left[\sin k_{0}(x + x_{0}) + h(x)\right]$$

=
$$\int_{0}^{\infty} dx \ e^{-kx} \frac{B}{2i} \left[e^{ik_{0}(x + x_{0})} - e^{-ik_{0}(x + x_{0})} + 2ih(x)\right]$$

=
$$\frac{B}{2i} \left(\frac{e^{ik_{0}x_{0}}}{k - ik_{0}} - \frac{e^{-ik_{0}x_{0}}}{k + ik_{0}} + 2iH(k)\right)$$

In the neighborhood of \pm ik_o, H(k) is finite. We expand log F(k) near these two poles,

$$\log F(ik_0 + \epsilon) = \log \frac{B}{2i} + i k_0 x_0 - \log \epsilon + 0(\epsilon)$$
$$\log F(-ik_0 + \epsilon) = \log \frac{-B}{2i} - i k_0 x_0 - \log \epsilon + 0(\epsilon)$$

$$\lim \left[\log F(ik_0 + \epsilon) - \log F(-ik_0 + \epsilon) \right] = \log (-1) + 2i k_0 x_0$$

$$\epsilon \rightarrow 0$$

 $\log F(k) = \log G(k) - \log P(k)$

$$= \frac{1}{2\pi i} \int_{\mathbf{R}} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \log \mathbf{P}(\mathbf{k}') - \log \mathbf{P}(\mathbf{k})$$
$$\lim_{\epsilon \to 0} \left[\log \mathbf{P}(i\mathbf{k}_{0} + \epsilon) - \log \mathbf{P}(-i\mathbf{k}_{0} + \epsilon) \right] = \log \left[\frac{\mathbf{P}'(i\mathbf{k}_{0})}{\mathbf{P}'(-i\mathbf{k}_{0})} \right] = \log (-1)$$

since K(y) is even, K(k) and P(k) are even; P'(k) is odd.

$$2 ik_{0}x_{0} = \frac{1}{2\pi i} \int \mathbf{R} \, d\mathbf{k}' \, \log \, \mathbf{P}(\mathbf{k}') \left[\frac{1}{\mathbf{k}' - i\mathbf{k}_{0}} - \frac{1}{\mathbf{k}' + i\mathbf{k}_{0}} \right]$$
$$x_{0} = \frac{1}{2\pi i} \int \mathbf{R} \, \frac{d\mathbf{k}'}{\mathbf{k}'^{2} + \mathbf{k}_{0}^{2}} \, \log \, \mathbf{P}(\mathbf{k}')$$
(1.9)

The two terms, log (-1), have been neglected since the form of the solution 1.8 is unchanged by the addition of a multiple of π to $k_0 x_0$. The evaluation of x_0 completes the determination of the asymptotic form of the solution equation 1.8. x_0 is expressed in equation 1.9 as a single integral, which in many cases must be evaluated numerically. To get the complete solution requires two integrations, one to evaluate log F(k) by equation 1.6, another to get f(x) by (1.7).

Two-Medium Problems

A more general problem that can be treated by the Wiener-Hopf technique is

$$n(x) = \int_{-\infty}^{\infty} dx' K'(x - x') n(x') + \int_{0}^{\infty} dx' K(x - x') n(x')$$

Breaking up n(x) as before and taking the Laplace transform of the resulting equation gives

$$F(k) + G(k) = K(k) F(k) + K'(k) G(k)$$

where the notation is the same as before. This may be written as

$$\mathbf{G}(\mathbf{k}) = \mathbf{F}(\mathbf{k}) \left(\frac{1 - \underline{\mathbf{K}}(\mathbf{k})}{\underline{\mathbf{K}}'(\mathbf{k}) - 1} \right) \stackrel{=}{=} \mathbf{F}(\mathbf{k}) \mathbf{P}(\mathbf{k})$$

This is now of the same form as equation 1.3. The rest of the treatment proceeds in the same way. With this more complicated form for P(k) there may be a greater number of singularities of log P(k), leading to a larger number of independent solutions. In particular it is no longer necessary to require that g(x) decay exponentially away from the boundary.

An important special case of this two-medium problem is that for which K(y) and K'(y) differ only by a multiplicative factor. This case will be treated extensively in the second chapter.

The Wiener-Hopf technique may be further extended to permit the solution of inhomogeneous displacement integral equations. This method is outlined in Appendix II.

Chapter II

APPLICATION TO NEUTRON PROBLEMS

In this chapter we treat the applications of the Wiener-Hopf method (combined with some approximations) to problems concerning the spatial distribution and time dependence of neutrons in spheres of multiplying and scattering materials. It will be shown that such problems, with suitable physical approximations, can be represented by integral equations closely analogous to the Wiener-Hopf equation. By making suitable mathematical approximations (the "end-point method") fairly accurate solutions to these equations can be gotten from the corresponding Wiener-Hopf solutions.

We make the following physical approximations:

A) We consider only one neutron velocity; hence for each material only one value for each cross section.

B) We treat all collision processes as isotropic. (Anisotropy of elastic scattering can be treated to a limited extent. It can be shown that if this anisotropy is neglected and the transport average used for the elastic scattering cross section quite accurate results will be obtained. Cf. LADC - 79 and MT - 26.)

C) The total mean free path will be taken to be the same for all materials involved.

D) The neutron distribution will be treated as a continuum. It will be taken to be spherically symmetric and of stable spatial distribution. These three conditions will certainly be good approximations if the neutron distribution has lived through many generations and consists of a sufficient number of neutrons to make statistical fluctuation negligible.

We adopt the following notation:

 $\sigma_{\rm f}$ is the fission probability per unit path length. (It is therefore the product of the fission cross section and the number of nuclei per unit volume.) Similarly,

 $\sigma_{\rm S}$ is the scattering probability per unit path length.

 σ_a is the absorption probability per unit path length.

 $\sigma = \sigma_{\rm f} + \sigma_{\rm s} + \sigma_{\rm a}$

v is the mean number of neutrons emerging from a fission process.

 $F = 1 + f = \frac{\nu \sigma_f + \sigma_S}{\sigma}$ is therefore the mean number of neutrons emerging from a collision.

v is the neutron velocity.

n(r, t) is the neutron density at point r at time t.

We express the neutron density at (\underline{r}, t) as an integral over all points at which these neutrons may have suffered their last collisions.

$$\mathbf{v} \mathbf{n}(\underline{\mathbf{r}}, \mathbf{t}) = \int d\underline{\mathbf{r}}' \quad \sigma \mathbf{v} \mathbf{F}(\underline{\mathbf{r}}') \mathbf{n}\left(\underline{\mathbf{r}}', \mathbf{t} - \frac{|\mathbf{r} - \mathbf{r}'|}{\mathbf{v}}\right) \frac{1}{4\pi(\mathbf{r} - \mathbf{r}')^2} \mathbf{e}^{-\sigma |\mathbf{r} - \mathbf{r}'|}$$
(2.1)

We look for solutions of the form

$$n(\underline{\mathbf{r}},t) = n(\underline{\mathbf{r}}) e^{\gamma_0 t}$$

The integral equation 2.1, then takes the form:

$$\mathbf{n}(\mathbf{r}) = \int d\mathbf{r} \, \boldsymbol{\sigma} \, \mathbf{F}(\mathbf{r}') \, \mathbf{n}(\mathbf{r}') \, \frac{1}{4\pi(\mathbf{r} - \mathbf{r}')^2} \, \mathbf{e}^{-(\sigma + \gamma_0/\mathbf{v}) |\mathbf{r} - \mathbf{r}'|}$$

We now rescale <u>r</u>, taking as the unit of length the mean attenuation distance, $1/(\sigma + \gamma_0/v)$.

$$\mathbf{x} = \mathbf{r} \left(\sigma + \gamma_0 / \mathbf{v} \right)$$

$$n(\underline{x}) = \frac{1}{1 + \gamma_{0}/\sigma_{v}} \int d\underline{x}' F(\underline{x}') n(\underline{x}') \frac{e^{-|x - x'|}}{4\pi(x - x')^{2}}$$

Defining $\gamma = \gamma_0 / \sigma v$ gives the three-dimensional integral equation.

$$\mathbf{n}(\underline{\mathbf{x}}) = \frac{1}{1+\gamma} \int d\underline{\mathbf{x}}' \ \mathbf{F}(\underline{\mathbf{x}}') \mathbf{n}(\underline{\mathbf{x}}') \frac{\mathbf{e}^{-|\mathbf{x}-\mathbf{x}'|}}{4\pi(\mathbf{x}-\mathbf{x}')^2}$$
(2.2)

- 10

If we now introduce polar coordinates, $\mathbf{x}' = (\mathbf{r}', \boldsymbol{\phi}', \boldsymbol{\theta}')$,

taking the point <u>x</u> on the polar axis we may make use of the assumed spherical symmetry of $n(\underline{x}')$ to reduce equation 2.2 to an equation in one dimension.

$$\mathbf{n}(\mathbf{r}) = \frac{1}{1+\gamma} \int \mathbf{r}'^2 \, \mathrm{d}\mathbf{r}' \, \mathbf{F}(\mathbf{r}') \, \mathbf{n}(\mathbf{r}') \, \iint \, \mathrm{d}\phi \sin \theta \, \mathrm{d}\theta \, \frac{\mathrm{e}^{-(\mathbf{r}^2 + \mathbf{r}'^2 - 2\mathbf{r}\mathbf{r}' \, \cos \theta)^{1/2}}{4\pi(\mathbf{r}^2 + \mathbf{r}'^2 - 2\mathbf{r}\mathbf{r}' \, \cos \theta)}$$

Taking $\mu = \cos \theta$, $l^2 = r^2 + r'^2 - 2rr' \cos \theta$

$$\int_{0}^{2\pi} d\phi \int_{0}^{\pi} \sin\theta \, d\theta \, \frac{e^{-(r^{2} + r'^{2} - 2rr'\cos\theta)^{\frac{1}{2}}}}{4\pi(r^{2} + r'^{2} - 2rr'\cos\theta)} = \frac{1}{2} \int_{-1}^{1} d\mu \frac{e^{-1}}{1^{2}}$$
$$= \frac{1}{2} \int_{|r - r'|}^{r + r'} \frac{1}{rr'} \frac{1}{r'} \frac{e^{-1}}{1^{2}}, \left(d\mu = -\frac{1}{rr'}\right)$$
$$= \frac{1}{2rr'} \left[E(|r - r'|) - E(r + r') \right]$$

where $E(s) = \int_{s}^{\infty} \frac{e^{-t} dt}{t}$

$$\mathbf{r} \ \mathbf{n}(\mathbf{r}) = \frac{1}{2(1+\gamma)} \int_{0}^{\infty} d\mathbf{r}' \ \mathbf{F}(\mathbf{r}') \ \mathbf{r}' \ \mathbf{n}(\mathbf{r}') \left[\mathbf{E}(|\mathbf{r} - \mathbf{r}'|) - \mathbf{E}(\mathbf{r} + \mathbf{r}') \right]$$
(2.3)

If we now define $u(r) \equiv r n(r)$ and treat u(r) as an odd function, and F(r) as an even function of r [no meaning has previously been assigned to negative values of r or to the corresponding <math>n(r) and F(r)], we may write equation 2.3 in the form:

$$u(\mathbf{r}) = \frac{1}{2(1+\gamma)} \int_{-\infty}^{\infty} d\mathbf{r}' \ F(\mathbf{r}') \ u(\mathbf{r}') \ E(|\mathbf{r} - \mathbf{r}'|)$$
(2.4)

If instead of assuming the material and neutron distribution spherically symmetric, we take both as functions of only one Cartesian coordinate, z, equation 2.2 may be reduced to an equation in one dimension as follows:

$$n(z) = \frac{1}{1+\gamma} \int dz' F(z') n(z') \int \int dx' dy' \frac{e^{-\left[(z-z')^2 + (y-y')^2 + (x-x')^2\right]^{\frac{1}{2}}}}{4\pi \left[(z-z')^2 + (y-y')^2 + (x-x')^2\right]}$$
$$= \frac{1}{1+\gamma} \int dz' F(z') n(z') \int_{0}^{2\pi} d\phi \int_{0}^{\infty} \rho d\rho \frac{e^{-1}}{4\pi l^2}$$

where $l^2 = (z - z')^2 + \rho^2$, $l dl = \rho d\rho$

$$n(z) = \frac{1}{2(1+\gamma)} \int dz' F(z') n(z') E(|z - z'|)$$
(2.5)

A comparison of equations 2.4 and 2.5 shows that the sphere problem 2.4 may be identified with a slab problem 2.5 in which the distribution of materials F(z) across the slab is the same as that along a diameter of the sphere. Any odd solution of the slab problem, n(z), may be identified with the quantity u(r) in the sphere problem and conversely. The "fundamental mode" of the sphere for which n(r) is everywhere positive corresponds to the "first harmonic" of the slab in which the neutron density takes on apparently meaningless negative values. For this reason, and because higher modes may be superimposed on the fundamental, we will treat the neutron density, n(z), as a real quantity which may have either sign.

For a tamped sphere of core radius a, outer tamper radius b, and mean attenuation distances, the integral equation 2.4 takes the form

$$u(r) = \frac{1 + f_t}{1 + \gamma} - \int_{-b}^{a} dr' u(r') \frac{1}{2} E(|r - r'|)$$

$$+ \frac{1 + f_{c}}{1 + \gamma} \int_{a}^{a} d\mathbf{r}' \ \mathbf{u}(\mathbf{r}') \frac{1}{2} \mathbf{E} (|\mathbf{r} - \mathbf{r}'|)$$
$$+ \frac{1 + f_{t}}{1 + \gamma} \int_{a}^{b} d\mathbf{r}' \ \mathbf{u}(\mathbf{r}') \frac{1}{2} \mathbf{E} (|\mathbf{r} - \mathbf{r}'|)$$

where f_c and f_t are the values of f in core and tamper respectively. This equation differs from the Wiener-Hopf equation in having four boundaries instead of one (or two for an untamped sphere). With more than one boundary no exact solution is known. We therefore resort to an approximation, namely to treat the behaviour of the solution near each boundary as if no other boundaries existed. It was shown in the first chapter that the solution of the one-boundary problem approaches, at large distances from the boundary, a solution of the problem with infinite limits. It is reasonable to expect that the solution of a two-boundary problem in which the boundaries are very far apart will behave in some middle region as a solution of the infinite-limits equation. If this is the case, we have only to combine two one-boundary solutions in such a way that their asymptotic components coincide. In a many-boundary problem, e.g., the tamped sphere, we apply this recipe in each region. This approximation method, the 'end-point method'', would seem, from the above argument, reasonably accurate only if the distances between boundaries are many mean attenuation distances. It is shown in Appendix I that the limit of reasonable accuracy is actually a few tenths of a mean attenuation distance. There is therefore good reason to believe that for sizes larger than that, the end-point method is sufficiently accurate.

In order to apply the end-point method we must first study the one-boundary problem with the "Milne kernel",

$$K(y) = c \frac{1}{2} \cdot E(|y|)$$

This kernel with c = 1 occurs in "the equation of E. A. Milne" describing the flow of radiation through the outermost layers of a star. We will, however, refer to it as the "Milne kernel" for all positive values of c. The general equation we have to study is then

$$n(x) = c' \int_{-\infty}^{0} dx' n(x') \frac{1}{2} E(|x - x'|) + c \int_{0}^{\infty} dx' n(x') \frac{1}{2} E(|x - x'|)$$

 $c = (1 + f)/(1 + \gamma).$

Several cases arise. For a free surface, either the outer surface of a tamper or the surface of an untamped sphere, we take c' = 0. For an interface we take both c and c' positive. For the core material, c must be greater than 1 (f> γ); in the tamper, c - 1 may be of either sign.

We first treat the free-surface case.

$$n(x) = c \int_{0}^{\infty} dx' n(x') \frac{1}{2} E (|x - x'|)$$

The characteristic equation is

c
$$\int_{-\infty}^{\infty} dy \frac{1}{2} E(|y|) e^{-ky} = (c/2) \int_{0}^{\infty} dy (e^{-ky} + e^{ky}) \int_{1}^{\infty} \frac{ds}{s} e^{-ys}$$

= $(c/2) \int_{1}^{\infty} \frac{ds}{s} \left(\frac{1}{s+k} + \frac{1}{s-k}\right)$
= $c \int_{1}^{\infty} \frac{ds}{s^2 - k^2}$

$$= \frac{c}{2k} \log\left(\frac{1+k}{1-k}\right) = \frac{c}{k} \tanh^{-1} k = 1$$

If c < 1 we have two real roots, $\pm k_0$ such that $c = k/tanh {}^1k_0$. If c > 1 we have two imaginary roots, $\pm i k_0$, such that $c = k_0/tan {}^{-1}k_0$. In either case it can be shown that the characteristic equation has only two roots. In the latter case the asymptotic solution is a sinusoidal function of k_0x , in the former, a hyperbolic function. We will represent the phase of the asymptotic solution by the "extrapolated endpoint," x_0 , such that the asymptotic solution is the sine or hyperbolic sine of $k_0(x + x_0)$. We now follow through, explicitly, the method of solution outlined in Chapter 1.

$$\begin{aligned} n(x) &\equiv f(x) + g(x) = c \quad \int_{-\infty}^{\infty} dx' f(x') \frac{1}{2} E(|x - x'|) \\ f(x) &= o \text{ for } x < o \\ g(x) &\equiv o \text{ for } x \ge o \\ F(k) + G(k) &= \int_{-\infty}^{\infty} dx n(x) e^{-kx} = \int_{-\infty}^{\infty} dx e^{-kx} \int dx' f(x') \frac{c}{2} E(|x - x'|) \\ &= \int_{-\infty}^{\infty} dx' f(x') e^{-kx'} \int_{-\infty}^{\infty} dy e^{-ky} \frac{c}{2} E(|y|) \\ &= F(k) \frac{c}{2k} \log\left(\frac{1+k}{1-k}\right) \\ G(k) &= F(k) \left\{ \frac{c}{2k} \log\left(\frac{1+k}{1-k}\right) - 1 \right\} = F(k) P(k) \end{aligned}$$

P(k) has singularities only at ± 1 . These singularities are branch points so that to make the function explicit we introduce cuts lying along the real axis from $-\infty$ to -1 and from +1 to $+\infty$. We treat first the case c > 1. The two roots of P(k) are then pure imaginary, $\pm ik_0$. The singularities of log P(k) are ± 1 and $\pm ik_0$. We look for a log F(k), analytic to the right of the imaginary axis [corresponding to the sinusoidal asymptotic solution, f(x)], and a log G(k), analytic to the left of +1 [corresponding to a g(x) decaying somewhat faster than e^{-x}], and satisfying

$$\log P(k) = \log G(k) - \log F(k)$$
(2.6)

The "sub-strip" in which all three of these quantities are analytic is 0 < R(k) < 1. We therefore break up log P(k) by means of a Cauchy integral along a contour running up and down in this strip and enclosing k, and (except for a common constant) identify log G(k) and -log F(k) with the two parts of the integral.

$$\log P_{R}(k) = \frac{1}{2\pi i} \int_{R} \frac{dk'}{k' - k} \log P(k') = \log G(k) + \text{constant},$$
$$\log P_{L}(k) = -\frac{1}{2\pi i} \int_{L} \frac{dk'}{k' - k} \log P(k') = \log F(k) + \text{constant}.$$

We simplify $\log P_{\mathbf{R}}(\mathbf{k})$ by deforming the right contour to enclose the right-hand cut.





Figure 3.

$$\log P_{\mathbf{R}}(\mathbf{k}) = \frac{1}{2\pi i} \int_{\infty}^{\infty} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \log \left[\frac{c}{2\mathbf{k}'} \left(\log \frac{1 + \mathbf{k}'}{1 - \mathbf{k}'} \right) - 1 \right] \left[I(\log) = \pi i \rightarrow 0 \right]$$
$$+ \frac{1}{2\pi i} \int_{1}^{\infty} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \log \left[\frac{c}{2\mathbf{k}'} \left(\log \frac{1 + \mathbf{k}'}{\mathbf{k}' - 1} + \pi i \right) - 1 \right] \left[I(\log) = 0 \rightarrow + \pi i \right]$$
$$= \frac{1}{\pi} \int_{1}^{\infty} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \tan^{-1} \left(\frac{\pi/2}{\frac{1}{2} \log \frac{\mathbf{k}' + 1}{\mathbf{k}' - 1} - \frac{\mathbf{k}'}{\mathbf{c}}} \right) \left[\tan^{-1} = 0 \rightarrow \pi \right]$$

Here the tan⁻¹ rises from 0 at k' = 1 to π at k' = + ∞ (as indicated by the bracketed expressions). Substituting k' = 1/s,

$$\log P_{R}(k) = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1 - ks)} T_{c'}$$

where

$$T_c = \tan^{-1} \left(\frac{\pi/2}{\tanh^{-1}s - 1/cs} \right) \begin{array}{c} T_c = \pi & s = 0 \\ = 0 & s = 1 \end{array}$$

$$\log P_{\rm R}(k) = \frac{1}{\pi} \int_0^1 \frac{{\rm ds}}{{\rm s}} T_{\rm C} + \frac{k}{\pi} \int_0^1 \frac{{\rm ds}}{1 - k{\rm s}} T_{\rm C}$$
(2.7)

Here and throughout this treatment we encounter logarithmically infinite constants. A slight modification of our procedure $\begin{bmatrix} to make P(k) \rightarrow 1 as |k| \rightarrow \infty \end{bmatrix}$ suffices to avoid this embarrassment. The present treatment is somewhat simpler, though formally less rigorous.

We simplify log $\mathtt{PL}(k)$ by a corresponding deformation of the left contour.



$$\log P_{L}(k) = \frac{1}{2\pi i} \left\{ \int_{-\infty}^{1} \log \left[\frac{c}{2k'} \left(\log \frac{|k'| - 1}{1 - k'} + \pi i \right) - 1 \right] \left[I(\log) = \pi i \rightarrow 2\pi i \right] \right. \\ \left. + \int_{-1}^{0} (2\pi i) + \int_{0}^{ik_{0}} (2\pi i) + \int_{-ik_{0}}^{0} (-2\pi i) + \int_{0}^{-1} (-2\pi i) \right. \\ \left. + \int_{-1}^{\infty} \log \left[\frac{c}{2k'} \left(\log \frac{|k'| - 1}{1 - k'} + \pi i \right) - 1 \right] \right\} \frac{dk'}{k' - k} \left[I(\log) = -2\pi i \rightarrow -\pi i \right] \\ \left. = \frac{1}{2\pi i} \int_{-\infty}^{-1} \frac{dk'}{k' - k} \log \frac{\frac{c}{2k'} \left(\log \frac{|k'| - 1}{1 - k'} + \pi i \right) - 1}{\frac{c}{2k} \left(\log \frac{|k'| - 1}{1 - k'} - \pi i \right) - 1} \left[\log = 2\pi \rightarrow 4\pi \right] \\ \left. + \log \frac{k}{1 + k} + \log \frac{k - ik_{0}}{k} - \log \frac{k}{k + ik_{0}} - \log \frac{1 + k}{k} \right]$$

Letting r = -k'

$$-\log P_{L}(k) = -\frac{1}{\pi} \int_{1}^{\infty} \frac{dr}{r+k} \tan^{-1} \frac{\pi/2}{\frac{r}{c} \frac{1}{2} \log \frac{r+1}{r-1}} \left[\tan^{-1} = 2\pi - \pi \right]$$
$$+ \log \frac{k^{2} + k_{0}^{2}}{(2+k)^{2}}$$

Letting
$$s = \frac{1}{r}$$
 we have

$$-\log P_{L}(k) = -\frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1+ks)} \left[2\pi + \tan^{-1} \frac{\pi/2}{1/cs - \tanh^{-1}s} \right] \left[\tan^{-1} = -T_{c} = -\pi - 0 \right] \\ + \log \frac{k^{2} + k_{0}^{2}}{(1+k)^{2}} \\ = -2 \int_{0}^{1} \frac{ds}{s} + \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1+ks)} T_{c} + \log (k^{2} + k_{0}^{2}) \\ \log P_{L}(k) = 2 \int_{0}^{1} \frac{ds}{s} \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s} T_{c} - \log (k^{2} + k_{0}^{2}) + \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1+ks} T_{c}$$
(2.8)

Combining those two expressions, 2.7 and 2.8, with

$$\log P(k) \equiv \log \left(\frac{c}{2k} \log \frac{1+k}{1-k} - 1\right) = \log P_R(k) - \log P_L(k)$$
(2.6)

gives

$$\log \left(\frac{c}{2k} \log \frac{1+k}{1-k} - 1\right) = \frac{2}{\pi} \int_{0}^{1} \frac{ds}{s} (T_{c} - \pi) + \log (k^{2} - k_{0}^{2}) + \frac{2k^{2}}{\pi} \int_{0}^{1} \frac{sds}{1-k^{2}s^{2}} T_{c} (2.9)$$

Taking the limit as k - 0 we get

$$\frac{1}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{\mathrm{s}} \left(\mathrm{T}_{\mathrm{c}} - \pi \right) = \frac{1}{2} \log \frac{\mathrm{c} - 1}{\mathrm{k}_{\mathrm{o}}^{2}}$$
(2.10)

and equation 2.9 becomes

$$\frac{k1}{\pi} \int_{0}^{1} \frac{sds}{1 - k^{2}s^{2}} = \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 - ks} T_{c} - \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 + ks} T_{c}$$

$$= -\frac{1}{2} \log \left(\frac{k^{2} + k_{0}^{2}}{k_{0}^{2}}\right) - \frac{1}{2} \log \frac{c - 1}{\frac{c}{2k} \log \frac{1 + k}{1 - k} - 1}$$
(2.11)

Dividing by k^2 and again letting k - 0,

$$\frac{1}{\pi} \int_0^1 \text{sds } T_c = - \frac{1}{2k_0^2} + \frac{c}{6(c-1)}$$

We now subtract the (infinite) constant, $2 \int_0^1 \frac{ds}{s} - \frac{1}{\pi} \int_0^1 \frac{ds}{s} T_c - \log B$, from $\log P_R(k)$ and $\log P_L(k)$ to give log G(k) and log F(k).

$$\log F(k) = -\log(k^{2} + k_{0}^{2}) + \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 + ks} T_{c} + \log B;$$

$$\log G(k) = \frac{2}{\pi} \int_{0}^{1} \frac{ds}{s} (T_{c} - \pi) + \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 - ks} T_{c} + \log B,$$
$$= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 - ks} T_{c} + \log \frac{B(c - 1)}{k_{0}^{2}}$$

We now determine x_0 and the value of B required to give the asymptotic sine wave in f(x) unit amplitude.

å

$$\begin{split} f(x) &= \sin k_0 (x + x_0) + h(x) & h(x) \longrightarrow 0 \text{ as } x \longrightarrow + \bullet \\ F(k) &= \frac{e^{ik_0 x_0}}{2i(k - ik_0)} - \frac{e^{-ik_0 x_0}}{2i(k + ik_0)} + H(k) = \frac{k \sin k_0 x_0 + k_0 \cos k_0 x_0}{k^2 + k_0^2} + H(k) \\ &\log F(ik_0 + \epsilon) = -\log(2i) + ik_0 x_0 - \log \epsilon + 0(\epsilon) \\ &\log F(-ik_0 + \epsilon) = -\log(-2i) - ik_0 x_0 - \log \epsilon + 0(\epsilon) \\ &\lim_{\epsilon \to 0} \left[\log F(ik_0 + \epsilon) - \log F(-ik_0 + \epsilon) \right] = \log(-1) + 2ik_0 x_0 \\ &\epsilon + 0 \\ &= \lim_{\epsilon \to 0} \left[\frac{ik_0 + \epsilon}{\pi} \int_0^1 \frac{ds T_c}{1 + (ik_0 + \epsilon)s} - \log (2ik_0 \epsilon + \epsilon^2) \\ &- \frac{ik_0 + \epsilon}{\pi} \int_0^1 \frac{ds T_c}{1 + (-ik_0 + \epsilon)s} + \log(-2ik_0 \epsilon + \epsilon^2) \right] \\ &= \frac{ik_0}{\pi} \int_0^1 ds T_c \left(\frac{1}{1 + ik_0 s} + \frac{1}{1 - ik_0 s} \right) + \log(-1) \\ &x_0 = \frac{1}{\pi} \int_0^1 \frac{ds}{1 + k_0^2 s^2} T_c \end{split}$$

Now adding the two values of log F gives

$$\log F(ik_{0} + \epsilon) + \log F(ik_{0} + \epsilon) = -2 \log (2\epsilon) + 0(\epsilon),$$

$$= 2 \log (2k_{0}\epsilon) + \frac{ik_{0}}{\pi} \int_{0}^{1} ds T_{c} \left(\frac{1}{1 + ik_{0}s} \cdot \frac{1}{1 - ik_{0}s}\right) + 2 \log B + 0(\epsilon)$$

$$= -2 \log (2k_{0}\epsilon) + \frac{2k_{0}2}{\pi} \int_{0}^{2} \frac{s ds}{1 + k_{0}^{2}s^{2}} T_{c} + 2 \log B + 0(\epsilon).$$

$$\log B = \log k_{0} = \frac{k_{0}^{2}}{\pi} \int_{0}^{1} \frac{s ds}{1 + k_{0}^{2}s^{2}} T_{c}$$

This integral may be evaluated by allowing $k \mbox{ to approach } ik_0$ in equation 2.11:

$$-\frac{k_{0}^{2}}{\pi} \int_{0}^{1} \frac{s \, ds}{1+k_{0}^{2} s^{2}} T_{c} = \lim_{\epsilon \to 0} \left[-\frac{1}{2} \log \left(\frac{2ik_{0}\epsilon}{k_{0}^{2}} \right) - \frac{1}{2} \log \frac{c-1}{-\frac{1}{ik_{0}}} \left(1 - \frac{c}{1+k_{0}^{2}} \right) \right] \epsilon$$

$$= -\frac{1}{2} \log \frac{2(c-1)}{1 - \frac{c}{1+k_{0}^{2}}}$$

$$\log B = \frac{1}{2} \frac{k_{0}^{2} \left(1 - \frac{c}{1+k_{0}^{2}} \right)}{2(c-1)}$$

$$\log F(k) = \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1+ks} T_{c} - \log \left(k^{2} + k_{0}^{2} \right) + \frac{1}{2} \log \frac{k_{0}^{2} \left(1 - \frac{e}{1+k_{0}^{2}} \right)}{2(c-1)}$$

$$F(k) = \frac{k_{0}}{k^{2} + k_{0}^{2}} \sqrt{\frac{1 - c/(1+k_{0}^{2})}{2(c-1)}} e^{\frac{k}{\pi}} \int_{0}^{1} \frac{ds}{1+ks} T_{c}.$$

$$H(k) \frac{1}{k^{2} + k_{0}^{2}} \left(k_{0} \sqrt{\frac{1 - c/(1+k_{0}^{2})}{2(c-1)}} e^{\frac{k}{\pi}} \int_{0}^{1} \frac{ds}{2+ks} T_{c} - k \sin k_{0} x_{0} - k_{0} \cos k_{0} x_{0} \right)$$

We can evaluate H (o), the total area of h(x), and $\frac{-H(0)}{H(0)}$, its "mean length",

$$H(o) = \frac{1}{k_0} \left(\sqrt{\frac{1 - c/(1 + k_0^2)}{2(c - 1)}} - \cos k_0 x_0 \right)$$
$$\frac{-H'(o)}{H(o)} = \frac{1}{H(o)k_0^2} \left(\sin k_0 x_0 - k_0 \sqrt{\frac{1 - c/(1 + k_0^2)}{2(c - 1)}} - \frac{1}{\pi} \int_0^1 ds T_c \right)$$

Making use of the formula

$$n(o) = \lim_{k \to \infty} k \int_{0}^{\infty} dx \ n(x) \ e^{-kx} = \lim_{k \to \infty} k \ F(k).$$

we get

$$n(0 = \lim_{k \to \infty} \frac{kk_0}{k^2 + k_0^2} \sqrt{\frac{1 - c/(1 + k_0^2)}{2(c - 1)}} e^{\frac{k}{\pi} \int_0^1 \frac{ds}{1 + ks} (T_c - \pi) + \log (1 + k)}$$

n(o) = k₀
$$\sqrt{\frac{1 - c/(1 + k_0^2)}{2(c - 1)}} e^{\frac{1}{\pi} \int_0^1 \frac{ds}{s}} (T_c - \pi) = \sqrt{\frac{1 - c/(1 + k_0^2)}{2}}.$$

We can derive an expression for h(x) suitable for numerical evaluation as follows:

$$h(x) = \frac{1}{2 i} \int_{-i\infty+\delta}^{i\infty+\delta} dk e^{kx} H(k), \quad 0 < \delta < 1$$

H(k) is not singular at $\pm ik_0$. The bracketed expression vanishes, thus the contour may be deformed to lie along the left cut. Only the integral

$$\frac{k}{\pi} \int_{0}^{1} \frac{ds}{1+ks} T_{c} = \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1-ks} T_{c} - \frac{2k^{2}}{\pi} \int_{0}^{1} \frac{s ds}{1-k^{2}s^{2}} T_{c}$$
$$= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1-ks} T_{c} - \log \left(\frac{k_{0}^{2}}{k^{2}+k_{0}^{2}} \frac{\frac{c}{2k} \log \frac{1+k}{1-k} - 1}{c-1} \right)$$

is double-valued across the cut. Thus only the first term in H(k) contributes.

$$h(\mathbf{x}) = \frac{1}{2\pi i} \int_{-\infty}^{1} d\mathbf{k} \ e^{\mathbf{k}\mathbf{x}} \frac{\mathbf{k}_{0}}{\mathbf{k}^{2} + \mathbf{k}_{0}^{2}} \sqrt{\frac{1 - c/(1 + \mathbf{k}_{0}^{2})}{2(c - 1)}} \ e^{\frac{\mathbf{k}}{\pi}} \int_{0}^{1} \frac{d\mathbf{s}}{1 - \mathbf{k}\mathbf{s}} \ ^{\mathrm{T}_{c}} \frac{(c - 1)(\mathbf{k}^{2} + \mathbf{k}_{0}^{2})}{\mathbf{k}_{0}^{2}} \left[\frac{1}{\frac{c}{2\mathbf{k}}} \left(\log \left| \frac{1 + \mathbf{k}}{1 - \mathbf{k}} \right| - \pi \mathbf{i} \right) - 1 - 1 \right] \\ - \frac{1}{\frac{c}{2\mathbf{k}}} \left(\log \left| \frac{1 + \mathbf{k}}{1 - \mathbf{k}} \right| + \pi \mathbf{i} \right) - 1 \right] \\ = \frac{c}{2\mathbf{k}_{0}} \sqrt{\frac{c - 1}{2}} \left(1 - \frac{c}{1 + \mathbf{k}_{0}^{2}} \right) - \frac{1}{\int_{-\infty}^{1}} \frac{d\mathbf{k}}{\mathbf{k}} \frac{\mathbf{k}}{\mathbf{k}} + \frac{\mathbf{k}}{\pi} \int_{0}^{1} \frac{d\mathbf{s}}{1 - \mathbf{k}\mathbf{s}} \ \mathrm{T}_{c} \\ = \frac{c}{2\mathbf{k}_{0}} \sqrt{\frac{c - 1}{2}} \left(1 - \frac{c}{1 + \mathbf{k}_{0}^{2}} \right) - \frac{1}{\int_{-\infty}^{1}} \frac{d\mathbf{k}}{\mathbf{k}} \frac{(c - 1)(\mathbf{k}^{2} + \mathbf{k}_{0}^{2})}{\mathbf{k}_{0}^{2} \left(1 - \frac{c}{2\mathbf{k}} \left(1 - \frac{c}{2\mathbf{k}} \right) - 1 \right)^{2} + \frac{\pi^{2}c^{2}}{4\mathbf{k}^{2}} \right]$$

Replacing k by -k gives

$$h(x) = -\frac{c}{2k_{O}}\sqrt{\frac{c-1}{2}\left(1-\frac{c}{1+k_{O}^{2}}\right)}\int_{1}^{\infty} \frac{k \, dk \, e}{\left(\frac{c}{2}\log\frac{k-1}{k+1}-k\right)^{2}+\left(\frac{\pi c}{2}\right)^{2}} e^{-kx}$$

(h(x) is negative for all x).

If c < 1 the roots of the characteristic equation are $\pm k_1$, where $c = k_1/\tanh^{-1}k_1$. The contours must now be taken as shown in Figure 5.





Proceeding in the same way as for c > 1 we get the analogous results:

$$n(x) = \sinh k_1(x + x_0) + h(x)$$

$$\frac{1}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{\mathrm{s}} (\mathrm{T}_{\mathrm{c}} - \pi) = \frac{1}{2} \log \frac{1 - \mathrm{c}}{\mathrm{k}_{1}^{2}}$$
(2.12)

$$\frac{k^2}{\pi} \int_{0}^{1} \frac{sds}{1-k^2s^2} T_c = -\frac{1}{2}\log\frac{k_0^2-k^2}{k_1^2} \cdot \frac{1-c}{1-\frac{c}{2k}\log\frac{1+k}{1-k}}$$
(2.13)

$$x_0 = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1-k_1^2s^2} T_c$$

$$T_c = \tan^{-1}\frac{\pi/2}{\tanh^{-1}s-1/cs} , \left[\tan^{-1}=\pi \to 0\right]$$

$$F(k) = \frac{k_1}{k^2-k_1^2} \sqrt{\frac{c/(1-k_1^2)-1}{2(1-c)}} e^{\frac{k}{F}} \int_{0}^{1} \frac{ds}{1+ks} T_c$$

$$H(0) = -\frac{1}{k_1} \left[\sqrt{\frac{c/(1-k_1^2)-1}{2(1-c)}} - \cosh k_1 x_0 \right]$$

$$\frac{-H'(0)}{H(0)} = -\frac{1}{H(0)k_1^2} \left[\sinh k_1 x_0 - k_1 \sqrt{\frac{c/(1-k_1^2)-1}{2(1-c)}} - \frac{1}{\pi} \int_{0}^{1} ds T_c \right]$$

$$n(0) = \sqrt{\frac{1}{2} \left(\frac{c}{1-k_1^2}-1\right)}$$

$$h(x) = -\frac{c}{2k_1} \sqrt{\frac{1-c}{2} \left(\frac{c}{1-k_1^2}-1\right)} \int_{1}^{\infty} \frac{kdk e^{-\frac{k}{\pi} \int_{0}^{1} \frac{ds}{1+ks} T_c}}{1 \left(\frac{c}{2}\log\frac{k+1}{k-1}-1\right)^2 + \left(\frac{\pi c}{2}\right)^2} e^{-kx}$$

Combining; these hyperbolic results (c < 1) with the elliptic results (c > 1) previously obtained shows the character of the solution and its numerically identifiable features to be continuous (as a function of c) across the parabolic (c = 1) boundary case.

We now treat the two-medium case, distinguishing the two materials (e.g., active material and tamper) only by their different values of c. Here four cases arise as the two c values are less than or greater than 1. We treat explicitly only the case: c > 1, c' < 1. The extension to other cases will then be obvious. Because of the applicability of the solution to the simple tamped sphere we refer to the one region, c > 1, x > 0, as "the core", and to the other, c < 1, x < 0, as "the tamper". We find two pertinent solutions, one belonging to a growing and the other to a decaying exponential asymptotic solution in the tamper. For the problem of the infinitely tamped sphere only the decaying solution will

figure (decaying as one moves away from the interface into the tamper). However, the "asymptotic solution" for a finite tamper will be a linear combination of the two solutions. The integral equation is:

$$n(x) = c' \int_{-\infty}^{0} dx' n(x') \frac{1}{2} E(|x - x'|) + c \int_{0}^{\infty} dx' n(x') \frac{1}{2} E(|x - x'|)$$
(2.14)

We use the same notation as before:

$$\begin{split} n(x) &= f(x) + g(x) \\ f(x) &= 0, \ x < 0 \\ g(x) &= 0, \ x \ge 0 \\ F(k) &= \int_{-\infty}^{\infty} dx \ f(x)e^{-kx} \\ G(k) &= \int_{-\infty}^{\infty} dx \ g(x)e^{-kx} \\ \hline \\ \frac{K}{2} (k) &= \int_{-\infty}^{\infty} dx \ \frac{1}{2} E(|x|) \ e^{-kx} &= \frac{1}{2k} \ \log \frac{1+k}{1-k} \\ F(k) + G(k) &= \int_{-\infty}^{\infty} dx \ n(x)e^{-kx} \\ &= \int_{-\infty}^{\infty} dx \ e^{-kx} \ \int_{-\infty}^{\infty} dx' \ \frac{1}{2} E(|x|) \ \int_{-\infty}^{\infty} dx' \ e^{-kx'} \ \left[c' \ g(x') + c \ f(x') \right] \\ &= \int_{-\infty}^{\infty} dy' \ e^{-ky} \ \frac{1}{2} E(|y|) \ \int_{-\infty}^{\infty} dx' \ e^{-kx'} \ \left[c' \ g(x') + c \ f(x') \right] \\ &= \frac{1}{2k} \ \log \frac{1+k}{1-k} \ \left[c' \ G(k) + c \ F(k) \right] \\ G(k) &= F(k) \ \frac{c}{2k} \ \log \frac{1+k}{1-k} \ -1 \\ 1 - \frac{c'}{2k} \ \log \frac{1+k}{1-k} \ \equiv F(k) \ P(k) \end{split}$$

The singularities of log P(k) now lie at:

$$\pm 1 \text{ (branch points)}$$

$$\pm ik_0 \left[\text{roots of } P(k), \frac{k_0}{\tan^{-1}k_0} = c \right]$$

$$\pm k_1 \left[\text{poles of } P(k) \frac{k_1}{\tanh^{-1}k_1} = c' \right]$$

F(k) and we assume also log F(k) must be analytic for R(k) > 0G(k) and we assume also log G(k) must be analytic for

 $R(K) < + k_1 \text{ for ''decaying solution'', i.e., } g(x) = 0(e^{k_1x})$ or R(k) < - k_1 for ''growing solution'', i.e., g(x) = 0(e^{-k_1x}) log P(k) is analytic for - 1 < R(k) < + 1, except at $\pm ik_0$, $\pm k_1$



For the two cases we choose contours as follows:





We treat first the decaying solution. As before we identify $\log F(k)$ and $\log G(k)$ with the left and right integrals (again excepting a constant).

$$\log P_{\mathbf{R}}(\mathbf{k}) = \frac{1}{2\pi i} \int_{\mathbf{R}} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \log P(\mathbf{k}') = \log G(\mathbf{k}) + \text{const.}$$
$$\log P_{\mathbf{L}}(\mathbf{k}) = -\frac{1}{2\pi i} \int_{\mathbf{L}} \frac{d\mathbf{k}}{\mathbf{k}' - \mathbf{k}} \log P(\mathbf{k}') = \log F(\mathbf{k}) + \text{const.}$$

We deform the contours as follows:



Figure 8.

$$\log \mathbf{P}_{\mathbf{R}}(\mathbf{k}) = \frac{1}{2\pi i} \int_{\mathbf{R}} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \left[\log \left(\frac{c}{2\mathbf{k}'} \log \frac{1 + \mathbf{k}'}{1 - \mathbf{k}'} - 1 \right) - \log \left(1 - \frac{c'}{2\mathbf{k}'} \log \frac{1 + \mathbf{k}'}{1 - \mathbf{k}'} \right) \right] \\ = \frac{1}{\pi} \int_{\mathbf{0}}^{1} \frac{d\mathbf{s}}{\mathbf{s}(1 - \mathbf{k}\mathbf{s})} \mathbf{T}_{\mathbf{C}} - \frac{1}{2\pi i} \int_{\mathbf{R}} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \log \left(1 - \frac{c'}{2\mathbf{k}'} \log \frac{1 + \mathbf{k}'}{1 - \mathbf{k}'} \right)$$
(2.15)

making use of the previous evaluation of the first term.

$$\log \mathbf{P}_{\mathbf{R}}(\mathbf{k}) = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1 - \mathbf{k}s)} \mathbf{T}_{\mathbf{C}} - \frac{1}{2\pi i} \int_{\mathbf{k}_{1}}^{\infty} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} (-2\pi i) - \frac{1}{2\pi i} \int_{\mathbf{R}'}^{\infty} \frac{d\mathbf{k}'}{\mathbf{k}' - \mathbf{k}} \log \left(\frac{c'}{2\mathbf{k}'} \log \frac{1 + \mathbf{k}'}{1 - \mathbf{k}'} - 1\right)$$

The last integral is now equivalent to that evaluated in equation 2.15 (and is identical with the rightcontour integral occurring in the one-medium problem for c < 1).

$$\log P_{R}(k) = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1-ks)} T_{c} + \int_{0}^{1/k_{1}} \frac{ds}{s(1-ks)} - \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1-ks)} T_{c'}$$
$$= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1-ks} (T_{c} - T_{c'}) + \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s} (T_{c} - T_{c'}) + \int_{0}^{1/k_{1}} ds \left(\frac{1}{s} + \frac{k}{1-ks}\right)$$

We choose the constant to make

$$\log G(k) = \log P_{R}(k) + \log B - \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s} (T_{c} - T_{c'}) - \int_{0}^{1/k_{1}} \frac{ds}{s}$$

$$= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 - ks} (T_{c} - T_{c'}) + \log \frac{Bk_{1}}{k_{1} - k}$$
(2.16)

Evaluating the left-contour integral gives

$$-\log P_{L}(k) = \frac{1}{2\pi i} \int_{L} \frac{dk'}{k' - k} \left[\log\left(\frac{c}{2k'} \log \frac{1 + k'}{1 - k'} - 1\right) - \log\left(1 - \frac{c'}{2k'} \log \frac{1 + k'}{1 - k'}\right) \right]$$
$$= \left\{ -2 \int_{0}^{1} \frac{ds}{s} + \log\left(k^{2} + k_{0}^{2}\right) + \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1 + ks)} T_{c} \right\}$$
$$- \frac{1}{2\pi i} \int_{L'} \frac{dk'}{k' - k} \log\left(1 - \frac{c'}{2k'} \log \frac{1 + k'}{1 - k'}\right)$$

Figure 10.

$$= \left\{ \dots \right\} - \frac{1}{2\pi i} \int_{-\infty}^{-k_1} \frac{dk'}{k' - k} (2\pi i) - \frac{1}{2\pi i} \int_{L''} \frac{dk'}{k' - k} \log \left(\frac{c'}{2k'} \log \frac{1 + k'}{1 - k'} - 1 \right)$$







$$\frac{1}{2\pi i} \int_{L''} \frac{dk}{k'-k} \log\left(\frac{c'}{2k'}\log\frac{1+k'}{1-k'}-1\right) = \frac{1}{2\pi i} \int_{R'} \frac{dk''}{k''+k} \log\left(\frac{c}{2k''}\log\frac{1+k''}{1-k''}-1\right),$$

$$= \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1+ks)} T_{c'}.$$

$$= \log P_{L}(k) = -2 \int_{0}^{1} \frac{ds}{s} + \log\left(k^{2}+k_{0}^{2}\right) + \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1+ks)} T_{c} + \int_{0}^{1/k_{2}} ds\left(\frac{1}{s}-\frac{k}{1+ks}\right) - \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1+ks)} T_{c'}.$$

$$= -\frac{k}{\pi} \int_{0}^{1} \frac{ds}{1+ks} \left(T_{c}-T_{c'}\right) + \log\frac{k_{1}\left(k^{2}+k_{0}^{2}\right)}{k_{1}+k} + \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s} \left(T_{c}-T_{c'}\right) - 2 \int_{0}^{1} \frac{ds}{s} + \int_{0}^{1/k_{1}} \frac{ds}{s}$$

$$\log F(k) = \log P_{L}(k) + \log B - \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s} \left(T_{c}-T_{c'}\right) - \frac{1/k_{1}}{s} \frac{ds}{s}$$

$$= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1+ks} \left(T_{c}-T_{c'}\right) + \log\frac{(k_{1}+k)B}{k_{1}(k^{2}+k_{0}^{2})} - \frac{2}{\pi} \int_{0}^{1} \frac{ds}{s} \left(T_{c}-T_{c'}\right) + 2 \int_{1/k_{1}}^{1} \frac{ds}{s}$$

$$-\frac{2}{\pi} \int_{0}^{1} \frac{ds}{s} \left[\left(\pi-T_{c'}\right) - \left(\pi-T_{c}\right)\right] = -\log\frac{k_{1}^{2}}{1-c'} + \log\frac{k_{0}^{2}}{c-1}$$

$$\log F(k) = \frac{k}{\pi} \int_{0}^{0} \frac{ds}{1 + ks} (T_{c} - T_{c'}) + \log \frac{(k_{1} + k)B}{k_{1}(k^{2} + k_{0}^{2})} + \log \left(\frac{1 - c'}{k_{1}^{2}} \cdot \frac{k_{0}^{2}}{c - 1}\right) + \log k_{1}^{2}$$
$$= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 + ks} (T_{c} - T_{c'}) + \log \frac{Bk_{0}^{2}(k_{1} + k)(1 - c')}{k_{1}(k^{2} + k_{0}^{2})(c - 1)} \cdot$$

We again determine x_0 and the value of B required to make the asymptotic sine solution of unit amplitude.

$$f(x) = \sin k_0 (x + x_0) + h(x), x > 0, h(x) \to 0 \text{ as } x \to +\infty$$
(2.17)
$$F(k) = \frac{1}{2i} \left(\frac{e^{ik_0 x_0}}{k - ik_0} - \frac{e^{-ik_0 x_0}}{k + ik_0} \right) + H(k)$$

 $\lim_{\epsilon \to 0} \left[\log \mathbf{F}(i\mathbf{k}_0 + \epsilon) - \log \mathbf{F}^{-}i\mathbf{k}_0 + \epsilon \right] = \log (-1) + 2i\mathbf{k}_0 \mathbf{x}_0$

$$= \frac{2ik_{0}}{\pi} \int_{0}^{1} \frac{ds}{1+k_{1}^{2}s^{2}} (T_{c} - T_{c'}) + \log\left(\frac{-2ik_{0}\epsilon}{+2ik_{0}\epsilon}\right) + \log\frac{k_{1} + ik_{0}}{k_{1} - ik_{0}}$$
$$x_{0} = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1+k_{0}^{2}s^{2}} (T_{c} - T_{c'}) + \frac{1}{k_{0}} \tan^{-1}\frac{k_{0}}{k_{1}} = x_{1} + \frac{1}{k_{0}} \tan^{-1}\frac{k_{0}}{k_{1}}$$
(2.18)

$$\lim_{\epsilon \to 0} \left[\log F(ik_0 + \epsilon) + \log F(-ik_0 + \epsilon) - 2 \log \epsilon \right] = -2 \log 2$$
$$= \frac{2k_0^2}{\pi} \int_0^1 \frac{s \, ds}{1 + k_0^2 s^2} \left(T_c - T_{c'} \right) + 2 \log \frac{Bk_0^2(1 - c')}{k_1(c - 1)} + \log \frac{k_1^2 + k_0^2}{4k_0^2} \right)$$

The first term may be evaluated by the use of equation 2.11 and equation 2.13.

$$\frac{2k_0^2}{\pi} \int_0^1 \frac{sds}{1+k_0^2 s^2} (T_c - T_{c'}) = \lim_{\epsilon \to 0} \left[\log \left\{ \frac{2ik_0 \epsilon(c-1)}{k_0^2 \frac{i}{k_0} \left(1 - \frac{c}{1+k_0^2}\right) \epsilon} \right\} \right] - \log \frac{(k_1^2 + k_0^2)(1 - c')}{k_1^2 \left(1 - \frac{c'}{k_0} \tan^{-1} k_0\right)} = \log \frac{2(c-1)k_1^2(1 - c'/c)}{\left(1 - \frac{c}{1+k_0^2}\right) (k_1^2 + k_0^2)(1 - c')}$$

$$(2.19)$$

 $\log \mathbf{B} = \log \frac{k_1 (c - 2)}{k_0^2 (1 - c')} - \frac{1}{2} \log \frac{k_1^2 + k_0^2}{k_0^2} \frac{1}{2} \log \frac{2(c - 1) k_1^2 (1 - c'/c)}{\left[1 - c/(1 + k_0^2)\right] (k_1^2 + k_0^2)(1 - c')}$

$$= \frac{1}{2} \log \frac{(c-1) \left[1 - c/(1+k_0^2) \right]}{2k_0^2 (1-c') (1-c'/c)}$$

$$\log F(h) = \frac{k}{\pi} \int_0^1 \frac{ds}{1+ks} \left(T_c - T_{c'} \right) + \frac{1}{2} \log \frac{k_0^2 (1-c') \left[1 - c/(1+k_0^2) \right]}{2k_1^2 (c-1) (1-c'/c)} + \log \left(\frac{k+k_1}{k^2+k_0^2} \right)$$

$$\begin{split} F(k) &= \frac{k_0}{k_1} \frac{k + k_1}{k^2 + k_0^2} \sqrt{\frac{(1 - c')\left[1 - c/(1 + k_0^2)\right]}{2(c - 1)(1 - c'/c)}} e^{\frac{k}{\pi}} \int_{0}^{1} \frac{ds}{1 + ks} (T_c - T_{c'}) \\ H(k) &= F(k) - \frac{k \sin k_0 x_0 + k_0 \cos k_0 x_0}{k^2 + k_0^2} \\ &= \frac{1}{k^2 + k_0^2} \left[\frac{k_0}{k_1} (k + k_1) \sqrt{\frac{(1 - c')\left[1 - c/(1 + k_0^2)\right]}{2(c - 1)(1 - c'/c)}} e^{\frac{k}{\pi}} \int_{0}^{1} \frac{ds}{1 + ks} (T_c - T_{c'}) \\ - k \sin k_0 x_0 - k_0 \cos k_0 x_0 \right] \\ H(0) &= \frac{1}{k_0} \left[\frac{(1 - c')\left[1 - c/(1 + k_0^2)\right]}{2(c - 1)(1 - c'/c)} - \cos k_0 x_0 \right] \\ H'(0) &= \frac{(1 - c')\left[1 - c/(2 + k_0^2)\right]}{2(c - 1)(1 - c'/c)} \left(\frac{1}{k_0 k_1} + \frac{1}{k_0 \pi} \int_{0}^{1} ds (T_c - T_c \cdot) \right) - \frac{1}{k_0^2} \sin k_0 x_0 \\ - \frac{H'(0)}{H(0)} &= -\frac{1}{H(0) k_0} \left[\sqrt{\frac{(1 - c')\left[1 - c/(1 + k_0^2)\right]}{2(c - 1)(1 - c'/c)}} \left(\frac{1}{k_1} + \frac{1}{\pi} \int_{0}^{1} ds (T_c - T_c \cdot) \right) - \frac{1}{k_0} \sin k_0 x_0 \right] \\ n(0) &= \lim_{k \to \infty} kF(k) = \lim_{k \to \infty} \frac{k}{k^2 + k_0^2} \frac{k_0}{k_1} (k + k_1) \sqrt{\frac{(1 - c')\left[1 - c/(1 + k_0^2)\right]}{2(c - 1)(1 - c'/c)}}} e^{\frac{\pi}{\pi}} \int_{0}^{1} \frac{ds}{s} (T_c' - \pi) \\ &= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 + ks} (T_c - \pi + \pi - T_{c'}) \\ &\to e^{\frac{1}{\pi}} \int_{0}^{1} \frac{ds}{s} (T_c - \pi) - \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s} (T_c' - \pi) \\ &= \sqrt{\frac{k_0^2(1 - c')}{k_0^2(1 - c')}} \end{array}$$

(using (2.10) and (2.12)).

$$n(0) = \sqrt{\frac{1 - c/(1 + k_0^2)}{2(1 - c'/c)}}$$
(2.20)
$$h(x) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} dk \ H(k) \ e^{kx}$$
$$= \frac{1}{2\pi i} \int_{-L''} \frac{dk \ e^{kx}(k + k_1)}{k^2 + k_0^2} \ Ce^{\frac{k}{\pi}} \int_{0}^{1} \frac{ds}{1 + ks} (T_c - T_{c'})$$





$$\begin{aligned} \text{where } \mathbf{C} &= \frac{\mathbf{k}_{0}}{\mathbf{k}_{1}} \sqrt{\frac{\left[1 - \mathbf{c}/(1 + \mathbf{k}_{0}^{2})\right](1 - \mathbf{c}')}{2(\mathbf{c} - 1)(1 - \mathbf{c}'/\mathbf{c})}} \\ &= \frac{\mathbf{k}}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{1 + \mathbf{k}_{S}} \left(\mathbf{T}_{C} - \mathbf{T}_{C'}\right) = \mathbf{e}^{\frac{\mathbf{k}}{\pi}} \int_{0}^{1} \frac{\mathrm{ds}}{1 - \mathbf{k}_{S}} \left(\mathbf{T}_{C} - \mathbf{T}_{C'}\right) \frac{(\mathbf{k}^{2} + \mathbf{k}_{0}^{2})(\mathbf{c} - 1)}{\mathbf{k}_{0}^{2} \left(\frac{\mathbf{c}}{2\mathbf{k}} \log \frac{1 + \mathbf{k}}{1 - \mathbf{k}} - 1\right)} \cdot \frac{\mathbf{k}_{1}^{2} \left(1 - \frac{\mathbf{c}'}{2\mathbf{k}} \log \frac{1 + \mathbf{k}}{1 - \mathbf{k}}\right)}{(\mathbf{k}_{1}^{2} - \mathbf{k}^{2})(1 - \mathbf{c}')} \\ &= \frac{1}{2\pi \mathbf{i}} \int_{-\infty}^{-1} d\mathbf{k} \ \mathbf{e}^{\mathbf{k}\mathbf{x}} \ \mathbf{C} \ \frac{\mathbf{k}_{1}^{2}(\mathbf{c} - 1)}{\mathbf{k}_{0}^{2}(\mathbf{k}_{1} - \mathbf{k})(1 - \mathbf{c}')} \ \mathbf{e}^{\frac{\mathbf{k}}{\pi}} \int_{0}^{1} \frac{d\mathbf{s}}{1 - \mathbf{k}_{S}} \left(\mathbf{T}_{C} - \mathbf{T}_{C'}\right) \left\{ \frac{1 - \frac{\mathbf{c}'}{2\mathbf{k}} \left(\log\left|\frac{1 + \mathbf{k}}{1 - \mathbf{k}}\right| - \pi\mathbf{i}\right)}{\frac{\mathbf{c}}{2\mathbf{k}} \left(\log\left|\frac{1 + \mathbf{k}}{1 - \mathbf{k}}\right| - \pi\mathbf{i}\right) - 1} \\ &- \frac{1 - \frac{\mathbf{c}'}{2\mathbf{k}} \left(\log\left|\frac{1 + \mathbf{k}}{1 - \mathbf{k}}\right| + \pi\mathbf{i}\right) - 1}{\frac{\mathbf{c}}{2\mathbf{k}} \left(\log\left|\frac{1 + \mathbf{k}}{1 - \mathbf{k}}\right| + \pi\mathbf{i}\right) - 1} \\ \end{aligned} \right\}.$$

Replacing k by -k gives

$$h(x) = \frac{1}{2\pi i} \int_{1}^{\infty} dk \ e^{-kx} \ \frac{k_1}{k_0(k_1 + k)} \sqrt[4]{\frac{\left[1 - c/(1 + k_0^2)\right](c - 1)}{2(1 - c')(1 - c'/c)}} \ e^{-\frac{k}{\pi}} \int_{0}^{1} \frac{ds}{1 + ks} \ (T_c - T_{c'}) \left[\dots\right]$$

where

$$\left\{ \cdots \right\} = -\frac{2\pi i}{2k} \frac{c\left(1 - \frac{c'}{2k}\log\frac{k+1}{k-1}\right) + c'\left(\frac{c}{2k}\log\frac{k+1}{k-1} - 1\right)}{\left(\frac{c}{2k}\log\frac{k+1}{k-1} - 1\right)^2 + \left(\frac{c\pi}{2k}\right)^2} = -\frac{\pi i}{k} \frac{c-c'}{\left(\frac{c}{2k}\log\frac{k+1}{k-1} - 1\right)^2 + \left(\frac{c\pi}{2k}\right)^2}$$
$$h(x) = -\frac{k_1^2}{2k_0} \sqrt{\frac{\left[1 - c/(1+k_0^2)\right](c-1)(1-c'/c)}{2(1-c')}} \int_{1}^{\infty} \frac{k \, dk}{k+k_1} \frac{e^{-\frac{k}{\pi} \int_{0}^{1} \frac{ds}{1+k_s}(T_c - T_{c'})}}{\left(\frac{c}{2}\log\frac{h+1}{k-1} - k\right)^2 + \left(\frac{c\pi}{2}\right)^2} e^{-kx}$$

Now returning to G(k) $\log G(k) = \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 - ks} (T_{c} - T_{c'}) + \log \frac{Bk_{1}}{k_{1} - k}$ (2.16) $= \frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 - ks} (T_{c} - T_{c'}) + \log \frac{k_{1}}{k_{1} - k} + \frac{1}{2} \log \frac{(c - 1) \left[1 - c/(1 + k_{0}^{2})\right]}{2k_{0}^{2}(1 - c')(1 - c'/c)}$

A check of this expression is afforded by evaluating

$$g(-c) = \lim_{k \to -\infty} -k \ G(k) = \sqrt{\frac{1 - c/(1 + k_0^2)}{2(1 - c'/c)}} = n(0), \qquad (ef. \ equation \ 2.20).$$

$$G(k) = \int_{-\infty}^{0} dx \ e^{-kx} \ g(x) = \int_{-\infty}^{0} dx \ e^{-kx} \left[Ae^{-k_1x} + j(x) \right], \qquad (ef. \ equation \ 2.20).$$

$$G(k) = \int_{-\infty}^{0} dx \ e^{-kx} \ g(x) = \int_{-\infty}^{0} dx \ e^{-kx} \left[Ae^{-k_1x} + j(x) \right], \qquad (ef. \ equation \ 2.20).$$

$$G(k) = \frac{A}{k_1 - k} + J(k), \ J(k_1) \ as \ x \to -\infty$$

$$G(k) = \frac{A}{k_1 - k} + J(k), \ J(k_1) \ is \ finite.$$

$$\log G(k_1 + \epsilon) = \log \left(\frac{-A}{\epsilon}\right) + 0(\epsilon)$$

$$= \log \left(\frac{-k_1}{s}\right) + \frac{k_1}{\pi} \int_{0}^{1} \frac{ds}{1 - k_1 s} (T_c - T_c') + \frac{1}{2} \log \frac{(c - 1)[1 - c/(1 + k_0^2)]}{2k_0^2(1 - c')(1 - c'/c)}$$

$$A = \frac{k_1}{k_0} \sqrt{\frac{(c - 1)[1 - c/(1 + k_0^2)]}{2(1 - c')(1 - c'/c)}} \ e^{\frac{k_1}{\pi}} \int_{0}^{1} \frac{ds}{1 - k_1 s} (T_c - T_{c'}).$$

$$\frac{k_1}{\pi} \int_{0}^{1} \frac{ds}{1 - k_1 s} (T_c - T_{c'}) = \frac{k_1}{\pi} \int_{0}^{1} \frac{ds}{1 - k_1^2 s^2} (T_c - T_{c'}) + \frac{k_1^2}{\pi} \frac{s \ ds}{1 - k_1^2 s^2} (T_c - T_{c'}).$$

The first term will be called k_1x_2 by analogy with the x_1 introduced in equation 2.18, the second can be evaluated by the use of equation 2.11 and 2.13.

$$e^{\frac{k_{1}}{\pi}\int_{0}^{1}\frac{ds}{1-k_{1}s}(T_{c}-T_{c'})} = e^{\frac{k_{1}x_{2}}{\sqrt{\frac{2k_{0}2(c/c'-1)(1-c')}{(k_{1}^{2}+k_{0}^{2})(c-1)[c'/(1-k_{1}^{2})-1]}}} (2.21)$$

so that

=

$$A = \frac{k_{1}}{\sqrt{k_{1}^{2} + k_{0}^{2}}} \frac{c \left[1 - c/(1 + k_{0}^{2})\right]}{c' \left[c'/(1 - k_{1}^{2}) - 1\right]}} e^{k_{1}x_{2}}$$

$$g(x) = \frac{k_{1}\sqrt{c \left[1 - c/(1 + k_{0}^{2})\right]}}{\sqrt{k_{1}^{2} + k_{0}^{2}} \sqrt{c' \left[c'/(1 - k_{1}^{2}) - 1\right]}} e^{k_{1}(x + x_{1})} + j(x) \qquad (2.22)$$

$$J(k) = G(k) - \frac{A}{k_{1} - k}$$

$$\frac{k_{1}}{k_{0}(k_{1} - k)} \sqrt{\frac{(c - 1)\left[1 - c/(1 + k_{0}^{2})\right]}{2(1 - c')(1 - c'/c)}} \left[e^{\frac{k}{\pi} \int_{0}^{1} \frac{ds}{1 - ks}} (T_{c} - T_{c'}) - e^{\frac{k_{1}}{\pi} \int_{0}^{1} \frac{ds}{1 - k_{1}s}} (T_{c} - T_{c'}) \right]$$

$$\begin{split} \mathbf{j}(\mathbf{x}) &= \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} d\mathbf{k} \ e^{\mathbf{k}\mathbf{x}} \ \mathbf{J}(\mathbf{k}), \\ &= \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} d\mathbf{k} \ \frac{e^{\mathbf{k}\mathbf{x}} \ \mathbf{k}_1}{\mathbf{k}_0(\mathbf{k}_1 - \mathbf{k})} \sqrt{\frac{(\mathbf{c} - 1)\left[1 - \mathbf{c}/(1 + \mathbf{k}_0 2)\right]}{2(1 - \mathbf{c}')(1 - \mathbf{c}'/\mathbf{c})}} \left[e^{\mathbf{k} \frac{\pi}{\pi}} \int_0^1 \frac{d\mathbf{s}}{1 - \mathbf{k}\mathbf{s}} (\mathbf{T}_{\mathbf{c}} - \mathbf{T}_{\mathbf{c}'}) - e^{\mathbf{k}_1} \frac{1}{\pi}} \int_0^1 \frac{d\mathbf{s}}{1 - \mathbf{k}\mathbf{s}\mathbf{s}} (\mathbf{T}_{\mathbf{c}} - \mathbf{T}_{\mathbf{c}'})} \right] \\ &= \frac{1}{2\pi i} \frac{\mathbf{k}_1}{\mathbf{k}_0} \sqrt{\frac{(\mathbf{c} - 1)\left[1 - \mathbf{c}/(1 + \mathbf{k}_0 2)\right]}{2(1 - \mathbf{c}')(1 - \mathbf{c}'/\mathbf{c})}} \int_{\mathbf{k}'}^\infty \frac{d\mathbf{k}}{\mathbf{k}_1 - \mathbf{k}} e^{\mathbf{k}\mathbf{x}} e^{\mathbf{k} \frac{\pi}{\pi}} \int_0^1 \frac{d\mathbf{s}}{1 + \mathbf{k}\mathbf{s}} (\mathbf{T}_{\mathbf{c}} - \mathbf{T}_{\mathbf{c}'}) \\ &= \frac{1}{2\pi i} \frac{\mathbf{k}_0}{\mathbf{k}_0} \sqrt{\frac{(1 - \mathbf{c}')\left[1 - \mathbf{c}/(1 + \mathbf{k}_0 2)\right]}{2(\mathbf{c} - 1)(1 - \mathbf{c}'/\mathbf{c})}} \int_{\mathbf{k}'}^\infty \frac{d\mathbf{k}}{\mathbf{k}_1 - \mathbf{k}} e^{\mathbf{k}\mathbf{x}} e^{\mathbf{k} \frac{\pi}{\pi}} \int_0^1 \frac{d\mathbf{s}}{1 + \mathbf{k}\mathbf{s}} (\mathbf{T}_{\mathbf{c}} - \mathbf{T}_{\mathbf{c}'}) \\ &= \frac{1}{2\pi i} \frac{\mathbf{k}_0}{\mathbf{k}_1} \sqrt{\frac{(1 - \mathbf{c}')\left[1 - \mathbf{c}/(1 + \mathbf{k}_0 2)\right]}{2(\mathbf{c} - 1)(1 - \mathbf{c}'/\mathbf{c})}}} \int_{\mathbf{k}'}^\infty \frac{d\mathbf{k}(\mathbf{k} + \mathbf{k}_1) e^{\mathbf{k}\mathbf{x}}}{\mathbf{k}^2 + \mathbf{k}_0^2} \frac{\pi}{\mathbf{k}^2 + \mathbf{k}_0^2} \left[\frac{\mathbf{c}}{2\mathbf{k}} \left(\log \frac{\mathbf{k} + 1}{\mathbf{k} - 1} + \pi \mathbf{i} \right) - 1}{1 - \frac{\mathbf{c}'}{2\mathbf{k}} \left(\log \frac{\mathbf{k} + 1}{\mathbf{k} - 1} + \pi \mathbf{i} \right)}{\mathbf{k}^2 + \mathbf{k}_0^2}} \right] \\ \mathbf{k}(\mathbf{x}) = \frac{\mathbf{k}_0}{2\mathbf{k}_1} \sqrt{\frac{(1 - \mathbf{c}')\left[1 - \mathbf{c}/(1 + \mathbf{k}_0^2)\right](1 - \mathbf{c}'/\mathbf{c})}{2(\mathbf{c} - 1)}} \int_{\mathbf{k}'}^\infty \frac{\mathbf{k}}{\mathbf{k}} \frac{d\mathbf{k}}{\mathbf{k}} \left(\mathbf{k} + \mathbf{k}_1\right)}{\mathbf{k}^2 + \mathbf{k}_0^2} \\ \frac{\mathbf{k}}{\mathbf{k}} \frac{1}{\mathbf{k}} \frac{d\mathbf{k}}{\mathbf{k}} \left(\mathbf{k} - \frac{\mathbf{k}}{\mathbf{k}} - \frac{\mathbf{k}$$

The second solution differs in having as an asymptotic solution in the tamper a growing exponential (growing for increasing negative x), e^{-k_1x} . The core solution is again sinusoidal, differing only in phase from the first solution. Thus, the left contour must still lie to the right of the roots of P(k) at \pm ik_0 . The tamper solution, g(x), is to grow as e^{-k_1x} . Thus G(k) must have a pole at $-k_1$. (It may also have a pole at $+k_1$, the corresponding asymptotic g(x), e^{k_1x} , will be dominated by the growing exponential.) To give G(k) a pole at $-k_1$ the right contour must pass to the left of the pole of P(k) at $-k_1$. Since the left-contour must always be to the left of the right contour, the two contours must be taken as in Figure 7. (Other contour arrangements are possible, e.g., but the solutions so obtained may be represented as linear combinations of the two solutions obtained from the contours of Figure 6 and 7.





Figure 13.

Deforming the contours of Figure 7 so as to permit simplification of the integrals gives this form:



Figure 14.

Taking as before:

$$\log P_{L}(k) = -\frac{1}{2\pi i} \int_{L} \frac{dk'}{k' - k} \log P(k') = \log F(k) + \text{constant}$$
$$\log P_{R}(k) = \frac{1}{2\pi i} \int_{R} \frac{dk'}{k' - k} \log P(k') = \log G(k) + \text{constant}$$

The integral, log $P_{R}(k)$, may be broken up into pieces which have been evaluated previously.

$$\log P_{R}(k) = \frac{1}{2\pi i} \int_{R} \frac{dk'}{k'-k} \log \left(\frac{c}{2k'} \log \frac{1+k'}{1-k'} - 1\right) - \frac{1}{2\pi i} \int_{R} \frac{dk'}{k'-k} \log \left(1 - \frac{c'}{2k'} \log \frac{1+k'}{1-k'}\right)$$
$$= \frac{1}{\pi} \int_{0}^{1} \frac{ds}{s(1-ks)} T_{c} - \frac{1}{2\pi i} \int_{-k_{1}}^{\infty} \frac{dk'}{k'-k} (-2\pi i)$$
$$- \frac{1}{2\pi i} \int_{R} \frac{dk'}{k'-k} \log \left(1 - \frac{c'}{2k'} \log \frac{1+k'}{1-k'}\right)$$

The last term has been evaluated in getting log $P_{\mathbf{R}}(\mathbf{k})$ for the decaying solution.

$$\log P_{\rm R}(k) = \frac{1}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{\mathrm{s}(1-\mathrm{ks})} T_{\rm c} + \int_{0}^{-1/k_1} \frac{\mathrm{ds}}{\mathrm{s}(1-\mathrm{ks})} + \int_{0}^{1/k_1} \frac{\mathrm{ds}}{\mathrm{s}(1-\mathrm{ks})} - \frac{1}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{\mathrm{s}(1-\mathrm{ks})} T_{\rm c}'$$
$$= \frac{k}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{1-\mathrm{ks}} (T_{\rm c} - T_{\rm c}') + \frac{1}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{\mathrm{s}} (T_{\rm c} - T_{\rm c}') + 2 \int_{0}^{1} \frac{\mathrm{ds}}{\mathrm{s}} - \log (\mathrm{k}^2 - \mathrm{k}_1^2).$$
$$\log G(k) = \frac{k}{\pi} \int_{0}^{1} \frac{\mathrm{ds}}{1-\mathrm{ks}} (T_{\rm c} - T_{\rm c}') - \log (\mathrm{k}_1^2 - \mathrm{k}^2) + \log B'$$
(2.24)

It may be observed that the G(k) here obtained differs by a factor of $\frac{B'}{k_1(k+k_1)B}$ from the G(k) previously obtained. Since the ratio of F(k) to G(k) is the same, the two F(k)'s must differ by the same factor. We may therefore write log F(k) immediately

$$\log F(k) = \frac{k}{\pi} \int_0^1 \frac{ds}{1 + ks} (T_c - T_{c'}) + \log \frac{B' k_0^2 (1 - c')}{k_1^2 (k^2 + k_0^2)(c - 1)}$$

B' is again to be evaluated to give the asymptotic sine solution unit amplitude.

$$f(x) = \sin k_{0}(x + x_{1}) + h(x), x > 0, h(x) \to 0 \text{ as } x \to \infty$$
(2.25)
$$F(k) = \frac{1}{2i} \left(\frac{e^{ik_{0}x_{1}}}{k - ik_{0}} - \frac{e^{-ik_{0}x_{1}}}{k + ik_{0}} \right) + H(k).$$

 $\lim_{\epsilon \to 0} \left[\log F(ik_0 + \epsilon) - \log F(-ik_0 + \epsilon) \right] = \log (-1) + 2ik_0 x_1,$

$$= \frac{2ik_0}{\pi} \int_0^1 \frac{ds}{1 + k_0^2 s^2} (T_c - T_{c'}) + \log(-1)$$

$$x_{1} = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1 + k_{0}^{2}s^{2}} (T_{c} - T_{c'}) \quad (x_{1} < 0 \text{ since } T_{c} < T_{c'} \text{ for } 0 < s < 1)$$
(2.26)

 $\lim_{\epsilon \to 0} \left[\log F(ik_0 + \epsilon) + \log F(-ik_0 + \epsilon) + 2 \log \epsilon \right] = -2 \log 2$

$$= \frac{2k_0^2}{\pi} \int_0^1 \frac{s \, ds}{1 + k_0^2 s^2} \left(T_c - T_{c'} \right) + 2 \log \frac{B' k_0^2 (1 - c')}{k_1^2 (c - 1)} - 2 \log (2k_0) \right)$$

$$\log B' = \log \frac{k_1^2 (c - 1)}{k_0 (1 - c')} - \frac{k_0^2}{\pi} \int_0^1 \frac{s \, ds}{1 + k_0^2 s^2} \left(T_c - T_{c'} \right)$$

$$= \log \frac{k_1^2 (c - 1)}{k_0 (1 - c')} - \frac{1}{2} \log \frac{2(c - 1)k_1^2 (1 - c'/c)}{[1 - c/(1 + k_0^2)] (k_1^2 + k_0^2)(1 - c')}$$
(cf. 2.19)

4

$$\begin{split} &= \frac{1}{2} \log \frac{k_1 2 (c - 1) \left[1 - c / (1 + k_0^2) \right] (k_1^2 + k_0^2)}{2k_0^2 (1 - c') (1 - c'/c)} \\ &\log F(k) = \frac{k}{\pi} \int_0^1 \frac{ds}{1 + ks} \left(T_c - T_{c'} \right) + \log B' + \log \frac{k_0^2 (1 - c')}{k_1^2 (k^2 + k_0^2) (c - 1)} \\ &= \frac{k}{\pi} \int_0^1 \frac{ds}{1 + ks} \left(T_c - T_{c'} \right) + \frac{1}{2} \log \frac{k_0^2 (1 - c') \left[1 - c / (1 + k_0^2) \right] (k_1^2 + k_0^2)}{2k_1^2 (c - 1) (k^2 + k_0^2)^2 (1 - c'/c)} \\ &F(k) = \frac{k_0 \sqrt{k_1^2 + k_0^2}}{k_1 (k^2 + k_0^2)} \sqrt{\frac{(1 - c') \left[1 - c / (1 + k_0^2) \right]}{2(c - 1) (1 - c'/c)}} e^{\frac{k}{\pi}} \int_0^1 \frac{ds}{1 + ks} \left(T_c - T_{c'} \right) \\ &H(k) = \frac{k_0 \sqrt{k_1^2 + k_0^2}}{k_1 (k^2 + k_0^2)} \sqrt{\frac{(1 - c') \left[1 - c / (1 + k_0^2) \right]}{2(c - 1) (1 - c'/c)}} e^{\frac{k}{\pi}} \int_0^1 \frac{ds}{1 + ks} \left(T_c - T_{c'} \right) \\ \end{split}$$

$$h(x) = \frac{1}{2\pi i} \int_{-i\infty+\delta}^{i\infty+\delta} dk \ H(k) \ e^{kx} = \frac{1}{2\pi i} \int_{-L''}^{\infty} dk \ F(k) \ e^{kx}, \quad (cf. \ Fig \ 12),$$

 $-\frac{k \sin k_0 x_1 + k_0 \cos k_0 x_1}{k^2 + k_0^2}$

since H(k) is regular at \pm ik₀ and F(k) - F(k) - H(k) is single-valued across the $-\infty \longrightarrow -1$ cut.

$$h(\mathbf{x}) = \frac{1}{2\pi i} \int_{-\infty}^{-1} d\mathbf{k} \ e^{\mathbf{k}\mathbf{x}} \ \frac{D(\mathbf{c} - 1)\mathbf{k}\mathbf{1}^2 \ e}{\mathbf{k}_0^2(\mathbf{k}\mathbf{1}^2 - \mathbf{k}^2)(1 - \mathbf{c}^{\,\prime})} \left\{ \frac{1 - \frac{\mathbf{c}^{\,\prime}}{2\mathbf{k}} \left(\log \left| \frac{1 + \mathbf{k}}{1 - \mathbf{k}} \right| - \pi \mathbf{i} \right)}{\frac{\mathbf{c}}{2\mathbf{k}} \left(\log \left| \frac{1 + \mathbf{k}}{1 - \mathbf{k}} \right| - \pi \mathbf{i} \right) - 1} \right. \\ \left. \frac{1 - \frac{\mathbf{c}^{\,\prime}}{2\mathbf{k}} \left(\log \left| \frac{1 + \mathbf{k}}{1 - \mathbf{k}} \right| - \pi \mathbf{i} \right) - 1}{\frac{\mathbf{c}^{\,\prime}}{2\mathbf{k}} \left(\log \left| \frac{1 + \mathbf{k}}{1 - \mathbf{k}} \right| + \pi \mathbf{i} \right) - 1} \right]$$

where
$$- \frac{\mathbf{k}_0 \sqrt{\mathbf{k}\mathbf{1}^2 + \mathbf{k}\mathbf{1}^2} \sqrt{(1 - \mathbf{c}^{\,\prime}) \left[1 - \mathbf{c}^{\,\prime}/(1 + \mathbf{k}_0^2) \right]}}$$

$$D = \frac{k_0 \sqrt{k_1^2 + k_1^2}}{k_1} \sqrt{\frac{(1 - c') \left[1 - c/(1 + k_0^2)\right]}{2(c - 1) (1 - c'/c)}}$$

$$h(x) = \frac{k_{1}c\sqrt{k_{1}^{2} + k_{0}^{2}}}{2k_{0}}\sqrt{\frac{\left[1 - c/(1 + k_{0}^{2})\right](c - 1)(1 - c'/c)}{2(1 - c')}} \int_{1}^{\infty} \frac{kdk}{k^{2} - k_{1}^{2}} \frac{e^{-\frac{k}{\pi}\int_{0}^{1} \frac{ds}{1 + ks}(T_{c} - T_{c'})}}{\left(\frac{c}{2}\log\frac{k + 1}{k - 1} - k\right)^{2} + \left(\frac{\pi c}{2}\right)^{2}} e^{-kx}$$

$$\log G(k) = \frac{k}{\pi}\int_{0}^{1} \frac{ds}{1 - ks}(T_{c} - T_{c'}) - \log (k_{1}^{2} - k^{2}) + \log B'.$$

$$G(k) = \frac{k_1 \sqrt{k_1 2 + k_0 2}}{k_0 (k_1 2 - k^2)} \sqrt{\frac{(c - 1) \left[1 - c/(1 + k_0^2)\right]}{2(1 - c') (1 - c'/c)}} e^{\frac{k}{\pi} \int_0^1 \frac{ds}{1 - ks} (T_c - T_{c'})}$$

= , say, $\frac{C}{k_1 2 - k^2} e^{\frac{k}{\pi} \int_0^1 \frac{ds}{1 - ks} (T_c - T_{c'})}$

G(k) has simple poles at $\pm k_1$ and a branch point at -1. We will therefore be able to write g(x) as

$$g(x) = Ae^{-k_1x} + Be^{k_1x} + j(x), j(x) = 0(e^x) as x - -\infty$$

$$G(k) = \frac{A}{-k - k_{1}} + \frac{B}{-k + k_{1}} + J(k),$$

$$A = -\frac{C}{2k_{1}} e^{-\frac{k_{1}}{\pi} \int_{0}^{1} \frac{ds}{1 + k_{1}s} (T_{c} - T_{c'})}$$

$$B = +\frac{C}{2k_{1}} e^{\frac{k_{1}}{\pi} \int_{0}^{1} \frac{ds}{1 - k_{1}s} (T_{c} - T_{c'})}$$

$$e^{\frac{t}{2}} \frac{k_1}{\pi} \int_0^1 \frac{ds}{1 + k_1 s} (T_c - T_{c'}) = e^{\frac{k_1 2}{\pi}} \int_0^1 \frac{s \, ds}{1 - k_1 2 s^2} (T_c - T_{c'}) + \frac{k_1}{\pi} \int_0^1 \frac{ds}{1 - k_1 2 s^2} (T_c - T_{c'})$$

$$\mathbf{J}(\mathbf{k}) = \frac{\sqrt{c\left[1 - c/(2 + k_0^2)\right]}}{k_1^2 - k^2} \left\{ \frac{k_1 \sqrt{k_1^2 + k_0^2}}{k_0} \sqrt{\frac{C - 1}{2(1 - c')(c - c')}} e^{\frac{K}{\pi} \int_0^1 \frac{ds}{1 - ks} (T_c - T_c^2)} \right\}$$

$$= \frac{c}{k_1} e^{\frac{k_1^2}{\pi} \int_0^1 \frac{s \, ds}{1 - k_1^2 s^2} (T_c - T_{c'}) \sinh k_1 (x + x_2) + j(x), }$$

where

where
$$x_{2} = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1 - k_{1}^{2}s^{2}} (T_{c} - T_{c'}), (x_{1} < x_{2} < 0)$$

$$\frac{k_{1}^{2}}{\pi} \int_{0}^{1} \frac{s \, ds}{1 - k_{1}^{2}s^{2}} (T_{c} - T_{c'}) = -\frac{1}{2} \log \frac{(k_{1}^{2} + k_{0}^{2})(c - 1)[c'/(1 - k_{1}^{2}) - 1]}{k_{0}^{2}(c/c' - 1)2(1 - c')}$$

$$g(x) = \sqrt{\frac{c\left[1 - c/(1 + k_0^2)\right]}{c'\left[c'/(1 - k_1^2) - 1\right]}} \sinh k_1(x + x_2) + j(x)$$
(2.27)

(cf. 2.21)

$$j(x) = \frac{1}{2\pi i} \int_{-j\infty}^{j\infty} dk \ e^{kx} \left\{ \frac{C}{k_1^2 - k^2} \ e^{\frac{k}{\pi} \int_0^1 \frac{ds}{1 - ks} (T_c - T_{c''})} - \frac{A}{-k - k_1} - \frac{B}{-k + k_1} \right\}$$

$$= \frac{C}{2\pi i} \int_{\mathbf{R}'} \frac{dke^{kx}}{k_1^2 - k^2} e^{\frac{k}{\pi} \int_0^1 \frac{ds}{2 + ks} (\mathbf{T}_{\mathbf{C}} - \mathbf{T}_{\mathbf{C}'}) \frac{k_0^2 (k_1^2 - k^2)(1 - \mathbf{C}')}{(k^2 + k_0^2)(c - 1)k_1^2}} \frac{\frac{c}{2k} \log \frac{1 + k}{1 - k} - 1}{1 - \frac{c'}{2k} \log \frac{1 + k}{1 - k}}$$

$$j(x) = \frac{(c - c') C k_0^2 (1 - c')}{2k_1^2 (c - 1)} \int_1^\infty \frac{kdk e^{\frac{kx}{e}} e^{-\int_0^1 \frac{1}{1 + ks} (\mathbf{T}_{\mathbf{C}} - \mathbf{T}_{\mathbf{C}'})}}{(k^2 + k_0^2) \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]}$$

$$j(x) = \frac{k_0 c \sqrt{k_1^2 + k_0^2}}{2k_1} \sqrt{\frac{(1 - c') (1 - c'/c) \left[1 - c/(1 + k_0^2)\right]}{2(c - 1)}}}{2(c - 1)} e^{\frac{k_0 c k}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]}}{e^{\frac{k_0 c}{2}} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]}} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]}}{e^{\frac{k_0 c}{2}} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]}} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2 \left[\left(k - \frac{c'}{2} \log \frac{k + 1}{k - 1}\right)^2 + \left(\frac{\pi c'}{2}\right)^2 \right]} e^{\frac{k_0 c}{2} k_0^2$$

We now have two solutions whose asymptotic forms are:

$$\sin k_0(x + x_1 + \frac{1}{k_0} \tan^{-1} \frac{k_0}{k_1} + \frac{k_1 \sqrt{c \left[1 - c/(1 + k_0^2)\right]}}{\sqrt{k_1^2 + k_0^2} \sqrt{c' \left[c'/(1 - k_1^2) - 1\right]}} e^{k_1(x + x_2)}$$

(cf. equations 2.17, 2.18, 2.22)

$$\sin k_0(x + x_1) - \frac{\sqrt{c \left[1 - c/(1 + k_0^2)\right]}}{\sqrt{c \cdot \left[c'/(1 - k_1^2) - 1\right]}} \quad \sinh k_1(x + x_2)$$

(cf. equations 2.25, 2.26, 2.27)

We introduce the notation,

$$\beta = \sqrt{c \left[1 - c/(1 + k_0^2)\right]}$$

$$\beta' = \sqrt{c' \left[c'/(1 - k_1^2) - 1\right]}$$

$$n_0(x) \xrightarrow{\sqrt{k_1^2 + k_0^2}}{k_1 \beta} \sin k_0 \left(x + x_1 + \frac{1}{k_0} \tan^{-1} \frac{k_0}{k_1}\right) \xrightarrow{e^{k_1(x + x_2)}}{\beta}$$

$$n_1(x) \xrightarrow{\sin k_0(x + x_1)}{\beta} \xrightarrow{\sinh k_1(x + x_2)}{\beta'}$$

 $n_0(x)$ is $\frac{\sqrt{k_1^2 + k_0^2}}{k_1\beta}$ times the "decaying solution" first obtained (2.14 to 2.23). $n_1(x)$ is $1/\beta$ times the "growing solution" next obtained (2.24 to 2.27). Subtracting $k_1n_1(x)$ from $k_1n_0(x)$ gives

$$n_2(x) = k_1 n_0(x) - k_1 n_1(x)$$

$$\frac{\sqrt{k_1^2 + k_0^2}}{\beta} \left(\sin k_0 (x + x_1) \sqrt{\frac{k_1}{k_1^2 + k_0^2}} + \cos k_0 (x + x_1) \sqrt{\frac{k_0}{k_1^2 + k_0^2}} \right) \\ - \frac{k_1}{\beta} \sin k_0 (x + x_1) \\ = \frac{k_0}{\beta} \cos k_0 (x + x_1) - \frac{k_1}{\beta'} \cosh k_1 (x + x_2)$$

If we now subtract $n_1(x)$ from $\frac{n_2(x)}{k_1}$ we get

$$n_{3}(x) = \frac{n_{2}(x)}{k_{1}} - n_{1}(x) - \frac{1}{\beta} \left[\cos k_{0}(x + x_{1}) \cdot \frac{k_{0}}{k_{1}} - \sin k_{0}(x + x_{1}) \right]$$
$$= -\sqrt{\frac{k_{1}^{2} + k_{0}^{2}}{k_{1}\beta}} \sin k_{0} \left(x + x_{1} - \frac{1}{k_{0}} \tan^{-1} \frac{k_{0}}{k_{1}} \right)$$
$$- \frac{1}{\beta} e^{-k_{1}(x + x_{2})}$$

We now have two simple pairs of linearly independent solutions, n(x) and $n_2(x)$; $n_0(x)$ and $n_3(x)$. For any one of these four solutions, hence also for any other solution made from them as linear combinations, the asymptotic solutions on the two sides and the derivatives of the asymptotic solutions have a constant ratio when evaluated at $x = -x_1$ and $x = -x_2$ for the core and tamper solutions respectively.

$$\frac{\text{asymptotic core solution } (x = -x_1)}{\text{asymptotic tamper solution } (x = -x_2)} = \frac{-k_0\beta'}{k_1\beta} = \frac{\text{core solution } (x = -x_1)}{\text{derivative of asymptotic tamper solution } (x = -x_2)}$$

the points, $-x_1$ and $-x_2$, are both on the core side of the interface, $-x_2$ being the farther from the interface. This property leads to the following recipe:

In each medium the asymptotic solution is one of the family of solutions of the equation: $(\Delta + k^2) n(x) = 0$, $\frac{k}{\tan^{-1}k} = c$ (k may be either real or imaginary). Each of the two asymptotic solutions to be joined at an interface is examined at its "fiducial point", distant Δx from the interface on the side of greater c.

$$\Delta x = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1 - k^{2}s^{2}} \left| T_{c} - T_{c'} \right| *$$

(The Δx for each solution uses its own k which may be either real or imaginary.) The two asymptotic solutions, each at its own fiducial point, have equal logarithmic derivatives. The magnitudes of the two solutions, evaluated at their fiducial points, have the same ratio as their values of the quantity,

$$\frac{k}{\beta} = \sqrt{\frac{k^2}{c \left[1 - c/(1 + k^2)\right]}} = \sqrt{\frac{k^2}{c \left[c/(1 - K^2) - 1\right]}} \quad \text{(for } K = ik\text{)}$$

* See Table 3, which gives $c \cdot \Delta X$.

This recipe paraphrases the connection-formulae given above identifying the two asymptotic solutions on the two-sides of an interface. It differs from a simple diffusion theoretic boundary condition connecting the asymptotic solutions only in so far as

1) Δx differs from 0

2) $\frac{k}{\beta}$ differs from a constant

This recipe connects only the asymptotic solutions. Detailed features of the solutions may be gotten from Table 1.

Symbols used in Table 1.

$$T_{c} = \tan^{-1}\left[\frac{n/2}{\tanh^{-1}s - 1/cs}\right]$$
, $T_{c}(0) = \pi$, $T_{c}(1) = 0$

In untamped solution

$$\begin{aligned} x_{0} &= \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1 + k_{0}^{2} s^{2}} T_{c}, \frac{k_{0}}{\tan^{-1} k_{0}} &= c, \beta = \sqrt{c \left[1 - c/(1 + k_{0}^{2})\right]}, c > 1 \\ x_{0} &= \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1 - k_{1}^{2} s^{2}} T_{c}, \frac{k_{1}}{\tanh^{-1} k_{1}} &= c, \beta' = \sqrt{c \left[c/(1 - k_{1}^{2}) - 1\right]}, c < 1. \end{aligned}$$

In tamped (two-medium) solutions the formulae have been written for the case c > 1, c' < 1. Other cases follow by analytic extensions.

$$\frac{k_{0}}{\tan^{-1}k_{0}} = c$$

$$k_{2} = \sqrt{k_{0}^{2} + k_{1}^{2}}$$

$$\frac{k_{1}}{\tanh^{-1}k_{1}} = c'$$

$$\beta = \sqrt{c\left[1 - c/(1 + k_{0}^{2})\right]}$$

$$\beta' = \sqrt{c'\left[c'/(1 - k_{1}^{2}) - 1\right]}$$

$$x_{1} = \frac{1}{\pi} \int_{0}^{1} \frac{ds}{1 + k_{0}^{2}s^{2}} (T_{c} - T_{c'})$$

for $(x_2 < x_1 < 0)$

$$x_2 = \frac{1}{\pi} \int_0^1 \frac{ds}{1 - k_1^2 s^2} (T_c - T_{c'})$$

Each of the four solutions is presented as an asymptotic solution in each medium (sinusoidal or hyperbolic) to which is added a discrepancy term (h(x) for x > 0, j(x) for x < 0). This discrepancy term may be of either sign.

APPENDIX I

ACCURACY OF TWO-BOUNDARY APPROXIMATION

To estimate the error introduced by neglecting the interaction of two boundaries we determine the effect of this neglect in the untamped sphere problem as a first order perturbation. The fundamental eigenvalue, c, of the equation,

$$n(x) = c \int_{-a}^{a} dx' n(x') \frac{1}{2} E(|x - x'|), n(-x) = -n(x).$$
 (i)

we write as $c = c_0/(1 + \epsilon) + 0(\epsilon^2)$, where $a = \frac{\pi}{k(c_0)} - x_0(c_0)$.

The integral operator

$$\int_{-\infty}^{\infty} dx' \frac{c}{2} E(|x - x'|)$$

we denote by Λ .

Write R = R(x) = 0 for x < -a= 1 for x > -a

$$L = L(x) = 0 \text{ for } x > a$$
$$= 1 \text{ for } x < a$$

Equation (i) becomes

$$(1 + \epsilon - \Lambda RL) n(x) = 0, \text{ valid for } -a \leq x \leq a$$

$$n(x) = n_0(x) + n_1(x) \qquad (ii)$$

$$n_0(x) = n_R(x) + n_L(x) - \sin k_0 x$$

where $n_{\mathbf{R}}(\mathbf{x})$ and $n_{\mathbf{L}}(\mathbf{x})$ are the exact one-boundary solutions satisfying

$$(1 - \Lambda R)n_{R} = (1 - \Lambda L)n_{L} = 0$$
$$n_{R}(x) = R \sin k_{O}x + h_{R}(x)$$
$$n_{L}(x) = L \sin k_{O}x + h_{L}(x)$$

Then

$$(1 + \epsilon - \Lambda \operatorname{RL})n_{1} = (\Lambda \operatorname{RL} - 1 - \epsilon)n_{0} = (\Lambda \operatorname{RL} - 1)(n_{R} + n_{L} - \sin k_{0}x) - \epsilon n_{0}$$
$$= \left[\Lambda \operatorname{R} - 1 - \Lambda \operatorname{R}(1 - L)\right]n_{R} + \left[\Lambda \operatorname{L} - 1 - \Lambda \operatorname{L}(1 - R)\right]n_{L}$$
$$-\left[\Lambda - 1 + \Lambda (\operatorname{RL} - 1)\right] \sin k_{0}x - \epsilon n_{0}$$
$$= -\Lambda \left[(1 - L)n_{R} + (1 - R)n_{L} + (\operatorname{RL} - 1) \sin k_{0}x\right] - \epsilon n_{0}$$

$$= -\Lambda \left[(1 - L)h_{R} + (1 - R(h_{L} + (R - RL + L - RL + RL - 1)sin k_{0}x] - \epsilon n_{0} \right]$$
$$= -\Lambda \left[(1 - L)h_{R} + (1 + R(h_{L}) - \epsilon n_{0}) \right]$$
(iii)

Since n_1 must be finite, the right side of (iii) must contain no component, n(x), satisfying (ii). Neglecting terms of order ϵ^2 we have

$$\int_{-a}^{a} dx n(x) \left\{ \Lambda \left[(1 - L)h_{R} + (1 - R)h_{L} \right] + \epsilon n_{0} \right\} = 0$$

$$\epsilon \int_{-a}^{a} dx n_{0}^{2}(x) = -\int_{-\infty}^{\infty} dx RL n(x) \Lambda \left[(1 - L)h_{R} + (1 - R)h_{L} \right]$$

$$= -\int_{-\infty}^{\infty} dx \left[(1 - L)h_{R} + (1 - R)h_{L} \right] \Lambda RL n(x)$$

$$= -\int_{-\infty}^{\infty} dx \left[(1 - L)h_{R} + (1 - R)h_{L} \right] n(x) \quad (iv)$$

The left term of (iv) is roughly 2a. The right term is minus twice the integral of the discrepancy term, $h_{\mathbf{R}}$ (>0) starting from a point distant 2a from its boundary, with n(x) beyond x = a. The character of n(x) in this region may be determined by taking c' = 0 in the decaying two-medium solution. Its value at the surface is

$$\sqrt{\frac{\beta}{2(c-0)}} = \sqrt{\frac{1-c/(1+k_0^2)}{2}}$$

The right term of (iv) will be approximately (-2) x $\frac{1 - c/(1 + k_0^2)}{2}$ h(2a) divided by their combined decay-rate, about 3-4.

For a tamped sphere we proceed in a similar way:

$$\begin{cases} 1 + \epsilon - \Lambda \left[RL + (1 - RL) \frac{c'}{c} \right] \right\} n(x) = 0 \\ n = n_0 + n_1 = n_R + n_L - \sin k_0 x + n_1 \\ \left\{ 1 - \Lambda \left[R + (1 - R) \frac{c'}{c} \right] \right\} n_R = \left\{ 1 - \Lambda \left[L + (1 - L) \frac{c'}{c} \right] \right\} n_L = 0 \\ 1 + \epsilon - \Lambda \left[\frac{c - c'}{c} RL + \frac{c'}{c} \right] \right\} n_1 = \left\{ \Lambda \left[\frac{c - c'}{c} RL + \frac{c'}{c} \right] - 1 \right\} \cdot (n_R + n_L - \sin k_0 x) - \epsilon n_0 \\ = \left\{ \Lambda \left[R + (1 - R) \frac{c'}{c} \right] - 1 \right\} n_R + \Lambda R(1 - L) \left(\frac{c'}{c} - 1 \right) n_R \\ + \left\{ \Lambda \left[L + (1 - L) \frac{c'}{c} \right] - 1 \right\} n_L + \Lambda L(1 - R) \left(\frac{c'}{c} - 1 \right) n_L \\ + \left\{ 1 - \Lambda \left[\frac{c - c'}{2} RL + \frac{c'}{c} \right] \right\} \sin k_0 x - \epsilon n_0 \\ = - \Lambda (1 - L) \left(\frac{c - c'}{c} \right) (R \sin k_0 x + h_R + g_R) \end{cases}$$

$$- \Lambda (1 - R) \left(\frac{c - c'}{c}\right) (L \sin k_0 x + h_L + g_L)$$
$$+ \left\{ 1 - \Lambda \left(\frac{c - c'}{c}\right) RL - \frac{c'}{c} \Lambda \right\} \sin k_0 x - \epsilon n_0$$
$$= (1 - \Lambda) \sin k_0 x - \frac{c - c'}{c} \Lambda \left\{ (1 - L)h_R + (1 - R)h_L \right\} - \epsilon n_0$$
$$= - \left(1 - \frac{c'}{c}\right) \Lambda \left\{ (1 - L)h_R + (1 - R)h_L \right\} - \epsilon n_0$$

Hence as before:

$$\epsilon \sim -\frac{2}{a} \left(1 - \frac{c'}{c}\right) \int dx \, n_0(x) \, \Lambda \left\{ (1 - L) \, h_R + (1 - R) h_L \right\}$$
$$\sim -\frac{2}{a} \left(1 - \frac{c'}{c}\right) \int_a^\infty dx \, n_0(x) \, h_R(x)$$

Estimating this integral in the same way as before gives, for example, for c = 2.0, c' = 1.0,

$$\epsilon - \frac{2}{.72} \times \frac{.5 \times .71 \times .003}{2} \sim .0015$$

For c' = 1 and various values of c, we obtain the estimates:

c	<u> </u>	$\frac{\%}{10}$ in critical radius
1.5	.0002	.09
2.0	.0015	.53
2.5	.003	1.0
3.0	.005	1.3
80	.02	2.0

The chief factor making these errors small is the rapid decay of h(x). Taking the untamped-solution values as typical (they will actually be somewhat too large) it would appear that ϵ will exceed .01 only for core diameters or tamper thicknesses considerably less than one mean free path.

Comparison with variation theory results gives about 0.3 as the limiting thickness for 1 per cent accuracy. (cf. Comparison of variation theory and end point results for tamped spheres, LADC - 77)

APPENDIX II

SOLUTION OF THE INHOMOGENEOUS WIENER-HOPF EQUATION

The Wiener-Hopf technique was shown by E. Reissner (Journal of Mathematics and Physics, Vol. XX (1941), pp 219-223) to permit extension to the inhomogeneous problem. We here treat only the one medium problem with the inhomogeneous term confined to $x \ge 0$. The extension to the two-medium problem with an unrestricted inhomogeneous term is immediately obvious. The equation we wish to solve is:

$$n(x) = \int_{0}^{\infty} dx' \ n(x') \ K(x - x') + f_{1}(x)$$
(a)

where $f_1(x)$ is known and vanishes for x < 0. The Laplace transform of (a), with the notation used previously is,

$$G(k) = F(k) (\underline{K}(k) - 1) + F_{1}(k) = F(k) P(k) + F_{1}(k),$$
(b)
$$F_{1}(k) = \int_{0}^{\infty} dx f_{1}(x) e^{-kx}$$

The solution of the corresponding homogeneous equation will be denoted by a subscript 0.

$$G_{0}(k) = F_{0}(k) P(k)$$
$$P(k) = G_{0}(k)/F_{0}(k)$$

We define $\underline{F}(k)$ such that

$$\mathbf{F}(\mathbf{k}) = \mathbf{F}_{\mathbf{0}}(\mathbf{k}) \mathbf{F}(\mathbf{k})$$

This introduces no singularities in $\underline{F}(k)$ in the right half-plane since $F_0(k)$ had no roots in the right half-plane. Then (b) becomes,

$$\mathbf{F}(\mathbf{k}) \mathbf{P}(\mathbf{k}) = \underline{\mathbf{F}}(\mathbf{k}) \mathbf{F}_{0}(\mathbf{k}) \left(\frac{\mathbf{G}_{0}(\mathbf{k})}{\mathbf{F}_{0}(\mathbf{k})} \right) = \underline{\mathbf{F}}(\mathbf{k}) \mathbf{G}_{0}(\mathbf{k}) = \mathbf{G}(\mathbf{k}) - \mathbf{F}_{1}(\mathbf{k})$$

Thus $-F_1(k)$ is the right-analytic component of $\underline{F}(k)$ $G_0(k)$, which we may write as

$$\left[\underline{F}(k) G_{0}(k)\right]_{R} = \frac{1}{2\pi i} \int_{L} \frac{dk'}{k' - k} \underline{F}(k') G_{0}(k'),$$

where the contour L lies to the left of k and of the singularities of $G_0(k)$ (which are entirely in the right half-plane) and to the right of the singularities of $\underline{F}(k)$ (in the left half-plane).

$$\left[\underline{\mathbf{F}}(\mathbf{k}) \ \mathbf{G}_{\mathbf{0}}(\mathbf{k})\right]_{\mathbf{R}} = -\mathbf{F}_{1}(\mathbf{k})$$
(c)

Making use of the fact that $\frac{1}{G_0(k)}$ as well as $G_0(k)$ is analytic in the left half-plane we can show that equation c is satisfied by

$$\underline{\mathbf{F}}(\mathbf{k}) = -\left[\mathbf{F}_{1}(\mathbf{k}) \frac{1}{\mathbf{G}_{0}(\mathbf{k})}\right]_{\mathbf{R}} \qquad (\mathbf{d})$$

since

$$\begin{bmatrix} G_{0}(k)\underline{F}(k) \end{bmatrix}_{R} = -\begin{bmatrix} G_{0}(k) \begin{bmatrix} F_{1}(k) \frac{1}{G_{0}(k)} \end{bmatrix}_{R} \end{bmatrix}_{R}$$
$$= \frac{-1}{(2\pi i)^{2}} \int_{L'} \frac{dk'}{k'-k} G_{0}(k') \int_{L''} \frac{dk''}{k''-k'} \frac{F_{1}(k'')}{G_{0}(k'')}$$
$$G_{0}(k) \underline{F}(k) \end{bmatrix}_{R} = -\frac{1}{(2\pi i)^{2}} \int_{L''} dk'' \frac{F_{1}(k'')}{G_{0}(k'')} \int_{L'} dk' G_{0}(k') \frac{1}{k''-k} \left(\frac{1}{k'-k} + \frac{1}{k''-k'} \right)$$



Displacing the contour L' to the left of L" picks up a residue at k' = k''. The remaining k' integral vanishes as it may be displaced indefinitely to the left, in which direction the integrand decays as

 $\frac{1}{|\mathbf{k}'|^2}$. This leaves:

$$\begin{bmatrix} G_0(k) \quad \underline{F}(k) \end{bmatrix}_R = -\frac{1}{(2\pi i)^2} \int_{\underline{J}_{\perp}''} dk'' \quad \frac{F_1(k'')}{G_0(k'')} \left(\frac{2\pi i}{k'' - k} \cdot G_0(k'') \right)$$
$$= - \begin{bmatrix} F_1(k) \end{bmatrix}_R = -F_1(k)$$

The particular integral of equation a has therefore the Laplace transform

$$\mathbf{F}(\mathbf{k}) = -\mathbf{F}_{O}(\mathbf{k}) \left[\frac{\mathbf{F}_{1}(\mathbf{k})}{\mathbf{G}_{O}(\mathbf{k})} \right]_{R}$$

To this may be added any multiple of the homogeneous solution, $F_0(k)$.

To extend this method of solution to the two-medium problem requires only the replacement of equation a by the corresponding two-medium equation. This leaves the form of equation b and the rest of the solution unchanged. To treat an inhomogeneous term existing for both x > 0 and x < 0 it suffices to break up the inhomogeneous term into a right and a left side part and treat each separately as above.

A particularly simple special case of the untamped inhomogeneous equation is that of the albedo problem —



Then

$$\begin{bmatrix} \underline{F_{1}(k)} \\ \overline{G_{0}(k)} \end{bmatrix}_{R} = \frac{1}{2\pi i} \int_{L} \frac{dk}{k' - k} \frac{1}{(k' + \alpha)G_{0}(k')}$$
$$= \frac{1}{G_{0}(-\alpha)(k + \alpha)} + \frac{1}{2\pi i} \int_{L'} \frac{dk'}{(k' - k)(k' + \alpha)G_{0}(k')}$$

In the second term the contour $L^{'}\,$ may be displaced indefinitely to the left. Its integrand may be written as

$$\frac{\text{Const.}}{\mathbf{k}'} + 0\left(\frac{1}{\mathbf{k}'}\right)$$

Thus the k-dependent part of the integral vanishes. The constant part represents an admixture of the homogeneous solution to $F_1(k)$ and therefore may be disregarded. The general solution is therefore

$$\mathbf{F}(\mathbf{k}) = -\mathbf{F}_{O}(\mathbf{k}) \left(\begin{bmatrix} \mathbf{F}_{1}(\mathbf{k}) \\ \mathbf{G}_{O}(\mathbf{k}) \end{bmatrix}_{\mathbf{R}} + \mathbf{A} \right) = -\mathbf{F}_{O}(\mathbf{k}) \left(\frac{1}{\mathbf{G}_{O}(-\alpha)(\mathbf{k} + \alpha)} + \mathbf{A} \right).$$

In an albedo problem c will be ≤ 1 and A should be chosen to make n(x) finite for all x, hence F(k) regular at $k = +k_1$, despite the pole of $F_0(k)$. Thus

$$A = -\frac{1}{G_0(-\alpha)(k_1 + \alpha)}$$

$$\mathbf{F}(\mathbf{k}) = \frac{(\mathbf{k} - \mathbf{k}_1)\mathbf{F}_0(\mathbf{k})}{(\mathbf{k} + \alpha)(\mathbf{k}_1 + \alpha)\mathbf{G}_0(-\alpha)}$$

The density of emergent neutrons in the albedo problem as a function of μ , the cosine of the angle of emergence, is

$$N(\mu) = c \int_0^\infty dx \ n(x) e^{-x/\mu}$$
$$= c F \frac{1}{\mu}$$

and is therefore given directly by the solution F(k).

	•
ш	ł
-10	

					IAB
SOLUTION	f(x) - f(x) Asymptotic Core solution	g(x) - j(x) Asymptotic TAMPER SOLUTION	n(o) = f(o) = g(o) Value of Solution At Interface	-H(O) = - J ^{oo} h(x)dx Area of discrepancy term in core	- <u>H'(o)</u> = <u>f</u> xh(x)dx H(o) f h(x)dx MEAN LENGTH OF DISCREF IN COPE
(>۱	Sin K _o (x+x _o)		<u>्र</u> उद्ध	$-\frac{1}{k_o} \left[\frac{\beta}{\sqrt{2c(c-1)}} - \cos \kappa_x \right]$	$\frac{1}{H(0)} \frac{1}{\kappa_{o}^{2}} \left[\frac{\sin \kappa_{o} x_{o}}{\sin \kappa_{o} x_{o}} - \frac{\beta}{\kappa_{o} \frac{\beta}{\sin \kappa_{o} x_{o}}} \frac{1}{\pi} \int_{0}^{1} dt \right]$
UNTAMPED					
CASE C<1	Sinfi K,(x+z.)		B'	$\frac{1}{\kappa_1} \left[\frac{\beta'}{\sqrt{2c(1-c)}} - \cosh \kappa_1 x_1 \right]$	$\frac{-1}{H(0)\kappa_{i}^{2}}\left[\begin{array}{c} \text{SLmh }\kappa_{i}x_{0}\\ -\kappa_{i}\frac{\beta'}{\sqrt{2c(1-c)}}\frac{1}{\pi}\int_{0}^{b}d^{2}\end{array}\right]$
n,	$\frac{k_{\lambda}}{K_{\beta}} \sin k_{0}(x+z_{0})$ $= \frac{1}{\beta} \sin k_{0}(x+x_{1})$ $+ \frac{K_{0}}{K_{\beta}} \cos k_{0}(x+x_{1})$	$\frac{1}{\beta'}e^{\kappa_i(z+x_a)}$	<u>κ</u> 2 Κ ₁ √2(c-c')	$\frac{K_{L}}{K_{1}K_{1}\beta}\left[\cos K_{0}\chi_{0}-\beta \sqrt{\frac{(1-c')}{2(c-1)(c-c')}}\right]$	$\frac{\kappa_{a}}{(-H(o))\kappa_{i}\kappa_{o}^{2}} \left[\frac{\kappa_{o} \left(\frac{1-c^{i}}{2} \right)}{\kappa_{i} \sqrt{2(c-i)(c-c^{i})}} \right] + \frac{1}{\beta} Sm \kappa_{o} \left(x_{i} + \frac{1}{\kappa_{o}} tan \right)$
$\mathcal{N}_{3} = \frac{n_{s}}{\kappa_{i}} - n_{i}$	$\frac{K_{\bullet}}{K_{i}\beta} \cos K_{\bullet}(z+z_{*})$ - $\frac{1}{\beta} \sin K_{\bullet}(z+z_{i})$ = $\frac{-K_{i}}{K_{i}\beta} \sin K_{i}(z+z_{i} - \frac{1}{K_{i}} \log \frac{K_{i}}{K_{i}})$	$\frac{1}{\beta'} e^{-\kappa_1(x+x_2)}$	<u> </u>	$\frac{K_{\lambda}}{K_{1}K_{0}\beta} \left[\beta \sqrt{\frac{(j-c')}{2(c-i)(c-c')}} - \cos \kappa_{0}(x_{1} - \frac{L}{K_{0}} \tan^{-1} \frac{K_{0}}{K_{1}}) \right]$	$\frac{K_{2}}{-H(o)K_{0}^{2}K_{1}}\left[\frac{K_{1}}{K_{1}}\sqrt{\frac{(1-c')}{2(c-1)(c-c')}}\left(1-\frac{K_{1}}{\pi}\right]_{0}^{2}\right]$ $+\frac{1}{\beta}Sun K_{1}\left(\chi_{1}-\frac{1}{K_{1}}+a\right)$
n,	$\frac{1}{\beta}$ Sim K.(2+2,)	<u>΄</u> , sinh κ _i (z+z ₂) β	0	$-\frac{1}{\kappa_0 \beta} \left[\frac{\kappa_1 \beta}{\kappa_1} \sqrt{\frac{(1-c')}{2(c-1)(c-c')}} - \cos \kappa_2 \right]$ (negative, i.e. H(o)>0)	$\frac{\frac{1}{K_{o}^{2}H(o)}\left[\frac{Sin K_{o} X_{i}}{\beta} - \frac{K_{o} K_{z}}{K_{i}} \sqrt{\frac{(i-c')}{2(c-i)(c-c')}} \frac{1}{\pi} \int_{0}^{t} ds \left(\frac{1}{2(c-i)(c-c')} - \frac{1}{\pi} \int_{0}^{t} ds \right) ds} ds$
N2 = K1(N-N1)	$\frac{k_{\circ}}{\beta}$ cor $k_{\circ}(z+z_{\circ})$	$\frac{\kappa_1}{\beta}$ cosh $\kappa_1(z+z_2)$	$\frac{k_z}{\sqrt{2(c \cdot c')}}$	- <u>Sim Kox</u> , B (positive, i.e. H(0)<0)	$\frac{-1}{SUn K_{o} \mathbf{x}_{i} K_{o}} \left[\frac{\beta \mathbf{x}_{\lambda}}{K_{i}} \sqrt{\frac{(1-\epsilon^{\prime})}{2(\epsilon-i)(\epsilon)}} - \cos K_{o} \mathbf{x}_{i} \right]$

TABLE	T	AECD - 2	2056	
0) = Jox h(x)dx b) = Joh(x)dx LENGTH OF DISCREPANCY IN COPE	$J(u) = \int_{0}^{u} j(x) dx$ AREA OF DISCREPANC IN TAMPER	$\frac{J'(\bullet)}{J(\bullet)} = \frac{\int_{-\infty}^{\infty} x_j(x) dx}{\int_{-\infty}^{\infty} j(x) dx}$ When LENGTH - OF DISCREPANCY IN TANK	K(x) DISCREPANCY (NEGATIVE) IN CORE (z≥0)	j(Z) DISCREPANCY (POSITIVE) IN TAMPER (ZSO)
$\frac{1}{2} \int_{K_{o}}^{R_{o}} \left[\frac{\sin \kappa_{o} \chi_{o}}{\sqrt{2 c(c-1)}} \frac{1}{\pi} \int_{0}^{t} ds T_{c} \right]$			$-\frac{\beta}{2\kappa_{v}}\sqrt{\frac{c(c-1)}{2}}\int_{1}^{\infty}\frac{\kappa d\kappa}{\left(\frac{c}{2}\log\frac{\kappa+1}{\kappa-1}-\kappa\right)^{2}+\left(\frac{\pi}{2}c\right)^{2}}}e^{\kappa d\kappa}$	K
$\frac{1}{\sigma)\kappa_{i}^{2}}\left[sinh \kappa_{i} x_{o} \\ \frac{\beta'}{\sqrt{2c(1-c)}} \frac{1}{\pi} \int_{0}^{1} ds T_{c} \right]$			$\frac{\beta'}{2\kappa}\sqrt{\frac{c(1-c)}{2}}\int_{\sqrt{\frac{c}{2}\log\frac{K+1}{K-1}-\kappa}}^{\infty}\frac{e^{-\frac{\kappa}{2}\int_{-\frac{K}{2}}^{\frac{1}{2}\frac{K}{K-1}}\frac{ds}{k}}}{\int_{-\frac{K}{2}}^{\infty}\frac{e^{-\kappa}}{k}}e^{-\kappa}$	
$= \frac{1}{\beta} Sim K_o \left(x_1 + \frac{1}{K_o} \tan^2 \frac{K_o}{K_i} \right) $	$\frac{k_{2}}{\kappa_{1}\kappa_{o}}\sqrt{\frac{C-1}{2(1-C')(C-C')}}$ $-\frac{1}{\kappa_{1}\beta'}e^{\kappa_{1}x_{2}}$	$+\frac{\frac{1}{\pi}\int_{0}^{1} ds(T_{c}-T_{c})}{1-\frac{K_{o}}{K_{A}\beta'}\sqrt{\frac{2(1-c')(cc')}{(c-1)}}}$	$\frac{K_{2} (c-1)(k-C')}{2 K_{1} (2(1-C'))} \int_{k+K_{1}}^{\infty} \frac{-\frac{k}{\pi} \left(\frac{1}{2} + K_{2} - T_{1} - T_{1}\right)}{k+K_{1} \left(\frac{2}{2} \log \frac{k+1}{k-1} - k\right) + \left(\frac{\pi}{2}\right)^{3}} e^{-Kx}$	$\frac{K_{0}K_{1}}{2K_{1}^{4}\sqrt{2(e^{-1})}}\int_{1}^{\infty} \frac{k_{0}(K_{1}+K_{1})}{(K^{2}+K_{2}^{2})\left(\frac{C}{2}\int_{0}^{1}\frac{ds}{K^{2}}\left(\frac{1}{2}\cdot\frac{T}{2}\right)\right]}$
$ \frac{K_{o}}{K_{i}} \frac{(1-C')}{2(C-i)(C-C')} \left(1 - \frac{K_{i}}{\pi} \int_{0}^{1} ds \left(T_{c} - \overline{L}\right) \right) $ $ + \frac{1}{\beta} \sin K_{o} \left(x_{i} - \frac{1}{K_{o}} + \cos^{-1} \frac{K_{o}}{K_{i}}\right) $	$\frac{e^{-\kappa_{i} z_{2}}}{\kappa_{i} \beta'}$ $-\frac{\kappa_{2}}{\kappa_{i} \kappa_{i}} \sqrt{\frac{(c-i)}{2(i-c')(c-c')}}$	$-\frac{1}{k_{1}}$ $+\frac{\frac{1}{\pi}\int_{a}^{b} ds(T_{c}-T_{c})}{\frac{1}{k_{2}}\sqrt{\frac{2(1-c)(-c)}{(c-1)}}e^{-k_{2}}}$	$\frac{k_{1}\left(\overline{k}-1\right)\left(\frac{1}{2}\right)}{2K_{0}\left(2\left(1-C^{2}\right)\right)}\left(\frac{kdK}{K}\right)\left(\frac{e}{2}\right)\left(\frac{kdK}{2}\right)\left(\frac{e}{2}\right)\left(\frac{kdK}{2}\right)\left(\frac{e}{2}\right)\left(\frac{kdK}{2}\right)\left(\frac{e}{2}\right)\left(\frac{kdK}{2}\right)\left(\frac{e}{2}\right)\left(\frac{kdK}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right)\left(\frac{e}{2}\right$	$\frac{k_{2}}{2k_{1}^{2}} \frac{\left(1-c^{2}\right)\left(c-c^{2}\right)\left(\frac{k_{2}}{2k_{1}^{2}}\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k_{2}}{2k_{1}^{2}}\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\right)\left(\frac{k_{2}}{2k_{1}^{2}}\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left(\frac{k-k_{2}}{2k_{1}^{2}}\right)\left$
$\overline{I}_{0}\left[\begin{array}{c} \frac{Sun K_{n}X_{1}}{\beta}\\ \sqrt{\frac{(1-c')}{2(c-1)(c-c')}} & \frac{1}{\pi}\int_{0}^{t}ds\left(T_{c}-T_{c'}\right)\right]$	$\frac{k_{2}}{K_{1}K_{2}}\sqrt{\frac{(c_{-i})}{2(i-c')(c-c')}} - \frac{\cosh k_{1}x_{2}}{k_{1}\beta'}$	$\frac{1}{J(0)} \frac{\left[\frac{K_{2}}{K_{1}} + \frac{(c-1)}{(c-1)}\right]}{\int_{0}^{1} \left[\frac{K_{2}}{K_{1}} + \frac{1}{2(1-c)} + \frac{1}{c}\right]} = \frac{1}{T} \frac{1}{d_{3}} \left[\frac{1}{(c-T_{c})} - \frac{Such K_{2}}{K_{1}^{2} \beta^{2}}\right]}{K_{1}^{2} \beta^{2}}$	$\frac{K_{k}K_{1}}{2K}\left[\frac{c_{*}}{(1-c')}\right]_{k}^{\infty} -\frac{\kappa}{4}\left[\frac{1}{2}\frac{k_{k}}{(1+\kappa)}\left(\frac{1}{2}-T_{c'}\right) +\kappa \left(\frac{K_{k}K_{1}}{2}\right)\right]_{k}^{\infty} -\frac{\kappa}{4}\left[\frac{K_{k}K_{1}}{2}\frac{K_{k}}{(1-\kappa)}\left(\frac{\pi}{2}\right)\right]_{k}^{\infty}\right]_{k}^{\infty}$	$\frac{k_{k_{k}}\left[1-c^{2}\right]\left(-c^{2}\right)\left[\frac{k_{k}dk}{k_{k}}\right]\left(\frac{k}{2}+\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}}{k_{k}}\right)\left(\frac{k_{k}$
$\frac{1}{\kappa_{1}} \left[\frac{\beta \kappa_{2}}{\kappa_{1}} \sqrt{\frac{(1-c')}{2(c-1)(c-c')}} - \cos \kappa_{0} x_{1} \right]$	_ <u>simh k,x.</u> ß'	$\frac{-1}{\sinh k_{i} z_{i} \cdot k_{i}} \left[\frac{\beta k_{i}}{k_{i}} \frac{(c-1)}{k_{i} \cdot k_{i} \cdot k_{i}} - \cosh k_{i} z_{i} \right]$	$\frac{k_{1}K_{1}}{2K_{0}} \sqrt{\frac{(c-1)(c-c')}{2(1-c')}} \left(\frac{\kappa^{2}dK}{\kappa^{2}dK} \frac{e}{e^{\frac{1}{2}\int_{0}^{1} \frac{dK}{1+K_{0}}(T_{c}-T_{c})}}{\frac{\kappa^{2}}{2(1-c')} \int_{0}^{\infty} \frac{\kappa^{2}}{\kappa^{2}} \frac{\kappa^{2}}{k_{0}} \frac{\kappa^{2}}{\kappa^{2}} \frac{\kappa^{2}}{\kappa^{2}} \frac{\kappa^{2}}{k_{0}} \frac{\kappa^{2}}{\kappa^{2}} \frac{\kappa^{2}}{k_{0}} \frac{\kappa^{2}}{\kappa^{2}} \frac{\kappa^{2}}{k_{0}} \frac{\kappa^{2}}{\kappa^{2}} \frac{\kappa^{2}}{k_{0}} \frac{\kappa^{2}}{\kappa^{2}} \frac$	$\frac{k_{L_{2}}}{2k_{1}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}}{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}}{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac{\left[1-\frac{k_{1}}{2}\right]^{2}} \frac$

1.2

2072

0	0.2	0.4	0.6	0.8	1.0	1.2	1.4	1.6	106 (1 + E) E
	Gitotte.	.86583	1008.	.74951	Stiot?	•67963	.65468	.63408	1.00000
	-94262	01173.	. 80785	. 75865	.720 ⁴⁴	60069.	.66541	.64493	19116.
	6£m16.	.87541	81418.	.76623	.72875	48869.	.67m	.6540 7	SIL43.
	-94588	.87903	13618.	•77264	.73584	.70633	.68215	46199 .	45537.
	61749.	.88220	41428.	.77823	.74202	.71268	.68893	.66885	.73473
	.94833	16433	.82818	418314	.74746	.71867	£6h169°	66 1 19.	.69315
	69056.	. 89056	611 953.	. 79323	.75873	010£1.	-707 ⁴⁶	.68784	.61086
	.95256	16tt68.	.84301	. 80120	.76767	62011.	SHLTT.	91369.	12 643.
	30426.	.89856	. ⁸⁴⁸ 33	.80773	.17502	.74820	.72579	.70674	.50111
	-95537	12106.	. 852 80	.81323	42187.	26427.	.13287	Totil.	.46210
	-95746	04906.	.85998	.82213	0£162.	.76583	- 74439	.72603	.140236
	51626 .	\$1016.	.86559	.82906	. 79922	77447	.75353	.73556	.35835
	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	0,0000

.61896 .60881 .599 .63221 .62204 .612 .64297 .63282 .62; .65200 .64187 .63;	.6432 6461	.65797 .6432 .67102 .6564
.65977 .64967 .64 жа. 73645 .645 базар		10+00. 00100. +0100. 65533. 15763. 01101. 16835. 01011.
67100 58370		#7100. 1876д. 1871017.

Table 2. (continued) $\frac{k}{1106(1+k)} \int_{0}^{1} \frac{40}{1+k^2} t^2$

AECD - 2056

	1.0	.28954 .26236	-23002	19298	.15537	.12058	12630.	.06269	.03906	.01831	0.0000	01628	03083	04393	05577	06654	07638	08541	09373	- 10142	- 10656	11519	12139	- 12715	- 13262	13774	14256	01741	15140	15545	- 15934	33043
	0.0	-28937 -26056	.22578	.18551	.14470	.10731	surve.	.011605	.02143	0.0000	01875	03537	05012	06331	07520	08597	8 1560° -	10μ7u	76211	12056	- 12758	11hft	610H1	- :158	- "TETT	15616	16084	16527	16944	17338	17713	33852
	0.8	.258674 .25802	.22025	.17595	13120	39060.	.05563	.02565	0.0000.0	02211	16140	05816	07304	08627	41360	10584	11854	12738	14561	16241	826h1	15615	16206	16756	17271	17753	18207	- 18632	- .19035	41461	12261	34855
1	1.0	.28732 .25435	.21280	.16326	.11350	.06922	.03162	0.0000	02669	14640	06896	46380	10084	- 11402	12576	13629	14581	15444	16233	16955	17619	18232	18802	- 19332	42391	20286	erros	21125	21507	21868	22209	36130
D	0.6	.23455 .24853	20229	.19241.	12680-	9110110.	0.0000	03328	68090	101180° -	10377	12072	13547	14841	- 15989	110/1	- 17932	18762	19516	- 20208	20841	21425	21964	22464	22929	23363	23770	24151	24509	74842	25166	37800
	0.5	.27934 .24020	.18661	34611.	.05434	0.0000		07818	10638	19621	21641	16576	18006	19252	20348	21321	22190	22973	23682	24326	- <u>1916</u>	25457	25951	26419	268119	27248	27621	01612	26295	28606	28897	060011
	7.0	.2691 19 .22591	16116	.07723	00000*0	06033	10618	14156	16951	- 19212	21080	64922	23988	25144	26154	27045	- 27538	28548	29165	29768	76208	30782	31226	31635	- 32020	32374	+0722	33012	33301	33572	33828	43383
	5.0	.25068	J1416	0.0000.0	06160	15625	20195	23571	26164	28217	29887	31272	J2443	33WF	- ,34318	35082	35757	36360	36901	37390	37834	382µ1	38613	38956	39272	- 39566	01366 -	40095	40334	40558	40769	48435
	0.2	.21486	0.0000.0	16125	25698	31537	35418	<u>38185</u>	140256	17814	43167	44235	45130	- 45892	46549	42124	47630	148080	48483	14884° -	9/164	- 19476	19751	50003	50236	50453	50653	50839	51014	51177	51331	56794
	1.0	.14611 0.00000	71945		53520	57007	59264	61809	62024	62935	63661	64256	64754	65176	 .65541 	65857	66136	66383	66604	66803	66983	74179	67297	67435	67561	67679	67788	67889	12629	68073	62156	71063
	0	0-00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1.00000	-1-00000	-1-00000	-1.00000	-1.00000	-1-00000	-1.00000	00000-1-	-1.00000	00000. [-	-1.00000
° /		001	0.2	0.3	# •0	0.5 0	0.6	0.7	0.5	6°0	1.0	1.1	1.2	2.3	1.L	1.5	1•6	1.7	1.6	1.9	5.0	2.1	2°3	2.3	2.4	2,5	2.6	2.7	2.8	2.9	0.5	8

Table 3. $c \Delta X(c,c^{\prime}) = \frac{c}{\pi} \int_{0}^{1} \frac{T_{c}^{\prime} - T_{c}}{1 - k^{2} s^{2}} ds$

AECD - 2056

°/-,	1.0	1.1	1.2	1.1	T .L	1.5	1.6	777	1.8	1.9	210
0	45635.	24682.	21695.	.28870	.25523	-28769	.28713	.28659	.28601	.28546	-28492
0.1	.26236	.26368	.26461	.26531	-26583	-26619	.26644	.26663	.26674	.26681	.26685
0.2	-23002	-23338	.23608	-23827	21012.	.24167	762#5.	01442.	.24506	.24592	.24667
ŗ.	36261.	36361.	-20389	20799	THILS.	11112.	10/15.	.21928	.22126	.22305	.22463
† .0	.15537	16401	.17115	11711.	.18232	.18675	-19062	confer.	70701.	87978	.2023
0.5	.12058	H1211	.14050	71841.	.15477	.16051	.16553.	.17000	19571.	.17755	.18078
0.6	11630.	10228	.11254	.12186	.12966	.13648	14241.	.14763	.15260	.15692	.16084
0.7	.06269	.07653	.08825	26360.	10706	42411.	HSISI.	.12762	.13308	13802	.14252
0.8	.03906	.05384	.06643	62170.	.08679	.09516	.10260	12601.	.11529	.12075	.12572
6°0	.01831	.03376	.04700	0584g	.06857	61220-	.08544	.09261	20660.	.10496	.11035
1.0	0,0000	-01594	. 029650	09Th0.	.05214	.06148	.069855	14270.	.08425	03060.	.09622
1.1	01628	00000	Lotto.	.02637	.03725	16910.	.05564	.06351	.07066	-07720	.08320
1.2	03083	01432	0,00000	.01257	.02372	.03367	.04263	.05076	.05816	.06193	.07115
 	04393	02727	01276	0.0000	.01135	.02151	.03068	10620.	.04662	.05360	.06003
1.4	05517	03903	01120	01150	0.0000	-01032	.0 <u>1</u> 966	.02316	.03593	707 NO.	.04966
	06654		03505	02205	01045	0.0000	91(600.	608TO-	.02600	.03328	10010-
1.6	07638			etten -	02010	12600	0.00000	£1800.	.01675	.02413	16010.
1.7 1.7	08541.	06841	05389	08040	02905	01846	00881	0,00000,0	.00810	.01558	.02250
1.6		07677	06226	416tio	03738	02674	40/10° -	00817	0.00000	.00755	.01456
1.9	54101	08452	- 0200 ⁴	05693	04515	84420	02 ¹ 175	01582	00761	0.0000	.00706
2°0	10856	09172	07729	06419	- ,052 ¹ 12	0 ¹¹⁷³	03198	02301	01476	11700	0.0000
2 •1	11519	09845	08405	66010	52630	p4825	03877	03620	02151	01383	00668
N N	12139	51401	01060		06563	05467	04519	03619	02789	02019	01300
2 • 3	12718	11062	09635	08338	07166	1/090	05123	04224	493394	02621	19610
ੜਾ ਨ	13262	11615	10195	08903	07736	06643	05697	36140° -	03966	49120	17450
ы Ц	+177t	12136	- "10724	09438	08274	48120	06240	05341	04510	03737	03013
2 - 6	14256	12627	42211° -	Et1660	08784	07697	06755	05858	05028	04255	03532
2.7	01741	1093	- 11696	10422	09267	48180	072W5	06350	05522	81740	04025
2.8	15140	- 13533	HH121	10877	09728	05648	11/10	06819	- °05992	05220	161110 -
6,5 6	15548	13950	12570	11309	10165	06060	08157	07266	14490	05671	Grifono
0,0	15934	14347	52621	11720	10582	09512	08582	+169L0° -	06871	06102	05351
8	33043	32352	31811	31331	30916	29525	30235	29953	10262	n2n62	29267

Table 3. (continued) $c \Delta \mathbf{x}(c,c') = \frac{c}{\pi} \int_{0}^{c} \frac{\mathbf{r}_{c'} - \mathbf{r}_{c}}{1 - \mathbf{r}^{2} \mathbf{r}^{2}} d\mathbf{r}$

aecd - 2056

1

,

i,

,

8	25000	25000	.25000	.25000	.25000	.25000	.25000	.25000	.25,000	-25000	.25000	25000	-25000	.25000	23000	.25000	.25000	25000	-25000	.25000	25000	-25000	.25000	5,000	25000	.25000	.25000	-25000	25000	800 K.	0.0000	
3.0	-28005	25080	23436	entra.	-20162	.18642	.17222	15898	.14665	13514	.12437	TSULL.	.10480	C05282	-08744	34670.	16170.	.06475	.05793	.05143	togeto.	CE660.	-03367	.02526	.02307	01808	.01330	00870	.00426	0.0000	27933	
2.9	64082.	.25055	.23372	.21664	41005.	.18459	10011.	.15656	66241.	.13227	.12132	SOLLI.	.10143	TE260.	.08384	.07576	.06813	.06038	.05399	.04745	12110.	.03525	.02955	11450.	.01855	.01388	10600.	Sutton.	0.0000.0	00428	26021	
2.8	28095.	25029	.23304	.21551	09861.	.18267	.16782	.151:02	61141.	.12925	01311.	.10767	06160.	.08871	.08006	.07188	.064 16	.05683	.04988	.04328	16960-	76070.	.02523	51610.	.01450	24600.	19100.	0.0000.0	00HHG	00875	28127	
2.7	28137	10040	23224	.21426	06961.	.18058	.16539	.15128	.13818	.12601	.11466	.10406	51460.	03450.	.07604	.06777	.05995	.05256	.04554	03887	-03253	.02648	17050.	.01519	16600.	.00 ¹⁴⁸⁵	0.0000	00465	J1600	111ETO" -	28236	
2.6	.28185	19046	14122	16212.	11261.	.17836	.16280	.14837	.13500	.12259	foill.	.10025	.09016	.08070	-07182	.06344	.05554	.04805	16010.	42420.	.02785	-02175	46510.	eto10-	.00508	0.0000	00 ¹ 87	alaoco -	hon -	01634	28350	
2.5	-28232	100002-	23054	.21151	9191.	.17600	.16003	14527	.13161	.11893	91701.	619 60.	46580.	.07634	.06733	.05885	.05085	62840.	.03613	JE 620.	0220.	.01676	16010.	1000.	0.0000	00510	00999	01467	T1910	02350	47485	
2.4	-28280	20002. 128240	22955	10000	19108	.17341	15704	16141.	.12795	.11502	10701.	78160.	.08143	.07168	.06255	.05396	.04587	03823	10110.	71420.	.01767	6tilo.	.00561	0.0000	00536	34010	01538	02007	02458	02841	28607	
2.3	-28331	-200/4	22850	20827	18884	.17066	.15384	.13835	12404	.11083	-09859	.08723	.07663	.06673	.05748	.04878	09010	.03287	02559	.01868	412I0.	-00592	0.0000	00563	10110	01616	02107	02578	P2070	01463	28752	
2.2	-26382	200790	22732	200	.18638	.16765	15037	13417	.11982	.10632	132.60	.08225	01110.	.06144	.05204	42540.	03146	02716	13910.	.01285	.00626	00000*0		01161	10/10	02216	02710	18150	07672	01066	28910	
2.1	-28H36	20053	20405	20 FH400	1877	01101	.14661	17028	.11528	THIOL.	.08573	07691	06597	.05577	.04625	11110-	.02895	02108	-01165	00664	0.0000.0	00629	01227	01795	02738	02854	34220 -	P1850	02.00	- 01704	03062	
2.0	26492.	-26685	10042.	10000	1K07K	16084	11252	.12572	21011.	09622	.08320	.07115	0090	04966	TOONO	10010	02250	01456	00100	00000-0	00668	01300	19610	17450	0101	03532	01025			- OFIRI	29267	
°/-,	0	1.0					2.0		0									- 1			1		2		i n			- 0	0.0		2 8)

END OF DOCUMENT

Table 3. (continued) $c \Delta x(c,c^{*}) = \frac{c}{\Pi} \int_{0}^{1} \frac{\tau_{c^{*}} - \tau_{c}}{1 - \kappa^{2} \kappa^{2}} ds$

3

AECD - 2056